YY2/BUB3 axis-mediated SAC hyperactivity determines tumor cell fate through chromosomal instability

Rendy Hosea[1,2], Wei Duan[1,2], Ian Timothy Sembiring Meliala[1,2], Wenfang Li[1,2],

Mankun Wei[1,2], Sharon Hillary[1,2], Hezhao Zhao[3], Makoto Miyagishi[4], Shourong Wu[1,2,5]*,

Vivi Kasim[1,2,5]*

[1]Key Laboratory of Biorheological Science and Technology, Ministry of Education, College of Bioengineering, Chongqing University, Chongqing 400044, China;
[2]The 111 Project Laboratory of Biomechanics and Tissue Repair, College of Bioengineering, Chongqing University, Chongqing 400044, China;
[3]Department of Gastrointestinal Surgery, Chongqing University Cancer Hospital, Chongqing University, Chongqing 400030, China;
[4]Life Science Innovation, School of Integrative and Global Majors, University of Tsukuba, Tsukuba, Ibaraki 305-0006, Japan;
[5]Chongqing Key Laboratory of Translational Research for Cancer Metastasis and Individualized Treatment, Chongqing University Cancer Hospital, Chongqing University, Chongqing 400030, China.

*Correspondence should be addressed to:
Vivi Kasim, College of Bioengineering, Chongqing University, Chongqing, China; Phone: +86-23-65112672, Fax: +86-23-65111802,
Email: vivikasim@cqu.edu.cn;

Shourong Wu, College of Bioengineering, Chongqing University, Chongqing, China; Phone: +86-23-65111632, Fax: +86-23-65111802,
Email: shourongwu@cqu.edu.cn

Running title: YY2/SAC activation determines tumor cell fate through CIN

Keywords: Chromosomal instability (CIN); drug resistance; mitosis; mitotic error; residual tumor cells; spindle assembly checkpoint (SAC); tumor cell fate; yin yang 2 (YY2)

Abstract

Spindle assembly checkpoint (SAC) is a crucial safeguard mechanism of mitosis fidelity, which is fundamental for equal division of duplicated chromosomes to the two progeny cells. Impaired SAC can lead to chromosomal instability (CIN), a well-recognized hallmark of cancer that facilitates tumor progression; paradoxically, high CIN levels are associated with better drug sensitivity and prognosis. However, the mechanism by which CIN determines tumor cell fates and drug sensitivity remain poorly understood. In this study, using a cross-omics approach, we identified YY2 as a mitotic regulator that peaks at M phase and promotes SAC activity by positively regulating the transcriptional activity of *budding uninhibited by benzimidazole 3* (*BUB3*), a component of SAC. While inducing CIN, YY2/SAC activity defect enhanced mitosis and tumor growth, whereas YY2/SAC hyperactivation, as a result of *YY2* overexpression, triggered mitotic delay and growth suppression. Furthermore, we revealed that excessive CIN, caused by either *YY2* overexpression or further inhibiting SAC activity in *YY2-* knocked out cells, leads to higher cell death rates. However, residual tumor cells that survived DNA damage-based therapy had moderate CIN and increased drug resistance; meanwhile *YY2* overexpression in these cells sensitizes them to DNA-damage agents. Hence, this study provides insights into the regulatory mechanism of SAC activity as well as the role of YY2/BUB3 axis, SAC activity, and CIN levels in determining tumor cell fate. Furthermore, this study also links up tumor cells drug resistance with moderate CIN, and suggest a novel anti-tumor therapeutic strategy that combines SAC activity modulators and DNA-damage agents.

Significance:

This study identifies the novel role of YY2/BUB3 axis as a SAC modulator, as well as emphasizing the role of YY2-mediated SAC activity and CIN levels in determining tumor cell fates.

Introduction

Chromosomal instability (CIN) is characterized by an increased rate of chromosomal structural and numerical abnormalities due to aberrant segregation. Chromosomal abnormalities cause significant fitness costs to normal cells, leading to various disorders, including metabolic alterations, proteotoxic stress, cell cycle arrest, and senescence (1,2). Furthermore, it is the most common cause of spontaneous abortion and severe developmental defects (3). Despite these deleterious effects, CIN is a well-recognized cancer hallmark, with approximately 90% of tumors displaying complex karyotypes, including structural and numerical CIN (4). Moreover, studies using mouse models support the notion that CIN significantly promotes tumor initiation (4). However, paradoxically, studies with human clinical tumor samples has revealed that high CIN levels are associated with improved prognosis and better responses to therapies, such as those using DNA-damage agents, thus demonstrating a tumor suppressive function (5-7).

Impaired activity of the spindle assembly checkpoint (SAC), one of the major cell cycle checkpoints, is a major cause of CIN (8). As a checkpoint at the metaphase–anaphase transition point, it ensures that sister chromatids are segregated equally into the two progeny cells and is thus the last checkpoint to guarantee mitotic fidelity through the passage of correct, intact genetic information to the progeny cells (9). SAC activity prevents premature cohesion cleavage, sister chromatid segregation, and mitotic exit by suppressing securin and cyclin B ubiquitination/proteasomal degradation before all sister chromatids are lined up at the equator and attached to spindle microtubules (9). Defect in SAC triggers premature sister chromatid segregation and mitotic exit, leading to increased CIN and faster cell proliferation, thereby

serving as a driving force for tumorigenesis (10,11). Similarly, SAC hyperactivation also

induces CIN; however, it leads to mitotic delay and tumor suppression (12,13). The role of

SAC hyperactivation in tumorigenesis remains poorly understood. Furthermore, the reasons

underlying the different outcomes related to SAC defects and hyperactivation remain to be

explored.

Although mutations in SAC genes are rare, alterations in SAC gene expression are

frequently found in tumor cells, suggesting a crucial role for transcriptional regulation in

aberrant SAC activity in tumor cells (14,15). However, the mechanisms underlying SAC

transcriptional regulation have not yet been fully elucidated. In this study, we utilized a cross-

omics approach to search for potential transcriptional regulators of SAC genes and identified

yin yang 2 (YY2) as a novel *budding uninhibited by benzimidazole 3* (*BUB3*) transcriptional

regulator. Our results showed that YY2/BUB3 axis positively modulates SAC activity,

prolongs mitotic time, and suppresses colorectal cancer (CRC) cells survival. However, despite

of their opposite effects on SAC activity and CRC cells survival, both YY2/BUB3 deficiency

and overexpression lead to increased CIN. We further revealed that different CIN degrees

induced by YY2/BUB3 alterations is crucial for determining CRC cell fate, notably, excessive

CIN level leads to cell death while moderate CIN is beneficial for CRC cells. Finally, our

results showed a strong correlation between moderate CIN and drug resistance, such as that

found in residual tumor cells resistant to DNA damage-inducing agents and/or SAC activation.

In contrast, the induction of excessive CIN in those cells by *YY2* overexpression sensitizes them

to DNA damage-inducing agents, suggesting a novel anti-tumor therapeutic approach by

combining SAC activator and DNA damage-inducing agents.

Materials and Methods

Plasmids and constructs

shRNA expression vectors targeting two different sites of BUB3 or p31/comet were constructed

using algorithm and method as described previously (16). Target sequences were predicted as

follows: 5′- GTC CAG AAG TGA ATG TGA T -3′ (shBUB31-1); 5′- AGA AGA AGA AGT

ATG CCT T -3′ (shBUB3-2); 5′- GGC AGA CCT TCC ATC CGA A -3′ (shp31-1); 5′- GGC

AGA ACT GGA GAG TGT C -3′ (shp31-2).

YY2 (NM_206923.4) overexpression vector (pcYY2), mutant *YY2* overexpression vectors,

and lentivirus vector overexpressing *YY2* (pLenti-YY2) were constructed as described

previously (17). For *BUB3* (NM_004725.4) overexpression vectors, human complementary

DNA (cDNA) obtained by reverse-transcribing the total RNA extracted from HCT116 cells

using the PrimeScript RT Reagent Kit with gDNA Eraser (Takara Bio, Dalian, China) was used

as the template for amplifying the corresponding regions using the PrimeSTAR Max DNA

Polymerase (Takara Bio). The amplicons were then cloned into the *Hind*III and *Eco*RI sites of

pcEF9-Puro vector (18). For doxycycline-inducible *YY2* overexpression vector (pTRIPZ-

YY2), *YY2* coding region was inserted into *Xho*I and *Mlu*I sites of tetracycline-inducible

lentiviral pTRIPZ vector (GE Dharmacon technology, Lafayette, CO).

For reporter vectors bringing different regions of *BUB3* promoter (Refseq No.

NC_000010.11; BUB3-Luc-1 with the -2,688 to +166 region; BUB3-Luc-2 with the -1,765 to

+166 region; BUB3-Luc-3 with the -1,224 to +166 region; BUB3-Luc-4 with the -723 to +166

region; and BUB3-Luc-5 with the -658 to +166 region), the corresponding regions of the *BUB3*

promoter were cloned into *Xho*I and *Hind*III sites of the pGL4.13 (Promega, Madison, WI).

Human genome DNA extracted from HCT116 cells using the TIANamp Genomic DNA Kit

(Tiangen Biotech, Beijing, China) was used as template for amplifying the promoter regions.

BUB3-luciferase vector with mutated predicted YY2 binding site (BUB3-Luc^mut) was

constructed based on the site-specific mutagenesis method using a Site-directed Gene

Mutagenesis Kit (Beyotime Biotechnology, Shanghai, China).

Cell lines and cell culture

HCT116 (catalog number: TCHu 99) and HEK293T (catalog number: GNHu17) cells were

purchased from the Cell Bank of Chinese Academy of Sciences (Shanghai, China) and cultured

in Dulbecco's modified Eagle's medium (Gibco, Life Technologies, Grand Island, NY) with

10% FBS (Biological Industries, Beith Haemek, Israel) and 1% penicillin-streptomycin. Cell

lines were verified using short-tandem repeat profiling method and were tested periodically for

mycoplasma contamination using Mycoplasma Detection Kit-QuickTest (Biotool, Houston,

TX). Transfection was performed using Lipofectamine 2000 (Invitrogen Life Technologies,

Carlsbad, CA) according to the manufacturer's instruction. HCT116^YY2KO cell with deletion of

nucleotides located in +95 to +151 region (56 bp) of *YY2* coding sequence was established

using CRISPR/Cas9 method as described previously.(17) For transfection, cells were seeded

in 6-well plates and transfected with 2 µg of indicated shRNA expression vectors and/or

overexpression vectors. Twenty-four hours after transfection, puromycin selection (final

concentration: 1 µg/ml) was performed for 36 h to eliminate untransfected cells.

Lentiviruses for establishing *YY2*-inducible cell lines and stable cell lines for xenograft

experiments were generated by co-transfecting HEK293T cells with 8 μg pLenti-YY2 or

pTRIPZ-YY2 vectors, 6 μg pCMVΔR, and 2 μg pCMV-VSVG in a 10 cm dish. Growth media

was changed the following day and lentivirus-containing supernatant was harvested and filtered

with a 0.45-μm filter after 48 h. For infection, HCT116 cells were cultured in 6-well plates.

Twenty-four hours later, the medium was changed to 1 ml fresh culture medium and 1 ml

corresponding lentivirus supernatant. Infected cells were then selected using 1 μg/ml

puromycin for 7 days.

For oxaliplatin treatment, cells were seeded in 6-well plates and cultured with medium

containing indicated concentration of oxaliplatin (MedChemExpress, Monmouth Junction, NJ).

Clinical human colon carcinoma specimen

Human colon carcinoma specimens were obtained from colon carcinoma patients undergoing

surgery at Chongqing University Cancer Hospital (Chongqing, China), and stored in Biological

Specimen Bank of Chongqing University Cancer Hospital. Patients did not receive

chemotherapy, radiotherapy, or other adjuvant therapies prior to the surgery. The specimens

were snap-frozen in liquid nitrogen. Prior patient's written informed consents were obtained.

The experiments were approved by the Institutional Research Ethics Committee of Chongqing

University Cancer Hospital, and conducted in accordance with Declaration of Helsinki.

Animal experiments

BALB/*c-nu*/*nu* mice (Strain number D000521, male, body weight: 18–22 g, 6 weeks old) were

purchased from the Chongqing Medical University (Chongqing, China), randomly divided into

4 groups (n = 7), and injected subcutaneously with 3×10^6 indicated stable cell lines.

Oxaliplatin (MedChemExpress) was administered intraperitoneally at a dose of 5 mg/kg twice

a week for 3 weeks. Treatment began on day 10, when the tumor volume reached 100 mm^3.

Tumor size (V) was evaluated by a caliper every 2 days using the following equation: $V = a \times$

$b^2/2$, where a and b are the major and minor axes of the tumor, respectively. The investigator

was blinded to the group allocation and during the assessment. Animal studies were approved

by the Institutional Ethics Committee of Chongqing Medical University, and carried out in the

Chongqing Medical University. All animal experiments conformed to the approved Guidelines

for the Care and Use of Laboratory Animals of Chongqing Medical University. All efforts were

made to minimize suffering.

Metaphase spread

Cells were prepared as described above and arrested at metaphase by 2 μg/ml colchicine

(Aladdin, Shanghai, China) treatment for 3 h before being harvested, re-suspended in

hypotonic solution (0.075 M KCl) and incubated for 15 min at 37 °C. After cell swelling, cells

were fixed using 2 ml of freshly prepared methanol-acetic acid fixative (3:1) and collected by

centrifugation. The pellet was re-suspended in 5 mL of 3:1 methanol-acetic acid for 30 min and

dropped onto pre-cooled slides. Images of mitotic chromosomes were acquired with

fluorescence microscope (Olympus BX53). Chromosome number per cell was quantified.

Karyotype and G-band analysis

Cells were treated with 2 μg/ml colchicine for 1 h and collected processed as described above. After dropped onto pre-cooled slides, the slides were allowed to dry overnight at room temperature, banded with trypsin and stained with 10% Giemsa stain (Biosharp, Shanghai, China). Banded metaphases were photographed using light microscopy with Olympus BX53 microscope and analyzed using SmartType software (Digital Scientific, Cambridge, UK). 10 metaphase spread images were analyzed for each sample.

Cell cycle enrichment and DNA content analysis

For cell cycle analysis, cells were prepared as described above and synchronized in G_0 phase by FBS starvation for 24 h. Cells were washed two times in phosphate-buffered saline (PBS) and either collected immediately (0 h) or further cultured in medium with 10% FBS for indicated time points. Percentages of each cell cycle phases were estimated using ModFit LT (Verity Software House, Topsham, ME).

For DNA content analysis, cells were prepared as described above and subjected to flow cytometry after staining the DNA with propidium iodide (PI) (NeoBiosciences). Percentage of cells with each DNA content (2N, 3N, 4N, 5-8N) were estimated using CytExpert software (Beckman Coulter). For DNA content analysis using sorted cells, cells were transfected or infected with indicated vectors and sorted for living and dying cells as described above before being stained with PI.

Chromosome bridge and micronucleus staining

Cells were prepared as described above and seeded in 3.5-cm confocal dishes (3×10^4 cells per

dish), incubated overnight, and synchronized at prometaphase by nocodazole treatment (final

concentration: 100 ng/ml; Beyotime Biotechnology) for 6 h. Cells were then fixed with 4%

paraformaldehyde for 30 min at room temperature and permeabilized with PBS containing

0.1% Triton X-100 for 5 min. Nuclei were stained with DAPI (Beyotime Biotechnology) for

15 min. For chromosome bridge observation, images were taken using laser scanning confocal

microscopy (Leica Microsystems TCS SP5, Heidelberg, Germany), and quantification was

performed by LAS X software (Leica Microsystems). For micronuclei observation, images

were taken using fluorescence microscope (Olympus IX71). Micronucleus rate was defined as

the ratio of total micronuclei to total cell number.

For chromosome bridge observation in xenografted tumor lesions, fresh xenografted tumor

lesions were fixed using 4 % paraformaldehyde for overnight, embedded in paraffin, and

sectioned at 4 μm thickness using a cryostat. After being dewaxed using xylene and rehydrated,

sections were stained with 0.5% hematoxylin-eosin (Sangon Bio, Shanghai, China). After

samples were dehydrated and mounted in coverslip, images were taken using light microscope

(Olympus BX53).

For micronucleus rate assay using sorted cells, cells were prepared and sorted for living

and dying cells as described above. Cells were then fixed on slides before being fixed,

permeabilized, and stained as described above. Quantification of nucleus size was performed

using ImageJ.

Live-cell imaging

For fluorescence imaging of mitotic division, cells were prepared as described above prior to

seeding in 3.5-cm glass bottom dish (Wuxi NEST Biotechnology, Wuxi, China;

1×10^5 cells/dish) and grown overnight. Nuclei was stained using Hoechst 33342 (Solarbio,

Beijing, China) 30 min prior to imaging. The culture was maintained at 37 °C under 5% CO_2 in

a stage-top incubator (Tokai Hit, Shizuoka, Japan). Images were acquired every 5 min for 6 h

with a 20-ms exposure time using inverted fluorescence microscope Olympus IX83 (Olympus,

Tokyo, Japan). Images were processed using FLUOVIEW (v.2.3). Mitotic time was quantified

as the time from nuclear envelope breakdown (NEBD) until the onset of anaphase.

To study cumulative mitotic exit, nocodazole (final concentration: 100 ng/ml; Beyotime

Biotechnology) was added, and the cells were imaged every 10 min for 10 h using Olympus

IX83 microscope. The number of cells that exited mitosis was quantified and analyzed over

time using FLUOVIEW software.

RNA extraction, quantitative real-time PCR (qRT-PCR), and western blotting

Detailed methods for performing western blotting and quantitative PCR analysis are described

in the Supplementary Materials and Methods. The sequences of the primers and antibodies

used are shown in Supplementary Tables S1 and S2, respectively. Images of uncropped blots

are shown in Supplementary Fig. S12A to S12K.

Statistical analysis

All values of the experimental results were presented as mean ± SD (n = 3; unless further

indicated). Quantification data were analyzed by one-way ANOVA conducted using SPSS

Statistics (v. 20.0.) A value of $P < 0.05$ was considered statistically significant. For each

experiment the number of technical replicates, and the number of independent experiments in

each group is reported in Figure legends.

Results

Identification of transcriptional regulators modulating SAC

To identify a novel modulator of the SAC and subsequently tumor cell CIN, we screened genes

involved in both "mitotic sister chromatid segregation" and "mitotic spindle assembly

checkpoint signaling" from the Gene Ontology (GO) database (QuickGO,

http://www.ebi.ac.uk/QuickGO; GO numbers: 0000070 and 0007094; respectively) and

obtained 195 genes. After excluding non-human-origin and hypothetical genes, we further

screened genes that have been reviewed and annotated using the manual-curation process in

UniProtKB and referenced based on a PubMed Unique Identifier (PMID). The 20 genes

identified in this screening were subjected to transcription factor (TF) enrichment analysis

using enrichment analysis version 3 (ChEA3, https://maayanlab.cloud/chea3/), a web server

application developed to conduct TF enrichment analysis utilizing gene set libraries from

published omics assay data extracted from multiple sources (19) (Fig. 1A). Among the top 17

TFs obtained, eight TFs were known SAC modulators, thus confirming the validity of this

screening method (20-23). whereas nine TFs were potential novel SAC modulators (Fig. 1B).

TFs that regulate the cell cycle usually exhibit oscillating levels during cell cycle

progression (24). Thus, we first determined the timing of each cell cycle phase by

synchronizing HCT116 CRC cells in the G_0/G_1 phase using serum starvation (Supplementary

Fig. S1A). The percentage of cells in the G_0/G_1 phase decreased from 78.12% immediately

after serum starvation release to 14.11% 16 h later, whereas cells in the S phase increased from

10.20% to a peak at 76.89% after 16 h, and cells in the G_2/M phase reached 55.69% at 24 h

after serum starvation release. Cell cycle-dependent expression analysis of the nine novel SAC

modulators revealed that significant expression of ZFP69, ZFP69B, ZNF93, ZNF730, and

ZNF878 could not be detected in any cell cycle state and that of ZNF215, ZNF280C, and

ZNF888 did not show significant differences during cell cycle progression. Meanwhile, YY2

mRNA expression clearly increased during cell cycle progression and peaked during the M

phase (Fig. 1C). A more detailed time-course investigation showed that YY2 mRNA

expression peaked at 20 h (Supplementary Fig. S1B), whereas its protein level started to

increase significantly at 16 h and peaked at 24 h after serum starvation release (Fig. 1D).

Meanwhile, the levels of G_1 and G_2/M phase cyclins, specifically cyclins D and B, peaked at 4

h and 24 h after serum starvation release, respectively, further confirming the relationship

between YY2 expression and the M phase.

SAC delays mitotic exit by stabilizing securin and suppressing APC/C activity. To further

explore the role of YY2 in regulating SAC activity and subsequently mitotic exit, we altered

YY2 expression in HCT116 cells, a near-diploid human CRC cell line with a mitotic time

similar to that of normal cells and considered CIN-negative cells (Supplementary Fig. S1C–

S1D) (25). As shown by time-lapse microscopic images, *YY2* overexpression delayed the

mitosis, whereas *YY2* knockout significantly accelerated it (Fig. 1E–1F, Supplementary Videos

S1–S4). Measurement of mitotic time revealed that *YY2* overexpression prolonged the mitotic

time from 53.6 min to 73.7 min, while *YY2* knockout significantly shortened it from 52.8 min

to 42.8 min (Fig. 1G–1H). Next, we modulated *YY2* expression and synchronized cells in the

M phase using nocodazole. *YY2* overexpression significantly slowed down the degradation of

securin and cyclin B, an event that marks mitotic exit (Supplementary Fig. S2A–S2B), whereas

knocking out *YY2* accelerated their degradation (Supplementary Fig. S2C–S2D). Moreover, to

validate *YY2* overexpression impact on SAC activity, we investigated the ability of *YY2*-

overexpressed cells treated with nocodazole, a metaphase-arresting SAC activator, to sustain

an extended mitotic arrest (26). *YY2* overexpression led to stronger SAC activity, as

demonstrated by fewer cells that exit mitosis under constant SAC activation; thereby

solidifying the role of YY2 in promoting SAC activity (Fig. 1I). These results revealed that

YY2, of which the expression oscillates during the cell cycle and reaches its peak in the M

phase, is a novel SAC modulator that controls mitotic progression.

YY2-mediated SAC regulation is crucial for its tumor suppressive effect

YY2 is a zinc-finger protein and has been reported to have a tumor suppressive function (17,27-

29). A comparative analysis using clinical CRC tissues and corresponding normal adjacent

tissues revealed a significant decrease in YY2 mRNA and protein levels in CRC tissues

(Supplementary Fig. S3A–S3B). Furthermore, *YY2* overexpression significantly suppressed the

viability of HCT116 cells (Supplementary Fig. S3C). As a decrease in cell viability might be

due to a decrease in proliferation, increase in cell death, or both, we next examined the effect

of YY2 on HCT116 cell proliferation and cell death. *YY2* overexpression robustly suppressed

CRC cell proliferation, as indicated by the decrease in EdU-positive cells and colony-formation

potential (Supplementary Fig. S3D–S3E), while increasing the cell death rate (Fig. 2A),

suggesting that YY2 may exert its tumor suppressive function most plausibly by decreasing

cell proliferation and promoting cell death. In contrast, the viability, proliferation, and colony-

formation potential of HCT116^{YY2KO} cells were significantly higher than those of wild-type

cells (Supplementary Fig. S3F–S3H). Interestingly, *YY2* knockout did not significantly affect

the cell death rate (Fig. 2B), most plausibly because of the overall low cell death rate.

To examine whether YY2 exerts its tumor suppressive effect by regulating SAC activity,

we treated *YY2*-overexpressed HCT116 cells with reversine, a SAC activity inhibitor.

Reversine treatment reduced cyclin B and securin accumulation induced by *YY2*

overexpression (Supplementary Fig. S3I) and re-suppressed mitotic time prolonged by *YY2*

overexpression from 72.7 min to 49.3 min (Fig. 2C and 2D, Supplementary Video S5).

Accordingly, reversine treatment restored the proliferation, viability, and colony-formation

potential of *YY2*-overexpressed HCT116 cells (Fig. 2E; Supplementary Fig. S3J–S3K) while

suppressing *YY2* overexpression-induced cell death (Fig. 2F). These results reveal that YY2-

mediated SAC activation and mitotic regulation are crucial for exerting its tumor suppressive

effect.

YY2 regulates SAC by directly activating *BUB3* transcription

To analyze the molecular mechanism underlying YY2-mediated regulation of SAC activity,

we performed a cross-omics analysis to identify its potential transcriptional target. To this end,

RNA-sequencing   data   obtained   in   our   previous   study

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE184138)   (17)   using   *YY2-*

overexpressed HCT116 cells were analyzed for differentially expressed genes (RNA-seq; Fig

3A). The differentially expressed genes (fold-change > 1.01; $P < 0.05$) were then analyzed for

enriched GO terms and Reactome pathways using the Database for Annotation, Visualization,

and Integrated Discovery (DAVID, https//david.ncifcrf.gov/). GO analysis indicated that *YY2*

overexpression enriched genes involved in cell division, mitotic nuclear division, sister

chromatid cohesion, and negative regulation of ubiquitin-protein ligase activity in mitosis (Fig.

3B); meanwhile, Reactome analysis revealed the enrichment of APC/C-Cdc20-mediated

degradation of securin, separation of sister chromatids, mitotic prometaphase, and resolution

of sister chromatid cohesion (Fig. 3C), indicating a possible prominent role of YY2 in

regulating sister chromatid segregation.

To identify potential YY2 transcriptional targets that regulate SAC activity, we

overlapped differentially expressed genes identified by RNA-seq with those identified by

chromatin immunoprecipitation (ChIP)-sequencing using an anti-YY2 antibody reported

previously (27), as well as genes obtained from QuickGO screening (Fig. 1A). Two genes,

BUB3 and *TEX14*, were identified as potential YY2 direct transcriptional targets (Fig. 3D).

BUB3 is a component of the SAC complex that regulates sister chromatid segregation and

mitotic progression (9); meanwhile, TEX14 is required for the formation of intercellular

bridges during meiosis (30). In accordance with previous studies (30,31), analysis of the

TEX14 expression profiles using GTEx v8 expression data across 29 tissues revealed its testis

specificity (Supplementary Fig. S4A). Furthermore, absolute qRT-PCR results showed that the

copy number of *TEX14* was very low in HCT116 cells (Supplementary Fig. S4B). Meanwhile,

BUB3 mRNA increased nearly two-fold upon *YY2* overexpression (Supplementary Fig. S4C),

whereas its protein level was positively correlated with YY2 in *YY2*-knock-down and *YY2*-

overexpressed HCT116 cells (Fig. 3E). Moreover, to avoid the effect of YY2 oscillation during

cell cycle, we arrested the HCT116 cells at G_0/G_1 phase using serum starvation and investigated

the effect of altering YY2 expression on BUB3. The results consistently demonstrated that *YY2*

alteration positively correlates with BUB3 expression level, further confirming the direct

regulation of YY2 on BUB3 expression (Supplementary Fig. S4D−S4E).

Next, we analyzed the *BUB3* promoter using JASPAR (http://jaspar.genereg.net/) (32),

and identified a potential YY2-binding site at −684 to −675 of its promoter region (Fig. 3F).

To confirm the role of this binding site in YY2-mediated regulation of *BUB3* transcriptional

activity, we constructed reporter vectors comprising the −2,688 to +166 region (BUB3-Luc-1),

−1,765 to +166 region (BUB3-Luc-2), −1,224 to +166 region (BUB3-Luc-3), −723 to +166

region (BUB3-Luc-4), and −658 to +166 region (BUB3-Luc-5) of the *BUB3* promoter (Fig.

3F). *YY2* knockout robustly suppressed the transcriptional activities of BUB3-Luc-1, BUB3-

Luc-2, BUB3-Luc-3, and BUB3-Luc4, which contained the predicted YY2-binding site, but

not that of BUB3-Luc-5, which lacked the predicted binding site (Fig. 3G). Concomitantly,

YY2 overexpression promoted the activities of BUB3-Luc-1, BUB3-Luc-2, BUB3-Luc-3, and

BUB3-Luc4, but not that of BUB3-Luc-5 (Fig. 3H). These results suggest that the −723 to

−659 region of the *BUB3* promoter is crucial for YY2-mediated regulation of its transcriptional

activity. Furthermore, the result of a ChIP assay using an anti-YY2 antibody revealed that YY2

could bind to the −721 to −589 region of the *BUB3* promoter (Fig. 3I). Subsequently, mutations

at the predicted YY2-binding site on the *BUB3* promoter (AAAGA*AG*TGG to

AAAGA*CT*TGG; BUB3-Luc^{mut}, Fig. 3J) abolished the YY2-mediated regulation of its

transcriptional activity (Fig. 3K–3L). To further confirm YY2 transcriptional regulator activity

on *BUB3*, we used mutant *YY2* overexpression vectors with a zinc finger mutation constructed

previously based on cancer genomics data set from the cBioportal database (Fig. 3M) (17,33).

Our results showed that these mutant YY2 failed to regulate *BUB3* promoter transcriptional

activity (Fig. 3N). Together, these results showed that YY2 is a transcriptional regulator of

BUB3 that binds to the −684 to −675 site of its promoter region.

To elucidate the role of BUB3 in YY2-induced SAC activity and prolonged mitosis, we

first analyzed the effect of *BUB3* overexpression on mitotic time. Similar to that of *YY2*

overexpression, *BUB3* overexpression in HCT116 cells prolonged the mitotic time from 51.3

min to 80.5 min (Supplementary Fig. S4F–S4H and Supplementary Video S6) and slowed the

degradation rates of cyclin B and securin proteins (Supplementary Fig. S4I–S4K). *BUB3*

overexpression restored the length of HCT116 cells' mitotic time, which was shortened to 43.3

min by *YY2* knockout, to a level similar to that observed in the control (50 min and 52.9 min,

respectively; Fig S5A–S5C and Supplementary Video S7). Next, we constructed two shRNA

expression vectors targeting different sites of BUB3 and selected shBUB3-2 (refers as shBUB3

hereafter), which had a higher suppressive effect, for further experiments (Supplementary Fig.

S5D–S5E). Similar to the trend observed after knocking out *YY2*, knocking down *BUB3*

shortened the mitotic time of HCT116 cells from 52.7 min to 40.5 min and resuppressed the

mitotic time, which was prolonged by *YY2* overexpression, from 71.4 min to 46.8 min

(Supplementary Fig. S5F–S5H and Supplementary Video S8–S9). Accordingly, *BUB3*

knockdown restored HCT116 cell viability and cell death, which was suppressed by *YY2*

overexpression (Supplementary Fig. S5I–S5J). Moreover, *BUB3* overexpression canceled the

increase of CRC cell viability in *YY2* knockout HCT116 cells (Supplementary Fig. S5K). These

results confirmed that YY2 regulates SAC and subsequently, mitotic progression and

tumorigenic potential by enhancing *BUB3* expression.

Both *YY2* knockout and overexpression induce CIN

Defects in SAC activity disrupt the mechanism that guarantees the proper arrangement of

chromosomes at the equator and attachment to the mitotic spindle, making them prone to

missegregation, and therefore, is a well-known causal factor of CIN induction. Thus, we

examined the effect of *YY2* silencing on CIN indicators. We observed significant increases in

the frequency of cells with DNA content > 4N (Fig. 4A) as well as in nuclear size (Fig. 4B) in

HCT116^{YY2KO} cells, indicating the increase of the frequency of polyploid cells. Metaphase

spread analysis results showed a significant increase in the karyotypic heterogeneity of

HCT116^{YY2KO} cells, as nearly 80% of HCT116 cells had 45–47 chromosomes and fewer than

4% had more than 47 chromosomes, whereas only approximately 50% of HCT116^{YY2KO} cells

had 45–47 chromosomes and the percentage of cells with more than 47 chromosomes increased

to nearly 35% (Fig. 4C). Single karyotype analysis further confirmed the increase of cells with

trisomy as well as karyotypic heterogeneity in HCT116^{YY2KO} (Fig. 4D).

Failed sister chromatid segregation leads to the formation of partly unsegregated sister

chromatids or chromosome bridges, of which DNA fragments are subsequently wrapped by a

nuclear membrane-like structure during cytokinesis to form a micronucleus, a small, nucleus-

like structure in the cytoplasm. *YY2* knockout significantly increased the chromosome bridge

frequency (Fig. 4E) and concomitantly increased the micronucleus rate (Fig. 4F), which could

lead to the formation of aneuploid cell. Meanwhile, *BUB3* overexpression could partially

abrogate the frequencies of polyploid cells and chromosome bridges, as well as the

micronucleus rate in HCT116^{YY2KO} cells (Supplementary Fig. S6A–S6C). Together, these

results suggest that defects in the YY2/BUB3 pathway could contribute to CIN induction in

tumor cells, most plausibly due to impaired SAC activity.

Surprisingly, *YY2* overexpression, which induced SAC hyperactivation and mitotic delay,

also increased the number of polyploid cells, nuclear size, karyotype heterogeneity,

chromosome bridge frequency, and micronucleus rates (Fig. 4G–4L). These results suggest

that *YY2* overexpression might also increase CIN.

Given that BUB3 is a component of SAC, we overexpressed *BUB3* and examined its effect

on CIN. *BUB3* overexpression induced cell death (Supplementary Fig. S7A) and increased CIN

phenotypes (Supplementary Fig. S7B–S7E). These results conform with previous studies

showing that SAC activation could also induce CIN and furthermore, cell death (12,34). To

further confirm the causal relation between *YY2* overexpression-induced SAC hyperactivation

and CIN, we suppressed SAC activity in HCT116 cells overexpressing *YY2*. Suppression of

BUB3 or treatment with reversine significantly decreased the number of polyploid cells

(Supplementary Fig. S8A–S8B), chromosome bridge frequency (Supplementary Fig. S8C–

S8D), and micronucleus rate (Supplementary Fig. S8E–S8F) in HCT116 cells overexpressing

YY2, which is likely due to the suppression of SAC activity to near wild-type level in these

cells as shown in Supplementary Fig. S5G-S5H. Together, our results clearly show that both

YY2 knockout and overexpression induced CIN.

CIN levels determine tumor cell fates

The aforementioned results showing that both *YY2* knockout and overexpression induced CIN were intriguing, as *YY2* knockout suppressed SAC activity, shortened the mitotic time, and promoted HCT116 cell proliferation, whereas YY2 overexpression induced SAC hyperactivation, prolonged the mitotic time, enhanced cell death, and suppressed HCT116 cell proliferation. Meanwhile, clinical studies have shown that patients with high CIN have a better prognosis (5-7). Using TCGA CRC data set, we next performed correlation analysis between relapse-free survival rate of CRC patients treated with oxaliplatin, a drug clinically used for treating CRC, and the pretreatment levels of CIN70, a 70-gene signature that has been established as surrogate of CIN (35). The results showed that the moderate CIN levels, *i.e.*, the second and third quartiles, were associated with poor prognosis; meanwhile, high levels of CIN, *i.e.*, the fourth quartile, were associated with better survival, suggesting that excessive CIN might be deleterious for tumors (Fig. 5A).

We next examined the fate of *YY2*-overexpressed cells. To control the timing of *YY2* overexpression, we established a Tet-On *YY2* overexpression system (Supplementary Fig. S9A–S9B). The cell death rate increased in a time-dependent manner after the induction of *YY2* overexpression (Fig. 5B); while treatment with reversine cancelled it (Fig. 5C), suggesting the correlation between *YY2* overexpression-induced SAC hyperactivation and cell death. Given that *YY2* overexpression also led to increased CIN (Fig. 4G–4L), these results indicated the possibility that YY2-induced cell death might be correlated with SAC hyperactivation-induced CIN. However, unlike the cell death rate, the micronucleus rate started to decline after reaching its peak at 48 h post-doxycycline treatment, leading to questions regarding the fate of cells with

high CIN (Fig. 5D). To examine the relationship between the CIN level and cell death, we

stained the cells with Annexin V and propidium iodide (PI), and sorted the "dying cells"

(Annexin V$^+$/PI$^-$), in which nuclear and chromosomal DNA fragmentation had not occurred

and the chromosome could still be observed, as well as "living cells" (Annexin V$^-$/PI$^-$) (Fig.

5E). Whereas there was no significant difference between the micronucleus rate of living and

dying control cells, the micronucleus rate, as well as the number of cells with more than one

micronucleus in dying *YY2*-overexpressed HCT116 cells, was significantly higher than that in

living *YY2*-overexpressed cells (Fig. 5F). Similarly, whereas there was no significant difference

in the percentage of polyploid cells between living and dying control cells (3.40% and 3.78%,

respectively), the percentage of polyploid cells in dying *YY2*-overexpressed HCT116 cells was

significantly higher than that in the corresponding living cells (34.29% and 10.16%,

respectively; Fig. 5G).

Furthermore, while there were only few dying cells in non-treated HCT116^{YY2KO} cells,

we confirmed that the micronucleus rate and percentage of polyploidy cells in those cells were

significantly higher than in living HCT116^{YY2KO} cells (Fig. 5H-5J). Moreover, while knocking

out *YY2* alone did not induce cell death and even benefits tumor cell proliferation, treatment

with reversine, which could further weaken SAC activity as indicated by the shortened mitotic

time (Supplementary Fig. S9C–S9D, Supplementary Video S10), increased CIN as well as cell

death in HCT116^{YY2KO} cells (Supplementary Fig. S9E–S9F). Hence, these results further

confirmed the association between cell death with high CIN. It is noteworthy that while

excessive CIN eventually leads to cell death, intrinsic characteristics of the cells, such as

different gene expression profiles, might also contribute to the capacity of the cells to tolerate

CIN stress, and eventually to their different outcomes such as in cell proliferation (36,37).

Indeed, we observed different expression profiles of genes related with DNA damage response

and negative regulation of apoptosis in *YY2*-overexpressed and *YY2*-knocked out HCT116 cells

(Supplementary Fig. S9G). Nevertheless, while further investigations are needed to

systematically analyze these differences, our results clearly suggest that cells with excessive

CIN induced by either YY2-mediated SAC hyperactivation or *YY2* knockout are more prone

to death.

We further noticed that while the level of CIN in the living *YY2*-overexpressed HCT116

cells was significantly lower than in the dying *YY2*-overexpressed HCT116 cells, it is slightly

higher than in living control cells, as indicated by the slightly increased micronucleus rate and

percentage of polyploid cells (Fig. 5F–5G, respectively). As described above, moderate CIN

level is correlated with poor prognosis (Fig. 5A). These results raised question regarding the

characteristics of these moderate CIN cells. To examine these characteristics, we reversed *YY2*

overexpression by withdrawing doxycycline after sorting the living cells to avoid continuous

SAC hyperactivation (TetOn-*YY2* Trans; Supplementary Fig. S10A–S10B). CIN indicator

analysis performed 2 days after doxycycline withdrawal showed that the slightly increased CIN

level was maintained in living TetOn-*YY2* Trans cells (Fig. 6A–6E and Supplementary Fig.

S10C), confirming that CIN is heritable and can be passed to the progeny cells as reported

previously (38). Meanwhile, while continuous *YY2* overexpression suppressed the viability and

colony-formation potential while increasing cell death rate (TetOn-*YY2* Conti), these

characteristics recovered to the levels similar to those of control in TetOn-*YY2* Trans cells (Fig.

6F–6G and Supplementary Fig. S10D). However, under DNA-damage stress induced by

oxaliplatin, TetOn-*YY2* Trans cells demonstrated robustly higher viability and colony-

formation potential (Fig. 6H–6I). Moreover, the cell death rate of TetOn-*YY2* Trans cells was

significantly lower than HCT116 cells, which were CIN-negative (Fig. 6J), suggesting that CIN

level before oxaliplatin treatment contribute to the observed differences in viability between

cells with continuous and transient *YY2* overexpression. Meanwhile, suppressing CIN in

TetOn-*YY2* Conti cells using reversine decreased cell death and improved cell viability

(Supplementary Fig. S10E–S10G), indicating that reducing excessive CIN level by inhibiting

SAC in SAC-hyperactivated cells can enhance tumor cell drug resistance. Together, these data

indicated the correlation between moderate CIN and tumor cell drug resistance.

Residual cells, which survived the anti-tumor drug treatment and undetectable by standard

morphological examination during the remission phase between the cycles of chemotherapy,

has attracted attention as one of the crucial factors of tumor drug resistance and tumor

recurrence, and thus as a hurdle for anti-tumor therapy (39). The fact that living cells with YY2-

induced SAC hyperactivation possessed moderate CIN suggests the possibility that moderate

CIN is also involved in drug resistance in residual tumor cells that survived treatment with

DNA damage-inducing agents. To test this possibility, we obtained residual tumor cells from

CRC cells infected with TetOn-*YY2* lentivirus by sorting living cells that survived oxaliplatin

treatment, treated them with or without doxycycline to induce *YY2* overexpression, and

examined their viability under oxaliplatin treatment (Fig. 6K). Micronucleus assay revealed

that without induction of *YY2* overexpression, residual cells that survived oxaliplatin treatment

had moderate CIN level (Fig. 6L, middle row); furthermore, these cells demonstrated improved

DNA-damage stress-resistance compared to that in control cells (Fig. 6M). Meanwhile, further

induction of CIN by activating *YY2* overexpression in these residual cells induced excessive

CIN (Fig. 6L, right row), leading to better drug sensitivity (Fig. 6M).

Our results demonstrated that YY2/SAC hyperactivation induces CIN in tumor cells,

leading to the mortality of cells with excessive CIN. Moreover, our results suggest that

moderate CIN, as observed in residual tumor cells surviving SAC hyperactivation or DNA

damage-inducing agent treatment, is crucial for drug resistance in these cells; while increasing

CIN in these cells could improve their drug sensitivity. Together, our findings highlight the

crucial role of CIN levels in determining tumor cell fate and drug resistance.

***YY2*-induced excessive CIN sensitizes tumor cells to DNA damage-inducing agent**

Given that *YY2* overexpression induces excessive CIN, we examined its synergistic effect with

a DNA damage-inducing agent. *YY2* overexpression significantly decreased the IC_{50} of

oxaliplatin; in contrast, *YY2* knockout robustly increased it (Fig. 7A–7B). Furthermore,

combinatorial index (CI) analysis using a previously described method (40,41) showed a

synergistic effect between *YY2* overexpression and oxaliplatin treatment (Fig. 7C). The results

of the EdU incorporation assay (Supplementary Fig. S11A–S11B) and cell death rate (Fig. 7D)

showed that *YY2* overexpression further enhanced the suppressive effect of oxaliplatin on

HCT116 cell proliferation potential while further promoting its effect on inducing cell death.

We then assessed the possibility of combining excessive CIN induced by *YY2*

overexpression with an anti-tumor drug inducing DNA damage for anti-tumor therapeutic

strategy. To this end, we established a *YY2*-overexpressed HCT116 stable cell line using a

lentivirus and performed xenograft experiments (Supplementary Fig. S11C). Whereas the

volume of xenografted tumors formed by control cells increased 16-fold within 4 weeks, *YY2*

overexpression or oxaliplatin treatment alone suppressed this increase to approximately eight-

fold. The suppressive effect was further enhanced by combining *YY2* overexpression and

oxaliplatin treatment, which suppressed the increase in tumor volume to only two-fold (Fig.

7E–7F). The tumor morphology further confirmed this tendency (Fig. 7E). Furthermore,

doubling time analysis showed that combining *YY2* overexpression and oxaliplatin treatment

resulted in more than 9 days of growth delay compared to that with *YY2* overexpression or

oxaliplatin treatment alone, thereby enhancing the therapeutic effect by approximately four-

fold (Fig. 7G). Western blotting and immunohistochemistry (IHC) staining results showed that

BUB3 expression was increased in tumors formed by *YY2*-overexpressed cells, whereas

oxaliplatin did not affect YY2 and BUB3 expression (Fig. 7H and Supplementary Fig. S11D).

Furthermore, IHC staining using γH2AX showed that DNA damage in tumor lesions, which

was increased by *YY2* overexpression or oxaliplatin treatment alone, was further enhanced by

combined oxaliplatin treatment and *YY2* overexpression (Supplementary Fig. S11D).

Finally, we examined CIN levels in the tumor lesions. Compared to that in the controls,

YY2 overexpression or oxaliplatin treatment increased the chromosome bridge frequency,

whereas in oxaliplatin-treated *YY2*-overexpressed HCT116 cells, this was further increased

(Fig. 7I). Collectively, these results demonstrated that *YY2* overexpression sensitizes tumor

cells to oxaliplatin by increasing CIN, suggesting that this combination might be a potential

anti-tumor therapeutic strategy.

Together, our results demonstrated that YY2 is a novel SAC modulator that activates *BUB3*

transcription, thereby influencing SAC activity. This subsequently determines tumor cell fate

and drug resistance by regulating the levels of CIN (Fig. 8).

Discussion

Impaired SAC is the main cause of chromosome segregation errors in mitosis, leading to

numerical and structural chromosome changes in tumor cells (8,10,11). Mouse models have

shown that impaired SAC promotes aneuploidy *in vivo* (42). Moreover, defective SAC activity

is closely related to diseases such as tumors and mosaic variegated aneuploidy, a rare disorder

with a high aneuploidy rate and increased tumor incidence (43). On the other hand, the role of

SAC hyperactivation in tumorigenesis is still poorly understood, with conflicting reports on its

pro- or anti-tumor effects (12,13,34). Nevertheless, mutations in SAC genes are rare in human

tumors, indicating that they are not major contributors to the mitotic errors observed in tumor

cells (14). In this study, we identified YY2, a transcription factor that oscillates during the cell

cycle and peaks during the M phase, as a novel SAC modulator. YY2 directly activates the

transcriptional activity of *BUB3*, a component of the SAC complex. We revealed that YY2

downregulation weakens SAC activity, thereby promoting tumorigenesis by accelerating sister

chromatid segregation, mitotic exit, and tumor cell proliferation. This suggests a mechanism

linking CIN to YY2 downregulation and loss-of-function mutations observed in various tumors

(27,44). Meanwhile, *YY2* overexpression hyperactivates SAC, leading to prolonged mitotic

time, decreased proliferation potential, and increased tumor cell death, thus suppressing

tumorigenesis. YY2 downregulation in various tumors, including colorectal, prostate, ovarian,

and liver cancers, is closely associated with poor survival and prognosis (44-46). While

previous studies have revealed that YY2 can trigger ferroptosis, ultraviolet damage response,

and p53-mediated cell cycle arrest (17,28,47), studies regarding its physiological and

pathological functions are still very limited, and the mechanisms underlying its tumor

suppressive effect have not been completely elucidated. Hence, our findings provide a new

perspective regarding a novel function of YY2 in regulating CIN and subsequently tumor cell

fate through its regulatory effect on SAC activity.

CIN is a hallmark of cancer and can drive tumorigenesis. Paradoxically, high CIN levels

before treatment are associated with improved overall survival outcomes, whereas intermediate

CIN levels are associated with poor clinical outcomes (5-7). A similar phenomenon was

observed in bacteria and viruses, where higher genomic instability benefits the population in

stressful environments through the development of mutations that provide a selective growth

advantage; however, cells with drastic instability never become dominant in a population, as

their instability levels lead to deleterious mutations exceeding the viability threshold (48). In

tumor cells, this concept of a 'just-right' level of instability was also observed based on

mathematical modeling of the evolutionary dynamics of genetically unstable populations

(49,50), suggesting that whereas moderate CIN benefits tumorigenesis as it can allow tumor

cells with varying genetic alterations to have greater chance of acquiring advantageous

characteristics, such as increased proliferation, excessive CIN might induce tumor cell death.

Yet, experimental evidence to support this hypothesis and the molecular mechanisms

underlying tumor cell CIN are still lacking.

Intriguingly, *YY2* overexpression, which hyperactivates the SAC and triggers tumor

suppressive effects, also induces CIN. Indeed, previous studies have revealed that SAC

hyperactivation, for example by overexpression of another SAC component, *MAD2*, or by

knocking down *p31*, a component of SAC silencing, could also induce CIN and cell death

(13,34,51,52). Here we revealed that in the *YY2*-induced SAC-hyperactivated cell population,

dying cells had significantly higher CIN than living cells, thus provides evidence that high CIN

induces cell death and is thus deleterious for tumor cells. Meanwhile, although significantly

lower than that in dying cells, living cells from this population also exhibited a certain degree

of CIN and improved DNA-damage resistance compared to those in control cells, suggesting

that moderate CIN is beneficial for tumorigenesis. Together with the fact that knocking out

YY2 alone is not sufficient for inducing cell death, while further inhibiting SAC activity in *YY2*-

knocked out cells could also lead to excessive CIN and cell death, our findings clearly showed

that different levels of CIN is crucial for determining tumor cell fate. Obviously, while

excessive CIN will eventually lead to cell death, intrinsic characteristics of the cells, such as

their gene expression profiles in responses to DNA damage stress and apoptotic stimuli, also

contribute greatly to their capacity in tolerating CIN stress, and subsequently, the threshold of

CIN-induced cell death. Hence, while further systematical investigation is needed to reveal the

way tumor cells explore their fitness landscape upon CIN induction, our findings provide

experimental evidence that conforms to the 'just-right' model.

DNA damage-inducing agents such as oxaliplatin, taxol, and radiotherapy, which have

been used clinically for treating tumors, can induce mitotic errors and CIN (6,53); while SAC

activator, such as MK-1775 and ZN-c3, are in clinical trials and are considered as potential

anti-tumor drugs (54). Meanwhile, increasing evidences suggest that a population of tumor

cells remain viable after exposure to various anti-tumor drugs. These residual cells display

reduced drug sensitivity, thereby providing a reservoir of cells that might seed the growth of

drug-resistant recurrent tumor. Therefore, targeting residual cells have attracted attention as a

crucial point for anti-tumor therapeutic strategy; however, the mechanisms by which these cells

are generated are still poorly understood. Our findings showed that residual tumor cells, that

survived DNA damage-inducing agents as well as YY2-induced SAC hyperactivation, possess

moderate CIN and are more resistant. Hence, while the underlying mechanism needs to be

further elucidated, our results link up for the first time the level of CIN and residual tumor cells

as described by the 'just-right' model (55,56), thus providing new perspective regarding the

generation of residual tumor cells as well as their drug resistance. Furthermore, although further

study is needed, these findings also point to possible complications, such as predisposing

normal or benign tumor cells to becoming more tumorigenic, owing to the induction of

moderate CIN mediated by DNA damage-based anti-tumor therapies. Moreover, these findings

indicate the possibility of applying the pretreatment CIN level as an indicator to identify

patients who would benefit from DNA damage-based anti-tumor therapies.

Subsequently, our results also demonstrated that *YY2* overexpression-induced SAC

hyperactivation, which triggers excessive CIN, significantly sensitizes CRC cells to oxaliplatin.

Combining *YY2* overexpression and oxaliplatin treatment significantly delayed tumorigenesis

and enhanced the therapeutic effect in a xenograft mouse model, most plausibly by increasing

CIN, as results showed synergism between YY2/SAC hyperactivation-induced excessive CIN

and DNA damage-inducing agents. Therefore, our findings suggest a potential anti-tumor

combinatory therapeutic strategy based on a DNA damage-inducing agent and YY2-induced

excessive CIN, to improve the efficacy of the former, and at the same time, preventing the

generation of residual tumor cells surviving DNA damage-inducing agents with moderate CIN,

which could trigger drug resistance.

Taken together, we identified YY2 as a novel modulator of SAC activity in which

alterations could induce different degrees of CIN and determine tumor cell fates. This provides

experimental evidence explaining the CIN paradox in cancer. Furthermore, our findings have

unraveled a mechanism of drug resistance in residual cells surviving anti-tumor therapies such

as those using DNA damage-inducing agents and SAC activation, and suggest a novel anti-

tumor therapeutic strategy that combines SAC activity regulators and DNA damage-inducing

agents.

Data availability:

All raw data generated in this study are available upon request from the corresponding author.

RNA-Seq data of *YY2*-overexpressed cells used here was obtained from our previous study.

The data was previously deposited in Gene Expression Omnibus and are available at

GSE184138. ChIP-Seq data analyzed in this study were obtained from Gene Expression

Omnibus at GSE76856. TEX14 tissues expression was obtained from Genotype-Tissue

Expression project at https://gtexportal.org/home/gene/TEX14. TCGA CRC data set was

obtained from cBioportal (coadread_tcga_pan_can_atlas_2018).

Acknowledgments

The authors would like to thank Professor Xia Zhang (Institute of Pathology, Southwest

Hospital, Third Military Medical University) for his helpful comments in preparing this

manuscript.

Fundings

This study was financially supported by the National Natural Science Foundation of China

(32070715 (V.K.), 82173029 (S.W.), 32270778 (V.K.), and 82372655 (S.W.)), and the Natural

Science Foundation of Chongqing (CSTB2022NSCQ-MSX0612 (V.K.) and

CSTB2022NSCQ-MSX0611 (SW)).

Author contributions

V.K. and S.W. conceptualized and supervised this study; R.H., I.T.S.M., W.D., and W.L.

performed the experiments; R.H., I.T.S.M., W.D., M.W., S.H., M.M., V.K., and S.W. analyzed

and interpreted the data; R.H., W.D., V.K., and S.W. wrote the paper; H.Z. collected human

clinical samples and performed clinical samples analysis; M.M. predicted and constructed

shRNA expression vectors.

Competing interests

Authors declare that they have no competing interests.

References

1.   Stingele S, Stoehr G, Peplowska K, Cox J, Mann M, Storchova Z. Global analysis of
genome, transcriptome and proteome reveals the response to aneuploidy in human cells.
Mol Syst Biol 2012;8:608.

2.   Williams BR, Prabhu VR, Hunter KE, Glazier CM, Whittaker CA, Housman DE, et al.
Aneuploidy Affects Proliferation and Spontaneous Immortalization in Mammalian
Cells. Science 2008;322:703-9.

3.   Chunduri NK, Storchova Z. The diverse consequences of aneuploidy. Nat Cell Biol
2019;21:54-62.

4.   Vasudevan A, Schukken KM, Sausville EL, Girish V, Adebambo OA, Sheltzer JM.
Aneuploidy as a promoter and suppressor of malignant growth. Nat Rev Cancer
2021;21:89-103.

5.   Birkbak NJ, Eklund AC, Li Q, McClelland SE, Endesfelder D, Tan P, et al. Paradoxical
relationship between chromosomal instability and survival outcome in cancer. Cancer
Res 2011;71:3447-52.

6.   Zaki BI, Suriawinata AA, Eastman AR, Garner KM, Bakhoum SF. Chromosomal
instability portends superior response of rectal adenocarcinoma to chemoradiation
therapy. Cancer 2014;120:1733-42.

7.   Jamal-Hanjani M, A'Hern R, Birkbak NJ, Gorman P, Gronroos E, Ngang S, et al.
Extreme chromosomal instability forecasts improved outcome in ER-negative breast
cancer: a prospective validation cohort study from the TACT trial. Ann Oncol
2015;26:1340-6.

8.   Levine MS, Holland AJ. The impact of mitotic errors on cell proliferation and
tumorigenesis. Genes Dev 2018;32:620-38.

9.   Musacchio A, Salmon ED. The spindle-assembly checkpoint in space and time. Nat
Rev Mol Cell Biol 2007;8:379-93.

10.   Li M, Fang X, Wei Z, York JP, Zhang P. Loss of spindle assembly checkpoint-mediated
inhibition of Cdc20 promotes tumorigenesis in mice. J Cell Biol 2009;185:983-94.

11.   Dai W, Wang Q, Liu T, Swamy M, Fang Y, Xie S, et al. Slippage of mitotic arrest and
enhanced tumor development in mice with BubR1 haploinsufficiency. Cancer Res
2004;64:440-5.

12.   de Cárcer G, Venkateswaran SV, Salgueiro L, El Bakkali A, Somogyi K, Rowald K, et
al. Plk1 overexpression induces chromosomal instability and suppresses tumor
development. Nat Commun 2018;9:3012.

13.   Sotillo R, Hernando E, Díaz-Rodríguez E, Teruya-Feldstein J, Cordón-Cardo C, Lowe
SW, et al. Mad2 Overexpression Promotes Aneuploidy and Tumorigenesis in Mice.
Cancer Cell 2007;11:9-23.

14.   Cosper PF, Copeland SE, Tucker JB, Weaver BA. Chromosome Missegregation as a
Modulator of Radiation Sensitivity. Semin Radiat Oncol 2022;32:54-63.

15.   Schvartzman J-M, Sotillo R, Benezra R. Mitotic chromosomal instability and cancer:
mouse modelling of the human disease. Nat Rev Cancer 2010;10:102-15.

16.   Miyagishi M, Taira K. Strategies for generation of an siRNA expression library directed
against the human genome. Oligonucleotides 2003;13:325-33.

17. Li Y, Li J, Li Z, Wei M, Zhao H, Miyagishi M, et al. Homeostasis Imbalance of YY2
and YY1 Promotes Tumor Growth by Manipulating Ferroptosis. Adv Sci (Weinh)
2022:e2104836.
18. Wu S, Wang H, Li Y, Xie Y, Huang C, Zhao H, et al. Transcription Factor YY1
Promotes Cell Proliferation by Directly Activating the Pentose Phosphate Pathway.
Cancer Res 2018;78:4549-62.
19. Keenan AB, Torre D, Lachmann A, Leong AK, Wojciechowicz ML, Utti V, et al.
ChEA3: transcription factor enrichment analysis by orthogonal omics integration.
Nucleic Acids Res 2019;47:W212-W24.
20. Pierantoni GM, Conte A, Rinaldo C, Tornincasa M, Gerlini R, Federico A, et al.
Deregulation of HMGA1 expression induces chromosome instability through
regulation of spindle assembly checkpoint genes. Oncotarget 2015;6:17342-53.
21. Rotelli MD, Policastro RA, Bolling AM, Killion AW, Weinberg AJ, Dixon MJ, et al.
A Cyclin A-Myb-MuvB-Aurora B network regulates the choice between mitotic cycles
and polyploid endoreplication cycles. PLoS Genet 2019;15:e1008253.
22. Naef V, Monticelli S, Corsinovi D, Mazzetto MT, Cellerino A, Ori M. The age-
regulated zinc finger factor ZNF367 is a new modulator of neuroblast proliferation
during embryonic neurogenesis. Sci Rep 2018;8:11836.
23. Nilsson MB, Sun H, Robichaux J, Pfeifer M, McDermott U, Travers J, et al. A
YAP/FOXM1 axis mediates EMT-associated EGFR inhibitor resistance and increased
expression of spindle assembly checkpoint components. Sci Transl Med
2020;12:eaaz4589.
24. Cuitino MC, Pecot T, Sun D, Kladney R, Okano-Uchida T, Shinde N, et al. Two
Distinct E2F Transcriptional Modules Drive Cell Cycles and Differentiation. Cell Rep
2019;27:3547-60 e5.
25. Choudhary A, Lera RF, Martowicz ML, Oxendine K, Laffin JJ, Weaver BA, et al.
Interphase cytofission maintains genomic integrity of human cells after failed
cytokinesis. Proc Natl Acad Sci USA 2013;110:13026-31.
26. Skowyra A, Allan LA, Saurin AT, Clarke PR. USP9X Limits Mitotic Checkpoint
Complex Turnover to Strengthen the Spindle Assembly Checkpoint and Guard against
Chromosomal Instability. Cell Rep 2018;23:852-65.
27. Wu X-N, Shi T-T, He Y-H, Wang F-F, Sang R, Ding J-C, et al. Methylation of
transcription factor YY2 regulates its transcriptional activity and cell proliferation. Cell
Discov 2017;3:17035.
28. Chen L, Shioda T, Coser KR, Lynch MC, Yang C, Schmidt EV. Genome-wide analysis
of YY2 versus YY1 target genes. Nucleic Acids Res 2010;38:4011-26.
29. Kakizaki F, Sonoshita M, Miyoshi H, Itatani Y, Ito S, Kawada K, et al. Expression of
metastasis suppressor gene AES driven by a Yin Yang (YY) element in a CpG island
promoter and transcription factor YY2. Cancer Sci 2016;107:1622-31.
30. Greenbaum MP, Yan W, Wu MH, Lin YN, Agno JE, Sharma M, et al. TEX14 is
essential for intercellular bridges and fertility in male mice. Proc Natl Acad Sci USA
2006;103:4982-7.

31. Gershoni M, Hauser R, Yogev L, Lehavi O, Azem F, Yavetz H, et al. A familial study
of azoospermic men identifies three novel causative mutations in three new human
azoospermia genes. Genet Med 2017;19:998-1006.
32. Castro-Mondragon JA, Riudavets-Puig R, Rauluseviciute I, Lemma RB, Turchi L,
Blanc-Mathieu R, et al. JASPAR 2022: the 9th release of the open-access database of
transcription factor binding profiles. Nucleic Acids Res 2022;50:D165-D73.
33. Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. The cBio cancer
genomics portal: an open platform for exploring multidimensional cancer genomics
data. Cancer Discov 2012;2:401-4.
34. Rowald K, Mantovan M, Passos J, Buccitelli C, Mardin BR, Korbel JO, et al. Negative
Selection and Chromosome Instability Induced by Mad2 Overexpression Delay Breast
Cancer but Facilitate Oncogene-Independent Outgrowth. Cell Rep 2016;15:2679-91.
35. Carter SL, Eklund AC, Kohane IS, Harris LN, Szallasi Z. A signature of chromosomal
instability inferred from gene expression profiles predicts clinical outcome in multiple
human cancers. Nat Genet 2006;38:1043-8.
36. Garribba L, De Feudis G, Martis V, Galli M, Dumont M, Eliezer Y, et al. Short-term
molecular consequences of chromosome mis-segregation for genome stability. Nat
Commun 2023;14:1353.
37. Thompson SL, Compton DA. Proliferation of aneuploid human cells is limited by a
p53-dependent mechanism. J Cell Biol 2010;188:369-81.
38. Reimann H, Stopper H, Hintzsche H. Long-term fate of etoposide-induced micronuclei
and micronucleated cells in Hela-H2B-GFP cells. Arch Toxicol 2020;94:3553-61.
39. Luskin MR, Murakami MA, Manalis SR, Weinstock DM. Targeting minimal residual
disease: a path to cure? Nat Rev Cancer 2018;18:255-63.
40. Chou TC. Drug combination studies and their synergy quantification using the Chou-
Talalay method. Cancer Res 2010;70:440-6.
41. Chou TC. Theoretical basis, experimental design, and computerized simulation of
synergism and antagonism in drug combination studies. Pharmacol Rev 2006;58:621-
81.
42. Simon JE, Bakker B, Foijer F. CINcere Modelling: What Have Mouse Models for
Chromosome Instability Taught Us? Recent Results Cancer Res 2015;200:39-60.
43. Suijkerbuijk SJ, van Osch MH, Bos FL, Hanks S, Rahman N, Kops GJ. Molecular
causes for BUBR1 dysfunction in the human cancer predisposition syndrome mosaic
variegated aneuploidy. Cancer Res 2010;70:4891-900.
44. Kaufhold S, Aziz N, Bonavida B. The Forgotten YY2 in Reported YY1 Expression
Levels in Human Cancers. Crit Rev Oncog 2017;22:63-73.
45. Li L, Li Y, Timothy Sembiring Meliala I, Kasim V, Wu S. Biological roles of Yin Yang
2: Its implications in physiological and pathological events. J Cell Mol Med
2020;24:12886-99.
46. Uhlen M, Zhang C, Lee S, Sjostedt E, Fagerberg L, Bidkhori G, et al. A pathology atlas
of the human cancer transcriptome. Science 2017;357:eaan2507.

47.   Kasim V, Xie Y-D, Wang H-M, Huang C, Yan X-S, Nian W-Q, et al. Transcription
factor Yin Yang 2 is a novel regulator of the p53/p21 axis. Oncotarget 2017;8:54694-
707.
48.   Cahill DP, Kinzler KW, Vogelstein B, Lengauer C. Genetic instability and darwinian
selection in tumours. Trends Cell Biol 1999;9:M57-60.
49.   Laughney AM, Elizalde S, Genovese G, Bakhoum SF. Dynamics of Tumor
Heterogeneity Derived from Clonal Karyotypic Evolution. Cell Rep 2015;12:809-20.
50.   Komarova NL, Wodarz D. The optimal rate of chromosome loss for the inactivation of
tumor suppressor genes in cancer. Proc Natl Acad Sci USA 2004;101:7017-21.
51.   Choi E, Zhang X, Xing C, Yu H. Mitotic Checkpoint Regulators Control Insulin
Signaling and Metabolic Homeostasis. Cell Rep 2016;166:567-81.
52.   Ma HT, Poon RYC. TRIP13 Regulates Both the Activation and Inactivation of the
Spindle-Assembly Checkpoint. Cell Rep 2016;14:1086-99.
53.   Lee HS, Lee NC, Kouprina N, Kim JH, Kagansky A, Bates S, et al. Effects of
Anticancer Drugs on Chromosome Instability and New Clinical Implications for
Tumor-Suppressing Therapies. Cancer Res 2016;76:902-11.
54.   Ghelli Luserna di Rora A, Cerchione C, Martinelli G, Simonetti G. A WEE1 family
business: regulation of mitosis, cancer progression, and therapeutic target. J Hematol
Oncol 2020;13:126.
55.   Bakhoum SF, Landau DA. Chromosomal Instability as a Driver of Tumor
Heterogeneity and Evolution. Cold Spring Harb Perspect Med 2017;7
56.   Pienta KJ, Hammarlund EU, Brown JS, Amend SR, Axelrod RM. Cancer recurrence
and lethality are enabled by enhanced survival and reversible cell cycle arrest of
polyaneuploid cells. Proc Natl Acad Sci USA 2021;118:e2020838118.
861
862

Figures

Figure 1. Identification of novel transcriptional regulators modulating SAC. **A,** Schematic diagram of screening strategy for potential novel transcriptional regulators modulating SAC. **B,** Top 17 potential TFs modulating SAC obtained from ChEA analysis. **C,** mRNA expression levels of potential novel SAC modulators at each cycle phase, as examined using qRT-PCR. **D,** YY2, cyclin D, and cyclin B protein expression levels in HCT116 cells at indicated time-points after serum starvation release, as determined by western blotting. **E–F,** Mitotic time of *YY2*-overexpressed HCT116 cells (**E**) and HCT116^{YY2KO} cells (**F**), as determined using time-lapse microscopy. Representative images (scale bars: 20 μm) are shown. **G–H,** Scatter plots showing the time-length from nuclear envelope breakdown (NEBD) to anaphase of *YY2*-overexpressed HCT116 cells (**G**) and HCT116^{YY2KO} cells (**H**) (n = 60, pooled from three independent experiments). **I,** Time-lapse analysis of duration of mitosis in *YY2*-overexpressed HCT116 cells arrested with nocodazole (final concentration: 0.2 μM; n = 150, pooled from three independent experiments). Cells transfected with pcCon, wild-type HCT116, or unsynchronized cells were used as controls. β-actin was used for qRT-PCR normalization and as western blotting loading control. Quantification data are shown as mean ± SD. All data were obtained from three independent experiments. *P* values were calculated by one-way analysis of variance (ANOVA). pcCon: pcEF9-Puro; Unsync: unsynchronized; **$P < 0.01$; NS: not significant; ND: not detected.

Figure 2. YY2 suppresses tumorigenesis by activating SAC activity. **A–B,** Cell death rate of *YY2*-overexpressed HCT116 (A) and HCT116[YY2KO] cells (B), as examined using Annexin V/PI staining and flow cytometry. **C–D,** Mitotic time of *YY2*-overexpressed, reversine-treated

HCT116 cells, as determined using time-lapse microscopy. Representative images (**C**; scale

bars: 20 μm) and scatter plot showing the time-length from NEBD to anaphase (**D**; n = 60,

pooled from three independent experiments) are shown. **E,** Proliferation potential of *YY2*-

overexpressed, reversine-treated HCT116 cells, as determined using EdU-incorporation assay.

Representative images (left; scale bars: 200 μm) and ratio of proliferative cells (right; each dot

represents the mean value of three technical replicates) are shown. **F,** Cell death rate of *YY2*-

overexpressed, reversine-treated HCT116 cells, as examined using Annexin V/PI staining.

Cells transfected with pcCon or wild-type HCT116 cells were used as controls. Quantification

data are shown as mean ± SD. All data were obtained from three independent experiments.

P values were calculated by one-way ANOVA. pcCon: pcEF9-Puro; Rev: reversine (final

concentration: 0.2 μM; **$**P < 0.01$; NS: not significant.

Figure 3. YY2 directly regulates *BUB3* transcription. **A,** Volcano plot of log2 fold-change versus adjusted *P* value for gene expression changes (fold-change > 1.01-fold; *P* value < 0.05) in *YY2*-overexpressed HCT116 cells, as analyzed by RNA-seq. **B–C,** GO (**B**) and Reactome

(**C**) enrichments of differentially expressed genes in *YY2*-overxpressed HCT116 cells. **D,** Overlapping genes upregulated by YY2 based on RNA-seq, predicted YY2 target genes identified by ChIP-seq, and QuickGO screening. **E,** BUB3 protein level in *YY2*-overexpressed HCT116 and HCT116^{YY2KO} cells, as determined using western blotting. **F–H,** Schematic diagram (**F**) as well as relative luciferase activities of *BUB3* reporter vectors in HCT116^{YY2KO} cells (**G**) and *YY2*-overexpressed HCT116 cells (**H**). **I,** Binding capacity of YY2 to the predicted region in the *BUB3* promoter, as determined using ChIP assay. Location of the primer pair used for PCR are shown. **J–L,** Schematic diagram (**J**) as well as relative luciferase activities of BUB3-Luc^{mut} in HCT116^{YY2KO} cells (**K**) and *YY2*-overexpressed HCT116 cells (**L**). **M–N,** Schematic diagram of *YY2* mutants overexpressing vectors (**M**) and relative luciferase activity of BUB3-Luc-1 in HCT116 cells overexpressing indicated *YY2* mutants (**N**). Cells transfected with pcCon or wild-type HCT116 cells were used as controls. β-actin was used as western blotting loading control. Quantification data are shown as mean ± SD. All data were obtained from three independent experiments. *P* values were calculated by one-way ANOVA. pcCon: pcEF9-Puro; **P < 0.01; NS: not significant.

Figure 4. Both *YY2* knockout and overexpression induces CIN. **A,** DNA content in HCT116$^{\text{YY2KO}}$ cells, as analyzed using PI staining and flow cytometry. **B,** Average nuclei size of DAPI-stained HCT116$^{\text{YY2KO}}$ cells (each dot represents average nuclei size from one independent experiment, with total 100 cells/group). **C,** Chromosome number per cell in HCT116$^{\text{YY2KO}}$ cells, as analyzed using metaphase spread. Representative images (scale bars: 10 μm) and percentage of cells with indicated chromosome number (total cells counted: 50 cells/group, pooled from three independent experiments) are shown. **D,** Single karyotype analysis of HCT116$^{\text{YY2KO}}$ cells. Representative images and percentage of cells with indicated chromosome number (total cells counted: 10 cells/group, pooled from three independent experiments) are shown. **E,** Chromosome bridge frequency in HCT116$^{\text{YY2KO}}$ cells. Representative images (scale bars: 5 μm) and quantification results (each dot represents chromosome bridge frequency from one independent experiment, with total 100 mitotic-cells/group) are shown. **F,** Micronucleus rate in HCT116$^{\text{YY2KO}}$ cells. Representative images of micronuclei (indicated by arrowheads; scale bars: 20 μm) and micronucleus rate (ratio of micronuclei number to total cell number; each dot represents micronucleus rate/slide with > 100 cells/slides; four technical replicates from three independent experiments) are shown. **G–L,** DNA content analysis (**G**), average nuclei size (**H**), chromosome number per cell (**I**), single karyotype analysis (**J**), chromosome bridge frequency (**K**), and micronucleus rate (**L**) in *YY2*-overexpressed HCT116 cells. Cells transfected with pcCon or wild-type HCT116 cells were used as controls. Quantification data are shown as mean ± SD. All data were obtained from three independent experiments. *P* values were calculated by one-way ANOVA. pcCon: pcEF9-Puro; *P < 0.05; **P < 0.01.

Figure 5. CIN level determines tumor cell fate. **A,** Kaplan-Meier relapse-free survival curves of 108 oxaliplatin-treated CRC patients stratified by CIN70 score quartile. **B–C,** Cell death rate of HCT116 cells after *YY2* induction by doxycycline (**B**) and after reversine treatment (final concentration: 0.2 μM) (**C**), as examined using Annexin V/PI staining. **D,** Micronucleus rate (scale bars: 20 μm; micronucleus rate: ratio of micronuclei number to total cell number; each dot represents micronucleus rate/slide with > 100 cells/slides; four technical replicates from three independent experiments) of HCT116 cells after *YY2* induction by doxycycline. **E–G,** Sorting (**E**), micronucleus rate (**F**), and DNA content (**G**) of living (L) and dying (D) *YY2*-overexpressed HCT116 cells. **H–J,** Sorting (**H**), micronucleus rate (**I**), and DNA content (**J**) of living (L) and dying (D) HCT116^{YY2KO} cells. Cells transfected with pcCon or wild-type HCT116 cells or infected with lentivirus generated using pTRIPZ-control (Con) were used as controls. Quantification data are shown as mean ± SD. All data were obtained from three independent experiments. *P* values were calculated by one-way ANOVA. Scale bars: 20 μm; pcCon: pcEF9-Puro; Dox: doxycycline (final concentration: 2 μg/ml); **$P < 0.01$; NS: not significant.

957 **Figure 6.** Residual tumor cells have moderate CIN and increased drug resistance. **A,** DNA
958 content in HCT116 cells transiently overexpressing *YY2* (TetOn-*YY2* Trans cells) 48 h after

sorting, as analyzed using PI staining and flow cytometry. **B,** Average nuclei size of TetOn-*YY2* Trans cells 48 h after sorting (each dot represents average nuclei size from one independent experiment, with total 100 cells/group). **C,** Chromosome number per cell in TetOn-*YY2* Trans cells 48 h after sorting, as analyzed using metaphase spread. Representative images (scale bars: 10 μm) and percentage of cells with indicated chromosome number (total cells counted: 50 cells/group, pooled from three independent experiments) are shown. **D,** Chromosome bridge frequency in TetOn-*YY2* Trans cells 48 h after sorting. Representative images (scale bars: 5 μm) and quantification results (each dot represents chromosome bridge frequency from one independent experiment, with total 100 mitotic-cells/group) are shown. **E,** Micronucleus rate of TetOn-*YY2* Trans cells 48 h after sorting. Representative images of micronuclei (indicated by arrowheads; scale bars: 20 μm) and micronucleus rate (ratio of micronuclei number to total cell number; each dot represents micronucleus rate/slide with > 100 cells/slides; four technical replicates from three independent experiments) are shown. **F–G,** Cell death rate at indicated time-points (**F**) and cell viability at 4 days after sorting (**G**) of TetOn-*YY2* Trans cells. **H–J,** Viability (**H**), colony-formation potential (**I**; each dot represents the mean value of three technical replicates), and cell death rate (**J**) of TetOn-*YY2* Trans cells at indicated times, 10 days, and 5 days after oxaliplatin treatment, respectively. **K–M,** Schematic diagram (**K**), micronucleus rate (**L**), and viability after oxaliplatin treatment (**M**) of residual HCT116 cells infected with TetOn-*YY2* lentivirus. Cells infected with control lentivirus generated from pTRIPZ-control (Con) were used as controls. TetOn-*YY2* Trans and Conti: cells infected with TetOn-*YY2* lentivirus and treated with doxycycline only before sorting or continuously, respectively. Pre-sort and post-sort: pre-sorting and post-sorting for living cells. Quantification data are shown as mean ± SD. All data were obtained from three independent experiments. *P* values were calculated by one-way ANOVA. Dox: doxycycline (final concentration: 2 μg/ml); **P < 0.01; NS: not significant.

C

Oxaliplatin	pcYY2	Effect	CI
0.16	2	0.26	0.65 **
0.31	2	0.42	0.35 **
0.63	2	0.54	0.26 *
1.25	2	0.62	0.25 *
2.5	2	0.68	0.27 *
5	2	0.78	0.21 *

CI value category

< 0.1: very strong synergism

0.1 – 0.3: strong synergism *

0.3 – 0.7: synergism **

0.7 – 0.85: moderate synergism

0.85 – 0.9: slight synergism

F

Treatment	Tumor volume Day 10	Tumor volume Day 28	Fold change
EV	102.36 ± 19.12	1629.34 ± 244.93	16.07
EV + Oxa	101.65 ± 20.09	884.63 ± 178.01	8.07
YY2	94.28 ± 12.42	704.13 ± 183.29	7.46
YY2 + Oxa	93.76 ± 14.03	197.98 ± 48.19	2.11

G

Treatment	Doubling time (days)	Growth delay (days) Absolute	Growth delay (days) Normalized	Enhancement factor
EV	5			
EV + Oxa	7	2		
YY2	8	3		
YY2 + Oxa	17	12	10 (for YY2)	3.3 (for YY2)
			9 (for Oxa)	4.5 (for Oxa)

Figure 7. Excessive CIN sensitizes tumor cells to DNA-damage agents. **A–B,** Viability of *YY2*-overexpressed HCT116 cells (**A**) and HCT116^{YY2KO} cells (**B**) treated with indicated concentrations of oxaliplatin for 24 h. IC_{50} was calculated using CompuSyn. **C,** Combination index (CI) between oxaliplatin and *YY2* overexpression, as calculated using CompuSyn. **D,** Cell death rate of oxaliplatin-treated *YY2*-overexpressed HCT116 cells, as examined using Annexin V/PI staining and flow cytometry. **E,** Tumor volume and morphological images of xenografted tumors formed by *YY2*-overexpressed HCT116 cells and treated with oxaliplatin at indicated time-points (n = 6/group). **F,** Fold-change of xenografted tumor volumes at day 28 compared to those at the starting point of treatment (day 10). **G,** Enhancement factor of combinatory treatment of oxaliplatin and *YY2* overexpression. **H,** YY2 and BUB3 protein levels in xenografted tumors, as determined using western blotting. **I,** Chromosome bridge frequency in tissue sections of xenografted tumors. Representative images (scale bars: 5 μm) and quantification results (each dot represents chromosome bridge frequency from one slide, with total 100 mitotic-cells/group) are shown. β-actin was used as western blotting loading control. Cells transfected with pcCon, wild-type HCT116 cells, or cells infected with empty virus (EV) were used as controls. Quantification data are shown as mean ± SD. All data were obtained from three independent experiments. *P* values were calculated by one-way ANOVA. pcCon: pcEF9-Puro; **P < 0.01.

1004

1005 **Figure 8.** Schematic diagram showing the role of SAC activity and CIN level in determining
1006 tumor cell fate and drug resistance.

1007