Biomolecular condensates form spatially inhomogeneous network fluids

Furqan Dar[1,8], Samuel R. Cohen[1,2,8], Diana M. Mitrea[3], Aaron H. Phillips[4], Gergely Nagy[5], Wellington C. Leite[5], Christopher B. Stanley[6], Jeong-Mo Choi[7,8,*], Richard W. Kriwacki[4, *], Rohit V. Pappu[1,*]

[1]Department of Biomedical Engineering and Center for Biomolecular Condensates, Washington University in St. Louis, St. Louis, MO 63130, USA

[2]Center of Regenerative Medicine, Washington University in St. Louis, St. Louis, MO 63130, USA

[3]Dewpoint Therapeutics Inc., 451 D Street, Boston, MA 02210, USA

[4]Department of Structural Biology, St. Jude Children's Research Hospital, Memphis, TN 38105, USA

[5]Neutron Scattering Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA

[6]Computational Sciences and Engineering Division, Oak Ridge National Laboratory, Oak Ridge, TN 37830

[7]Department of Chemistry and Chemistry Institute for Functional Materials, Pusan National University, Busan 46241, Republic of Korea

[8]These authors contributed equally: Furqan Dar, Samuel R. Cohen, and Jeong-Mo Choi

[*] e-mail: jmchoi@pusan.ac.kr, richard.kriwacki@stjude.org, pappu@wustl.edu

Abstract

The functions of biomolecular condensates are thought to be influenced by their material properties, and these will be determined by the internal organization of molecules within condensates. However, structural characterizations of condensates are challenging, and rarely reported. Here, we deploy a combination of small angle neutron scattering, fluorescence recovery after photobleaching, and coarse-grained molecular dynamics simulations to provide structural descriptions of model condensates that are formed by macromolecules from nucleolar granular components (GCs). We show that these minimal facsimiles of GCs form condensates that are network fluids featuring spatial inhomogeneities across different length scales that reflect the contributions of distinct protein and peptide domains. The network-like inhomogeneous organization is characterized by a coexistence of liquid- and gas-like macromolecular densities that engenders bimodality of internal molecular dynamics. These insights suggest that condensates formed by multivalent proteins share features with network fluids formed by systems such as patchy or hairy colloids.

Biomolecular condensates are compositionally-distinct membraneless bodies that enable spatiotemporal organization and control over a range of biochemical reactions in cells [1, 2, 3, 4, 5, 6]. Condensates are often thought of as spatially homogeneous liquids that form via liquid-liquid phase separation [7, 8]. However, a more nuanced view is emerging. This is being driven by realization of the importance of multivalence of associative motifs or domains as being crucial for driving condensate formation [9, 10]. Reversible, physical crosslinks formed among multivalent macromolecules give rise to networked molecules that underlie the viscoelasticity of condensates [11, 12, 13, 14, 15, 16, 17, 18, 19, 20]. The phase transitions that give rise to condensates involve coupled associative and segregative phase transitions of associative macromolecules [21, 22].

Proteins that are exemplars of associative macromolecules have distinct molecular grammars, typically featuring oligomerization domains, ligand binding domains, and intrinsically disordered regions (IDRs) with distinctive sequence features [21, 23, 24, 25, 26, 27]. The coupling of associative and segregative phase transitions, referred to as COAST [21], and the driving forces for these transitions derive from the molecular grammars of associative macromolecules [20, 10, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37]. Complex coacervation is the clearest illustration of the coupling between associative and segregative phase transitions [27, 38, 39, 40, 41, 42, 43, 44]. Here, the complexation of polyelectrolytes is driven by a combination of enthalpically favorable associations and entropically favored release of counterions [38, 41, 43, 45]. If the complexes are higher-order clusters of undefined stoichiometry, then we arrive at the Ogston limit where size and hydration details lower the solubility and the complexes undergo a segregative transition to generate coexisting dilute and dense phases [21, 46, 47]. The dilute phase will comprise dissociated polyions, complexed polyions that form higher-order oligomers [48, 49] and pre-percolation clusters [50, 51], all of which must be electroneutral and hence will involve different degrees of ion associations. The dense phase will be a percolated network of polyions, whereby each polyion has multiple partners, and formally this will be limited by the number of uncompensated charges on each polyion in the dense phase. Accordingly, complex coacervation involves electrostatically-driven associations, which can give rise to higher-order complexes, and segregation into dense and dilute phases driven by a combination of counterion release and the lower solubility, due to altered hydration profiles, of higher order complexes. Other instantiations of COAST-like processes include the coupling of percolation and phase separation [9, 10, 52, 53]. Percolation, specifically bond percolation, also known as physical gelation [36, 53, 54, 55], is a continuous associative phase transition whereby motifs or domains form reversible, physical crosslinks to enable the formation of sequence- and architecture-specific networks that span the length scale of the system of interest [33, 52]. As the networks grow, phase separation can be driven by the balance of inter-macromolecule, macromolecule-solvent, and solvent-solvent interactions, controlled largely by the sequence and structural characteristics of spacers, which are regions outside the associative domains and motifs [46, 56].

COAST-like processes give rise to condensates with network-like internal organization [29, 57, 58, 59]. The networks will be defined by the architectures of the constituent molecules and the extents of crosslinking among the molecules [33, 52, 57, 60]. The internal viscosity of condensates and the elasticity of the networks will be governed by the interplay between the timescales for molecular transport within and into / out of condensates and the timescales for making and breaking physical crosslinks. Furthermore, network-like internal organization will engender spatial inhomogeneities of physically crosslinked macromolecules [21, 28, 29, 58].

Viscoelastic materials have time-dependent properties and network structures contribute directly to viscoelastic moduli of condensates [61]. Even if condensates are dominantly viscous fluids

as opposed to elastic solids, there will be timescales where the materials are dominantly elastic [58].
Condensates can age, and if they undergo equilibrium fluid-to-solid transitions, they transform
from dominantly viscous to dominantly elastic viscoelastic materials [58]. Alternatively, some aged
condensates can behave like viscoelastic network glasses [62, 63]. While the network-like
organization within condensates has been inferred from viscoelastic measurements and validated
by the reproduction of measured moduli using computed network structures [58], there is a paucity
of measurements that directly test the hypothesis of network structures within condensates.

The approach we pursue here is rooted in its historical use in the study of simple and
complex fluids, and is based on scattering measurements, specifically small angle neutron
scattering (SANS) [64, 65, 66, 67]. A key advantage of SANS is that one can investigate the presence of
spatial inhomogeneities that range from a few angstroms to hundreds of nanometers [68]. Here, we
investigate the structures of condensates that are mimics of nucleolar sub-phases. The nucleolus is
a spatially organized condensate featuring at least three coexisting sub-phases. The GC, which is
the outermost layer, is scaffolded by nucleophosmin (NPM1) [69, 70]. Condensates formed by
complexation of NPM1 and Arginine-rich (R-rich) peptides and proteins such as rpL5 and SURF6
have been used to test postulates of the molecular handoff model for ribosomal subunit assembly
within the nucleolar GC [71, 72, 73, 74, 75, 76]. Measurements of internal structure within condensates
that are based on SANS were first reported by Mitrea et al., [71]. They studied condensates formed
via heterotypic interactions of cationic, arginine-rich peptides (rpL5) and N130, the N-terminal
130-residues of NPM1. The N130 construct includes the oligomerization domain (OD) and at least
three short regions that are rich in acidic residues [77].

Here, we revisit the SANS data collected by Mitrea et al.,[71] updating these with new
measurements and combining these with simulations to answer the following question: how might
descriptors from theories of simple and complex fluids be adapted for describing condensates? To
answer this question, we adapt approaches that integrate scattering data with computer simulations
[78, 79, 80, 81, 82, 83, 84, 85]. We combine traditional approaches based on pair distribution functions with
graph-theoretic methods to arrive at descriptions of network structures of condensates formed by
N130 and rpL5. The simulations we use are based on bespoke, sequence-specific coarse-grained
models. The latter were developed using a machine-learning approach that is bootstrapped against
atomistic simulations [86].

We find that condensates formed by N130 and rpL5 (N130 + rpL5) have short-range order,
long-range disorder, and a layering at intermediate length scales. These are the hallmarks of fluid-
like organization. Quantitative analysis of structure factors and computed radial distribution
functions combined with graph-theoretic analysis establishes the concept of a network fluid
defined by distinct sub-graphs. One of the sub-graphs is gas-like, and the other is liquid-like, and
this gives rise to two distinct classes of motions within condensates.

RESULTS

N130 and rpL5 form condensates via complexation

Following the work of Mitrea et al., [71] the N130 construct corresponds to residues 1-130
of mouse NPM1 that includes the OD interspersed by short, disordered regions that encompass
acidic residues (Fig. 1a). Previous work showed that N130 forms condensates with R-rich peptides.
Two regions within N130 are enriched in acidic residues. One encompasses a flexible loop
(residues 35-44, termed A1). The other is located at the C-terminus (residues 120-133, termed A2).
These regions were shown to mediate interactions with R-rich peptides and promote condensate

formation (Fig. 1a) [77]. There also is an N-terminal acidic region (residues 1-16, termed A0) that
we discuss below.

Fig. 1: Complexation between acidic regions within N130 and R-motifs of rpL5 is required for condensation.
(a) Schematic representation of N130 including the different acidic regions. The amino acid sequence of N130 is also
shown. The three acidic regions A0, A1, and A2 span residues 4-18, 35-44, and 120-133, respectively. Non-native N-
terminal residues remaining after protease cleavage are underlined. On the right, we show the overall structure of the
hairy colloid generated by superposition of 50 distinct conformations from atomistic simulations. These simulations
use the ABSINTH implicit solvent model [87] and explicit representations of solution ions [88] (which are not shown in
the figure). The pentamerized OD (PDB ID 4N8M), in gray, was modeled as a rigid molecule in the atomistic
simulations. **(b)** The amino acid sequence of rpL5. The panel on the right shows a superposition of 50 different
conformations extracted from ABSINTH-based simulations. **(c)** Confocal microscopy images of phase separation of
100 μM N130 upon titrating the concentration of rpL5 in buffer and 150 mM NaCl. N130 is labeled with
AlexaFluor488. **(d)** Two-component phase boundary for N130 + rpL5, showing the result of concentration titrations.
(e) SANS curve showing the intensity $I(q)$ plotted against q, the scattering vector, for condensates formed by a solution
of N130 (200 μM): rpL5 at 1:3 stoichiometry. Multi-peak analysis, as described by Mitrea et al., [71] leads to the
identification of two major peaks corresponding to 55 Å (right arrow) and 77 Å (left arrow) that are annotated on the
figure. The SANS curve for N130 pentamers, in the absence of rpL5, is shown for comparison. In the interest of clarity,
this curve is shifted upwards vis-à-vis the curve for the N130 + rpL5 system. We computed the scattering curve for
individual N130 pentamers. These computed profiles show qualitative resemblance to the SANS curve shown here
for N130 pentamers (see Supplementary Fig. S1).

We performed atomistic simulations using the ABSINTH implicit solvation model and
forcefield paradigm [87]. In these simulations, the pentamerized OD was modeled as a rigid domain,
the conformations adopted by the IDRs were sampled using Monte Carlo moves, and the
simulations were performed at low salt concentrations of 20 mM. From the simulations, we
obtained an overall structure of the N130 pentamer and for an ensemble of conformations formed
by N130 complexed rpL5 (Fig. 1b) [71]. The rpL5 peptide was taken from the ribosomal protein L5
and its sequence corresponds to the region that has been shown via experiments to interact with
NPM1 [77]. Simulations show that rpL5 adopts ensembles of expanded conformations that maximize
the favorable solvation of Arg and Lys residues [89].

Titrating in rpL5 at a fixed N130 concentration of 100 μM in the absence of crowders gives
a threshold rpL5 concentration for phase separation that is between 250 and 300 μM in 150 mM
NaCl (Fig. 1c). The phase boundary (Fig. 1d) is consistent with previous experimental studies [71,
90]. Next, we performed SANS measurements to probe the molecular organization within
condensates formed by N130 complexed with rpL5 (Fig. 1e). The SANS intensity is a convolution
of the form factor and structure factor. The former quantifies scattering that results from the
average shapes of the scatterers, whereas the latter quantifies how the particles scatter neutrons
due to spatial correlations caused by intra- and intermolecular interactions. Specifically, the
structure factor measures density correlations in reciprocal space, whereas the form factor is the
Fourier transform of the density distribution.[91].

The importance of complexation as a driver of internal organization is made clear by the
lack of peaks in the scattering profile for N130 pentamers when rpL5 is absent from the solution
(also see Supplementary Fig. S1). The inhomogeneities in spatial densities that are evident in the
scattering profile (shown by the arrows of Fig. 1d) are indicative of order on these length scales.
The multi-peak fitting analysis, developed by Mitrea et al.,[71] combined with analysis of derivatives
of the scattering profile show that the most reliable, high signal-to-noise peaks correspond to length
scales of ~55 Å and ~77 Å (Fig. 1e). In the derivative analysis, the signal-to-noise is found to
decrease at low q values, and this makes the unambiguous assignments of peaks beyond the second
one more unreliable (Supplementary Fig. S2).

The molecular diameter of the pentamerized OD (~53 Å) is a useful ruler for calibrating
the different length scales. To further characterize the nature of the ordering and the interactions
that contribute to ordering, we turned to computational approaches.

Systematic coarse-graining and predictions

To investigate the internal structure of fluid-like condensates, we performed coarse-grained
(CG) simulations of the N130 + rpL5 condensates. In the CG model, the pentamerized OD of the
N130 pentamer, referred to hereafter as PD, was modeled as a single, spherical bead defined by
excluded-volume interactions. We used a single-bead-per-residue representation for residues in
the IDRs of N130. Accordingly, in addition to the acidic regions, A1 and A2, we also explicitly
modeled the N-terminal region of N130 (termed A0). The architectures of the coarse-grained N130
molecules are reminiscent of hairy colloids [92], featuring disordered, acidic regions that protrude
from one side of the sphere that mimics the PD. Hairy colloids are known to form network fluids
through anisotropic interactions engendered by the architectures of the constituent molecules [93, 94,
95, 96, 97]. All residues in rpL5 were modeled as single beads.

The systematic coarse-graining procedure was initiated by bootstrapping against
information generated using atomistic simulations based on the ABSINTH implicit solvation
model and forcefield paradigm [87] (Fig. 2a). Having prescribed the resolution for the CG model, we
then used ensembles from atomistic simulations of N130 pentamers with 15 copies of rpL5 to
generate forcefield parameters for the CG model. For this, we use the CAMELOT algorithm [29, 86,
98] that combines a Gaussian Process Bayesian Optimization [99] module, with an appropriate
architecture and CG model. The parameters of the CG model minimize the difference between the
atomistic conformational ensembles and the CG representation. This affords the dual advantages
of computational efficiency afforded by the CG model and the sequence-specific effects learned
via the CAMELOT algorithm. Using the CG representation, we simulate a dense phase with 108
copies of N130 and 1620 copies of the rpL5 peptide.

Fig. 2: Coarse-grained simulations of N130 + rpL5 condensates highlight the importance of an N-terminal acidic region (A0) within N130. (a) Coarse-graining procedure for N130 and rpL5 systems. To generate sequence- and system-specific CG models for N130 and rpL5 peptides, we start with atomistic Monte Carlo simulations of individual molecules that are based on the ABSINTH implicit solvation model and forcefield paradigm [87]. Ions are modeled explicitly in these simulations. We then prescribe a CG model for the system. Here, the residues of the PD are collectively modeled as one large bead depicted in gray. Next, for regions outside the PD, the residues are single beads, and the bead types are organized into three groups: 1: E,K,R,D; 2: V,F,M,I,L,Y,Q,N,W,H; and 3: A,P,G,S,T,C, respectively. To determine the optimal interaction parameters for the CG model, we used the CAMELOT algorithm, [86]. (b) Bead-to-bead contact maps from coarse-grained simulations of dense-phases comprising a 1:3 ratio of N130 and rpL5. The A1 and A2 regions interact favorably with rpL5 peptides. The contact maps reveal interactions involving a region we refer to as A0. (c) Confocal microscopy images of phase separation at a fixed concentration of N130^{+A2} upon titrating with rpL5. The A0 tract in the wild type is replaced with the reversed sequence of the A2 tract. (d) Two-component phase boundary for N130^{+A2} + rpL5, showing the result of concentration titrations. (e) Comparative SANS curves for the WT N130 + rpL5 and +A2 mutant + rpL5. SANS data were collected with 200 μM N130 and 600 μM rpL5. For clarity, the curve for N130^{+A2} + rpL5 is shifted upwards. (f) FRAP curves for the condensates containing N130 and N130^{+A2}. FRAP curves were collected at 100 μM N130 and 400 μM rpL5 with twelve replicates. Error bars represent the standard deviation of the mean.

Results from the coarse-grained simulations of dense phases were used to compute inter-
residue contact maps between the disordered regions of N130 and the rpL5 peptide (Fig. 2b). The
A1 and A2 regions make favorable contacts with the basic residues in rpL5. We also observed that
the A0 region makes contacts with the basic residues in rpL5. The frequency of contacts suggests
that this region forms stronger interactions with rpL5 than A1. The contacts involve acidic residues
within A0. The contact maps derived from the coarse-grained simulations suggest a rank ordering
of interactions between acidic regions and rpL5, with A2 being the most favorable and A1 the least.

Our predictions motivated the generation of a new mutant construct where we replaced A0
with the residues from A2, in reverse order, to increase the linear charge density (see the sequence
of the new A0 region in Fig. 2c). We refer to this construct as $N130^{+A2}$. It has more acidic residues
than the wild type. We hypothesized, based on simulations, that the +A2 mutant should form
condensates with a lower threshold concentration of rpL5 for a given N130 concentration. Indeed,
titrating in rpL5 at a fixed concentration of $N130^{+A2}$ leads to a lower rpL5 threshold concentration
(Fig. 2d) when compared to the threshold concentration that is required for condensation with
wild-type N130 (Fig. 1c). Increasing the strength of the electrostatic interactions in A0 reduces the
threshold concentration for rpL5 from 300 µM to below 250 µM at 100 µM N130. Note that the
designs were chosen to ensure that the stoichiometric ratio required for condensation does not
change.

Next, we investigated the impact of the +A2 mutant using SANS (Fig. 2e). We observed
similar pairs of peaks at intermediate q-values for both N130 + rpL5 and the +A2 mutant + rpL5.
Small shifts in the locations of the peaks are likely a combination of inherent noise and a
contribution from electrostatic repulsions in the disordered N- and C-termini of N130 emanating
from the same face of the PD [77]. The C-terminus of the wild-type protein contains nine negatively
charged residues corresponding to A2, and the +A2 mutant increases the net charge on the
pentamer by 25.

We also measured the impact of the +A2 mutant on the internal dynamics of N130 + rpL5.
For this, we performed measurements of fluorescence recovery after photobleaching (FRAP) on
the condensates (Fig. 2f). The FRAP curve for N130 + rpL5 indicates dynamical exchange with
the bulk solution with the recovery time constant being 53 ± 2 s. Increasing the total charge on
N130 via the +A2 mutant decreases the overall extent of FRAP, resulting in a longer recovery time
of 103 ± 8 s. Similarly, we observe that $N130^{+A2}$ + rpL5 displays slower overall dynamics at
shorter timescales, and the dynamics of the two systems approach one another at longer times. The
average recovery times were obtained by fitting the data, for both constructs, to a single species
model. This does ignore the prospect of there being an immobile fraction. However, since FRAP
data are a convolution of contributions from physical crosslinks and molecular transport, we chose
a parsimonious, single species model to avoid over-fitting and over-interpretations of the data.

Condensates formed by N130 + rpL5 are network fluids.

As observed in the SANS data (Fig. 1e), N130 + rpL5 condensates display correlations at
length scales that are consistent with dimensions of the PD of N130. Therefore, we focus our
analysis on the spatial correlations formed by N130 within condensates. Obtaining the
experimental structure factor by deconvolution of the SANS spectrum would require modeling the
form factor. This becomes intractable given the geometry of the molecules [100]. Instead of solving
an inverse problem, we computed pairwise correlations via the radial distribution function $g(r)$.
This is the real-space analog of the experimentally measurable structure factor [101]. It describes how

spatial densities change as a function of distance from an arbitrary reference particle. Normalized
to an ideal gas, where the distance between particle pairs is completely uncorrelated, $g(r)$ is the
standard descriptor of liquid structure in theory, experiment, and simulations.

There are accounts of condensates being akin to simple liquids [7]. However, in the physical
literature, the term "simple liquids" refers to fluids formed by Lennard-Jones (LJ) particles.
Accordingly, we calibrated our expectations regarding the organization of N130 and rpL5
molecules within condensates, using the radial distribution function, $g(r)$, for the LJ fluid as a
touchstone (Supplementary Fig. S3). In an LJ fluid, structure is defined purely by packing
considerations [102].

In any $g(r)$, the first peak corresponds to the nearest neighbors in the vicinity of the
reference particle of diameter σ, and the additional peaks correspond to higher-order neighboring
shells. As a measure of the density correlations, $g(r)$ quantifies how the average density at a
separation r from the center of any particle varies with respect to the average density of the fluid.
In a fluid, the density correlations are large in the vicinity of the reference particle, and the relative
probability, vis-à-vis the ideal gas, decays as a function of distance until the density becomes
indistinguishable from the average density of the fluid [103]. Structure can be further characterized
by the volume integral over $g(r)$ up to defined positions such as the first minimum. This quantifies
the nearest-neighbor coordination number [101]. For the LJ fluid, the coordination number is ~12-13
due to optimal packing of the spherical particles. In contrast, complex fluids have open, network-
like organization due in part to less efficient packing.

Fig. 3: Radial distribution functions point to network fluid structure of N130 + rpL5 condensates. (a) Radial
distribution function $g_{PD\text{-}PD}(r)$ in black and $n_{PD\text{-}PD}(r)$ in red quantifying the correlations between N130 PDs in the
simulated N130 + rpL5 system. There are local maxima at 53 Å, 95 Å, and 144 Å. On average, approximately four
PDs are found within the first coordination shell. **(b)** Radial distribution functions $g_{+-}(r)$ quantifying the correlations
between positive charges in rpL5 and negative charges in acidic regions of N130. The inset shows a zoom in to troughs
in $g(r)$ in a spatial range between 75 Å and 140 Å that is between 1.5σ and 2.5σ, where σ is the diameter of the N130
PD. Averages were calculated over three replicates for the LJ system and 5 replicates for N130 + rpL5.

From the coarse-grained simulations, we computed $g_{PD\text{-}PD}(r)$, which quantifies spatial
correlations between pairs of PDs of different N130 molecules (Fig 3a). The profile for $g_{PD\text{-}PD}(r)$
is consistent with liquid-like organization, featuring short-range order and long-range disorder
with $g_{PD\text{-}PD}(r)$ approaching unity at large distances. Here, disorder, which refers to the length scale
at which $g_{PD\text{-}PD}(r)$ approaches unity, is evident beyond an inter-PD distance of 3σ, where σ ≈ 53
Å is the diameter of an N130 pentamer.   Integration of $g_{PD\text{-}PD}(r)$ up to the first minimum yields a
coordination number of approximately four. This suggests that the average structure of the fluid,
as interrogated from the vantage point of N130, is not determined purely by packing considerations,

as would be the case for an LJ fluid. Instead, rather like liquid water, which has a coordination number of approximately four, defined by networks of hydrogen bonds, the N130 molecules appear to be incorporated into a network fluid.

The peaks in $g_{PD-PD}(r)$ occur at 53 Å, 95 Å, and 144 Å. The second and third peaks correspond to ordering beyond the molecular length scale. The ratios of the computed peaks to those estimated based on SANS measurements are 0.96 and 1.25 for the first and second, peaks, respectively. Note that the estimates of higher-order peaks from SANS data are less reliable given lower signal-to-noise as quantified using analysis of the derivatives (Supplementary Fig. S2). Further, the parameters of the CG model, especially the parameters for Van der Waals interactions, which are governed by the inter-residue and inter-domain distances, will depend on the screening length and ion-mediated correlations in atomistic simulations. The ABSINTH model includes explicit representations of solution ions, and these simulations were performed at low ionic strength, with the salt concentration set at 20 mM given the explicit representations of ions and large droplet sizes. The inclusion of explicit representations of ions leads to exponential increases in simulation time because of the way electrostatic interactions are handled in the ABSINTH model [87]. In the SANS measurements, the salt concentrations were 150 mM. Therefore, given the parameterization of the CG model using atomistic simulations, the differences in peak positions that correspond to intermediate and longer-range ordering are due to differences in effective Debye lengths between the simulations and SANS measurements. Because the Van der Waals parameters are learned from atomistic simulations, one cannot achieve perfect congruence by simply changing Debye lengths in the coarse-grained simulations. Instead, we need salt concentration dependent parameters within the CG model. This requires a model for how the salt-dependent interactions change at different length scales. The remainder of the discussion focuses on insights we can glean from the CG simulations. In doing so, we presume semi-quantitative rather than fully quantitative congruence with SANS experiments.

Next, we computed the $g(r)$ between basic residues in rpL5 and acidic residues in the three different regions of N130 (Fig. 3b). Note that these $g(r)$ profiles were computed as a linear superposition of pair distributions between all the basic residues in the peptide and all the acidic residues in a specific region. Each of these $g(r)$ profiles has distinct peak positions and heights for the first maximum. The heights of the peaks, realized in a range of $r < 50$ Å, are highest for A2 and lowest for A1. These trends are observed in the corresponding potentials of mean force (Supplementary Fig. S4). The most favorable interactions in the distance range of $r < 50$ Å are realized between A2 and rpL5. The hierarchy of interactions encoded in the different acidic regions of N130 agrees with the contact maps (Fig. 2b).

Interactions between acidic regions and rpL5

Next, we investigated the effects of *in silico* mutations where we neutralized the charges within each of acidic region while keeping the density of the simulated dense phases fixed to that of the wild type N130 + rpL5 condensates. These simulations were designed to assess how mutations that affect electrostatic interactions mediated by one region affect the totality of the network structure. We computed $g(r)$ between pairs of PDs and between the acidic regions on N130 and basic residues of rpL5. Neutralizing the acidic residues on any of the three regions leads to a reduction in the first maximum of $g_{PD-PD}(r)$ (Fig. 4a). The magnitude of the reduction in the first maximum is greatest for the A2 mutant, followed by the A0 and A1 mutants, for which the values of the peak heights are statistically similar within error (Supplementary Fig. S5). This indicates that mutations to A2 affect the overall structure more than mutations to the other regions.

The potentials of mean force (Supplementary Fig. S6) corroborate this inference showing that the
least favorable interactions involve the N130 PD of the A2 mutant. The interactions between basic
residues of rpL5 and acidic residues within A0, A1, and A2 are modular. This is clear from the
$g_{AX-rpL5}(r)$ profiles (Fig. 4b-d), where X = 0, 1, or 2, which we compute from simulations where
one of A0, A1, or A2 is neutralized. The $g_{AX-rpL5}(r)$ profile deviates from that of the WT only for
the region in which the charges are neutralized. Otherwise, the profiles remain roughly equivalent
to those obtained from the wild-type N130 + rpL5 condensates. This suggests that the acidic
regions make modular, and seemingly independent interactions with rpL5 peptides (Fig. 4b-d).

Fig. 4: Interactions between acidic regions and rpL5 are modular and independent of one another. (a) $g_{PD-PD}(r)$
computed after neutralizing the charges in each of the acidic regions indicated in the legend. **(b)** $g_{A0-rpL5}(r)$ quantifies
the pair correlations between acidic residues in A0 and basic residues in rpL5. Results are shown for the WT (black),
when acidic residues are neutralized in A0 (blue), A1 (green), and A2 (magenta). Neutralizing charges within A0
weakens the interactions between A0 and rpL5. However, neutralizing the charges within A1 and A2 does not
significantly affect the interactions between A0 and rpL5. **(c)** $g_{A0-rpL5}(r)$ quantifies the pair correlations between acidic
residues in A1 and basic residues in rpL5. Results are shown for the WT (black), when acidic residues are neutralized
in A0 (blue), A1 (green), and A2 (magenta). Neutralizing charges within A1 weakens the interactions between A1 and
rpL5 (green curve). However, neutralizing the charges within A0 and A2 does not significantly affect the interactions
between A1 and rpL5. **(d)** $g_{A2-rpL5}(r)$ quantifies the pair correlations between acidic residues in A2 and basic residues
in rpL5. Results are shown for the WT (black), when acidic residues are neutralized in A0 (blue), A1 (green), and A2
(magenta). Neutralizing charges within A2 weakens the interactions between A2 and rpL5 (magenta curve). However,
neutralizing the charges within A0 and A1 does not significantly affect the interactions between A2 and rpL5. In all
cases, averages were calculated over five replicates.

Graph-theoretic analyses of network structures of condensates.

The simulation results suggest that N130 + rpL5 condensates are network fluids as opposed
to simple liquids. To put the network fluid concept on a quantitative footing, we turned to graph-
theoretic analysis. These approaches have been used to analyze network fluids such as hydrogen-
bonding networks[104, 105, 106, 107, 108, 109] and network glasses [110].

Fig. 5: Flowchart describing the graph-theoretic analysis of simulations of dense-phases of N130 and rpL5. We start by selecting a set of residues of interest. As an illustrative example, we pick the acidic residues in N130 and the basic residues in rpL5. To determine an edge, we compute the $g(r)$ between sets of beads where the first minimum serves as the distance cutoff for bead adjacency. Given this cutoff, we construct the bead-to-bead adjacency matrices. The system includes the N130 PD. Performing appropriate block-summations of the bead-to-bead adjacency matrix generates the molecular adjacency matrix. The latter is analyzed using standard graph-theoretic analyses. As an example, a random embedding is shown where the size of a node corresponds to the degree centrality of that node.

Following previous work on molecular fluids [111], we construct unweighted graphs in which two molecules are considered adjacent if any of the constituent beads are within the cutoff distance defined by the first minimum in the corresponding $g(r)$. Using this criterion, we constructed appropriate adjacency matrices via block summations (Fig. 5). We then analyzed the network structure formed by the molecular neighbors for the set of beads considered.

To provide a suitable prior of a non-networked fluid where structure is dominated by packing considerations alone, we performed graph-theoretic analyses on systems of LJ particles. For this, we quantified the degree distributions for the vapor, liquid, and solid phases of spherical particles interacting via LJ potentials (Fig. 6a). The degree reflects the number of connections or edges emanating from a node. Here, a node is an individual LJ particle. For an ideal gas, the degree is zero. However, since LJ particles have finite size and there are attractive dispersion interactions, the vapor phase is not ideal. Instead, the degree distribution is skewed to the right. For the LJ liquid, we observe a broad distribution that is roughly symmetrical about a mean degree value of 13. As the density is further increased to obtain a solid, we see that the degree distribution shows a sharp

387 peak at twelve, corresponding to the number of neighbors expected for a 3D hexagonal close-
388 packed lattice [112]. The locally inhomogeneous nature of a liquid allows for interactions with more
389 neighbors than the true ground-state number seen in the solid phase.

Fig. 6: Each acidic region of N130 in the N130 + rpL5 dense phase imparts a different network structure onto the system. (a) Degree distributions, $P(k)$, for pure LJ systems. Because the density of the vapor ($\rho = 0.01$, $T = 1.000$) is low, the most likely degree is zero. For the liquid ($\rho = 0.80$, $T = 1.000$), the degree distribution shifts towards higher values with a mean around twelve, where the widths of the distribution correspond to the inherent variation in the number of bonds that particles can make in the locally spatially inhomogeneous environment of a liquid. For the solid ($\rho = 1.5$, $T = 0.758$), the degree distribution is peaked at twelve. (b) Degree distributions, $P(k)$, for the complementary charge interactions between the different acidic regions of N130 and the rpL5 peptides. Unlike the graphs in panel (a), the distributions display bimodality, which is an indication of a bipartite graph. Consistent with the radial distribution functions in Fig. 3, we see that A2 has the largest degrees, followed by A0 and A1. In all cases, averages were calculated across five replicates.

Next, we constructed graphs using acidic residues in N130 and the basic residues in rpL5
as nodes. In contrast to the LJ systems, the computed degree distributions are bimodal (Fig. 6b),
and this is suggestive of a bipartite network structure. The multimeric nature of the N130 pentamer
allows the acidic regions to interact with multiple rpL5 peptides, as seen in the broad second peaks
in the degree distributions. Consistent with the radial distribution functions for N130 + rpL5, we
also observe a hierarchy of degrees, with A2-rpL5 having the largest degree and A1-rpL5 having
the smallest. However, for the first peaks near $k = 0$, which correspond to the smaller rpL5, we see
that the different acidic regions do not show appreciable differences. Comparison to the LJ system
suggests that the N130 + rpL5 system features both liquid- and gas-like interactions. Here, the
term "gas" refers to the presence of unassociated, freely diffusing rpL5 that coexists with a liquid
comprising associated rpL5 molecules.

Dynamics within network fluids show two distinct regimes.

In spatially inhomogeneous systems, there can be regions that are locally dense or dilute.
This is made clear in the graph-theoretic analysis, which shows two interaction modalities. Similar
results have been reported for fluids formed by patchy particles, especially near the liquid-gas
coexistence region [93, 94, 95]. We reasoned that the coexistence of liquid- and gas-like organization
within the condensates should have dynamical consequences. To test for this, we analyzed the
simulations to compute mean square displacements (MSDs) of the PDs. The MSD is calculated as
a function of lag time. This involves a double average, where the inner average is a cumulative
sum along the time axis, starting from zero, and progressing in increments of $t + \Delta$, where the mean
square displacement is computed over times t and $t + \Delta$ and averaged over the motions of individual
molecules. The outer average is over all molecules. A characteristic timescale corresponds to the
time it takes for the PD to diffuse across a distance corresponding to its diameter. We rescaled the

abscissa by t_D, which is the timescale over which the motion of the PD fits best to a purely diffusive
model with MSD being proportional to t. We find that there is a timescale below t_D where the
motion is super-diffusive with an exponent greater than one, and a timescale above t_D where the
motion of the PD is sub-diffusive with an exponent less than one (Fig. 7a). Based on the observed
length scales, the super-diffusive motion reflects the contributions of short-range steric repulsions
among the PDs and the electrostatic repulsions between acidic residues. Conversely, the sub-
diffusive motions reflect contributions from physical crosslinks between acidic residues and rpL5
peptides. Histograms of the exponents that we compute for the MSDs show a bimodal distribution
(Fig. 7b). The distribution of sub-diffusive exponents is broader and reflects the heterogeneities of
motions impacted by associative interactions between acidic regions and rpL5 peptides. The MSDs
calculated for the PD and for charged residues in each of the acidic regions and the basic residues
in the rpL5 peptides contrast with the MSDs computed in terms of the PD alone (Fig. 7c). The
acidic regions and the peptides show sub-diffusive motions on all timescales, reflecting the fact
that these moieties are influenced mainly by associative intermolecular interactions.

Fig. 7: Motions within dense phases of N130 + rpL5 show bimodality. (a) MSD of the PD plotted against the lag
time shows that N130 has super-diffusive and sub-diffusive regimes. Here, the abscissa is a unitless parameter $t' = t/t_D$
where t_D is the timescale over which the motion of the PD is purely diffusive. The red region in the panel indicates
the timescales that fit best to purely diffusive motion. There is a timescale below t_D where the motion is super-diffusive
and a timescale above t_D where the motion is sub-diffusive, with the regimes and corresponding exponents indicated
in the panel. **(b)** Histograms of the exponents calculated for the mean square displacements of individual PDs. The
bimodal distribution reflects the presence of super-diffusive and sub-diffusive regimes. **(c)** MSDs of the PD and
charged residues in each of the acidic regions and the rpL5 peptides. In contrast to the PD, the acidic regions and the
peptides show only sub-diffusive motions on all timescales. In all cases, averages were calculated over five replicates.

Discussion

Condensates have been referred to as simple liquids [7] or as structureless entities
characterized by non-specific interactions [113]. Systems of LJ particles form simple liquids, and
macromolecules that drive condensation are not LJ particles. Further, liquids are not structureless
entities. And instead of *ad hoc* criteria used to define liquids in the condensate literature [8], structure
in liquids is characterized by short-range order and long-range disorder. The order parameter for
describing liquid-state structure is the radial distribution function. The extent and range of ordering
that can be quantified using radial distribution functions is directly connected to the molecular
architectures, the spatial range, types, and strengths of intermolecular interactions [103]. SANS
measurements are particularly useful for gleaning quantitative insights regarding radial
distribution functions.

Here, we deployed a combination of experimental and computational techniques to
demonstrate that condensates formed by N130 and rpL5 are network fluids. This was established
by observing peaks in SANS curves of condensates that are indicative of molecular order on the

length scale of the N130 pentamers. The SANS data and computations show short- and
intermediate-range ordering versus long-range disorder. From the computed $g_{\text{PD-PD}}(r)$ profiles, we
find that N130 pentamers have four nearest neighbors on average. Complexation between the
acidic regions and the rpL5 peptides also contributes to the overall structure of the condensates.
The acidic regions of N130 function as independent interaction modules. This explains why the
valence of such cohesive motifs is an important driver of condensation and material properties [33,
34, 35, 36, 71].

The sequence-specific CG model allowed us to identify a new acidic region, termed A0, in
the N-terminal end of N130. We find that A0 interacts more strongly with the disordered peptide
rpL5 than was hitherto appreciated. Mutations that increase the charge in A0 help lower the
threshold concentration of rpL5 that is needed to observe condensation driven by heterotypic
interactions with N130.

We find that there are two types of sub-graphs that underlie the structure of the N130 +
rpL5 condensates. One of the sub-graphs corresponds to gas-like organization, and the other
corresponds to that of a liquid. Note that "gas-like" implies that there are regions within
condensates where the concentrations of macromolecules are ultra-dilute, and hence solvent filled.
This is akin to the empty liquid concept [114] reported for patchy colloids. Conversely, what we refer
to as "liquid-like" refers to regions that are dense in macromolecules. The bipartite graphs also
have dynamical fingerprints, which are manifest as the bimodality we observe for the MSDs of the
PDs. Super- and sub-diffusive behaviors that we report here have been observed in MSDs
computed from simulations of oligomer-grafted nanoparticles [115]. They are also consistent with
data from nuclear magnetic resonance experiments where Gibbs et al., found that the PDs of NPM1
form an immobilized scaffold in NPM1+p14ARF mixtures [116]. Taken together, our findings place
the N130 + rpL5 system, and other such systems, in the same category as patchy and / or hairy
colloids [93, 94, 95, 97, 114, 115, 117, 118].

Here, we focused mainly on the effects of heterotypic interactions between N130 and R-
rich rpL5 peptides on the network structure of the N130 + rpL5 condensates. Previous work has
shown that the homotypic interactions within NPM1, uncovered in the presence of crowders, can
also affect both the phase behavior and the mesoscopic structure of condensates formed with
SURF6N [73, 74]. A new method that leveraged the Edmond-Ogston formalism [119], allows for the
intrinsic strengths of homotypic interactions to be uncovered using crowder titrations [47].
Knowledge of the strengths of homotypic interactions, and the relative interplay with heterotypic
interactions, will allow for simulations of binary and higher-order mixtures that mimic nucleolar
GCs. An application of graph-theoretic analysis, guided by SANS measurements, to condensates
that form mainly under the competing interplay of homotypic and heterotypic interactions should
be feasible through the combination of methods deployed in this work. The interplay between
network structures defined by the whole range of homotypic and heterotypic interactions should
illuminate the relationship between rheological properties and network structure [58], for mapping
intra-condensate spatial organizational preferences[28], and for dynamical control over
compositional identities of protein-RNA condensates [27].

Since the nucleolus is a multicomponent and multiphasic condensate [70], we expect that
varying the stoichiometries of different components will affect the overall structural properties of
nucleoli. Future studies that apply the combination of experimental, computational, and analytical
techniques deployed here to more complex systems will enrich our understanding of the
relationship between the spatial organization of condensed systems and the network properties.

METHODS

Cloning

All NPM1-derived N130 constructs were subcloned into a pET28b plasmid vector, in frame with an N-terminal 6x His tag, followed by a TEV protease recognition sequence. rpL5 peptides with variable linkers (rpL5-10L, rpL5-16L, rpL5-20L) were subcloned into a pET28b vector, in frame with an N-terminal 6x His tag and the GB1 protein [120], followed by a TEV protease recognition sequence.

Protein expression and purification

The plasmid constructs were used to transform in *E. coli* strain BL21(DE3) (Millipore Sigma, Burlington, MA, USA), followed by incubation with shaking at 37 °C. When bacterial cultures reached an optical density at 600 nm of ~0.8, the temperature was reduced to 20 °C and protein expression was induced with the addition of IPTG (GoldBio, St. Louis, MO, USA) to a final concentration of 1 mM, and further incubated with shaking overnight. Cells were harvested by centrifugation and lysed by sonication in buffer A (25 mM Tris, 300 mM NaCl, 5 mM β-mercaptoethanol, pH 7.5). The soluble fraction was further separated by centrifugation for 30 min at 30,000 x g and loaded on a Ni-NTA column, pre-equilibrated in buffer A. Bound protein was eluted with a gradient of buffer B (25 mM Tris, 300 mM NaCl, 500 mM Imidazole, 5 mM β-mercaptoethanol, pH 7.5). The fractions containing the protein of interest were identified by SDS-PAGE, pooled and the 6x His affinity tag was removed by proteolytic cleavage, in the presence of TEV protease, while dialyzing against 4 L of 10 mM Tris, 200 mM NaCl, 5 mM β-mercaptoethanol, pH 7.5. To remove the cleaved affinity tag and any un-cleaved material, the protein was applied to an orthogonal Ni-NTA column, and the flow-through loaded on a C4 HPLC column, in 0.1% Trifluoroacetic acid, and eluted with a linear gradient of 0.1 % Trifluoroacetic acid in acetonitrile. The fractions containing the proteins of interest were identified by SDS-PAGE, pooled and lyophilized. Lyophilized N130 and N130^{+A2} proteins were resuspended in 6 M Guanidine hydrochloride, 25 mM Tris, pH 7.5 and reduced by the addition of 10 mM dithiothreitol. The proteins were refolded by dialysis, using 3 exchanges of 10 mM Tris, 150 mM NaCl, 2 mM mM dithiothreitol, pH 7.5, at 4 °C. The protein concentration during refolding was maintained at or below 100 μM N130 monomer. Protein identities were verified by determining their molecular weight using mass spectrometry in the Center for Proteomics and Metabolomics at St. Jude Children's Research Hospital.

Fluorescence labeling of proteins

N130 and N130^{+A2} were labeled with Alexa-488 (ThermoFisher, Waltham, MA, USA) at Cys104 by incubating a molar excess of Alexa-488 maleimide with freshly reduced N130 proteins overnight at 4 °C with oscillation. Excess dye was removed by successive rounds of dialysis against 10 mM Tris, pH 7.5, 150 mM NaCl, and 2 mM DTT. Labeled proteins were then unfolded in the presence of 10 mM Tris, pH 7.5, 2 mM DTT, and 6 M GdmHCl and combined with unlabeled protein to a final concentration of 10% labeled protein and refolded by successive rounds of dialysis against 10 mM Tris, pH 7.5, 150 mM NaCl, and 2 mM DTT.

Fluorescence microscopy

Microscopy plates (Greiner Bio, Kremsmünster, Austria) and slides (Grace BioLabs, Bend, OR, USA) were coated with PlusOne Repel Silane ES (GE Healthcare, Pittsburgh, PA, USA) and Pluronic F-127 (Sigma-Aldrich, St. Louis, MO, USA) and washed with water before the transfer of protein solutions. Fluorescent microscopy experiments were performed using a 3i Marianas spinning disk confocal microscope (Intelligent Imaging Innovations Inc., Denver, CO, USA) with a 100X oil immersion objective (N.A. 1.4). The phase diagram depicted in Fig. 2d was generated by computing the average of the index of dispersion of fluorescent microscopy images of 5 images per well. The threshold for positive phase separation has been set to 10% of the maximum value. Fluorescent recovery after photobleaching (FRAP) experiments were performed by bleaching a circular area with a diameter of 1 μm in the center of droplets (n=12) to approximately 50% of initial fluorescence intensity. The observed fluorescence intensities were then normalized to global photobleaching during data acquisition and fitted as a group to determine recovery times according to [121] :

$$I_t = \frac{I_0 + I_\infty \frac{t}{t_{1/2}}}{1 + \frac{t}{t_{1/2}}}$$

Here, I_0 is the pre-bleach intensity, I_∞ the steady-state, post-bleach intensity, t is the time at which FRAP is measured, and $t_{1/2}$, the time elapsed before half the pre-bleach intensity has been recovered following photobleaching. Uncertainty in the reported half-lives represent the standard error of the fit to the data. FRAP experiments were performed one hour after the mixing of components.

Peptides

The rpL5 peptide was synthesized in the Macromolecular Synthesis lab at the Hartwell Center, St. Jude Children's Research Hospital. The lyophilized powder was directly reconstituted in buffer, and the pH was adjusted to 7.5 using 1 M Tris base.

SANS

N130 and N130^{+A2} were buffer exchanged into 10 mM Tris, 150 mM NaCl, 2 mM DTT, in D_2O (measured pH, 7.5). Lyophilized rpL5 peptides were resuspended in dialysis buffer. Monodisperse samples of protein only and phase separated samples with rpL5 were prepared in the dialysis buffer. SANS experiments were performed on the extended Q-range small-angle neutron scattering (EQ-SANS, BL-6) beam line at the Spallation Neutron Source (SNS) located at Oak Ridge National Laboratory (ORNL). In 30 Hz operation mode, a 4 m sample-to-detector distance with 2.5-6.1 and 9.8-13.4 Å wavelength bands was used [122] covering a combined scattering vector range of $0.006 < q < 0.44$ Å^{-1}. $q = 4\pi \sin(\theta)/\lambda$, where 2θ is the scattering angle and λ is the neutron wavelength. Samples were loaded into 1 or 2 mm pathlength circular-shaped quartz cuvettes (Hellma USA, Plainville, NY, USA) and sealed. SANS measurements were performed at 25 °C using the EQ-SANS rotating tumbler sample environment to counteract condensate settling. Data reduction followed standard procedures using MantidPlot[123] and

582 drtsans.[124] The measured scattering intensity was corrected for the detector sensitivity and
583 scattering contribution from the solvent and empty cells, and then placed on absolute scale using
584 a calibrated standard [125].

Atomistic Monte Carlo simulations using the ABSINTH model

For the first step of systematic coarse-graining, we performed atomistic Monte Carlo
simulations to obtain a robust description of the conformational ensembles of N130. For this, we
employed the ABSINTH implicit solvation model and forcefield paradigm [87]. In this model, all
polypeptide atoms and solution are modeled explicitly, and the degrees of freedom are the
backbone and sidechain dihedral angles as well as the translational motions of the solution ions,
which are spheres. All simulations were performed using version 2.0 of the CAMPARI modeling
package (http://campari.sourceforge.net) and the abs_opls_3.2.prm parameter set. The initial
structure of N130 was modelled as a pentamer, where the structure of the oligomerization domains
is based on the coordinates deposited in the protein data bank (PDB ID: 4N8M). The structures of
each disordered N-terminal tail (residues 1-18, GSHMEDSMDMDMSPL) and disordered A2 tract
(residues 124-133, EDAESEDEDE) were built using CAMPARI. The degrees of freedom internal
to the oligomerization domains were held fixed during the ABSINTH simulations, reflecting the
fact that the domains are well folded and tightly bound to each other. The system is placed in a
soft-wall spherical potential with radius 70 Å. We included sodium and chloride ions to mimic the
salt concentration of ~20 mM, in addition to neutralizing ions. The simulation temperature was set
to 300 K.

For efficient sampling of the conformational ensemble, we first performed simulations
based on the so-called excluded volume or EV limit. In this limit, all terms other than the steric
repulsions and any dihedral angle terms in the potential functions are switched off. Note that the
ABSINTH model uses fixed bond lengths and bond angles. These initializing simulations were
performed for 10^8 Monte Carlo (MC) steps. We sampled 100 different structures from the EV limit
simulations and used each of them as initial structures simulations based on the full potential. Each
simulation consists of 10^8 MC steps, and the structural information was stored every 5,000 steps.
Hence, we collected 20,000 snapshots per trajectory, from 100 independent trajectories of
atomistic simulations. Next, we performed ABSINTH-based MC simulations for 2.1×10^7 MC
steps with sampling frequency of $(5,000 \text{ steps})^{-1}$, where the first 10^6 MC steps were discarded as
equilibration.

Computations of scattering profiles using CAMPARI

We used the computed the scattering form factor $P(q)$ from snapshots generated using the
ABSINTH-based simulations. We excised the conformations of N130 pentamers, and computed
Kratky profiles using the `scattercalc` functionality within the CAMPARI package
(http://campari.sourceforge.net). In these calculations, the scattering cross-sections of all atoms
are set to be unity. The results we obtain for $P(q)$ are plotted as $\log(P(q))$ versus $\log(q)$ in
Supplementary Fig. S1.

Systematic coarse-graining

Our approach to developing coarse-grained models involves three steps: First, we choose
the resolution for the coarse-grained (CG) model. Second, we choose the form for the potential
functions that describe interactions among pairs of coarse-grained sites. And third, we use a
Gaussian process Bayesian optimization (GPBO) module to parameterize the model to ensure that

the CG model recaptures conformational statistics of the atomistic simulations. In our choice of
the model, each residue is modelled as a bead, except for the oligomerization domain of N130,
which by itself forms a large bead with excluded volume. The mass of each bead is determined by
the total mass belonging to the specific bead. For example, the oligomerization domain bead has
mass of 47544.8 amu. The position of each bead is set equal to the position of the center of mass
in its atomistic representation.

The potential function used for the simulations is decomposed into five different terms:

632
$$W = W_{\text{LJ}} + W_{el} + W_b + W_\theta + W_\phi \tag{1}$$

Each term contains several interaction parameters, which were parameterized using the
CAMELOT algorithm [86]. This uses a GPBO module to minimize an objective function, which is
defined as the difference between the site-to-site distance distributions generated by atomistic
Monte Carlo simulations based on the ABSINTH model and by coarse-grained molecular
dynamics (MD) simulations. Within the CAMELOT algorithm, we change the parameters of the
potential function for the CG model, perform CG MD simulations to obtain conformational
statistics, specifically inter-site distance distributions, quantify the objective function, and iterate
until a stationary state is reached for the objective function. In this work, we used conformational
statistics derived from the atomistic simulations of structural ensembles of N130 and R-rich
peptides as the reference against which the CG model was parameterized. The optimized
parameters are given in Supplementary Tables 1-8. The bead types and corresponding amino acid
residues are given in Supplementary Fig. S7.

To reduce the computational cost of scanning the parameter space, we grouped several
amino acid types into one bead type, following the previous work. For the rpL5 peptide, we
grouped different amino acids into three different bead types: *charged* (K, R, E, D), *large* (V, F,
M, I, L, Y, Q, N, W, H), and *small* (A, P, G, S, T, C). Each group has its own ϵ_i value to be
determined. For each residue in either the large or small group, we used σ_i as twice of the radius
of gyration of the specific residue in the atomistic simulations (and consequently σ_i is not identical
for all residues in the same group; it is residue-dependent). For charged residues, its σ_i was left as
a free parameter to be determined by the optimization module. Hence, we have four unknown
parameters, three ϵ_i values for the charged, large, and small bead types plus one σ_i value for the
charged bead type.

Coarse-grained model for MD simulations

The coarse-grained (CG) model is summarized in Fig. 1, and the potential function used
for the simulations can be decomposed into five different terms given in Eq. 1. Here,

658
$$W_{LJ,ij} = 4\epsilon_{ij} \left[\left(\frac{\sigma_{ij}}{r_{ij}} \right)^{12} - \left(\frac{\sigma_{ij}}{r_{ij}} \right)^{6} \right], \qquad r_{ij} < r_c$$

659 is the standard LJ potential with cutoff $r_c = 2.5\sigma$. While we decomposed the two-body interaction
660 parameters into one-body parameters: $\sigma_{ij} = (\sigma_i + \sigma_j)/2$ and $\epsilon_{ij} = \sqrt{\epsilon_i \epsilon_j}$. All energies have units
661 of kcal / mol.

662   The electrostatic interactions were modeled using a Debye-Hückel potential given by

663
$$W_{el,ij} = C \frac{q_i q_j}{\epsilon r_{ij}} \exp(-\kappa\, r_{ij}) , \qquad r_{ij} < r_c$$

664 implemented as the `lj/cut/coul/debye` pair-style in LAMMPS [126] with $\epsilon = 80.0$, $\kappa = 0.1$, $\sigma_1 =$
665 24.1 and $\sigma_2 = 15.0$ Å. with constant C, charges q, dielectric constant ϵ, and inverse Debye length
666 κ. In this work, we used $\epsilon = 80.0$ and $\kappa = 0.1$ Å^{-1}. The charges were assigned manually; beads
667 for R and K have +1, beads for D and E have -1, and other beads (including the PD bead) have 0.

668   The bond and angle terms are modeled as harmonic potentials,

669
$$W_{b,i} = K_i(b_i - b_{0i})^2$$
670 is a quadratic bonded potential implemented as the harmonic bond-style in LAMMPS [126].
671
$$W_{\theta,i} = K_i(\theta_i - \theta_{0i})^2$$

is a quadratic angular term implemented as the harmonic angle-style in LAMMPS [126]. The bond
parameters $K_{b,i}$ and b_{0i} were obtained by fitting the normal distribution to the distribution of the
distance between two adjacent residues. The angle parameters $K_{a,i}$ and θ_{0i} were also obtained by
fitting the normal distribution to the distribution of the angle between three adjacent residues. For
the oligomerization domain bead, we assigned arbitrarily high values for the energy parameters:
$K_{b,i} = 60000$ kcal / mol-Å^2 and $K_{a,i} = 6000000$ kcal / mol-radian^2, keeping the bond extremely
rigid. The equilibrium length and angle were determined by the PDB structure.

Except for the oligomerization domain bead, the dihedral term is given by a Fourier series
potential,

681
$$W_{\phi,i} = \sum_{n=1}^{3} K_{ni}(1 - \cos(n\,\phi_{0i} - \phi_{ni}))$$

is a Fourier series dihedral term implemented as part of the class2 dihedral style in LAMMPS.
with arbitrarily high value of $K_{d,i} = 6000000$ kcal / mol-radian^2 and experimentally determined
ϕ_{0i}. The parameters corresponding to the potentials, derived from CAMELOT, for the different
systems considered are in Supplementary Tables S1-S8. Lastly, $W_{\phi,i,\text{quad}} = K_i(\phi_i - \phi_{0i})^2$ is used
to constrain the five arms of N130 with a very high K-values.

Simulations of LJ systems using MD

Using the `lj/melt` example in the LAMMPS documentation as a starting point, we
performed standard NVT simulations, with 10^4 Lennard-Jones particles, to understand the network
structure of different LJ phases. To access different pure phases, we used the NIST parameters for
a pure LJ gas, a pure LJ fluid, and a pure LJ solid. In LJ units, the densities and temperatures for
the different systems are in Table 1.

693
694 **Table 1:** Density and temperature for different LJ-phases in LJ-units.

Phase	ρ^*	T^*
Vapor	0.01	1.0
Fluid	0.8	1.0
Solid	1.5	0.758

695
With the standard $dt = 0.005\tau$, where τ is the dimensionless Lennard-Jones time unit, an initial
$t_{\text{therm}} = 1 \times 10^5$ timesteps were run to let the systems settle, after which for $t = 1 \times 10^7$
timesteps were run for data production. Trajectory snapshots were output every 10000 timesteps.
The last 500 frames were used to calculate the radial distribution functions (RDFs) and degree

distributions. Three independent replicates were run for each condition, and the standard error of
the mean between replicates is used as the measure of uncertainty.

Simulations of the N130 wild type and rpL5 peptides

Given the initial configuration generated using CAMELOT [86], we used the replication
command in LAMMPS to generate 108 copies of N130 pentamers 1620 copies of the rpL5 peptide.
Following the replication, `fix deform` was used with `fix nve/limit` to reduce the box-sizes for
the simulations to 250 nm. The final configuration served as initial conditions for NPT simulations
to prepare systems at the correct intrinsic density. NPT simulations were run for $t = 1 \times 10^7$
timesteps. The final configurations from these NPT simulations served as the starting configuration
for NVT simulations which were run for $t = 1 \times 10^9$ timesteps. Trajectory snapshots were output
every 50000 steps, and we only considered the last 1000 frames for the different analyses. Five
independent NVT replicates were run for each condition, and the standard error of the mean
between replicates is used as the measure of uncertainty.

Simulations of the N130 mutants

714   Using the final box-size from the NPT simulations with N130 and the native rpL5 peptide,
715 we ran NVT simulations for $t = 2 \times 10^9$ timesteps. Like the peptide simulations, trajectory
716 snapshots were output every 50000 timesteps in the last 1×10^9 steps, and only the last 1000
717 frames were considered for analyses. Five independent NVT replicates were run for each condition,
718 and the standard error of the mean between replicates is used as the measure of uncertainty.

Calculation of the radial distribution functions $g(r)$

We used VMD[127] to calculate $g(r)$ for the different sets of beads considered in this work.
The specific function used was `measure gofr`. To calculate the $g(r)$ for the N130 + rpL5 system,
we used a bin size, dr, of 0.5 Å. Note that the $g(r)$ profiles for the N130 mutants were computed
from the pair distributions between all the acidic residues in the select acidic region, and all basic
residues in the peptide. For the LJ-fluid simulations, we used a bin size of 0.025 in units of dr/σ.
As with our analysis of the N130 + rpL5 system, we averaged over all replicates.

Graph-building methods.

First, we identify the regions on different molecules whose contributions to the network
structure of the fluid we wish to investigate. We focused our analysis on the network formed by
acidic residues in a particular acidic region in N130 and the basic residues in rpL5. This defines
two sets of residues. Given the two sets of residues / beads, we calculate the inter-set $g(r)$ as
explained in the methods. For each $g(r)$ we compute the location of the first minimum. The
locations of these minima serve as the cut-off radii for defining the presence of an edge between
beads.

Given two sets of beads, a particular trajectory snapshot, and the computed cut-off radius,
we generate a bead-adjacency matrix. A basic bead is considered adjacent to an acidic residue if
the distance between the selected acidic and basic beads is within the cutoff radius. This calculation
is performed for all pairs defined by the two sets of beads that are chosen for the analysis. This
generates a bead adjacency matrix where an edge is drawn between beads in the chose set if the
inter-bead distances are within the computed cutoff radius. We can either consider the total bead
adjacency matrix where every bead in the system is included and where all the bead-types not in

the initially chosen set are non-adjacent by construction, or we can generate a bead adjacency
matrix where only the considered beads are included. We choose the latter.

In more detail, suppose that set-1 has m-bead *types* and set-2 has n-bead *types*. Then, given
α N130 molecules, set-1 has $\alpha \cdot m$ beads in total. Similarly, given β rpL5 molecules, set-2 has
$(\beta \cdot n)$ beads in total. Therefore, the bead-adjacency matrix will be an $(\alpha \cdot m + \beta \cdot n) \times (\alpha \cdot$
$m + \beta \cdot n)$ matrix. As an example, suppose we have 1 N130 molecule, and 1 rpl5 molecule. This
would then give us an $(m + 15n) \times (m + 15n)$ matrix. Furthermore, suppose that the beads are
ordered such that the first m-rows correspond to the N130 beads, and so the next n-rows are the
rpL5 beads (since adjacency matrices are symmetric the first m-columns would be for N130 and
the next n-columns would be for rpL5). To go from this bead adjacency matrix to a molecular
adjacency matrix we would look at blocks of this bead adjacency matrix. Let the bead-adjacency
matrix be $\hat{B}$. Using an indexing, B[0:m, 0:m] corresponds to the sub-graph of N130 adjacent beads.
In our case, no beads will be adjacent since the graph is intentionally constructed between the
acidic residues in N130 and the basic residues in rpL5. Moving on, B[0:m, m:m+n] (or B[m:m+n,
0:m] due to symmetry) corresponds to the sub-graph between the N130 beads and the beads of the
first rpL5 molecule. Similarly, if we had more than 1 rpL5 molecule, B[0:m, m+i*n:m+(i+1)*n]
gives us the subgraph between the N130 beads and the beads of the i-th rpL5 molecule. To
generate the molecular adjacency matrix, $\hat{A}$ we check if any of the sub-graphs from the bead-
adjacency matrix are non-empty (or that there are edges in that graph) or that there is a 1 in the
B[0:m, m+i*n:m+(i+1)*n] block. Therefore, A[0,0]=0 in our case by construction, or that
molecule-0 and molecule-0 are *not* adjacent.

Returning to the more general case, we have the bead adjacency matrix that corresponds to
a matrix of size $(\alpha \cdot m + \beta \cdot n) \times (\alpha \cdot m + \beta \cdot n)$ where the first $\alpha \cdot m$ rows (or columns)
correspond to the beads in N130 molecules, and the last $\beta \cdot n$ rows (or columns) correspond to the
beads in rpL5 molecules. We check all the blocks of the matrix which correspond to the beads
between the different molecules in the system. These correspond to the subgraphs between the
beads of different molecules. For any sub-graph or block that is non-empty, we consider the two
corresponding molecules to be adjacent. This gives the molecular adjacency matrix which should
have the shape $(\alpha + \beta) \times (\alpha + \beta)$. This is the final graph that is analyzed. Graph properties are
calculated per-snapshot and then averaged over the total set of frames considered.

The molecular adjacency graph is constructed by considering the adjacency between any
of the beads from the initially selected set. Since we only care about the acidic and basic beads, by
construction this graph avoids self-loops. Furthermore, since we only care about specific blocks
of the bead adjacency matrix being non-empty, we only get an unweighted graph. If, however, we
wanted to get the weighted graph, we would simply take the sum of the number of edges in a
particular sub-graph or take the sum of that block from the bead adjacency matrix.

On an implementation level, we can skip most of the block reductions by simply asking
first: given the two sets of beads and the cutoff radius, which pairs of beads are adjacent. Then,
from the pairs of adjacent beads, we ask which molecules the beads in the pairs come from. More
specifically we identify the molecule-ID of each bead. From this set of molecule-ID pairs, we
simply find the unique pairs. Given these unique pairs of adjacent molecule-IDs, we construct the
molecular adjacency matrix since the molecule-IDs directly correspond to the indices in the
adjacency matrix. We set those elements of the $(\alpha + \beta) \times (\alpha + \beta)$ matrix to 1, and we get our
molecular adjacency matrix.

Degree distributions

From a given trajectory snapshot, we generate molecular graphs, $G(V, E)$, where individual molecules are represented as nodes, V. To calculate the edges between nodes in a generalizable way, we use the first minima from the RDFs of the given sets of beads. We use signal.find_peaks from the SciPy package[128] to find these minima. We generated RDFs for the acidic / negatively charged beads in N130 from a particular acidic region, and the basic residues in the rpL5 peptide. Given the RDFs, we then use the first minimum as the cut-off radius for the definition of an edge. An edge, E, is therefore drawn between two nodes if any of the beads from the considered sets are within the distance corresponding to the first minimum of the $g(r)$ of interest. We use MDAnalysis[129] to analyze the trajectories to find adjacent molecules. Given these adjacency matrices, we then calculate the degree of each node by calculating the total number of edges for each node, or by calculating the sum of each row (or column) of the adjacency matrix. The python package NumPy[130] is used to generate these degrees. From the degrees, we calculate the degree distribution. The degree distributions from the last 10^3 frames are averaged over, and the average over the five simulation replicates are reported here.

Mean square displacements.

The mean square displacement (MSD) was calculated by averaging the displacements in particle positions over all windows of length m and over all particles N:

$$MSD(m) = \frac{1}{N}\sum_{i=1}^{N} \frac{1}{N-m}\sum_{k=0}^{N-m-1}(\vec{r}_i(k+m) - \vec{r}_i(k))^2 \qquad (2)$$

for $m = 1, \dots, N-1$. For the acidic regions and rpL5, the MSD was calculated with respect to all acidic and basic residues, respectively. For all MSDs, we fit the first 30 ps and everything past 200 ps using single exponents to describe the two different regimes. For the MSD with respect to the PD, the region from approximately 80 to 120 ps was found to fit best to a simple diffusion model with $R^2 = 0.9984$. The crossover t_D between the super- and sub-diffusive regimes is the median value within the diffusive region of the MSD with respect to the PD. In all cases, the MSD was averaged over five replicates. To calculate the histograms of the exponents of the individual molecules, we modified Eq. (2) to calculate the MSD with respect to each PD rather than averaging over all PDs. We then fit each MSD as before.

Plotting

To generate the plots, we use Matplotlib[131] along with Seaborn[132]. Adobe Illustrator® is used to generate the final figures shown.

REFERENCES

1.   Banani SF, Lee HO, Hyman AA, Rosen MK. Biomolecular condensates: organizers of cellular biochemistry. *Nat Rev Mol Cell Bio* **18**, 285-298 (2017).

2.   Brangwynne CP, *et al.* Germline P granules are liquid droplets that localize by controlled dissolution/condensation. *Science* **324**, 1729-1732 (2009).

825 3.   Brangwynne CP, Mitchison TJ, Hyman AA. Active liquid-like behavior of nucleoli
826   determines their size and shape in Xenopus laevis oocytes. *PNAS* **108**, 4334-4339 (2011).
827

828 4.   Feric M*, et al.* Coexisting Liquid Phases Underlie Nucleolar Subcompartments. *Cell* **165**,
829   1686-1697 (2016).
830

831 5.   Shin Y, Berry J, Pannucci N, Haataja MP, Toettcher JE, Brangwynne CP. Spatiotemporal
832   Control of Intracellular Phase Transitions Using Light-Activated optoDroplets. *Cell* **168**,
833   159-171.e114 (2017).
834

835 6.   Shin Y, Brangwynne CP. Liquid phase condensation in cell physiology and disease.
836   *Science* **357**, eaaf4382 (2017).
837

838 7.   Taylor N, Elbaum-Garfinkle S, Vaidya N, Zhang H, Stone HA, Brangwynne CP.
839   Biophysical characterization of organelle-based RNA/protein liquid phases using
840   microfluidics. *Soft Matter* **12**, 9142-9150 (2016).
841

842 8.   Hyman AA, Weber CA, Jülicher F. Liquid-liquid phase separation in biology. *Annu Rev*
843   *Cell Dev Biol* **30**, 39-58 (2014).
844

845 9.   Li P*, et al.* Phase transitions in the assembly of multivalent signalling proteins. *Nature* **483**,
846   336-340 (2012).
847

848 10.   Mittag T, Pappu RV. A conceptual framework for understanding phase separation and
849   addressing open questions and challenges. *Molecular Cell* **82**, 2201-2214 (2022).
850

851 11.   Alshareedah I, Moosa MM, Pham M, Potoyan DA, Banerjee PR. Programmable
852   Viscoelasticity in Protein-RNA Condensates with Disordered Sticker-Spacer Polypeptides.
853   *Nat Commun* **12**, 6620 (2021).
854

855 12.   Keizer VIP*, et al.* Live-cell micromanipulation of a genomic locus reveals interphase
856   chromatin mechanics. *Science* **377**, 489-495 (2022).
857

858 13.   Feric M, Sarfallah A, Dar F, Temiakov D, Pappu RV, Misteli T. Mesoscale structure–
859   function relationships in mitochondrial transcriptional condensates. *PNAS* **119**,
860   e2207303119 (2022).
861

862 14.   Böddeker TJ*, et al.* Non-specific adhesive forces between filaments and membraneless
863   organelles. *Nature Physics* **18**, 571-578 (2022).
864

865 15.   Zhou H-X. Viscoelasticity of biomolecular condensates conforms to the Jeffreys model.
866   *The Journal of Chemical Physics* **154**, 041103 (2021).
867

868 16.   Ghosh A, Kota D, Zhou H-X. Shear relaxation governs fusion dynamics of biomolecular
869   condensates. *Nat Commun* **12**, 5995 (2021).
870

17. Bergeron-Sandoval LP*, et al.* Endocytic proteins with prion-like domains form viscoelastic condensates that enable membrane remodeling. *PNAS* **118**, e2113789118 (2021).

18. Alshareedah I, Kaur T, Banerjee PR. Methods for characterizing the material properties of biomolecular condensates. *Methods in Enzymology* **646**, 143-183 (2021).

19. Roberts S*, et al.* Injectable tissue integrating networks from recombinant polypeptides with tunable order. *Nature Materials* **17**, 1154-1163 (2018).

20. Berry J, Brangwynne CP, Haataja M. Physical principles of intracellular organization via active and passive phase transitions. *Rep Prog Phys* **81**, 046601 (2018).

21. Pappu RV, Cohen SR, Dar F, Farag M, Kar M. Phase Transitions of Associative Biomacromolecules. *Chem Rev* **123**, 8945-8987 (2023).

22. Zhang Z, Chen Q, Colby RH. Dynamics of associative polymers. *Soft Matter* **14**, 2961-2977 (2018).

23. Choi JM, Holehouse AS, Pappu RV. Physical Principles Underlying the Complex Biology of Intracellular Phase Transitions. *Annu Rev Biophys* **49**, 107-133 (2020).

24. Choi JM, Dar F, Pappu RV. LASSI: A lattice model for simulating phase transitions of multivalent proteins. *PLoS Comput Biol* **15**, e1007028 (2019).

25. Powers SK*, et al.* Nucleo-cytoplasmic Partitioning of ARF Proteins Controls Auxin Responses in Arabidopsis thaliana. *Molecular Cell* **76**, 177-190.e175 (2019).

26. Sanders DW*, et al.* Competing Protein-RNA Interaction Networks Control Multiphase Intracellular Organization. *Cell* **181**, 306-324 e328 (2020).

27. Lin AZ*, et al.* Dynamical control enables the formation of demixed biomolecular condensates. *Nat Commun* **14**, 7678 (2023).

28. Farag M, Borcherds WM, Bremer A, Mittag T, Pappu RV. Phase separation of protein mixtures is driven by the interplay of homotypic and heterotypic interactions. *Nat Commun* **14**, 5527 (2023).

29. Farag M, Cohen SR, Borcherds WM, Bremer A, Mittag T, Pappu RV. Condensates formed by prion-like low-complexity domains have small-world network structures and interfaces defined by expanded conformations. *Nat Commun* **13**, 7722 (2022).

30. Bremer A*, et al.* Deciphering how naturally occurring sequence features impact the phase behaviours of disordered prion-like domains. *Nature Chemistry* **14**, 196-207 (2022).

915   31.   Zeng X, Holehouse AS, Chilkoti A, Mittag T, Pappu RV. Connecting Coil-to-Globule
916   Transitions to Full Phase Diagrams for Intrinsically Disordered Proteins. *Biophysical*
917   *Journal* **119**, 402-418 (2020).
918
919   32.   Yang P*, et al.* G3BP1 Is a Tunable Switch that Triggers Phase Separation to Assemble
920   Stress Granules. *Cell* **181**, 325-345 e328 (2020).
921
922   33.   Schmit JD, Bouchard JJ, Martin EW, Mittag T. Protein Network Structure Enables
923   Switching between Liquid and Gel States. *Journal of the American Chemical Society* **142**,
924   874-883 (2020).
925
926   34.   Martin EW*, et al.* Valence and patterning of aromatic residues determine the phase
927   behavior of prion-like domains. *Science* **367**, 694-699 (2020).
928
929   35.   Wang J*, et al.* A Molecular Grammar Governing the Driving Forces for Phase Separation
930   of Prion-like RNA Binding Proteins. *Cell* **174**, 688-699.e616 (2018).
931
932   36.   Choi JM, Hyman AA, Pappu RV. Generalized models for bond percolation transitions of
933   associative polymers. *Physical Review E* **102**, 042403 (2020).
934
935   37.   Guillen-Boixet J*, et al.* RNA-Induced Conformational Switching and Clustering of G3BP
936   Drive Stress Granule Assembly by Condensation. *Cell* **181**, 346-361 e317 (2020).
937
938   38.   Pak Chi W*, et al.* Sequence Determinants of Intracellular Phase Separation by Complex
939   Coacervation of a Disordered Protein. *Molecular Cell* **63**, 72-85 (2016).
940
941   39.   Priftis D, Megley K, Laugel N, Tirrell M. Complex coacervation of poly(ethylene-
942   imine)/polypeptide aqueous solutions: Thermodynamic and rheological characterization.
943   *Journal of Colloid and Interface Science* **398**, 39-50 (2013).
944
945   40.   Neitzel AE, Fang YN, Yu B, Rumyantsev AM, de Pablo JJ, Tirrell MV. Polyelectrolyte
946   Complex Coacervation across a Broad Range of Charge Densities. *Macromolecules* **54**,
947   6878-6890 (2021).
948
949   41.   Sing CE, Perry SL. Recent progress in the science of complex coacervation. *Soft Matter*
950   **16**, 2885-2914 (2020).
951
952   42.   Adhikari S, Leaf MA, Muthukumar M. Polyelectrolyte complex coacervation by
953   electrostatic dipolar interactions. *The Journal of Chemical Physics* **149**, (2018).
954
955   43.   Galvanetto N*, et al.* Extreme dynamics in a biomolecular condensate. *Nature* **619**, 876-883
956   (2023).
957
958   44.   Margossian KO, Brown MU, Emrick T, Muthukumar M. Coacervation in polyzwitterion-
959   polyelectrolyte systems and their potential applications for gastrointestinal drug delivery
960   platforms. *Nat Commun* **13**, 2250 (2022).

45. Brangwynne CP, Tompa P, Pappu RV. Polymer physics of intracellular phase transitions. *Nature Physics* **11**, 899-904 (2015).

46. Ogston AG. On the interaction of solute molecules with porous networks. *The Journal of Physical Chemistry* **74**, 668-669 (1970).

47. Chauhan G, Bremer A, Dar F, Mittag T, Pappu RV. Crowder titrations enable the quantification of driving forces for macromolecular phase separation. *Biophysical Journal* **123**, 1-17.

48. Chowdhury A*, et al.* Driving forces of the complex formation between highly charged disordered proteins. *PNAS* **120**, e2304036120 (2023).

49. Veis A. A review of the early development of the thermodynamics of the complex coacervation phase separation. *Adv Colloid Interfac* **167**, 2-11 (2011).

50. Kar M*, et al.* Phase separating RNA binding proteins form heterogeneous distributions of clusters in subsaturated solutions. *PNAS* **119**, e2202222119 (2022).

51. Lan C*, et al.* Quantitative real-time in-cell imaging reveals heterogeneous clusters of proteins prior to condensation. *Nat Commun* **14**, 4831 (2023).

52. Harmon TS, Holehouse AS, Rosen MK, Pappu RV. Intrinsically disordered linkers determine the interplay between phase separation and gelation in multivalent proteins. *eLife* **6**, 30294 (2017).

53. Semenov AN, Rubinstein M. Thermoreversible gelation in solutions of associative polymers. 1. Statics. *Macromolecules* **31**, 1373-1385 (1998).

54. Choi J-M, Dar F, Pappu RV. LASSI: A lattice model for simulating phase transitions of multivalent proteins. *PLOS Computational Biology* **15**, e1007028 (2019).

55. Flory PJ. Molecular Size Distribution in Three Dimensional Polymers. I. Gelation1. *Journal of the American Chemical Society* **63**, 3083-3090 (1941).

56. Flory PJ. Thermodynamics of High Polymer Solutions. *The Journal of Chemical Physics* **10**, 51-61 (1942).

57. Shillcock JC, Lagisquet C, Alexandre J, Vuillon L, Ipsen JH. Model biomolecular condensates have heterogeneous structure quantitatively dependent on the interaction profile of their constituent macromolecules. *Soft Matter* **18**, 6674-6693 (2022).

58. Alshareedah I*, et al.* Sequence-encoded grammars determine material properties and physical aging of protein condensates. *bioRxiv*, 2023.2004.2006.535902 (2023).

59. Vilgis TA. 8 – Polymer Networks. *Comprehensive Polymer Science and Supplements* **8**, 227-279 (1989).

60. Bhandari K, Cotten MA, Kim J, Rosen MK, Schmit JD. Structure–Function Properties in Disordered Condensates. *The Journal of Physical Chemistry B* **125**, 467-476 (2021).

61. Wróbel JK, Cortez R, Fauci L. Modeling viscoelastic networks in Stokes flow. *Physics of Fluids* **26**, (2014).

62. Jawerth L*, et al.* Protein condensates as aging Maxwell fluids. *Science* **370**, 1317-1323 (2020).

63. Jabbari-Farouji S*, et al.* High-bandwidth viscoelastic properties of aging colloidal glasses and gels. *Physical Review E* **78**, 061402 (2008).

64. Elstone NS*, et al.* Understanding the Liquid Structure in Mixtures of Ionic Liquids with Semiperfluoroalkyl or Alkyl Chains. *The Journal of Physical Chemistry B* **127**, 7394-7407 (2023).

65. Hirosawa K*, et al.* SANS study on the solvated structure and molecular interactions of a thermo-responsive polymer in a room temperature ionic liquid. *Physical Chemistry Chemical Physics* **18**, 17881-17889 (2016).

66. Tanaka H, Tong H, Shi R, Russo J. Revealing key structural features hidden in liquids and glasses. *Nature Reviews Physics* **1**, 333-348 (2019).

67. Malenkov GG. Structure and dynamics of liquid water. *Journal of Structural Chemistry* **47**, S1-S31 (2006).

68. Mühlbauer S*, et al.* Magnetic small-angle neutron scattering. *Reviews of Modern Physics* **91**, 015004 (2019).

69. Pederson T. The Nucleolus. *Cold Spring Harbor Perspectives in Biology* **3**, (2011).

70. Lafontaine DLJ, Riback JA, Bascetin R, Brangwynne CP. The nucleolus as a multiphase liquid condensate. *Nat Rev Mol Cell Bio* **22**, 165-182 (2021).

71. Mitrea DM*, et al.* Nucleophosmin integrates within the nucleolus via multi-modal interactions with proteins displaying R-rich linear motifs and rRNA. *Elife* **5**, (2016).

72. Mitrea DM, Kriwacki RW. Phase separation in biology; functional organization of a higher order. *Cell Communication and Signaling* **14**, 1 (2016).

73. Ferrolino MC, Mitrea DM, Michael JR, Kriwacki RW. Compositional adaptability in NPM1-SURF6 scaffolding networks enabled by dynamic switching of phase separation mechanisms. *Nat Commun* **9**, 5064 (2018).

74. Mitrea DM*, et al.* Self-interaction of NPM1 modulates multiple mechanisms of liquid-liquid phase separation. *Nat Commun* **9**, 842 (2018).

75. Riback JA*, et al.* Composition-dependent thermodynamics of intracellular phase separation. *Nature* **581**, 209-214 (2020).

76. Riback JA*, et al.* Viscoelasticity and advective flow of RNA underlies nucleolar form and function. *Molecular Cell* **83**, 3095-3107.e3099 (2023).

77. Mitrea DM*, et al.* Structural polymorphism in the N-terminal oligomerization domain of NPM1. *PNAS* **111**, 4466-4471 (2014).

78. Clark GNI, Hura GL, Teixeira J, Soper AK, Head-Gordon T. Small-angle scattering and the structure of ambient liquid water. *PNAS* **107**, 14003-14007 (2010).

79. Borodin O*, et al.* Liquid Structure with Nano-Heterogeneity Promotes Cationic Transport in Concentrated Electrolytes. *ACS Nano* **11**, 10462-10471 (2017).

80. Maier EE, Krause R, Deggelmann M, Hagenbuechle M, Weber R, Fraden S. Liquid like order of charged rodlike particle solutions. *Macromolecules* **25**, 1125-1133 (1992).

81. Londono JD, Annis BK, Turner JZ, Soper AK. The intermolecular hydrogen–hydrogen structure of chain–molecule liquids from neutron diffraction. *The Journal of Chemical Physics* **101**, 7868-7872 (1994).

82. Cousin F, Gummel J, Ung D, Boué F. Polyelectrolyte−Protein Complexes: Structure and Conformation of Each Specie Revealed by SANS. *Langmuir* **21**, 9675-9688 (2005).

83. Fujii K, Kumai T, Takamuku T, Umebayashi Y, Ishiguro S-i. Liquid Structure and Preferential Solvation of Metal Ions in Solvent Mixtures of N,N-Dimethylformamide and N-Methylformamide. *The Journal of Physical Chemistry A* **110**, 1798-1804 (2006).

84. Troitzsch RZ, Martyna GJ, McLain SE, Soper AK, Crain J. Structure of Aqueous Proline via Parallel Tempering Molecular Dynamics and Neutron Diffraction. *The Journal of Physical Chemistry B* **111**, 8210-8222 (2007).

85. Schöttl S*, et al.* Combined molecular dynamics (MD) and small angle scattering (SAS) analysis of organization on a nanometer-scale in ternary solvent solutions containing a hydrotrope. *Journal of Colloid and Interface Science* **540**, 623-633 (2019).

86. Ruff KM, Harmon TS, Pappu RV. CAMELOT: A machine learning approach for coarse-grained simulations of aggregation of block-copolymeric protein sequences. *J Chem Phys* **143**, 243123 (2015).

87. Vitalis A, Pappu RV. ABSINTH: a new continuum solvation model for simulations of polypeptides in aqueous solutions. *Jounral of Computational Chemistry* **30**, 673-699 (2009).

88. Mao AH, Pappu RV. Crystal lattice properties fully determine short-range interaction parameters for alkali and halide ions. *The Journal of Chemical Physics* **137**, (2012).

89. Fossat MJ, Zeng X, Pappu RV. Uncovering Differences in Hydration Free Energies and Structures for Model Compound Mimics of Charged Side Chains of Amino Acids. *J Phys Chem B* **125**, 4148-4161 (2021).

90. Banerjee PR, Milin AN, Moosa MM, Onuchic PL, Deniz AA. Reentrant Phase Transition Drives Dynamic Substructure Formation in Ribonucleoprotein Droplets. *Angewandte Chemie International Edition* **56**, 11354-11359 (2017).

91. Guinier A, Fournet Gr. *Small-Angle Scattering of X-rays*. Wiley (1955).

92. Chen YM. Shaped Hairy Polymer Nanoobjects. *Macromolecules* **45**, 2619-2631 (2012).

93. de las Heras D, Tavares JM, Telo da Gama MM. Phase diagrams of binary mixtures of patchy colloids with distinct numbers of patches: the network fluid regime. *Soft Matter* **7**, 5615-5626 (2011).

94. Dias CS, Araújo NAM, Telo da Gama MM. Dynamics of network fluids. *Adv Colloid Interfac* **247**, 258-263 (2017).

95. Dias CS, Tavares JM, Araújo NAM, Telo da Gama MM. Dynamics of a network fluid within the liquid–gas coexistence region. *Soft Matter* **14**, 2744-2750 (2018).

96. Speedy RJ, Debenedetti PG. Persistence time for bonds in a tetravalent network fluid. *Mol Phys* **86**, 1375-1386 (1995).

97. Espinosa JR, Joseph JA, Sanchez-Burgos I, Garaizar A, Frenkel D, Collepardo-Guevara R. Liquid network connectivity regulates the stability and composition of biomolecular condensates with many components. *PNAS* **117**, 13238-13247 (2020).

98. Bai W, Sargent CJ, Choi J-M, Pappu RV, Zhang F. Covalently-assembled single-chain protein nanostructures with ultra-high stability. *Nat Commun* **10**, 3317 (2019).

99. Seeger M. Gaussian processes for machine learning. *Int J Neural Syst* **14**, 69-106 (2004).

100. Pedersen JS. Analysis of small-angle scattering data from colloids and polymer solutions: modeling and least-squares fitting. *Adv Colloid Interfac* **70**, 171-210 (1997).

101. Hansen J-P, McDonald IR. *Theory of simple liquids: with applications of soft matter*, Fourth edition. edn. Elsevier (2013).

1144
1145   102.   Chandler D, Weeks JD, Andersen HC. Van der waals picture of liquids, solids, and phase
1146   transformations. *Science* **220**, 787-794 (1983).
1147

1148   103.   Widom B. Intermolecular Forces and the Nature of the Liquid State: Liquids reflect in their
1149   bulk properties the attractions and repulsions of their constituent molecules. *Science* **157**,
1150   375-382 (1967).
1151

1152   104.   Choi JH, Lee H, Choi HR, Cho M. Graph Theory and Ion and Molecular Aggregation in
1153   Aqueous Solutions. *Annu Rev Phys Chem* **69**, 125-149 (2018).
1154

1155   105.   Bako I, Pusztai L, Pothoczki S. Topological descriptors and Laplace spectra in simple
1156   hydrogen bonded systems. *J Mol Liq* **363**, (2022).
1157

1158   106.   Pusztai L, Bako I, Pothoczki S. Connecting Diffraction Experiments and Network Analysis
1159   Tools for the Study of Hydrogen-Bonded Networks. *J Phys Chem B* **127**, 3109-3118 (2023).
1160

1161   107.   Agayan GM, Balabaev NK, Rodnikova MN. Description of Mixed Networks of H-Bonds
1162   in a Water-Ethylene Glycol System by Methods of Graph Theory and Delaunay Simplices.
1163   *Russ J Phys Chem A* **95**, 1283-1290 (2021).
1164

1165   108.   Faccio C, Benzi M, Zanetti-Polzi L, Daidone I. Low- and high-density forms of liquid
1166   water revealed by a new medium-range order descriptor. *J Mol Liq* **355**, (2022).
1167

1168   109.   de Oliveira PMC, de Souza JIR, da Silva JAB, Longo RL. Temperature Dependence of
1169   Hydrogen Bond Networks of Liquid Water: Thermodynamic Properties and Structural
1170   Heterogeneity from Topological Descriptors. *J Phys Chem B* **127**, 2250-2257 (2023).
1171

1172   110.   Tan AR, Urata S, Yamada M, Gomez-Bombarelli R. Graph theory-based structural
1173   analysis on density anomaly of silica glass. *Comp Mater Sci* **225**, (2023).
1174

1175   111.   Choi JH, Cho M. Ion aggregation in high salt solutions. II. Spectral graph analysis of water
1176   hydrogen-bonding network and ion aggregate structures. *Journal of Chemical Physics* **141**,
1177   (2014).
1178

1179   112.   Kihara T, Koba S. Crystal Structures and Intermolecular Forces of Rare Gases. *J Phys Soc
1180   Jpn* **7**, 348-354 (1952).
1181

1182   113.   Musacchio A. On the role of phase separation in the biogenesis of membraneless
1183   compartments. *The EMBO Journal* **41**, e109952 (2022).
1184

1185   114.   Russo J, Leoni F, Martelli F, Sciortino F. The physics of empty liquids: from patchy
1186   particles to water. *Rep Prog Phys* **85**, (2022).
1187

1188   115.   Chremos A, Panagiotopoulos AZ, Koch DL. Dynamics of solvent-free grafted
1189   nanoparticles. *Journal of Chemical Physics* **136**, (2012).

1190
1191 116. Gibbs E, Perrone B, Hassan A, Kummerle R, Kriwacki R. NPM1 exhibits structural and
1192   dynamic heterogeneity upon phase separation with the p14ARF tumor suppressor. *J Magn*
1193   *Reson* **310**, (2020).
1194
1195 117. Bianchi E, Largo J, Tartaglia P, Zaccarelli E, Sciortino F. Phase Diagram of Patchy
1196   Colloids: Towards Empty Liquids. *Physical Review Letters* **97**, 168301 (2006).
1197
1198 118. Sciortino F, Zaccarelli E. Reversible gels of patchy particles. *Current Opinion in Solid*
1199   *State and Materials Science* **15**, 246-253 (2011).
1200
1201 119. Edmond E, Ogston AG. An approach to the study of phase separation in ternary aqueous
1202   systems. *Biochemical Journal* **109**, 569-576 (1968).
1203
1204 120. Cheng Y, Patel DJ. An efficient system for small protein expression and refolding.
1205   *Biochemical and Biophysical Research Communications* **317**, 401-405 (2004).
1206
1207 121. Feder TJ, Brust-Mascher I, Slattery JP, Baird B, Webb WW. Constrained diffusion or
1208   immobile fraction on cell surfaces: a new interpretation. *Biophys J* **70**, 2767-2773 (1996).
1209
1210 122. Heller WT*, et al.* The suite of small-angle neutron scattering instruments at Oak Ridge
1211   National Laboratory. *J Appl Crystallogr* **51**, 242-248 (2018).
1212
1213 123. Arnold O*, et al.* Mantid-Data analysis and visualization package for neutron scattering and
1214   mu SR experiments. *Nucl Instrum Meth A* **764**, 156-166 (2014).
1215
1216 124. Heller WT*, et al.* drtsans: The data reduction toolkit for small-angle neutron scattering at
1217   Oak Ridge National Laboratory. *Softwarex* **19**, (2022).
1218
1219 125. Wignall GD, Bates FS. Absolute Calibration of Small-Angle Neutron-Scattering Data. *J*
1220   *Appl Crystallogr* **20**, 28-40 (1987).
1221
1222 126. Thompson AP*, et al.* LAMMPS - a flexible simulation tool for particle-based materials
1223   modeling at the atomic, meso, and continuum scales. *Computer Physics Communications*
1224   **271**, 108171 (2022).
1225
1226 127. Humphrey W, Dalke A, Schulten K. VMD: Visual molecular dynamics. *J Mol Graph*
1227   *Model* **14**, 33-38 (1996).
1228
1229 128. Virtanen P*, et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python.
1230   *Nat Methods* **17**, 261-272 (2020).
1231
1232 129. Michaud-Agrawal N, Denning EJ, Woolf TB, Beckstein O. Software News and Updates
1233   MDAnalysis: A Toolkit for the Analysis of Molecular Dynamics Simulations. *Journal of*
1234   *Computational Chemistry* **32**, 2319-2327 (2011).
1235

1236   130.   Harris CR*, et al.* Array programming with NumPy. *Nature* **585**, 357-362 (2020).

1237

1238   131.   Hunter JD. Matplotlib: A 2D graphics environment. *Comput Sci Eng* **9**, 90-95 (2007).

1239

1240   132.   Waskom M. seaborn: statistical data visualization. *Journal of Open Source Software* **6**,
1241   (2021).

1242

Acknowledgments

This work was supported by the St. Jude Children's Research Hospital Research Collaborative on the Biology and Biophysics of RNP granules (to R.V.P and R.W.K), NIGMS (R01 GM115634 and R35 GM131891 to R.W.K.), NCI (P30 CA021765 to St. Jude Children's Research Hospital), ALSAC (supporting studies at St. Jude Children's Research Hospital), and National Research Foundation (NRF) grants of Korea (2021R1C1C1010943 and 2022R1A4A1033471 to J.-M.C.). A portion of this research, conducted at the Oak Ridge National Laboratory (ORNL) Spallation Neutron Source, was sponsored by the Scientific User Facilities Division, Office of Basic Energy Sciences, U.S. Department of Energy. S. R. Cohen acknowledges financial support via T32 EB028092 from the US National Institutes of Health. We thank Jared M. Lalmansingh for technical assistance with CAMPARI. Fluorescence microscopy images were acquired at the St. Jude Cell & Tissue Imaging Center at St. Jude Children's Research Hospital (supported by P30 CA021765); we thank V. Frohlich, J. Peters, A. Taylor, A. Pitre, and G. Campbell for technical assistance.

Author contributions

R.V.P., J-M.C., and R.W.K came up with the project idea. D.M.M. and A.H.P. prepared samples for measurements. D.M.M., A.H.P, and C.B.S., performed SANS measurements and analyzed SANS data. D.M.M. and A.H.P., designed and characterized the phase behaviors of mutants. J-M.C designed the simulations, and prototyped the CAMELOT-based coarse-graining, and the original simulations using LAMMPS. F.D. performed all the LAMMPS simulations reported in this work. F.D., S.R.C., and R.V.P. designed and iterated on the structure of the analysis with inputs from J-M.C and R.W.K. F.D. and S.R.C. deployed the entirety of analyses, including of the SANS data, and integrated the findings with experimental work. F.D., S.R.C., and A.H.P. made the figures. F.D., S.R.C., and R.V.P. wrote the manuscript.

Competing interests

R.V.P. is a member of the scientific advisory board and shareholder of Dewpoint Therapeutics Inc. D.M.M. is an employee and shareholder of Dewpoint Therapeutics. The work reported here was not influenced by these affiliations. The remaining authors have no competing interests to declare.