1

Genetic and pharmacologic alterations of claudin9 levels suffice to induce functional

and mature inner hair cells.

Yingying Chen*[1,2], Jeong Han Lee[*1], Jin Li[3], Seojin Park[1,4], Maria C. Perez Flores[1], Braulio

Peguero[5], Jennifer Kersigo[6], Mincheol Kang[1,4], Jinsil Choi[1], Lauren Levine[7], Michael Anne

Gratton[7], Bernd Fritzsch[6], Ebenezer N. Yamoah[1±]

[1]University of Nevada, Reno, School of Medicine, Department of Physiology and Cell Biology,
Reno NV 89557
[2] Indiana University School of Medicine, Department of Pharmacology and Toxicology,
Indianapolis, IN, 46202, USA
[3]Department of Otolaryngology, University of Washington Seattle, WA, USA
[4]Prestige Biopharma, 11-12F, 44, Myongjigukje7-ro, Gangseo-gu, Busan, South Korea 67264
[5]Otolaryngology-Head, Neck Surgery, St. Louis University, St. Louis, Missouri 63108,
[6]University of Iowa, Department of Biology Iowa City, IA
18

The number of figures: 7

Supplement figures: 10

The number of pages: 50

Abstract: 125

Text:

#Corresponding author with complete address, including email address:
*Authors contributed equally and are co-first authors.
Ebenezer N. Yamoah
Department of Physiology and Cell Biology
School of Medicine
1664 N Virginia St
Reno, NV 89557
enyamoah@gmail.com
32

34 *In memory of the late Dr. Bruce Tempel, who helped to generate the mouse model.*

Hearing loss is the most common form of sensory deficit. It occurs predominantly due to hair cell (HC) loss. Mammalian HCs are terminally differentiated by birth, making HC loss challenging to replace. Here, we show the pharmacogenetic downregulation of *Cldn9*, a tight junction protein, generates robust supernumerary inner HCs (IHCs) in mice. The ectopic IHC shared functional and synaptic features akin to typical IHCs and were surprisingly and remarkably preserved for at least fifteen months >50% of the mouse's life cycle. *In vivo*, *Cldn9* knockdown using shRNA on postnatal days (P) P2-7 yielded analogous functional ectopic IHCs that were equally durably conserved. The findings suggest that Cldn9 levels coordinate embryonic and postnatal HC differentiation, making it a viable target for altering IHC development pre- and post-terminal differentiation.

Introduction

Mammalian cochlear hair cells (HCs) comprise a single row of inner hair cells (IHCs) and three rows of outer hair cells (OHCs). HCs transduce sound-mediated mechanical force into neural electrical codes for ear-brain intercommunication. The IHCs are the predominant afferent transducers, while the OHCs amplify low-level sound. Mammalian HCs are terminally differentiated by birth, and they are susceptible to damage by ototoxic drugs, noise-overexposure, aging, and environmental insults, resulting in hearing loss, the most common sensory deficit (Neitzel and Fligor, 2019; Rybak and Ramkumar, 2007; Wilson and Tucci, 2021; Wu et al., 2020). Emerging understanding of the mechanisms of transcription factors that induce HC differentiation (Atkinson et al., 2014; Bermingham et al., 1999; Garcia-Anoveros et al., 2022; Zheng and Gao, 2000), and potential induction for regeneration are promising, but none have produced new HCs with sustained functions (Iyer et al., 2022b; Zine et al., 2021). Thus, the current treatment for hearing loss ensuing from HC loss is cochlear implants, despite the potential advantages of HC-replacement therapy (Hinton et al., 2021; Sullivan et al., 2020).

In the mammalian cochlea, each HC is separated from the next by intervening supporting cells (SCs), forming an invariant and alternating mosaic along the cochlea's length. Cochlear SCs can divide and trans-differentiate into HCs in the neonatal stage, serving as a potential resource for HC differentiation using transcription and developmental signaling factors (White et al., 2006). At the neonatal stage, *Atoh1*, a basic helix-loop-helix factor, induces SCs trans-differentiation into HCs. Upregulation of *Atoh1, GFI1,* and *POU4F3* triggers HC differentiation, but the fledgling HCs invariably degenerate, suggesting pre-maturity (Iyer et al., 2022a; Liu et al., 2012). Additionally, inhibition of the Notch signaling or upregulation of the wnt1 pathway suffices to drive HC formation from SCs (Tang et al., 2023), but the functional features of the newly developed HCs are circumspect (Mizutari et al., 2013; Waqas et al., 2016). While transcription factors are potent targets for developmental regulation, a cadre of these essential DNA-binding proteins and the precise timing of expression are required to complete the

developmental cascade (Jahan et al., 2015). Thus, the key is identifying HC-developing bands

of transcription and signaling factors to treat hearing loss. Recent studies identified the

transcription factors INSM1 and IKZF2 as the regulators of OHC fate, while the transcription

factor TBX2 specifies and maintains HC and SC fate (Kaiser et al., 2022; Li et al., 2023),

advancing understanding of HC-subtype developmental specification mechanisms.

Contact-mediated lateral inhibition is among the final developmental events, where once

a cell fate is determined, it inhibits neighboring cells from becoming that cell type. The HC-SC

interphase is laced with tight junction proteins (TJPs), which may mediate lateral inhibition

mechanisms in nonmammalian vertebrates, though their additional function is unclear in

mammals (Chrysostomou et al., 2012; Goodyear and Richardson, 1997; Petrovic et al., 2014).

In addition, TJPs were also found to regulate cell proliferation (Bhat et al., 2020; Díaz-

Coránguez et al., 2019). The TJP scaffold cingulin regulates lateral inhibition in HC-SC

rearrangements in the avian basilar papilla (Goodyear and Richardson, 1997), and claudin b

(*cldnb*), an ortholog of the *cldn4* in humans, upregulation controls cellular patterning during HC

regeneration in zebrafish (Montalbano et al., 2021). Moreover, damaged HC extrusion and the

breaking of intercellular junctional adhesions may trigger differentiation and regenerative

proliferation (Corwin et al., 2007). In mammals, E-cadherin's junctional expression negatively

alters HC regenerative capacity (Burns et al., 2008; Burns et al., 2013), while *Cldn9* is a positive

regulator of cell proliferation (Hong et al., 2014; Zavala-Zendejas et al., 2011). Although the high

expression of *Cldn9* in the organ of Corti (OC) is well-established (Kudo et al., 2018; Nakano et

al., 2009) (**Fig. 1A-C**), it is unknown whether regulation of the TJP contributes to sensory cell

differentiation.

To determine the roles in vivo of Cldn9, we generated doxycycline (dox)-tet-OFF-Cldn9

transgenic mice to regulate expression levels of Cldn9. The downregulation of Cldn9 resulted in

functional supernumerary (ectopic) putative IHCs along the cochlear contour. Auditory neurons

innervated ectopic mechanically transducing IHCs with synaptic features resembling normal

IHCs. Analogous additional putative IHCs differentiation was observed when Cldn9-shRNA was

injected through the round window to postnatal (P) days (P2, P4, and P7) mice, suggesting that

regulation of Cldn9 levels coordinates embryonic and postnatal development differentiation of

SC into IHCs. Notably, the ectopic IHCs at the apical and middle-frequency contour of the

cochlea were preserved for over half the mouse's life cycle (15 months), making Cldn9 a viable

target for generating supernumerary IHCs.

Results

To control *Cldn9* levels *in vivo*, we generated a mouse model with a site-specific genetic

switch regulated by dietary doxycycline (dox) and dox-containing drinking water without

interfering with the typical profile of *Cldn9* expression ((Supplement figure 1 (**S1**)). **Figures 1D-F**

show the design constructs and Southern blot analysis to confirm the insertion cassette in the

ES cells. The genotyping was performed using PCR with the tail tissue. Transgene was

generated on mixed C57/B6 and backcrossed into a CBA-CaJ (CBA) background after 12

generations to reduce accelerated progressive hearing loss (Peguero and Tempel, 2015;

Sergeyenko et al., 2013; Willott and Erway, 1998). Results of quantitative RT-PCR from the

three groups of animals, including wild-type littermates (*Cldn9$^{+/+}$*), heterozygote *(Cldn9$^{+/T}$)* and

homozygous (*Cldn9$^{T/T}$*) with (1 mg/ml) and without dox treatment. Without dox treatment,

Cldn9$^{T/T}$ and *Cldn9$^{+/T}$* mice demonstrate ~55 and a 40-fold increase in *Cldn9* mRNA expression

in cochlear tissue compared with *Cldn9$^{+/+}$* cochleae (**S2**). Treatment of *Cldn9$^{+/T}$* mice with dox (1

mg/ml) resulted in an ~0.4-0.6-fold decline in mRNA levels compared to *Cldn9$^{+/+}$* cochleae (**S2)**,

translating to a marked difference in Cldn9 protein expression (**S2**). Immunoelectron

microscopic analysis showed that Cldn9 levels reduced by ~8-fold in the *Cldn9$^{+/T}$* cochlea (**S3**).

The *Cldn9$^{T/T}$* mice had a reduced survival rate (1.5±0.3 months) relative to the *Cldn9$^{+/T}$*

littermates (23±2 months*).* Thus, all experiments were restricted to *Cldn9$^{+/T}$ and Cldn9$^{+/+}$*

littermates fed on dox-water (1 mg/ml). There were no recognizable differences in body weight

between *Cldn9$^{+/T}$* and *Cldn9$^{+/+}$* mice for both females and males (**S2**). All animals were in the

CBA background. *Cldn9* downregulation in the *Cldn9$^{+/T}$* cochlea showed a qualitative decrease

in the Cldn6 and an increase in ILDR1 TJP levels but no comparative differences in others

(Kitajiri et al., 2004; Kitajiri and Katsuno, 2016) (**S4**).

Downregulation of *Cldn9* induces the production of ectopic cochlear HCs.

5-week-old mice *Cldn9$^{+/T}$* cochleae displayed a notable row of ectopic HCs (**Fig. 2A-E)**.

The ectopic HCs were observed along the cochlear contour, ranged in abundance from base to

apex (**S5**), and had contact with innervating neurons, shown in cochlear sections (**Fig. 2A)**. The

ectopic HCs were positively labeled with anti-myosin VIIa antibody and phalloidin-labeled

stereocilia bundles, which are features of typical HCs. Moreover, the ectopic HCs were not

labeled with antibodies to prestin (**S6**), a marker for OHCs, suggesting the new HCs were likely

derived from the IHC lineage. A distinctive "U"-shaped IHC bundle was apparent (**Fig. 2D)**.

Scanning electron microscopic (SEM) was used for high-resolution analyses to evaluate IHC

and their bundle morphology. The original and ectopic IHCs in *Cldn9$^{+/T}$* mice had normal

morphology, including intact hair bundles (**Fig. 2E**). However, the stereocilia bundle orientation

was less orderly when compared to those in *Cldn9$^{+/+}$* cochlea (**Fig. 2D-E**). In addition to their

shape, the ectopic HCs expressed multiple CtBP2 labeled synaptic structures in contrast to

typical OHCs (**Figs. 3B, 5B**), and reacted positively to otoferlin antibodies (Pangrsic et al., 2010;

Roux et al., 2006; Strenzke et al., 2016) (data not shown). We denote the new HCs as "ectopic"

IHCs. IHC counts at different ages (P2-P21) and the cochlear frequency segments (8 kHz and

32 kHz) demonstrate that the *Cldn9*-induced ectopic IHCs were most prominent at the cochlear

apex but remained statistically significant at the base (**Figs 2E-F**). Viable ectopic IHCs were

identified in 15-month *Cldn9$^{+/T}$* mice in the apical cochlea (**Fig. 3**). For OHCs, the numbers

along the cochlear contour, apex, middle, and base were not significantly different among the

two genotypes at P14 (**S7A)** and at the apical turn of organ of Corti in 15-month-old *Cldn9$^{+/T}$*

mice (**S7B**).

Functional features of *Cldn9*-induced ectopic IHCs resemble normal IHCs.

Motor responses of both mouse genotypes' ($Cldn9^{+/+}$ and $Cldn9^{+/T}$) to auditory stimuli

(Preyer's test) were normal. To evaluate the status of IHC function, we analyzed auditory

brainstem responses (ABR) to various sound-pressure levels. $Cldn9^{+/T}$ mice responses

exhibited similar characteristic responses to broadband clicks and pure tones at 8, 16, and 32

kHz stimuli (**Fig. 4A**), with ~5-15 dB threshold elevation in the $Cldn9^{+/T}$ mice [F (12, 579) =2.07,

P=0.02]. The pattern of hearing threshold remained virtually constant from 2-8 months of

monitoring [F (12, 319) =0.50, P=0.91]. Moreover, $Cldn9^{+/+}$ and $Cldn9^{+/T}$ mice yielded similar

distortion products [F (6, 357) =0.45, P=0.84] (**Fig. 4B**), suggesting normal OHC function.

Besides, a significant drop of ABR wave 1 amplitude was observed in the 2 months $Cldn9^{+/T}$

mice comparing to the wild-type littermates (8 kHz F (10, 235)=2.91, P=0.02; 16 kHz F (10,

237)=1.99, P=0.04; 32 kHz F (10, 233)=7.80, P=0.00; click F (10, 233)=3.99, P=0.00 (**S8A-D**).

However, wave 1-latency shows no difference between $Cldn9^{+/T}$ and $Cldn9^{+/+}$ mice at 2 months

(**S8E-H**). The apparent normal morphology of the ectopic IHCs led to the hypothesis that the

IHCs may exhibit functional mechano-electrical transducer (MET) currents. Since active MET

channels are partially open at rest, the rapid uptake of the channel permeable lipophilic dye

FM1-43 (Gale et al., 2001), was assessed. Results were determined from FM1-43 dye loading

of apically located original and ectopic IHCs in 4-week-old $Cldn9^{+/T}$ cochlear site representing

~4-6-kHz characteristic frequencies (CFs). Local perfusion of 10-μM FM1-43 resulted in intense

dye labeling at the hair bundle level of original and ectopic IHCs. The dye-membrane

partitioning and diffusion across the IHCs' basal aspects occurred within seconds. Z-stacked-

time-lapse images were taken below hair bundle level 2-sec post dye exposure and at the

basolateral compartment (**Fig. 4C**). The time constants (τ) of dye loading at the bundle and

supra-nuclear basal membrane levels for original IHCs were (data from 4 mice); 19.4±1.1 sec

(n=27) and 29.0±3.8 sec (n=27), and for ectopic IHCs were 23.1±4.5 sec (n=27) and 47.0±8.4

sec (n=27) (**Fig. 4D**). These results are consistent with functioning IHCs at rest. We conclude

that the ectopic IHC bundles are set at the optimum dynamic range to transduce MET current at

rest like the original counterparts. IHC MET current magnitudes and kinetic profiles from the

original and ectopic rows at the ~3-4-kHz cochlear-place map in P21 mice were comparable, as

summarized in figures **4E-G**. The normalized current-displacement relationships were well-fitted

with a two-state Boltzmann function, portraying marked similarities between the original and the

ectopic IHC MET currents (**Fig. 4H**). However, the displacement-response relationship for the

ectopic IHCs was right-shifted, indicative of reduced sensitivity.

Synaptic features of ectopic IHCs and original IHCs.

To determine whether the ectopic IHCs had additional properties in terms of systemic

functions, we examined features such as neuronal innervation in both the original and ectopic

IHCs. We labeled auditory neurons and IHCs with calretinin (Calb2) antibodies (Sun et al.,

2018). Results show Calb2-positive-subtype neurites at the modiolar aspects of both IHCs (**Fig.**

5A). The synapses between the IHCs and auditory neurons at the apical, middle, and basal

cochlear locations from 5-week-old *Cldn9$^{+/+}$* and *Cldn9$^{+/T}$* mice show substantial differences. The

organization of afferent synapses was significantly different, identified as paired presynaptic-

CtBP2 (red) and postsynaptic-Homer1 (green) immunopuncta. Contrasting *Cldn9$^{+/+}$* from

Cldn9$^{+/T}$ cochlear samples, results showed reduced synaptic numbers in the *Cldn9$^{+/T}$* (**Fig. 5B-**

C). Moreover, quantifying the mean number of synapses per IHC among the original and

ectopic IHCs showed variations along the cochlear axis (**Fig. 5D**). This data suggests that

ectopic IHCs are equipped to serve as afferent receptors capable of transducing and

transmitting mechanical displacement into neural codes, while synaptic transmission from

ectopic IHCs might be compromised because significant differences in synapse numbers were

identified.

Postnatal induction of ectopic IHCs by shRNA knockdown.

Because pregnant mothers were fed on dox-water from gestation, the ectopic IHCs in

Cldn9$^{+/T}$ cochlea were of embryonic origin. We designed an efficient shRNA construct to knock

down *Cldn9* postnatally. Viral transduction *in vivo*, through round window injection into the cochlear tissue, was monitored with a GFP reporter gene (**Fig. 6A**). We first assessed the efficiency and specificity of the shRNA knockdown of *Cldn9* using quantitative RT-PCR (**S9A**) and immunofluorescence microscopy (**Fig. 6A**). Four days post-injection, there was an ~20-fold reduction in *Cldn9* mRNA relative to nontargeting scrambled shRNA injected cochlea (**Fig. 6A, S9**). Transduction *in vivo* of *Cldn9* shRNA into P2-7 inner ears yielded cochleae with ectopic IHCs compared to internal controls, consisting of opposite cochlea injected with nontargeting scrambled shRNA injected, which did not display ectopic IHCs. By contrast, the P14 inner ear injected with *Cldn9*-shRNA produced no detectable increase in ectopic IHCs as counted by three independent blinded observers (**Fig. 6B-D**)**.** Ultrastructural SEM analysis of *Cldn9*-shRNA transduced P2-7 inner ears show ectopic IHC with hair bundles resembling original IHCs (**Fig. 6C**). The MET currents invoked from the ectopic-IHCs induced by postnatal *Cldn9* knockdown were in keeping with functional HCs.

The endocochlear potential and K$^+$ concentrations in *Cldn9*$^{+/T}$ mice

The cochlear duct is furnished with cellular syncytia, K$^+$ channels, and transporters/pumps that operate to orchestrate a unidirectional (basal to apical) flux of K$^+$ at the lateral wall to produce the endocochlear potential (EP, ~+80 mV), an extracellular potential, subserving the proverbial powerhouse for HC functions (Von Bekesy, 1952). A remnant of K$^+$ flux is a high K$^+$ endolymph (~140 mM) restricted from leakage into the basolateral aspects of HCs by TJ between HCs and SCs. At four months of age, the magnitude of the EP and the K$^+$ concentration of the endolymph and perilymph of *Cldn9*$^{+/T}$ mice were variable significantly from age-matched littermates. A slight decline in the amplitude of the EP and a substantial rise in perilymph K$^+$ was detected in 8-month-old *Cldn9*$^{+/T}$, but there is no statistical difference compared with the detection in 4-month-old Cldn9$^{+/T}$(**S10**). It is unclear whether the modest changes in the EP and K$^+$ concentration of perilymph can account for the threshold increase in the *Cldn9*$^{+/T}$ mice.

Discussions

Fate determination is typically completed by birth in cochlear HCs, the primary receptors for mechanosensory sound detection. Generating functional HCs in the mammalian cochlea, with a proper cellular organization that allows for cochlear sound frequency selectivity, has been a demanding yet unsolved challenge. Consequently, deafness resulting from HC loss, which constitutes a significant portion of sensorineural hearing loss, is incurable. Multiple aspects must be considered in inducing HC differentiation since HCs have distinct functional and structural features along the cochlear axis. For optimal sensing of different sound frequencies, cochlear apical-to-basal HCs have diverse structural configurations and ion channel configurations and densities that sculpt and preserve low-to-high frequency sound processing (Kimitsuki et al., 2001). Additionally, HCs are connected to SCs by TJ proteins, enhancing the sensitivity of cochlear OHC sound amplification and maintaining high K^+ concentration in the endolymph at the apical surface and low K^+ concentration in the perilymph milieu surrounding HCs (Cohen-Salmon et al., 2002). This process establishes a unidirectional K^+ flux in the cochlear duct to generate the EP, which boosts the receptor potential of HC by ~80 mV (Hudspeth, 2008; Von Bekesy, 1952). Thus, besides overcoming the insurmountable terminal differentiation of HCs, newly differentiated HCs *in vivo* should be equipped with features that resemble the original primary HCs to integrate into the specialized cochlear environment.

Among the multiple approaches used with limited success in HC replacement are overexpressing transcription factors involved in prosensory cell differentiation and silencing inhibitory factors in the induction of HC fate. These methods include 1) Transfecting proneural genes, such as *Atoh1*, in embryonic, newborn, and damaged mature cochlear tissue. However, HC-competent cells invariably lose their responsiveness post-birth and in adult animals. In the case of the damaged cochlea, it is challenging to distinguish between transdifferentiated and repaired HCs (Gubbels et al., 2008; Izumikawa et al., 2005; White et al., 2006). 2) Simulating

cell division using cell-cycle inhibitors, this strategy activates apoptotic genes, leading to cell

death and deafness (Lowenheim et al., 1999). 3) Inhibiting Notch signaling using a γ-secretase

inhibitor to stimulate HC differentiation from potential inner ear resident stem cells after noise

trauma (Jeon et al., 2011; Mizutari et al., 2013). The scarcity of resident stem cells in adult

cochlea may limit this strategy. Thus, multiple therapeutic armamentariums are required to

restore hearing in the translational setting.

261   Previous studies demonstrated that claudin-9 is essential for hearing function and the

maintenance of auditory HCs, using an ethyl nitrosourea-induced *Cldn9* mutant mouse model

(Nakano et al., 2009), which resulted in OHC degeneration.

The current results demonstrate that embryonic regulation of *Cldn9* levels, but not null deletion

using the dox-tet-OFF-*Cldn9* transgenic strategy induces functional ectopic IHCs, but not OHCs,

along the cochlear contour with increasing numbers from base to apex. These *Cldn9*

downregulated-induced IHCs mature, acquiring robust MET currents and neural innervation with

synaptic structures markedly like resident IHCs. Results also revealed that postnatal

downregulation of *Cldn9* levels *in vivo*, using shRNA, suffice to coordinate SC differentiation into

IHCs. Because the ectopic IHCs remain viable for a sizable duration of the mouse's lifespan, the

Cldn9 regulatory strategy to induce IHC differentiation subserves a feasible approach to replace

lost HCs. The downregulation of Cldn9-mediated selective IHC-increase indicates Cldn9's role

during the latter phase of HC differentiation, perhaps post-OHC-fate determination (Garcia-

Anoveros et al., 2022). The findings suggest Cldn9-mediated effects may be upstream of the

transcriptional factor-mediated trans-differentiation of HCs since the ectopic HCs had features of

IHCs, in contrast to primordial HCs generated by *Atoh1* (Yang et al., 2012). It deepens our

understanding of the importance of tapping into later stages of HC differentiation, which will

likely result in end-organ-specific HCs.

279   Moreover, a pragmatic strategy requires titrated levels of the TJP to render new HCs

without compromising the sensory epithelial cellular syncytial, a decline in the EP, and,

significantly, a gradual extracellular K$^+$ increase that mediates undue HC depolarization and

death (Nakano et al., 2009). The critical period at which alteration of TJP level can induce

ectopic and new IHCs remains unclear, although, in the current report, we have demonstrated

that functional and viable mature ectopic IHCs can be generated by regulating Cldn9 levels.

Our findings that downregulation in the TJP, Cldn9, can regulate IHC differentiation are

in conceptual agreement with reports demonstrating that lateral inhibition can affect HC

specification (Lanford et al., 1999; Stone and Rubel, 1999). Similar accounts are described

where newly formed HCs express delta1-like (Dll1) and jagged 2 (Jag2) ligands to mediate

Notch1-receptor activation in adjacent antecedent cells, thereby inducing the expression of hairy

and enhancer of split (Hes1/5), which suppresses pro-HC transcription factors (Bermingham et

al., 1999; Chrysostomou et al., 2012). Consistent with this scheme, interruption of Notch1

signaling during HC development leads to HC overproduction (Brooker et al., 2006; Lanford et

al., 1999; Zine et al., 2000). In lateral inhibition, the foremost developing cells adopting HC fate

antagonize the neighboring cells from differentiating into HCs through direct cell-to-cell

communication (Brown and Groves, 2020; Mizutari et al., 2013). A possible explanation for the

current findings is that TJ proteins, mainly Cldn9, are signaling in Notch-mediated lateral

induction (Daudet and Lewis, 2005; Lewis, 1998). Canonical Notch signaling is activated when a

Notch ligand, such as Delta-like1 (Dl1) in an adjacent HC, binds to a receptor in the SC,

resulting in the release of the intracellular domain of the Notch receptor (NICD), which

translocate to the nucleus to activate the Notch-targeted gene transcriptions. The findings also

confirm that the Notch signaling pathway is responsible for homeostatic TJP expression *in vitro*

and promotes barrier function *in vivo* in the RAG1-adoptive transfer model of colitis (Mathern et

al., 2014). Indeed, occluding junction depletion disrupts Notch and mitogen-activated protein

kinase (MAPK) signaling in intestinal tissue (Fairchild et al., 2016). In the scheme provided in

Figure 7, Cldn9 may subserve the signaling catalyst to activate NICD cascades that suppress

neighboring SCs from trans-differentiation. A limitation of the model is that if *Cldn9*-induced

effects were solely dependent on the Notch signaling, ectopic OHCs and IHCs would have

ensued. Alternatively, *Cldn9* levels disruption may alter the mechanical properties of the

developing and maturing organ of Corti that may trigger ectopic IHC differentiation, an

epiphenomenon independent of the Notch signaling. Future studies and emerging findings on

HC differentiation will likely address these shortcomings (Kaiser et al., 2022; Li et al., 2023). In

cochlear tissue, downregulation of Cldn9 led to concomitant reduced expression of Cldn6 and

increased ILDR1. It is unclear whether the induction of ectopic resulted from reduced

expression of *Cldn9* alone or combined TJP alterations. It is conceivable that targeting a

different TJP may have similar effects on OHC differentiation, requiring impending studies.

357 **Materials and Methods**
358
All procedures were performed under research guidelines of the institutional animal care

and use committee of the University of Nevada, Reno. Mice of either sex were studied.

Doxycycline (dox)-tet-OFF-Cldn9 transgenic mice were generated. In the mouse line, dox

concentration can regulate the level of Cldn9 gene expression. The construct consisted of a

tetracycline-based genetic switch (tTA cassette) made of three main modules: 1) The

tetracycline-controlled transcriptional activator (tTA); 2) The neomycin resistance gene flanked

by LoxP sites; and 3) Six copies of the tet operator (tetO) fused to the minimal CMV

promoter. The tTA cassette was inserted at the -110 -nucleotide position upstream of the

translational start of *Cldn9* to generate the targeting vector. The targeting vector was

electroporated into B6 mouse embryonic stem cells. Following the selection in G418, DNA

samples from the neomycin-resistant ES cell clones were prepared for short-arm

PCR/sequencing analysis and Southern blot analysis to confirm the insertion of the tTA cassette

into ES cells. Genetically modified ES cells containing one copy of the tTA cassette were

injected into healthy albino B6 blastocysts, and the injected blastocysts were transplanted into

the uterus of an albino B6 mouse to generate the chimeric mouse. The chimeric mouse was

then bred with albino B6 mice to produce the F1 heterozygous mouse, and the germline

transmission was confirmed by tail DNA genotyping. The selection marker was deleted in the

tTA cassette by crossing the F1 mouse with the embryonic Cre line (B6.129S4-*Meox2*[tm1(cre)Sor]/J).

We backcrossed the B6/129S4 background unto the CBA/CaJ mouse background for 12

generations to prevent the masking of age-related hearing loss effects. 1.0 mg/ml of dox water

were fed to *Cldn9* breeding pairs from breeding day one, and heterozygote *(Cldn9$^{+/T}$)* and

Homozygous (*Cldn9^{T/T}*) mice and wild-type littermates (*Cldn9^{+/+}*), through the time until for the

sample collections. The body weights of mice were recorded. Genotyping was performed by

PCR using a set of primers that flank the knockin in the *Cldn9* gene: forward primer Cldn9

knockin-F (knockin-sequence): 5'–ATCCACGCTGTTTTGACCTC–3', Cldn9 R3 (Reverse): 5'–

TCTGGACCACACAGGACATC– 3'. PCR fragments were separated with 2% agarose gel for an

800 bp product in wild-type, 365 bp in homozygous mutants, and two in heterozygous littermate

mice (**Fig. 1**).

Auditory brainstem recordings (ABR) and Distortion product otoacoustic emissions

(DPOAE) measurements.

Cldn9 mice (*Cldn9^{+/+}*, *Cldn9^{+/T}*, and *Cldn9^{T/T}*) littermates were tested at 2-8 months of

age. Mice were anesthetized with ketamine and xylazine by intraperitoneal (IP) injection

(25 mg/kg). Body temperature was monitored using a rectal probe and maintained at

36.8±1.0°C using a homeothermic device (Harvard Apparatus). ABR and DPOAE

measurements were described previously (Dou et al., 2004). For ABR assays, thresholds were

obtained by presenting tone bursts at 4, 8, 16, and 32 kHz and a clicking sound from 0 dB to 90

dB sound pressure levels (SPL) in 5 dB intervals. Tones were 2.5 ms, while click was 0.1 ms in

duration, with a repetition rate of 21/s. Electrodes were placed sub-dermally behind the tested

ear (reference), the vertex (active), and the back (ground). Evoked potentials were averaged

over 512 repetitions and collected using a Tucker Davis Technology (TDT) RZ6 processor and

BioSigRZ software. The threshold was defined as the lowest intensity of stimulation that yielded

a repeatable waveform based on an identifiable ABR wave.

DPOAE measurements were performed using the same TDT system with two calibrated

MF1 speakers connected to an ER10B+ microphone. Data was collected every 21 ms and

averaged 512 times. DPOAEs were recorded using two pure tones with frequencies f1 and f2,

using an f2/f1 ratio of 1.2. Input/output (I/O) functions were obtained by increasing the primary

tone L1 (and corresponding L2) in 5-dB steps from 20 to 80 dB SPL at 8, 16, and 32 kHz

frequencies. During DPOAE testing, the probe assembly was placed in the mouse's left ear

canal after visual inspection to ensure no ear infection or inflammation of the tympanic

membrane. DPOAE thresholds were defined as the lowest level of f1 required to produce a

DPOAE ≥ –5 dB SPL(Chen et al., 2021).

Cochlear mapping and hair cells and synaptic counts

The cochlea was micro-dissected into three to five pieces following the method

described (Montgomery and Cox, 2016). Cochlear pieces were measured, and a frequency map

was computed based on a 3D reconstruction of the sensory epithelium for HCs and synapse

count of associated structures to relevant frequency regions using a custom plug-in to ImageJ

(Muniak et al., 2013). Confocal z-stacks of the 4, 8, 16, and 32 kHz areas were collected using a

Leica Stellaris8 (Leica) and Nikon A1R laser scanning confocal microscope (Nikon Instruments

Inc.). Images were gathered in a 512 x 512 raster using a high-resolution oil immersion

objective (60x). IHCs and OHCs at the frequency locations were quantified using myosin-VIIa-

positive as an HC-marker within a 70-100-µm field (Chen et al., 2021). Synaptic ribbons and

homer-labeled puncta could be counted manually or automatically by setting the filters for the

spots function using 3D (x-y-z axis) representations of each confocal z-stack with the

microscopic image analysis software Imaris (Oxford Instruments, USA).

Inner ear histological analysis

The cochleae were intra-labyrinthine perfused through the oval and round windows with

4% paraformaldehyde (PFA). The samples were decalcified in 10% EDTA up to 72-96 hrs,

depending on the age, at 4°C. Micro dissected pieces were immunostained with antibodies to

the following: (1) mouse anti-C-terminal binding protein 2 (pre-synaptic-marker, BD Biosciences,

1:200, Cat # 612044), (2) rabbit anti-myosin-VIIa (HC-marker, Proteus Biosciences, Inc,1:600,

Cat # 25-6790), (3) mouse anti-sox2 (supporting cell (SC)-marker, Santa Cruz Biotechnology,

Inc, 1:200, Cat # sc-365823), and (4) rabbit anti-Homer 1 (post-synaptic marker, Synaptic

Systems, 1:250, Cat # 160 003), (5) rabbit anti-immunoglobulin like domain containing receptor

1 (ILDR1) (Antibodies-online.com, 1:200, Cat # ABIN1386369), (6) mouse anti-Cldn9 (Santa

Cruz, 1:200, Cat # sc-398836), (7) mouse anti-Cldn6 (Santa Cruz, 1:200, Cat # sc-393671), (8)

rabbit anti-Sox2 (Abcam, 1:200, Cat # ab97959), (9) rabbit anti-Tuj1 and chicken anti-Tuj1

(Abcam,1:500, Cat # ab18207, ab41489), (10) goat anti-calretinin (Swant Inc., 1:500, code #

CG1), (11) mouse anti-calretinin (Millipore Sigma, 1:200, Cat # MAB1568), (12) rabbit anti-

Prestin (Santa Cruz, 1:200, Cat # sc-22692), and (13) rabbit anti-calbindin (Cell signaling

technology, 1:200, Cat # 13176S) with appropriate secondary antibodies coupled to Alexa-405,

-488, -568, and -647 fluorophores.. DAPI labeled the cell nucleus after secondary antibody

incubation. Samples were stained with phalloidin and mounted with Fluoro-Gel (Electron

Microscopy Sciences). Images were captured with a confocal microscope.

RNA extraction and quantitative RT-PCR of Cochlear tissue

The cochlea was dissected from the mouse and homogenized on ice. Because of limited

tissue, we combined 10-15 mice cochleae for the study. Total RNA was isolated using the

RNeasy Plus Mini Kit (Qiagen), and cDNA was generated using the RT2 First Strand Kit

(Qiagen). cDNA was combined with RT2 SYBR Green Master Mix (Qiagen), specific qRT-PCR

primers, and qRT-PCR analysis was run using the ViiaTM 7 Real-Time PCR System (ABI).

Primer efficiencies were determined by standard dilution curve analysis. Three separate

samples were used from 10 animals for each group. The experiments from each sample were

performed in triplicate, and average cycle threshold (Ct) values were normalized to GAPDH

expression. ΔΔCt values were determined relative to $Cldn9^{+/+}$ cochlear samples. Fold change

was defined as 2(–ΔΔCt). Primers used include Gapdh (SA Biosciences) and Cldn9

(ThermoFisher).

Electron Microscopy

Transmission electron microscopy (TEM) and scanning electron microscopy (SEM) of

cochlear sensory epithelia were performed as described (Schwander et al., 2007; Senften et al.,

2006). Five to eight-week-old *Cldn9* mice and littermates were sacrificed for TEM, and the

cochleae were fixed in 2.5% glutaraldehyde in 0.1 M cacodylate buffer at 4°C overnight. After

several washes with buffer alone, cochleae were fixed in 1% osmium tetroxide at RT for 1-hr.

After that, the fixed cochleae were decalcified in 10% EDTA for 3–4 days. Fixed and decalcified

cochleae were dehydrated using a graded ethanol series and embedded in epoxy resin.

Ultrathin sections were cut with a diamond knife. Specimens were examined using an electron

microscope.

For SEM, mice were perfused with 4% PFA in 1x PBS, inner ears were isolated, and the

stapes footplate was removed. Ears were flushed and fixed overnight in 4% PFA and 2.5%

glutaraldehyde in 1x PBS. After washing in ddH$_2$O 3X for 1 hour, the samples were post-fixed

with 1% osmium tetroxide for approximately 1 hour. Samples were washed before decalcifying

for 3-4 days in 0.25 M EDTA at 4°C with daily solution changes. The cochleae were micro

dissected, the tectorial membranes removed, and gradually dehydrated in 30%, 50%, 70%, 80%,

90%, 100% ethanol, 2:1 ethanol/hexamethyldisilazane (HMDS, Thermo Scientific #A15139.AE),

1:2 ethanol/HMDS, and finally 100% HMDS. Samples were transferred to an open well plate in

HMDS and allowed to air dry overnight in a fume hood. They were then mounted on aluminum

stubs (Ted Pella #16111) using double-sided carbon tape (EMS #77817-12) and stored in a

specimen mount holder (EMS #76510) sealed in a desiccator until sputter-coated with Au/Pd

(Emitech Sputter Coater K550) and viewed. Images were captured utilizing the Hitachi S-4800

SEM. An accelerating voltage of 1kV and 5kV was used. Images were compiled using

CorelDRAW X7 graphic suite software.

Measurements of the endocochlear potential (EP) and K$^+$ concentration

Cldn9 heterozygote mice and wild-type littermates were anesthetized using ketamine

and xylazine (100/25 mg/kg, i.p.) and K$^+$ concentration, and the EP was measured using

double-barreled microelectrodes. An incision was made along the midline of the neck, and soft

tissues were bluntly dissected laterally to expose the trachea and the animal's left bulla. A

tracheostomy was done, and the musculature over the bulla was cut posteriorly to expose the

bone. A small hole was made in the cochlear capsule directly over the scala media of the lower

basal turn. The EP electrode was filled with 300 mM NaCl, the K^+-selective barrel was silanized,

and the tip was filled with a liquid ion exchange (Fluka 60398, K^+ ionophore I- Cocktail B) that

was backfilled with 150 mM KCl. A round-window approach made measurements in the basal

turn of the cochlea through the basilar membrane of the first turn. The K^+-selective electrode

was calibrated in solutions with known cation (K^+ and Na^+) concentrations *in situ* at 37°C.

shRNA CLDN9 knockdown

siRNAs were designed, using siRNA at WHITEHEAD software, and cloned under U6

promoter in pSilencer5.1-U6 vector to produce hairpin siRNAs (shRNAs; Vector Biolabs,

NM_020293). shRNA was packaged into adeno-associated virus constructs: AAV/Anc80L65-

GFP-U6-mCLDN9-shRNA (4) at 1.0×10^{12} GC/mL titer. shRNA sequence for mouse CLDN9 is

5'-CACC GTGCTTCGGGACTGGATAAGACTCGAGTCTTATCCAGTCCCGAAGCAC TTTTT-3'.

1.0 µl shRNA or scrambled shRNA was injected into the round window of mice at P2 – 7 and

P14. Mice were anesthetized by induced hypothermia and kept on a cold surface during the

injection procedure. After disinfecting the skin with 70% ethanol and Povidone iodine, an

incision was made in only the left ear (experiment) and right ear (scramble shRNA injection).

Underlying fat and soft tissue were carefully dissected to expose the round window of the

cochlea. The glass pipette was pulled with a P-2000 (Sutter Instrument, Novato, CA) and

sharpened with a BV-10 Micropipette Beveler (Sutter Instrument, Novato, CA). The shRNA was

injected using a nanoinjector (Sutter Instrument, Novato, CA). After all the shRNA was injected

into the round window, the pipette was left in place for 30 seconds before removal. The muscles

and fat tissue were covered, and the skin was closed with a polypropylene suture. The mice

were kept on the warm bedding heated by a heating pad for recovery before returning to their

mother. Total surgery time did not exceed 15 min.

Voltage-clamp recording of hair cell mechanoelectrical transducer (MET) current.

Patch-clamp experiments were performed in the standard whole-cell mode using an

Axopatch 200B amplifier (Axon Instruments). Patch electrodes were pulled with a horizontal

puller (Sutter Ins. Navato, CA) and had a resistance of 2–3 MΩ when filled with pipette solution

consisting of (in mM) 135 CsCl, 10 HEPES, 2.5 EGTA, 0.25 $CaCl_2$, $MgCl_2$, 4 MgATP, and 0.4

Na_2GTP (pH adjusted to 7.3 with CsOH). The bath solution consisted of (in mM) 130 NaCl, 3

KCl, 1 $MgCl_2$, 10 HEPES, 2.5 $CaCl_2$, and 10 glucose (pH was adjusted to 7.3 using NaOH).

Currents were sampled at 20 kHz and filtered at 2 kHz. Voltages were not corrected for a liquid

junction potential. No leak current subtraction was performed. Cells were held at -80 mV. All

electrophysiological experiments were performed at RT (21-22°C). We used stepwise and

sinewave mechanical stimulation of IHC bundles through a piezo-driven fluid-jet stimulator to

record IHC mechanoelectrical transducer (MET) currents. We represented the bundle

displacement in the form of applied piezo-driven voltage. Bundle displacement was not

calibrated for each cell because of variations in stimulating probe positions relative to the

stimulated hair bundle.

Data analysis

The ABR and DPOAE data were analyzed using GraphPad Prism 7 (GraphPad

Software, San Diego, CA, US) and OriginPro 2020 (OriginLab Corp., Northampton, Mass, US).

Three-way ANOVA was used to analyze the ABR and DPOAE threshold. Two-way ANOVA was

applied for the ABR wave 1 amplitude/latency data, synapse count, and HCs count. Significance

was assumed at a *p*-value of 0.05 in all statistical analyses.

530
531

532 **Acknowledgments**

533 We thank members of our laboratory for their comments on this manuscript. This work was

534 supported by grants from the National Institutes of Health (DC016099, DC05135, AG051443,

535 DC015252, and AG060504) to E.N.Y.

Author Contributions

538   E.N.Y and BF designed the research, Y.C. M.C.P.F., J.H.L, J.L, S.P, M.K, B.P, J.K, J.C, L.L,

539   MAG, and E.N.Y., analyzed data and wrote the manuscript. All authors read and approved the

540   final manuscript.

Author Information

542   Correspondence should be addressed to E.N.Y., the Program in Communication Science, the

543   Department of Translational Neuroscience, the College of Medicine, Phoenix AZ, University of

544   Arizona Phoenix, AZ. Email: enyamoah@arizona.edu

References
Atkinson, P.J., Wise, A.K., Flynn, B.O., Nayagam, B.A., and Richardson, R.T. (2014). Hair cell
regeneration after ATOH1 gene therapy in the cochlea of profoundly deaf adult guinea pigs.
PloS one *9*, e102077.
Bermingham, N.A., Hassan, B.A., Price, S.D., Vollrath, M.A., Ben-Arie, N., Eatock, R.A., Bellen,
H.J., Lysakowski, A., and Zoghbi, H.Y. (1999). Math1: an essential gene for the generation of
inner ear hair cells. Science *284*, 1837-1841.
Bhat, A.A., Syed, N., Therachiyil, L., Nisar, S., Hashem, S., Macha, M.A., Yadav, S.K.,
Krishnankutty, R., Muralitharan, S., and Al-Naemi, H. (2020). Claudin-1, a double-edged sword
in cancer. International journal of molecular sciences *21*, 569.
Brooker, R., Hozumi, K., and Lewis, J. (2006). Notch ligands with contrasting functions: Jagged1
and Delta1 in the mouse inner ear. Development *133*, 1277-1286.
Brown, R., and Groves, A.K. (2020). Hear, Hear for Notch: Control of Cell Fates in the Inner Ear
by Notch Signaling. Biomolecules *10*.
Burns, J., Christophel, J.J., Collado, M.S., Magnus, C., Carfrae, M., and Corwin, J.T. (2008).
Reinforcement of cell junctions correlates with the absence of hair cell regeneration in
mammals and its occurrence in birds. Journal of Comparative Neurology *511*, 396-414.
Burns, J.C., Collado, M.S., Oliver, E.R., and Corwin, J.T. (2013). Specializations of intercellular
junctions are associated with the presence and absence of hair cell regeneration in ears from
six vertebrate classes. Journal of Comparative Neurology *521*, 1430-1448.
Chen, Y., Bielefeld, E.C., Mellott, J.G., Wang, W., Mafi, A.M., Yamoah, E.N., and Bao, J. (2021).
Early physiological and cellular indicators of cisplatin-induced ototoxicity. Journal of the
Association for Research in Otolaryngology *22*, 107-126.
Chrysostomou, E., Gale, J.E., and Daudet, N. (2012). Delta-like 1 and lateral inhibition during
hair cell formation in the chicken inner ear: evidence against cis-inhibition. Development *139*,
3764-3774.
Cohen-Salmon, M., Ott, T., Michel, V., Hardelin, J.P., Perfettini, I., Eybalin, M., Wu, T., Marcus,
D.C., Wangemann, P., Willecke, K., *et al.* (2002). Targeted ablation of connexin26 in the inner
ear epithelial gap junction network causes hearing impairment and cell death. Curr Biol *12*,
1106-1111.
Corwin, J.T., Jones, J.E., Katayama, A., Kelley, M.W., and Warchol, M.E. (2007). Hair cell
regeneration: the identities of progenitor cells, potential triggers and instructive cues. Paper
presented at: Ciba Foundation Symposium 160-Regeneration of Vertebrate Sensory Receptor

Cells: Regeneration of Vertebrate Sensory Receptor Cells: Ciba Foundation Symposium 160
(Wiley Online Library).
Daudet, N., and Lewis, J. (2005). Two contrasting roles for Notch activity in chick inner ear
development: specification of prosensory patches and lateral inhibition of hair-cell
differentiation. Development *132*, 541-551.
Díaz-Coránguez, M., Liu, X., and Antonetti, D.A. (2019). Tight junctions in cell proliferation.
International Journal of Molecular Sciences *20*, 5972.
Dou, H., Vazquez, A.E., Namkung, Y., Chu, H., Cardell, E.L., Nie, L., Parson, S., Shin, H.S., and
Yamoah, E.N. (2004). Null mutation of alpha1D Ca2+ channel gene results in deafness but no
vestibular defect in mice. J Assoc Res Otolaryngol *5*, 215-226.
Fairchild, M.J., Yang, L., Goodwin, K., and Tanentzapf, G. (2016). Occluding Junctions Maintain
Stem Cell Niche Homeostasis in the Fly Testes. Curr Biol *26*, 2492-2499.
Gale, J.E., Marcotti, W., Kennedy, H.J., Kros, C.J., and Richardson, G.P. (2001). FM1-43 dye
behaves as a permeant blocker of the hair-cell mechanotransducer channel. J Neurosci *21*,
7013-7025.
Garcia-Anoveros, J., Clancy, J.C., Foo, C.Z., Garcia-Gomez, I., Zhou, Y., Homma, K., Cheatham,
M.A., and Duggan, A. (2022). Tbx2 is a master regulator of inner versus outer hair cell
differentiation. Nature *605*, 298-303.
Goodyear, R., and Richardson, G. (1997). Pattern formation in the basilar papilla: evidence for
cell rearrangement. Journal of Neuroscience *17*, 6289-6301.
Gubbels, S.P., Woessner, D.W., Mitchell, J.C., Ricci, A.J., and Brigande, J.V. (2008). Functional
auditory hair cells produced in the mammalian cochlea by in utero gene transfer. Nature *455*,
537-541.
Hinton, A.S., Yang-Hood, A., Schrader, A.D., Loose, C., Ohlemiller, K.K., and McLean, W.J. (2021).
Approaches to Treat Sensorineural Hearing Loss by Hair-Cell Regeneration: The Current State of
Therapeutic Developments and Their Potential Impact on Audiological Clinical Practice. Journal
of the American Academy of Audiology *32*, 661-669.
Hong, L., Wu, Y., Feng, J., Yu, S., Li, C., Wu, Y., Li, Z., Cao, L., Wang, F., and Zhang, Y. (2014).
Overexpression of the cell adhesion molecule claudin-9 is associated with invasion in pituitary
oncocytomas. Human Pathology *45*, 2423-2429.
Hudspeth, A.J. (2008). Making an effort to listen: mechanical amplification in the ear. Neuron
59, 530-545.
Iyer, A.A., Hosamani, I., Nguyen, J.D., Cai, T., Singh, S., Beyer, L., Zhang, H., Jen, H.-I., Yousaf, R.,
and Birol, O. (2022a). Cellular reprogramming with ATOH1, GFI1, and POU4F3 implicate
epigenetic changes and cell-cell signaling as obstacles to hair cell regeneration in mature
mammals. bioRxiv.
Iyer, A.A., Hosamani, I., Nguyen, J.D., Cai, T., Singh, S., McGovern, M.M., Beyer, L., Zhang, H.,
Jen, H.-I., and Yousaf, R. (2022b). Cellular reprogramming with ATOH1, GFI1, and POU4F3
implicate epigenetic changes and cell-cell signaling as obstacles to hair cell regeneration in
mature mammals. Elife *11*, e79712.
Izumikawa, M., Minoda, R., Kawamoto, K., Abrashkin, K.A., Swiderski, D.L., Dolan, D.F., Brough,
D.E., and Raphael, Y. (2005). Auditory hair cell replacement and hearing improvement by Atoh1
gene therapy in deaf mammals. Nat Med *11*, 271-276.

Jahan, I., Pan, N., Kersigo, J., and Fritzsch, B. (2015). Neurog1 can partially substitute for Atoh1
function in hair cell differentiation and maintenance during organ of Corti development.
Development *142*, 2810-2821.
Jeon, S.J., Fujioka, M., Kim, S.C., and Edge, A.S. (2011). Notch signaling alters sensory or
neuronal cell fate specification of inner ear stem cells. J Neurosci *31*, 8351-8358.
Kaiser, M., Ludtke, T.H., Deuper, L., Rudat, C., Christoffels, V.M., Kispert, A., and Trowe, M.O.
(2022). TBX2 specifies and maintains inner hair and supporting cell fate in the Organ of Corti.
Nat Commun *13*, 7628.
Kimitsuki, T., Nishida, M., Kawano, H., Haruta, A., Matsuda, K., and Komune, S. (2001).
Membrane current possessing the properties of a mechano-electric transducer current in inner
hair cells of guinea-pig cochlea. Brain Res *915*, 101-103.
Kitajiri, S.-i., Furuse, M., Morita, K., Saishin-Kiuchi, Y., Kido, H., Ito, J., and Tsukita, S. (2004).
Expression patterns of claudins, tight junction adhesion molecules, in the inner ear. Hearing
research *187*, 25-34.
Kitajiri, S., and Katsuno, T. (2016). Tricellular Tight Junctions in the Inner Ear. Biomed Res Int
2016, 6137541.
Kudo, T., Wangemann, P., and Marcus, D.C. (2018). Claudin expression during early postnatal
development of the murine cochlea. BMC physiology *18*, 1-8.
Lanford, P.J., Lan, Y., Jiang, R., Lindsell, C., Weinmaster, G., Gridley, T., and Kelley, M.W. (1999).
Notch signalling pathway mediates hair cell development in mammalian cochlea. Nature
genetics *21*, 289-292.
Lewis, J. (1998). Notch signalling and the control of cell fate choices in vertebrates. Semin Cell
Dev Biol *9*, 583-589.
Li, S., He, S., Lu, Y., Jia, S., and Liu, Z. (2023). Epistatic genetic interactions between Insm1 and
Ikzf2 during cochlear outer hair cell development. Cell Rep *42*, 112504.
Liu, Z., Dearman, J.A., Cox, B.C., Walters, B.J., Zhang, L., Ayrault, O., Zindy, F., Gan, L., Roussel,
M.F., and Zuo, J. (2012). Age-dependent in vivo conversion of mouse cochlear pillar and Deiters'
cells to immature hair cells by Atoh1 ectopic expression. Journal of Neuroscience *32*, 6600-6610.
Lowenheim, H., Furness, D.N., Kil, J., Zinn, C., Gultig, K., Fero, M.L., Frost, D., Gummer, A.W.,
Roberts, J.M., Rubel, E.W.*, et al.* (1999). Gene disruption of p27(Kip1) allows cell proliferation in
the postnatal and adult organ of corti. Proc Natl Acad Sci U S A *96*, 4084-4088.
Mathern, D.R., Laitman, L.E., Hovhannisyan, Z., Dunkin, D., Farsio, S., Malik, T.J., Roda, G., Chitre,
A., Iuga, A.C., Yeretssian, G.*, et al.* (2014). Mouse and human Notch-1 regulate mucosal immune
responses. Mucosal Immunol *7*, 995-1005.
Mizutari, K., Fujioka, M., Hosoya, M., Bramhall, N., Okano, H.J., Okano, H., and Edge, A.S. (2013).
Notch inhibition induces cochlear hair cell regeneration and recovery of hearing after acoustic
trauma. Neuron *77*, 58-69.
Montalbano, G., Olivotto, I., Germanà, A., and Randazzo, B. (2021). Evaluation of the hair cell
regeneration and claudin b and phoenix gene expression during exposure to low concentrations
of cadmium and zinc in early developing zebrafish larvae. Comparative Biochemistry and
Physiology Part C: Toxicology & Pharmacology *248*, 109116.
Montgomery, S.C., and Cox, B.C. (2016). Whole mount dissection and immunofluorescence of
the adult mouse cochlea. JoVE (Journal of Visualized Experiments), e53561.

Muniak, M.A., Rivas, A., Montey, K.L., May, B.J., Francis, H.W., and Ryugo, D.K. (2013). 3D model of frequency representation in the cochlear nucleus of the CBA/J mouse. Journal of Comparative Neurology *521*, 1510-1532.

Nakano, Y., Kim, S.H., Kim, H.-M., Sanneman, J.D., Zhang, Y., Smith, R.J., Marcus, D.C., Wangemann, P., Nessler, R.A., and Bánfi, B. (2009). A claudin-9–based ion permeability barrier is essential for hearing. PLoS genetics *5*, e1000610.

Neitzel, R.L., and Fligor, B.J. (2019). Risk of noise-induced hearing loss due to recreational sound: Review and recommendations. The Journal of the Acoustical Society of America *146*, 3911-3921.

Pangrsic, T., Lasarow, L., Reuter, K., Takago, H., Schwander, M., Riedel, D., Frank, T., Tarantino, L.M., Bailey, J.S., Strenzke, N., *et al.* (2010). Hearing requires otoferlin-dependent efficient replenishment of synaptic vesicles in hair cells. Nat Neurosci *13*, 869-876.

Peguero, B., and Tempel, B.L. (2015). A chromosome 17 locus engenders frequency-specific non-progressive hearing loss that contributes to age-related hearing loss in mice. Journal of the Association for Research in Otolaryngology *16*, 459-471.

Petrovic, J., Formosa-Jordan, P., Luna-Escalante, J.C., Abelló, G., Ibañes, M., Neves, J., and Giraldez, F. (2014). Ligand-dependent Notch signaling strength orchestrates lateral induction and lateral inhibition in the developing inner ear. Development *141*, 2313-2324.

Roux, I., Safieddine, S., Nouvian, R., Grati, M., Simmler, M.C., Bahloul, A., Perfettini, I., Le Gall, M., Rostaing, P., Hamard, G., *et al.* (2006). Otoferlin, defective in a human deafness form, is essential for exocytosis at the auditory ribbon synapse. Cell *127*, 277-289.

Rybak, L., and Ramkumar, V. (2007). Ototoxicity. Kidney international *72*, 931-935.

Schwander, M., Sczaniecka, A., Grillet, N., Bailey, J.S., Avenarius, M., Najmabadi, H., Steffy, B.M., Federe, G.C., Lagler, E.A., and Banan, R. (2007). A forward genetics screen in mice identifies recessive deafness traits and reveals that pejvakin is essential for outer hair cell function. Journal of Neuroscience *27*, 2163-2175.

Senften, M., Schwander, M., Kazmierczak, P., Lillo, C., Shin, J.-B., Hasson, T., Géléoc, G.S., Gillespie, P.G., Williams, D., and Holt, J.R. (2006). Physical and functional interaction between protocadherin 15 and myosin VIIa in mechanosensory hair cells. Journal of Neuroscience *26*, 2060-2071.

Sergeyenko, Y., Lall, K., Liberman, M.C., and Kujawa, S.G. (2013). Age-related cochlear synaptopathy: an early-onset contributor to auditory functional decline. J Neurosci *33*, 13686-13694.

Stone, J.S., and Rubel, E.W. (1999). Delta1 expression during avian hair cell regeneration. Development *126*, 961-973.

Strenzke, N., Chakrabarti, R., Al-Moyed, H., Muller, A., Hoch, G., Pangrsic, T., Yamanbaeva, G., Lenz, C., Pan, K.T., Auge, E., *et al.* (2016). Hair cell synaptic dysfunction, auditory fatigue and thermal sensitivity in otoferlin Ile515Thr mutants. EMBO J *35*, 2519-2535.

Sullivan, C.B., Al-Qurayshi, Z., Zhu, V., Liu, A., Dunn, C., Gantz, B.J., and Hansen, M.R. (2020). Long-term audiologic outcomes after cochlear implantation for single-sided deafness. Laryngoscope *130*, 1805-1811.

Sun, S., Babola, T., Pregernig, G., So, K.S., Nguyen, M., Su, S.M., Palermo, A.T., Bergles, D.E., Burns, J.C., and Muller, U. (2018). Hair Cell Mechanotransduction Regulates Spontaneous Activity and Spiral Ganglion Subtype Specification in the Auditory System. Cell *174*, 1247-1263 e1215.

Tang, P.C., Chen, L., Singh, S., Groves, A.K., Koehler, K.R., Liu, X.Z., and Nelson, R.F. (2023). Early
Wnt Signaling Activation Promotes Inner Ear Differentiation via Cell Caudalization in Mouse
Stem Cell-Derived Organoids. Stem Cells *41*, 26-38.
Von Bekesy, G. (1952). Resting potentials inside the cochlear partition of the guinea pig. Nature
169, 241-242.
Waqas, M., Zhang, S., He, Z., Tang, M., and Chai, R. (2016). Role of Wnt and Notch signaling in
regulating hair cell regeneration in the cochlea. Frontiers of medicine *10*, 237-249.
White, P.M., Doetzlhofer, A., Lee, Y.S., Groves, A.K., and Segil, N. (2006). Mammalian cochlear
supporting cells can divide and trans-differentiate into hair cells. Nature *441*, 984-987.
Willott, J.F., and Erway, L.C. (1998). Genetics of age-related hearing loss in mice. IV. Cochlear
pathology and hearing loss in 25 BXD recombinant inbred mouse strains. Hear Res *119*, 27-36.
Wilson, B.S., and Tucci, D.L. (2021). Addressing the global burden of hearing loss. The Lancet
397, 945-947.
Wu, P.-z., O'Malley, J.T., de Gruttola, V., and Liberman, M.C. (2020). Age-related hearing loss is
dominated by damage to inner ear sensory cells, not the cellular battery that powers them.
Journal of Neuroscience *40*, 6357-6366.
Yang, J., Bouvron, S., Lv, P., Chi, F., and Yamoah, E.N. (2012). Functional features of trans-
differentiated hair cells mediated by Atoh1 reveals a primordial mechanism. J Neurosci *32*,
3712-3725.
Zavala-Zendejas, V.E., Torres-Martinez, A.C., Salas-Morales, B., Fortoul, T.I., Montaño, L.F., and
Rendon-Huerta, E.P. (2011). Claudin-6, 7, or 9 overexpression in the human gastric
adenocarcinoma cell line AGS increases its invasiveness, migration, and proliferation rate.
Cancer Investigation *29*, 1-11.
Zheng, J.L., and Gao, W.-Q. (2000). Overexpression of Math1 induces robust production of extra
hair cells in postnatal rat inner ears. Nature Neuroscience *3*, 580-586.
Zine, A., Messat, Y., and Fritzsch, B. (2021). A human induced pluripotent stem cell-based
modular platform to challenge sensorineural hearing loss. Stem Cells *39*, 697-706.
Zine, A., Van De Water, T.R., and de Ribaupierre, F. (2000). Notch signaling regulates the
pattern of auditory hair cell differentiation in mammals. Development *127*, 3373-3383.

Figure legend.

Figure 1. The cochlear partition and construct of doxycycline-tet-OFF-*Cldn9* mouse line.

A. Schematic diagram of the radial section of the cochlear partition. The organ of Corti (OC) is seated on the basilar membrane (BM). The inner and outer pillar cells (IPC and OPC) are separated from the inner and outer hair cells (IHC, OHCs) by tight junction proteins (TJPs). The red lines indicate the expression outline of Cldn9. TM: tectorial membrane; IHC: inner hair cell; OHCs: outer hair cells; IBC: inner border cell; IPhC: inner phalangeal cell; IPC: inner pillar cell; DCs: Deiters cells; HCs: Hensen's cells. **B.** The immunostaining of *Cldn9* in the 8-wk old WT (*Cldn9$^{+/+}$*) mouse cochlea showing IHC stained with myosin7a (cyan) and claudin9 (red)antibodies, phalloidin, actin (green) for stereocilia bundles and Dapi (blue) for nuclear stain. **C**. Surface view of the cochlea **B**. Scale = 10 μm. **D.** The construct of Doxycycline-tet-OFF-*Cldn9* in the ES cells. Generation of a genetically modified mouse line in which the level of *Cldn9* gene expression can be regulated by doxycycline: A tetracycline-based genetic switch (tTA cassette) consists of three main modules, the tetracycline-controlled transcriptional activator (tTA), the neomycin resistance gene flanked by LoxP sites, and the final module that contains six copies of the tet operator (tetO) fused to the minimal CMV promoter. The tTA cassette was inserted at the -110-nucleotide position upstream of the translational start of *Cldn9* to generate the targeting vector. The targeting vector was electroporated into B6 mouse embryonic stem cells. Following the selection in G418, DNA samples from the neomycin-

resistant ES cell clones were prepared for short-arm PCR/sequencing analysis and Southern

blot analysis to confirm the insertion of the tTA cassette in the ES cells. The genetically modified

ES cells containing one copy of the tTA cassette were injected into the healthy albino B6

blastocysts, and the injected blastocysts were transplanted into the uterus of an albino B6

mouse to generate the chimeric mouse. The chimeric mouse was then bred with albino B6 mice

to produce the F1 heterozygous mouse, and the germline transmission was confirmed by tail

DNA genotyping. The deletion of the selection marker in the tTA cassette by crossing the F1

mouse with the embryonic Cre line (B6.129S4-Meox2 tm1(cre)Sor /J). **E**. Genomic DNA

samples were prepared from neomycin-resistant ES cell clones and digested with BstAPI or

BamHI. The tTA cassette insertion was confirmed by the Southern blot analysis. **F**. Mouse tail

DNA genotyping was carried out using primers P1/P3 for WT and P2/P3 for targeted alleles

showing *Cldn9$^{T/T}$*, *Cldn9$^{T/+}$* and *Cldn9$^{+/+}$* samples.

Figure 2. *Cldn9* knockdown-induced supernumerary cochlear IHCs. A. Radial section of

the cochlea partition of 8-wk-old WT (*Cldn9$^{+/+}$*) (left Panel) and *Cldn9*+/T (right Panel) mice.

Myosin 7a (cyan) stained IHCs and phalloidin (green) labeled actin stereocilia bundles and other

cellular structures, and Tuj1 (red) stained neurons. The yellow asterisks (*) denote ectopic-IHC.

Scale bar = 5 μm. **B.** 3-D rendition of cochlear surface preparation (aged 8-wks old), showing a

single row of IHCs and three rows of OHCs in the WT (top Panel: *Cldn9$^{+/+}$*) and for the *Cldn9$^{+/T}$*

(lower Panel). The ectopic IHCs are marked with (*) Scale bar = 10 μm. **C.** Mid-modiolar section

of the 8-week-old cochlea of *Cldn9$^{+/+}$* (top) and *Cldn9$^{+/T}$* (lower) panels. The red arrows show

one and two IHC bundles in the *Cldn9$^{+/+}$* and *Cldn9$^{+/T}$* cochlear sections, respectively. SG =

spiral ganglion, TM= tympanic membrane IHC = inner hair cell, OHCs = outer hair cells Scale

bar = 10 μm. **D.** Surface preparation of cochlea of *Cldn9$^{+/T}$*, showing two-rows of IHCs (the 2nd-

row IHCs are marked with (*) and higher magnification of a tilted section revealing the nuclei

(Dapi stain in blue, and arrow points to the nuclei of the 2nd IHC row). The dotted yellow line

shows the hair bundles of the 1st row, and the red lines denote the orientations of the hair

bundles. Scale bar = 10 μm. **E.** Scanning electron photomicrographs of the apical (Apex),

middle (MT), and basal (Base) cochlear turn of 6-week-old *Cldn9*$^{+/+}$ and *Cldn9*$^{+/T}$ mice. The

ectopic IHCs are marked with (*). **F.** Summary data of Quantification of IHCs count at 8- and 32-

kHz placed cochlear map at postnatal days (P) 2, 7, 14, 21 in *Cldn9*$^{+/+}$ compared with *Cldn9*$^{+/T}$

mice. Each data point is the means of 4-blind counts. At 8 kHz cochlear placed-map, the *p*

values were P2 = 7.4×10^{-3}, P7 = 3.0×10^{-4}, P14 = 2×10^{-3}, and P21 = 1.5×10^{-4}. At 32 kHz

cochlear placed-map, the *p* values were P2 = 4.0×10^{-3}, P7 = 1.0×10^{-3}, P14 = 8.5×10^{-6}, and P21

= 1.7×10^{-2}.

Figure 3. *Cldn9* knockdown-induced supernumerary IHCs remain intact in older mice. A.

Surface cochlear preparation of a 15-months (P450) old *Cldn9*$^{+/+}$ mouse shows IHCs and OHCs

labeled with Myosin7a (Myo7a) antibody (green) and actin with phalloidin (cyan) and

presynaptic maker CtBP2 antibody (red), which is nuclear-positive. The white dash line marks

the single row of IHCs. Note that the missing OHCs are marked with white asterisks at 15

months. **B.** Similar preparation as in **A**, from a 15-month-old *Clnd9*$^{+/T}$ mouse. Viable ectopic IHC

can be seen with yellow arrows pointing to the hair bundles and nuclei, stained with CtBP2

antibody (*). **C.** Summary data of Quantification of IHCs count at 8- and 32-kHz placed cochlear

map at postnatal days (P) 21, 30, 300, and 450 in *Cldn9*$^{+/+}$ compared with *Cldn9*$^{+/T}$ mice. Each

data point is the mean of 3-blind counts. At 8 kHz cochlear placed-map, the *p* values were P30

= 2.9×10^{-7}, P300 = 3.7×10^{-6}, and P450 = 1.4×10^{-7}. At 32 kHz cochlear placed-map, the *p* values

were P30 = 1.1×10^{-2}, P300 = 4.6×10^{-4}, and P450 = 6.7×10^{-5}.

Figure 4. Auditory brainstem recordings (ABR) and distortion product otoacoustic

emissions (DPOAE) assessment and mechanoelectrical transducer channel-FM1-43

permeation and currents elicited from original (O; IHC$_O$) and ectopic (E; IHC$_E$) IHC stereocilia bundles. A, ABR thresholds from the $Cldn9^{+/+}$ (open symbols, das lines) and $Cldn9^{+/T}$ (solid symbols, solid lines) monitored from 2-8 months. The sound pressure levels (SPL) in dB of pip tones (in kHz) 4, 8, 16, and 32 and broadband clicks delivered to the ear are indicated. ABR thresholds for $Cldn9^{+/+}$ (n = 12) and $Cldn9^{+/T}$ (n = 17) mice. On average, the $Cldn9^{+/T}$ had a 5-15 dB elevated threshold compared to $Cldn9^{+/+}$ mice. Within each group of mice, there was no significant shift in ABR threshold from 2-8 months of monitoring. The data means ± SD. **B,** Mean DPOAE threshold for 2-8 months ($Cldn9^{+/+}$ group: n=9; $Cldn9^{+/T}$ group: n = 14) was tested, measuring the 2f1 - f2 DPOAE over a geometric-mean frequency range at 8, 16, and 32 kHz. The lines and symbols are the same as the labels in Fig. 4A. On average, the $Cldn9^{+/T}$ had a 5-7 dB elevated threshold compared to $Cldn9^{+/+}$ mice. **C,** Fluorescent images of FM1-43 were taken at different time points at two focal planes: the base of IHC bundle (BHB) and the below cuticular plate (BHC) levels of postnatal day (P) 14 $Cldn9^{+/T}$ cochlea. The ectopic IHC bundles are marked with (*). **D.** Time 0 indicates the onset of dye application (10 mM for the 5-sec duration). Consecutive images at BHB and BHC were taken at 5-s intervals. Dye loaded into this region remained sustained. The dye enters the apical aspects of the cell before being visualized at the basal pole. (Scale bar, 10 μm). The change in fluorescence at focal levels as a function of time (adjusted for the interval between frame capture at each level). Densitometric data of mean pixel intensity were measured in arbitrary grayscale units. Summary data from IHC$_O$ and IHC$_E$ (n = 27 from 3 mice). Frames were taken from 3-4 kHz placed map of the mouse cochlea. The change in fluorescence was fitted with an exponential function, and the time constants (t, in s) of FM1-43 dye loading in IHC$_O$ (in black circles) at BHB and BHC levels,19.4±1.1 (n = 27) and 29.0±3.8 (n = 27). Similar analyses for IHC$_E$ (in red circles) at BHB and BHC levels were 23.1±4.5 (n = 27) and 47.0±8.4 (n = 27). **E,** Typical MET current traces from IHC$_O$ and IHC$_E$ elicited sinusoidal hair bundle deflection at 40 Hz. **F.** MET current elicited a

series of ~200-ms hair bundle displacements, using fluid-jet deflection towards the taller

stereocilia. Hair cells were held at -80 mV. All recordings were made from apical IHCs. Bundle

deflection was elicited with 0.1-0.9 V pressure clamps in 0.1-V steps. Estimates for the exact

bundle displacement were not determined. For clarity, a few traces were omitted. **G,** Summary

of group data of the maximum IHC MET current measured from IHC_O, 335±181 pA (n = 18) and

IHC_E, 268±192 (n = 14) (mean ± SD) **H,** Normalized displacement response relationships fitted

with a two-state Boltzmann function. The half-maximum displacement voltage (in V) for IHC_O

and IHC_E were 0.20±0.02 and 0.30±0.01 (n= 7).

Figure 5.

Pre- and post-synaptic features of original and ectopic IHCs in *Cldn9$^{+/+}$* and *Cldn9$^{+/T}$*

mice. A, The innervation of calretinin (Calb2)-positive afferent neurons (red) of an original and

ectopic IHCs (marked *, and **) from a 5-wks-old *Cldn9$^{+/T}$* mouse cochlea. Dapi (blue) labeled

the nuclei. Arrows point to the nerve terminals. Scale bar = 5 μm. **B,** CtBP2 labeled (red) pre-

synapse and Homer1 labeled (green) post-synapse in the cochlear IHCs in the *Cldn9$^{+/+}$* and

Cldn9$^{+/T}$ mice. The original IHC (IHC_O) and ectopic IHC (IHC_E) outlines are marked with purple

and cyan dashed lines. The two right panels show a 3-D rendition of the pre-and post-synaptic

markers in *Cldn9$^{+/+}$* and *Cldn9$^{+/T}$* cochleae. Scale bar = 5 μm. **C,** Quantification of the number of

pre- (red) and post-synaptic (green) immunopuncta at three tonotopic cochlear locations: apex

(3-5 kHz), middle (12-16 kHz), and base (32-40 kHz) from 8-wk-old *Cldn9$^{+/+}$* (open circles) and

Cldn9$^{+/T}$ (solid circles) mice. Colocalized pre- and post-synaptic marker counts per IHC are

shown in yellow. Values (mean ± SD) are illustrated ($p <$ 0.05, 0.01, 0.001 = *, **, ***).

Comparing pre-, post-synaptic and colocalized markers for apical IHCs, between *Cldn9$^{+/+}$* (n =

56 from 3 mice) and *Cldn9$^{+/T}$* (n = 117 from 4 mice) the *p* values were pre- p = 2.2X10^{-42}, post-

=3.5X10^{-30}, co-expression, 4.3X10^{-34}. For mid-cochlea IHCs, between *Cldn9$^{+/+}$* (n = 29 from 3

mice) and *Cldn9$^{+/T}$* (n = 79 from 4 mice) the *p* values were pre- p = 1.7X10^{-19}, post- =1.3X10^{-14},

co-expression, $2.5\text{X}10^{-18}$. For basal-cochlea IHCs, between $Cldn9^{+/+}$ (n = 97 from 3 mice) and

$Cldn9^{+/T}$ (n = 94 from 4 mice) the p values were pre- $p = 3.5\text{X}10^{-27}$, post- $=8.2\text{X}10^{-27}$, co-

expression, $2.9\text{X}10^{-26}$. **D,** analysis of synaptic features among IHC_O and IHC_E in $Cldn9^{+/T}$

cochlea. At the cochlea apex, comparing pre- post- and colocalized markers between IHC_O (n =

76 from 3 mice) and IHC_E (n = 41 from 3 mice), the p values were pre- $p = 7.0\text{X}10^{-2}$, post- =

$9.0\text{X}10^{-3}$, co-expression = $7.0\text{X}10^{-2}$. For mid-cochlea comparing pre-, post- and colocalized

markers between IHC_O (n = 48 from 3 mice) and IHC_E (n = 30 from 3 mice) the p values were

pre- $p = 1.4\text{X}10^{-4}$, post- = $8.9\text{X}10^{-5}$, co-expression, $1.7\text{X}10^{-5}$. For basal-cochlea comparing pre-,

post- and colocalized markers between IHC_O (n = 65 from 3 mice) and IHC_E (n =22 from 3 mice)

the p values were pre- $p = 9.0\text{X}10^{-2}$, post- = $1.0\text{X}10^{-2}$, co-expression, $5.0\text{X}10^{-2}$.

Figure 6. Postnatal induction of ectopic IHCs with the Cldn9 *shRNA* knockdown

A, Ectopic IHCs observed in P21 cochlea after *Cldn9 shRNA* injection at P2. Downregulation of

Cldn9 levels by ~20-fold compared to scrambled shRNA is shown in S7A, and a reduced

protein expression (**Fig. 6A**) is shown in the 3[rd] panel (*Cldn9*, red). GFP (green) expression is a

marker for adeno (AD)-virus-transduced cells. The right panel shows the merged image with two

rows of IHCs stained with HC marker Myo7a (cyan) and Ad-GFP-mCldn9 shRNA (green)

expression in the adjacent row of supporting cells. The Ad-GFP-mCldn9 shRNA (green) is

expressed in HCs. **B,** Examples of phalloidin stained whole-mount organ of Corti samples from

cochleae of the wild-type mice injected at P2, P4, P7, and P14 with Cldn9 shRNA. The tissues

were harvested 20 days after viral injection. Sections at or near the cuticular-plate level show

actin labeled with phalloidin (red). Several ectopic IHCs are marked with asterisks in yellow (*).

Where nuclei were labeled with Dapi (blue), we marked the ectopic IHC nuclei with a white

arrow. **C,** Scanning electron photomicrographs of the apical (apex), middle (mid), and basal

(base) cochlear turn of 5-6-wks-old mice after scrambled shRNA and *Cldn9* shRNA injection at

P2, showing original and ectopic IHCs (marked with white arrows). Images were captured at the

cochlea's apical, middle, and basal contour. **D,** Summary data of quantification of ectopic IHCs

count at cochlear apex, middle, and base for samples where *Cldn9* shRNA was injected at P2,

P4, P7, and P14. Each data point (n=11) is the mean of 3-blind counts from 3 mice. Ectopic

IHCs at-P2 injection (mean/100-mm±SD) for apical, middle, and basal cochlea were 7.5±4.5,

5.0±2.4, and 3.2±1.6; at-P4; 8.5±3.8, 6.1±2.5, and 3.2±1.6; at-P7; 2.8±2.4, 2.4±1.7, and 1.1±1.0;

at-P14, 0.2±0.5, 0.1±0.3, and 0.0±0.0. Data are plotted to show individual replicates (n=11;

mean□±□SD). There were significant differences at *p*<0.05 level for comparison between P2-7

and P14 F (11,120) = (17) *p = 4.2X10^{20}. Post hoc* comparisons using the Tukey HSD test

indicate that post-shRNA injection-induced ectopic IHCs are significantly different at P2 apex vs

P14 apex (*p*=2.4X10^{-8}); P2 mid vs. P14 mid (*p*=1.3X10^{-4}); and P2 base vs. P14 base (*p*=7.0X10^{-2}).

Figure 7. Study design and summary of findings.

A, The study design schematic included the doxycycline-treated *Cldn9$^{+/T}$* and *Cldn9$^{+/+}$* mice and

postnatal mice transduced with *Cldn9* shRNA. Whereas ectopic IHCs were generated by inner

ear injection of Cldn9 shRNA from P2-7, we did not detect significant ectopic IHCs at P14. **B,**

The contact lateral inhibition in the late-differentiation stage of IHCs and SCs. The solid red line

showed the intact Cldn9 sealing between HCs and SCs, while the dashed red line showed the

downregulated *Cldn9* level between cells.

Supplement S1

Schematic diagram of the construct used to regulate translation of CLDN9. We will use the tTA ("tet-

off") cassette developed by Seeburg and Adelman for expressing the SK3 channel {Bond, 2000

#182} (Bond et al., 2000) and modified to include unique *Not* I and *Xho* I cloning sites at each

end of the cassette. The cassette contains three motifs: i.) tTA preceded by the tri-partite leader

sequence that confers highly efficient translational initiation followed by the SV40 polyA and

transcriptional termination sequence, ii.) bacterial neomycin (G418) resistance gene driven by

the herpes simplex virus thymidine kinase (TK) promoter and the yeast URA3 gene; these

selectable markers are flanked by loxP sites and followed by the human growth hormone (hGH)

polyA and transcriptional termination sequence, iii.) six copies of the tet operator (tetO6) fused

to the minimal cytomegalovirus (CMV) promoter. The native *Cldn9* promoter drives

the expression of tTA, which in CLDN9 from the tetO6/CMV promoter. In the presence of Dox

(crossed circles), tTA is bound and cannot bind to the tetO6/CMV promoter, thereby turning off

the transcription of CLDN9.

S2

A. Quantitative RT-PCR of *Cldn9* transcripts from cochlear tissue from six groups of animals,

including *Cldn9$^{T/T}$*, *Cldn9$^{+/T}$* with and without dox treatment compared with WT littermates with

and without dox treatment. **B.** Body weight measurements of female and male mice from dox-

treated *Cldn9$^{+/T}$* and *Cldn9$^{+/+}$* groups were recorded at 4, 6, and 8 wks old. **C.** The

immunostaining of *Cldn9* in 8-wk old WT (*Cldn9$^{+/+}$*) mouse cochlea*Cldn9* (red), IHC stained

myosin7a (cyan) and supporting cells stained Sox2 (green) and Dapi (blue) for nuclear stain.

Scale = 10 μm.

S3

Immunogold localization of Cldn9 in *Cldn9$^{+/T}$* and *Cldn9$^{+/+}$* mice. A. Cldn9 expression sites

between inner hair cells (IHCs) and supporting cells (SCs) were examined with immunogold

electron microscopy with the post-embedding technique. Secondary antibodies are conjugated

to 16-nm colloidal gold particles (arrows). Gold particles were noted at the junctions between

SC and IHCs in *Cldn9$^{+/+}$* mice cochlea (indicated). *Cldn9$^{+/T}$* mice had reduced gold particles. **B.**

Summary of gold particle counts between IHC and SC between Cldn9$^{+/+}$ and Cldn9$^{+/T}$. Mean

gold particles (mean±SD) for *Cldn9$^{+/+}$* = 25±14 (n = 41 from 3 cochleae) and for *Cldn9$^{+/T}$* = 3±2

(n = 41, from 4 cochleae), *p = 2.8X10^{-15}*.

1008

S4

The expression of cldn6 and ILDR1 in the organ of Corti. A-B, The immunostaining of Cldn6

(red) in the mouse cochlea from *Cldn9$^{+/+}$* (wildtype, WT) and C*ldn9$^{+/T}$*. The lower Panel is the

side view of the cochlear section. HCs were labeled with Myo7a (cyan) supporting cells with

Sox2 (green) and Dapi-stained (blue) nuclei. The levels of Cldn6 were reduced in the *Cldn9$^{+/T}$*

cochlea. **C-D**, The immunostaining of ILDR1 (red) in the mouse cochlea from *Cldn9$^{+/+}$* and

Cldn9$^{+/T}$. The lower Panel is the side view of the cochlear section. HCs were labeled with Myo7a

(cyan) supporting cells with Sox2 (green) and Dapi-stained (blue) nuclei. The levels of ILDR1

were increased in the *Cldn9*$^{+/T}$ cochlea. Scale bar = 10 μm.

1018

S5

The ectopic HCs can be seen along the cochlear apicobasal contour.

A, The immunostaining of Myosin 7a (red) and F-actin (green) in C*ldn9*$^{+/T}$ mouse cochlear apical

segment. **B**, Panel shows smaller segments magnified to clarify and detect multiple ectopic HCs.

HCs were labeled with Myo7a (red). Scale bar = 10 μm.

S6

The ectopic HCs stain positively to Myosin 7a but negatively to prestin antibodies

A-B, The immunostaining of prestin (red) in the mouse cochlea from *Cldn9*$^{+/+}$ (wildtype, WT)

and C*ldn9*$^{+/T}$. The lower Panel is the side view of the cochlear section. HCs were labeled

with Myo7a (cyan) phalloidin-stained actin (green) and Dapi-stained (blue) nuclei. Scale

bar = 10 μm.

1030
S7

A. Summary data of Quantification of OHCs count at ~4, 8, 16, 32, and 45-kHz placed cochlear

map at postnatal days (P14) in *Cldn9*$^{+/+}$ (WT, in black symbols) compared with *Cldn9*$^{+/T}$ (in red

symbols) mice. Each data point is the mean of 4-blind counts. **B.** At 8 kHz cochlear placed-map,

OHC counts were changed for P14, 21, 30, 300, and 450 mice. OHC numbers remained

relatively constant (P14-30) until P300-450) when they were reduced.

1037

1038 **S8**

1039 **The ABR wave one amplitude and latency in *Cldn9$^{+/T}$* and *Cldn9$^{+/+}$* mice**. **A-D.** ABR wave 1

1040 amplitude was decreased, while the latency varied in *Cldn9$^{+/T}$* mice compared to *Cldn9$^{+/+}$* mice

1041 when measured at 8 kHz, 16 kHz, 32 kHz, and click **(E-H).** * P < 0.05, ** P < 0.01, *** P < 0.001.

1042

1043 **S9**

1044 **A. Quantitative RT-PCR of *Cldn9*** transcripts from cochlear tissue from WT un-injected,

1045 scrambled (scram), and shRNA P2- and P14-old-injected mice. **B.** Examples of SEM of the

1046 shRNA-injected cochlea (age of samples at the harvesting time (P24) for imaging and the age at

1047 shCldn9 injection P2). Yellow arrows show ectopic IHC bundles.

1048

S10

The endocochlear potential (EP) and perilymph K$^+$ in *Cldn9$^{+/+}$* and *Cldn9$^{+/T}$* mice. A.

Measurement of the EP at the basal turn of the cochlea, in 4-month-old *Cldn9$^{+/+}$* = 99±11 mV

(n=13), *Cldn9$^{+/T}$* = 83±11 mV (n=13) and at 8-month-old for *Cldn9$^{+/T}$* = 80±12 (mV) (n=11). There

were significant differences at *p*<0.05 level for comparison between *Cldn9$^{+/+}$* vs. *Cldn9$^{+/T}$*

F(2,34)=(9.3) *p* =5.9$X10^{-4}$. *Post hoc* comparisons using the Tukey HSD test indicate *Cldn9$^{+/+}$* vs.

Cldn9$^{+/T}$ at 4 months (*p*=4.0$X10^{-3}$); *Cldn9$^{+/+}$* (4 months) vs. *Cldn9$^{+/T}$* (8 months) (*p*=1.2$X10^{-3}$); are

significantly different. **B.** Recordings of perilymph K$^+$, in 4-month-old *Cldn9$^{+/+}$* = 2.2±0.5 mM

(n=14), *Cldn9$^{+/T}$* = 4.4±0.8 mM (n=15) and at 8-month-old for *Cldn9$^{+/T}$* = 4.4±1.4 (mM) (n=12).

There were significant differences at *p*<0.05 level for comparison between *Cldn9$^{+/+}$* vs. *Cldn9$^{+/T}$*

F (2,36)=(24.4) *p* =1.6$X10^{-7}$. *Post hoc* comparisons using the Tukey HSD test indicate *Cldn9$^{+/+}$*

vs. *Cldn9$^{+/T}$* at 4 months (*p*=1.1$X10^{-6}$); *Cldn9$^{+/+}$* (4 months) vs. *Cldn9$^{+/T}$* (8 months) (*p*=2.3$X10^{-6}$);

are significantly different.

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

1101 **Figure 1**

1102

1122 **Figure 2**
1123

1148 **Figure 3**
1149

1150
1151

Figure 4

1171

1172
1173

1174
1175
1176
1177

1197   **Figure 5**

Figure 6

1249 **Figure 7**
1250

1251
1252
1253

1273   **Supplement S1**
1274

1310 **S2**
1311

1312
1313

Supplement S3

1358 **Supplement S4**
1359

1380   **Supplement S5**
1381

1403   **Supplement S6**
1404

1426 **Supplement S7**
1427

1449 **Supplement S8**
1450

1481 **Supplement S9**
1482

1504 **Supplement S10**
1505