Evolution of genotypic and phenotypic diversity in multispecies biofilms

Cristina I. Amador[1], Sofia Zoe Moscovitz[1], Lorrie Maccario[1], Jakob Herschend[1 2], Isabel-Sophie Kramer[1], Hannah Jeckel[3], Vaughn Cooper[4], Knut Drescher[3] Thomas R. Neu[5], Mette Burmølle[1] and Henriette L. Røder[1 6 #]

[1]Section of Microbiology, Department of Biology, University of Copenhagen, Copenhagen, Denmark

[2]Present address: Functional Food, Novozymes A/S, 2800, Lyngby, Denmark

[3]Biozentrum, University of Basel, Switzerland

[4]University of Pittsburgh, Pittsburgh, USA

[5]Helmholtz Centre for Environmental Research, Magdeburg, Germany

[6]Pressent address: Section of Microbiology and Fermentation, Department of Food Science, University of Copenhagen, Copenhagen, Denmark

Author correspondence to h.lyng.roeder@food.ku.dk

Keywords: multispecies, bacterial interactions, biofilm, matrix, adaptive evolution, sporulation

Abstract

Interactions occurring in microbial communities can affect the fitness and adaptability of their individual members when facing changing environmental conditions. This study investigated the impact of interspecies interactions in selecting *Bacillus thuringiensis* variants emerging in biofilm and planktonic environments. During evolution experiments, a *B. thuringiensis* distinct phenotypic variant of *B. thuringiensis* frequently occurred, despite the presence of other species or culture setup. Remarkably, selection of this variant was significantly favored over its ancestor in biofilm settings and when coexisting with other species co-isolated from a wastewater facility, namely *Pseudomonas defluvii* and *Pseudomonas brenneri*. Interestingly, the evolved phenotype did not show higher biofilm productivity than its ancestor under any condition, while it was indeed reduced in mixed-species biofilms. Such observation aligned with the reduced abundance of matrix proteins TasA and SpoVG, which are major biofilm matrix components in *Bacillus* species, as revealed by matrix proteomics analysis. Furthermore, the variant showed shorter generation time and a lack of sporulation compared to its ancestor, consistent with mutations in key genes for regulating sporulation. Our results indicate that interspecies interactions within biofilms promote *B. thuringiensis* diversification and alter traits such as biofilm matrix production. Despite sporulation being a survival mechanism, this study provides evidence that sporulation does not confer a fitness advantage in *in vitro* biofilm environments, even within mixed cultures.

Introduction

When encountering new environmental challenges, bacterial populations can adapt through different mechanisms: altering their physiological state (phenotype), i.e. expressing specific traits, acquiring new advantageous mutations (*de novo* genetic variation), or utilizing the pre-existing genetic diversity within the population (standing genetic variation) [1–3]. Besides intraspecific variation, natural communities contain many species that might either constrain adaptation by competing for resources, e.g., via niche exclusion [4] or facilitate adaptation by making new niches available [5, 6]. In biofilms, microbial cells are in close proximity and can influence each other´s phenotypes and evolutionary trajectories through social interactions [7]. Spatial segregation in biofilms increases the frequency of interactions between cells of the same genotype and favors cooperative behaviors but can also support variants outcompeted in unsegregated environments [8–10] and facilitate coexisting populations [11, 12]. Several studies have reported that competitive interactions within biofilms are dominant, including competition for limited space and resources [10, 13–16]. However, there is evidence that competition can subside locally over time, often leading to stable coexistence [17]. Thus, to better understand bacterial adaptation mechanisms in natural environments, investigations of adaptation mechanisms in the presence of spatial structure and multiple species that co-occur in natural communities are crucial.

We have previously described that interspecies interactions can lead to evolved variants that display enhanced mutualism in biofilms [18–20]. The evolution of social interactions in biofilms has been examined for multiple Gram-negative bacteria, including *Pseudomonas spp*. and *Burkholderia cenocepacia* [21–23]. However, comparatively less attention has been dedicated to investigating evolutionary processes in Gram-positive bacteria under mixed-species conditions. Although some studies have explored the adaptive evolution of Gram-positive bacteria in diverse biofilm systems, their focus has primarily been on adaptive genotypes [24–26] or co-culture [20].

In this study, we specifically investigate the influence of cultivation settings and the presence of multiple species on the evolutionary pathways of *Bacillus thuringiensis*, a spore-forming bacterium with entomopathogenic properties [27]. *B. thuringiensis* was cultivated alone or in combination with *Pseudomonas defluvii* and/or *Pseudomonas brenneri,* co-isolated strains with *B. thuringiensis* from a wastewater facility. In contrast to other studies where cells needed to

disperse from biofilms and recolonize new surfaces during the short-term evolution assays, our

objective was to identify cells that remained attached within biofilms [20, 28]. We observed that a

phenotypic variant of *B. thuringiensis*, that showed reduced Congo red binding, emerged with

higher frequency than its ancestor. This phenotype was linked to mutations in sporulation genes

and, consequently, lack of sporulation. The evolved phenotype invaded and displaced the ancestral

strain under all tested conditions, but cocultivation with other species and biofilm setup enhanced

selection.

Results

Regular emergence of a *B. thuringiensis* morphotype variant after short-term evolution assay

We investigated whether the presence of other species affected selection of new phenotypic variants of *B. thuringiensis* (BT) in biofilm or planktonic conditions. The biofilm condition required cells to attach to polycarbonate slides for 24 hours of growth in either monoculture, co-culture or multispecies culture, followed by transferring the polycarbonate slide to new media for 8 consecutive cycles. At the end of the experiment, cultures were plated and counted based on distinct colony morphology (Figure 1a).

After 8 days, in all replicate biofilm cultivation experiments, we detected a particular *B. thuringiensis* (BT) phenotypic variant, which we termed "light variant" as it displayed reduced Congo red binding compared to the ancestor (Figure 1b). The population size of BT did not differ among treatments despite the culture condition or composition (mono-culture, two-species culture, or multispecies-culture, Figure 1c). However, the BT light variant independently emerged in each short-term biofilm evolution experiment and displayed significantly higher CFU counts than the ancestral BT strain in all conditions (Figure 1c).

Analogous 8-day evolution experiments in liquid culture showed no significant difference in CFU counts between the ancestral BT and BT light variant in monoculture conditions. However, in the presence of either *P. brenneri* or *P. defluvii* the BT light variant achieved higher CFU level than the ancestral BT strain, though no significant within combination (Figure 1c).

CFU counts were also compared using phenotype as a main factor (ANOVA (CFU~Phenotype), regardless of the species combination. Detection of the variant phenotype was significantly increased compared to its ancestor regardless of the settings ($p_{Biofilm} = 0$; $p_{Plank} = 8.82$ x 10^{-5}, Figure 1c). Together, the results of the short-term evolution experiments in biofilm and planktonic conditions indicate that the BT light variant has a higher fitness in biofilm conditions and in the presence of *P. brenneri* and *P. defluvii*, compared with the ancestral wild-type BT strain.

Figure 1. Short-term evolution assay of *Bacillus thuringiensis* in the presence/ absence of *Pseudomonas* species. (a) Planktonic/ Biofilm transfer model procedure utilized for the evolution experiment. *B. thuringiensis* (BT) was cultured either individually or together with either or both *P. brenneri* (PB) and *P. defluvii* (PL) in TSB medium in planktonic or biofilm settings and allowed to grow for 24 hours. Then, a polycarbonate chip with attached biofilm was washed thrice and moved to a new well with fresh medium. This was done every day for eight consecutive days, when the biofilms and planktonic samples were plated out and CFU counted. **(b)** Wild-type and variant phenotypes differ in colony morphology (left) and streaks (center) when cultivated on TSA supplemented with 40 µg/mL Congo red and 20 µg/mL Coomassie brilliant blue (G). The rightmost panel shows wild type and variant sedimentation phenotypes when the strains are cultivated in liquid TSB. In TSB, sedimentation can be observed only for the wild-type strain. **(c)** *B. thuringiensis* cell counts (CFU/mL) of the variant (red) and wild-type (blue) phenotypes in different strain combinations in biofilm (left panel) or planktonic (right panel) setups. Horizontal lines denote the mean value for each dataset while the box width indicates the number of replicates (n) and dots indicate outlier data points. Dissimilar letters indicate significance per facet (Tukey post-host test, $p < 0.05$).

The selected *B. thuringiensis* variant displayed distinct phenotypic traits than its ancestor

We isolated 150 BT light variants from different experiments, and all showed the same phenotypic traits. The ancestral BT and the BT light variant displayed distinct colony morphology but also showed a visibly different phenotype during growth in liquid cultures: the ancestral wild-type BT cells sink to the bottom of the culture in non-shaking conditions, whereas the BT light variant remained in suspension (Figure 1b, right panel).

Besides these phenotypic traits, the BT light variant failed to produce spores compared with the ancestral BT strain, as evidence by optical microscopy (not shown). Given that the visible phenotypic traits of all 150 isolated light variants were identical, this was tested on four of the selected isolated light variants to test their ability to form spores: V1, V2, V3 and V6. V1 and V2 were isolated from co-cultures with *P. brenneri,* while V3 and V6 were isolated from BT monocultures (Table S1). The spore formation ability of BT strains was quantified in planktonic settings over four days and compared to control conditions where only vegetative cells would be able to form colonies. These assays showed that the BT light variants completely lacked spore formation, whereas the ancestral BT strain was able to produce spores despite the incubation time (Figure 2a). Next, we investigated the effect of other species on spore formation in the ancestor. The ability to form spores was assessed in mono-, two-species-, or multispecies biofilm conditions for 24-hour biofilms, however no significant difference was observed between the tested combinations (Figure 2b).

As the variants failed to produce spores, we hypothesized that this might impact their growth rate since sporulation is an energetically costly process for the cell. Thus, growth curves were performed for the BT wild-type and the BT light variants, and different growth parameters were analyzed. We found that the four BT light variants consistently showed a significantly faster doubling time (t_{gen}) and growth rate (r) compared to their ancestor (Table S2). Consequently, the time required to reach half of the maximum culture yield was significantly lower (t_{mid} or ½ k). This suggests that the BT light variants may have a growth advantage compared to wild-type BT when competing for resources with the *Pseudomonas* species, which could lead to the observed higher CFU counts in mixed-species cultures in our short-term evolution experiments in biofilms and in planktonic cultures.

140
Figure 2. Ability of spore formation for *B. thuringiensis* wild-type or variant phenotypes. *B.*
thuringiensis: BT; *P. brenneri*: PB; *P. defluvii* (PL) **(a)** Spore forming ability of BT wild-type and variants
in planktonic monocultures after 1, 2, 3 or 4 days of incubation, expressed in the same way as a). Box plots
show 5 biological replicates, represented as individual points. Outliers are depicted as bigger black points.
Dissimilar letters in black indicate significance between strains, or facets ($p < 0.05$, Tukey post-hoc test),
while turquoise dissimilar letters denote significant comparisons only within BT (heated vs. non-heated).
(b) BT wild-type spore formation ability in biofilm settings in different strain combinations, after one day
of incubation. Box plots of BT CFU counts in control conditions (vegetative cells, non-heated, blue boxes)
and after boiling (germinating spores, heated, white boxes), based on ≥10 biological replicates indicated as
independent points, and 3 technical replicates. Dissimilar letters $p < 0.05$ (Tukey post-hoc test).

Nutrient reduction selects for higher *B. thuringiensis* diversification in mixed-species biofilms

The presence of other species selected for the BT light variant with higher frequency in full-strength TSB and we thus hypothesized that competition for resources may be a driver of diversification. We reproduced the evolution experiment in a biofilm setting but lowered the nutrients 5-fold (20% TSB, Figure S1a). CFU counts of BT and variant significantly decreased in 20 % TSB, however the variant exhibited a more pronounced reduction in these conditions (Abundance TSB100 *vs.* TSB20, variant = 13.7-fold, $p = 0$; wt = 2.4-fold, $p = 0.0428$). Unlike 100 % TSB, the wild-type was significantly more abundant in monoculture, suggesting a fitter wild-type phenotype under reduced nutrient availability in the absence of competing species (Abundance variant vs. wt) = -11-fold, $p = 8.47 \times 10^{-5}$). However, the BT light variant still emerged with higher abundance than the wild-type phenotype when either or both *Pseudomonas* species were present, although not significant (Abundance variant-wt 20 %: 2.47-fold, $p = 0.0614$). Interestingly, comparison of mono-species vs. mixed species within phenotype evidenced significantly enhanced selection of the variant in the presence of either or both *Pseudomonas*, as opposed to significant counter selection of the wild-type phenotype (mixed species vs. monospecies: variant: 9.7 to 19.1-fold; wt: -3.3 to -14.6-fold). These results indicate that the BT light variant has a growth advantage in the presence of other species, even when the nutrients have been reduced 5-fold.

For the CFU counts of *Pseudomonas*, different trends were observed for both strains depending on the media used (Figure S1b-c). PL was negatively affected by the presence of only BT or in the presence of BT and PB in 100 % TSB, but PL abundance did not change significantly in 20 % TSB (Figure S1b; $p_{20-100} = 0.8292$). On the contrary, PB CFU levels remained stable in full-strength TSB, but decreased significantly in 20 % TSB (Figure S1c; $p_{20-100} = 0.0083$).

We conclude that, during the evolution experiment, *P. brenneri* abundance significantly decreased in 20 % TSB. Since *P. brenneri* enhanced the diversification of *B. thuringiensis* in full-strength TSB, we hypothesize that a lower abundance of *P. brenneri* in 20 % TSB may account for the reduced selection of *B. thuringiensis* variants.

*The **B**. thuringiensis variant produced less biofilm matrix than its ancestor in mixed species*
biofilms

An eight-day evolution assay selected for the variant phenotype despite the nutrient
concentration and setup (planktonic or biofilm), but its frequency was enhanced in biofilm
conditions. In this evolution experiment, our focus was on promoting the retention of cells within
the biofilm rather than their dispersal. Consequently, we conducted an investigation to determine
if the evolved phenotype exhibited enhanced biofilm formation. We quantified total biofilm
formation (cells and matrix) in PC slides after 24 hours using a crystal violet assay but also the
contribution of each strain by CFU counts for both the BT wild-type and V1 variant (Figure 3).

Figure 3. Biofilm formation of *Bacillus thuringiensis* wild-type and V1 variant in mono- or mixed-
species cultures in TSB. BT: B. *thuringiensis*; V1: V1 variant; PL: *P. defluvii*; PB: *P. brenneri*. Box plot
height shows data variance and the mean value is indicated by a horizontal black line. **(a)** Total biofilm
measured as crystal violet stained biomass (A_{590}) after 24 hours. Data is based on at least 4 biological
replicates and 6 technical replicates. **(b)** Biofilm measured as CFU counts per mL after 24-hour incubation.
Monospecies: cell counts for each species in monoculture. Wild-type mix: BT counts in dual- or
multispecies mixtures. V1 mix: V1 counts in dual- or multispecies mixtures. Data is based on at least 4

biological replicates, indicated as black dots, and 9 technical replicates. Dissimilar letters indicate significant differences, with p< 0.05 (Tukey post-hoc test).

In monoculture, biofilm formation (A_{590}) was similar for BT wild-type and variant, but significantly induced in co-culture or mixed-species cultures despite the BT phenotype (Tukey HSD, p < 0.0001) (Figure 3a). Interestingly, species combinations containing the variant phenotype showed significantly decreased biofilm productivity compared to its ancestor ($CV_{wt-variant}$ = 1.8-fold, $p_{(wt-variant)}$ = 0.0071; ANOVA(CV~Phenotype)). However, no significant difference was found when comparing wild-type and variant within each culture condition (coculture PL, coculture PB or multispecies). Similar results were observed for V2, V3 and V6 variants, indicating that the source of isolation did not alter their biofilm phenotype (Figure S2a).

CFU counts showed no significant difference between wild-type and variant or culture condition (Figure 3b; $p_{phenotype}$ = 0.563; $p_{culture}$ = 0.186), two-way ANOVA). Wild-type and variant cell counts increased in combination with PB compared to monocultures, although not significantly. Comparisons of BT wild-type and variant across all culture conditions (t-test, factor = BT phenotype) resulted in significantly reduced biofilm biomass but not cell counts ($p_{CV\_wt-variant}$ = 0.0071, $p_{CFU\_PL\ wt-variant}$ = 0.204, $p_{CFU\_PB\ wt-variant}$ = 0.9991, pointing towards differential matrix production/induction. Similar results were observed for the other variants despite the medium strength used (Figure S2b and S2c, 100 % TSB and 20 % TSB, respectively). Furthermore, all variants consistently replicated the results of the evolution experiment in 20 % TSB, with reduced abundance compared to the wild-type in monoculture conditions ($p_{wt-variant}$ = 4.97 x 10^{-7} (t-test). Discrepancies in the biofilm biomass measurements between the CV and CFU methods imply that there could be compensation for adhesion by the *Pseudomonas* species or enhanced production of the biofilm matrix under these specific conditions.

Pseudomonas species showed similar biofilm biomass (A_{590}, crystal violet) in monoculture, which was significantly induced in co-culture or multispecies (Figure S3a). However, CFU counts were not significantly changed despite the culture condition, except for PL counts in multispecies biofilms with the variant phenotype (Figure S3b PL monospecies vs. multispecies: wt = -2.3-fold, *p* = 0.2394; variant = -9.2-fold, *p* = 2.85 x 10^{-4}; Figure S4b).

224   Based on all these observations, we can conclude that the variant exhibited comparable
225   colonizing ability to its ancestor in nutrient-rich media. However, there might be a deficiency in
226   its own matrix production or the ability to induce matrix production in the *Pseudomonas* species.

Matrix glycan composition altered by cocultivation and *B. thuringiensis* adaptive phenotype

BT wild-type and variants showed different phenotypic features and biofilm formation
after 24 hours, suggesting differential matrix production in mixed-species biofilms. Thus, we
aimed to characterize matrix glycan composition by applying a fluorescent lectin screening in
multispecies biofilms (BT+PL+PB), including 78 different lectins binding to carbohydrates of
varying nature (Table S3). Binding results were filtered by fluorescent signal (data not shown) and
three lectins with strong binding in multispecies biofilms were selected for further analysis: AAL,
WGA and RCA. AAL binds α-fucose-containing polymers, WGA N-acetylglucosamine
(GlucNAc) and RCA galactose and N-acetylgalactosamine (Gal/GalNAc) glycans [42]. The
binding of such lectins was tested for BT wild-type and variant strains in mono- and dual-species
biofilms with PB.

AAL lectin showed differential binding in monoculture biofilms (Figure 4a). AAL stained
BT cells and also some matrix-like structures in BT biofilms, while the variants showed reduced
lectin signal. This suggests that glycan composition of the cell wall/surface is distinct in the
variants and, thereby, also the AAL binding. Image quantification of lectin binding supported
visual observations, with a significant reduction of lectin binding in the variants (Figure 4b);
Lectin/biomass ratio −amount of lectin binding per biovolume unit of biofilm biomass− showed a
reduced tendency in the variants, though not significant probably due to replicate variation. (Figure
4c). In coculture with PB, AAL binding was similar despite the BT phenotype, ancestral or variant
(Figure S4), suggesting a change of glycan production triggered by interspecies interactions.
However, PB also showed AAL binding in monoculture and, thus, association of lectin signal with
the specific matrix producer results cumbersome (Figure S4, BT and PB panels).

249
Figure 4. Monoculture biofilms of *B. thuringiensis* wild-type and variant strains stained with AAL lectin and Syto60. (a) Maximum intensity projection (MIP) images of Z-stacks recorded by confocal laser scanning microscopy for 24-hour biofilms. Lectin AAL: fluorescent images of biofilms stained with the fluorescent lectin AAL-FITC. Biomass: fluorescent images of biofilms stained with the biomass stain Syto60. Merge: combined images of biomass and lectin stains. BT: *B. thuringiensis* wild-type; V1: variant1, V2: variant 2; V3: variant 3; V4: variant 4; PB: *P. brenneri*. Images represent 124 µm x 124 µm field view. **(b)** Biofilm biovolume (µm³) of lectin fluorescence (AAL-FITC, green) and biomass fluorescence (red, Syto60). Biofilm biovolume was calculated using BiofilmQ per channel (10), based on ≥ three biological replicates and two images per replicate. Bars show median values per channel and error bars the standard error of the median. Dissimilar letters denote significant p-value ($p < 0.05$) per channel (green = lectin; red = biomass). **(c)** Ratios of lectin biovolume/ biomass biovolume for each strain. Each dot represents a biological replicate and crossbars indicate the median ratio for each strain. Dissimilar letters denote significant p-value ($p < 0.05$).

Despite the results observed for AAL, RCA lectin did not result in any differential binding of BT or variants in either mono or coculture with PB. This lectin bound thin filaments are produced by PB, revealing a galactose/N-Acetylgalactosamine polymer network in PB biofilms

(Figure S5). WGA bound to BT cell surfaces (Figure S6), likely the cell wall since this lectin binds

the sugar GlcNAc, a major component of peptidoglycan [43], but no difference was observed in

neither the variants nor mixed-species combinations (data not shown).

***B. thuringiensis* protein abundance decreased in response to cocultivation and genotype**

Biofilm CFU counts as well as crystal violet assays evidenced a biofilm-inducing

phenotype in cocultures of *B. thuringiensis* and *P. brenneri*. BT variants, although more abundant

after a short-term evolution assay, displayed reduced biofilm formation and/or matrix production

than BT wild-type. Thus, we investigated the influence of interspecies interactions and/or acquired

mutations on biofilm matrix proteomes. It is worth highlighting that we employed a chemical

extraction of biofilm matrix as input material for our proteomic experiments and, thus, the number

of expected proteins is much lower than for total biofilm samples. First, we performed a

comparative analysis of proteomes of *B.thuringiensis* proteomes by phenotype, variants or wild-

type, in combination with *P. brenneri* (coculture) or without (monoculture). Probabilistic principal

component analysis (PPCA) evidenced that BT wild-type monoculture samples were the most

distinct from the rest, clustering together at the right of PC1 (Figure 4a). PC1 variation seemed to

be driven by *B. thuringiensis* phenotype (wild-type vs variant), whereas PC2 by cultivation

condition (monoculture vs. coculture).   Variant coculture or monoculture samples could not be

discriminated, while BT wild-type coculture samples were grouped in a distinct cluster compared

to monoculture. Interestingly, BT coculture proteomes overlapped with variant mono- and

coculture proteomes, suggesting a change in protein abundance in the presence of *P. brenneri*.

286
Figure 5. Differential protein abundance in *B. thuringiensis* wild-type or variants in monoculture vs.
coculture biofilms with *P. brenneri*. BT = *B. thuringiensis*; PB = *P. brenneri*; BT_V1 = *B. thuringiensis*
variant 1; BT_V6 = *B. thuringiensis* variant 6. **(a)** Probabilistic PCA plot of all monoculture and co-culture
proteomes based on 5 biological replicates. Color indicates the sample type and shape the wild-type
(triangles) or variant phenotypes (circles). Ellipses indicate sample group clustering based on the principal
components. **(b)** Number of proteins differentially expressed in *B. thuringiensis* wild type monoculture
(BT) vs. coculture (BT+PB). Green bars indicate proteins with higher abundance in monoculture while red
bars indicate proteins decreased in monoculture and thus, increased in coculture. Each category role is
indicated with a letter code corresponding to the COG classification (Galperin et al., 2021), (A-Z), **(c)**
Protein abundance in BT vs BT+PB biofilms. Protein abundance is indicated in the x axis as log2 fold
change (log$_2$FC). Proteins more abundant in coculture are shown in the grey box (negative values), while
more abundant in monoculture in white (positive values). The y axis indicate –log(P-value). Dashed lines
indicate the threshold for significance (y axis) and protein abundance difference (x axis). Proteins with no
significant p-value and/ or with difference < 1 and >-1 are depicted as grey dots. Significantly changed
proteins are shown as colored points, with each COG category represented by a different color. Black dots
denote not annotated category (unknown). Analysis was done using data from 5 biological replicates and
pairwise comparisons with ≥60 % valid values (different to 0) in both co-culture and monoculture.

Additionally, different comparisons were made to investigate the impact of such factors on *B. thuringiensis* biofilm proteins: i) monoculture wild-type vs. variants and ii) coculture wild-type vs. variants and iii) monoculture vs. coculture for each pair (Figure S7b-f). We also inspected iv) *P. brenneri* proteomes comparing monoculture vs. coculture per BT strain (Figure S8).

Comparison of BT wild-type vs. variants in monoculture resulted in 187 proteins detected, 150 uniquely detected in the wild-type (Table S5 and Figure S7b), indicating major downregulation in both variants. Among those, different proteins involved in sporulation or regulated by the Stage 0 sporulation protein, Spo0A, were only detected in the wild-type, consistent with sporulation-defective phenotypes observed in the variants. To discard unique expressions due to detection limits, we calculated the mean and 0.05 percentile of each proteome to use as a cutoff (Table S4 and Figure S8a). However, DivIVA, SpoVG, immune inhibitor A (CDS_04360 and CDS_05099), neutral protease B (CDS_00401) and TasA major biofilm matrix component (CDS_ 04363 and CDS_04365) were still above the expression cutoff in BT wild-type, compared to the variants. Only 20 proteins were common to all three strains (Table S6 and Figure S7b) and 11 significantly changed, related to amino acid and carbohydrate metabolism energy production, among others. We hypothesize that the acquired mutations switch off the expression of multiple genes involved in sporulation, a highly complex and tiered-regulated process. This would be consistent with the faster doubling time observed for the variants, given that fewer proteins, and consequently energy, are required with this expression.

The proteomes of BT variants and wild-type were also compared in co-culture. Only a few proteins were detected in the variants, and protein detection in the wild-type was reduced by 70% compared to the monoculture proteome. This indicates that *P. brenneri* diminished the disparity between the three proteomes, making them more similar (Figure 4c). In the wild-type co-culture, 15 proteins were uniquely expressed (Table S7), four of which were mentioned earlier (CDS_00401, CDS_04360, CDS_04363, and CDS_04365). Interestingly, these proteins were significantly more abundant in co-cultures (Table S8), suggesting that *P. brenneri* induces their expression, which is dependent on a wild-type phenotype.

Monoculture vs. co-culture comparisons rendered highly different results depending on the phenotype compared. In the wild-type pairwise analysis, 163 proteins were detected and 111 were only expressed in monoculture (n = 111), supporting the previous observation that *P. brenneri*

reduces abundance of biofilm matrix-associated proteins in *B. thuringiensis*. Fifty proteins were co-expressed in both conditions (Figure 5b) and 50 % of which presented significantly different abundance ($p < 0.05$) and log2 fold change greater than 1 (or < -1) (Table S8 and Figures S9a), with amino acid transport and metabolism as the most represented category (Figure S9b). Analysis within the V1 and V6 variants resulted in reduced matrix proteomes, with 15 and 5 proteins detected, respectively, indicating 91-97 % downregulation in the variants compared to wild-type proteomes (Tables S9, S10 and Figures S7e-f).

P. brenneri proteomes were subjected to analysis similar to that of BT. A total of 169 proteins were detected and all mono- and co-culture proteomes showed highly similar mean expression and 0.05 quantiles (Table S4 and Figure S8a). Unlike BT results, the presence of other species did not strongly affect the number of proteins detected and 64 % of all proteins were shared among all samples (n = 108) (Figure S8b). However, PPCA of all proteomes evidenced a main culture effect, driving PC1, with PB monoculture samples clustering together and separated from the rest (Figure S8c). Interestingly, seventeen proteins were uniquely expressed in co-culture with BT wild-type (Table S11). However, co-culture samples with BT, V1, or V6 could not be discriminated in the PPCA. Volcano plots of each pairwise comparison supported similar expression patterns of *P. brenneri* in combination with *B. thuringiensis* regardless of the BT phenotype (Figure 8a). More specifically, cocultivation resulted in an increased abundance of multiple amino acid transporters in *P. brenneri*, BT despite its phenotype. (Figures 9b).

Based on all proteomics results, it can be concluded that both BT phenotype and co-cultivation impacted protein expression, though with different patterns for the species involved. For *B. thuringiensis*, different proteins involved in sporulation were only detected in the wild-type, but co-cultivation also changed expression significantly and to a similar extent (80 % downregulation). However, an abundance of *P. brenneri* proteins was highly influenced by the presence of BT regardless of its phenotype, leading to increased and decreased protein abundance.

B. thuringiensis variants detected mutations in sporulation-related genes

Whole genome sequences of the variants and wild-type revealed mutations in sporulation-related genes in the different variants correlating with the lack of sporulation observed in all the phenotypes.

Variants V1, V3 and V6 showed mutations that led to the loss of function of the *spo0A* gene, but of a different nature (Figure 6). The *spo0A* gene encodes a response regulator that, in its phosphorylated active form, activates the transcription of early sporulation genes [44, 45]. Large deletions were found in this region for variants V1 and V3: V1 lacked a 1.5 Kb fragment including most of *spo0A* and CDS0080 genes, while V3 with a 4.5 Kb deletion also covered the *spoIVB* and *recN* genes (Figure 6). V6 only showed a non-synonymous mutation in the *spo0A* gene (C151>T) resulting in a change in the amino acid sequence Q51>stop leading to early termination and, thus, loss of function of Spo0A. These mutations were confirmed by Sanger sequencing using specific primers (data not shown). However, for V2 it was not possible to confirm any clear link to the expected mutations or junction links to the expected phenotype. The mutations found in *spo0A* undoubtedly explain the non-sporulating phenotype for variants V1, V3 and V6 given the central role of *spo0A* and *spoIVB* in the sporulation process.

Figure 6. Mutations in *B. thuringiensis* variants compared to wild type and their effect on sporulation pathway. (a) Mutations found in the selected variants (BT Vx, circled pink) compared to their ancestor (BT, circled blue). Dashed boxes indicate large deletions in variants V1 and V3. Q51>* in variant V6 indicates a change in amino acid 51 of the *spo0A* gene from Q to stop codon (*). **(b)** Role of SpoA for activation of the sporulation pathway in *Bacillus* sp. Phosphorylated Spo0A activates transcription of genes involved in sporulation stages 0, I, and II. This figure was created with BioRender.com based on literature (Sonenshein, 2000; Tan & Ramamurthi, 2014).

Discussion

Microbial interactions can affect fitness and survival of the interacting species and their evolutionary trajectories, for instance, to better exploit available resources, avoid competition, improve mutual dependence or optimize the exchange of nutrients. However, adaptive evolution experiments in multispecies biofilms are still scarce. This study focused on investigating the impact of interspecies interactions and spatial structure on the selection of traits in the Gram-positive bacterium *B. thuringiensis*.

Here, we used short-term evolution assays where *B. thuringiensis, P. defluvii* and *P. brenneri* were cultured alone or combined in planktonic and biofilm settings (Fig. 1a). A novel *B. thurgingiensis* phenotypic variant, which we termed "light variant", emerged in all assays, and displayed a higher abundance than the ancestral wild-type (Fig. 1b-c). We observed that interspecies interactions with *Pseudomonas* species increased the abundance of such variant, suggesting a fitter phenotype than its ancestor in such conditions. Previous studies of experimental evolution in biofilm settings have reported the emergence of adapted variants in response to other species. However, they usually showed improved biofilm properties compared to their ancestor [18, 20, 46, 47]. Moreover, Lin and collaborators reported the appearance of a fuzzy spreader phenotype of *B. thuringiensis* in a bead-transfer evolution study, where all adapted lineages resulted in enhanced biofilm production compare to its ancestor [26]. In contrast, the *B. thuringiensis* variants selected in this study displayed reduced biofilm productivity, linked to decreased production biofilm matrix components. Still, the adapted phenotype was favored regardless of the setting (planktonic or biofilm, Fig. 1c) and nutrient availability (Fig. S1). This points towards a gained fitness advantage of the non-sporulating variants over the wild-type, consistent with their faster generation time.

Whole-genome sequencing of the variants linked lack of sporulation with mutations in key genes regulating sporulation genes (Fig. 5). Some of the sequenced isolates of the light variant (V1, V3, V6) harbored mutations in *spo0A* and *spoIVB*, genes with a central role in the sporulation process. The *spo0A* gene encodes a response regulator that, in its phosphorylated active form, activates transcription of early sporulation genes [44, 45]. It has been shown that mutations in *spoIVB* or *spo0A* successfully blocks sporulation in *Bacillus subtilis*. Consequently, mutations in

these genes will result in failure to initiate sporulation or cortex formation and, consequently,
explain the non-sporulation phenotype of variants V1 and V3-V6.

Irrespective of the particular mutations, all *B. thuringiensis* light variants showed faster
generation time than the ancestor (Table S2), non-sporulating phenotype (Figure 2b), and reduced
AAL lectin binding (Figure 4), pointing towards a convergent evolutionary pathway for all light
variants. Quantification of biofilm biomass by crystal violet assays proved reduced matrix
production of the *B. thuringiensis* light variants (Fig. 3a) in mixed culture biofilms suggesting a
potential impairment of matrix production, or reduced triggering of matrix production in
Pseudomonas species. Consistent with this observation, proteomic analysis evidenced unique
expression of different matrix proteins in the wild-type. Two TasA/CalY orthologues and SpoVG
were expressed in BT wild-type but not in the variants. TasA is a major matrix component in
Bacillus biofilms and its absence results in significant defects in biofilm and pellicle formation in
B. subtilis [48, 49] while its paralog is needed for root colonization in *B. cereus* [50]. In *B.*
thuringiensis, the TasA-orthologue CalY has been identified as a cell-surface adhesion and
described as a major virulence factor and biofilm matrix protein [51]. SpoVG is a regulator of both
sporulation and biofilm formation in *B. cereus* 0-9 that activates the levels of Spo0A and, thus, the
fate of *B. cereus* cells: biofilm or sporulation [52, 53]. We speculate that the lack of expression of
TasA and the biofilm regulatory protein SpoVG could contribute to the differences observed
between BT CFU counts and total biofilm biomass for the variant phenotype (CV). Proteomic
results also evidenced the unique expression of two metalloproteases only in *B. thuringiensis* wild-
type: NprB and InhA (immune inhibitor A). These proteins have been linked with protein recycling
during sporulation and are directly or indirectly regulated by Spo0A [44, 54, 55].

From a community perspective, all species were preserved after the 8-day evolution
experiment in multispecies combinations (Figure S1), suggesting a stable interaction between the
three species. Interestingly, interspecies interactions impacted differently protein abundance
depending on the species. We observed that cocultivation significantly altered the proteome of *B.*
thuringiensis but only in the ancestor (Figure 4a). Meanwhile, for *P. brenneri* the presence of BT
greatly impacted protein abundance regardless of the phenotype (Figure S9a). Previous studies
have shown that coexistence can remain stable over time if they are not exploitable, for instance,
altering the pH or secretion of costless by-products to the other species' benefit [56–61]. Despite

its phenotype, we observed a higher abundance of amino acid transporters in *P. brenneri* when cocultivated with *B. thuringiensis*, suggesting potential cross-feeding in such conditions. We have previously reported that cooperative interactions between community members can facilitate metabolic cross-feeding on specific amino acids, though in a different multispecies biofilm model [37, 62]. Thus, we postulate a non-exploitable interaction between the variant phenotype and *P. brenneri* species, that is, improved growth but not at the cost of the other species.

Biofilm formation and spore production are two different survival strategies of bacteria facing adverse environmental stresses. However, intertwined regulatory pathways between biofilm formation and sporulation have been proposed for both *B. subtilis* and *B. cereus* [63, 64]. Recent work connects spore formation with biofilm dispersal in *Bacillus subtilis* [65]. When conditions become unfavorable, spores can detach from the biofilm and be carried away by air, water, or other vectors. Dispersed spores can initiate new biofilm formation in different niches, aiding in bacterial survival and colonization. This study used an experimental evolution setup that selected for biofilm-associated. In such conditions, we hypothesize that non-sporulating variants would undergo limited passive dispersal and vegetative cells may be forced to stay within the biofilm, dead or alive. Thus, sporulation in such a setting would be dispensable for *B. thuringiensis.*

Our results support that non-sporulating variants can be selected at a higher rate in *in vitro* biofilm setups and in the presence of co-isolated species; thus, spatial structure, diversity and complexity within microbial communities should be considered as drivers of bacterial diversification in such environments.

Methods

Strains and materials

The three bacterial strains *Pseudomonas defluvii* (PL), *Pseudomonas brenneri* (PB) and *Bacillus thuringiensis* (BT) were isolated from water samples taken at a wastewater treatment plant in Kalundborg, Denmark, and genome sequenced earlier. All strains were grown at 24 °C in Tryptic Soy Broth (TSB; 17 g pancreatic digest of casein, 3 g papaic digest of soybean meal, 5 g sodium chloride, 2.5 g dextrose, 2.5 g dibasic potassium phosphate in 1 L distilled water, pH 7.3), supplemented with 1.5 % agar when needed. To distinguish the bacterial strains and/or species variants, 40 µg/mL of Congo Red and 20 µg/mL Coomassie Brilliant Blue G were added to the TSB base.

Short-term evolution assay with spatial structure

Overnight cultures in TSB medium were adjusted to an optical density at 600 nm (OD_{600}) of 0.15 prior to usage. The experiment was conducted in 24-well plates, where each well contained a polycarbonate chip (12 x 12 x 0.78 mm^3) (PC) that served as a surface for establishing the biofilm. In each well, 2 mL of OD_{600} 0.15 cultures were added, either as mono-species culture or mixed-species culture. After 24 hours, the floating pellicle was gently removed before the polycarbonate chip was transferred to a new well with fresh media. This was done for 8 days, followed by washing the polycarbonate chip 3 times in 1x phosphate-buffered saline (PBS) and placing it in a 5 mL Eppendorf tube with 2 ml 1x PBS. The biofilm was then removed from the polycarbonate surface and the biofilm structure was homogenized by shaking at 600 rpm for 5 minutes. The resulting suspension was then plated on TSB solid medium, complemented with 40 µg/mL Congo Red and 20 µg/mL Coomassie Brilliant Blue with three technical replicates to estimate colony forming units (CFUs) and to screen for new variants with different colony morphology. The same setup was utilized to evaluate interactions between strains and new variants, with growth on polycarbonate chips for 24 hours.

Short-term evolution assay without spatial structure

To test selection pressure in conditions not favoring spatial structure, a short-term evolution assay was also performed with well-mixed planktonic cultures. Cultures were adjusted to $OD_{600} = 0.15$ as before, and 5 mL of this culture was incubated overnight, with shaking at 250 rpm. After 24

493 hours, 5 µL of this culture was transferred to new fresh media. The transfer was repeated for 8

494 days, followed by plating on TSB solid medium with Congo Red and Coomassie Brilliant Blue,

495 with three technical replicates.

Quantification of biofilm formation

Assessment of the biofilm formation on the polycarbonate surfaces was performed using a

modified crystal violet assay [29]. Biofilms were allowed to grow on a polycarbonate surface for

24 hours as described above, washed three times in 1x PBS and then transferred to a well of a 24-

well plate containing 2 mL of 1 % crystal violet aqueous solution and incubated for 20 min in the

dark. Then, polycarbonate chips were carefully removed and rinsed five times in 1x PBS. Crystal

violet stained polycarbonate chips were then incubated in 2 mL of 96 % ethanol for 30 min. The

biofilm biomass was then quantified as the absorbance of crystal violet at 590 nm (A_{590}). To avoid

measurement saturation, samples were diluted in 96% ethanol when A_{590} was higher than 1.

Growth curves in liquid cultures

Cultures of the wild-type and variants were allowed to grow overnight in liquid TSB medium.

Then, 1:100 dilutions of the overnight cultures in TSB were prepared, and these cultures were used

to inoculate 96-well plates, using 3 biological and 3 technical replicates. Growth was monitored

for 24 hours at 23 °C with 282 rpm double orbital shaking and OD_{600} readings every 10 minutes.

Growth data was integrated to calculate different growth parameters using the Growthcurver R

package [30] to determine the following parameters: t_mid, the point of inflection of the growth

curve; K, carrying capacity or maximum size of the population; t_gen, generation time of a strain;

r, intrinsic growth rate of a population; sigma, the fit goodness of the model.

Optical microscopy

515 Liquid cultures of the different strains were incubated for 24 h in TSB, at 24 °C with shaking at

516 250 rpm. From these 24-h old cultures, 10 µL were pipetted onto a microscope glass slide and a

517 coverslip was applied on top. Samples were observed with an Olympus BH2 optical microscope

518 with 10x and 40x magnification objectives.

Sporulation assay

To compare the sporulation capacity of *Bacillus thuringiensis* and its variants, a sporulation assay was performed. Firstly, the cell concentration in liquid cultures was measured by enumerating CFUs per mL. Next, the culture was heat-shocked at 80 ˚C for 20 minutes before CFUs were enumerated again. The following calculation was used to estimate the percentage of spores in the culture: colonies per mL after heat shock divided by colonies per mL before heat shock. Furthermore, the sporulation assay was performed for liquid cultures, but also for the biofilms grown on polycarbonate chips, to estimate the number of spores formed in mono-strain cultures and multi-strain cultures.

Biofilm matrix staining with fluorescent lectins

Matrix glycoconjugate characterization in the three species community was performed using 78 different fluorescently labeled lectins (using FITC, AlexaFluor488, or Fluorescein) in combination with confocal laser scanning microscopy (TCS SP5X, Leica Germany, controlled by the software program LAS-AF vers. 2.41) as described previously by Neu and Kuhlicke [31, 32]. The lectins and fluorescent conjugates that were used are listed in Supplementary Table S3. Briefly, biofilms were grown on polycarbonate chips for 24 hours, followed by washing three times with 1x PBS, as described above. Staining solutions containing fluorescent lectin was prepared at concentration 100 µg/mL (in filter-sterilized ultrapure water) and 100 then added to a biofilm on a polycarbonate chip, followed by incubation for 30 min in the dark. Excess staining solution was washed twice with dH_2O before adding SYTO60 to the sample, prior to confocal imaging. Excitation of FITC, AlexaFluor488, and Fluorescein was performed at 488 nm, and the emission signals were recorded from 500–580 nm. Excitation of SYTO60 was performed at 652 nm and emission was recorded from 665-750 nm. For lectins conjugated to Rhodamine, excitation was done at 525 nm and emission was recorded from 570-650 nm. Image data sets were recorded using 63x NA 0.9 and 63x NA 1.2 objective lenses with a step size of 0.5 µm.

Based on the screening, three lectins: *Aleuria Aurantia* Lectin (AAL), Wheat Germ Agglutinin (WGA) and *Ricinus Communis* Agglutinin (RCA) were chosen for further testing. Here, a dual species combination of *Pseudomonas brenneri* (PB) and *Bacillus thuringiensis* (BT) was used to estimate better the effects between, and on *Bacillus thuringiensis* and its variant when co-cultured with *Pseudomonas*. Images of AAL-lectin-stained biofilms of BT and variants V1, V2 and V3

550 were analyzed using the biofilm image analysis software suite BiofilmQ [33]. Biofilm volume

551 (biovolume) was calculated for each fluorescent channel: lectin (FITC channel) and biomass

552 (SYTO60 channel) with a minimum of 6 images per sample (3 biological replicates and 2 images

553 per replicate). Moreover, biovolume ratios (AAL-lectin-stained biovolume divided by the total

554 biovolume) were calculated per biological replicate.

Biofilm matrix proteome measurements

Biofilms were allowed to grow for 24 hours in TSB medium on polycarbonate chips as described

above, followed by washing 3 times in 1x PBS to remove planktonic cells and biofilm pellicle.

Five polycarbonate chips were combined to extract a sufficient amount of biofilm matrix material

for proteomics analysis. The biofilm was then resuspended in 5 mL of 1x PBS by shaking at 1000

rpm for 10 min. The biofilm matrix was extracted using the formaldehyde and NaOH extraction

method [34, 35]. Briefly, 0.006 V formaldehyde 37% (30 µL) was added to the biofilm samples

and incubated for 1 hour at 4 °C while mixing, followed by addition of 0.4 V NaOH 1 N (2 mL)

and incubation for 3 hours at 4 °C while mixing. Samples were then dialyzed using 3500 MWCO

22 mm SnakeSkin™ Dialysis Tubing (ThermoFisher Scientific) against 2.5 L milliQ water,

overnight at 4 °C with stirring. Protein content in dialyzed samples was quantified using the Bio-

Rad Protein assay, a modified Bradford assay [36], before freezing with liquid nitrogen. Finally,

extracted biofilm matrix was lyophilized using an ALPHA 1-4 LDplus freeze dryer (Christ, VWR)

and kept at -80 °C until sample preparation for proteomics.

For the proteomics sample preparation, digestion, mass spectrometry, and data analysis were

conducted as described in Herschend et al. [37] with some modifications, described in the

supplementary methods. Reference genomes of *B. thuringiensis, P. defluvii*, and *P. brenneri* were

prepared as part of a previous study and are available online (NCBI) with IDs JARTVI000000000,

CP123067.1 and CP122540.1, respectively.

Whole-genome sequencing and analysis of variants

Genomes of *Bacillus thuringiensis* wild-type and variants V1, V2, V3 and V6 (Table S1) were

sequenced by extracting the DNA using DNeasy Blood & Tissue Kit (Qiagen) from overnight

cultures. The samples were sequenced with short reads using Truseq libraries on the Illumina

NovaSeq 6000 and 2x150 cycles (Novogene, UK).

Data was analyzed using breseq (v0.35.0) to call variants using the –p polymorphism flag when

analyzing population sequences [38]. As the reference genome, we used the previously constructed

hybrid assembly of long and short reads from *B. thuringiensis,* available online (NCBI) with ID

JARTVI000000000. Mutations that were detected between our ancestral strain and the reference

genome were not considered. The breseq software also predicts new junctions within the genome

which may occur due to mobile element insertions, large deletions, structural rearrangements and

prophage excision. These analyses were also included. Unfiltered breseq output can be accessed

in the supplementary material.

Statistical analysis and visualization

All analysis and plotting was done in RStudio v4.1.1 [39]. Unless otherwise specified, log10

transformed data was fitted to linear models and tested for normality and equal variance. Statistical

significance was then tested with ANOVA using the same parameters as the linear models. A

general expression was ANOVA(CFU ~Combination*Phenotype), however the quantitative

variable and categorical variables changed depending on the dataset. A Tukey's post-hoc test was

performed to consider multiple comparisons when the ANOVA test was significant, unless

otherwise specified. Compact letter display for multiple comparisons was calculated using R

package "multicompView" v0.1-8 [40], while package "ggplot2" v3.4.0 was used for data

visualization [41].

Acknowledgments

The authors thank Sara Olsen and Anette Løth for technical assistance, the DTU Proteomics core facility for technical assistance in proteomics sample preparation. Support by Ute Kuhlicke with lectin analysis and confocal laser scanning microscopy is highly appreciated. We thank the Villum Foundation for grants to MB (no. 10098) and HR (no. 34434) and the US Office of Naval Research, for grant N629091812174 to MB, as well as the Swiss National Science Foundation Consolidator Grant TMCG-3_213801 to KD, and the German Bundesministerium für Bildung und Forschung grant TARGET-Biofilms 16GW0245 to KD, funding this work.

Conflict of interests

The authors declare that no conflicts of interest exist.

References

1. Agrawal AA. Phenotypic plasticity in the interactions and evolution of species. *Science* 2001; **294**: 321–326.

2. Barrett RDH, Schluter D. Adaptation from standing genetic variation. *Trends Ecol Evol* 2008; **23**: 38–44.

3. Barrick JE, Lenski RE. Genome dynamics during experimental evolution. *Nat Rev Genet 2013 1412* 2013; **14**: 827–839.

4. Scheuerl T, Hopkins M, Nowell RW, Rivett DW, Barraclough TG, Bell T. Bacterial adaptation is constrained in complex communities. *Nat Commun 2020 111* 2020; **11**: 1–8.

5. Calcagno V, Jarne P, Loreau M, Mouquet N, David P. Diversity spurs diversification in ecological communities. *Nat Commun* 2017; **8**.

6. Emerson BC, Kolm N. Species diversity can drive speciation. *Nature* 2005; **434**: 1015–1017.

7. Nadell CD, Xavier JB, Foster KR. The sociobiology of biofilms. *FEMS Microbiol Rev* 2009; **33**: 206–224.

8. Gross K, Cardinale BJ. Does species richness drive community production or vice versa? Reconciling historical and contemporary paradigms in competitive communities. *Am Nat* 2007; **170**: 207–220.

9. Cardinale BJ, Wright JP, Cadotte MW, Carroll IT, Hector A, Srivastava DS, et al. Impacts of plant diversity on biomass production increase through time because of species complementarity. *Proc Natl Acad Sci U S A* 2007; **104**: 18123–18128.

10. Nadell CD, Drescher K, Foster KR. Spatial structure, cooperation and competition in biofilms. *Nat Rev Microbiol 2016 149* 2016; **14**: 589–600.

11. Rainey PB, Travisano M. Adaptive radiation in a heterogeneous environment. *Nat 1998 3946688* 1998; **394**: 69–72.

12. Habets MGJL, Rozen DE, Hoekstra RF, De Visser JAGM. The effect of population structure on the adaptive radiation of microbial populations evolving in spatially structured environments. *Ecol Lett* 2006; **9**: 1041–1048.

13. Foster KR, Bell T. Competition, not cooperation, dominates interactions among culturable microbial species. *Curr Biol* 2012; **22**: 1845–1850.

14. Xavier JB, Foster KR. Cooperation and conflict in microbial biofilms. *Proc Natl Acad Sci U S A* 2007; **104**: 876–881.

15. Rao D, Webb JS, Kjelleberg S. Competitive interactions in mixed-species biofilms containing the marine bacterium Pseudoalteromonas tunicata. *Appl Environ Microbiol* 2005; **71**: 1729–1736.

16. Hibbing ME, Fuqua C, Parsek MR, Peterson SB. Bacterial competition: surviving and
thriving in the microbial jungle. *Nat Rev Microbiol* 2010; **8**: 15.

17. Ghoul M, Mitri S. The Ecology and Evolution of Microbial Competition. *Trends
Microbiol* 2016; **24**: 833–845.

18. Røder HL, Herschend J, Russel J, Andersen MF, Madsen JS, Sørensen SJ, et al. Enhanced
bacterial mutualism through an evolved biofilm phenotype. *ISME J* 2018.

19. Liu W, Jacquiod S, Brejnrod A, Russel J, Burmølle M, Sørensen SJ. Deciphering links
between bacterial interactions and spatial organization in multispecies biofilms. *ISME J*
2019; **13**: 3054–3066.

20. Henriksen NNSE, Hansen MF, Kiesewalter HT, Russel J, Nesme J, Foster KR, et al.
Biofilm cultivation facilitates coexistence and adaptive evolution in an industrial bacterial
community. *NPJ biofilms microbiomes* 2022; **8**.

21. McDonald MJ, Gehrig SM, Meintjes PL, Zhang XX, Rainey PB. Adaptive Divergence in
Experimental Populations of Pseudomonas fluorescens. IV. Genetic Constraints Guide
Evolutionary Trajectories in a Parallel Adaptive Radiation. *Genetics* 2009; **183**: 1041.

22. Traverse CC, Mayo-Smith LM, Poltak SR, Cooper VS. Tangled bank of experimentally
evolved Burkholderia biofilms reflects selection during chronic infections. *Proc Natl Acad
Sci U S A* 2013; **110**: E250–E259.

23. Thöming JG, Tomasch J, Preusse M, Koska M, Grahl N, Pohl S, et al. Parallel
evolutionary paths to produce more than one Pseudomonas aeruginosa biofilm phenotype.
NPJ Biofilms Microbiomes 2020; **6**.

24. Leiman SA, Arboleda LC, Spina JS, McLoon AL. SinR is a mutational target for fine-
tuning biofilm formation in laboratory-evolved strains of Bacillus subtilis. *BMC Microbiol*
2014; **14**.

25. Dragoš A, Lakshmanan N, Martin M, Horváth B, Maróti G, García CF, et al. Evolution of
exploitative interactions during diversification in Bacillus subtilis biofilms. *FEMS
Microbiol Ecol* 2018; **94**.

26. Lin Y, Xu X, Maróti G, Strube ML, Kovács ÁT. Adaptation and phenotypic
diversification of Bacillus thuringiensis biofilm are accompanied by fuzzy spreader
morphotypes. *npj Biofilms Microbiomes 2022 81* 2022; **8**: 1–14.

27. Ehling-Schulz M, Koehler TM, Lereclus D. The Bacillus cereus Group: Bacillus species
with Pathogenic Potential. *Microbiol Spectr* 2019; **7**.

28. Poltak SR, Cooper VS. Ecological succession in long-term experimentally evolved
biofilms produces synergistic communities. *ISME J* 2011; **5**: 369.

29. O'Toole G, Kolter R. Initiation of biofilm formation in Pseudomonas fluorescens
WCS365 proceeds via multiple, convergent signalling pathways: a genetic analysis. *Mol
Microbiol* 1998; **28**: 449–461.

681 30. Sprouffske K, Wagner A. Growthcurver: An R package for obtaining interpretable metrics
682   from microbial growth curves. *BMC Bioinformatics* 2016; **17**: 1–4.

683 31. Neu T, Kuhlicke U. Fluorescence Lectin Bar-Coding of Glycoconjugates in the
684   Extracellular Matrix of Biofilm and Bioaggregate Forming Microorganisms.
685   *Microorganisms* 2017.

686 32. Neu TR, Kuhlicke U. Matrix glycoconjugate characterization in multispecies biofilms and
687   bioaggregates from the environment by means of fluorescently-labeled lectins. *Front
688   Microbiol* 2022; **13**.

689 33. Hartmann R, Jeckel H, Jelli E, Singh PK, Vaidya S, Bayer M, et al. Quantitative image
690   analysis of microbial communities with BiofilmQ. *Nat Microbiol 2021 62* 2021; **6**: 151–
691   156.

692 34. Liu H, Fang HHP. Extraction of extracellular polymeric substances (EPS) of sludges. *J
693   Biotechnol* 2002; **95**: 249–256.

694 35. Pan X, Liu J, Zhang D, Chen XI, Li L, Song W, et al. A comparison of five extraction
695   methods for extracellular polymeric substances (EPS) from biofilm by using three-
696   dimensional excitation-emission matrix (3DEEM) fluorescence spectroscopy. *undefined*
697   2010; **36**: 111–116.

698 36. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities
699   of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976; **72**: 248–254.

700 37. Herschend J, Damholt ZBV, Marquard AM, Svensson B, Sørensen SJ, Hägglund P, et al.
701   A meta-proteomics approach to study the interspecies interactions affecting microbial
702   biofilm development in a model community. *Sci Rep* 2017.

703 38. Deatherage DE, Barrick JE. Identification of mutations in laboratory-evolved microbes
704   from next-generation sequencing data using breseq. *Methods Mol Biol* 2014; **1151**: 165–
705   188.

706 39. Team. RStudio: Integrated Development for R. *Rstudio, BMC* 2021.

707 40. Piepho HP. An Algorithm for a Letter-Based Representation of All-Pairwise
708   Comparisons. *http://dx.doi.org/101198/1061860043515* 2012; **13**: 456–466.

709 41. Wickham H. ggplot2: Elegant Graphics for Data Analysis. 2016. Springer-Verlag, New
710   York.

711 42. Neu TR, Kuhlicke U. Fluorescence Lectin Bar-Coding of Glycoconjugates in the
712   Extracellular Matrix of Biofilm and Bioaggregate Forming Microorganisms.
713   *Microorganisms* 2017; **5**.

714 43. Angeles DM, Scheffers DJ. The Cell Wall of Bacillus subtilis. *Curr Issues Mol Biol 2021,
715   Vol 41, Pages 539-596* 2020; **41**: 539–596.

716 44. Fawcett P, Eichenberger P, Losick R, Youngman P. The transcriptional profile of early to
717   middle sporulation in Bacillus subtilis. *Proc Natl Acad Sci U S A* 2000; **97**: 8063–8068.

45. Molle V, Fujita M, Jensen ST, Eichenberger P, González-Pastor JE, Liu JS, et al. The
Spo0A regulon of Bacillus subtilis. *Mol Microbiol* 2003; **50**: 1683–1701.

46. Hansen SK, Rainey PB, Haagensen JAJ, Molin S. Evolution of species interactions in a
biofilm community. *Nat 2006 4457127* 2007; **445**: 533–536.

47. Cosetta CM, Niccum B, Kamkari N, Dente M, Podniesinski M, Wolfe BE. Bacterial–
fungal interactions promote parallel evolution of global transcriptional regulators in a
widespread Staphylococcus species. *ISME J 2023 179* 2023; **17**: 1504–1516.

48. Romero D, Aguilar C, Losick R, Kolter R. Amyloid fibers provide structural integrity to
Bacillus subtilis biofilms. *Proc Natl Acad Sci U S A* 2010; **107**: 2230–2234.

49. Branda SS, Chu F, Kearns DB, Losick R, Kolter R. A major protein component of the
Bacillus subtilis biofilm matrix. *Mol Microbiol* 2006; **59**: 1229–1238.

50. Lakshmanan V, Kitto SL, Caplan JL, Hsueh Y-H, Kearns DB, Wu Y-S, et al. Microbe-
Associated Molecular Patterns-Triggered Root Responses Mediate Beneficial
Rhizobacterial Recruitment in Arabidopsis 1[C][W][OA]. *Plant Physiol Ò* 2003; **160**:
1642–1661.

51. Candela T, Fagerlund A, Buisson C, Gilois N, Kolstø AB, Økstad OA, et al. CalY is a
major virulence factor and a biofilm matrix protein. *Mol Microbiol* 2019; **111**: 1416–1429.

52. Huang Q, Zhang Z, Liu Q, Liu F, Liu Y, Zhang J, et al. SpoVG is an important regulator
of sporulation and affects biofilm formation by regulating Spo0A transcription in Bacillus
cereus 0–9. *BMC Microbiol* 2021; **21**: 1–17.

53. Lin Y, Briandet R, Kovács ÁT. Bacillus cereus sensu lato biofilm formation and its
ecological importance. *Biofilm* . 2022. , **4**

54. Livingston ET, Mursalin MH, Coburn PS, Astley R, Miller FC, Amayem O, et al. Immune
inhibitor a metalloproteases contribute to virulence in Bacillus endophthalmitis. *Infect
Immun* 2021; **89**.

55. Harwood CR, Kikuchi Y. The ins and outs of Bacillus proteases: activities, functions and
commercial significance. *FEMS Microbiol Rev* 2022; **46**: 1–20.

56. Ratzke C, Gore J. Modifying and reacting to the environmental pH can drive bacterial
interactions. *PLoS Biol* 2018; **16**.

57. Røder HL, Herschend J, Russel J, Andersen MF, Madsen JS, Sørensen SJ, et al. Enhanced
bacterial mutualism through an evolved biofilm phenotype. *ISME J* 2018.

58. Herschend J, Koren K, Røder HL, Brejnrod A, Kühl M, Burmølle M. In Vitro Community
Synergy between Bacterial Soil Isolates Can Be Facilitated by pH Stabilization of the
Environment. *Appl Environ Microbiol* 2018; **84**.

59. Pacheco AR, Segrè D. A multidimensional perspective on microbial interactions. *FEMS
Microbiol Lett* 2019; **366**: 125.

60.   Aranda-Díaz A, Obadia B, Dodge R, Thomsen T, Hallberg ZF, Güvener ZT, et al.
Bacterial interspecies interactions modulate pH-mediated antibiotic tolerance. *Elife* 2020;
9.

61.   Fritts RK, McCully AL, McKinlay JB. Extracellular Metabolism Sets the Table for
Microbial Cross-Feeding. *Microbiol Mol Biol Rev* 2021; **85**.

62.   Hansen LBS, Ren D, Burmølle M, Sørensen SJ. Distinct gene expression profile of
Xanthomonas retroflexus engaged in synergistic multispecies biofilm formation. *ISME J*
2017.

63.   Vishnoi M, Narula J, Devi SN, Dao HA, Igoshin OA, Fujita M. Triggering sporulation in
Bacillus subtilis with artificial two-component systems reveals the importance of proper
Spo0A activation dynamics. *Mol Microbiol* 2013; **90**: 181–194.

64.   Huang Y, Flint SH, Palmer JS. Bacillus cereus spores and toxins – The potential role of
biofilms. *Food Microbiol* 2020; **90**: 103493.

65.   Nishikawa M, Kobayashi K. Calcium Prevents Biofilm Dispersion in Bacillus subtilis. *J*
Bacteriol 2021; **203**.

66.   Galperin MY, Wolf YI, Makarova KS, Alvarez RV, Landsman D, Koonin E V. COG
database update: focus on microbial diversity, model organisms, and widespread
pathogens. *Nucleic Acids Res* 2021; **49**: D274–D281.

67.   Sonenshein AL. Control of sporulation initiation in Bacillus subtilis. *Curr Opin Microbiol*
2000; **3**: 561–566.

68.   Tan IS, Ramamurthi KS. Spore formation in Bacillus subtilis. *Environ Microbiol Rep*
2014; **6**: 212.
776