Using a modular massively parallel reporter assay to discover context-dependent regulatory

activity in type 2 diabetes-linked noncoding regions

Adelaide Tovar[1], Yasuhiro Kyono[1,4], Kirsten Nishino[1], Maya Bose[1], Arushi Varshney[1], Stephen C.J.

Parker[1,2,3,5], and Jacob O. Kitzman[1,2,5]

[1]Gilbert S. Omenn Department of Computational Medicine and Bioinformatics,

[2]Department of Human Genetics,

[3]Department of Biostatistics,

University of Michigan, Ann Arbor, MI 48109,

[4]Current affiliation: Seattle Hub for Synthetic Biology, Allen Institute, Seattle, WA 98109,

[5]Co-corresponding authors.

Address correspondence to: scjp@umich.edu (S.C.J.P.) and kitzmanj@umich.edu (J.O.K.)

Running head: MODULAR MPRA TO DISCOVER REGULATORY GRAMMARS

Keywords: transcription factors, enhancers, diabetes, gene regulation

Abstract

Most genome-wide association signals for complex disease reside in the noncoding genome, where

defining function is nontrivial. MPRAs (massively parallel reporter assays) offer a scalable means to

identify functional regulatory elements, but are typically conducted without regard to cell type, pairing

cloned fragments with a generic housekeeping promoter. To explore the context-sensitivity of MPRAs,

we screened enhancer activity across a panel of nearly 12,000 198-bp fragments spanning over 300

type 2 diabetes- and metabolic trait-associated regions in the 832/13 rat insulinoma beta cell line, a

relevant model of pancreatic beta cells. We explored these fragments' context sensitivity by comparing

their activities when placed up- or downstream of a reporter gene, and in combination with either a

synthetic housekeeping promoter (SCP1) or a more biologically relevant promoter corresponding to the

human insulin (*INS*) gene. We identified clear effects of MPRA construct design on enhancer activity.

Specifically, a subset of fragments (n = 702/11,656) displayed positional bias, evenly distributed across

up- and downstream preference. Promoter choice also influenced MPRA activity (n = 698/11,656),

mostly biased towards the cell-specific *INS* promoter (73.4%). To identify sequence features associated

with promoter preference, we used Lasso regression with 562 genomic annotations and discovered that

fragments with *INS* promoter-biased activity are enriched for HNF1 motifs. HNF1 family transcription

factors are key regulators of glucose metabolism disrupted in maturity onset diabetes of the young

(MODY), suggesting genetic convergence between rare coding variants that cause MODY and

common T2D-associated regulatory regions. We designed a follow-up MPRA containing HNF1 motif-

enriched fragments and observed several instances where deletion or mutation of HNF1 motifs

disrupted the *INS* promoter-biased enhancer activity, specifically in the beta cell model but not in a

skeletal muscle cell line, another diabetes-relevant cell type. Together, our study suggests that cell-

specific regulatory activity is partially influenced by enhancer-promoter compatibility and indicates that

careful attention should be paid when designing MPRA libraries to capture context-specific regulatory

processes at disease-associated genetic signals.

Introduction

Complex diseases arise from individual and interactive effects of genetic, environmental, and lifestyle factors. Recent genome-wide association studies have revealed that the majority of complex disease-associated loci (~90%) are found in noncoding regions where they are expected to alter gene regulation (Watanabe et al. 2019). Progress to identify causal variants and define their role(s) in disease pathogenesis has been slow due to the lack of a clear variant-to-function framework for noncoding variants. Massively parallel reporter assays (MPRAs) have emerged as a useful method for screening large collections of putative regulatory elements, learning principles of disease-associated regulatory variation, and efficiently prioritizing candidate causal variants at disease loci (Khetan et al. 2021; Tewhey et al. 2016; Arnold et al. 2013; Bergman et al. 2022).

A wide array of MPRA designs exist, primarily differing by the position where fragments of interest are cloned relative to a basal promoter, and by reporter gene (summarized in (Klein et al. 2020)). Enhancers are classically defined as acting at a distance in an orientation-independent manner (Banerji et al. 1981), though how distinct they are from promoters remains unclear. Klein et al. reported modest impacts of fragment position on enhancer activity, which could nevertheless mask the often subtle regulatory effects being measured (Ulirsch et al. 2016; Choi et al. 2020; Tewhey et al. 2016; Lu et al. 2021; Myint et al. 2020; McAfee et al. 2023), clouding downstream interpretation.

In addition to fragment positioning, another key design consideration is the choice of promoter. To date, most MPRAs used to study disease-associated genetic variants rely on a standard design featuring one of several housekeeping promoters (e.g., minimal promoter [minP], super core promoter [SCP1]) (Tewhey et al. 2016; McAfee et al. 2022; Melnikov et al. 2012; Patwardhan et al. 2012). There is conflicting evidence as to the influence of promoter-enhancer compatibility rules: some studies suggest that unique sequence features including transcription factor motifs dictate how strongly an interaction will occur (Zabidi et al. 2015; Martinez-Ara et al. 2022) while others indicate that there is minimal specificity with a simple additive model combining intrinsic enhancer and promoter strength (Hong and Cohen 2022; Bergman et al. 2022; Sahu et al. 2022). There have been limited comparisons of MPRA design parameters including fragment position and enhancer-promoter compatibility, and how

these choices interact at disease-associated loci has not yet been systematically examined. We

therefore speculated that using a promoter of a physiologically relevant gene may improve our ability to

detect disease-associated regulatory perturbations in a tractable and well-matched cell model.

Here, we used a modular MPRA to assess enhancer activity across multiple promoter and

position contexts to evaluate how enhancer-promoter compatibility contributes to context-specific gene

regulation in type 2 diabetes.

Results

Massively parallel reporter design and cloning

To explore regulatory activity across loci associated with type 2 diabetes (T2D) and related

metabolic traits, we designed a library of 13,226 targets spanning regions in high linkage disequilibrium

($R^2 > 0.8$) with index signals reported by the DIAMANTE (Mahajan et al. 2022, 2018; Spracklen et al.

2020) and MAGIC (Chen et al. 2021) consortia (**Figure 1**; **Table S1**). We added short 16 bp anchor

sequences on both sides of each fragment to enable cloning into the MPRA constructs. We cloned this

fragment library into four different MPRA constructs: upstream or downstream of a portion of the human

insulin (*INS*) promoter or the synthetic super core (SCP1) promoter (Juven-Gershon et al. 2006)

(**Figure S1A**). We simultaneously delivered the resulting plasmid libraries to the 832/13 rat insulinoma

cell line via electroporation, then collected cells 24 hours later to isolate RNA. We performed this

experiment in triplicate and prepared barcode sequencing libraries from the cDNA and plasmid input to

estimate enhancer activity of each fragment (**Figure S1C**).

Regulatory activities measured across fragments

Our MPRA library contained both reference and alternate alleles but was not barcoded to

sufficient complexity for allelic effect detection. Accordingly, we focused on patterns of regulatory

activity across sequence contexts rather than allelic bias in activity. After filtering for fragments

represented by greater than two barcodes and with at least ten input DNA counts across all four

configurations, we proceeded with a set of 11,656 fragments for downstream analysis.

To qualitatively examine the influence of position and promoter on fragment activity, we used

principal component analysis (PCA; **Figure 2A**) on $\log_2$(RNA/DNA) activity estimates using library size-

normalized counts. We identified that the four constructs separate on principal component 2 (13.25% of

variance) based on the position into which fragments were cloned, where both downstream constructs

cluster together separately from the two upstream constructs. The two upstream constructs further

separate on principal component 1 (15.68% of variance) based on the promoter that was used. To

evaluate whether other technical factors confounded the group separations we saw using PCA, we

calculated pairwise Spearman correlation between the first five principal components, RNA library size,

DNA library size, and ratio of RNA library size/DNA library size (library size ratio) (**Figure S2A**). We

identified strong negative correlation ($R^2 = 0.972$) between PC1 and library size ratio, suggesting a

relationship between PC1 and overall enhancer activity for a given library. Additionally, we observed a

strong relationship between RNA library size and DNA library size ($R^2 = 0.972$), which is expected

given that RNA counts are partially dependent on the plasmid DNA counts. Otherwise, we observed no

strong correlations between any of the PCs and technical factors.

To systematically test all fragments in each library for enhancer activity, we used the R package

MPRAnalyze (Ashuach et al. 2019), which estimates activity by comparing the RNA counts for each

barcode to DNA counts for the same barcodes (**Tables S2-5**). Within each of the four different

constructs, we calculated the proportion of fragments categorized as significantly active by

MPRAnalyze (**Figure S2B**; FDR < 0.05): 7,724 (of 11,656 total fragments tested, 66.27%) downstream

of the *INS* promoter, 5,596 (48.01%) upstream of the *INS* promoter, 6,337 (54.37%) downstream of the

SCP1 promoter, and 4,680 (40.15%) upstream of the SCP1 promoter. These proportions are

comparable to those calculated when using all fragments without requiring their representation across

all four configurations (**Table S6**). Of the set of 11,656 fragments, only 826 (7.09%) were significantly

active across all four configurations (**Figure S2C**). We also defined sets of fragments that were

uniquely active in a single configuration representing a collective 2,743 fragments (23.53%): 1,114

(9.56%) downstream of the *INS* promoter, 543 (4.66%) upstream of the *INS* promoter, 686 (5.89%)

downstream of the SCP1 promoter, and 400 (3.43%) upstream of the SCP1 promoter (**Figure S2C**).

Taken together, these initial analyses indicate that each MPRA construct influences measured

enhancer activity in a unique manner where most fragments were significantly active across only one or

two constructs of the four tested.

Quantifying promoter and position effects

We systematically quantified effects of each promoter and position on an individual fragment

basis using a linear modeling approach in MPRAnalyze. We modeled the activity of each fragment with

position and promoter as covariates, then performed a Wald test between each contrast (SCP1 vs. *INS*

and upstream vs. downstream) to extract model coefficients (**Tables S7-8**). Overall, we identified

698/11,656 fragments (5.99%) with significant promoter effects (**Figure 2B**; *FDR* < 0.05) and

703/11,656 fragments (6.03%) with significant upstream/downstream position effects. Between the two

groups, only 74 fragments display both promoter and position effects. While fragments with significant

position effects were roughly divided between upstream- and downstream-bias, we observed that

73.35% of fragments with significant promoter effects were biased towards the *INS* promoter (n =

512/698; p = 2.2 x 10^{-16}, binomial test with 55% expectation, or the proportion of fragments active with

the *INS* promoter). The sequence content of the SCP1 promoter is markedly different from the *INS*

promoter used here. SCP1 is an 81-bp sequence composed of four highly conserved core promoter

motifs (Juven-Gershon et al. 2006): a TATA box, the initiator element (Inr), a motif ten element (MTE),

and the downstream promoter element (DPE). SCP1 is known to act as a strong promoter for

enhancers in conventional reporter constructs, and was used in one of the first genome-wide pooled

reporter assays, STARR-seq (Arnold et al. 2013). By contrast, we used a 408-bp fragment of the *INS*

promoter spanning -364 to +44 bp of the human *INS* gene (German et al. 1995) which contains a

battery of pancreas-expressed TF binding sites (*e.g.,* RFX6, NEUROD1, INSM1) (Chandra et al. 2014)

and designated glucose-responsive elements (Melloul et al. 2002). We hypothesized that specific

sequence features of the *INS* promoter may enable regulatory compatibility with a broader subset of

diabetes and related trait-associated fragments.

Sequence features associated with position- or promoter-bias

Next, we were interested in defining specific sequence features associated with promoter- or position-driven activity bias using a set of genomic annotations, including transcription factor motifs, accessible chromatin peaks measured in human pancreatic islets, and ChromHMM-based chromatin state inferences derived from genomic profiling in islets (**Figure 3A**). Given that this set of annotations is large (~3,000 features), we anticipate that only a small subset is associated with activity bias. To address this challenge, we used LASSO regression, an extension of linear regression that penalizes and thereby removes uninformative predictors, resulting in a simple, interpretable model. For each unique fragment sequence, we calculated overlap scores across all annotations which we standardized for use as model predictors using rank-based inverse normalization. We performed LASSO regression with 10-fold cross validation, where the annotation overlap scores were the predictors and the position or promoter bias scores (i.e., signed Wald statistics from the previous joint analysis) were the outcome variables. To account for redundancy in TF motifs, we first performed LASSO using a clustered set of 540 motifs (D'Oliveira Albanus et al. 2021) and compared to results using the full set of annotations (**Tables S9-12**). Only four features showed significant association with promoter bias, with all four enriched in fragments with *INS* promoter-biased activity versus SCP1 (**Figure 3B**). Notably, two of these were motifs for HNF family transcription factors, which play important roles in pancreatic beta cell development, differentiation, and homeostasis, and when disrupted cause maturity onset diabetes of the young (MODY), an early onset monogenic form of diabetes (Nkonge et al. 2020; Zhang et al. 2021). Concordantly, many noncoding variants associated with complex, later-onset diabetes (i.e., T2D) are enriched in HNF1 motifs (Quang et al. 2015). Additionally, the *INS* promoter itself overlaps a putative HNF1 motif; however, there is conflicting evidence as to whether HNF1A/HNF1B directly mediates *INS* transcription (Okita et al. 1999; Párrizas et al. 2001; Emens et al. 1992). Together this regulatory convergence suggests that these two TFs serve as critical regulators of transcriptional activity central to diabetes pathogenesis.

When performing LASSO with the unclustered set of motifs, we additionally found that motifs for NF-κB constituents (RELA/p65, NFKB) and the ubiquitously expressed activator TFs SP1 and SRF

were associated with SCP1 promoter-biased fragment enhancer activity (**Figure S3A**). SP1 functions

by interacting with the transcription factor II D complex ($TF_{II}D$) (Pugh and Tjian 1990), a constituent of

the RNA polymerase II preinitiation complex that binds to core sequences found in the SCP1 promoter

(Louder et al. 2016). HINFP (also known as MIZF), a transcriptional repressor (Sekimata and Homma

2004; Sekimata et al. 2001), was the strongest coefficient associated with SCP1 promoter-biased

activity. In addition to the annotations associated with *INS* promoter-biased activity in the reduced set,

we found that several GATA family TFs and the SOX family member SRY motifs were strongly

predictive of fragments' bias towards the *INS* promoter, as was the motif for NKX6.1, another key

pancreatic beta cell identity TF. Similar to *HNF1A*/*HNF1B*, coding variants in *NKX6-1* have more

recently been explored as very rare causes of MODY (<1%) (Mohan et al. 2018; Taylor et al. 2013) and

T2D-associated variants are also enriched in NKX6.1 motifs (Chiou et al. 2021; Khetan et al. 2021,

2018).

We applied the same approach to define features associated with upstream/downstream

position bias (**Figure 3C; Figure S3B**). Of the significantly associated features, most (32/46) exhibited

bias towards upstream-cloned fragments. Features predictive of upstream bias included overlap with

open chromatin peaks in pancreatic islets ("atac_seq"), motifs for the transcriptional activator NF-Y

(which binds to CCAAT motifs in promoters) (Dolfini et al. 2012) and several TFs (e.g., CUX1, ISX,

POU family members, and PAX family members) known to bind to promoters of target genes important

for developmental processes and fate specification (Vadnais et al. 2013). The strongest predictor of

upstream-biased activity was presence of the "Active_TSS," an annotation denoting active promoters

as defined by ChromHMM (Ernst and Kellis 2012). Conversely, many fewer annotations were

associated with downstream-biased enhancer activity, though among these included the

"typicalEnhancer" ChromHMM state, which overlaps conventional enhancer marks such as p300 or

H3K27ac (Vahedi et al. 2015).

Evaluating effects of HNF1 motif perturbations on regulatory activity

To explore the putative interaction between the HNF1 motif within enhancer regions and the *INS*

promoter, we designed and constructed a targeted MPRA library (**Figure 4A; Figure S1B**). We

selected corresponding variants in 47 fragments with significant *INS* promoter-biased activity (FDR <

0.05) containing HNF1 motifs. To generate motif deletion or shuffled fragments, we first identified the

most likely motif position using SNP-aware position weight matrix scanning. For motif deletion

fragments, we removed the matched sequence and added (*n* motif bp)/2 of surrounding sequence to

each end of the fragment to keep the fragment length constant. For motif shuffled fragments, we

performed a dinucleotide shuffle on the matched motif sequence to disrupt the motif while maintaining

the same sequence content. In cases where the variant was adjacent to the HNF1 motif (rather than

contained within), we synthesized motif deletion and motif shuffled fragments with both alleles. We

cloned this set of fragments into an MPRA vector upstream of the *INS* promoter then delivered this

library to 832/13 cells (n = 6), as described previously.

Of the 47 fragment sets we synthesized and tested, 31 sets had at least one fragment with

significant activity (FDR < 0.05) and were retained for downstream analysis (**Table S13**). One such

variant was rs1635852, which is found in the first intron of *JAZF1*, a well-characterized transcriptional

regulator of the pancreatic beta cell response to metabolic stress (Kobiita et al. 2020). This variant falls

in a high-information content position of the HNF1 consensus motif (JASPAR record MA0046.2), where

changing from the reference allele (T, risk allele for T2D) to the alternate allele (C, non-risk) is predicted

to strengthen the motif (**Figure 4B**). We did not observe a statistically significant allelic difference in

activity between the reference and alternate alleles, likely owing to high variance in the alternate allele

fragment (Wilcoxon rank-sum test p = 0.22); however, the effect is in the expected direction (Fogarty et

al. 2013). We did observe that shuffling the HNF1 motif led to a significant decrease in activity

compared to either the risk (Wilcoxon rank-sum test p = 0.01587) or non-risk allele fragments (**Figure**

4C; Wilcoxon rank-sum test p = 0.007937). Deleting the HNF1 motif led to significantly decreased

activity compared to the non-risk allele alone (Wilcoxon rank-sum test p = 0.01587). To evaluate the

extent to which these effects were restricted to the *INS* promoter, we also tested these fragments with

the SCP1 promoter. We observed no significant differences in activity based on HNF1 motif status

(**Figure S4A**). A previous study reported higher activity for the alternative compared with the reference allele using reporter and DNA affinity assays (Fogarty et al. 2013), consistent with the direction of effect we observed. Interestingly, this study concluded that regulatory activity at this locus was mediated by differential interaction with another key transcription factor expressed in beta cells, PDX1. In contrast to the HNF1 motif, the rs1635852 risk T allele is slightly preferred over the non-risk C allele in the PDX1 motif (JASPAR record MA0132.2). Because our scrambled HNF1 motif oligo also partially disrupts the PDX1 motif, this assay may also be detecting changes to PDX1 binding. Our results point to a possible complementary regulatory mechanism at this locus mediated by HNF1A/B. We again observed HNF1 motif-dependent activity at rs11819995, a T2D signal variant recently identified in a multi-ancestry GWAS by the DIAMANTE consortium (Mahajan et al. 2022). This variant is located in the first intron of the canonical *ETS1* transcript and overlaps an alternative transcription start site active in pancreatic islets (Varshney et al. 2017; Mahajan et al. 2022). The 14-bp HNF1 motif matched by position weight matrix scanning (*GTTAAACTTTAAAC*) starts 11 bp downstream of the variant in the fragment sequence (**Figure 4D**). Here, we observed a modest allelic effect in fragments with intact HNF1 motifs (Wilcoxon rank-sum $p = 0.09524$). Deleting the HNF1 motif significantly decreased the regulatory activity of fragments, in the presence of either the reference (C, non-risk allele for T2D; Wilcoxon rank-sum test $p = 0.007937$) or alternate allele (T, risk; Wilcoxon rank-sum test $p = 0.03175$) of this variant (**Figure 4E**). Additionally, shuffling the HNF1 motif in fragments with either allele decreased activity in comparison to the original fragment (Wilcoxon rank-sum test $p = 0.007937$). As with the previously described variant, we tested fragments spanning rs11819995 in a construct with the SCP1 promoter. None of these fragments were differentially active based on whether the motif was intact, deleted, or shuffled (**Figure S4B**).

To further explore HNF1 motif-*INS* promoter interaction, we considered two related but distinct questions: (1) does perturbing the HNF1 motif have similar effects when cloned in proximity to a different tissue-specific promoter, and (2) is the interaction between the HNF1 motif and *INS* promoter restricted to a pancreatic beta cell line? To these ends, we cloned the same library of original, deleted, or shuffled fragments in one of two new promoter contexts, with either the SCP1 promoter or with the

promoter for the skeletal muscle-specific gene *MYBPC2*. We then delivered these two libraries along

with the *INS* promoter MPRA library to the LHCN-M2 human skeletal muscle myoblast line (n = 6) after

differentiation to myotubes (**Figure 4F**). In this line, *MYBPC2* is abundantly expressed, and *INS* is

transcriptionally silent.

In the LHCN-M2 cell line, none of the HNF1 motif fragment sets displayed significant enhancer

activity ($\log_2$(RNA/DNA) > 0 and FDR < 0.05) when paired with any of the three different promoters

(**Figure 4G**). Additionally, we observed that their activity was modulated by promoter context, where the

sequences tended to be more active when cloned with the muscle-active *MYBPC2* promoter versus

SCP1 or *INS1*. Finally, we observed only one instance in which disrupting the HNF1 motif led to

modulated regulatory activity. Compared to the original fragment, scrambling the HNF1 motif caused a

modest increase in enhancer activity when paired with the *INS* promoter (Wilcoxon rank-sum test p =

0.015); however, HNF1A and HNF1B are not expressed in skeletal muscle, so enhancer activity of this

fragment is likely driven by a separate transcription factor. In all other cases across all three promoter

pairings, we no longer observed any dependence upon the HNF1 motif. Taken together, this indicates

HNF1A/B-orchestrated regulatory activity is beta cell-specific.

Discussion

Massively parallel reporter assays (MPRAs) enable large-scale variant-to-function studies of cis-

regulatory elements. However, MPRA implementations differ in key technical aspects including the

cloning and configuration of the reporter construct, whether it is integrated or episomal, and which basal

promoter it carries. Differences in sequence and cellular context can influence the activity of the cloned

fragments (Klein et al. 2020; Bergman et al. 2022); however, given these screens' scale, it is not routine

practice to examine how each of these factors influence individual fragments' activity or how they may

modulate the effects of genetic variation.

In this study, we used a modular MPRA construct to explore the effects of sequence context

and proximity upon regulatory elements' activity. We found that for a subset of disease-associated

genomic sequences, enhancer activity was sensitive to proximity and identity of the promoter used,

independent of overall activity. We found specific genomic annotations that were associated with these

biases and noted that direction of effect was concordant with those interpretations. For instance,

fragments annotated as proximal transcription start sites by ChromHMM tended to be more active when

cloned upstream, and conversely, those annotated as enhancers were more active when cloned

downstream. Of note, we discovered that sequences preferentially active with the human insulin (*INS*)

promoter were enriched in motifs for the important pancreatic beta cell transcription factors including

HNF1A/B and NKX6.1. In a follow-up MPRA, we demonstrated the necessity of HNF1A/B motifs for

activity of a subset of these fragments and confirmed that the dependence upon those motifs was

specific to both the cell type and the paired promoters.

Our observation that some sequences' activity depends upon proximity to a promoter is

consistent with many previous reports. Enhancers are known to work at a range of distances, including

1,000s of kb away from their targets. Here, we identified that sequences that were more active when

placed downstream (distal to) the promoter and reporter gene tended to be enriched for features of

typical enhancers while those that were more active in the upstream position (proximal) contained

promoter-like elements including enrichment for active TSSs. It is important to note that the MPRA

construct we used in this study to explore positional effects is distinct from the STARR-seq vector in

that fragments are cloned downstream of the poly-A signal and thus are not transcribed (Klein et al.

2020; Das et al. 2023). Therefore, we do not suspect that interaction with RNA-binding proteins or

accelerated RNA decay influenced the results observed here.

Previous studies using reporter assays have indicated that core promoter sequences display

some level of tissue-specificity (Mattioli et al. 2019), where observed activity in an MPRA vector is

strongly correlated with endogenous transcriptional activity from the same loci. This has also been

observed by others working to design minimal synthetic promoters that can function in a context-

dependent manner, including with respect to cell type (Zahm et al. 2024). Together with our work, this

suggests that promoter sequences contain a unique syntax that interacts with a cell's regulatory milieu

to produce highly tuned gene expression beyond the standard elements common to many promoters

(e.g., TATA box, downstream promoter element, etc.). It is important to note that we used a well-

delimited promoter for the highly cell type-specific human insulin gene that was previously defined

through deletion mapping and motif analysis (German et al. 1995; Melloul et al. 2002). Selecting

suitable promoters for other cell types will require a similar level of inspection.

Though not explicitly considered in the work presented here, the choice of using an episomal or

integrated MPRA vector has also been examined previously (Inoue et al. 2017; Klein et al. 2020;

Guzman et al. 2023). Here, we used an integrated construct to compare activity at a specific locus in

the LHCN-M2 human skeletal muscle cell line compared to an electroporated episomal construct in the

832/13 rat beta cell line (**Figure 4D-E**), due to technical limitations that make electroporation or

transfection of differentiated LHCN-M2 cells unfeasible. While we see differences in enhancer activity

across the two lines, we believe that this is a biological distinction and not a methodological one. An

additional source of concern for episomal MPRAs is preferential usage of the bacterial origin of

replication (ORI) compared to the SCP1 promoter (Muerdter et al. 2018; Klein et al. 2020).

Consequently, researchers have advised to use other synthetic promoters (e.g., minP) or ORI-less

versions of the MPRA backbone to avoid this issue. Given the systematic discrepancies measured

between the two methods in the studies referenced above, this design choice may also require careful

forethought.

Although our initial, large MPRA library contained both reference and alternate alleles across a

large set of T2D-associated variants, it was not cloned at sufficient complexity to allow for sensitive

detection of allelic effects, which are often subtle. We therefore focused our analyses on fragment-level

activity across position and promoter contexts. This approach complements prior allele-focused

MPRAs, such as one published in 2021 by Khetan and colleagues (Khetan et al. 2021) that

characterized over 2,000 T2D-associated variants. Given that the number of T2D-associated genomic

loci has more than quadrupled since these libraries were constructed (Suzuki et al. 2024; Mahajan et

al. 2022; Vujkovic et al. 2020), it is a high priority to develop new MPRA libraries to profile the

expanded set of loci.

The work presented here indicates that in MPRA experiments, proximity to the promoter and the

choice of promoter each have measurable impacts which may vary across different fragments. Though

it takes consideration to select appropriate promoters for a cell type of interest, we show that for

pancreatic islet beta cells using the *INS* promoter was sufficient to capture regulatory activity at many

T2D-associated loci. Given the inherent sparsity of MPRA datasets, employing these libraries across a

variety of design configurations and cell types will enhance our ability to understand regulatory

processes that are disrupted in complex diseases.

Methods

Cell culture

We obtained INS-1 832/13 rat insulinoma cells from Dr. Christopher Newgard (Sarah W. Stedman

Nutrition and Metabolism Center, Duke University, Durham, NC) and LHCN-M2 human skeletal muscle

myoblasts from Evercyte. We cultured 832/13 cells in RPMI-1640 containing 10% fetal bovine serum,

0.05 mM β-mercaptoethanol, 1 mM sodium pyruvate, 2 mM L-glutamine, 10 mM HEPES, and 1,000

U/mL penicillin/streptomycin. We maintained LHCN-M2 cells on 0.1% gelatin-coated flasks in 4:1

DMEM (4.5 g/L glucose, glutamine, bicarbonate):Medium 199 (with bicarbonate) containing 15% FBS,

20 mM HEPES, 30 ng/mL zinc sulfate, 1.4 ug/mL vitamin B12, 55 ng/mL dexamethasone, 2.5 ng/mL

recombinant human hepatocyte growth factor, 10 ng/mL basic fibroblast growth factor, and 600 U/mL

penicillin/streptomycin.

Large (first) MPRA library design, construction, and delivery

We compiled regions of interest for the large MPRA library from loci associated with type 2 diabetes

(T2D) and related metabolic traits, drawing on signal variants reported by the DIAMANTE Consortium

(Mahajan et al. 2022) and signal variants for HbA1C, fasting insulin adjusted for BMI, fasting glucose

adjusted for BMI, and 2-hr glucose adjusted for BMI reported by the MAGIC Consortium (Chen et al.

2021). To provide broader disease context, we also included loci for non-T2D traits including

rheumatoid arthritis, systemic lupus erythematosus, and type 1 diabetes. We used UK Biobank imputed

genotypes to calculate linkage disequilibrium and fetch proxies with an R^2 greater than 0.8. We

removed any variants that were not SNVs (i.e., indels, CNVs) resulting in a set of 13,226 sites. For

each site, we designed 198-bp fragment sequences in three positional configurations relative to the

focal base: offset left (position 50 of 198 bp), centered (position 99), or offset right (position 149). In

cases where a variant had two alternate alleles, we designed both sets of sequences. We extracted all

sequences from the hg19 reference genome. We added 16-bp flanking adapter sequences for PCR

amplification and cloning. We obtained a pool of 79,511 full-length 230-bp fragments from Agilent.

We constructed the large MPRA library as described previously (Varshney et al. 2021) (**Figure S1A**).

We added homology arms via PCR, then cloned amplified fragments into either a KpnI- or EcoRV-

digested (upstream or downstream, respectively) MPRA backbone via HiFi assembly. We transformed

this first assembly into electrocompetent 10-beta cells. We post-barcoded by digesting the plasmid

library with PmeI then used HiFi assembly to insert 16-bp random barcodes at the PmeI restriction site.

We transformed this reaction into electrocompetent 10-beta cells and prepared final plasmid libraries

for transfection using the ZymoPURE Plasmid Maxiprep Kit.

For the first library, we electroporated 50 ug of plasmid into 25 million 832/13 rat insulinoma cells for

each biological replicate and lysed cells with TRIzol 24 hours later. After phase separation, we used the

Direct-zol RNA Miniprep Kit to isolate total RNA. We enriched for mature mRNA transcripts with

fragment(dT) beads then treated 2 ug of mRNA with DNase I to remove plasmid or genomic DNA

contamination. We reverse transcribed 1 ug of mRNA using SuperScript III with a custom primer

containing a 6-bp unique molecular identifier (UMI) targeting transcripts derived from the MPRA

plasmids (**Table S14; Figure S1C**). We treated cDNA samples with DpnI to eliminate any residual

plasmid DNA.

Motif (second) MPRA library design, construction, and delivery

We constructed the motif MPRA library using a modified version of the methods detailed in Tewhey, et

al. 2016. While the cloning strategy differed slightly from what we used for the first library, the assay

design and functional readout between the two are comparable (**Figure S1B**). We selected fragment

sequences from the first library with strong *INS* promoter-biased activity that contained HNF1 motifs.

We obtained fragments spanning the same amount of genomic sequence as the previous library but

containing different adapters (IDT). We added barcodes and a promoter cloning scaffold via PCR, then

cloned the amplified fragments into a modified pMPRA1 vector (a gift from Tarjei Mikkelsen; Addgene

#49349) using Golden Gate assembly with PaqCI. We digested this assembly with SfiI to remove

empty backbones then transformed into electrocompetent 10-beta cells. We inserted clonal promoter

fragments using Golden Gate assembly and digested with AsiSI to remove promoterless constructs.

We transformed this final assembly into electrocompetent 10-beta cells, expanded in 150 mL cultures,

and isolated plasmid using the ZymoPURE Plasmid Maxiprep Kit.

For the second library, we electroporated 30 ug of plasmid into 30 million 832/13 cells. We lysed cells

24 hours later with Buffer RLT Plus and isolated total RNA with the Qiagen RNeasy Midi Kit. We

reverse transcribed 30 ug of DNAse I-treated total RNA using SuperScript IV (**Figure S1B**). We

performed a cDNA amplification PCR and sample indexing with all cDNA made.

For use with the LHCN-M2 skeletal muscle myocyte cell line, we ported the assembled MPRA block

(fragment, barcode, promoter, GFP) to a lentiviral transfer vector (a gift from Nadav Ahituv; Addgene

#137725) via restriction cloning (Gordon et al. 2020; Inoue et al. 2017). The University of Michigan Viral

Vector Core produced infectious lentiviral particles with this transfer vector and third generation

lentiviral plasmids in HEK293T cells. Per replicate, we infected 4×10^6 LHCN-M2 human skeletal

myoblasts with our MPRA library at an MOI of ~10. After infection, we passaged the cells for one week

to remove any unincorporated virus or contaminating transfer plasmid, then differentiated the cells for

one week. We isolated RNA and gDNA from each replicate using the Qiagen AllPrep DNA/RNA mini

kit.

For all samples generated through use of the second set of MPRA libraries, we reverse transcribed

RNA into cDNA with a GFP-specific primer containing a 6-bp UMI, then constructed indexed

sequencing libraries for both the cDNA and gDNA libraries using Illumina-compatible primers (**Table**

S15).

Sequencing data collection

We sequenced pairing and cDNA/input sequencing libraries for the initial set of MPRA studies on an

Illumina Hiseq. We sequenced libraries generated for the second set of MPRA on an Illumina Novaseq

6000. For both sets of studies, we received paired-end 150-bp reads.

To create the barcode-fragment pairing dictionary, we used a custom pipeline. First, we merged the

paired-end 150-bp reads using bwa v0.7.17 then extracted barcode and fragment sequences using

cutadapt v4.3 (Li and Durbin 2009; Martin 2011). To align the merged fragment reads against our

reference FASTA file, we used minimap2 v2.24 (Li 2018). After filtering for mapping quality and

sequencing depth, we created a final table with fragment-barcode pairs and removed any duplicate

barcodes.

We processed MPRA sequencing data using a custom pipeline that quantifies barcode counts while

accounting for possible sequencing errors. We identified reads with the barcode sequence and trimmed

constant sequences using cutadapt v4.3. We used UMI-tools v1.1.2 to identify and cluster UMIs, then

used starcode-umi to cluster UMI-barcode and perform deduplication (Zorita et al. 2015; Smith et al.

2017). Finally, we used starcode to cluster and count barcodes.

Activity estimation and joint modeling

Prior to model fitting, we merged cDNA and input barcode counts with the corresponding pairing

dictionary requiring an exact match. For both MPRA libraries, we used the R package MPRAnalyze to

estimate enhancer activity and promoter or position bias (Ashuach et al. 2019). This modeling approach

fits two separate models for cDNA and input counts, then relates the two to yield a per-fragment

transcription rate. For all analyses, resulting enhancer activity scores were converted from a natural log

scale to binary log scale, though values are reported on both scales in Tables S2-5.

For the first MPRA library, we used this method to estimate enhancer activity within each construct type

and then performed *post hoc* comparisons outside of the analysis framework. To evaluate quantitative

effects of promoter and position on fragment activity in the first MPRA library, we jointly estimated

enhancer activity across all four constructs and included promoter and position as covariates.

For the second MPRA library, we used this method to estimate enhancer activity across the entire

library within the two cell types (832/13 and LHCN-M2) and nominate candidate fragments for further

inspection. Then, on a per fragment basis, we compared activity differences across fragment types

using a Wilcoxon rank-sum test.

LASSO regression to select features associated with promoter or position bias

Here we used least absolute shrinkage and selection operator (LASSO) regression to model MPRA

activity bias (promoter or position) and select genomic features associated with this bias (Tibshirani

1996). LASSO regression uses a penalty term to shrink coefficients of less informative variables to

zero, thereby removing them from the model and enhancing interpretability. We quantified activity bias

by using the Wald statistics from our joint estimates of enhancer activity and multiplying them by the

sign of the log-fold change to indicate direction of bias (INS versus SCP1 for promoter and up- versus

downstream for position). As features in our models, we used a set of annotations and TF motifs

described previously (Varshney et al. 2021; D'Oliveira Albanus et al. 2021). For each fragment, we

quantified motif occurrences or annotation overlap. To generate overlap scores that could be used as

parameters for LASSO regression and compared uniformly across annotations, we inverse-normalized

the $-\log_{10}$(p-value) for each motif or annotation overlap using R/RNOmni (McCaw et al. 2020). We ran

the regression using R/glmnet with default parameters ($\alpha = 1$ corresponding to LASSO) and 10-fold

cross validation using the signed Wald statistic as the outcome and annotation scores as the predictors

(Friedman et al. 2010). We performed regression analyses for the promoter and position bias scores

separately and glmnet automatically determined the appropriate lambda value by minimizing the mean

cross-validated error.

Data and code access

All processed sequencing data generated in this study have been submitted to the NCBI Gene

Expression Omnibus (GEO) under accession numbers GSE279057 (first library, GSE279071 (second

library), and GSE247455 (LHCN-M2 data). Corresponding raw sequencing data is available on the

NCBI Sequencing Read Archive under accession numbers PRJNA1142481 and PRJNA1142558.

Complete results files are available as part of this manuscript as Supplementary Tables.

Custom scripts used to preprocess and analyze data from the second MPRA library and other auxiliary

analyses are available at https://github.com/adelaidetovar/modular-t2d-mpra.

Competing Interest Statement

The authors declare they have no conflicts of interest.

Acknowledgements

The authors would like to acknowledge members of the Kitzman and Parker labs (University of

Michigan) for their critical feedback. The authors would also like to thank the University of Michigan

Viral Vector Core for producing lentivirus and the University of Michigan Advanced Genomics Core for

sequencing services.

Funding

The authors acknowledge support from the National Institute of Diabetes and Digestive and Kidney

Diseases, grants 1UM1DK126185-01 (S.C.J.P.), R01 DK117960 (S.C.J.P.), Opportunity Pool Funding

(A.T., S.C.J.P., J.O.K.), and T32DK101357 (A.T.); the National Institute of General Medical Studies

Contributions

5 Conceptualization by A.T., Y.K., J.O.K., and S.C.J.P. Fragment design and MPRA library cloning by

6 Y.K., A.T., and K.N. Data processing and analysis by A.T., Y.K., and M.B., with contributions from A.V.

7 A.T. wrote the manuscript with input from and editing by J.O.K. and S.C.J.P.

References

Arnold CD, Gerlach D, Stelzer C, Boryń ŁM, Rath M, Stark A. 2013. Genome-wide quantitative enhancer activity maps identified by STARR-seq. *Science* **339**: 1074–1077.

Ashuach T, Fischer DS, Kreimer A, Ahituv N, Theis FJ, Yosef N. 2019. MPRAnalyze: statistical framework for massively parallel reporter assays. *Genome Biol* **20**: 183.

Banerji J, Rusconi S, Schaffner W. 1981. Expression of a beta-globin gene is enhanced by remote SV40 DNA sequences. *Cell* **27**: 299–308.

Bergman DT, Jones TR, Liu V, Ray J, Jagoda E, Siraj L, Kang HY, Nasser J, Kane M, Rios A, et al. 2022. Compatibility rules of human enhancer and promoter sequences. *Nature* **607**: 176–184.

Chandra V, Albagli-Curiel O, Hastoy B, Piccand J, Randriamampita C, Vaillant E, Cavé H, Busiah K, Froguel P, Vaxillaire M, et al. 2014. RFX6 regulates insulin secretion by modulating Ca2+ homeostasis in human β cells. *Cell Rep* **9**: 2206–2218.

Chen J, Spracklen CN, Marenne G, Varshney A, Corbin LJ, Luan J, Willems SM, Wu Y, Zhang X, Horikoshi M, et al. 2021. The trans-ancestral genomic architecture of glycemic traits. *Nat Genet* 1–21.

Chiou J, Zeng C, Cheng Z, Han JY, Schlichting M, Miller M, Mendez R, Huang S, Wang J, Sui Y, et al. 2021. Single-cell chromatin accessibility identifies pancreatic islet cell type- and state-specific regulatory programs of diabetes risk. *Nat Genet* **53**: 455–466.

Choi J, Zhang T, Vu A, Ablain J, Makowski MM, Colli LM, Xu M, Hennessey RC, Yin J, Rothschild H, et al. 2020. Massively parallel reporter assays of melanoma risk variants identify MX2 as a gene promoting melanoma. *Nat Commun* **11**: 2718.

Das M, Hossain A, Banerjee D, Praul CA, Girirajan S. 2023. Challenges and considerations for reproducibility of STARR-seq assays. *Genome Res* **33**: 479–495.

Dolfini D, Gatta R, Mantovani R. 2012. NF-Y and the transcriptional activation of CCAAT promoters. *Crit Rev Biochem Mol Biol* **47**: 29–49.

D'Oliveira Albanus R, Kyono Y, Hensley J, Varshney A, Orchard P, Kitzman JO, Parker SCJ. 2021. Chromatin information content landscapes inform transcription factor and DNA interactions. *Nat Commun* **12**: 1307.

Emens LA, Landers DW, Moss LG. 1992. Hepatocyte nuclear factor 1 alpha is expressed in a hamster insulinoma line and transactivates the rat insulin I gene. *Proc Natl Acad Sci U S A* **89**: 7300–7304.

Ernst J, Kellis M. 2012. ChromHMM: automating chromatin-state discovery and characterization. *Nat Methods* **9**: 215–216.

Fogarty MP, Panhuis TM, Vadlamudi S, Buchkovich ML, Mohlke KL. 2013. Allele-specific transcriptional activity at type 2 diabetes-associated single nucleotide polymorphisms in regions of pancreatic islet open chromatin at the JAZF1 locus. *Diabetes* **62**: 1756–1762.

Friedman JH, Hastie T, Tibshirani R. 2010. Regularization Paths for Generalized Linear Models via Coordinate Descent. *J Stat Softw* **33**: 1–22.

German M, Ashcroft S, Docherty K, Edlund H, Edlund T, Goodison S, Imura H, Kennedy G, Madsen O, Melloul D. 1995. The insulin gene promoter. A simplified nomenclature. *Diabetes* **44**: 1002–

1004.

Gordon MG, Inoue F, Martin B, Schubach M, Agarwal V, Whalen S, Feng S, Zhao J, Ashuach T, Ziffra
R, et al. 2020. lentiMPRA and MPRAflow for high-throughput functional characterization of gene
regulatory elements. *Nat Protoc* **15**: 2387–2412.

Guzman C, Duttke S, Zhu Y, De Arruda Saldanha C, Downes NL, Benner C, Heinz S. 2023. Combining
TSS-MPRA and sensitive TSS profile dissimilarity scoring to study the sequence determinants
of transcription initiation. *Nucleic Acids Res*. http://dx.doi.org/10.1093/nar/gkad562.

Hong CKY, Cohen BA. 2022. Genomic environments scale the activities of diverse core promoters.
Genome Res **32**: 85–96.

Inoue F, Kircher M, Martin B, Cooper GM, Witten DM, McManus MT, Ahituv N, Shendure J. 2017. A
systematic comparison reveals substantial differences in chromosomal versus episomal
encoding of enhancer activity. *Genome Res* **27**: 38–52.

Juven-Gershon T, Cheng S, Kadonaga JT. 2006. Rational design of a super core promoter that
enhances gene expression. *Nat Methods* **3**: 917–922.

Khetan S, Kales S, Kursawe R, Jillette A, Ulirsch JC, Reilly SK, Ucar D, Tewhey R, Stitzel ML. 2021.
Functional characterization of T2D-associated SNP effects on baseline and ER stress-
responsive β cell transcriptional activation. *Nat Commun* **12**: 5242.

Khetan S, Kursawe R, Youn A, Lawlor N, Jillette A, Marquez EJ, Ucar D, Stitzel ML. 2018. Type 2
Diabetes-Associated Genetic Variants Regulate Chromatin Accessibility in Human Islets.
Diabetes **67**: 2466–2477.

Klein JC, Agarwal V, Inoue F, Keith A, Martin B, Kircher M, Ahituv N, Shendure J. 2020. A systematic
evaluation of the design and context dependencies of massively parallel reporter assays. *Nat
Methods* **17**: 1083–1091.

Kobiita A, Godbersen S, Araldi E, Ghoshdastider U, Schmid MW, Spinas G, Moch H, Stoffel M. 2020.
The Diabetes Gene JAZF1 Is Essential for the Homeostatic Control of Ribosome Biogenesis
and Function in Metabolic Stress. *Cell Rep* **32**: 107846.

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**: 3094–3100.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics **25**: 1754–1760.

Louder RK, He Y, López-Blanco JR, Fang J, Chacón P, Nogales E. 2016. Structure of promoter-bound
TFIID and model of human pre-initiation complex assembly. *Nature* **531**: 604–609.

Lu X, Chen X, Forney C, Donmez O, Miller D, Parameswaran S, Hong T, Huang Y, Pujato M, Cazares
T, et al. 2021. Global discovery of lupus genetic risk variant allelic enhancer activity. *Nat
Commun* **12**: 1611.

Mahajan A, Spracklen CN, Zhang W, Ng MCY, Petty LE, Kitajima H, Yu GZ, Rüeger S, Speidel L, Kim
YJ, et al. 2022. Multi-ancestry genetic study of type 2 diabetes highlights the power of diverse
populations for discovery and translation. *Nat Genet* **54**: 560–572.

Mahajan A, Taliun D, Thurner M, Robertson NR, Torres JM, Rayner NW, Payne AJ, Steinthorsdottir V,
Scott RA, Grarup N, et al. 2018. Fine-mapping type 2 diabetes loci to single-variant resolution
using high-density imputation and islet-specific epigenome maps. *Nat Genet* **50**: 1505–1513.

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal **17**: 10–12.

Martinez-Ara M, Comoglio F, van Arensbergen J, van Steensel B. 2022. Systematic analysis of intrinsic
enhancer-promoter compatibility in the mouse genome. *Mol Cell* **82**: 2519-2531.e6.

Mattioli K, Volders P-J, Gerhardinger C, Lee JC, Maass PG, Melé M, Rinn JL. 2019. High-throughput
functional analysis of lncRNA core promoters elucidates rules governing tissue specificity.
Genome Res **29**: 344–355.

McAfee JC, Bell JL, Krupa O, Matoba N, Stein JL, Won H. 2022. Focus on your locus with a massively
parallel reporter assay. *J Neurodev Disord* **14**: 50.

McAfee JC, Lee S, Lee J, Bell JL, Krupa O, Davis J, Insigne K, Bond ML, Zhao N, Boyle AP, et al.
2023. Systematic investigation of allelic regulatory activity of schizophrenia-associated common
variants. *Cell Genom* **3**: 100404.

McCaw ZR, Lane JM, Saxena R, Redline S, Lin X. 2020. Operating characteristics of the rank-based
inverse normal transformation for quantitative trait analysis in genome-wide association studies.
Biometrics **76**: 1262–1272.

Melloul D, Marshak S, Cerasi E. 2002. Regulation of insulin gene transcription. *Diabetologia* **45**: 309–
326.

Melnikov A, Murugan A, Zhang X, Tesileanu T, Wang L, Rogov P, Feizi S, Gnirke A, Callan CG Jr,
Kinney JB, et al. 2012. Systematic dissection and optimization of inducible enhancers in human
cells using a massively parallel reporter assay. *Nat Biotechnol* **30**: 271–277.

Mohan V, Radha V, Nguyen TT, Stawiski EW, Pahuja KB, Goldstein LD, Tom J, Anjana RM, Kong-
Beltran M, Bhangale T, et al. 2018. Comprehensive genomic analysis identifies pathogenic
variants in maturity-onset diabetes of the young (MODY) patients in South India. *BMC Med*
Genet **19**: 22.

Muerdter F, Boryń ŁM, Woodfin AR, Neumayr C, Rath M, Zabidi MA, Pagani M, Haberle V, Kazmar T,
Catarino RR, et al. 2018. Resolving systematic errors in widely used enhancer activity assays in
human cells. *Nat Methods* **15**: 141–149.

Myint L, Wang R, Boukas L, Hansen KD, Goff LA, Avramopoulos D. 2020. A screen of 1,049
schizophrenia and 30 Alzheimer's-associated variants for regulatory potential. *Am J Med Genet*
B Neuropsychiatr Genet **183**: 61–73.

Nkonge KM, Nkonge DK, Nkonge TN. 2020. The epidemiology, molecular pathogenesis, diagnosis,
and treatment of maturity-onset diabetes of the young (MODY). *Clin Diabetes Endocrinol* **6**: 20.

Okita K, Yang Q, Yamagata K, Hangenfeldt KA, Miyagawa J, Kajimoto Y, Nakajima H, Namba M,
Wollheim CB, Hanafusa T, et al. 1999. Human insulin gene is a target gene of hepatocyte
nuclear factor-1alpha (HNF-1alpha) and HNF-1beta. *Biochem Biophys Res Commun* **263**: 566–
569.

Párrizas M, Maestro MA, Boj SF, Paniagua A, Casamitjana R, Gomis R, Rivera F, Ferrer J. 2001.
Hepatic nuclear factor 1-alpha directs nucleosomal hyperacetylation to its tissue-specific
transcriptional targets. *Mol Cell Biol* **21**: 3234–3243.

Patwardhan RP, Hiatt JB, Witten DM, Kim MJ, Smith RP, May D, Lee C, Andrie JM, Lee S-I, Cooper
GM, et al. 2012. Massively parallel functional dissection of mammalian enhancers in vivo. *Nat*

Biotechnol **30**: 265–270.

Pugh BF, Tjian R. 1990. Mechanism of transcriptional activation by Sp1: evidence for coactivators. *Cell* **61**: 1187–1197.

Quang DX, Erdos MR, Parker SCJ, Collins FS. 2015. Motif signatures in stretch enhancers are enriched for disease-associated genetic variants. *Epigenetics Chromatin* **8**: 23.

Sahu B, Hartonen T, Pihlajamaa P, Wei B, Dave K, Zhu F, Kaasinen E, Lidschreiber K, Lidschreiber M, Daub CO, et al. 2022. Sequence determinants of human gene regulatory elements. *Nat Genet* **54**: 283–294.

Sekimata M, Homma Y. 2004. Sequence-specific transcriptional repression by an MBD2-interacting zinc finger protein MIZF. *Nucleic Acids Res* **32**: 590–597.

Sekimata M, Takahashi A, Murakami-Sekimata A, Homma Y. 2001. Involvement of a novel zinc finger protein, MIZF, in transcriptional repression by interacting with a methyl-CpG-binding protein, MBD2. *J Biol Chem* **276**: 42632–42638.

Smith T, Heger A, Sudbery I. 2017. UMI-tools: modeling sequencing errors in Unique Molecular Identifiers to improve quantification accuracy. *Genome Res* **27**: 491–499.

Spracklen CN, Horikoshi M, Kim YJ, Lin K, Bragg F, Moon S, Suzuki K, Tam CHT, Tabara Y, Kwak S-H, et al. 2020. Identification of type 2 diabetes loci in 433,540 East Asian individuals. *Nature* **582**: 240–245.

Suzuki K, Hatzikotoulas K, Southam L, Taylor HJ, Yin X, Lorenz KM, Mandla R, Huerta-Chagoya A, Melloni GEM, Kanoni S, et al. 2024. Genetic drivers of heterogeneity in type 2 diabetes pathophysiology. *Nature* **627**: 347–357.

Taylor BL, Liu F-F, Sander M. 2013. Nkx6.1 is essential for maintaining the functional state of pancreatic beta cells. *Cell Rep* **4**: 1262–1275.

Tewhey R, Kotliar D, Park DS, Liu B, Winnicki S, Reilly SK, Andersen KG, Mikkelsen TS, Lander ES, Schaffner SF, et al. 2016. Direct Identification of Hundreds of Expression-Modulating Variants using a Multiplexed Reporter Assay. *Cell* **165**: 1519–1529.

Tibshirani R. 1996. Regression shrinkage and selection via the lasso. *Journal of the royal statistical society series b-methodological* **58**: 267–288.

Ulirsch JC, Nandakumar SK, Wang L, Giani FC, Zhang X, Rogov P, Melnikov A, McDonel P, Do R, Mikkelsen TS, et al. 2016. Systematic Functional Dissection of Common Genetic Variation Affecting Red Blood Cell Traits. *Cell* **165**: 1530–1545.

Vadnais C, Awan AA, Harada R, Clermont P-L, Leduy L, Bérubé G, Nepveu A. 2013. Long-range transcriptional regulation by the p110 CUX1 homeodomain protein on the ENCODE array. *BMC Genomics* **14**: 258.

Vahedi G, Kanno Y, Furumoto Y, Jiang K, Parker SCJ, Erdos MR, Davis SR, Roychoudhuri R, Restifo NP, Gadina M, et al. 2015. Super-enhancers delineate disease-associated regulatory nodes in T cells. *Nature* **520**: 558–562.

Varshney A, Kyono Y, Elangovan VR, Wang C, Erdos MR, Narisu N, Albanus RD, Orchard P, Stitzel ML, Collins FS, et al. 2021. A transcription start site map in human pancreatic islets reveals functional regulatory signatures. *Diabetes*. http://dx.doi.org/10.2337/db20-1087.

Varshney A, Scott LJ, Welch RP, Erdos MR, Chines PS, Narisu N, Albanus RD, Orchard P, Wolford BN, Kursawe R, et al. 2017. Genetic regulatory signatures underlying islet gene expression and type 2 diabetes. *Proc Natl Acad Sci U S A* **114**: 2301–2306.

Vujkovic M, Keaton JM, Lynch JA, Miller DR, Zhou J, Tcheandjieu C, Huffman JE, Assimes TL, Lorenz K, Zhu X, et al. 2020. Discovery of 318 new risk loci for type 2 diabetes and related vascular outcomes among 1.4 million participants in a multi-ancestry meta-analysis. *Nat Genet* **52**: 680–691.

Watanabe K, Stringer S, Frei O, Umićević Mirkov M, de Leeuw C, Polderman TJC, van der Sluis S, Andreassen OA, Neale BM, Posthuma D. 2019. A global overview of pleiotropy and genetic architecture in complex traits. *Nat Genet* **51**: 1339–1348.

Zabidi MA, Arnold CD, Schernhuber K, Pagani M, Rath M, Frank O, Stark A. 2015. Enhancer-core-promoter specificity separates developmental and housekeeping gene regulation. *Nature* **518**: 556–559.

Zahm AM, Owens WS, Himes SR, Fallon BS, Rondem KE, Gormick AN, Bloom JS, Kosuri S, Chan H, English JG. 2024. A massively parallel reporter assay library to screen short synthetic promoters in mammalian cells. *Nat Commun* **15**: 10353.

Zhang H, Colclough K, Gloyn AL, Pollin TI. 2021. Monogenic diabetes: a gateway to precision medicine in diabetes. *J Clin Invest* **131**. http://dx.doi.org/10.1172/JCI142244.

Zorita E, Cuscó P, Filion GJ. 2015. Starcode: sequence clustering based on all-pairs search. *Bioinformatics* **31**: 1913–1919.

Figure Titles and Legends

Figure 1. Study design. We synthesized a library of 198-bp fragments encompassing 13,226 sites in

high linkage disequilibrium with type 2 diabetes (T2D) and related trait-associated signals. We cloned

this library into each of four MPRA constructs: fragment upstream or downstream of the human insulin

(*INS*) promoter or upstream or downstream of the super core promoter (SCP1). We simultaneously

delivered all four libraries to 832/13 rat insulinoma cells (n = 3) and collected DNA and RNA for

sequencing.

Figure 2. Promoter and fragment position relative to the promoter and reporter gene influence

MPRA activity. (A) After fragment activity quantification in R/MPRAnalyze, we performed principal

components analysis to estimate broad influence of promoter (captured primarily by PC1) and position

(captured by PC2) on MPRA activity. Each point represents an individual replicate, points are color-

coded based on plasmid configuration, and point shape indicates position of the cloned fragment. (B)

To jointly estimate effects of promoter and position across fragments in all four configurations, we

included promoter and position as covariates in the MPRAnalyze model. This scatterplot displays the

coefficients from Wald testing for the promoter (pink points along x-axis, *INS* promoter vs. SCP1

promoter with local FDR < 0.05) and position (orange points along y-axis, upstream versus downstream

with local FDR < 0.0). PC: principal component, FC: fold-change

Figure 3. Position and promoter bias in enhancer activity is partially attributed to presence of

relevant annotations. (A) To select sequence features associated with position or promoter bias, we

performed LASSO regression with 10-fold cross validation on position or promoter bias scores (signed

Wald statistics) as a function of several thousand transcription factor motifs and genomic annotations.

(B) Significant predictors of promoter bias are displayed. With the clustered set of annotations, we

identified only features enriched in fragments that are more active with the *INS* promoter. **(C)** Top 25

predictors (i.e., nonzero coefficients) associated with position bias effects are shown. Positive

coefficients indicate annotations that are enriched in fragments that are more active in the upstream

position while negative coefficients are enriched in fragments that are more active in the downstream

position.

Figure 4. HNF1 transcription factor motifs contribute to enhancer activity near selected T2D-

associated variants. (A) For 47 HNF1 motif-overlapping fragments with significant *INS* promoter bias

effects on activity, we designed three versions: original (motif intact), deleted (motif removed and

sequence adjusted), and shuffled (dinucleotide-shuffled motif). When the tested variant was adjacent to

the motif, we synthesized both reference and alternate alleles for each version. For variants directly

overlapping the motif, we generated only one deletion and one shuffled fragment. **(B)** We synthesized

four fragments corresponding to the variant rs1635852, which overlaps an HNF1 motif at a high

information content position. The T2D risk allele (T) disrupts this motif, while the non-risk allele (C)

matches the consensus. **(C)** Shuffling the motif significantly decreased enhancer activity compared to

intact fragments with either the risk T (Wilcoxon rank-sum test $p = 0.016$) or non-risk C allele ($p =$

0.008). Motif deletion also significantly decreased enhancer activity compared to the non-risk C allele (p

$= 0.016$). **(D)** For the variant rs11819995, located 11 bp upstream of an HNF1 motif, we synthesized six

fragments. **(E)** Deletion of the motif significantly decreased enhancer activity for both the reference (C,

non-risk) and alternate (T, risk) alleles (p for reference allele $= 0.008$ and for alternate allele $= 0.03$).

Shuffling the motif likewise reduced activity for both alleles ($p = 0.008$). **(F)** To assess context-specific

effects, we cloned these fragments into MPRA vectors with the SCP1 or skeletal muscle-specific

MYBPC2 promoter and delivered all three to LHCN-M2 human skeletal muscle myotubes (n = 6). **(G)**

When paired with the *INS* promoter, the shuffled rs11819995-containing fragment showed increased

activity relative to the original fragment ($p = 0.015$); however, none of the fragments containing

rs11819995 functioned as enhancers in LHCN-M2 myotubes, regardless of promoter context. Overall,

their activity is highest when paired with the skeletal muscle-specific promoter.

design oligos in regions of interest
(T2D/related trait GWAS signals)

clone into one
of four MPRA vectors

deliver to
832/13 rat insulinoma cells
(pancreatic beta cells)

collect & sequence

SCP1 +
Upstream

SCP1 +
Downstream

INS +
Upstream

INS +
Downstream

input DNA BCs

GFP mRNA BCs

A

O1
O2
O3
O4

Promoter or position
bias score

TF motif/annotation
overlap scores

TF1 TF2 TF3 TF4 ... AnnotN

B Predictors of Promoter Bias

POU1F1_2

HNF1_2

FOXJ3_3

HNF1_3

Coefficient

−0.12   −0.09   −0.06   −0.03   0.0

TF Motif or Genomic Annotation

INS

C Predictors of Position Bias

Upstream

X1_Active_TSS
ETS_known7
BATF_known1
CUX1_5
NFY_known4
ISX_2
atac_seq
NFY_known2
MYC_known20
ZSCAN26_1
HOMEZ_2
ATF2_1
GATA_known5
E2F_known18
ZNF423_1
PAX6_1
SOX17_2
PAX4_5
POU3F3_1
FOXL1_2
PAX2_3
RARA_3
PPARA_2
typicalEnhancer
ZBTB7A_known2

Coefficient

−0.05   0.0   0.10   0.20   0.30

TF Motif or Genomic Annotation

Downstream