Complete suspension culture of human induced pluripotent stem cells supplemented with
suppressors of spontaneous differentiation

Mami Matsuo-Takasaki[1], Sho Kambayashi[2], Yasuko Hemmi[1], Tamami Wakabayashi[1], Tomoya
Shimizu[1], Yuri An[1], Kazuhiro Takeuchi[2], Masato Ibuki[2], Yoshikazu Kawai[2], Masafumi
Umekage[3], Tomoaki M Kato[3], Michiya Noguchi[4], Koji Nakade[5], Yukio Nakamura[4], Tomoyuki
Nakaishi[2], Masayoshi Tsukahara[3], Yohei Hayashi[1,6]*.

[1]iPS Cell Advanced Characterization and Development Team, RIKEN BioResource Research
Center, Ibaraki, 305-0074, Japan; [2]Regenerative Medicine and Cell Therapy Laboratories,
KANEKA CORPORATION, Kobe, 650-0047, Japan; [3]Research and Development Center, CiRA
Foundation, Kyoto, 606-8507, Japan; [4]Cell Engineering Division, RIKEN BioResource Research
Center, Ibaraki, 305-0074, Japan; [5]Gene Engineering Division, RIKEN BioResource Research
Center, Ibaraki, 305-0074, Japan; [6]Faculty of Medicine, University of Tsukuba, Ibaraki, 305-
8575, Japan.
14

*For corresponding:

yohei.hayashi@riken.jp (YHa)

Abstract
Human induced pluripotent stem cells (hiPSCs) are promising resources for producing various
type of tissues in regenerative medicine; however, a scalable culture system that can precisely
control the cell status for hiPSCs is not developed yet. Utilizing suspension culture without
microcarriers or special materials allows for massive production, automation, cost effectiveness,
and safety assurance in industrialized regenerative medicine. Here, we found that hiPSCs cultured
in suspension conditions with continuous agitation without any microcarriers or extracellular
matrix components were more prone to spontaneous differentiation than those cultured in
conventional adherent conditions. Adding PKCβ and Wnt signaling pathway inhibitors in the
suspension conditions suppressed the spontaneous differentiation of hiPSCs into ectoderm and
mesendoderm, respectively. In these conditions, we successfully completed culture processes of
hiPSCs including the generation of hiPSCs from peripheral blood mononuclear cells with the
expansion of bulk population and single-cell sorted clones, long-term culture with robust self-
renewal characteristics, single cell cloning, direct cryopreservation from suspension culture and
their successful recovery, and efficient mass production of a clinical-grade hiPSC line. Our
results demonstrate that precise control of hiPSC status in suspension culture conditions paves the
way for their stable and automated clinical application.
34

Introduction

Human induced pluripotent stem cells (hiPSCs) are promising resources for various types of tissues in regenerative medicine (Takahashi *et al*, 2007; Yu *et al*, 2007). To enable cell therapy from hiPSCs, the development of a large-scale manufacturing system is essential because massive cell numbers are required to compose transplantable cells which are enough to rescue the desired physiological function (Chen *et al*, 2014; Kim & Kino-Oka, 2020; Tannenbaum & Reubinoff, 2022). In general, hiPSCs are believed to possess their scaffold dependency, thus are cultured under adhesion and monolayer culture conditions (Hayashi & Furue, 2016; Xu *et al*, 2001). However, utilizing suspension culture without microcarriers or special materials allows for massive production, automation, cost-effectiveness, and safety assurance in industrialized regenerative medicine.

Several attempts have been made to develop suspension culture technologies enabling rapid and large-scale preparation of hiPSCs (Amit *et al*, 2010; Amit *et al*, 2011; Dang *et al*, 2004; Elanzew *et al*, 2015; Horiguchi & Sakai, 2016; Hunt *et al*, 2014; Ibuki *et al*, 2019; Kehoe *et al*, 2010; Kim *et al*, 2019; Krawetz *et al*, 2010; Kwok *et al*, 2018; Lam *et al*, 2016; Nath *et al*, 2017; Oh *et al*, 2009; Olmer *et al*, 2012; Rohani *et al*, 2020; Shafa *et al*, 2012; Singh *et al*, 2010; Steiner *et al*, 2010; Wang *et al*, 2013; Zweigerdt *et al*, 2011) (Summarized in Table 1). These studies have achieved long-term culture and/or mass expansion of hiPSCs and/or human embryonic stem cells (hESCs) in suspension conditions. However, completed processes from clonal hiPSC generation to mass production of hiPSCs based on the precise control of cell status have not yet been achieved.

In this study, we have investigated what hampers the stable maintenance of undifferentiated cell states in suspension conditions. HiPSCs cultured in suspension conditions with continuous agitation without any microcarriers or extracellular matrix (ECM) components were more prone to spontaneous differentiation than those cultured in conventional adherent conditions. From screening of candidate molecules to suppress the spontaneous differentiation of hiPSCs, we have identified that inhibitors of PKCβ and Wnt signaling pathways suppress their differentiation into ectoderm and mesendoderm, respectively. In these conditions, we aimed to complete the processes of handling hiPSCs including the generation of hiPSCs with the expansion of bulk population and single-cell sorted clones, long-term culture with robust self-renewal characteristics, single-cell cloning, direct cryopreservation from suspension culture and their successful recovery, and efficient mass production of a clinical-grade hiPSC line.

Results

Suspension cultured hiPSCs are prone to spontaneous differentiation.

First, we investigated whether the quality of hiPSCs in suspension and adherent conditions are equivalent or not. HiPSCs cultured in a conventional medium with continuous agitation (90 rpm) in non-adhesive cell culture plates for 2 passages (5 days during passages) were examined (Figure 1A). In suspension culture on day 5 and 10, hiPSCs formed round cell assemblies with slightly uneven surfaces (Figure 1B). Gene expression analysis with RT-qPCR revealed that the expression of differentiation markers, such as PAX6 (ectoderm), SOX17 (endoderm), and T (mesoderm) increased in suspension-cultured hiPSCs for 10 days (Figure 1C). To monitor the spontaneous differentiation at single cell resolution, we established knock-in reporter hiPSC lines of PAX6-tdTomato and SOX17-tdTomato to visualize and quantify the expression of PAX6 and SOX17 at protein level, respectively (Figure 1—figure supplement 1A–H). tdTomato-positive cells were clearly observed in day 10 samples in suspension-cultured hiPSCs, whereas no positive cells were observed in adherent culture conditions (Figure. 1D). Flow cytometric analysis revealed that hiPSCs in suspension-culture conditions contained non-negligible percentages of PAX6-tdTomato-positive and SOX17-tdTomato-positive cells (Figure. 1E and 1F). Western blot analysis revealed that protein expression of PAX6 and SOX17 in suspension-culture was significantly increased (Figure 2—figure supplement 1A–D). The ratio of positive cells for a cell surface marker for undifferentiated hiPSCs, TRA1-60, was significantly lower in suspension-culture conditions (Figure 2—figure supplement 1E and F). These results suggest that a portion of hiPSCs was spontaneously differentiated in suspension conditions when cultured in conventional medium. To examine global changes in gene expression patterns between suspension and adherent conditions, whole-transcriptomic RNA-seq experiments with statistical tests were performed. Gene Set Enrichment Analysis (GSEA) of all the genes and Gene Ontology Enrichment Analysis (GOEA) on differentially-regulated genes revealed that, in suspension conditions, number of genes involved in differentiation toward various tissues and cell-cell adhesions were significantly up-regulated, whereas genes involved in nucleotide metabolism, hypoxic responses, and extracellular matrix organization were significantly-down-regulated (Figure 1G–J). These results suggest that hiPSCs in suspension conditions are in the process of spontaneous differentiation into various cell lineages and are characterized with specific signatures of gene expression patterns.

Wnt signaling inhibitors suppress spontaneous mesendodermal differentiation in suspension culture condition.

We next investigated the factors having an inhibitory effect on hiPSCs spontaneous differentiation in suspension conditions (Figure 2A). Since Wnt signaling induces the differentiation of human pluripotent stem cells into mesendoderm lineages (Nakanishi *et al*, 2009;

Sumi *et al*, 2008; Tran *et al*, 2009; Vijayaragavan *et al*, 2009; Woll *et al*, 2008), and is

endogenously involved in the regulation of mesendoderm differentiation of pluripotent stem cells

(Dziedzicka *et al*, 2021), we hypothesized that adding Wnt signaling inhibitors/activator may

alter spontaneous differentiation of hiPSCs into mesendoderm. Therefore, Wnt signaling

inhibitors, IWP2 or IWR-1-endo (Chen *et al*, 2009), or an activator, CHIR99021 (Ring *et al*,

2003), were added to the culture medium under suspension conditions. HiPSC aggregates treated

with and without inhibitors showed similar shapes (Figure 2B). In contrast, hiPSC aggregates

treated with CHIR99021 formed heterogeneously shaped cyst-like structures, suggesting that they

were largely differentiated. In samples treated with inhibitors, both T and SOX17 expression

levels were significantly reduced to the level of adherent-cultured hiPSCs; however, there was no

reduction in PAX6 expression (Figure 2C). In contrast, CHIR99021-treated cell aggregates

showed markedly decreased OCT4 and increased T and SOX17 expression. Additionally, SOX17

protein expression was suppressed in hiPSCs treated with IWR-1-endo in suspension-culture

conditions, although its expression increased in hiPSCs in suspension conditions with

conventional culture medium compared to adherent conditions (Figure 2—figure supplement 1A

and B). These results indicate that Wnt signaling inhibitors effectively suppressed

mesendodermal differentiation in suspension-culture, but was insufficient to suppress ectodermal

differentiation.

PKC signal inhibitors suppress spontaneous neuroectododermal differentiation in

suspension conditions.

To identify molecules with inhibitory activity on neuroectodermal differentiation, hiPSCs were

treated with candidate molecules in suspension conditions. Out of the candidate molecules tested

(listed in Supplementary Table 1), Go6983, a pan-PKC inhibitor (Gschwendt *et al*, 1996), showed

the strongest inhibition against PAX6 expression (Figure 2D). Further, simultaneous treatment

with Go6983 and IWR-1-endo decreased PAX6, T, and SOX17 expression, while maintaining

OCT4 expression. To confirm these screening test results, the dose dependent effect of Go6983 or

another PKCα, β, γ inhibitor GF109203X (GFX) (Toullec *et al*, 1991) on the inhibition of

PAX6 expression was observed at constant concentration of IWR-1-endo (Figure 2E and F).

These results demonstrated that the inhibition of PKC signaling pathway effectively suppress

neuroectoderm differentiation in hiPSCs cultured in suspension conditions. Furthermore, we

tested various PKC inhibitors to suppress neuroectodermal differentiation from hiPSCs in

suspension culture using the 201B7 hiPSC line (Figure 2—figure supplement 2A). The

suppression of PAX6 expression in hiPSCs cultured in suspension conditions was observed with

PKC inhibitors, enzastaurin (Ly317615), sotrastaurin (AEB071), Ro-32-0432, Go6983,

GF109203X, and LY333531, which possessed PKCβ inhibition activity (Figure 2—figure

supplement 2B-D); however, sotrastaurin and Ro-32-0432 also showed growth inhibition. Thus,

for the rest of the study, a PKCβ specific inhibitor, LY333531 (Jirousek *et al*, 1996) was used. As

compared to controls, another hiPSC line, WTc11, cultured in suspension conditions treated with

IWR-1-endo and LY333531 formed homogenous, round, smooth-surfaced aggregates (Figure.
3A). PAX6 expression was strongly suppressed by the addition of LY333531 (Figure. 3B).
Furthermore, after adding IWR-1-endo, the inhibitory effect of PAX6 expression was further
enhanced, and simultaneously, OCT4 expression was restored to the same level as in the
adherent-culture. PAX6 protein expression was also suppressed in hiPSCs treated with IWR-1-
endo and LY333531 in suspension-culture conditions while its expression increased in
suspension conditions with conventional culture medium compared to adherent-conditions
(Figure 2—figure supplement 1C and D). The ratio of TRA-1-60-positive cells was higher in
suspension-culture conditions supplemented with IWR-1-endo and LY333531 than in control
conditions without these inhibitors (Figure 2—figure supplement 1E and F). These results
indicate that the maintenance of suspension-cultured hiPSCs is specifically facilitated by the
combination of PKCβ and Wnt signaling inhibition.

We examined the effect of PKCβ and Wnt inhibitors on global gene expression of hiPSCs under
suspension conditions. RNA-seq data were obtained from suspension conditions in absence of
inhibitors (Sus), supplemented with IWR-1-endo (IWR), LY333531 (LY), IWR-1-endo and
LY333531 (IWRLY), and adherent conditions (Ad). Hierarchical clustering obtained from these
data showed that LY and IWRLY were grouped closely with Ad (Figure 3C). In contrast, Sus and
IWR were both grouped as discrete populations from Ad. Additionally, hierarchical clustering in
gene expression among these conditions was supported by principal component 1 (PC1) in the
principal component analysis (PCA) (Figure 3D). In contrast, PC2 represented genes related to
the effects of specific inhibitors under these conditions. Next, we investigated the effect of
LY333531 and IWR-1-endo in suspension culture. Many genes involved in pluripotency: KLF4
and ID1, and epithelial cell-cell interactions: CDH1 (E-cadherin), were significantly up-regulated
in IWRLY, while many transcription factors involved in differentiation: PAX2, 3, 5, and 8, SP5,
DBX1, and TFAP2B, were down-regulated in IWRLY (Figure 3E). GSEA and GOEA on down-
regulated genes in IWRLY showed that the expression of developmentally associated genes,
whose expression was elevated in Sus, was generally reduced in IWRLY (Figure 3F and G).
GOEA on up-regulated genes revealed gene sets involved in epithelial cell types (Figure 3H). As
compared to Ad, genes involved in sensory system development, cell-cell adhesion, and Wnt and
PI3K signaling pathways were up-regulated in IWRLY, and genes involved in nucleotide
metabolism and hypoxic responses were down-regulated under IWRLY conditions (Figure 3—
figure supplement 1A-D). These results suggest that PKCβ and Wnt signaling inhibitors in
suspension conditions regulate global gene expression patterns to suppress spontaneous
differentiation, albeit remaining expression signatures of suspension culture, possibly due to the
microenvironment within the formed aggregates and physiological differences.

Further, we examined the expression pattern changes in specific isoforms of PKCs in hiPSCs
cultured in adherent/suspension conditions. RNA expression of PKCα (PRKCA) and PKCβ
(PRKCB) was significantly up-regulated under suspension-culture conditions compared to

adhesion conditions (Figure 3—figure supplement 2A). Moreover, phospho-PKCβ protein

expression was significantly elevated (Figure 3—figure supplement 2B and C). These results

suggest that elevated expression of PKCβ in suspension-cultured hiPSCs could affect the

spontaneous differentiation.

Simultaneous inhibition of PKCβ and Wnt signals maintains undifferentiated hiPSCs in

suspension for long-term culture.

We next performed long-term culture for 10 passages in suspension conditions and compared

hiPSC growth in presence of LY333531 or Go6983. When hiPSCs were seeded at 4×10^5

cells/well, the average cell number reached approximately 12-fold after 5 days under both

conditions (Figure 4A and B). After 10 passages, aggregates of hiPSCs cultured in the presence

of LY333531 showed a uniform spherical shape, whereas aggregates cultured in the presence of

Go6983 were heterogeneously spherical (Figure 4C). Notably, in LY333531-treated cells, OCT4-

positive cell numbers were significantly higher than in Go6983-treated samples, as determined by

immunostaining (Figure 4D and E). To evaluate whether hiPSCs cultured in suspension

conditions with PKCβ and Wnt signaling inhibitors for 10 passages maintain pluripotency, we

performed embryoid body (EB) formation assay. These EBs contained positive cells for TUJ1,

SMA, and AFP as ectodermal, mesodermal, and endodermal marker, respectively (Figure 4F).

Comparative genomic hybridization (CGH) array analysis showed that hiPSCs cultured long-term

in the presence of PKCβ and Wnt inhibitors retained their normal human karyotype (Figure 4G).

These results indicate that, for long-term culture, inhibition of Wnt signaling and PKCβ in

suspension conditions was sufficient to maintain the self-renewal, pluripotency, and genomic

integrity of hiPSCs.

We further investigated whether the effects of PKCβ and Wnt inhibitors on suppressing hiPSCs

spontaneous differentiation in suspension-culture conditions are applicable to other culture

medium. Morphologies and gene expression profile of hiPSCs, cultured in suspension conditions

with another commercially available maintenance medium, StemScale (ThermoFisher Scientific,

Inc., MA, USA), were examined (Figure 4—figure supplement 1A). HiPSCs cultured in

suspension treated with IWR-1-endo and LY333531 formed homogenously round and smooth-

surfaced aggregates compared to controls (Figure 4—figure supplement 1B). The differentiation

markers expression, which was elevated under suspension conditions, was suppressed by the

simultaneous addition of PKCβ and Wnt signal inhibitors, as observed with StemFit medium

(Figure 4—figure supplement 1C and D). These results suggest that conventional suspension

culture methods might contain spontaneous differentiating cells and that the addition of inhibitors

of PKCβ and Wnt signaling pathways to culture medium generally suppresses spontaneous

differentiation.

Mass and clonal expansion of hiPSCs in suspension conditions supplemented with inhibitors

of PKCβ and Wnt signaling pathways.

For the clinical applications of hiPSCs, its homogenous mass production is required to obtain
sufficient quantities. To test the feasibility of mass production under suspension-culture
conditions supplemented with PKCβ and Wnt signal inhibitors, we first performed suspension-
culture using healthy-donor derived hiPSC lines, 1383D6 and 1231A3, in a 30 mL bioreactor
with stirring blade and demonstrated that undifferentiated hiPSCs were efficiently maintained in
this condition (Figure 5—figure supplement 1A). These cell lines showed higher cell density in
the bioreactor at each passage after 3 days (Figure 5—figure supplement 1B). Under these
conditions, the concentration of glucose decreased daily (Figure 5—figure supplement 1C),
whereas that of L-lactic acid increased (Figure 5—figure supplement 1D). These results indicate
that the cell lines proliferated with active energy consumption. Further, RT-qPCR analysis
showed no marked differences in the undifferentiated and differentiation markers expression
between hiPSCs cultured in adhesion vs. suspension with PKCβ and Wnt inhibitors (Figure 5—
figure supplement 1E and F). Flow cytometric analysis showed that more than 90% of these cells
were positive for pluripotency marker proteins: NANOG, OCT4, and SOX2 (Figure 5—figure
supplement 1G). G-band analysis of suspension-cultured hiPSCs after five passages revealed
normal karyotype (Figure 5—figure supplement 1H). These results demonstrate that
undifferentiated hiPSCs were efficiently maintained in the bioreactor using the culture medium
supplemented with inhibitors of PKCβ and Wnt signaling pathways.

Next, we expanded hiPSCs in a large-scale culture system under perfusion conditions in the
presence of IWR-1-endo and LY333531. In this experiment, clinical-grade hiPSC line, Ff-l14s04,
which is derived from PBMCs of a donor carrying homozygous alleles for major HLA loci
(HLA-A, HLA-B, and HLA-DR) was used (Kitano *et al*, 2022). Large-scale hiPSCs preparation
using a perfusion-culture system with 320 mL bioreactor having stirred-wing was performed in
GMP-compliant, clinical grade StemFit AK03N medium containing IWR-1-endo and LY333531
(Figure 5A). When the culture was started at 1×10^5 cells/mL in 320 mL medium, hiPSCs
proliferated nearly 10-fold after 3 to 4 days and reached approximately 1×10^9 cells in total to
produce ~300 stock vials (1×10^6 cells/vial). This large-scale culture was repeated three times
(Passages 1, 2, and 3), and hiPSCs obtained at each passage were characterized. These samples
showed similar or higher viability (>90%) to that of adherent-culture derived cells (Figure 5B).
When compared to adherent conditions, these samples in suspension-culture showed a similar or
higher proliferation rate (Figure 5C). Cells obtained from large-suspension cultures formed
typical hiPSC-like colonies, after vials were thawed and seeded in adherent-culture conditions
(Figure 5D). Flow cytometry detected over 90% of cells positive for pluripotent cell markers
expression: TRA-1-60, SSEA4, and OCT4 (Figure 5E and F). Further, G-band analysis revealed
that hiPSCs retained their normal karyotype even after three passages under large-scale
suspension conditions (Figure 5G). When large-scale suspension-culture derived hiPSCs were
incubated in each germ layer-specific differentiation medium for 4 - 7 days, the expression of
early differentiation markers for ectoderm (PAX6 and SOX1), mesoderm (T and PDGFRA) and

endoderm (SOX17 and CXCR4) was significantly induced (Figure 5H). These cells were then
directly differentiated into dopaminergic neural progenitors, cardiomyocytes, and hepatocytes to
evaluate their differentiation capacity and propensity. There were no differences in the
differentiation efficiency toward these lineages (Figure 5I-M). Hence, these results indicate that
the characteristics and quality of clinical-grade hiPSCs cultured in large-scale suspension
conditions in the presence of PKCβ and Wnt inhibitors were equivalent to those of hiPSCs
maintained under adherent conditions. Taken together, we were successful in mass suspension-
culture conditions supplemented with Wnt and PKCβ inhibitors.

Single cell sorting and expansion of hiPSC subclones cultured in suspension conditions.
We sorted a hiPSC line, 201B7, with TRA-1-60 antibody into individual wells of a 96-well plate
and attempted to establish subclones by expanding them with serial passages (Figure 6A). By day
28 after this single-cell sorting, we expanded single-cell-sorted hiPSC subclones in suspension
culture supplemented with IWR1-endo and LY3333531 (Figure 6B). On day 28, we examined
seven subclones for their cell growth and expression of OCT4 and TRA-1-60. The subclones
showed more than three million cells and high ratios of OCT4- and TRA1-60-positive cells
(Figure 6C-E). These results indicate that we have successfully derived single-cell-cloned
sublines in suspension culture.

Direct freeze and thaw of hiPSCs cultured in suspension conditions.
We froze a hiPSC line, 201B7, in suspension culture conditions supplemented with IWR1-endo
and LY3333531. Then, we thawed these frozen vials and directly reseeded the cells in suspension
conditions (Figure 6—figure supplement 1A). By day 10 after reseeding, we expanded the
hiPSCs in suspension culture supplemented with IWR1-endo and LY3333531 (Figure 6—figure
supplement 1B). On day 10, we examined three vials for their cell growth and expression of
OCT4 and TRA-1-60. The subclones showed more than three million cells and high ratios of
OCT4- and TRA1-60-positive cells (Figure 6—figure supplement 1C-E). These results indicate
that we have successfully frozen and thawed hiPSCs in suspension culture directly.

Together, these results suggest that the suspension culture conditions supplemented with Wnt and
PKCβ inhibitors are robust in harsh conditions such as single-cell expansion and freezing-and-
thaw processes.

Establishment of hiPSC lines in complete suspension-culture conditions supplemented with
inhibitors of PKCβ and Wnt signaling pathways.
For the clinical use of hiPSCs, more secure and efficient establishment of hiPSCs in closed
systems is strongly desired. To achieve this, we aimed to establish hiPSCs in suspension-culture
conditions. Using PBMCs as a starting material, we generated hiPSCs using a novel replication-
defective and persistent Sendai virus vector (SeVdp) infection or episomal vector electroporation,
as these methods are well-known for producing transgene-free, clinical-grade hiPSCs (Fusaki *et*

al, 2009; Nishimura *et al*, 2011) (Figure 7A). PMBCs were infected with SeVdp carrying OCT4,
SOX2, KLF4, and L-MYC. Infected cells were kept cultured by repeating passages every 5 - 6
days. Cell aggregates cultured with IWR1-endo and LY33351 showed uniform spherical structure
(Figure 7B), and most of the cells were positive for OCT4 (Figure 7C) and TRA-1-60 (Figure
7D) on day 56. These bulk cells were able to differentiate into three germ layers in an in vitro EB
formation assay (Figure 7E) and in teratomas that were transplanted into immunodeficient
NOD.Cg-Prkdcscid Il2rgtm1Wjl/SzJ (NSG) mice (Figure 7F). These bulk-hiPSCs maintained
normal karyotype (Figure 7G). By single cell sorting with fluorescent labeled TRA-1-60
antibody, we expanded single-cell-derived clones in suspension-culture conditions. A clone (F-
10) showed OCT4 and TRA1-60 expressions (Figure 7H-J). In addition, the established clone
showed potency to differentiate into derivatives of three germ layers *in vitro* as EBs (Figure 7K)
and *in vivo* as teratoma transplanted into NSG mice (Figure. 7L). A normal karyotype was
observed in this clone (Figure 7M). SeVdp was nearly extinct in these bulk populations and F-10
clone (Figure 7N). These results demonstrate that, we were successful in establishing transgene-
free hiPSC lines using SeVdp infection in suspension-culture conditions in the presence of PKCβ
and Wnt inhibitors.

Furthermore, we were successful in establishing single-cell-derived hiPSC lines using the
transfection of episomal plasmid vectors in suspension-culture conditions in the presence of
PKCβ and Wnt signaling inhibitors. PMBCs were transfected with episomal plasmid vector mix
carrying OCT4, SOX2, KLF4, L-MYC, LIN28, and mp53DD and cultured in suspension
conditions in the presence or absence of IWR-1-endo and LY33351 with repeated passages to
reach enough cell numbers to make cell stocks and characterization (Figure 7—figure supplement
1A). After 36 days of culture, 59% double positive cells for TRA-1-60 and SSEA4 were observed
in control, whereas 78% were seen in the presence of IWR-1-endo and LY333531 (Figure 7—
figure supplement 1B and C). We performed cell sorting of TRA-1-60 and SSEA4 double
positive cells and further repeated 5 more passages in the absence or presence of IWR-1-endo and
LY333531. Cell aggregates cultured in culture medium supplemented with IWR-1-endo and
LY333531 showed uniform spherical structures (Figure 7—figure supplement 1D), and the most
of these cells were TRA-1-60 and SSEA4 double positive (Figure 7—figure supplement 1E). In
contrast, cell aggregates cultured without these inhibitors showed lumpy, heterogeneous shapes,
and decreased undifferentiated population. Immunostaining of OCT4 on dissociated cell
aggregates resulted in higher OCT4 positive cells in IWR-1-endo and LY333531 treated cells
(Figure 7—figure supplement 1F and G). We further performed single cell sorting of TRA-1-60-
positive cells from bulk-cultured populations at 10 passages (Figure 7—figure supplement 1H
and I). We selected 3 clones to be characterized further. These clones showed round spherical
shapes in their aggregates and expression of OCT4, NANOG, and TRA-1-60 proteins (Figure 7—
figure supplement 1J-L). Pluripotency into three germ layers was confirmed in vitro with EB
formation assay (Figure 7—figure supplement 1M) and teratoma formation analysis (Figure.
Figure 7—figure supplement 1N). These clones retained a normal karyotype even after a long-

culture (Figure 7—figure supplement 1O). These results demonstrate that single-cell-derived

hiPSC lines were established using the transfection of episomal plasmid vectors in suspension-

culture conditions in the presence of PKCβ and Wnt signaling inhibitors.

Discussion

In this study, we have developed a series of methods to generate and maintain hiPSCs in suspension conditions. First, we have identified compounds that suppress spontaneous differentiation of hiPSCs in suspension cultures. Based on these findings, we have newly achieved a complete series of culture processes including hiPSC establishment, long-term culture, mass culture, single-cell cloning, and direct freeze and thaw (summarized in Table 1). These methods require no special materials including microcarriers or dialysis bags, and are effective with various existing culture media and many hiPSC lines. Therefore, this suspension-culture of hiPSCs is anticipated to have novel industrial applications, which have been never achieved with conventional methods because suspension-culture conditions, which precisely control cellular states, have advantages in terms of mass culture, automation, and safety assurance.

We have identified two compounds as factors that ameliorate the disadvantages of the suspension culture of hiPSCs. Wnt signaling inhibitors specifically inhibited differentiation toward mesendoderm, whereas, the PKCβ inhibitor, LY333531, inhibited ectodermal differentiation and, when combined with IWR-1-endo, maintained the undifferentiated nature of hiPSCs with high efficiency. Go6983, a pan-PKC inhibitor, has been used to promote the self-renewal of mammalian PSCs (Dutta *et al*, 2011; Kinehara *et al*, 2013; Rajendran *et al*, 2013; Takashima *et al*, 2014); however, in these cases, the role of specific isoforms in PSC self-renewal and differentiation are not fully elucidated yet. In this study, we demonstrated that PKCβ-specific inhibitor LY333531 as well as Go6983 showed the suppression of neuroectodermal differentiation. Interestingly, these combinations of chemical inhibitors against WNT and PKC signaling pathways are also being used in the induction and maintenance of naive human pluripotent stem cells (Bayerl *et al*, 2021; Bredenkamp *et al*, 2019; Guo *et al*, 2017; Khan *et al*, 2021). While, in suspension-culture, we did not aim to drive hiPSCs to a naïve state by the use of these combinations of chemical inhibitors conditions, it would be worthwhile to investigate the effect of adaptation of suspension-culture supplemented with PKCβ and Wnt signaling inhibitors on the transition of the naïve PSCs types.

We successfully generated hiPSC lines from PBMCs in suspension conditions for the first time by adding the abovementioned compounds to the culture medium. Both transduction methods with SeVdp and episomal plasmid vectors enabled the establishment of hiPSCs under suspension conditions in the presence of IWR-1-endo and LY333531. Clonal expansion of hiPSCs was also performed by single-cell sorting with flow cytometry and were successfully characterized for their self-renewal, pluripotency, and genomic integrity. These results indicate that hiPSC lines generated in suspension conditions have the same quality in terms of their pluripotency, self-renewal, and genomic integrity as those generated with conventional adherent conditions. Precise control of hiPSC status in suspension culture could pave the way for their stable and automated clinical application toward autologous cell therapy of hiPSCs.

Methods

Human subjects

The generation and use of hiPSCs were approved by the Ethics Committee of RIKEN

BioResource Research Center, Graduate School of Medicine of Kyoto University, and KANEKA

CORPORATION.

Animal experiments

All animal experiments were approved by the Animal Experimentation Committee of the RIKEN

Tsukuba branch and were performed according to the committee's guiding principles and the

"Guide for the Care and Use of Laboratory Animals" published by the National Institute of

Health.

Data availability

All data are available in the main text or the supplementary materials. RNA-seq data are available

in the Gene Expression Omnibus as GSE222833. All the data supporting the findings of this

study are available from the lead contact upon reasonable request. Source data is provided with

this paper. This paper does not report original code.

HiPSCs cultured in adherent conditions

WTC11 (GM25256 from Coriell Institute) (Hayashi *et al*, 2016), 201B7 (HPS0063 from RIKEN

Cell Bank) (Takahashi *et al*., 2007), 454E2 (HPS0077 from RIKEN Cell Bank) (Okita *et al*,

2011), 1383D6 (HPS1006 from RIKEN Cell Bank) (Okita *et al*., 2011), 1231A3 (HPS0381 from

RIKEN Cell Bank) (Okita *et al*., 2011), and Ff-I14s04 (CiRA foundation, Kyoto University)

(Kitano *et al*., 2022) were used as healthy-donor hiPSC lines. With tdTomato-reporter hiPSC

lines, we generated knock-in hiPSC lines for PAX6-TEZ (constructed from 454E2 line) and

SOX17-TEZ (constructed from 1383D6 line) using CRISPR-Cas9 genome editing (Tsukamoto *et*

al, 2021) (Figure. S1A). These hiPSCs were cultured in StemFit AK02N medium (Cat#AK02N,

Ajinomoto, Co., Ltd., Tokyo, Japan). Medium change was performed every day and passaged at

80%–90% confluency after 6-7 days of culture. At passage, PBS/EDTA solution (0.5 mM,

Cat#06894-14, Nacalai Tesque) was used to dissociate hiPSC colonies, and these cells were

seeded at a density of 2,500 cells/cm^2. 10 μM Y-27632 (Cat#HY-10071, FUJIFILM Wako Pure

Chemical Corporation, Osaka, Japan) and 0.25 μg/cm^2 iMatrix-511 silk (Cat#892021, Matrixome

Inc, Osaka, Japan) were added to the culture dish on seeding day.

Suspension-culture of hiPSCs

Suspension culture with rotation at 90 rpm was performed with a plate shaker (Cat#WB-101SRC,

WAKENBTECH, Kyoto, Japan or #0081704-000, TAITEC, Tokyo, Japan) installed in a CO_2

incubator (Cat#Steri-Cycle i160, Thermo Fisher Scientific, MA, USA) and operated under high

humidity continuously during the whole culture period. To start the culture hiPSCs in suspension

conditions, 4×10^5 cells were seeded in one well of a low-attachment 6-well plate (Cat#MS-

80060, Sumitomo Bakelite, Tokyo, Japan) with 4 mL of StemFit AK02N medium,

StemFitAK03N (Cat#AK03N, Ajinomoto, Co., Ltd., Tokyo, Japan), or StemScale PSC

suspension medium (A4965001, Thermo Fisher Scientific, MA, USA) supplemented with 10 μM

Y-27632. This plate was placed onto the plate shaker in the CO_2 incubator. The medium without

Y-27632 was changed every day unless otherwise specified. On days 3 - 5, the hiPSC aggregates

were dissociated with Accutase (Cat#12679-54, Nacalai Tesque) at 37°C for 10 min and used for

the next suspension-culture. This passage was performed every 5 days unless otherwise specified.

To screen for factors that inhibit spontaneous differentiation of hiPSCs, chemicals or recombinant

proteins were added to the culture medium (listed in Supplementary Table S1).

Bioreactor culture of hiPSCs

A frozen stock of hiPSCs was pre-cultured twice in adherent conditions to prepare enough cell

numbers. To prepare enough hiPSCs to start large-scale culture, hiPSCs were pre-cultured in

iMatrix-511MG (Cat#892 005, Matrixome, Inc, Osaka, Japan) or Vitronectin (VTN-N)

Recombinant Human Protein, Truncated (Cat#A14700, Thermo Fisher Scientific, MA, USA)-

coated cell culture flasks with StemFit AK03N medium including 20 μM IWR-1-endo and 1 μM

LY333531. Perfusion-culture was started with 3.2 - 4.8×10^7 cells in 320 mL of StemFit AK03N

medium (Cat#AK03N, Ajinomoto, Tokyo, Japan) with IWR-1-endo and LY333531. The reactor

system BioFlo320 (Eppendorf, Hamburg, Germany) was used for large-scale hiPSCs culture. To

maintain the lactate concentration below a certain level and to regulate the pH, the culture was

carried out by increasing the amount of medium perfusion per unit time in accordance with the

cell proliferation transition. To prevent pH decrease, CO_2 concentration was regulated by

feedback control in the reactor system. After 3-4 days of culture, the formed hiPSC aggregates

were dissociated with TrypLE SELECT (Cat#A12859-01, Thermo Fisher Scientific, MA, USA)

collected for making cell stocks (~300 tubes). This perfusion culture was repeated three times

(P1, P2, and P3) and the cells were prepared at each expansion step.

HiPSC generation in suspension culture conditions

HiPSCs were generated from healthy donor-derived PBMCs (Precision for Medicine). Thawed

PMBCs from a vial containing around 1×10^7 cells were pre-cultured in one well of low

attachment 6-well plate including 4 mL of StemSpan-AOF (Cat#ST100-0130, STEMCELL

Technologies, Vancouver, Canada) supplemented with recombinant human IL-6 (100 ng/mL),

IL-3 (10 ng/mL), SCF (300 ng/mL), TPO (300 ng/mL), and FLT3 ligand (300 ng/mL) (all from

FUJIFILM Wako Pure Chemical Corporation, Tokyo Japan). After 24 h of incubation with

continuous stirring at 37°C and 5% CO_2, PBMCs were spun down with centrifugation at 200 × g

for 10 min at low deceleration speed and resuspended in StemSpan ACF for cell counting. 2.5 ×

10^6 cells of PMBCs were centrifuged at $200 \times$ g for 10 min with low deceleration speed, and

electroporated using Nucleofector 2b device (Lonza, Basel, Switzerland) with iPS cell Generation

Episomal vector Mix (Takara Bio, Shiga, Japan) and Amaxa Human CD34$^+$ Cell Nucleofector kit

(Lonza, Basel, Switzerland) according to manufacturer's protocol. 5×10^5 of electroporated cells

were seeded in one well of a low attachment 6-well plate in 4 mL StemSpan ACF with the

cytokines mentioned above. Suspension-culture was performed with continuous agitation at 90

rpm. Stem span ACF medium were gradually replaced with StemFit AK02N medium. Formed

cell aggregates were passaged with Accutase on day 16 in the presence of 10 μM Y27632, and

suspension-culture was continued until the cell numbers reached a sufficient amount for

characterization. 10 μM IWR-1-endo and 1 μM LY333531 were added from day 3.

SeVdp infection was performed with Sendai Reprogramming Kit (CytoTune EX-iPS of

virus solution, ID Pharma, Tsukuba, Japan) according to the manufacturer's protocol with some

modifications. Briefly, pre-cultured 1×10^6 PMBCs were centrifuged at $200 \times$ g for 10 min, with

low deceleration speed, and resuspended in 2 mL of StemSpan ACF with cytokines. PMBCs were

gently mixed with 2 mL of virus solution prepared at MOI = 5 per 1×10^6 cells. 5×10^5 infected

cells were seeded in one well of low-attachment 6-well plate at a total volume of 4 mL with

StemSpan ACF plus cytokines. Suspension-culture was initiated with continuous agitation at 90

rpm. Stem span ACFs were gradually replaced with StemFit AK02N as mentioned above, and the

cell aggregates were passaged with Accutase on day 16 in the presence of 10 μM Y-27632.

Suspension-culture and passages continued until cell reached a sufficient number for

characterization. 10 μM IWR-1-endo and 1 μM LY333531 were added from day 23.

Three germ layer differentiation *in vitro*

For Embryoid body (EB) formation assay, suspension-cultured hiPSC lines were dissociated with

Accutase and 1.0×10^4 cells were seeded in each well of an EZ-BindShut 96-well-V plate

(Cat#4420-800SP AGC TECHNO GLASS CO., LTD, Shizuoka, Japan) with 100 μL Stem Fit

AK02N medium supplemented with 10 μM Y-27632. Before culture, the 96-well-V plate was

centrifuged at 200 x g for 3 min for efficient cell mass formation. The next day, the culture

medium was switched to DMEM high Glucose (Cat#08458-16, Nacalai tesque, Kyoto, Japan)

supplemented with 10% fetal bovine serum (here after referred as EB medium). On day 8, EBs

were transferred into 0.1% (w/v) Gelatin Solution (Cat#190–15805, FUJIFILM Wako Pure

Chemical Corporation, Tokyo Japan) -coated 12-well plate and further cultured in EB medium for

another 8 days. The medium was changed every day. EBs were fixed with 4% paraformaldehyde

in Dulbecco's Phosphate Buffered Saline (D-PBS) (Cat#14249-24, Nacalai tesque, Kyoto, Japan)

for 10 min at room temperature and used for immunostaining. Differentiation was validated by

immunostaining against each germ layer markers.

The differentiation potency of suspension-cultured hiPSCs toward three germ layers was

also evaluated by culturing in germ layer specific differentiation medium. For ectodermal

differentiation, suspension-cultured hiPSCs spheroids were cultured with Stem Fit AK02N

without C medium supplemented with 10 μM SB431542 (Cat#198-16543, FUJIFILM Wako Pure

Chemical Corporation, Tokyo, Japan) and 10 μM DMH1 (Cat#041-33881FUJIFILM Wako Pure

Chemical Corporation, Tokyo, Japan) for 7 days. For mesodermal differentiation, suspension-

cultured hiPSCs spheroids were cultured with RPMI1640 medium (FUJIFILM Wako Pure

Chemical Corporation, Tokyo, Japan) supplemented with B-27 supplement, minus insulin

(A1895601, Thermo Fisher Scientific, MO, USA) and 7.5 μM CHIR99021 (Cat#034-23103,

FUJIFILM Wako Pure Chemical Corporation, Tokyo, Japan) for 1 day, and continuously with

RPMI1640 medium supplemented with B-27 supplement for 2 days. For endodermal

differentiation, suspension-cultured hiPSCs spheroids were cultured with RPMI1640 medium

supplemented with 1 mM Sodium pyruvate (FUJIFILM Wako Pure Chemical Corporation,

Tokyo, Japan), 1 x NEAA (FUJIFILM Wako Pure Chemical Corporation, Tokyo, Japan), 80

ng/mL Activin A (R&D Systems, MN, USA), 55 μM 2-mercaptethanol (FUJIFILM Wako Pure

Chemical Corporation, Tokyo, Japan), 50 ng/mL FGF2 (R&D Systems, MN, USA), 20 ng/mL

BMP4, and 3 μM CHIR99021 for 2 days, and continuously with RPMI1640 medium

supplemented with 1 mM Sodium pyruvate, 1× NEAA, 80 ng/mL Activin A, 55 μM 2-

mercaptethanol, and 0.5% Knockout serum replacement (KSR; Thermo Fisher Scientific, MO ,

USA) for 2 days.

Neuroectoderm and endoderm differentiation of reporter hiPSC lines

Reporter hiPSC lines, PAX6-TEZ and SOX17-TEZ were used as positive controls for in vitro

neuroectoderm and endoderm differentiation, respectively. PAX6-TEZ was seeded in each well

of a 24 well-plate coated with 0.25 μg/cm^2 iMatrix-511 silk with 1 mL StemFit AK02N medium

supplemented with 10 μM Y-27632. Next day, the culture medium was switched to Stem Fit

AK02N medium without supplement C, instead of containing 10 μM SB431542 and 10 μM

DMH1. SOX17-TEZ was seeded in each well of a 24 well-plate (same as above), and on the next

day, the culture medium was switched to StemFit AK02N medium without supplement C instead

of containing 3 μM CHIR99021 and 10 ng/mL Activin A (Cat#014-23961, FUJIFILM Wako

Pure Chemical Corporation, Tokyo, Japan). The medium was changed every day. On day 7,

tdTomato expression was observed under all-in-one fluorescent microscope (BZ-X800;

KEYENCE).

Cardiomyocyte differentiation

Cardiomyocyte differentiation was performed according to a modified method described

previously (Funakoshi *et al*, 2016). Both large-scale suspension cultured hiPSCs and typical

adherent-cultured hiPSC were collected by centrifugation after dissociation into single cell with

TrypLE Select. The cells were suspended in 1.5 mL cardiomyocyte differentiation media (CDM),

consisting of StemPro34 medium (Thermo Fisher Scientific) supplemented with 2 mM Glutamax

(Thermo Fisher Scientific), 50 μg/mL ascorbic acid, 4×10^{-4} M monothioglycerol (Sigma-

Aldrich), 150 μg/mL transferrin (Roche), and with Matrigel (Cat#354277, Corning, NY, USA)),

10 μM Y-27632 and 2 ng/mL human recombinant BMP4 (R&D Systems) and then cultured in
ultra-low attachment 6 well plate (Corning) in $5\%CO_2/5\%$ O_2. After 24 h, 1.5 mL CDM with
6 ng/mL human recombinant activin A (R&D Systems) and 5 ng/mL bFGF (R&D Systems) were
added into the wells. After 3 days, medium was changed to 3 mL CDM with 10 ng/mL VEGF
(R&D Systems), SB431542 (Sigma-Aldrich), Dorsomorphin (Sigma-Aldrich) and 1 μM IWP-3
(Stemgent). After 7 days, medium was changed to 3 mL CDM with 10 ng/mL VEGF (R&D
Systems). The medium was changed every 2 days. The cells were collected at 15 days after
differentiation and analyzed by flow cytometry.
536
Dopaminergic progenitor cells differentiation
Dopaminergic progenitor cells differentiation was performed according to a modified method
described previously (Doi *et al*, 2020). Both large-scale suspension cultured hiPSCs and typical
adherent-cultured hiPSC were collected by centrifugation after dissociation into single cell with
TrypLE Select. The cells were suspended in 1 mL dopaminergic progenitor cells differentiation
medium-1 (DPM-1), consisting of Glasgow's minimum essential medium (Thermo Fisher)
supplemented with 8% knockout serum replacement (Thermo Fisher Scientific), 1% MEM Non-
Essential Amino Acids Solution (Thermo Fisher Scientific), 1 mM sodium pyruvate (Sigma-
Aldrich), 0.1 mM 2-mercaptoethanol (FujiFilm Wako) and 100 nM LDN193189 (Stemgent), and
were then seeded in iMatrix-coated 24-well plates at a density of 1×10^6 cells/well with 10 μM Y-
27632 and 500 nM A-83-01 (FujiFilm Wako). After 1 and 2 days, medium was changed to DPM-
1 with 500 nM A-83-01, 100 ng/mL recombinant human FGF8 (FujiFilm Wako), and 2 μM
purmorphamine (FujiFilm Wako). After 3 to 6 days, medium was changed every day to DPM-1
with 500 nM A-83-01, 100 ng/mL recombinant human FGF8, 2 μM purmorphamine, and 3 μM
CHIR99021 (FujiFilm Wako). After 7 to 11 days, medium was changed every day to DPM-1
with 3 μM CHIR99021. On differentiation day 12, cells were dissociated using TrypLE select
and suspended in the dopaminergic progenitor cells differentiation medium-2 (DPM-2),
consisting Neurobasal medium (Thermo Fisher Scientific), 2% B27 supplement (without Vitamin
A, Thermo Fisher), 1% GlutaMAX (Thermo Fisher Scientific), 10 ng /mL human recombinant
glial cell-derived neurotrophic factor (FujiFilm Wako), 200 mM ascorbic acid, 20 ng / mL human
recombinant brain-derived neurotrophic factor (FujiFilm Wako), and 400 μM dibutyryl cAMP
(FujiFilm Wako), and were then plated on U-shaped 96-well plates (Thermo Fisher Scientific) at
a density of 2×10^4 cells/150 μL/well with 10 μM Y-27632. After 15 to 26 days, half of medium
was changed every 2 days to DPM-2. Dopaminergic progenitor cells differentiation efficiency
was analyzed by flow cytometry at 12 days after differentiation. On differentiation day 12, cells
were labeled with anti-CORIN antibody (Clone 5B6, Sigma-Aldrich) and secondary antibody
(A11001, Thermo Fisher).
564
Hepatocyte differentiation

Hepatocyte differentiation was performed according to a modified method described previously
(Si-Tayeb *et al*, 2010). Both large-scale suspension-cultured and typical adherent-cultured hiPSCs
were collected by centrifugation after dissociation into single cell with TrypLE Select. These
hiPSCs were cultured on Matrigel-coated 6-well plates until 80% confluency with StemFit
medium (Ajinomoto). To initiate differentiation, the medium was changed to 1 mL Hepatocyte
differentiation medium (HDM) consisting of RPMI1640 medium (Thermo Fisher) supplemented
with 1× GlutaMax (Thermo Fisher) and 1× B27 Supplement minus vitamin A, with the addition
of 100 ng/mL Activin A (R&D Systems). After 1 to 3 days, medium was changed every day to
HDM with 100 ng/mL Activin A. After 4 to 8 days, medium was changed every day to HDM
with 20 ng/mL human recombinant BMP-4 (R&D Systems) and 20 ng/mL human recombinant
FGF-4 (R&D Systems). After 9 to 13 days, medium was changed every day to HDM with 20
ng/mL human recombinant HGF (R&D Systems). After 14 to 24 days, medium was changed
every 2 days to Hepatocyte Culture Medium (HCM; Lonza) with 20 ng/mL human recombinant
Oncostatin M (R&D Systems). On differentiation day 25, the albumin concentration in the culture
supernatant was determined by ELISA using an anti-human albumin antibody (Betyl
Laboratories, Catalog # E88-129).
582

Teratoma formation
Suspension-cultured hiPSCs were dissociated with Accutase and then resuspended with 1 mL of
StemFit AK02N medium supplemented with 10 μM Y-27632. 1 x 10^6 cells were collected by
centrifugation at $200 \times g$ for 3 m and suspended in ice-cold 50% Matrigel solution: StemFit
AK02N with 10 μM Y-27632 = 1:1). Cells were injected into testis or leg of NOD.Cg-Prkdcscid
Il2rgtm1Wjl/SzJ (NSG) mice (The Jackson Laboratory, Bar Harbor, ME, USA) using an 18-G
needle. Two to three months later, teratomas were collected and fixed with 4% paraformaldehyde
in D-PBS. Paraffin-embedded sectioning and hematoxylin-eosin (HE) staining was performed.
Three germ layer derivatives were observed under CKX53 microscope with a DP22 camera and
CellSens software (EVIDENT, Tokyo, Japan).
593

Karyotyping
For virtual karyotyping, genomic DNA (gDNA) was extracted from hiPSCs using a DNeasy
Blood & Tissue Kit (Cat#69504, QIAGEN, Hulsterweg, Netherland) and was used for microarray
assay. Virtual karyotyping was performed with GeneChip Scanner System 3000 using Karyostat
Assay arrays (Cat#905403, both from Thermo Fisher Scientific, MA, US) according to the
manufacturer's protocol. Data were analyzed using the Chromosome Analysis Suite (ChAS) and
Affymetrix GeneChip Command Console software programs.
G-band analysis was performed using the common Giemsa staining method with hiPSCs
fixed by Carnoy's fixtative (3:1 ratio of methanol: glacial acetic acid).
603

qRT-PCR

Total RNA was extracted with a FastGene RNA premium kit (Cat#FG-81250, NIPPON Genetics,

Tokyo, Japan) and used for reverse transcription reaction. cDNA was synthesized by using a

ReverTra Ace qPCR RT kit (Cat#FSQ-101, TOYOBO, Osaka, Japan) with random primers.

Real-Time qPCR reactions were performed with a QuantStudio 3 System (Thermo Fisher

Scientific) using THUNDERBIRD Probe qPCR Mix (Cat#QPS-101, TOYOBO) with TaqMan

probes (listed in Supplementary Table 2) (Thermo Fisher Scientific) according to the

manufacturer's instructions. Gene expression was described as the fold change relative to the

control sample value (ΔΔCt method) after normalization to the corresponding GAPDH or β-

Actin values, unless otherwise specified. For the residual SeV detection, data were normalized to

the expression of GAPDH and displayed as relative fold increase to hiPSC line established with

episomal vector (WTc11). SeVdp-infected fibroblasts on passage 1 (SeV-34 and -39) were used

as positive controls.

Immunocytochemistry

Immunocytochemistry was performed on adherent cells. Suspension-cultured hiPSCs were

dissociated with Accutase and transiently cultured as adherent in iMatrix-coated culture dishes for

3 h before immunocytochemistry. The cells were fixed with 4% paraformaldehyde in D-PBS for

10 min at room temperature, then permeabilized in D-PBS containing 0.1% Triton X-100 for

another 10 min. Cells were incubated with primary antibodies in D-PBS containing 0.1% Bovine

Serum Albumin (BSA; Cat#017-22231, FUJIFILM Wako Pure Chemical Corporation, Tokyo,

Japan) overnight at 4 °C. The secondary antibodies were incubated for 1 hour at room

temperature in D-PBS containing 0.1% BSA. Fluoro-KEEPER Antifade Reagent, Non-Hardening

Type with DAPI (Cat#12745-74, Nacalai tesque, Kyoto, Japan) was used for nuclear

counterstaining. Fluorescence images were taken with an all-in-one fluorescent microscope (BZ-

X800; KEYENCE, Osaka, Japan). The primary and secondary antibodies used in this study are

listed in Supplementary Tables 3 and 4, respectively.

Automatic capillary Western blot (Simple Western Assays)

To detect the expression of PAX6 or SOX17, adherent- or suspension-cultured PAX6-TEZ or

SOX17-TEZ hiPSCs were collected on day 10 (Passage 2). The cell lysate was prepared from $1 \times$

10^6 cells with sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) sample

buffer solution without 2-ME (2x) (Cat#30567-12, Nacalai Tesque) supplemented with 100 mM

Dithiothreitol (DTT) (Cat#14130-41, Nacalai Tesque). To detect phosphorylated PKCβ, cell

lysates were collected on day 5 of suspension-cultured hiPSCs with extraction buffer mentioned

above, containing 1% phosphatase inhibitor cocktail (Cat#07574-61, Nacalai Tesque). Samples

were denatured at 95°C for 5 min. Western blotting was performed with a capillary automatic

western blotting device (Simple Western, Wes; Bio-Techne, CA, USA). Jes/Wes 12- to 230-kDa

separation module for Wes, 8 × 25 capillary cartridges (Cat#SM-W004, ProteinSimple), and

Anti-Rabbit/Goat/Mouse Detection Module kit (Cat#DM-002/DM-001/DM/006, ProteinSimple)

were used. Preparation of reagents and sample loading were conducted according to the

manufacturer's instructions. The data were analyzed and quantified by compass for simple

Western software program (ProteinSimple). GAPDH was used as the reference for normalization.

The primary antibodies used in this study are listed in Supplementary Table 3.

Flow cytometry

Flow cytometry of self-renewal markers, TRA-1-60 and SSEA4 was performed as described in

our previous study. Briefly, adherent- or suspension-cultured hiPSCs were dissociated with 0.5

mM EDTA solution in D-PBS or Accutase, respectively. Dissociated cells (0.5 or 1×10^6 cells)

were collected and centrifuged at $200 \times$ g for 3 minutes and resuspended in 500 or 1000 μL PBS

containing 0.1% BSA and 0.5 mM EDTA, then incubated with or without anti-TRA-60 antibody

or anti-SSEA4 antibody for 1 hour at 4 °C under oblique light conditions. After washing with

PBS, cells were resuspended in 500 mL D-PBS containing 0.1% BSA and 0.5 mM EDTA, and

filtrated with 35-mm cell strainer (Cat#352235, FALCON, Thermo Fisher Scientific). For PAX6-

TEZ and SOX17-TEZ hiPSC lines, adherent- or suspension-cultured hiPSCs were dissociated

into single cells and collected cells were resuspended in 500 mL PBS containing 0.1% BSA and

0.5 mM EDTA. After passing through the 35-mm cell strainer, cells were analyzed for tdTomato.

The primary and secondary antibodies used in this study are listed in Supplementary Table 3 and

4, respectively. Flow cytometry was performed with SH800 cell Sorter (SONY, Tokyo, Japan).

RNA-Seq

Total RNA was extracted using the FastGene RNA premium kit, and strand-specific library

preparation was performed. The prepared library was sequenced using a NovaSeq6000 (Illumina,

Inc, CA, USA). Sequencing was performed in a 150 bp × 2 paired-end conFigureuration with a

data output of about 6 Gb per sample (~ 20 million paired reads). Library preparation and

sequencing was performed in GENEWIZ (Azenta, MA, USA). To identify differentially

regulated genes, sequencing data was analyzed with a CLC Genomics Workbench (QIAGEN,

Hulsterweg, the Neterlands) and R package, edgeR (v3.30.3) in R language (v 4.1.0). An MA plot

(log2 fold change versus mean average expression) comparing transcriptomes between

suspension- and adherent-culture from RNA-seq data. Transcripts with log2 fold change $\geqq$ 2 or $\leqq$

-2 (FDR <0.01) are highlighted with red and blue dots, respectively. The extracted genes were

analyzed for gene ontology enrichment and biological pathways using the R package,

clusterProfiler (v3.16.1). GSEA was also performed using clusterProfiler. An enrichment map

was used to visualize the GSEA results with enrichment score (ES) and false discovery rate

(FDR) values. Software used for the analysis is listed in Supplementary Table 5.

Statistical analysis

Statistical analysis was performed using Student's t-test or Dunnett's test for multiple

comparisons with a single control condition. P values < 0.05 were considered statistically

significant. *, ** or *** in the graphs indicate P<0.05, P<0.01 or P<0.001, respectively. No
statistical methods were used to predetermine sample size. The experiments were not randomized
and the investigators were not blinded to allocation during experiments and outcome assessment.
686

Acknowledgments
We thank Dr. Bruce Conklin for providing us with the WTC11 hiPSC line; Dr. Shinya
Yamanaka, Dr. Kazutoshi Takahashi, Dr. Masato Nakagawa, and Dr. Keisuke Okita for providing
us with 201B7, 454E2, 1231A3, 1383D6, and Ff-I14s04 hiPSC lines. We would like to express
our sincere gratitude to Dr. Kaoru Saijo, Mr. Daiki Kondo and Mr. Masahiko Yamada for their
technical assistance, and Ms. Kumiko Omori for her administrative support. This work was
funded by a collaborative research grant from Kaneka corporation. We would like to thank
Editage [http://www.editage.com] for editing and reviewing this manuscript for English language.

Author Contributions: Conceptualization: YHA, Methodology: MMT, SK, CW, KT,MI, YHA,
Investigation: MMT, SK, YHE, TW, TS, YA, KT, MI, MN, MU, TMK, MN, KN, YN, YHA,
Visualization: MMT, SK, TS, YH, Funding acquisition: MMT, TN, YK, YHA, Project
administration: MMT, TN, YK, YHA, Supervision: MMT, TN, YN, NT, YK, YHA, Writing –
original draft: MMT, KS, MT, YHA, Writing – review & editing: MMT, YHA

Competing Interest Statement: Y. HA received collaborative research grant from KANEKA
corporation during this study. S. K., K.T., M. I., Y.K., and T.N. were employed with KANEKA
corporation during this study. M.M.T., S.K. K.T., M. I., Y.K., and Y.HA. are inventors of patents
arising from this work (WO/2021/162090 and PCT/JP2020/005255). The other authors declare no
competing interests.

Classification: Applied Biological Sciences, Cell Biology, and Developmental Biology in
BIOLOGICAL SCIENCES

Keywords: induced pluripotent stem cells (iPSCs), suspension culture, spontaneous
differentiation, Protein Kinase C (PKC), WNT

References

Amit M, Chebath J, Margulets V, Laevsky I, Miropolsky Y, Shariki K, Peri M, Blais I, Slutsky
G, Revel M *et al* (2010) Suspension culture of undifferentiated human embryonic and induced
pluripotent stem cells. *Stem Cell Rev Rep* 6: 248-259
Amit M, Laevsky I, Miropolsky Y, Shariki K, Peri M, Itskovitz-Eldor J (2011) Dynamic
suspension culture for scalable expansion of undifferentiated human pluripotent stem cells. *Nat*
Protoc 6: 572-579
Bayerl J, Ayyash M, Shani T, Manor YS, Gafni O, Massarwa R, Kalma Y, Aguilera-Castrejon A,
Zerbib M, Amir H *et al* (2021) Principles of signaling pathway modulation for enhancing human
naive pluripotency induction. *Cell Stem Cell* 28: 1549-1565 e1512
Bredenkamp N, Yang J, Clarke J, Stirparo GG, von Meyenn F, Dietmann S, Baker D, Drummond
R, Ren Y, Li D *et al* (2019) Wnt Inhibition Facilitates RNA-Mediated Reprogramming of Human
Somatic Cells to Naive Pluripotency. *Stem Cell Reports* 13: 1083-1098
Chen B, Dodge ME, Tang W, Lu J, Ma Z, Fan CW, Wei S, Hao W, Kilgore J, Williams NS *et al*
(2009) Small molecule-mediated disruption of Wnt-dependent signaling in tissue regeneration
and cancer. *Nat Chem Biol* 5: 100-107
Chen KG, Mallon BS, McKay RD, Robey PG (2014) Human pluripotent stem cell culture:
considerations for maintenance, expansion, and therapeutics. *Cell Stem Cell* 14: 13-26
Dang SM, Gerecht-Nir S, Chen J, Itskovitz-Eldor J, Zandstra PW (2004) Controlled, scalable
embryonic stem cell differentiation culture. *Stem Cells* 22: 275-282
Doi D, Magotani H, Kikuchi T, Ikeda M, Hiramatsu S, Yoshida K, Amano N, Nomura M,
Umekage M, Morizane A *et al* (2020) Pre-clinical study of induced pluripotent stem cell-derived
dopaminergic progenitor cells for Parkinson's disease. *Nat Commun* 11: 3369
Dutta D, Ray S, Home P, Larson M, Wolfe MW, Paul S (2011) Self-renewal versus lineage
commitment of embryonic stem cells: protein kinase C signaling shifts the balance. *Stem Cells*
29: 618-628
Dziedzicka D, Tewary M, Keller A, Tilleman L, Prochazka L, Ostblom J, Couvreu De
Deckersberg E, Markouli C, Franck S, Van Nieuwerburgh F *et al* (2021) Endogenous suppression
of WNT signalling in human embryonic stem cells leads to low differentiation propensity towards
definitive endoderm. *Sci Rep* 11: 6137
Elanzew A, Sommer A, Pusch-Klein A, Brustle O, Haupt S (2015) A reproducible and versatile
system for the dynamic expansion of human pluripotent stem cells in suspension. *Biotechnol J*
10: 1589-1599
Funakoshi S, Miki K, Takaki T, Okubo C, Hatani T, Chonabayashi K, Nishikawa M, Takei I,
Oishi A, Narita M *et al* (2016) Enhanced engraftment, proliferation, and therapeutic potential in
heart using optimized human iPSC-derived cardiomyocytes. *Sci Rep* 6: 19111

Fusaki N, Ban H, Nishiyama A, Saeki K, Hasegawa M (2009) Efficient induction of transgene-
free human pluripotent stem cells using a vector based on Sendai virus, an RNA virus that does
not integrate into the host genome. *Proc Jpn Acad Ser B Phys Biol Sci* 85: 348-362

Gschwendt M, Dieterich S, Rennecke J, Kittstein W, Mueller HJ, Johannes FJ (1996) Inhibition
of protein kinase C mu by various inhibitors. Differentiation from protein kinase c isoenzymes.
FEBS Lett 392: 77-80

Guo G, von Meyenn F, Rostovskaya M, Clarke J, Dietmann S, Baker D, Sahakyan A, Myers S,
Bertone P, Reik W *et al* (2017) Epigenetic resetting of human pluripotency. *Development* 144:
2748-2763

Hayashi Y, Furue MK (2016) Biological Effects of Culture Substrates on Human Pluripotent
Stem Cells. *Stem Cells Int* 2016: 5380560

Hayashi Y, Hsiao EC, Sami S, Lancero M, Schlieve CR, Nguyen T, Yano K, Nagahashi A, Ikeya
M, Matsumoto Y *et al* (2016) BMP-SMAD-ID promotes reprogramming to pluripotency by
inhibiting p16/INK4A-dependent senescence. *Proc Natl Acad Sci U S A* 113: 13057-13062

Horiguchi I, Sakai Y (2016) Serum replacement with albumin-associated lipids prevents excess
aggregation and enhances growth of induced pluripotent stem cells in suspension culture.
Biotechnol Prog 32: 1009-1016

Hunt MM, Meng G, Rancourt DE, Gates ID, Kallos MS (2014) Factorial experimental design for
the culture of human embryonic stem cells as aggregates in stirred suspension bioreactors reveals
the potential for interaction effects between bioprocess parameters. *Tissue Eng Part C Methods*
20: 76-89

Ibuki M, Horiguchi I, Sakai Y (2019) A novel tool for suspension culture of human induced
pluripotent stem cells: Lysophospholipids as a cell aggregation regulator. *Regen Ther* 12: 74-82

Jirousek MR, Gillig JR, Gonzalez CM, Heath WF, McDonald JH, 3rd, Neel DA, Rito CJ, Singh
U, Stramm LE, Melikian-Badalian A *et al* (1996) (S)-13-[(dimethylamino)methyl]-10,11,14,15-
tetrahydro-4,9:16, 21-dimetheno-1H, 13H-dibenzo[e,k]pyrrolo[3,4-
h][1,4,13]oxadiazacyclohexadecene-1,3(2H)-d ione (LY333531) and related analogues: isozyme
selective inhibitors of protein kinase C beta. *J Med Chem* 39: 2664-2671

Kehoe DE, Jing D, Lock LT, Tzanakakis ES (2010) Scalable stirred-suspension bioreactor culture
of human pluripotent stem cells. *Tissue Eng Part A* 16: 405-421

Khan SA, Park KM, Fischer LA, Dong C, Lungjangwa T, Jimenez M, Casalena D, Chew B,
Dietmann S, Auld DS *et al* (2021) Probing the signaling requirements for naive human
pluripotency by high-throughput chemical screening. *Cell Rep* 35: 109233

Kim MH, Kino-Oka M (2020) Bioengineering Considerations for a Nurturing Way to Enhance
Scalable Expansion of Human Pluripotent Stem Cells. *Biotechnol J* 15: e1900314

Kim MH, Takeuchi K, Kino-Oka M (2019) Role of cell-secreted extracellular matrix formation in
aggregate formation and stability of human induced pluripotent stem cells in suspension culture. *J
Biosci Bioeng* 127: 372-380

Kinehara M, Kawamura S, Tateyama D, Suga M, Matsumura H, Mimura S, Hirayama N, Hirata M, Uchio-Yamada K, Kohara A *et al* (2013) Protein kinase C regulates human pluripotent stem cell self-renewal. *PLoS One* 8: e54122

Kitano Y, Nishimura S, Kato TM, Ueda A, Takigawa K, Umekage M, Nomura M, Kawakami A, Ogawa H, Xu H *et al* (2022) Generation of hypoimmunogenic induced pluripotent stem cells by CRISPR-Cas9 system and detailed evaluation for clinical application. *Mol Ther Methods Clin Dev* 26: 15-25

Krawetz R, Taiani JT, Liu S, Meng G, Li X, Kallos MS, Rancourt DE (2010) Large-scale expansion of pluripotent human embryonic stem cells in stirred-suspension bioreactors. *Tissue Eng Part C Methods* 16: 573-582

Kwok CK, Ueda Y, Kadari A, Gunther K, Ergun S, Heron A, Schnitzler AC, Rook M, Edenhofer F (2018) Scalable stirred suspension culture for the generation of billions of human induced pluripotent stem cells using single-use bioreactors. *J Tissue Eng Regen Med* 12: e1076-e1087

Lam AT, Chen AK, Ting SQ, Reuveny S, Oh SK (2016) Integrated processes for expansion and differentiation of human pluripotent stem cells in suspended microcarriers cultures. *Biochem Biophys Res Commun* 473: 764-768

Nakanishi M, Kurisaki A, Hayashi Y, Warashina M, Ishiura S, Kusuda-Furue M, Asashima M (2009) Directed induction of anterior and posterior primitive streak by Wnt from embryonic stem cells cultured in a chemically defined serum-free medium. *FASEB J* 23: 114-122

Nath SC, Nagamori E, Horie M, Kino-Oka M (2017) Culture medium refinement by dialysis for the expansion of human induced pluripotent stem cells in suspension culture. *Bioprocess Biosyst Eng* 40: 123-131

Nishimura K, Sano M, Ohtaka M, Furuta B, Umemura Y, Nakajima Y, Ikehara Y, Kobayashi T, Segawa H, Takayasu S *et al* (2011) Development of defective and persistent Sendai virus vector: a unique gene delivery/expression system ideal for cell reprogramming. *J Biol Chem* 286: 4760-4771

Oh SK, Chen AK, Mok Y, Chen X, Lim UM, Chin A, Choo AB, Reuveny S (2009) Long-term microcarrier suspension cultures of human embryonic stem cells. *Stem Cell Res* 2: 219-230

Okita K, Matsumura Y, Sato Y, Okada A, Morizane A, Okamoto S, Hong H, Nakagawa M, Tanabe K, Tezuka K *et al* (2011) A more efficient method to generate integration-free human iPS cells. *Nat Methods* 8: 409-U452

Olmer R, Lange A, Selzer S, Kasper C, Haverich A, Martin U, Zweigerdt R (2012) Suspension culture of human pluripotent stem cells in controlled, stirred bioreactors. *Tissue Eng Part C Methods* 18: 772-784

Rajendran G, Dutta D, Hong J, Paul A, Saha B, Mahato B, Ray S, Home P, Ganguly A, Weiss ML *et al* (2013) Inhibition of protein kinase C signaling maintains rat embryonic stem cell pluripotency. *J Biol Chem* 288: 24351-24362

Ring DB, Johnson KW, Henriksen EJ, Nuss JM, Goff D, Kinnick TR, Ma ST, Reeder JW,

Samuels I, Slabiak T *et al* (2003) Selective glycogen synthase kinase 3 inhibitors potentiate

insulin activation of glucose transport and utilization in vitro and in vivo. *Diabetes* 52: 588-595

Rohani L, Borys BS, Razian G, Naghsh P, Liu S, Johnson AA, Machiraju P, Holland H, Lewis

IA, Groves RA *et al* (2020) Stirred suspension bioreactors maintain naive pluripotency of human

pluripotent stem cells. *Commun Biol* 3: 492

Shafa M, Sjonnesen K, Yamashita A, Liu S, Michalak M, Kallos MS, Rancourt DE (2012)

Expansion and long-term maintenance of induced pluripotent stem cells in stirred suspension

bioreactors. *J Tissue Eng Regen Med* 6: 462-472

Si-Tayeb K, Noto FK, Nagaoka M, Li J, Battle MA, Duris C, North PE, Dalton S, Duncan SA

(2010) Highly efficient generation of human hepatocyte-like cells from induced pluripotent stem

cells. *Hepatology* 51: 297-305

Singh H, Mok P, Balakrishnan T, Rahmat SN, Zweigerdt R (2010) Up-scaling single cell-

inoculated suspension culture of human embryonic stem cells. *Stem Cell Res* 4: 165-179

Steiner D, Khaner H, Cohen M, Even-Ram S, Gil Y, Itsykson P, Turetsky T, Idelson M,

Aizenman E, Ram R *et al* (2010) Derivation, propagation and controlled differentiation of human

embryonic stem cells in suspension. *Nat Biotechnol* 28: 361-364

Sumi T, Tsuneyoshi N, Nakatsuji N, Suemori H (2008) Defining early lineage specification of

human embryonic stem cells by the orchestrated balance of canonical Wnt/beta-catenin,

Activin/Nodal and BMP signaling. *Development* 135: 2969-2979

Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, Yamanaka S (2007)

Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 131:

861-872

Takashima Y, Guo G, Loos R, Nichols J, Ficz G, Krueger F, Oxley D, Santos F, Clarke J,

Mansfield W *et al* (2014) Resetting transcription factor control circuitry toward ground-state

pluripotency in human. *Cell* 158: 1254-1269

Tannenbaum SE, Reubinoff BE (2022) Advances in hPSC expansion towards therapeutic entities:

A review. *Cell Prolif*: e13247

Toullec D, Pianetti P, Coste H, Bellevergue P, Grand-Perret T, Ajakane M, Baudet V, Boissin P,

Boursier E, Loriolle F *et al* (1991) The bisindolylmaleimide GF 109203X is a potent and

selective inhibitor of protein kinase C. *J Biol Chem* 266: 15771-15781

Tran TH, Wang X, Browne C, Zhang Y, Schinke M, Izumo S, Burcin M (2009) Wnt3a-induced

mesoderm formation and cardiomyogenesis in human embryonic stem cells. *Stem Cells* 27: 1869-

1878

Tsukamoto S, Nakade K, Wakabayashi T, Nakashima K, Takami M, Hemmi Y, Kuramochi Y,

Shimizu T, Arai Y, Matsuo-Takasaki M *et al* (2021) Generation of two ISL1-tdTomato reporter

human induced pluripotent stem cell lines using CRISPR-Cas9 genome editing. *Stem Cell Res* 53:

102363

Vijayaragavan K, Szabo E, Bosse M, Ramos-Mejia V, Moon RT, Bhatia M (2009) Noncanonical
Wnt signaling orchestrates early developmental events toward hematopoietic cell fate from
human embryonic stem cells. *Cell Stem Cell* 4: 248-262
Wang Y, Chou BK, Dowey S, He C, Gerecht S, Cheng L (2013) Scalable expansion of human
induced pluripotent stem cells in the defined xeno-free E8 medium under adherent and
suspension culture conditions. *Stem Cell Res* 11: 1103-1116
Woll PS, Morris JK, Painschab MS, Marcus RK, Kohn AD, Biechele TL, Moon RT, Kaufman
DS (2008) Wnt signaling promotes hematoendothelial cell development from human embryonic
stem cells. *Blood* 111: 122-131
Xu C, Inokuma MS, Denham J, Golds K, Kundu P, Gold JD, Carpenter MK (2001) Feeder-free
growth of undifferentiated human embryonic stem cells. *Nat Biotechnol* 19: 971-974
Yu J, Vodyanik MA, Smuga-Otto K, Antosiewicz-Bourget J, Frane JL, Tian S, Nie J, Jonsdottir
GA, Ruotti V, Stewart R *et al* (2007) Induced pluripotent stem cell lines derived from human
somatic cells. *Science* 318: 1917-1920
Zweigerdt R, Olmer R, Singh H, Haverich A, Martin U (2011) Scalable expansion of human
pluripotent stem cells in suspension culture. *Nat Protoc* 6: 689-700

Figures and Tables

878 **Figure. 1. HiPSCs maintained under suspension-culture conditions undergo spontaneous**

879 **differentiation.** (A) Schematics representing hiPSCs in suspension-culture conditions. (B) Phase-

880 contrast images of adherent- or suspension-cultured hiPSCs on day 5 (Passage 1 (P1)) and day 10

(Passage 2 (P2)). Scale bars: 400 μm. (C) Gene expression in hiPSCs cultured under adherent or
suspension conditions on P2. Bar graphs show the mean ± SE (n=3). P-values were statistically
analyzed by Student's t-test. (D) Phase-contrast and fluorescent images of adherent or suspension-
cultured reporter hiPSCs on P2. White arrowheads indicate spontaneous expression of SOX17
and PAX6 in suspension conditions. Scale bars: 400 μm. (E) Quantification of hiPSCs
spontaneous differentiation with flow cytometry. (F) Averaged tdTomato-positive cell ratio (%)
from flow cytometry data (mean ± SE from n=3). P-values were statistically analyzed by
Student's t-test. (G) An MA plot comparing transcriptomes between suspension- and adherent-
culture from RNA-seq data. The representative gene name is shown in the plot. (H) GSEA on the
gene sets of suspension-cultured hiPSCs to adherent cultures. Adjusted P-values are shown as
blue to red from low to high values. (I and J) GOEA on the gene sets of suspension-cultured
hiPSCs to adherent culture. Results are ranked by significance (p-adjusted value) and/or counted
gene numbers.

Figure 2. PKC inhibitors suppress spontaneous differentiation of hiPSCs into neural ectoderm in suspension-culture conditions. (A) Schematics. (B) Phase-contrast pictures of suspension-cultured hiPSCs in the presence of Wnt signaling inhibitors (IWP-2 and IWR-1-endo) or activator (CHIR99021). Scale bars: 400 μm. (C) Gene expression of hiPSCs in suspension-

culture conditions with or without IWP2, IWR-1-endo or CHIR99021. RT-qPCR was performed

on day 10 samples (P2). Gene expressions were normalized to GAPDH and displayed as relative

fold increase to adherent-cultured samples. Bar graphs show the mean ± SE. P-values were

statistically analyzed using Dunnett's multiple comparisons test. (D) Screening of inhibitory

activity of candidate molecules on neuroectoderm differentiation in suspension-cultured hiPSCs.

Candidate molecules were added in combination as shown. Results are displayed as relative fold

increase to suspension-cultured samples without pharmacological treatment. n=1. (E) Phase-

contrast images of suspension-cultured hiPSCs on P2 in the presence of PKC inhibitors (Go6983

or GF109203X) alone, or in combination with IWR-1-endo. Scale bars: 400 μm. (F) Suspension-

cultured hiPSCs gene expression in the presence of IWR-1-endo or with combined IWR-1-endo

and different doses of PKC inhibitors. Results are displayed as relative fold increase to adherent-

culture. Data are presented as mean ± SE (n=3).

Figure 3. The inhibitors of WNT and PKCβ efficiently maintain the self-renewal of hiPSCs in suspension conditions. (A) Phase-contrast images of suspension-cultured hiPSCs on P2 in the presence of IWR-1-endo, LY333531, or both. Scale bars: 400 μm. (B) Gene expression of suspension-cultured hiPSCs on P2 in the presence of IWR-1-endo, LY333531, or both. Data are

presented as mean ± SE. P-values were statistically analyzed with Dunnett's test. (C) Hierarchical

clustering of adherent and suspension-cultured hiPSCs on P2 using Ward's method from RNA-

seq data (n=3 in each condition). Ad, Adherent; Sus, Suspension; IWR, IWR-1-endo; LY,

LY333531; IWRLY, IWR-1-endo and LY333531. (D) Principal component analysis (PCA) plot

showing clusters of samples based on similarity. Gene expression variance are displayed as PC1

and PC2. (E) MA plot comparing transcriptomes between IWRLY and Sus. (F) GSEA on the

gene sets of IWRLY to Sus from these RNA-seq data. Statistically significant enrichment is

shown. P-values are represented in blue to red from low to high values. (G and H) GOEA for the

gene sets of IWRLY to Sus from these RNA-seq data. Analysis was performed on down-

regulated genes in (G) and up-regulated genes in (H).

Figure 4. Long-term suspension culture of hiPSCs is maintained by simultaneous suppression of PKCβ and Wnt signals. (A) The number of hiPSCs at passage during long-term suspension-culture every 5 days. Cell culture was performed in the presence of IWR-1-endo plus Go6983 or LY333531. (B) Bar graph indicating average cell numbers for 10 passages (mean ± SE). (C) Phase-contrast images of suspension-cultured hiPSCs on passage 10. Scale bars: 400 μm. (D) Immunocytochemistry of OCT4 on passage 10 samples. Scale bars: 100 μm. (E) Bar graph showing the percentages of OCT4 positive cells. Values were calculated from randomly selected three regions from immunofluorescence images. Data are presented as mean ± SE. P-value was statistically analyzed by Student's t-test. (F) Immunocytochemistry of differentiated cells in EBs from suspension-cultured hiPSCs in the presence of IWR-1-endo; and Go6983 or LY333531. Anti-TUJ1, -SMA and -AFP antibodies were used to detect ectoderm, mesoderm and endoderm differentiation, respectively. Scale bars:100 μm. (G) Chromosomal copy numbers detected with CGH array analysis (Karyostat assay) of suspension-cultured hiPSCs in the presence of IWR-1-endo; and Go6983 (Upper panel) or LY333531 (Lower panel) at passage 10.

Figure 5. Mass suspension-culture of clinical-grade hiPSCs in the presence of PKCβ and Wnt signaling inhibitors. (A) Schematics. (B) Cell viability at seeding. Data are presented as mean ± SE (n=3). P-values were statistically analyzed with Dunnett's multiple comparisons test. (C) Total cell numbers were counted at day 5. Data were presented as mean ± SE (n=3). P-values were statistically analyzed with Dunnett's test. (D) Representative phase-contrast images of these hiPSCs. Scale bar: 1 mm. (E) Representative flow cytometry data of pluripotent markers in these hiPSCs. (F) Quantification of flow cytometry data for the pluripotent markers. Data are presented as mean ± SE (n=3). (G) Karyotypes of these hiPSCs. Left: Representative pictures of G-band

analysis. Right: Table of karyotype results (n=3). The numbers in brackets indicate the cell

numbers examined. (H) *In vitro* differentiation from hiPSCs was assessed using RT-qPCR (mean

± SE) (n=3). P-value was statistically analyzed by Student's t-test. (I) Representative flow

cytometry data for CORIN. (J) Quantification of CORIN-positive cells (mean ± SE, n=3). (K)

Representative flow cytometry data for cardiac Troponin T (cTnT). (L) Quantification of cTnT-

positive cells (mean ± SE, n=3). (M) Albumin secretion levels of hepatocytes differentiated from

hiPSCs (mean ± SE, n=3).

Figure 6. Establishment of single-cell-sorted hiPSC subclones cultured in suspension

conditions supplemented with IWR1-endo and LY3333531. (A) Schematics representing the

establishment of single-cell-derived hiPSC subclones from 201B7 line. Single-cell-sorted cells

were expanded in the culture medium supplemented with IWR-1-endo and LY333531. Formed

colonies were picked on day 14, and expanded by repeating passage every 4-5 days under

suspension conditions. Characteristic analysis was performed on day 28 after single-cell-sorting.

(B) Phase-contrast images of 201B7 on day 28 after single-cell sorting (Clone #2). Scale bars:

500 μm. (C) Represented flow cytometry of TRA-1-60 (Clone #2). (D) Represented

immunocytochemistry of OCT4 (Clone #2). Scale bars: 200 μm. (E) Summary of the

characterization of single cell-sorted clones. Total cell numbers on day 28, the ratio of TRA-1-60-

positive cells (%), and the ratio of OCT4-positive cells (%) are shown (mean ± SE) (n=7).

Figure 7. Establishment of hiPSCs in complete suspension conditions using SeVdp. (A)
Schematics. (B) Phase-contrast images of PBMCs on day 56 after infection. Scale bars: 400 μm.
(C) Immunocytochemistry of OCT4 on bulk-hiPSCs on day 56. Scale bars: 200 μm. (D) Flow
cytometry of TRA-1-60 in bulk-hiPSCs on day 61. (E) Immunocytochemistry of TUJ1, SMA,
and AFP on differentiated cells in EBs from bulk-hiPSCs on day 56. Scale bars: 100 μm. (F) HE
staining of teratoma sections derived from bulk-hiPSCs. White arrowheads indicate
representative tissue structures derived from ectoderm, mesoderm, and endoderm. Scale bars: 100
μm. (G) Chromosomal copy numbers detected with CGH array analysis on bulk-hiPSCs. (H)
Phase-contrast image of an established hiPSC clone at passage 7. Scale bars: 400 μm. (I)
Immunocytochemistry of established clone, F-10, with anti-OCT4 antibody. Scale bars: 200 μm.

(J) The percentage of TRA-1-60-positive cells with flow cytometry on established clone F-10 line. (K) HE staining of teratoma sections derived from F-10 clone line. The details are the same as in (F). (L) Immunocytochemistry in EBs from established F-10 clone. Scale bars: 100 μm. (M) Chromosomal copy numbers of F-10 clone line. (N) Residual SeVdp genomic RNA in established hiPSCs with RT-qPCR.

Figure 1−figure supplement 1. Characterization of PAX6-tdTomato and SOX17-tdTomato reporter lines. (**A**) Schematic of construction of reporter hiPSC lines. CRISPR/Cas9 was employed to knock-in the 2A-tdTomato gene and EF1a promoter-driven Zeocin (Bleomycin) resistance gene sequence (TEZ) to the 3' terminal of SOX17 or PAX6 open reading frame (ORF), respectively. (B) Genotyping of selected clones by genomic PCR. Arrows indicate alleles of wildtype or insertion in the PAX6 (left panel) or SOX17 (right panel) genes. (C) Phase-contrast images of representative colonies and immunocytochemistry of pluripotency markers: OCT4 and NANOG. Left panels: PAX6-TEZ. Right panels: SOX17-TEZ. (D) Flow cytometry of cell surface markers: TRA-1-60 and SSEA4. Left: PAX6-TEZ. Right: SOX17-TEZ. (E) Chromosomal copy numbers detected with CGH array analysis of PAX6-TEZ (top) and SOX17-

TEZ (bottom). (F) Immunocytochemistry of differentiated cells in EBs from established bulk-

hiPSCs on day 56. Anti-TUJ1, -SMA and -AFP antibodies were used. Scale bars: 100 μm. (G)

Differentiation potency of PAX6-TEZ in the neural ectoderm. Left: Flow cytometry of tdTomato

using PAX6-TEZ cultured in neuroectodermal differentiation medium for 7 days. Middle:

Overlay images of phase-contrast and tdTomato fluorescence of PAX6-TEZ cultured in

neuroectodermal differentiation medium (upper) or maintenance medium (lower). Right: Overlay

images of tdTomato fluorescence and endogenous PAX6 protein detected with

immunocytochemistry with anti-PAX6 antibody. (H) Differentiation potency of SOX17-TEZ in

the endoderm. Left: Flow cytometry of tdTomato using SOX17-TEZ cultured in endodermal

differentiation medium for 7 days. Middle: Overlay images of phase-contrast and tdTomato

fluorescence of SOX17-TEZ cultured in endoderm differentiation medium (upper) or

maintenance medium (lower). Right: Overlay images of tdTomato and endogenous SOX17

protein detected by immunocytochemistry with anti-SOX17 antibody.

1011

Figure 2‒figure supplement 1. Effects of PKCβ and IWR-1-endo on suspension-cultured

hiPSCs at the protein level. (A) Simple western blotting of adherent- or suspension-cultured

hiPSCs (SOX17-TEZ) in the presence or absence of IWR-1-endo. SOX17-TEZ cultured in

endoderm differentiation medium (Actinin A + CHIR) was used as a positive control for SOX17

expression. (B) Quantification of SOX17 expression. The protein expression was normalized to

GAPDH. Data are presented as mean ± SE (n=3). (C) Simple western blotting of adherent- or

suspension-cultured hiPSCs (PAX6-TEZ) in the presence or absence of IWR-1-endo and

LY333531. PAX6-TEZ cultured in neurectoderm differentiation medium (SB431542+DMH1)

was used as the positive control for PAX6 expression. (D) Quantification of expression of PAX6.

The protein expression was normalized to GAPDH. Data are presented as mean ± SE (n=3). (E)

Phase-contrast images and representative flow cytometry data of pluripotency marker, TRA-1-60,

in adherent- or suspension-cultured hiPSCs (1383D6 line) on day10 (Passage 2). Suspension-

culture was performed in the presence or absence of IWR-1-end and LY333531. The percentage

of TRA-1-60 positive cells is shown in the right corner. (F) Quantified data for TRA-1-60

positive ratio (%). The percentage of TRA-1-60 positive cells is shown as mean ± SE (n=3).

1027

Figure 2–figure supplement 2. Activity of different type of PKC inhibitors on spontaneous

differentiation of suspension-cultured hiPSCs. (A) Schematic representing suspension culture

of hiPSCs supplemented with PKC inhibitors. Healthy donor-derived hiPSCs (201B7 line) were

grown under continuous agitation in the presence of Y-27632, IWR-1-endo and PKC inhibitors.

Passage was performed every four days. (B) Phase-contrast images of suspension cultured

hiPSCs on day 8 (P2) in the absence or presence of IWR-1-endo and each PKC inhibitors. Scale

bars, 500 μm. (C) Cell growth of hiPSCs (201B7). Graph shows live cell numbers counted on day

4 and day 8 in suspension-culture (n=1). The PKC inhibitors added are labelled in the graphs. (D)

Gene expression in hiPSCs cultured in adherent or suspension conditions with each PKC

inhibitor. RT-qPCR was performed on cDNAs prepared from day 8 (passage 2) samples (n=1).

qPCR was performed for pluripotency (OCT4, SOX2, NANOG) and early differentiation markers

(PAX6, T, SOX17). Gene expression was normalized to beta-ACTIN.

1040

Figure 3–figure supplement 1. Global gene expression of suspension-cultured hiPSCs with IWR-1-endo and LY333531 in comparison to adherent-cultured hiPSCs. (A) MA plot (log2 fold change versus mean average expression) comparing the transcriptomes between adherent- and suspension-cultured hiPSCs treated with IWR-1-endo and LY33353 (IWRLY). Transcripts with log2 fold change ≧ 2 or ≦ -2 (FDR <0.01) are highlighted with red or blue dots,

respectively. The names of the representative genes are shown in the plot. (B) GSEA of the gene

sets of suspension-cultured hiPSCs with IWRLY compared to adherent-cultured hiPSCs.

Analysis was performed on both up-regulated and down-regulated genes. Statistically significant

enrichment is shown. Adjusted P-values are illustrated from blue to red as low to high. (C and D)

GOEA of the gene sets of suspension-cultured hiPSCs with IWRLY compared to adherent-

cultured hiPSCs. Analysis was performed on up-regulated genes in (C) and down-regulated genes

in (D), respectively. Results are ranked by significance (adjusted P values) and/or counted gene

numbers.

1055

Figure 3—figure supplement 2. Up-regulation of PKC genes in suspension-cultured

hiPSCs. (A) RNA expression of conventional PKC isoform genes in hiPSCs as detected by RT-

qPCR. cDNA was prepared from adherent- or suspension-cultured hiPSCs on day 5. The gene

expression was normalized to GAPDH. Results are displayed as relative fold increase to

adherent-culture. Data are presented as mean ± SE (n=3). P-values were statistically analyzed by

Student's t-test. (B) Detection of phosphorylated PKCβ protein (pPKCβ) in suspension cultured

hiPSCs on day 5 using an automated capillary western blot (Simple Western) assay. Adherent-

cultured hiPSCs were used as a control. (C) Quantification of pPKCβ expression. The protein

expression was normalized to GAPDH. Data are shown as mean ± SE (n=3). P-value was

statistically analyzed using Student's t-test. Scale bars, 200 μm.

1066

1067

Figure 4—figure supplement 1. Generality of the inhibitory effects of IWR-1-end and

LY333531 on spontaneous differentiation of hiPSCs cultured in suspension condition. (A)

Schematic representing suspension culture of hiPSCs using StemScale medium (Thermo Fisher
Scientific) as the basic culture medium. hiPSCs were grown under continuous agitation (90 rpm)
in the presence or absence of IWR-1-endo and LY333531. Passage was performed every four
days according to the manufacture's protocol. cDNAs were prepared from day 8 (P2) and used
for RNA seq. (B) Phase-contrast images of suspension cultured hiPSCs on day 8 in the presence
or absence of IWR-1-endo and LY333531. Scale bars, 400 μm. (C) RNA expression of
pluripotency and early differentiated genes in hiPSCs, as detected by RT-qPCR. cDNA was
prepared from suspension-cultured hiPSCs on day 8 in the presence or absence of IWR-1-endo
and LY33353. The gene expression was normalized to GAPDH. Data are presented as mean ± SE
(n=3). (D) Heat map of marker genes for pluripotency and differentiation to three germ layers
from RNA-seq data in suspension-cultured iPSCs without compounds (Sus_1, Sus_2 and Sus_3)
or with IWR-1-endo and LY333531 (IWRLY_1, IWRLY_2 and IWRLY_3). The listed genes
were classified as pluripotent, mesendoderm, mesoderm, endoderm, and ectoderm, as defined in
the Scorecard Assay (Thermo Fisher Scientific).
1084

1085

Figure 5−figure supplement 1. Suspension-culture of hiPSCs supplemented with IWR-1-

1086

endo and LY333531 using bioreactors with continuous agitation. (A) Schematic representing

1087

suspension culture of hiPSCs in a 30 mL bioreactor. Healthy donor-derived hiPSCs (1231A3 and

1088

1383D6 lines) were grown with continuous agitation in the presence or absence of IWR-1-endo

1089

and LY333531. Passages were performed every three days. (B) Graphs showing cell numbers at

1090

each passage (n=1). Upper: suspension-culture of 1231A3 and 1383D6 lines. Lower: adherent-

culture of 1231A3 and 1383D6 lines. (C) Graphs showing glucose concentration in the culture

medium at each culture day. Upper: suspension-culture of 1231A3 and 1383D6 lines. Lower:

adherent-culture of 1231A3 and 1383D6 lines. (D) Graphs showing lactic acid concentration in

the culture medium at each culture day. Upper: suspension-culture of 1231A3 and 1383D6 lines.

Lower: adherent-culture of 1231A3 and 1383D6 lines. (E and F) RT-qPCR analysis of hiPSCs

cultured in 30 mL bioreactor or adherent conditions. cDNA samples were prepared from passage

3 and 5 of the 1231A3 in (E) and 1383D6 in (F). qPCR was performed for *OCT4* and *NANOG*

(pluripotent), *CDX2* (mesoderm), *SOX17* (endoderm) and *PAX6* (neuroectoderm). The gene

expression was normalized to *β-ACTIN*. Bar graph showing the average gene expressions of

passage 3 and 5 (mean ± SE). (G) Flow cytometric analysis of suspension- or adherent-cultured

1231A3 and 1383D6 lines. Positive cells (%) of pluripotency marker proteins: NANOG, OCT4,

and SOX2 are shown. (H) Karyotyping of hiPSCs cultured in 30 mL bioreactor. Cell samples

were prepared on passage 5, and used for G-band karyotyping analysis. A short deletion on

chromosome 9 of 1231A3 (indicated with arrow) was derived from the original line.

1106

1107

Figure 6–figure supplement 1. Re-suspension from frozen stocks of hiPSCs under
suspension culture conditions supplemented with IWR-1-endo and LY333531. (A)
Schematics of re-suspension culture of frozen stocks of single-cells-derived 201B7 clone (Vial 1-
3). (B) Represented phase-contrast images on day 10 (Vial 3). Scale bars: 500 μm. (C)
Represented flow cytometry of TRA-1-60 (Vial 3). (D) Represented immunocytochemistry of
OCT4 (Vial 3). Scale bars: 200 μm. (E) Summary of the characterization of re-suspension-
cultured iPSCs from frozen stocks. Total cell numbers on day 10, the ratio of TRA-1-60-positive
cells (%), and the ratio of OCT4-positive cells (%) are shown (mean ± SE) (n=3).

1116

Figure 7−figure supplement 1. Establishment of hiPSCs in completed suspension conditions using episomal vectors. (A) Flow chart representing the establishment of hiPSCs from PBMCs by electroporation of episomal vectors. (B) Phase-contrast images of transfected cells on day 36 after electroporation of episomal vectors in the absence or presence of IWR-1-endo and

LY333531. Scale bars: 400 μm. (C) Flow cytometry of TRA-1-60 and SSEA4. TRA-1-60 and

SSEA4-psotive cells were sorted as the bulk fraction and further suspended in the absence or

presence of IWR-1-endo and LY333531. The percentages of TRA-1-60 and SSEA4-psotive cells

are shown in the right corner. (D) Phase-contrast images of suspension-cultured cell aggregates

after bulk sorting on passage 5. Scale bars: 400 μm. (E) Flow cytometry of TRA-1-60 and

SSEA4. The percentage of TRA-1-60- and SSEA4-psotive cells are shown in the right corner. (F)

Immunocytochemistry of OCT4 on sorted-cells cultured in the absence (top) or presence (bottom)

of IWR-1-endo and LY333531 for 5 passages. Scale bars: 400 μm. (G) Quantified data for

OCT4-positive cell percentage. Bar graph indicating average OCT4-positive cell percentages

(mean ± SE) (n=3). P-value was statistically analyzed by Student's t-test. (H) Single-cell sorting

and cloning of TRA-1-60 positive cells. Flow cytometry was performed for cell fraction that was

cultured in the presence of IWR-1-endo and LY333531. TRA1-60-positive cells were collected as

single cells in a 96-well plate and expanded in the culture medium supplemented with Y-27632,

IWR-1-endo and LY333531. (I) Representative phase-contrast images of single-sorted cell

aggregates on day 5 and 12. Scale bars, 400 μm. (J) Phase-contrast images of established clones

(B-10, E-4, and E-5). Scale bars: 400 μm. (K) Immunocytochemistry of OCT4 and NANOG in

these clones. The percentage of positive cell are shown. Scale bars: 100 μm. (L) Flow cytometry

of TRA-1-60 in these clones. TRA-1-60 positive percentages are shown in the right corner. (M)

Immunocytochemistry of differentiated cells in EBs from these clones. Anti-TUJ1, -SMA and -

AFP antibodies were used. Scale bars: 100 μm. (N) Hematoxylin-Eosin staining of paraffin-

embedded sections of teratoma derived from these clones. White arrowheads indicate

representative tissue structures of the ectoderm, mesoderm, or endoderm. Scale bars, 100 μm. (O)

Chromosomal copy numbers detected by CGH array analysis (Karyostat assay) of these clones.

1145

1146 **Table 1. The list of published study on scalable suspension culture without microcarriers**
1147 **for hPSCs.**

Article	Medium	Additives	Cell lines tested	Scalability	Transcriptome	Single cell cloning	Direct freeze and thaw	iPSC generation
Matsuo-Takasaki et al. 2023 (This study)	AK02N, AK03N, StemScale	Y-27632, IWR-1-endo, LY333531	hiPSC (WTc11, 201B7, 1383D6, 1231A3, Ff-I14s04)	$\sim10^9$	Yes	Yes	Yes	Yes
Amit et al., 2010, 2011	DMEM/F12 + KSR	Y-27632, LIF, bFGF	hESC (I3, Ie, I5, H9.2, H9, H7, H14) hiPSC (iF4, J1.2.3, C3, C2, KTN7, KTN13)	$\sim10^7$	No	No	No	No
Krewetz et al., 2010	mTeSR1	Y-27632, Rapamycin	hESC (H9)	$\sim10^8$	No	No	No	No
Singh et al., 2010 Zweigerdt et al., 2011 Olmer et al., 2012	KnockOut DMEM + KSR, mTeSR1	Y-27632, bFGF	hESC (hES2, hES3, and ESI049) hiPSC (hCBiPS2, hiPSOCT4eGFP)	$\sim10^8$	No	No	No	No
Steiner et al., 2010	Neurobasal medium + KSR	Y-27632, bFGF, Activin A, Fibronectin, Gelatin, BDNF, NT3, NT4, Nutridoma-CS	hESC (HES1, HES2, H7)	$\sim10^6$	No	No	No	No (Of note, 3 hESC lines derivation)
Wang et al., 2013	TeSR-E8	Y-27632	hiPSC (BC1, TNC1)	$\sim10^8$	No	No	No	No
Hunt et al., 2014	mTeSR1	Y-27632	hESC (H1, H9)	$\sim10^7$	No	No	No	No
Elanzew et al., 2015	mTeSR1, TeSR-E8	Y-27632	hESC (H9) and hiPSC (iLB-C-31f-r1)	$\sim10^7$	No	No	No	No
Horiguchi et al., 2016; Ibuki et al., 2019	TeSR-E8, mTeSR1	Y-27632, KSR, Albumax, LPA, SIP,	hiPSC (TkDN4-M, TkDA3-4, 201B7)	$\sim10^6$	No	No	No	No
Nath et al., 2017	Dialyzed DMEM/F-12	Y-27632, bFGF, TGF-b1	hiPSC (Tic)	$\sim10^8$	No	No	No	No
Kwok et al, 2018	mTeSR1, StemMACS	Y-27632	hiPSC (AR1034ZIMA hiPSC clone1, FS hiPSC clone2)	$\sim10^9$	No	No	No	No
Rohani et al., 2020	mTeSR1, RSeT	Y-27632, Rapamycin	hESC (H1, H9)	$\sim10^7$	Yes	No	No	No

1148

1149 **Supplementary Table 1. The list of chemicals and cytokines used in the screening assay.**

Name	Effect	Manufacturer	Identifier	Concentration
CHIR99021	Wnt signal activator	Fujifilm Wako Pure Chemical	034-23103	$3\,\mu$M
IWP-2	Wnt signaling inhibitor	Fujifilm Wako Pure Chemical	034-24301	$5\,\mu$M
IWR-1-endo	Wnt signaling inhibitor	Fujifilm Wako Pure Chemical	037-25131	$10\,\mu$M
Go6976	PKC (α, β) inhibitor	Fujifilm Wako Pure Chemical	071-06551	$5\,\mu$M
Go6983	PKC (α, β, γ, δ) inhibitor	Fujifilm Wako Pure Chemical	078-06443	$5\,\mu$M
GF109203X	PKC (α, β, γ) inhibitor	Tocris Bioscience	0741	$5\,\mu$M
Enzastaurin	PKC (α, β, γ, ε) inhibitor	Sigma-Aldrich	SML0762	$5\,\mu$M
K-252a	PKC (μ) inhibitor	Sigma-Aldrich	K1639	$5\,\mu$M
LY333531 (Ruboxistaurin)	PKC (β) inhibitor	Sigma-Aldrich	SML0693	$5\,\mu$M
Myricitrin	PKC (α, ε) inhibitor	Sigma-Aldrich	91255	$5\,\mu$M
Sotrastaurin	PKC (α, β, δ, ε, η, θ) inhibitor	Cayman	16726	$5\,\mu$M
ZIP	PKC (ζ) inhibitor	Tocris Bioscience	2549	$5\,\mu$M
Ro-32-0432	PKC (α, β, γ, ε) inhibitor	Tocris Bioscience	1587	$5\,\mu$M
FK506	Calcineurin inhibitor	Fujifilm Wako Pure Chemical	063-06071	100 ng/mL
Cyclosporine A (CsA)	Calcineurin inhibitor	Fujifilm Wako Pure Chemical	031-24931	1 mg/mL
DAPT	γ-secretase inhibitor	Fujifilm Wako Pure Chemical	043-33581	$10\,\mu$M
BMP4 (Human, recombinant)	BMP-SMAD signaling activator	Fujifilm Wako Pure Chemical	020-18851	40 ng/mL
LIF (Human, recombinant)	LIF-STAT signaling activator	Nacalai Tesque	NU0013-1	2000 units/mL
Valproic acid (VPA)	Epigenetic modifier	Fujifilm Wako Pure Chemical	227-01071	0.5 mM
Ascorbic acid	Epigenetic modifier	Nacalai Tesque	13048-42	50 ng/mL

1150

1151 **Supplementary Table 2. The list of Taqman probes and primer sets used in this study.**

Target gene	Taqman Assay ID or sequence
ACTB	F: cctcatgaagatcctcaccga R: ttgccaatggtgatgacctgg
CDX2	F: cacccacagccatagacctac R: gtcagtccaggcaatgcttc
CXCR4	F: actgagaagcatgacggacaag R: aggtagcggtccagactgatg
GAPDH	Hs02786624_g1
GATA6	Hs00232018_m1
NANOG	Hs02387400_g1
OCT4 (POU5F1)	Hs04260367_gH
PAX3	Hs00240950_m1
PAX6	Hs01088114_m1
PDGFRa	F: gctgagcctaatcctctgcc R: actgctcacttccaagaccg
PKCα (PRKCA)	Hs00925200_m1
PKCβ (PRKCB)	Hs00176998_m1
PKCγ (PRKCG)	Hs00177010_m1
SOX1	Hs01057642_s1
SOX2	F: caccaatcccatccacactcac R: gcaaagctcctaccgtaccac
SOX17	Hs00751752_s1
T	Hs00610080_m1
SeV detection	F: accaacaggcggtggtgcaa R: tccaccccaacccctagcgt
PAX6 genotyping	F: ggaattccagtacttcacgtgaaggcatct R: tagaactgaagcggctctaacagccatttt
SOX17 genotyping	F: agaaaggggtgcctttagaggacgggtgtt R: aggaagtgtgtaacactgcttctggcctgc

1152

1153 **Supplementary Table 3. The list of primary antibodies used in this study.**

Name	Host species	Dilutions or Concentrations	SOURCE	IDENTIFIER
Alpha-Fetoprotein (AFP)	Mouse	1 μg/mL	R&D systems	Cat#MAB1368
Alpha-SMA	Mouse	1 μg/mL	R&D systems	Cat#MAB1420
Cardiac Troponin T	Mouse	10 μg/mL	Thermo Fisher Scientific	Cat#MA5-12960
CORIN	Mouse	10 μg/mL	Sigma-Aldrich	Cat#WH0010699M1
GAPDH	Mouse	1 μg/mL	R&D systems	Cat#MAB5718
NANOG	Rabbit	0.5 μg/mL	Reprocell	Cat#RCAB004P-F
OCT-3/4	Mouse	0.4 μg/mL	Santa Cruz	Cat#sc-5279
Oct-4A (C30A3) Rabbit mAb (Alexa Fluor 488 Conjugate)	Rabbit	1:30	Cell Signaling Technology	Cat# 5177S
PAX6	Rabbit	1:400 (Immunocytochemistry)	MBL	Cat#PD022
PAX6	Sheep	5 μg/mL (Wes)	R&D Systems	Cat#AF8150
Anti-PKC beta1 + PKC beta2 (phospho T641)	Rabbit	1:50	Abcam	Cat#ab194749
SOX17	Goat	0.4 μg/mL (Immunocytochemistry) 1 μg/mL (Wes)	R&D systems	Cat#AF1924
FITC Mouse anti-SSEA-4	Mouse	1:5	BD Bioscience	Cat#560126
APC anti-human SSEA4	Mouse	1:20	BioLegend	Cat#330418
Alexa Fluor 488 anti-human TRA1-60-R	Mouse	1:20	BioLegend	Cat#330614
Alexa Fluor 488 Mouse anti-Human TRA-1-60 Antigen	Mouse	1:10	BD Bioscience	Cat#560173
TUJ1	Mouse	1 μg/mL	R&D systems	Cat#MAB1195

1154   **Supplementary Table 4. The list of secondary antibodies used in this study.**

Name	Host species	Dilutions	SOURCE	IDENTIFIER
Anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor Plus 488	Donkey	1:1000	Thermo Fisher Scientific	Cat#A-21202
DyLight 488 anti-rabbit IgG (minimal x-reactivity)	Donkey	1:1000	BioLegend	Cat#406404
Anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor Plus 555	Donkey	1:1000	Thermo Fisher Scientific	Cat#A-31570
Anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 555	Goat	1:1000	Thermo Fisher Scientific	Cat#A-21428
Anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 488	Goat	1:1000	Thermo Fisher Scientific	Cat#A-11001
Donkey Anti-Sheep IgG HRP Affinity Purified PAb	Sheep	1:50	R&D systems	Cat# HAF016

1155

1156 **Supplementary Table 5. Software used in this study.**
1157

Name	Source	URL
clusterProfiler (v3.16.1)	R package	https://bioconductor.org/packages/release/bioc/html/clusterProfiler.html
edgeR (v3.30.3)	R package	https://bioconductor.org/packages/release/bioc/html/edgeR.html
R (version4.1.0)	R package	https://www.r-project.org/
CLC Genomics Workbench (v20)	QIAGEN	https://digitalinsights.qiagen.com/products-overview/discovery-insights-portfolio/analysis-and-visualization/qiagen-clc-genomics-workbench/
Cell Sorter software	Sony	https://www.sonybiotechnology.com/us/instruments/sh800s-cell-sorter/software/
Compass software for Simple Western	Protein Simple	https://www.bio-techne.com/resources/instrument-software-download-center/compass-software-simple-western
BZ-X800 microscope analysis software	Keyence	https://www.keyence.com/landing/microscope/lp_fluorescence.jsp
QuantStudio 3 and 5 Real-Time PCR System Software	Thermo Fisher Scientific	https://www.thermofisher.com/jp/ja/home/global/forms/life-science/quantstudio-3-5-software.html
Chromosome Analysis Suite (ChAS) and Affymetrix GeneChip Command Console software programs	Thermo Fisher Scientific	https://www.thermofisher.com/jp/ja/home/life-science/microarray-analysis/microarray-analysis-instruments-software-services/microarray-analysis-software/chromosome-analysis-suite.html
CellSens microscope software	Evident (Olympus)	https://www.olympus-lifescience.com/en/support/downloads/

1158