Nitrosative stress under microaerobic conditions triggers inositol metabolism in

Pseudomonas extremaustralis

Esmeralda C. Solar Venero[1]; Lucia Giambartolomei[2]; Ezequiel Sosa[3]; Darío Fernández

do Porto [2,3]; Nancy I. López[1,2–] & Paula M. Tribelli [1,2,–,*]

[1]IQUIBICEN-CONICET, Intendente Guiraldes 2160, 1428EGA, Buenos Aires,

Argentina.

[2]Departamento de Química Biológica, Facultad de Ciencias Exactas y Naturales,

Universidad de Buenos Aires, 1428EGA, Buenos Aires, Argentina.

[3]Instituto de Cálculo, Facultad de Ciencias Exactas y Naturales, UBA, Buenos Aires,

Argentina

–: contributed equally to this work

*: Correspondence to: paulatrib@qb.fcen.uba.ar

Abstract

Bacteria are exposed to reactive oxygen and nitrogen species that provoke oxidative

and nitrosative stress which can lead to macromolecule damage. Coping with stress

conditions involves the adjustment of cellular responses, which helps address

metabolic challenges. In this study, we performed a global transcriptomic analysis of

the response of *Pseudomonas extremaustralis* to nitrosative stress, induced by S-

nitrosoglutathione (GSNO), a nitric oxide donor, under microaerobic conditions. The

analysis revealed the upregulation of genes associated with inositol catabolism; a

compound widely distributed in nature whose metabolism in bacteria has aroused

interest. The RNA-seq data also showed heightened expression of genes involved in

essential cellular processes like transcription, translation, amino acid transport and

biosynthesis, as well as in stress resistance including iron-dependent superoxide

dismutase, alkyl hydroperoxide reductase, thioredoxin, and glutathione S-transferase in

response to GSNO. Furthermore, GSNO exposure differentially affected the expression

of genes encoding nitrosylation target proteins, encompassing metalloproteins and

proteins with free cysteine and /or tyrosine residues. Notably, genes associated with

iron metabolism, such as pyoverdine synthesis and iron transporter genes, showed

activation in the presence of GSNO, likely as response to enhanced protein turnover.

Physiological assays demonstrated that *P. extremaustralis* can utilize inositol

proficiently under both aerobic and microaerobic conditions, achieving growth

comparable to glucose-supplemented cultures. Moreover, supplementing the culture

medium with inositol enhances the stress tolerance of *P. extremaustralis* against

combined oxidative-nitrosative stress. Concordant with the heightened expression of

pyoverdine genes under nitrosative stress, elevated pyoverdine production was

observed when *myo*-inositol was added to the culture medium. These findings highlight

the influence of nitrosative stress on proteins susceptible to nitrosylation and iron

metabolism. Furthermore, the activation of *myo*-inositol catabolism emerges as a

protective mechanism against nitrosative stress, shedding light on this pathway in

bacterial systems, and holding significance in the adaptation to unfavorable conditions.

Introduction

Bacterial survival and adaptability in changing environments depend on different

resistance mechanisms against several stress agents. Free radicals are highly reactive

species, mainly originated from energy producing processes like aerobic and anaerobic

respiration, causing damage to different macromolecules. Reactive oxygen species

(ROS) are produced during aerobic respiration, while anaerobic respiration, utilizing

nitrate as electron acceptor, generates reactive nitrogen species (RNS). ROS and RNS

trigger oxidative or nitrosative stress, respectively. RNS include nitric oxide (NO) and

its derivatives peroxynitrite ($ONOO^-$), nitrosothiols formed through reaction with thiol

groups, and nitrotyrosine from nitration of tyrosine by NO, $ONOO^-$ or NO_2^- [1–3].

Nitrosative and oxidative stress can also be induced from various environmental

compounds, like pollutants, xenobiotics, and antimicrobials, and from UV radiation [4].

NO is also produced by the mammalian immune system and acts as a modulator

molecule in both mammals and plants [5–7]. Nitrosative stress can lead to DNA and

macromolecular damage, including protein oxidation. Some residues or structures are

more susceptible to the RNS or ROS attack such as iron-sulfur (Fe-S) clusters, metal

centers, tyrosine and cysteine residues, and thiols [8]. The radical attack to these

residues or clusters results in both reversible and irreversible damage, leading to

alterations in function and structure[9].

Pseudomonas species exhibit a wide variety of energy-generation metabolisms

spanning from aerobic to anaerobic, involving NO_3^- reduction, complete denitrification

process to N_2 and even fermentation of arginine or pyruvate [10,11]. These energy-

generating pathways enable *Pseudomonas* species to thrive in diverse environmental

niches, including soil, water, and host-associated habitats, and they play an integral

role in the ecological and physiological success of these microorganisms.

Pseudomonas extremaustralis is an Antarctic bacterium, capable to growth under

different temperatures and oxygen availability. The denitrification process in this

bacterium is incomplete, due to the absence of *nir* genes which encode the enzymes

required for the reduction of NO_2^- to NO, but interestingly, harbors all the *nor* and *nos*

genes associated with nitric and nitrous oxide reduction, respectively [12].

Bacterial adaptation to different stress conditions entails changes leading to

solve metabolic and structural challenges, such us alterations or damage in DNA, RNA

and proteins and envelopes, among others. Some strategies can involve the utilization

of alternative pathways to exploit different carbon sources. Thus, when *P. putida* was

grown at 10°C, the uptake and assimilation of branched-chain amino acids alongside

the activation of the 2-methylcitrate pathway to generate succinate and pyruvate has

been identified as mechanism employed to cope with stress when central metabolism

is downregulated[13]. Likewise, in cold conditions genes involved in primary

metabolism were downregulated in *P. extremaustralis* whereas genes linked to ethanol

oxidation were activated, showing that this secondary pathway resulted essential for

cold growth [14]. In *P. aeruginosa* has been reported that under oxidative stress the

glyoxylate shunt, an alternative to the tricarboxylic acid cycle, increases bacterial

survival allowing the utilization of acetate and fatty acids as carbon sources [15].

Considering the importance of RNS derived from both anaerobic respiration and

other biotic and abiotic environmental reactions, we evaluated the global transcriptomic

response of *P. extremaustralis* to nitrosative stress under low O_2 tensions using S-

nitrosoglutation (GSNO), as a NO donor compound. Our findings show that exposure

to GSNO under microaerobic conditions prompted adjustments in central metabolic

pathways including iron metabolism along with the upregulation of genes associated

with the inositol catabolism. Notably, *myo*-inositol arises as a carbon source supporting

P. extremaustralis growth and increased nitro-oxidative stress resistance and

pyoverdine production. These findings highlight the versatility and adaptability of *P.*

extremaustralis to withstand nitrosative stress, by displaying an alternative metabolism,

thus contributing to the understanding of the survival mechanisms employed by

microorganisms in extreme environments.

Materials and methods

Strains and culture conditions

P. extremaustralis 14-3b (DSM 25547), a species isolated from the Antarctica [16] was

used through the experiments. Bacterial cultures were grown in Lysogeny Broth

medium (LB) at 30 °C. Microaerobic cultures were incubated in sealed bottles with a

1:2 medium-to-flask volume ratio and 50 rpm agitation. Microaerobic culture's medium

was supplemented with 0.8 g/l KNO_3.

GSNO effect on *P. extremaustralis*' survival and growth was determined in

microaerobic cultures exposed to 1, 10 and 100 µM of GSNO. Bacterial growth was

evaluated by OD_{600nm} measurement after adding the different concentrations of GSNO

at T=0. Optical density was monitored over 24 h at 30min intervals using an automated

plate reader (BMG OPTIMA FLUOstar). For survival experiments, cultures were grown

for 24h and further exposed to the different concentrations of GSNO for 1h. Afterwards,

appropriate dilutions of control cultures (m-C) or cultures exposed to GSNO were

plated in LB agar and incubated at 30°C. Colony-forming units per ml (CFU / ml) was

determined and the survival percentage was calculated with respect to control cultures.

RNA extraction and RNA library preparation

Cultures were microaerobically grown in LB medium for 24 h and then incubated for 1 h

with 100 μM S-nitrosoglutatione (GSNO, Sigma Aldrich) (m-NS condition) or with the

addition of sterile water (m-C). Total RNA was isolated from 6 ml of *P. extremaustralis*

m-C and m-NS cultures using the Trizol method. Samples were treated with DNAse I

and were validated using an Agilent 2100 Bioanalyzer (Agilent Technologies). To

improve the quality of the readings, ribosomal RNA was depleted from the samples,

using the RiboZERO Kit (Illumina), following the manufacturer's instructions. Libraries

were prepared using the TruSeq RNA Library Prep Kit v2 (Illumina). Mass sequencing

was performed using NextSeq 550 platform with a single-end protocol. For each

condition duplicated independent RNA extraction and libraries were used.

RNA-seq data analysis

Reads were preprocessed using the Trimmomatic tool [17] by eliminating adapters and

low-quality sequences. Reads' quality was evaluated using the Fast QC tool

(www.bioinformatics.babraham.ac.uk/projects/fastqc/).

Reads alignment and assembly to the *P. extremaustralis* genome, transcript

identification and abundance quantification was carried out using the Rockhopper

software that use the Bowtie2, Bayesian and Anders and Huber approaches

[18].Reads were normalized per kilobase per million mapped reads (RPKM).

Differential gene expression was considered only with $P < 0.05$ and $Q < 0.05$.

Concordance between the independent replicates for each of the analyzed conditions

was verified by performing a Spearman correlation analysis of normalized counts (Fig.

S1).

Genes were assorted into functional classes using KEGG [19], MetaCyc [20] and String

[21] tools.

Prediction of oxidation and nitrosylation targets

The Target-Pathogen tool [22] was used to identify proteins with possible oxidation and

/ or nitrosylation target sites (Free Cys or Tyr residues or metal biding clusters) within

the *P. extremaustralis* genome.

Comparative genome analysis

General genome sequence analysis was performed using the bioinformatics tools

available on National Center for Biotechnology Information (www.ncbi.nlm.nih.gov),

including BLAST (Basic Local Alignment Search Tool) [23]. We also used RAST (Rapid

Annotation using Subsystem Technology) server [24] and Pseudomonas Genome

Database (Pseudomonas.com, [25]). Genomes used, with corresponding GenBank

accession numbers, were: *P. extremaustralis:* 14-3b (AHIP00000000.1), USBA 515

(FUYI01), DSM 17835^T (LT629689.1), 2E-UNGS (CP091043.1),

CSW01(JAQKGS01),1906 (JARIXU01), NQ5 (JARBJR01) strains, *P. protegens* Pf-5

(CP000076.1), and *P. syringae* pv *syringae* B728a (CP000075.1)

Quantitative Real Time PCR Experiments (RT qPCR)

Total RNA of *P. extremaustralis* microaerobic and m-NS cultures was isolated using

the Total RNA Extraction Kit (RBC Biosciences). After treatment with DNaseI, cDNA

was obtained using random hexamers (Promega) and Revert Aid reverse transcriptase

(ThermoFisher Scientific) according to the manufacturer's instructions. RT qPCR was

performed using a MyiQ2 Real-Time PCR Detection System (Bio-Rad Laboratories,

Hercules, USA) and Real-Time PCR mix (EvaGreen qPCR Mix Plus no ROX). The

expression of target genes was evaluated using the following primers: *motB* forward 5′

CAACGCCAGCAACAAAGACA 3′ and reverse 5′ CTTCGGAAAACCGCGCATC 3′; *hbo*

forward   5′   CGAGCCTAAGAGCAACACCA   3′   and   reverse

5′TGAATAGGCTGTCGGCACTG   3′   and   *flaB*   forward   5'

CGTAACCAGCGCTGACATGGCTC   3'   and   reverse   5'

GACAGGATCGCAGTGGAAGCCGA 3'. Expression of the 16S rRNA gene was used

for normalization of target genes expression levels in each condition, obtained using

primers   forward   5′   AGCTTGCTCCTTGATTCAGC   3′   and   reverse   5′

AAGGGCCATGATGACTTGAC' 3. The cycling conditions were as follows:

denaturation at 95 °C for 15 min, 40 cycles at 95 °C for 25 s, 60 °C for 15 s, and 72 °C

for 15 s, with fluorescence acquisition at 80 °C in single mode. The $2^{-\Delta\Delta CT}$ method [26]

was used to calculate the relative fold gene expression of individual genes.

Bacterial growth using *myo*-inositol as sole carbon source

Myo-inositol metabolism was evaluated in E2 medium [27] supplemented with 10 g/l of

myo-inositol or glucose under aerobic and microaerobic conditions. Microaerobic

cultures were performed as described above while for aerobic experiments, cultures

were incubated in Erlenmeyer flasks using 1:10 medium-to-flask volume ratio and 200

rpm agitation. Aerobic and microaerobic cultures were incubated for 24h and 48h,

respectively.

Biofilm formation was assayed in E2 medium supplemented with *myo*-inositol or

glucose and KNO_3 in polystyrene microtiter plates with an initial OD_{600nm} of 0.025. After

48h of static incubation at 30°C, OD_{600nm} of planktonic cells (Planktonic cells

absorbance, PCA) was measured and biofilms were quantified using the standard

crystal violet method [28]. Briefly, attached cells were stained with 200 µl of 0.1%

crystal violet, further washed and the colorant was solubilized with absolute ethanol.

Crystal violet solution was transferred to flat bottom microtiter plates and OD_{570nm}

(Crystal violet absorbance CVA) was measured with a BMG OPTIMA FLUOstar

microplate reader. Biofilm formation index was defined as CVA/PCA.

Pyoverdine production

Pyoverdine production of microaerobic cultures was analyzed in iron-limited E2

medium, without microelements, supplemented with *myo*-inositol or glucose as carbon

source and KNO_3. After 48h pyoverdines in the culture supernatants were determined

by measuring integrated fluorescence emission between 445-460 nm after excitation at

420 nm [29,30]. The values were expressed relative to cell dry weight per ml.

Survival and stress experiments

P. extremaustralis was cultured for 48h under microaerobic conditions in E2 cultures

supplemented with *myo*-inositol or glucose and KNO_3 (0.8 g/l). Cultures were exposed

to 100 µM GSNO (m-NS) for 1h and cell viability was determined in plate assay.

Growth inhibition in response to oxidative or combined nitro-oxidative stress was

evaluated by filter disk assay by growing *P. extremaustralis* in E2 medium

supplemented with glucose or *myo*-inositol and increasing KNO_3 concentrations of 0.8,

1.5 and 2.5 g/l that led to nitrite accumulation. Control cultures without nitrate were

performed (oxidative stress only). LB plates were seeded with the different cultures and

Whatman n° 1 filter discs (6 mm) were impregnated with 5 µl of 30% (v/v) H_2O_2 (Merck)

as previously described [31]. Plates were incubated overnight, and the diameter of the

halo was determined using the software ImageJ [32]. Nitrite concentration in the

supernatant of cultures was determined following the method described by [33] and

values were normalized to cell dry weight.

Data availability

RNA-seq data were deposited in the European Molecular Biology Laboratory under

accession number E-MTAB-11689.

Statistical analysis

Differences between means were determined through the Student's t test with

confidence levels at > 95% in which $P < 0.05$ was considered as statistically significant.

A Fisher's exact test was performed to analyze the nitrosylation targets proteins. 1 or 2-

ways ANOVA with multiple comparison were used when correspond.

Results

Effects of GSNO exposure in *P. extremaustralis*

Nitrosative stress effect on *P. extremaustralis* survival and growth under microaerobic

conditions was evaluated using different S-nitrosoglutation (GSNO) concentrations.

Growth in presence of different concentrations of GSNO was similar to the control

culture except for 100 µM that showed a significant decrease ($P<0.05$) at the stationary

phase (Fig. S2). Survival assays showed that when the bacterial culture was exposed

to GSNO for 1h a significant decrease in bacterial viable counts (CFU/ml) was

observed comparing with the control culture and provoked a 75% drop in survival in

microaerobic cultures (Fig. 1). These results indicate that GSNO exposure affected *P.*

extremaustralis survival in short time, thus we chose 100 µM GSNO (m-NS) exposure

for 1h for further experiments.

Figure 1.   Survival of microaerobic cultures to GSNO exposure. Cultures were

exposed during 1h and survival in LB plates was determine. Circles represent

individual values of independent experiments. Error bars represent the standard

deviation of the mean. Different letters denote significant differences using 1-way

ANOVA with Tukey's multiple comparisons test (P < 0.05)

Transcriptomic profile of *P. extremaustralis* under low aeration and nitrosative

stress conditions.

P. extremaustralis response to nitrosative stress was evaluated by comparing

transcriptomic data obtained by RNA sequencing (RNA-Seq) in microaerobic cultures

m-C and m-NS. RNA-seq expression profile revealed 5888 transcripts. Rockhopper

analysis showed 249 transcripts with differential expression in m-NS compared to m-C

(P <0.05, Q <0.05). Among them, 86 genes were found to be repressed and 163 over-

expressed, representing 1.46% and 2.77% of the total, respectively (Fig. S3a, Table

S1). Several genes with differential expression did not have defined associated

functions and were classified as "hypothetical", representing 11.6% and 21% of the

over-expressed and repressed genes, respectively. Among genes with a known

predicted function, *P. extremaustralis* under m-NS showed an increased expression of

genes involved in carbon, RNA, DNA and iron metabolism (Table S1).

RNA-seq results were validated by analyzing the expression of three selected genes,

motB, *flaB*, and *hbo* using qRT PCR (Fig. S2b). In concordance with the RNA-seq

results, we found higher expression levels of *motB,* encoding a flagellar motor rotation

protein, in m-NS vs. m-C, whereas we observed a decrease in *hbO,* encoding the

hemoglobin-like protein HbO (Fig. S2b, Table S1) and no differences in *flaB* expression

between conditions.

P. extremaustralis' response to nitrosative stress

To elucidate possible functional relationships, differentially expressed genes were

categorized into functional groups (Fig. 2a). GSNO exposure caused increased

expression of genes encoding different RNS detoxifying enzymes such as an iron-

dependent superoxide dismutase (PE143B_0125285) and an alkyl hydroperoxide

reductase (*ahpC*), a thioredoxin (PE143B_0111145) and a glutathione S-transferase

(PE143B_0109905). A gene encoding another glutathione S-transferase

(PE143B_0100180) was repressed in m-NS along with *hbO* (Fig. 2a, Table S1).

Transcriptomic analysis showed an increase in mRNA expression of iron and heme

transporters coding genes such as *ctaB, ybaN, irpA* and *afuA* (Fig 2b, Table S1). In m-

NS we found upregulation in genes related to pyoverdine siderophore biosynthesis, a

key function for iron uptake, including *pdvA, pdvM, pdvF, fpvA, fpvE* and *fpvK* (Fig. 2b,

Table S1). Moreover, the arginase coding gene, was also upregulated in m-NS

comparing with m-C. Arginase catalyzes the conversion of arginine to ornithine, which

can then serve as a substrate for pyoverdine biosynthesis (Fig. 2b, Table S1).

Iron and other metals are key components of iron-sulfur proteins that could be target of

RNS, leading to protein inactivation and triggering cellular signaling [34]. Therefore, we

analyzed other cellular functions related with transcription and protein turn-over. GSNO

treated cultures showed an increase in expression of RNA helicase

(PE143B_0107010), RNA polymerase associated protein RapA, transcriptional

terminator *nusB* (PE143B_0123570) and 11 ribosomal proteins coding genes (Table

S1). In concordance, a gene encoding translation initiation inhibitor was repressed

(Table S1). To deeply analyze the protein turnover related with RNS attack we

performed an analysis using Target Pathogen platform to detect genes encoding

proteins with free tyrosine or cysteine as well as metal binding sites which are reactive

to RNS [1]. Among differentially expressed genes in m-NS we found that nitrosylable

proteins were over-represented in comparison to non-nitrosylable when total genes of

P. extremaustralis were considered (Fisher's exact test, $P = 0.0323$). Overall, our

results suggest that nitrosative stress under microaerobic conditions provokes an

increase in the expression of genes involved in transcription and translation processes,

probably to compensate the damage to proteins with Fe-S clusters or metal binding

(Fig. 2c).

Figure 2. Effect of GSNO on gene expression profile. **a.** Classification of

differentially expressed genes after GSNO exposure in microaerobiosis (m-NS vs. m-

C) into functional categories. Hypothetical transcripts (with no inferred function) are not

displayed. Green and red bars represent up- and down-regulated genes, respectively.

b. Over-expressed genes related to iron metabolism. **c**. Nitrosylable proteins

determined with Target Pathogens in the differential expressed genes compared to

total nytrosilable proteins in *P. extremaustralis* genome. DE: differentially expressed

genes. Fisher's exact test, P = 0.0323.

Inositol metabolism in *P. extremaustralis*

Remarkably, RNAseq results showed increased expression of genes related to inositol

catabolism which involves enzymes that convert inositol into acetyl-CoA and

dihydroxyacetone phosphate (DHAP) (Fig. 3a, Table S1).

Our results showed a significant up-regulation (P and Q<0.05) in mRNAs encoded in

the *iol* operon in response to GSNO stress, including *iolB* (2-fold), *iolD* (1.75-fold), *iolE*

(1.83-fold) *iolG2* (also known as *iolW* or *mocA*) (1.86-fold), *iolR* (2.62-fold), and the *iatA*

gene (1.75-fold) (Fig. 3a; Table S1). No significant differences were observed in the

expression of *iolA* (1-fold)*, iolG1*(1.54-fold) and *iolC/J* (1.25-fold) in both conditions,

neither for *iatP* (1.20-fold) and *ibpA* (1.44-fold) (Fig. 3a). It was found that *iolC*

possesses the DUF2090 domain that was recently proposed to encode 2-deoxy-5-

keto-gluconic acid-6-phosphate aldolase [35]. This domain fulfills function originally

attributed to *iolJ* which is not present in *P. extremaustralis* genome (Fig. 3b)*.

When we compared the inositol catabolic cluster of *P. extremaustralis* with *P.*

protegens Pf-5, a well-known plant growth promoter species, and *P. syringae* pv.

syringae B728a, a plant pathogen, we found some differences in this genomic region.

Unlike *P. protegens* Pf-5, the genome of *P. extremaustralis* did not exhibit the presence

of *iolL*. Conversely, we identified a duplication of *iolG* within this bacterial species. *P.*

syringae pv. syringae B728a presented *iolH*, which is absent in both *P. protegens* and

P. extremaustralis. Similar to *P. extremaustralis, P. syringae* pv. syringae B728a

presented two copies of *iolG* but separated by *iolH* (Fig. 3b). The *iolA* gene in *P.*

extremaustralis is located in a different genomic region like in *P. protegens* and other

analyzed *Pseudomonas* [36]. However, in P. *syringae* pv. syringae B728a is included in

the *iol* catabolic cluster (Fig. 3b). Furthermore, we found that the *iol* gene cluster of *P.*

extremaustralis exhibited a high degree of intraspecific conservation showing the same

organization as 14-3b in all genomes analyzed, including strains USBA 515,

DSM17835^T, 2E-UNGS, CSW01,1906, NQ5.

Figure 3. Metabolic pathway associated with inositol catabolism in *Pseudomonas*

***extremaustralis* 14-3b: Expression under nitrosative stress and genetic**

organization. **a**. Gene expression and metabolic route related to inositol catabolism

under m-NS Genes and arrows shown in green indicate up-regulated functions under

m-NS, whereas those in blue represent functions with no expression differences

between both conditions. **b**. Analysis of genes involved in inositol in *P. extremaustralis*

in comparison with *P. protegens* Pf-5 and *P. syringae* pv. syringae B728a. (*) The

same genetic organization was observed in USBA 515, DSM17835^T, 2E-UNGS,

CSW01,1906 and NQ5 *P. extremaustralis* strains. Compounds: KMI, 2-keto-*myo*-

inositol; THcHDO, 3, 3D-(3,4/5) trihydroxycyclohexane-1,2-dione; 5DG, 5-deoxy

glucuronic acid; DKG, 2-deoxy-5-keto-d-gluconic acid; DKGP, DKG 6-phosphate;

DHAP, dihydroxyacetone phosphate; MSA, malonic semialdehyde; acetyl-CoA, acetyl

coenzyme A; DHAP, dihydroxyacetone phosphate; G3P, glyceraldeyde-3-phosphate;

TCA, tricarboxylic acid cycle. Genes: *iolG1*, *myo*-inositol dehydrogenase; *iolG2*, *scyllo*-

inositol dehydrogenase; *iolE*, 2KMI dehydratase; *iolD*, THcHDO hydrolase; *iolB*, 5DG

isomerase; *iolC/J*, DKG kinase + aldolase; *iolA*, MSA dehydrogenase; *iatA*, sugar ABC

transporter ATP-binding protein; *iatP*, ABC transporter permease; *ibpA*, sugar ABC

transporter substrate-binding protein.

As we detected the presence and expression of *iol* genes, we first tested *P.*

extremaustralis capability to grow under aerobic conditions using *myo*-inositol as sole

carbon source. Growth was similar between *myo*-inositol and glucose supplemented

cultures reaching an OD$_{600nm}$ value of 2.02 ± 0.15 and 2.22 ± 0.15, respectively. Under

microaerobic conditions *P. extremaustralis* developed an evident biofilm in *myo*-inositol

supplemented cultures (Fig. 4a). Therefore, OD$_{600nm}$ and cellular dry weight (DW) were

determined (Fig. 4b). Similar results were obtained for both carbon sources regardless

of the approach employed for growth estimation. Biofilm formation was assayed in E2

medium supplemented with glucose or inositol. In concordance, we also found that with

myo-inositol supplemented cultures *P. extremaustralis* showed a higher biofilm index

compared to glucose (Fig. 4c).

Additionally, considering the upregulation of coding genes related with pyoverdine

biosynthesis after treatment with GSNO, we investigated siderophores production in *P.*

extremaustralis using *myo*-inositol as carbon source. Pyoverdine production was higher

for *myo*-inositol supplemented cultures comparing to glucose (Fig. 4d)

Figure 4. Physiological features of *P. extremaustralis* using *myo*-inositol o

glucose as carbon source. a. Photograph of microaerobic *myo*-inositol or glucose

supplemented cultures showing the biofilm in the surface. **b.** Microaerobic growth

measured by optical density (OD_{600nm}) or Cellular dry weight (DW). **c.** Biofilm formation

index (ACV/APC) in *myo*-inositol or glucose. **d.** Effect of carbon sources on pyoverdine

production. RFU: Relative Fluorescence Units. Fluorescence units were normalized to

the cell dry weight/ml. Error bars represent the standard deviation of the mean.

***Myo*-inositol impact on stress response**

Further, we investigated stress resistance when *myo*-inositol was used as sole carbon

source under low oxygen conditions. To analyze the nitrosative stress response, we

performed a survival test using microaerobic cultures in E2 medium supplemented with

myo-inositol or glucose exposed to GSNO treatment. We found no differences in

survival between those treated with GSNO and control cultures despite of the carbon

source (Fig. 5a).

P. extremaustralis is only capable to reduce nitrate to nitrite which accumulates in the

extracellular medium. Therefore, the resistance to nitro-oxidative stress derivate from

the combination of the nitrite accumulated and H_2O_2 was analyzed in *myo*-inositol and

glucose supplemented cultures, following the scheme shown in Fig. S4. Nitrite

production was detected in cultures supplemented with *myo*-inositol and glucose as

sole carbon source and different nitrate concentrations reaching a maximum of 13.553±

1.624, confirming also the *myo*-inositol utilization in a respiratory pathway (Fig. 5b).

When *P. extremaustralis* was grown in *myo*-inositol or glucose supplemented media

without nitrate, the oxidative stress resistance was similar for both carbon sources

while the nitro-oxidative stress resistance was higher for cultures grown in *myo*-inositol

in all tested KNO_3 concentrations (Fig. 5c).

Figure 5. Nitrosative and combined nitro-oxidative stress resistance in *P. extremaustralis*. a. Survival after GSNO exposure. **b.** Nitrite production in cultures supplemented with *myo*-inositol or glucose and different nitrate concentrations. **c.** Inhibition halos of cultures performed in (**b**) exposed to H_2O_2 (Combined nitro-oxidative stress). Error bars represent the standard deviation of the mean.

Discussion

Bacterial species are subjected to different stress conditions including lack of nutrients, changes in temperature or oxygen availability and oxidative and nitrosative stress, among others. Oxidative and nitrosative stress can be produced by the imbalance between the ROS or RNS generation and the incapability to detoxify them [37] .

In response to these stress conditions, bacteria may activate various response mechanisms, such as the production of chaperons and detoxicant enzymes, the adjustment of gene expression, and the modification of their cell envelope composition.

A common effect during stress is the reprograming of transcription and translation

processes including the reduction of ribosome biogenesis, the modification of the

translational machinery and the regulation of initiation and elongation of certain

proteins [38]. Interestingly, gene expression during stress involves not only the

transcription processes but also the modification of mRNA stability and in overall with

translation provokes a transcriptome stability [39,40]. In this work, we found an up-

regulation of the transcription and translation machinery probably to compensate the

damage caused by RNS particularly on nitrosylable proteins. The Fe-S clusters, that

can be found in three different forms, exhibit high capacity of accepting or donating

electrons thus are important for redox response and act as redox sensors like the Fnr

regulator in *Escherichia coli* or Anr in *Pseudomonas* species [41–43].These proteins

and those containing metals are involved in several essential cellular functions like

respiration, central carbon catabolism and RNA and DNA processes and are a target of

RNS damage [41] Particularly, it has been reported that the reaction between nitric

oxide (NO) with Fe-S clusters lead to protein degradation and breakdown of the cluster

generating dinitrosyl iron complexes [44]. Our analysis using Target Pathogen platform

showed the enrichment of genes encoding nitrosylable proteins in the differentially

expressed ones, particularly in those upregulated. Our hypothesis is that to cope with

GSNO derived stress the Fe-S and metal binding proteins need to be replaced. This

also could explain the upregulation of Fe-uptake related genes observed in presence of

GSNO. Iron is involved in the Fenton reaction in presence of H_2O_2, derived from

endogenous or exogenous sources, that could exacerbate the oxidative damage but

paradoxically this metal is also necessary for Fe-S and metal protein biogenesis [45,46]

. This complex scenario additionally requires the involvement of enzymes and

antioxidant cycles. In presence of GSNO we found upregulated the coding genes of

glutathione S-transferase, superoxide dismutase (SOD), thiol-disulfide isomerase,

thioredoxins and alkyl hydroperoxide reductase subunit C-like protein. The alkyl

hydroperoxide reductase, important during non-lethal stress, SOD as well as the

oxidation of thioredoxins in response to oxidative stress depends on NAD(P)H

availability and in overall reduction power is needed for the essential process for

survival and to cope with stress [47] .Interestingly, the needed of NADH and/or NADPH

production during stress could represent a challenge for bacterial cells. Glycolysis and

TCA provides the NADH and energy necessaries for bacterial survival but also creates

an oxidative scenario through the respiration chain [48]. It has been reported several

metabolic strategies to obtain energy and reducing power besides glycolysis such as

aminoacid catabolism, ethanol oxidation and in *P. fluorescens* an entire metabolic

reprogramming has been reported in presence of oxidative stress to generate NADPH

[49]. In this previous work the authors showed an increase in the activity and

expression of malic enzyme (similar to this work in presence of GSNO), glucose-6-

phosphate dehydrogenase and NADP+-isocitrate dehydrogenase [49].

In addition, we found that in presence of GSNO under microaerobic conditions *P.*

extremaustralis over-expressed several *iol* catabolic genes including the transcriptional

regulator and the *iatA* gene. We also demonstrated that this bacterium was effectively

capable of using inositol as its sole carbon source. Inositol is a sugar alcohol present in

all domains of life and presents various structural isomers [50], with *myo*-inositol being

the most abundant in nature. *Myo*-inositol is found in high quantities in plants, including

root exudates, and may also originate from the dephosphorylation of phytate, one of

the major phosphorus storage molecules for plants [51]. Inositol catabolism has been

experimentally demonstrated in several bacterial species, both Gram positive and

Gram negative, belonging to genera such as Rhizobium*, Lactobacillus, Klebsiella,*

Corynebacterium, Legionella, Bacillus, and Salmonella [52–58].   Furthermore, an

extensive analysis of inositol catabolic genes in thousands of bacterial genomes

revealed a widespread distribution within different ecological niches [51].

While inositol metabolism in bacteria has garnered recent attention, its role in

stress resistance remains unexplored. Remarkably, our findings revealed that inositol

protects against nitro-oxidative stress, and increases biofilm and pyoverdine

production, which are relevant traits for environmental adaptability. Inositol catabolism leads to the production of glyceraldehyde-3 phosphate, and acetyl-CoA both intermediaries of central energy generation metabolism. In addition, *iolG2,* which was found overexpressed after GSNO exposure, is related to NAPH generation, necessary for the function of antioxidative defenses.

It is worth noting that previous transcriptome analysis of *P. extremaustralis* performed under cold conditions or under microaerobic conditions with or without oxidative stress did not show the upregulation of inositol catabolic genes [14,59,60], indicating a fine-tuning response to nitrosative stress.

Some RNS also play a crucial role in cell signaling. For instance, in plants, NO mediates for various processes including root hair growth, stomatal closure, programmed cell death and other responses that help them adapt to changing environmental conditions [8] whereas in bacteria, NO plays a role as intermediary in denitrification processes, but also in biofilm formation and quorum sensing [61]. The *iol* gene cluster was identified in both plant-pathogenic bacteria, and rhizospheric plant growth promoting bacteria forming symbiotic root nodules [51].

Recently, the *iol* gene cluster in *Pseudomonas* was identified as an important trait in root colonizers, by increasing swimming motility and siderophore production in response to plant derived inositol [36]. Increased pyoverdine siderophore production in presence of *myo*-inositol was also observed in this work. *P. extremaustralis* possesses plant growth-promotion traits, including the ability to solubilize phosphate and produce indole acetic acid [62]. Since NO is also a signaling molecule released by plants [63], we hypothesize that NO released by GSNO mimic the NO production of plants, triggering the differential expression of the inositol metabolic pathway conferring an adaptive advantage.

In conclusion, our findings reveal a multifaceted and finely tuned response mechanism to nitrosative stress that encompasses transcriptional reprogramming, the upregulation of genes encoding key proteins involved in iron homeostasis, and the activation of

inositol catabolism. Notably, experimental results point to inositol metabolism as a

novel mechanism to cope with stress.

Acknowledgments

This work was funded by grants from Universidad de Buenos Aires and Consejo

Nacional de Investigaciones Científicas y Técnicas (CONICET). We are thankful to Dr.

Jörg Vogel (Institut für Molekulare Infektionsbiologie (IMIB) – University of Würzburg)

for his guidance and for providing resources for conducting the RNAseq experiments.

E.C.S.V. received a Deutscher Akademischer Austauschdienst (DAAD) fellowship for

short-term research stays at the RNA biology group (IMIB – University of Würzburg,

Germany).

References

1.   Crack JC, Thomson AJ, Brun NE Le. Iron – Sulfur Clusters as Biological
Sensors⬜: The Chemistry of Reactions with Molecular Oxygen and Nitric Oxide.
2014.

2.   Czaja AJ. Immunopathogenesis of Autoimmune Liver Damage. 2017. pp. 19–48.
doi:10.1016/B978-0-444-63707-9.00002-7

3.   Tharmalingam S, Alhasawi A, Appanna VP, Lemire J, Appanna VD. Reactive
nitrogen species (RNS)-resistant microbes: adaptation and medical implications.
Biol Chem. 2017;398: 1193–1208. doi:10.1515/hsz-2017-0152

4.   Aranda-Rivera AK, Cruz-Gregorio A, Arancibia-Hernández YL, Hernández-Cruz
EY, Pedraza-Chaverri J. RONS and Oxidative Stress: An Overview of Basic
Concepts. Oxygen. 2022;2: 437–478. doi:10.3390/oxygen2040030

5.   Lelieveld S, Wilson J, Dovrou E, Mishra A, Lakey PSJ, Shiraiwa M, et al.
Hydroxyl Radical Production by Air Pollutants in Epithelial Lining Fluid Governed
by Interconversion and Scavenging of Reactive Oxygen Species. Cite This:
Environ Sci Technol. 2021;55: 14069–14079. doi:10.1021/acs.est.1c03875

6.   Li H, Zhou X, Huang Y, Liao B, Cheng L, Ren B. Reactive Oxygen Species in
Pathogen Clearance: The Killing Mechanisms, the Adaption Response, and the
Side Effects. Front Microbiol. 2020;11: 622534. doi:10.3389/fmicb.2020.622534

7.   Del Río LA. ROS and RNS in plant physiology: an overview. 2015 [cited 23 Aug
2023]. doi:10.1093/jxb/erv099

8.   Zaffagnini M, Bedhomme M, Marchand CH, Morisse S, Trost P, Lemaire SD.
Redox Regulation in Photosynthetic Organisms: Focus on Glutathionylation.

https://home.liebertpub.com/ars. 2012;16: 567–586.
doi:10.1089/ARS.2011.4255

9.   Robinson JL, Adolfsen KJ, Brynildsen MP. Deciphering nitric oxide stress in
bacteria with quantitative modeling. Curr Opin Microbiol. 2014;19: 16–24.
doi:10.1016/j.mib.2014.05.018

10. Arai H. Regulation and Function of Versatile Aerobic and Anaerobic Respiratory
Metabolism in *Pseudomonas aeruginosa*. Front Microbiol. 2011;2: 103.
doi:10.3389/fmicb.2011.00103

11. Tribelli PM, Lujan AM, Pardo A, Ibarra JG, Fernández Do Porto D, Smania A, et
al. Core regulon of the global anaerobic regulator Anr targets central metabolism
functions in *Pseudomonas* species. Sci Rep. 2019;9: 9065. doi:10.1038/s41598-
019-45541-0

12. Raiger Iustman LJ, Tribelli PM, Ibarra JG, Catone M V, Solar Venero EC, López
NI. Genome sequence analysis of *Pseudomonas extremaustralis* provides new
insights into environmental adaptability and extreme conditions resistance.
Extremophiles. 2015;19: 207–220. doi:10.1007/s00792-014-0700-7

13. Fonseca P, Moreno R, Rojo F. Growth of Pseudomonas putida at low
temperature: Global transcriptomic and proteomic analyses. Environ Microbiol
Rep. 2011. doi:10.1111/j.1758-2229.2010.00229.x

14. Tribelli PM, Solar Venero EC, Ricardi MM, Gómez-Lozano M, Raiger Iustman
LJ, Molin S, et al. Novel Essential Role of Ethanol Oxidation Genes at Low
Temperature Revealed by Transcriptome Analysis in the Antarctic Bacterium
Pseudomonas extremaustralis. PLoS One. 2015;10: e0145353.
doi:10.1371/journal.pone.0145353

15. da Cruz Nizer WS, Inkovskiy V, Versey Z, Strempel N, Cassol E, Overhage J.
Oxidative Stress Response in *Pseudomonas aeruginosa*. Pathogens. 2021;10:
1187. doi:10.3390/pathogens10091187

16. López NI, Pettinari MJ, Stackebrandt E, Tribelli PM, Põtter M, Steinbüchel A, et
al. *Pseudomonas extremaustralis* sp. nov., a Poly(3-hydroxybutyrate) producer
isolated from an antarctic environment. Curr Microbiol. 2009;59: 514–519.
doi:10.1007/s00284-009-9469-9

17. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics. 2014;30: 2114–2120.
doi:10.1093/bioinformatics/btu170

18. Tjaden B. A computational system for identifying operons based on RNA-seq
data. Methods. 2020;176: 62–70. doi:10.1016/j.ymeth.2019.03.026

19. Ogata H, Goto S, Sato K, Fujibuchi W, Bono H, Kanehisa M. KEGG: Kyoto
Encyclopedia of Genes and Genomes. Nucleic Acids Res. 1999;27: 29–34.
doi:10.1093/nar/27.1.29

20. Caspi R, Billington R, Ferrer L, Foerster H, Fulcher CA, Keseler IM, et al. The
MetaCyc database of metabolic pathways and enzymes and the BioCyc
collection of pathway/genome databases. Nucleic Acids Res. 2016;44: D471-
480. doi:10.1093/nar/gkv1164

21.   Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, et al.
589   STRING v11: protein–protein association networks with increased coverage,
590   supporting functional discovery in genome-wide experimental datasets. Nucleic
591   Acids Res. 2019;47: D607–D613. doi:10.1093/nar/gky1131

592   22.   Sosa EJ, Burguener G, Lanzarotti E, Defelipe L, Radusky L, Pardo AM, et al.
593   Target-Pathogen: a structural bioinformatic approach to prioritize drug targets in
594   pathogens. Nucleic Acids Res. 2018;46: D413–D418. doi:10.1093/nar/gkx1015

595   23.   Altschul S. Gapped BLAST and PSI-BLAST: a new generation of protein
596   database search programs. Nucleic Acids Res. 1997;25: 3389–3402.
597   doi:10.1093/nar/25.17.3389

598   24.   Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, et al. The RAST
599   Server: Rapid Annotations using Subsystems Technology. BMC Genomics.
600   2008;9: 75. doi:10.1186/1471-2164-9-75

601   25.   Winsor GL, Griffiths EJ, Lo R, Dhillon BK, Shay JA, Brinkman FSL. Enhanced
602   annotations and features for comparing thousands of *Pseudomonas* genomes in
603   the Pseudomonas genome database. Nucleic Acids Res. 2016;44: D646-53.
604   doi:10.1093/nar/gkv1227

605   26.   Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using
606   Real-Time Quantitative PCR and the $2-\Delta\Delta CT$ Method. Methods. 2001;25: 402–
607   408. doi:10.1006/meth.2001.1262

608   27.   Lageveen RG, Huisman GW, Preusting H, Ketelaar P, Eggink G, Witholt B.
609   Formation of Polyesters by *Pseudomonas oleovorans*: Effect of Substrates on
610   Formation and Composition of Poly-(R)-3-Hydroxyalkanoates and Poly-(R)-3-
611   Hydroxyalkenoates . Appl Environ Microbiol. 1988;54: 2924–2932.
612   doi:10.1128/AEM.54.12.2924-2932.1988

613   28.   O'Toole GA, Kolter R. Flagellar and twitching motility are necessary for
614   *Pseudomonas aeruginosa* biofilm development. Mol Microbiol. 1998;30: 295–
615   304. doi:10.1046/j.1365-2958.1998.01062.x

616   29.   Ruiz JA, Bernar EM, Jung K. Production of Siderophores Increases Resistance
617   to Fusaric Acid in *Pseudomonas protegens* Pf-5. Devireddy lax, editor. PLoS
618   One. 2015;10: e0117040. doi:10.1371/journal.pone.0117040

619   30.   Olmo A Del, Caramelo C, Sanjose C. Fluorescent complex of pyoverdin with
620   aluminum. J Inorg Biochem. 2003;97: 384–387. doi:10.1016/S0162-
621   0134(03)00316-7

622   31.   Ayub ND, Pettinari MJ, Ruiz JA, López NI. A polyhydroxybutyrate-producing
623   *Pseudomonas* sp. isolated from antarctic environments with high stress
624   resistance. Curr Microbiol. 2004;49: 170–174. doi:10.1007/s00284-004-4254-2

625   32.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of
626   image analysis. Nat Methods. 2012;9: 671–675. Available:
627   http://www.ncbi.nlm.nih.gov/pubmed/22930834

628   33.   Gerhard P, Murray R, Costilow N, Nester W, Wood N, Krieg N, et al. Manual of
629   Method of General Bacteriology. Washington: American Society for
630   microbiology,; 1981.

34. Seth D, Hausladen A, Stamler JS. Anaerobic Transcription by OxyR: A Novel
Paradigm for Nitrosative Stress. Antioxid Redox Signal. 2020;32: 803–816.
doi:10.1089/ars.2019.7921

35. Price MN, Deutschbauer AM, Arkin AP. Filling gaps in bacterial catabolic
pathways with computation and high-throughput genetics. PLoS Genet. 2022;18:
e1010156. doi:10.1371/journal.pgen.1010156

36. Sánchez-Gil JJ, Poppeliers SWM, Vacheron J, Zhang H, Odijk B, Keel C, et al.
The conserved *iol* gene cluster in *Pseudomonas* is involved in rhizosphere
competence. Current Biology. 2023;33: 3097-3110.e6.
doi:10.1016/j.cub.2023.05.057

37. Chautrand T, Souak D, Chevalier S, Duclairoir-Poc C. Gram-Negative Bacterial
Envelope Homeostasis under Oxidative and Nitrosative Stress. Microorganisms.
2022;10. doi:10.3390/microorganisms10050924

38. Advani VM, Ivanov P. Translational Control under Stress: Reshaping the
Translatome. BioEssays. 2019;41. doi:10.1002/bies.201900009

39. Kristoffersen SM, Haase C, Weil MR, Passalacqua KD, Niazi F, Hutchison SK,
et al. Global mRNA decay analysis at single nucleotide resolution reveals
segmental and positional degradation patterns in a Gram-positive bacterium.
Genome Biol. 2012;13: R30. doi:10.1186/gb-2012-13-4-r30

40. Vargas-Blanco DA, Shell SS. Regulation of mRNA Stability During Bacterial
Stress Responses. Front Microbiol. 2020;11: 2111.
doi:10.3389/fmicb.2020.02111

41. Vernis L, El Banna N, Baïlle D, Hatem E, Heneman A, Huang M-E. Fe-S
Clusters Emerging as Targets of Therapeutic Drugs. Oxid Med Cell Longev.
2017;2017: 3647657. doi:10.1155/2017/3647657

42. Golinelli-Cohen M-P, Bouton C. Fe-S Proteins Acting as Redox Switch: New Key
Actors of Cellular Adaptive Responses. Curr Chem Biol. 2017;11.
doi:10.2174/2212796811666170406163809

43. Galimand M, Gamper M, Zimmermann A, Haas D. Positive FNR-like control of
anaerobic arginine degradation and nitrate respiration in *Pseudomonas*
aeruginosa. J Bacteriol. 1991;173: 1598–606. Available:
http://www.ncbi.nlm.nih.gov/pubmed/1900277

44. Harrop TC, Tonzetich ZJ, Reisner E, Lippard SJ. Reactions of synthetic [2Fe-2S]
and [4Fe-4S] clusters with nitric oxide and nitrosothiols. J Am Chem Soc.
2008;130: 15602–10. doi:10.1021/ja8054996

45. Reniere ML. Reduce, Induce, Thrive: Bacterial Redox Sensing during
Pathogenesis. J Bacteriol. 2018;200. doi:10.1128/JB.00128-18

46. Winterbourn CC. Toxicity of iron and hydrogen peroxide: the Fenton reaction.
Toxicol Lett. 1995;82–83: 969–74. doi:10.1016/0378-4274(95)03532-x

47. Lemire J, Alhasawi A, Appanna VP, Tharmalingam S, Appanna VD. Metabolic
defence against oxidative stress: the road less travelled so far. J Appl Microbiol.
2017;123: 798–809. doi:10.1111/jam.13509

48.   Zhang J, Li X, Olmedo M, Holdorf AD, Shang Y, Artal-Sanz M, et al. A Delicate
674   Balance between Bacterial Iron and Reactive Oxygen Species Supports Optimal
675   *C. elegans* Development. Cell Host Microbe. 2019;26: 400-411.e3.
676   doi:10.1016/j.chom.2019.07.010

677   49.   Singh R, Mailloux RJ, Puiseux-Dao S, Appanna VD. Oxidative stress evokes a
678   metabolic adaptation that favors increased NADPH synthesis and decreased
679   NADH production in *Pseudomonas fluorescens*. J Bacteriol. 2007;189: 6665–75.
680   doi:10.1128/JB.00555-07

681   50.   Michell RH. Inositol derivatives: evolution and functions. Nat Rev Mol Cell Biol.
682   2008;9: 151–161. doi:10.1038/nrm2334

683   51.   Weber M, Fuchs TM. Metabolism in the Niche: a Large-Scale Genome-Based
684   Survey Reveals Inositol Utilization To Be Widespread among Soil, Commensal,
685   and Pathogenic Bacteria. Microbiol Spectr. 2022;10: e0201322.
686   doi:10.1128/spectrum.02013-22

687   52.   Fry J, Wood M, Poole PS. Investigation of *myo* -Inositol Catabolism in
688   *Rhizobium leguminosarum* bv. *viciae* and Its Effect on Nodulation
689   Competitiveness. Molecular Plant-Microbe Interactions®. 2001;14: 1016–1025.
690   doi:10.1094/MPMI.2001.14.8.1016

691   53.   Yebra MJ, Zúñiga M, Beaufils S, Pérez-Martínez G, Deutscher J, Monedero V.
692   Identification of a Gene Cluster Enabling *Lactobacillus casei* BL23 To Utilize
693   *myo* -Inositol. Appl Environ Microbiol. 2007;73: 3850–3858.
694   doi:10.1128/AEM.00243-07

695   54.   Krings E, Krumbach K, Bathe B, Kelle R, Wendisch VF, Sahm H, et al.
696   Characterization of *myo* -Inositol Utilization by *Corynebacterium glutamicum* :
697   the Stimulon, Identification of Transporters, and Influence on l -Lysine
698   Formation. J Bacteriol. 2006;188: 8054–8061. doi:10.1128/JB.00935-06

699   55.   Morinaga T, Ashida H, Yoshida K-I. Identification of two scyllo-inositol
700   dehydrogenases in *Bacillus subtilis*. Microbiology (Reading). 2010;156: 1538–
701   1546. doi:10.1099/mic.0.037499-0

702   56.   Yuan C, Yang P, Wang J, Jiang L. Myo-inositol utilization by *Citrobacter koseri*
703   promotes brain infection. Biochem Biophys Res Commun. 2019;517: 427–432.
704   doi:10.1016/j.bbrc.2019.07.112

705   57.   Manske C, Schell U, Hilbi H. Metabolism of *myo* -Inositol by *Legionella*
706   *pneumophila* Promotes Infection of Amoebae and Macrophages. Appl Environ
707   Microbiol. 2016;82: 5000–5014. doi:10.1128/AEM.01018-16

708   58.   Kröger C, Rothhardt JE, Brokatzky D, Felsl A, Kary SC, Heermann R, et al. The
709   small RNA RssR regulates myo-inositol degradation by *Salmonella enterica*. Sci
710   Rep. 2018;8: 17739. doi:10.1038/s41598-018-35784-8

711   59.   Tribelli PM, Rossi L, Ricardi MM, Gomez-Lozano M, Molin S, Raiger Iustman LJ,
712   et al. Microaerophilic alkane degradation in *Pseudomonas extremaustralis*: a
713   transcriptomic and physiological approach. J Ind Microbiol Biotechnol. 2018;45:
714   15–23. doi:10.1007/s10295-017-1987-z

715   60.   Solar Venero EC, Ricardi MM, Gomez-Lozano M, Molin S, Tribelli PM, López NI.
716   Oxidative stress under low oxygen conditions triggers hyperflagellation and

motility in the Antarctic bacterium *Pseudomonas extremaustralis*. Extremophiles.
2019;23. doi:10.1007/s00792-019-01110-x

61.   Williams DE, Boon EM. Towards Understanding the Molecular Basis of Nitric
Oxide-Regulated Group Behaviors in Pathogenic Bacteria. J Innate Immun.
2019;11: 205–215. doi:10.1159/000494740

62.   Ibarra JG. Estudios microbiológicos relacionados con el mejoramiento de
cultivos vegetales en zonas desfavorables. . Doctoral Thesis, Facultad de
Ciencias Exactas y Natural. 2017.

63.   Sasaki Y, Oguchi H, Kobayashi T, Kusama S, Sugiura R, Moriya K, et al.
Nitrogen oxide cycle regulates nitric oxide levels and bacterial cell signaling. Sci
Rep. 2016;6: 22038. doi:10.1038/srep22038

728