Nitrosative stress under microaerobic conditions triggers inositol metabolism in

Pseudomonas extremaustralis

Esmeralda C. Solar Venero[1#]; Lucia Giambartolomei[2]; Ezequiel Sosa[3]; Darío Fernández

do Porto [2,3]; Nancy I. López[1,2†] & Paula M. Tribelli [1,2,†,*]

[1]IQUIBICEN-CONICET, Intendente Guiraldes 2160, 1428EGA, Buenos Aires,

Argentina.

[2]Departamento de Química Biológica, Facultad de Ciencias Exactas y Naturales,

Universidad de Buenos Aires, 1428EGA, Buenos Aires, Argentina.

[3]Instituto de Cálculo, Facultad de Ciencias Exactas y Naturales, UBA, Buenos Aires,

Argentina

[#]Current address: Department of Biochemistry, School of Medicine, Universidad

Autónoma de Madrid and Instituto de Investigaciones Biomédicas Alberto Sols

(Universidad Autónoma de Madrid-Consejo Superior de Investigaciones Científicas),

Madrid, Spain

[†]: contributed equally to this work

*: Correspondence to: paulatrib@qb.fcen.uba.ar

Abstract

Bacteria are exposed to reactive oxygen and nitrogen species that provoke oxidative and

nitrosative stress which can lead to macromolecule damage. Coping with stress

conditions involves the adjustment of cellular responses, which helps address metabolic

challenges. In this study, we performed a global transcriptomic analysis of the response

of *Pseudomonas extremaustralis* to nitrosative stress, induced by S-nitrosoglutathione

(GSNO), a nitric oxide donor, under microaerobic conditions. The analysis revealed the

upregulation of genes associated with inositol catabolism; a compound widely distributed

in nature whose metabolism in bacteria has aroused interest. The RNA-seq data also

showed heightened expression of genes involved in essential cellular processes like transcription, translation, amino acid transport and biosynthesis, as well as in stress resistance including iron-dependent superoxide dismutase, alkyl hydroperoxide reductase, thioredoxin, and glutathione S-transferase in response to GSNO. Furthermore, GSNO exposure differentially affected the expression of genes encoding nitrosylation target proteins, encompassing metalloproteins and proteins with free cysteine and /or tyrosine residues. Notably, genes associated with iron metabolism, such as pyoverdine synthesis and iron transporter genes, showed activation in the presence of GSNO, likely as response to enhanced protein turnover. Physiological assays demonstrated that *P. extremaustralis* can utilize inositol proficiently under both aerobic and microaerobic conditions, achieving growth comparable to glucose-supplemented cultures. Moreover, supplementing the culture medium with inositol enhances the stress tolerance of *P. extremaustralis* against combined oxidative-nitrosative stress. Concordant with the heightened expression of pyoverdine genes under nitrosative stress, elevated pyoverdine production was observed when *myo*-inositol was added to the culture medium. These findings highlight the influence of nitrosative stress on proteins susceptible to nitrosylation and iron metabolism. Furthermore, the activation of *myo*-inositol catabolism emerges as a protective mechanism against nitrosative stress, shedding light on this pathway in bacterial systems, and holding significance in the adaptation to unfavorable conditions.

Introduction

Bacterial survival and adaptability in changing environments depend on different resistance mechanisms against several stress agents. Free radicals are highly reactive species, mainly originated from energy producing processes like aerobic and anaerobic respiration, causing damage to different macromolecules. Reactive oxygen species (ROS) are produced during aerobic respiration, while anaerobic respiration, utilizing

nitrate as electron acceptor, generates reactive nitrogen species (RNS). ROS and RNS trigger oxidative or nitrosative stress, respectively. RNS include nitric oxide (NO) and its derivatives peroxynitrite ($ONOO^-$), nitrosothiols formed through reaction with thiol groups, and nitrotyrosine from nitration of tyrosine by NO, $ONOO^-$ or NO_2^- (Crack et al., 2014; Czaja, 2017; Tharmalingam et al., 2017). Nitrosative and oxidative stress can also be induced from various environmental compounds, like pollutants, xenobiotics, and antimicrobials, and from UV radiation (Aranda-Rivera et al., 2022). NO is also produced by the mammalian immune system and acts as a modulator molecule in both mammals and plants (Del Río, 2015; Lelieveld et al., 2021; Li et al., 2020). Nitrosative stress can lead to DNA and macromolecular damage, including protein oxidation. Some residues or structures are more susceptible to the RNS or ROS attack such as iron-sulfur (Fe-S) clusters, metal centers, tyrosine and cysteine residues, and thiols (Zaffagnini et al., 2012). The radical attack to these residues or clusters results in both reversible and irreversible damage, leading to alterations in function and structure(Robinson et al., 2014).

Pseudomonas species exhibit a wide variety of energy-generation metabolisms spanning from aerobic to anaerobic, involving NO_3^- reduction, complete denitrification process to N_2 and even fermentation of arginine or pyruvate (Arai, 2011; Tribelli et al., 2019). These energy-generating pathways enable *Pseudomonas* species to thrive in diverse environmental niches, including soil, water, and host-associated habitats, and they play an integral role in the ecological and physiological success of these microorganisms. *Pseudomonas extremaustralis* is an Antarctic bacterium, capable to growth under different temperatures and oxygen availability. The denitrification process in this bacterium is incomplete, due to the absence of *nir* genes which encode the enzymes required for the reduction of NO_2^- to NO, but interestingly, harbors all the *nor* and *nos* genes associated with nitric and nitrous oxide reduction, respectively (Raiger Iustman et al., 2015).

The bacterial adaptation to different stress conditions entails changes leading to solve metabolic and structural challenges, such us alterations or damage in DNA, RNA and proteins and envelopes, among others. Some strategies can involve the utilization of alternative pathways to exploit different carbon sources. Thus, when *P. putida* was grown at 10°C, the uptake and assimilation of branched-chain amino acids alongside the activation of the 2-methylcitrate pathway to generate succinate and pyruvate has been identified as mechanism employed to cope with stress when central metabolism is downregulated (Fonseca et al., 2011). Likewise, in cold conditions genes involved in primary metabolism were downregulated in *P. extremaustralis* whereas genes linked to ethanol oxidation were activated, showing that this secondary pathway resulted essential for cold growth (Tribelli et al., 2015). In *P. aeruginosa* has been reported that under oxidative stress the glyoxylate shunt, an alternative to the tricarboxylic acid cycle, increases bacterial survival allowing the utilization of acetate and fatty acids as carbon sources (da Cruz Nizer et al., 2021).

Considering the importance of RNS derived from both anaerobic respiration and other biotic and abiotic environmental reactions, we evaluated the global transcriptomic response of *P. extremaustralis* to nitrosative stress under low O_2 tensions using S-nitrosoglutation (GSNO), as a NO donor compound. Our findings show that exposure to GSNO under microaerobic conditions prompted adjustments in central metabolic pathways including iron metabolism along with the upregulation of genes associated with the inositol catabolism. Notably, *myo*-inositol arise as a carbon source supporting *P. extremaustralis* growth and increased nitro-oxidative stress resistance and pyoverdine production. These findings highlight the versatility and adaptability of *P. extremaustralis* to withstand nitrosative stress, by displaying an alternative metabolism, thus contributing to the understanding of the survival mechanisms employed by microorganisms in extreme environments.

Materials and methods

Strains and culture conditions

P. extremaustralis 14-3b (DSM 25547), a species isolated from the Antarctica (López et al., 2009) was used through the experiments. Bacterial cultures were grown in Lysogeny Broth medium (LB) at 30 °C. Microaerobic cultures were incubated in sealed bottles with a 1:2 medium-to-flask volume ratio and 50 rpm agitation. Microaerobic culture's medium was supplemented with 0.8 g/l KNO_3.

GSNO effect on *P. extremaustralis*' survival and growth was determined in microaerobic cultures exposed to 1, 10 and 100 µM of GSNO. Bacterial growth was evaluated by OD_{600nm} measurement after adding the different concentrations of GSNO at T=0. Optical density was monitored over 24 h at 30min intervals using an automated plate reader (BMG OPTIMA FLUOstar). For survival experiments, cultures were grown for 24h and further exposed to the different concentrations of GSNO for 1h. Afterwards, appropriate dilutions of control cultures (m-C) or cultures exposed to GSNO were plated in LB agar and incubated at 30°C. Colony-forming units per ml (CFU / ml) was determined and the survival percentage was calculated with respect to control cultures.

RNA extraction and RNA library preparation

Cultures were microaerobically grown in LB medium for 24 h and then incubated for 1 h with 100 µM S-nitrosoglutatione (GSNO, Sigma Aldrich) (m-NS condition) or with the addition of sterile water (m-C). Total RNA was isolated from 6 ml of *P. extremaustralis* m-C and m-NS cultures using the Trizol method. Samples were treated with DNAse I and were validated using an Agilent 2100 Bioanalyzer (Agilent Technologies). To improve the quality of the readings, ribosomal RNA was depleted from the samples, using the RiboZERO Kit (Illumina), following the manufacturer's instructions. Libraries were prepared using the TruSeq RNA Library Prep Kit v2 (Illumina). Mass sequencing

was performed using NextSeq 550 platform with a single-end protocol. For each

condition duplicated independent RNA extraction and libraries were used.

RNA-seq data analysis

Reads were preprocessed using the Trimmomatic tool (Bolger et al., 2014) by eliminating

adapters and low-quality sequences. Reads' quality was evaluated using the Fast QC

tool (www.bioinformatics.babraham.ac.uk/projects/fastqc/).

Reads alignment and assembly to the *P. extremaustralis* genome, transcript

identification and abundance quantification was carried out using the Rockhopper

software (Tjaden, 2020). Reads were normalized per kilobase per million mapped reads

(RPKM). Differential gene expression was considered only with $P < 0.05$ and $Q < 0.05$.

Concordance between the independent replicates for each of the analyzed conditions

was verified by performing a Spearman correlation analysis of normalized counts (Fig.

S1).

Genes were assorted into functional classes using KEGG (Ogata et al., 1999), MetaCyc

(Caspi et al., 2016) and String (Szklarczyk et al., 2019) tools.

Prediction of oxidation and nitrosylation targets

The Target-Pathogen tool (Sosa et al., 2018) was used to identify proteins with possible

oxidation and / or nitrosylation target sites (Free Cys or Tyr residues or metal biding

clusters) within the *P. extremaustralis* genome.

Comparative genome analysis

General genome sequence analysis was performed using the bioinformatics tools

available on National Center for Biotechnology Information (www.ncbi.nlm.nih.gov),

including BLAST (Basic Local Alignment Search Tool) (Altschul, 1997). We also used

RAST (Rapid Annotation using Subsystem Technology) server (Aziz et al., 2008) and

Pseudomonas Genome Database (Pseudomonas.com, (Winsor et al., 2016)). Genomes

used, with corresponding GenBank accession numbers, were: *P. extremaustralis:* 14-3b

(AHIP00000000.1), USBA 515 (FUYI01), DSM 17835^T (LT629689.1), 2E-UNGS

(CP091043.1), CSW01(JAQKGS01),1906 (JARIXU01), NQ5 (JARBJR01) strains, *P.*

protegens Pf-5 (CP000076.1), and *P. syringae* pv *syringae* B728a (CP000075.1)

Quantitative Real Time PCR Experiments (RT qPCR)

Total RNA of *P. extremaustralis* microaerobic and m-NS cultures was isolated using the

Total RNA Extraction Kit (RBC Biosciences). After treatment with DNaseI, cDNA was

obtained using random hexamers (Promega) and Revert Aid reverse transcriptase

(ThermoFisher Scientific) according to the manufacturer's instructions. RT qPCR was

performed using a MyiQ2 Real-Time PCR Detection System (Bio-Rad Laboratories,

Hercules, USA) and Real-Time PCR mix (EvaGreen qPCR Mix Plus no ROX). The

expression of target genes was evaluated using the following primers: *motB* forward 5′

CAACGCCAGCAACAAAGACA 3′ and reverse 5′ CTTCGGAAAACCGCGCATC 3′; *hbo*

forward 5′ CGAGCCTAAGAGCAACACCA 3′ and reverse

5′TGAATAGGCTGTCGGCACTG 3′ and *flaB* forward 5'

CGTAACCAGCGCTGACATGGCTC 3' and reverse 5'

GACAGGATCGCAGTGGAAGCCGA 3'. Expression of the 16S rRNA gene was used for

normalization of target genes expression levels in each condition, obtained using primers

forward 5′ AGCTTGCTCCTTGATTCAGC 3′ and reverse 5′

AAGGGCCATGATGACTTGAC' 3. The cycling conditions were as follows: denaturation

at 95 °C for 15 min, 40 cycles at 95 °C for 25 s, 60 °C for 15 s, and 72 °C for 15 s, with

fluorescence acquisition at 80 °C in single mode. The $2^{-\Delta\Delta CT}$ method (Livak & Schmittgen,

2001) was used to calculate the relative fold gene expression of individual genes.

Bacterial growth using *myo*-inositol as sole carbon source

193 *Myo*-inositol metabolism was evaluated in E2 medium (Lageveen et al., 1988)

194 supplemented with 10 g/l of *myo*-inositol or glucose under aerobic and microaerobic

conditions. Microaerobic cultures were performed as described above while for aerobic experiments, cultures were incubated in Erlenmeyer flasks using 1:10 medium-to-flask volume ratio and 200 rpm agitation. Aerobic and microaerobic cultures were incubated for 24h and 48h, respectively.

Biofilm formation was assayed in E2 medium supplemented with myo-inositol or glucose and KNO_3 in polystyrene microtiter plates with an initial OD_{600nm} of 0.025. After 48h of static incubation at 30°C, OD_{600nm} of planktonic cells (Planktonic cells absorbance, PCA) was measured and biofilms were quantified using the standard crystal violet method (O'Toole & Kolter, 1998). Briefly, attached cells were stained with 200 µl of 0.1% crystal violet, further washed and the colorant was solubilized with absolute ethanol. Crystal violet solution was transferred to flat bottom microtiter plates and OD_{570nm} (Crystal violet absorbance CVA) was measured with a BMG OPTIMA FLUOstar microplate reader. Biofilm formation index was defined as CVA/PCA.

Pyoverdine production

Pyoverdine production of microaerobic cultures was analyzed in iron-limited E2 medium, without microelements, supplemented with myo-inositol or glucose as carbon source and KNO_3. After 48h pyoverdines in the culture supernatants were determined by measuring integrated fluorescence emission between 445-460 nm after excitation at 420 nm (Olmo et al., 2003; Ruiz et al., 2015). The values were expressed relative to cell dry weight per ml.

Survival and stress experiments

P. extremaustralis was cultured for 48h under microaerobic conditions in E2 cultures supplemented with myo-inositol or glucose and KNO_3 (0.8 g/l). Cultures were exposed to 100 µM GSNO (m-NS) for 1h and cell viability was determined in plate assay.

Growth inhibition in response to oxidative or combined nitro-oxidative stress was evaluated by filter disk assay by growing *P. extremaustralis* in E2 medium supplemented

with glucose or *myo*-inositol and increasing KNO_3 concentrations of 0.8, 1.5 and 2.5 g/l

that led to nitrite accumulation. Control cultures without nitrate were performed (oxidative

stress only). LB plates were seeded with the different cultures and Whatman n° 1 filter

discs (6 mm) were impregnated with 5 µl of 30% (v/v) H_2O_2 (Merck) as previously

described (Ayub et al., 2004). Plates were incubated overnight and the diameter of the

halo was determined using the software ImageJ (Schneider et al., 2012). Nitrite

concentration in the supernatant of cultures was determined following the method

described by (Gerhard et al., 1981) and values were normalized to cell dry weight.

Data availability

RNA-seq data were deposited in the European Molecular Biology Laboratory under

accession number E-MTAB-11689.

Statistical analysis

Differences between means were determined through the Student's t test with

confidence levels at > 95% in which $P < 0.05$ was considered as statistically significant.

A Fisher's exact test was performed to analyze the nitrosylation targets proteins. 1 or 2-

ways ANOVA with multiple comparison were used when correspond.

Results

Effects of GSNO exposure in *P. extremaustralis*

Nitrosative stress effect on *P. extremaustralis* survival and growth under microaerobic

conditions was evaluated using different S-nitrosoglutation (GSNO) concentrations.

Growth in presence of different concentrations of GSNO was similar to the control culture

except for 100 µM that showed a decrease at the stationary phase (Fig. S2). Moreover,

survival experiments showed that after 1h GSNO exposure, a significant decrease in

bacterial viable counts (CFU/ml) was observed comparing with the control culture (Fig.

1). These results indicate GSNO exposure affected *P. extremaustralis* survival. Thus, we

chose 100 µM GSNO (m-NS) exposure for 1h which provoked a 75% drop in survival in

microaerobic cultures for RNAseq experiments.

Figure 1.   Survival of microaerobic cultures to GSNO exposure. Circles represent

individual values of independent experiments. Error bars represent the standard

deviation of the mean. Different letters denote significant differences using 1-way

ANOVA with Tukey's multiple comparisons test (P < 0.05)

Transcriptomic profile of *P. extremaustralis* under low aeration and nitrosative

stress conditions.

P. extremaustralis response to nitrosative stress was evaluated by comparing

transcriptomic data obtained by RNA sequencing (RNA-Seq) in microaerobic cultures m-

C and m-NS. Data analysis showed 249 transcripts with differential expression in m-NS

compared to m-C (Rockhopper, P <0.05, Q <0.05). Among them, 86 genes were found

to be repressed and 163 over-expressed, representing 1.4% and 2.7% of the total,

respectively (Fig. S3a, Table S1a). Several genes with differential expression did not

have defined associated functions and were classified as "hypothetical", representing 11.6% and 21% of the over-expressed and repressed genes, respectively. Among genes with a known predicted function, *P. extremaustralis* under m-NS showed an increased expression of genes involved in carbon, RNA, DNA and iron metabolism (Table S1a). RNA-seq results were validated by analyzing the expression of three selected genes, *motB*, *flab*, and *hbo* using qRT PCR (Fig. S2b). In concordance with the RNA-seq results, we found higher expression levels of *motB,* encoding a flagellar motor rotation protein, in m-NS vs. m-C, whereas we observed a decrease in *hbo* (Fig. S2b, Table S1a) and no differences in *flaB* expression between conditions.

P. extremaustralis' response to nitrosative stress

To elucidate possible functional relationships, differentially expressed genes were categorized into functional groups (Fig. 2a). GSNO exposure caused increased expression of genes encoding different RNS detoxifying enzymes such as an iron-dependent superoxide dismutase (PE143B_0125285) and an alkyl hydroperoxide reductase (*ahpC*), a thioredoxin (PE143B_0111145) and a glutathione S-transferase (PE143B_0109905). A gene encoding another glutathione S-transferase (PE143B_0100180) was repressed in m-NS along with *hbO* (Fig. 2a, Table S1a).

Transcriptomic analysis showed an increase in mRNA expression of iron and heme transporters coding genes such as *ctaB, ybaN, irpA* and *afnA* (Fig 2b, Table S1a). In m-NS we found upregulation in genes related to pyoverdine siderophore biosynthesis, a key function for iron uptake, including *pdvA, pdvM, pdvF, fpvA, fpvE* and *fpvK* (Fig. 2b, Table S1a). Moreover, the arginase coding gene, was also upregulated in m-NS comparing with m-C. Arginase catalyzes the conversion of arginine to ornithine, which can then serve as a substrate for pyoverdine biosynthesis (Fig. 2b, Table S1a).

Iron and other metals are key components of iron-sulfur proteins that could be target of RNS, leading to protein inactivation and triggering cellular signaling (Seth et al., 2020). Therefore, we analyzed other cellular functions related with transcription and protein

turn-over. GSNO treated cultures showed an increase in expression of RNA helicase

(PE143B_0107010), RNA polymerase associated protein RapA, transcriptional

terminator *nusB* (PE143B_0123570) and 11 ribosomal proteins coding genes (Table

S1a). In concordance, a gene encoding translation initiation inhibitor was repressed

(Table S1a). To deeply analyze the protein turnover related with RNS attack we

performed an analysis using Target Pathogen platform to detect genes encoding proteins

with free tyrosine or cysteine as well as metal binding sites which are reactive to RNS

(Crack et al., 2014). Among differentially expressed genes in m-NS we found that

nitrosylable proteins were over-represented in comparison to non-nitrosylable when total

genes of *P. extremaustralis* were considered (Fisher's exact test, P = 0.0323). Overall,

our results suggest that nitrosative stress under microaerobic conditions provokes an

increase in the expression of genes involved in transcription and translation processes,

probably to compensate the damage to proteins with Fe-S clusters or metal binding (Fig.

2c).

Figure 2. Effect of GSNO on gene expression profile. a. Classification of differentially

expressed genes after GSNO exposure in microaerobiosis (m-NS vs. m-C) into

functional categories. Hypothetical transcripts (with no inferred function) are not

displayed. Green and red bars represent up- and down-regulated genes, respectively.

b. Over-expressed genes related to iron metabolism. **c**. Nitrosylable proteins determined

with Target Pathogens in the differential expressed genes compared to total nytrosilable

proteins in *P. extremaustralis* genome. DE: differentially expressed genes. Fisher's exact

test, P = 0.0323.

Inositol metabolism in *P. extremaustralis*

Remarkably, RNAseq results showed increased expression of genes related to inositol

catabolism which involves enzymes that convert inositol into acetyl-CoA and

dihydroxyacetone phosphate (DHAP) (Fig. 3a, Table S1a).

Our results showed an up-regulation in mRNAs encoded in the *iol* operon in response to

GSNO stress, including *iolB* (2-fold), *iolD* (1.75-fold), *iolE* (1.83-fold) *iolG2* (also known

as *iolW* or *mocA*) (1.86-fold), *iolR* (2.62-fold), and the *iatA* gene (1.75-fold) (Fig. 3a; Table

S1a, b). No differences were observed in the expression of *iolA, iolG1* and *iolC* in both

conditions, neither for *iatP* and *ibpA* (Fig. 3a, Table S1b). It was found that *iolC*

possesses the DUF2090 domain that was recently proposed to encode 2-deoxy-5-keto-

gluconic acid-6-phosphate aldolase (Price et al., 2022). This domain fulfills function

originally attributed to *iolJ* which is not present in *P. extremaustralis* genome (Fig. 3b)*.*

When we compared the inositol catabolic cluster of *P. extremaustralis* with *P. protegens*

Pf-5, a well-known plant growth promoter species, and *P. syringae* pv. syringae B728a,

a plant pathogen, we found some differences in this genomic region. Unlike *P. protegens*

Pf-5, the genome of *P. extremaustralis* did not exhibit the presence of *iolL*. Conversely,

we identified a duplication of *iolG* within this bacterial species. *P. syringae* pv. syringae

B728a presented *iolH*, which is absent in both *P. protegens* and *P. extremaustralis.*

Similar to *P. extremaustralis, P. syringae* pv. syringae B728a presented two copies of

iolG but separated by *iolH* (Fig. 3b). The *iolA* gene in *P. extremaustralis* is located in a

different genomic region like in *P. protegens* and other analyzed *Pseudomonas*

(Sánchez-Gil et al., 2023). However, in P. *syringae* pv. syringae B728a is included in the

iol catabolic cluster (Fig. 3b). Furthermore, we found that the *iol* gene cluster of *P.*

extremaustralis exhibited a high degree of intraspecific conservation showing the same

organization as 14-3b in all genomes analyzed, including strains USBA 515,

DSM17835^T, 2E-UNGS, CSW01,1906, NQ5.

Figure 3. Metabolic pathway associated with inositol catabolism in *Pseudomonas*

***extremaustralis* 14-3b: Expression under nitrosative stress and genetic**

organization. **a**. Gene expression and metabolic route related to inositol catabolism

under m-NS Genes and arrows shown in green indicate up-regulated functions under m-

NS, whereas those in blue represent functions with no expression differences between both conditions. **b**. Analysis of genes involved in inositol in *P. extremaustralis* in comparison with *P. protegens* Pf-5 and *P. syringae* pv. syringae B728a. (*) The same genetic organization was observed in USBA 515, DSM17835^T, 2E-UNGS, CSW01,1906 and NQ5 *P. extremaustralis* strains. Compounds: KMI, 2-keto-*myo*-inositol; THcHDO, 3, 3D-(3,4/5) trihydroxycyclohexane-1,2-dione; 5DG, 5-deoxy glucuronic acid; DKG, 2-deoxy-5-keto-d-gluconic acid; DKGP, DKG 6-phosphate; DHAP, dihydroxyacetone phosphate; MSA, malonic semialdehyde; acetyl-CoA, acetyl coenzyme A; DHAP, dihydroxyacetone phosphate; G3P, glyceraldeyde-3-phosphate; TCA, tricarboxylic acid cycle. Genes: *iolG1*, *myo*-inositol dehydrogenase; *iolG2*, *scyllo*-inositol dehydrogenase; *iolE*, 2KMI dehydratase; *iolD*, THcHDO hydrolase; *iolB*, 5DG isomerase; *iolC/J*, DKG kinase + aldolase; *iolA*, MSA dehydrogenase; *iatA*, sugar ABC transporter ATP-binding protein; *iatP*, ABC transporter permease; *ibpA*, sugar ABC transporter substrate-binding protein.

As we detected the presence and expression of *iol* genes, we first tested *P. extremaustralis* capability to grow under aerobic conditions using *myo*-inositol as sole carbon source. Growth was similar between *myo*-inositol and glucose supplemented cultures reaching an OD_{600nm} value of 2.02 ± 0.15 and 2.22 ± 0.15, respectively. Under microaerobic conditions *P. extremaustralis* developed an evident biofilm in *myo*-inositol supplemented cultures (Fig. 4a). Therefore, OD_{600nm} and cellular dry weight (DW) were determined (Fig. 4b). Similar results were obtained for both carbon sources regardless of the approach employed for growth estimation. Biofilm formation was assayed in E2 medium supplemented with glucose or inositol. In concordance, we also found that with *myo*-inositol supplemented cultures *P. extremaustralis* showed a higher biofilm index compared to glucose (Fig. 4c).

Additionally, considering the upregulation of coding genes related with pyoverdine biosynthesis after treatment with GSNO, we investigated siderophores production in *P.*

extremaustralis using *myo*-inositol as carbon source. Pyoverdine production was higher

for *myo*-inositol supplemented cultures comparing to glucose (Fig. 4d)

Figure 4. Physiological features of *P. extremaustralis* using *myo*-inositol o glucose

as carbon source. a. Photograph of microaerobic *myo*-inositol or glucose supplemented

cultures showing the biofilm in the surface. **b.** Microaerobic growth measured by optical

density (OD_{600nm}) or Cellular dry weight (DW). **c.** Biofilm formation index (ACV/APC) in

myo-inositol or glucose. **d.** Effect of carbon sources on pyoverdine production. RFU:

Relative Fluorescence Units. Fluorescence units were normalized to the cell dry

weight/ml. Error bars represent the standard deviation of the mean.

***Myo*-inositol impact on stress response**

Further, we investigated stress resistance when *myo*-inositol was used as sole carbon

source under low oxygen conditions. To analyze the nitrosative stress response, we

performed a survival test using microaerobic cultures in E2 medium supplemented with

myo-inositol or glucose exposed to GSNO treatment. We found no differences in survival

between those treated with GSNO and control cultures despite of the carbon source (Fig.

5a).

P. extremaustralis is only capable to reduce nitrate to nitrite which accumulates in the

extracellular medium. Therefore, the resistance to nitro-oxidative stress derivate from the

combination of the nitrite accumulated and H_2O_2 was analyzed in *myo*-inositol and

glucose supplemented cultures, following the scheme shown in Fig. S4. Nitrite

production was detected in cultures supplemented with *myo*-inositol and glucose as sole

carbon source and different nitrate concentrations reaching a maximum of 13.553±

1.624, confirming also the *myo*-inositol utilization in a respiratory pathway (Fig. 5b).

When *P. extremaustralis* was grown in *myo*-inositol or glucose supplemented media

without nitrate, the oxidative stress resistance was similar for both carbon sources while

the nitro-oxidative stress resistance was higher for cultures grown in *myo*-inositol in all

tested KNO_3 concentrations (Fig. 5c).

Figure 5. Nitrosative and combined nitro-oxidative stress resistance in *P. extremaustralis*. a. Survival after GSNO exposure. **b.** Nitrite production in cultures supplemented with *myo*-inositol or glucose and different nitrate concentrations. **c.** Inhibition halos of cultures performed in (**b**) exposed to H_2O_2 (Combined nitro-oxidative stress). Error bars represent the standard deviation of the mean.

Discussion

Bacterial species are subjected to different stress conditions including lack of nutrients, changes in temperature or oxygen availability and oxidative and nitrosative stress, among others. Oxidative and nitrosative stress can be produced by the imbalance between the ROS or RNS generation and the incapability to detoxify them (Chautrand et al., 2022).

In response to these stress conditions, bacteria may activate various response mechanisms, such as the production of chaperons and detoxicant enzymes, the

adjustment of gene expression, and the modification of their cell envelope composition.

A common effect during stress is the reprograming of transcription and translation

processes including the reduction of ribosome biogenesis, the modification of the

translational machinery and the regulation of initiation and elongation of certain proteins

(Advani & Ivanov, 2019). Interestingly, gene expression during stress involves not only

the transcription processes but also the modification of mRNA stability and in overall with

translation provokes a transcriptome stability (Kristoffersen et al., 2012; Vargas-Blanco

& Shell, 2020). In this work, we found an up-regulation of the transcription and translation

machinery probably to compensate the damage caused by RNS particularly on

nitrosylable proteins. The Fe-S clusters, that can be found in three different forms, exhibit

high capacity of accepting or donating electrons thus are important for redox response

and act as redox sensors like the Fnr regulator in *Escherichia coli* or Anr in

Pseudomonas species (Galimand et al., 1991; Golinelli-Cohen & Bouton, 2017; Vernis

et al., 2017).These proteins and those containing metals are involved in several essential

cellular functions like respiration, central carbon catabolism and RNA and DNA

processes and are a target of RNS damage (Vernis et al., 2017) Particularly, it has been

reported that the reaction between nitric oxide (NO) with Fe-S clusters lead to protein

degradation and breakdown of the cluster generating dinitrosyl iron complexes (Harrop

et al., 2008). Our analysis using Target Pathogen platform showed the enrichment of

genes encoding nitrosylable proteins in the differentially expressed ones, particularly in

those upregulated. Our hypothesis is that to cope with GSNO derived stress the Fe-S

and metal binding proteins need to be replaced. This also could explain the upregulation

of Fe-uptake related genes observed in presence of GSNO. Iron is involved in the Fenton

reaction in presence of H_2O_2, derived from endogenous or exogenous sources, that

could exacerbate the oxidative damage but paradoxically this metal is also necessary for

Fe-S and metal protein biogenesis (Reniere, 2018; Winterbourn, 1995) . This complex

scenario additionally requires the involvement of enzymes and antioxidant cycles. In

presence of GSNO we found upregulated the coding genes of glutathione S-transferase,

superoxide dismutase (SOD), thiol-disulfide isomerase, thioredoxins and alkyl

hydroperoxide reductase subunit C-like protein. The alkyl hydroperoxide reductase,

important during non-lethal stress, SOD as well as the oxidation of thioredoxins in

response to oxidative stress depends on NAD(P)H availability and in overall reduction

power is needed for the essential process for survival and to cope with stress (Lemire et

al., 2017) .Interestingly, the needed of NADH and/or NADPH production during stress

could represent a challenge for bacterial cells. Glycolysis and TCA provides the NADH

and energy necessaries for bacterial survival but also creates an oxidative scenario

through the respiration chain (Zhang et al., 2019). It has been reported several metabolic

strategies to obtain energy and reducing power besides glycolysis such as aminoacid

catabolism, ethanol oxidation and in *P. fluorescens* an entire metabolic reprogramming

has been reported in presence of oxidative stress to generate NADPH (Singh et al.,

2007). In this previous work the authors showed an increase in the activity and

expression of malic enzyme (similar to this work in presence of GSNO), glucose-6-

phosphate dehydrogenase and NADP+-isocitrate dehydrogenase (Singh et al., 2007).

In addition, we found that in presence of GSNO under microaerobic conditions *P.*

extremaustralis over-expressed several *iol* catabolic genes including the transcriptional

regulator and the *iatA* gene. We also demonstrated that this bacterium was effectively

capable of using inositol as sole carbon source. Inositol is a sugar alcohol present in all

domains of life and presents various structural isomers (Michell, 2008), with *myo*-inositol

being the most abundant in nature. *Myo*-inositol is found in high quantities in plants,

including root exudates, and may also originate from the dephosphorylation of phytate,

one of the major phosphorus storage molecules for plants (Weber & Fuchs, 2022).

Inositol catabolism has been experimentally demonstrated in several bacterial species,

both Gram positive and Gram negative, belonging to genera such as   *Rhizobium,*

Lactobacillus, Klebsiella, Corynebacterium, Legionella, Bacillus, and Salmonella (Fry et

al., 2001; Krings et al., 2006; Kröger et al., 2018; Manske et al., 2016; Morinaga et al.,

2010; Yebra et al., 2007; Yuan et al., 2019). Furthermore, an extensive analysis of

inositol catabolic genes in thousands of bacterial genomes revealed a widespread
distribution within different ecological niches (Weber & Fuchs, 2022).

Although recently inositol metabolism in bacteria has received many interest, its
role in stress resistance has not been studied. Remarkably, our findings revealed that
inositol protects against nitro-oxidative stress, and increases biofilm and pyoverdine
production, which are relevant traits for environmental adaptability. Inositol catabolism
leads to the production of glyceraldehyde-3 phosphate, and acetyl-CoA both
intermediaries of central energy generation metabolism. In addition, *iolG2,* which was
found overexpressed after GSNO exposure, is related to NAPH generation, necessary
for the function of antioxidative defenses.

It is worth noting that previous transcriptome analysis of *P. extremaustralis* performed
under cold conditions or under microaerobic conditions with or without oxidative stress
did not show the upregulation of inositol catabolic genes (Solar Venero et al., 2019;
Tribelli et al., 2015, 2018), indicating a fine-tuning response to nitrosative stress.

Some RNS also play a crucial role in cell signaling. For instance, in plants, NO mediates
for various processes including root hair growth, stomatal closure, programmed cell
death and other responses that help them adapt to changing environmental conditions
(Zaffagnini et al., 2012) whereas in bacteria, NO plays a role as intermediary in
denitrification processes, but also in biofilm formation and quorum sensing (Williams &
Boon, 2019). The *iol* gene cluster was identified in both plant-pathogenic bacteria, and
rhizospheric plant growth promoting bacteria forming symbiotic root nodules (Weber &
Fuchs, 2022).

Recently, the *iol* gene cluster in *Pseudomonas* was identified as an important trait in root
colonizers, by increasing swimming motility and siderophore production in response to
plant derived inositol (Sánchez-Gil et al., 2023). Increased pyoverdine siderophore
production in presence of *myo*-inositol was also observed in this work. *P. extremaustralis*
possesses plant growth-promotion traits, including the ability to solubilize phosphate and
produce indole acetic acid (Ibarra, 2017). Since NO is also a signaling molecule released

by plants (Sasaki et al., 2016), we hypothesize that NO released by GSNO mimic the
NO production of plants, triggering the differential expression of the inositol metabolic
pathway conferring an adaptive advantage.

In conclusion, our findings reveal a multifaceted and finely-tuned response mechanism
to nitrosative stress that encompasses transcriptional reprogramming, the upregulation
of genes encoding key proteins involved in iron homeostasis, and the activation of
inositol catabolism. Notably, experimental results point to inositol metabolism as a novel
mechanism to cope with stress.

Acknowledgments

This work was funded by grants from Universidad de Buenos Aires and Consejo
Nacional de Investigaciones Científicas y Técnicas (CONICET). We are thankful to Dr.
Jörg Vogel (Institut für Molekulare Infektionsbiologie (IMIB) – University of Würzburg) for
his guidance and for providing resources for conducting the RNAseq experiments.
E.C.S.V. received a Deutscher Akademischer Austauschdienst (DAAD) fellowship for
short-term research stays at the RNA biology group (IMIB – University of Würzburg,
Germany).

References

Advani, V. M., & Ivanov, P. (2019). Translational Control under Stress: Reshaping the
Translatome. *BioEssays*, *41*(5). https://doi.org/10.1002/bies.201900009

Altschul, S. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein
database search programs. *Nucleic Acids Research*, *25*(17), 3389–3402.
https://doi.org/10.1093/nar/25.17.3389

Arai, H. (2011). Regulation and Function of Versatile Aerobic and Anaerobic Respiratory
Metabolism in Pseudomonas aeruginosa. *Frontiers in Microbiology*, *2*(2), 103.
https://doi.org/10.3389/fmicb.2011.00103

Aranda-Rivera, A. K., Cruz-Gregorio, A., Arancibia-Hernández, Y. L., Hernández-Cruz,
E. Y., & Pedraza-Chaverri, J. (2022). RONS and Oxidative Stress: An Overview of
Basic Concepts. *Oxygen*, *2*(4), 437–478. https://doi.org/10.3390/oxygen2040030

Ayub, N. D., Pettinari, M. J., Ruiz, J. A., & López, N. I. (2004). A polyhydroxybutyrate-
producing Pseudomonas sp. isolated from antarctic environments with high stress

resistance. *Current Microbiology, 49*(3), 170–174. https://doi.org/10.1007/s00284-
004-4254-2

Aziz, R. K., Bartels, D., Best, A. A., DeJongh, M., Disz, T., Edwards, R. A., Formsma, K.,
Gerdes, S., Glass, E. M., Kubal, M., Meyer, F., Olsen, G. J., Olson, R., Osterman,
A. L., Overbeek, R. A., McNeil, L. K., Paarmann, D., Paczian, T., Parrello, B., …
Zagnitko, O. (2008). The RAST Server: Rapid Annotations using Subsystems
Technology. *BMC Genomics, 9*(1), 75. https://doi.org/10.1186/1471-2164-9-75

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for
Illumina sequence data. *Bioinformatics, 30*(15), 2114–2120.
https://doi.org/10.1093/bioinformatics/btu170

Caspi, R., Billington, R., Ferrer, L., Foerster, H., Fulcher, C. A., Keseler, I. M., Kothari,
A., Krummenacker, M., Latendresse, M., Mueller, L. A., Ong, Q., Paley, S.,
Subhraveti, P., Weaver, D. S., & Karp, P. D. (2016). The MetaCyc database of
metabolic pathways and enzymes and the BioCyc collection of pathway/genome
databases. *Nucleic Acids Research, 44*(D1), D471-480.
https://doi.org/10.1093/nar/gkv1164

Chautrand, T., Souak, D., Chevalier, S., & Duclairoir-Poc, C. (2022). Gram-Negative
Bacterial Envelope Homeostasis under Oxidative and Nitrosative Stress.
Microorganisms, 10(5). https://doi.org/10.3390/microorganisms10050924

Crack, J. C., Thomson, A. J., & Brun, N. E. Le. (2014). *Iron − Sulfur Clusters as Biological
Sensors : The Chemistry of Reactions with Molecular Oxygen and Nitric Oxide*.

Czaja, A. J. (2017). *Immunopathogenesis of Autoimmune Liver Damage* (pp. 19–48).
https://doi.org/10.1016/B978-0-444-63707-9.00002-7

da Cruz Nizer, W. S., Inkovskiy, V., Versey, Z., Strempel, N., Cassol, E., & Overhage, J.
(2021). Oxidative Stress Response in Pseudomonas aeruginosa. *Pathogens, 10*(9),
1187. https://doi.org/10.3390/pathogens10091187

Del Río, L. A. (2015). *ROS and RNS in plant physiology: an overview.*
https://doi.org/10.1093/jxb/erv099

Fonseca, P., Moreno, R., & Rojo, F. (2011). Growth of Pseudomonas putida at low
temperature: Global transcriptomic and proteomic analyses. *Environmental
Microbiology Reports*. https://doi.org/10.1111/j.1758-2229.2010.00229.x

Fry, J., Wood, M., & Poole, P. S. (2001). Investigation of *myo* -Inositol Catabolism in
Rhizobium leguminosarum bv. *viciae* and Its Effect on Nodulation Competitiveness.
Molecular Plant-Microbe Interactions®, 14(8), 1016–1025.
https://doi.org/10.1094/MPMI.2001.14.8.1016

Galimand, M., Gamper, M., Zimmermann, A., & Haas, D. (1991). Positive FNR-like
control of anaerobic arginine degradation and nitrate respiration in Pseudomonas
aeruginosa. *Journal of Bacteriology, 173*(5), 1598–1606.
http://www.ncbi.nlm.nih.gov/pubmed/1900277

Gerhard, P., Murray, R., Costilow, N., Nester, W., Wood, N., Krieg, N., & Phillips, G.
(1981). *Manual of Method of General Bacteriology.* American Society for
microbiology.

Golinelli-Cohen, M.-P., & Bouton, C. (2017). Fe-S Proteins Acting as Redox Switch: New
Key Actors of Cellular Adaptive Responses. *Current Chemical Biology*, *11*(2).
https://doi.org/10.2174/2212796811666170406163809

Harrop, T. C., Tonzetich, Z. J., Reisner, E., & Lippard, S. J. (2008). Reactions of synthetic
[2Fe-2S] and [4Fe-4S] clusters with nitric oxide and nitrosothiols. *Journal of the*
American Chemical Society, *130*(46), 15602–15610.
https://doi.org/10.1021/ja8054996

Ibarra, J. G. (2017). *Estudios microbiológicos relacionados con el mejoramiento de*
cultivos vegetales en zonas desfavorables. [Doctoral Thesis]. Facultad de Ciencias
Exactas y Natural.

Krings, E., Krumbach, K., Bathe, B., Kelle, R., Wendisch, V. F., Sahm, H., & Eggeling,
L. (2006). Characterization of *myo* -Inositol Utilization by *Corynebacterium*
glutamicum : the Stimulon, Identification of Transporters, and Influence on
<scp>l</scp> -Lysine Formation. *Journal of Bacteriology*, *188*(23), 8054–8061.
https://doi.org/10.1128/JB.00935-06

Kristoffersen, S. M., Haase, C., Weil, M. R., Passalacqua, K. D., Niazi, F., Hutchison, S.
K., Desany, B., Kolstø, A.-B., Tourasse, N. J., Read, T. D., & Økstad, O. A. (2012).
Global mRNA decay analysis at single nucleotide resolution reveals segmental and
positional degradation patterns in a Gram-positive bacterium. *Genome Biology*,
13(4), R30. https://doi.org/10.1186/gb-2012-13-4-r30

Kröger, C., Rothhardt, J. E., Brokatzky, D., Felsl, A., Kary, S. C., Heermann, R., & Fuchs,
T. M. (2018). The small RNA RssR regulates myo-inositol degradation by
Salmonella enterica. *Scientific Reports*, *8*(1), 17739.
https://doi.org/10.1038/s41598-018-35784-8

Lageveen, R. G., Huisman, G. W., Preusting, H., Ketelaar, P., Eggink, G., & Witholt, B.
(1988). Formation of Polyesters by Pseudomonas oleovorans : Effect of Substrates
on Formation and Composition of Poly-(R)-3-Hydroxyalkanoates and Poly-(R)-3-
Hydroxyalkenoates . *Applied and Environmental Microbiology*, *54*(12), 2924–2932.
https://doi.org/10.1128/AEM.54.12.2924-2932.1988

Lelieveld, S., Wilson, J., Dovrou, E., Mishra, A., Lakey, P. S. J., Shiraiwa, M., Pöschl, U.,
& Berkemeier, T. (2021). Hydroxyl Radical Production by Air Pollutants in Epithelial
Lining Fluid Governed by Interconversion and Scavenging of Reactive Oxygen
Species. *Cite This: Environ. Sci. Technol*, *55*, 14069–14079.
https://doi.org/10.1021/acs.est.1c03875

Lemire, J., Alhasawi, A., Appanna, V. P., Tharmalingam, S., & Appanna, V. D. (2017).
Metabolic defence against oxidative stress: the road less travelled so far. *Journal of*
Applied Microbiology, *123*(4), 798–809. https://doi.org/10.1111/jam.13509

Li, H., Zhou, X., Huang, Y., Liao, B., Cheng, L., & Ren, B. (2020). Reactive Oxygen
Species in Pathogen Clearance: The Killing Mechanisms, the Adaption Response,
and the Side Effects. *Frontiers in Microbiology*, *11*, 622534.
https://doi.org/10.3389/fmicb.2020.622534

Livak, K. J., & Schmittgen, T. D. (2001). Analysis of Relative Gene Expression Data
Using Real-Time Quantitative PCR and the 2−ΔΔCT Method. *Methods*, *25*(4), 402–
408. https://doi.org/10.1006/meth.2001.1262

López, N. I., Pettinari, M. J., Stackebrandt, E., Tribelli, P. M., Põtter, M., Steinbüchel, A.,
& Méndez, B. S. (2009). *Pseudomonas extremaustralis* sp. nov., a Poly(3-
hydroxybutyrate) producer isolated from an antarctic environment. *Current
Microbiology*, *59*(5), 514–519. https://doi.org/10.1007/s00284-009-9469-9

Manske, C., Schell, U., & Hilbi, H. (2016). Metabolism of *myo* -Inositol by Legionella
pneumophila Promotes Infection of Amoebae and Macrophages. *Applied and
Environmental Microbiology*, *82*(16), 5000–5014.
https://doi.org/10.1128/AEM.01018-16

Michell, R. H. (2008). Inositol derivatives: evolution and functions. *Nature Reviews
Molecular Cell Biology*, *9*(2), 151–161. https://doi.org/10.1038/nrm2334

Morinaga, T., Ashida, H., & Yoshida, K.-I. (2010). Identification of two scyllo-inositol
dehydrogenases in Bacillus subtilis. *Microbiology (Reading, England)*, *156*(Pt 5),
1538–1546. https://doi.org/10.1099/mic.0.037499-0

Ogata, H., Goto, S., Sato, K., Fujibuchi, W., Bono, H., & Kanehisa, M. (1999). KEGG:
Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Research*, *27*(1), 29–
34. https://doi.org/10.1093/nar/27.1.29

Olmo, A. Del, Caramelo, C., & Sanjose, C. (2003). F luorescent complex of pyoverdin
with aluminum. *Journal of Inorganic Biochemistry*, *97*, 384–387.
https://doi.org/10.1016/S0162-0134(03)00316-7

O'Toole, G. A., & Kolter, R. (1998). Flagellar and twitching motility are necessary for
Pseudomonas aeruginosa biofilm development. *Molecular Microbiology*, *30*(2),
295–304. https://doi.org/10.1046/j.1365-2958.1998.01062.x

Price, M. N., Deutschbauer, A. M., & Arkin, A. P. (2022). Filling gaps in bacterial catabolic
pathways with computation and high-throughput genetics. *PLOS Genetics*, *18*(4),
e1010156. https://doi.org/10.1371/journal.pgen.1010156

Raiger Iustman, L. J., Tribelli, P. M., Ibarra, J. G., Catone, M. V, Solar Venero, E. C., &
López, N. I. (2015). Genome sequence analysis of *Pseudomonas extremaustralis*
provides new insights into environmental adaptability and extreme conditions
resistance. *Extremophiles. 19*(1), 207–220. https://doi.org/10.1007/s00792-014-
0700-7

Reniere, M. L. (2018). Reduce, Induce, Thrive: Bacterial Redox Sensing during
Pathogenesis. *Journal of Bacteriology*, *200*(17). https://doi.org/10.1128/JB.00128-
18

Robinson, J. L., Adolfsen, K. J., & Brynildsen, M. P. (2014). Deciphering nitric oxide
stress in bacteria with quantitative modeling. *Current Opinion in Microbiology*, *19*,
16–24. https://doi.org/10.1016/j.mib.2014.05.018

Ruiz, J. A., Bernar, E. M., & Jung, K. (2015). Production of Siderophores Increases
Resistance to Fusaric Acid in Pseudomonas protegens Pf-5. *PLOS ONE*, *10*(1),
e0117040. https://doi.org/10.1371/journal.pone.0117040

Sánchez-Gil, J. J., Poppeliers, S. W. M., Vacheron, J., Zhang, H., Odijk, B., Keel, C., &
de Jonge, R. (2023). The conserved iol gene cluster in Pseudomonas is involved in
rhizosphere competence. *Current Biology*, *33*(15), 3097-3110.e6.
https://doi.org/10.1016/j.cub.2023.05.057

Sasaki, Y., Oguchi, H., Kobayashi, T., Kusama, S., Sugiura, R., Moriya, K., Hirata, T., Yukioka, Y., Takaya, N., Yajima, S., Ito, S., Okada, K., Ohsawa, K., Ikeda, H., Takano, H., Ueda, K., & Shoun, H. (2016). Nitrogen oxide cycle regulates nitric oxide levels and bacterial cell signaling. *Scientific Reports*, *6*(1), 22038. https://doi.org/10.1038/srep22038

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nature Methods*, *9*(7), 671–675. http://www.ncbi.nlm.nih.gov/pubmed/22930834

Seth, D., Hausladen, A., & Stamler, J. S. (2020). Anaerobic Transcription by OxyR: A Novel Paradigm for Nitrosative Stress. *Antioxidants & Redox Signaling*, *32*(12), 803–816. https://doi.org/10.1089/ars.2019.7921

Singh, R., Mailloux, R. J., Puiseux-Dao, S., & Appanna, V. D. (2007). Oxidative stress evokes a metabolic adaptation that favors increased NADPH synthesis and decreased NADH production in Pseudomonas fluorescens. *Journal of Bacteriology*, *189*(18), 6665–6675. https://doi.org/10.1128/JB.00555-07

Solar Venero, E. C., Ricardi, M. M., Gomez-Lozano, M., Molin, S., Tribelli, P. M., & López, N. I. (2019). Oxidative stress under low oxygen conditions triggers hyperflagellation and motility in the Antarctic bacterium Pseudomonas extremaustralis. *Extremophiles*, *23*(5). https://doi.org/10.1007/s00792-019-01110-x

Sosa, E. J., Burguener, G., Lanzarotti, E., Defelipe, L., Radusky, L., Pardo, A. M., Marti, M., Turjanski, A. G., & Fernández Do Porto, D. (2018). Target-Pathogen: a structural bioinformatic approach to prioritize drug targets in pathogens. *Nucleic Acids Research*, *46*(D1), D413–D418. https://doi.org/10.1093/nar/gkx1015

Szklarczyk, D., Gable, A. L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., Simonovic, M., Doncheva, N. T., Morris, J. H., Bork, P., Jensen, L. J., & Mering, C. von. (2019). STRING v11: protein–protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Research*, *47*(D1), D607–D613. https://doi.org/10.1093/nar/gky1131

Tharmalingam, S., Alhasawi, A., Appanna, V. P., Lemire, J., & Appanna, V. D. (2017). Reactive nitrogen species (RNS)-resistant microbes: adaptation and medical implications. *Biological Chemistry*, *398*(11), 1193–1208. https://doi.org/10.1515/hsz-2017-0152

Tjaden, B. (2020). A computational system for identifying operons based on RNA-seq data. *Methods*, *176*, 62–70. https://doi.org/10.1016/j.ymeth.2019.03.026

Tribelli, P. M., Lujan, A. M., Pardo, A., Ibarra, J. G., Fernández Do Porto, D., Smania, A., & López, N. I. (2019). Core regulon of the global anaerobic regulator Anr targets central metabolism functions in Pseudomonas species. *Scientific Reports*, *9*(1), 9065. https://doi.org/10.1038/s41598-019-45541-0

Tribelli, P. M., Rossi, L., Ricardi, M. M., Gomez-Lozano, M., Molin, S., Raiger Iustman, L. J., & Lopez, N. I. (2018). Microaerophilic alkane degradation in *Pseudomonas extremaustrali*s: a transcriptomic and physiological approach. *Journal of Industrial Microbiology & Biotechnology*, *45*(1), 15–23. https://doi.org/10.1007/s10295-017-1987-z

Tribelli, P. M., Solar Venero, E. C., Ricardi, M. M., Gómez-Lozano, M., Raiger Iustman, L. J., Molin, S., & López, N. I. (2015). Novel Essential Role of Ethanol Oxidation Genes at Low Temperature Revealed by Transcriptome Analysis in the Antarctic Bacterium Pseudomonas extremaustralis. *PloS One*, *10*(12), e0145353. https://doi.org/10.1371/journal.pone.0145353

Vargas-Blanco, D. A., & Shell, S. S. (2020). Regulation of mRNA Stability During Bacterial Stress Responses. *Frontiers in Microbiology*, *11*, 2111. https://doi.org/10.3389/fmicb.2020.02111

Vernis, L., El Banna, N., Baïlle, D., Hatem, E., Heneman, A., & Huang, M.-E. (2017). Fe-S Clusters Emerging as Targets of Therapeutic Drugs. *Oxidative Medicine and Cellular Longevity*, *2017*, 3647657. https://doi.org/10.1155/2017/3647657

Weber, M., & Fuchs, T. M. (2022). Metabolism in the Niche: a Large-Scale Genome-Based Survey Reveals Inositol Utilization To Be Widespread among Soil, Commensal, and Pathogenic Bacteria. *Microbiology Spectrum*, *10*(4), e0201322. https://doi.org/10.1128/spectrum.02013-22

Williams, D. E., & Boon, E. M. (2019). Towards Understanding the Molecular Basis of Nitric Oxide-Regulated Group Behaviors in Pathogenic Bacteria. *Journal of Innate Immunity*, *11*(3), 205–215. https://doi.org/10.1159/000494740

Winsor, G. L., Griffiths, E. J., Lo, R., Dhillon, B. K., Shay, J. A., & Brinkman, F. S. L. (2016). Enhanced annotations and features for comparing thousands of Pseudomonas genomes in the Pseudomonas genome database. *Nucleic Acids Research*, *44*(D1), D646-53. https://doi.org/10.1093/nar/gkv1227

Winterbourn, C. C. (1995). Toxicity of iron and hydrogen peroxide: the Fenton reaction. *Toxicology Letters*, *82–83*, 969–974. https://doi.org/10.1016/0378-4274(95)03532-x

Yebra, M. J., Zúñiga, M., Beaufils, S., Pérez-Martínez, G., Deutscher, J., & Monedero, V. (2007). Identification of a Gene Cluster Enabling *Lactobacillus casei* BL23 To Utilize *myo* -Inositol. *Applied and Environmental Microbiology*, *73*(12), 3850–3858. https://doi.org/10.1128/AEM.00243-07

Yuan, C., Yang, P., Wang, J., & Jiang, L. (2019). Myo-inositol utilization by Citrobacter koseri promotes brain infection. *Biochemical and Biophysical Research Communications*, *517*(3), 427–432. https://doi.org/10.1016/j.bbrc.2019.07.112

Zaffagnini, M., Bedhomme, M., Marchand, C. H., Morisse, S., Trost, P., & Lemaire, S. D. (2012). Redox Regulation in Photosynthetic Organisms: Focus on Glutathionylation. *Https://Home.Liebertpub.Com/Ars*, *16*(6), 567–586. https://doi.org/10.1089/ARS.2011.4255

Zhang, J., Li, X., Olmedo, M., Holdorf, A. D., Shang, Y., Artal-Sanz, M., Yilmaz, L. S., & Walhout, A. J. M. (2019). A Delicate Balance between Bacterial Iron and Reactive Oxygen Species Supports Optimal C. elegans Development. *Cell Host & Microbe*, *26*(3), 400-411.e3. https://doi.org/10.1016/j.chom.2019.07.010