Variability in radiocesium activity concentration in growing hardwood shoots in Fukushima, Japan

Hiroki Itô[1*], Satoru Miura[2], Masabumi Komatsu[2], Tsutomu Kanasashi[3], Junko Nagakura[2], Keizo Hirai[2]

1 Hokkaido Research Center, Forestry and Forest Products Research Institute, Sapporo, Japan

2 Forestry and Forest Products Research Institute, Tsukuba, Japan

3 Institute of Environmental Radioactivity, Fukushima University, Fukushima, Japan

* abies.firma@gmail.com

Abstract

The radiocesium contamination caused by the Fukushima Daiichi Nuclear Power Plant accident has made it difficult to use coppice woods as bed logs for mushroom cultivation. Evaluating the variability in the radiocesium activity concentration of logs is necessary in order to predict how many coppice woodlands are available for producing mushroom bed logs. To clarify the variability in radiocesium activity concentrations and to estimate the sample size required to estimate these concentrations with sufficient accuracy, we modeled the log-transformed radiocesium activity concentrations in growing shoots of hardwoods. We designed two models: (1) a model with mean concentrations that varied among stands with a standard deviation that was the same among stands, and (2) a model with varying means and standard deviations. We fit the data pertaining to only *Quercus serrata* to both models and calculated the widely applicable information criterion values. Consequently, we adopted the simpler model (1). Applying the selected model to data for all species, we examined the relationship between the number of measurement individuals and the predictive distribution of the expected concentration. Based on previous recommendations and measurement costs,

we proposed that five individuals would be appropriate for estimating radiocesium activity concentration in a stand.

Introduction

In Japan, coppice woodlands, consisting mainly of oaks, have long been managed for obtaining wood for fuel, charcoal, bed logs for mushroom cultivation, and for other purposes. Woodlands in Japan are a major component of what is referred to locally as "satoyama" landscape, which are areas that comprised of woodlands, grasslands, paddy fields, ponds, and related land use types [2, 3]. These areas, which are characterized by periodic human disturbance, have relatively high levels of biodiversity [1] However, although managed coppice woodlands can maintain biodiversity [4], large areas of them have been abandoned due to socio-economic changes [5].

More than any other areas in Japan, the traditional "satoyama" landscape is most prevalent in the southern Tohoku and northern Kanto regions of Japan. In Fukushima Prefecture, which is located in southern Tohoku, large area of coppice woodlands have been managed to produce the bed logs that are used for mushroom cultivation [6, 7]. For this purpose, oak *Quercus serrata* and other hardwood tree species have been cultivated in the coppice woodland. The production of these bed logs previously benefited forestry activities and the regional economy of the area, as well as supporting the biodiversity of the region.

However, the Fukushima Daiichi Nuclear Power Plant accident, which occurred in March 2011, polluted the area with radioactive materials. Most of the fallout consisted of the radiocesium isotopes, ^{134}Cs and ^{137}Cs. Though, as a result of its longer 30-year half-life, only ^{137}Cs remains in the environment. This contamination has had an enormous impact on the communities and industries in the region, including the production of bed logs for mushroom cultivation in the oak coppices

In order to meet the required safety standards limits for food, $100\,\mathrm{Bq\,kg^{-1}}$, bed logs used for mushroom cultivation must have a radioactivity concentration lower than $50\,\mathrm{Bq\,kg^{-1}}$ [8]. This is because the maximum transfer factor from the bed log to cultivated mushrooms has been estimated as being approximately two [7]. Resuming bed log production is considered to be necessary for the forestry and regional economy

and would also contribute towards maintaining the biodiversity of the region. 29

Recently, studies have been conducted to facilitate the resumption of coppice 30
management. For example, Kanasashi et al. [9] established a strong and negative 31
correlation between the exchangeable potassium present in soil and the radiocesium 32
(^{137}Cs) activity concentrations in the growing shoots of *Q. serrata*. The observation of a 33
constant relationship between radiocesium activity concentration of growing shoots 34
(new branches) and that in woody stem means that the concentration in growing shoots 35
are well suited for use as a proxy for the concentration in woody stems [10,11]. Based 36
on this assumption, coppices situated on the potassium-rich soils may be well suited for 37
the production of bed logs with low radiocesium contamination even in areas that were 38
experienced radioactive fallout. 39

The application of potassium fertilizer to reduce the absorption of radiocesium is a 40
widely used measure in agricultural fields [12]. In woodlands, which are considerably 41
larger than agricultural lands, searching for coppices with low radiocesium activity 42
concentrations could also be a promising measure for sourcing bed logs with 43
radiocesium activity concentrations that are within permissible limits. Indeed, for 44
mushroom cultivation on sawdust media, the relationship between mushroom fruiting 45
body and medium radiocesium activity concentrations have been clarified [13,14]. To 46
accurately predict the proportion of available logs, estimating not only the mean, but 47
also the variances in the radiocesium activity concentrations present in the coppice 48
stems is crucial. It has been shown that a definitive range exists in the radiocesium 49
activity concentrations of growing shoots within and among stands of coppices and 50
plant species [9]. However, this variance is seldom known a priori [15,16]. Moreover, if 51
the variance in the radiocesium activity concentrations varies among stands, then it will 52
be difficult to accurately predict the concentrations from small sample sizes. It is 53
therefore necessary to clarify the distribution of the variance in the radiocesium activity 54
concentration within and among coppice stands. In addition, the variance would be 55
important to determine the required sample size to predict the concentration. 56

Two questions arise regarding the measurement of radiocesium activity concentration 57
in growing shoots: (1) Can the variances of the measurement values of the radiocesium 58
activity concentration in growing shoots be regarded as constant or variable among 59
different stands? (2) What number of individual samples are required to measure the 60

stand-level distribution of radiocesium activity concentration? In this study, we ⁠ ₆₁
evaluated the stand-level variation in radiocesium activity concentrations in the growing ⁠ ₆₂
shoots of coppice species, and predicted the concentration distribution within a stand. ⁠ ₆₃
The findings will enable the estimation of the mean radiocesium activity concentration ⁠ ₆₄
in growing shoots within a specified area, thereby ensuring that the concentration ⁠ ₆₅
distribution is under a predetermined threshold. ⁠ ₆₆

Materials and Methods

Data collection

We established 40 study stands in Miyakoji, Tamura City, Fukushima Prefecture, Japan, ⁠ ₆₉
and sampled growing shoots of hardwood species in each stand between December 2016 ⁠ ₇₀
and March 2017. The radiocesium activity concentration is known to be stable during ⁠ ₇₁
the dormant stage (November to April) [17]. In this study, we assumed that the stands ⁠ ₇₂
comprised areas of several tens of square meters in size. Permission was obtained from ⁠ ₇₃
the Fukushima Chuo Forestry Co-operative to collect samples from the study area. The ⁠ ₇₄
Q. serrata data used in this analysis were essentially the same as those in Kanasashi et ⁠ ₇₅
al. [9], except that some measurements were updated. The data file is provided in the ⁠ ₇₆
supporting information (S1 File). In addition, other hardwood species that were present ⁠ ₇₇
in the stands with *Q. serrata*, as well as regenerated coppice sprouts and new ⁠ ₇₈
plantation, were also included in the analysis. A total of 453 individuals of hardwood ⁠ ₇₉
tree species were collected. Of these, 418 individuals of five species had radiocesium ⁠ ₈₀
activity concentrations that could be detected by the method described below were ⁠ ₈₁
included in the analysis (i.e., 35 individuals (7.7%) were excluded) (Table 1). ⁠ ₈₂

Table 1. Number of individuals of each species that were sampled

Species	No.
Quercus serrata Murray	184
Cerasus jamasakura (Siebold ex Koidz.) H.Ohba	111
Castanea crenata Siebold et Zucc.	86
Quercus acutissima Carruth.	30
Zelkova serrata (Thunb.) Makino	7

The collected samples were processed as described in a previous study [9], and the ⁠ ₈₃
radioactivity concentrations were measured using a NaI(Tl) scintillation detector (2480 ⁠ ₈₄

WIZARD^2; PerkinElmer Japan Co., Ltd., Yokohama, Japan) or high-purity germanium detectors (GEM40P4-76 and GEM-FX7025P4-ST; ORTEC, Oak Ridge, TN, USA). All of the detectors were calibrated using the radioactivity standard gamma volume sources, MX035 and MX033U8PP (Japan Radioisotope Association, Tokyo, Japan). For the NaI(Tl) scintillation detector measurements, samples were placed into 20-mL vials and measured from 2,000 to 172,800 s, after which the gamma-ray spectra were analyzed using the Code Fukushima software package [18]. The 662 keV peaks of Cs-137 were considered to be a distinct peak when the full width at half maximum (FWHM) was greater than 30 keV and the relative counting error was less than 33.3%; peaks meeting these criteria were used for the analysis. For the germanium semiconductor detector, measurements were performed from 1,800 to 86,400 s, and data with relative counting errors of 33.3% or less were used for analysis. The activity concentrations were decay-corrected to the values on December 1, 2016.

Statistical modeling

We assumed that the natural logarithm (hereafter written as logarithm) of the radiocesium activity concentration Y_i (Bq kg^{-1}) in the growing shoots followed a normal distribution [15, 19], namely

$$\log_e(Y_i) \sim \text{Normal}(\mu_{s[i]}, \sigma^2_{s[i]}), \tag{1}$$

where $\mu_{s[i]}$ denotes the mean, and $\sigma_{s[i]}$ denotes the standard deviation for stand $s[i]$ containing the i-th individual. We excluded censored data from the model analysis, as the detection limits of such data were indefinite and varied depending on the measurement time.

We further assumed that μ_s, the mean for stand s, was composed of the overall mean $\bar{\mu}$ and a random effect for the stand $\epsilon_{M[s]}$,

$$\mu_s = \bar{\mu} + \epsilon_{M[s]}$$
$$\epsilon_M \sim \text{Normal}(0, \sigma^2_M).$$

where σ_{M} denotes the standard deviation of ϵ_{M}.

We constructed two models predicated on distinct methodologies for formulating the standard deviation, denoted as σ_s, as outlined in Eq 1 (Table 2).

1. σ_s is constant for all stands, namely $\sigma_s = \bar{\sigma}$.

2. σ_s can vary due to a random stand effect, namely $\sigma_s = \bar{\sigma}\exp(\epsilon_{\mathrm{V}[s]})$.

The random stand effect ϵ_{V} was assumed to follow a normal distribution, and can be represented as follows:

$$\epsilon_{\mathrm{V}} \sim \mathrm{Normal}(0, \sigma_{\mathrm{V}}^2).$$

Table 2. Models to be compared

Model	Structure
Model 1	$\log_{\mathrm{e}}(Y_i) \sim \mathrm{Normal}(\mu_{s[i]}, \sigma_{s[i]}^2), \mu_s \sim \mathrm{Normal}(\bar{\mu}, \sigma_{\mathrm{M}}^2),$ $\sigma_s = \bar{\sigma}$
Model 2	$\log_{\mathrm{e}}(Y_i) \sim \mathrm{Normal}(\mu_{s[i]}, \sigma_{s[i]}^2), \mu_s \sim \mathrm{Normal}(\bar{\mu}, \sigma_{\mathrm{M}}^2),$ $\sigma_s = \bar{\sigma}\exp(\epsilon_{\mathrm{V}[s]}), \epsilon_{\mathrm{V}} \sim \mathrm{Normal}(0, \sigma_{\mathrm{V}}^2)$

Y_i: measured value of the radiocesium activity concentration in the growing shoots (Bq kg^{-1}) of the i-th individual, $\mu_{s[i]}$: mean of $\log_{\mathrm{e}}(Y_i)$ in the stand $s[i]$ containing the i-th individual, $\sigma_{s[i]}$: standard deviation of $\log_{\mathrm{e}}(Y_i)$ in the stand $s[i]$ (fixed in model 1), $\bar{\mu}$: overall mean of $\log_{\mathrm{e}}(Y_i)$ in the study area, $\bar{\sigma}$: overall standard deviation, ϵ_{V}: random stand effect on σ_s, σ_{M}: standard deviation of the random effect ϵ_{M}, σ_{V}: standard deviation of the random effect ϵ_{V}.

The prior distribution of the overall mean was defined as a vague prior as follows:

$$\bar{\mu} \sim \mathrm{Normal}(0, 100^2).$$

The priors of the standard deviations ($\bar{\sigma}$, σ_{M}, and σ_{S}) were defined as weakly informative priors [20, 21], as follows:

$$\sigma \sim \mathrm{HalfNormal}(10^2).$$

The parameters in the models were estimated using a Markov chain Monte Carlo [118]
(MCMC) approach. The MCMC calculation was conducted using Stan 2.32.2 [22] via [119]
the CmdStanR 0.5.3 interface [23] for R [24]. We obtained the MCMC samples from [120]
four independent Markov chains, each consisting of 8,000 MCMC iterations after a [121]
warmup stage comprising 8,000 iterations. We then evaluated the convergence using the [122]
value of $\hat{R}$, or the Gelman-Rubin statistics [25, 26] and a visual census of the degree of [123]
chain mixture. The Stan model code is included in the supporting information (S2 File, [124]
S3 File). [125]

In addition, we also calculated the widely applicable information criterion [126]
(WAIC) [27] values for each model. The WAIC values were calculated under the [127]
stipulation of forecasting the radiocesium activity concentration in growing shoots in a [128]
new stand. Analogous to the Akaike information criterion (AIC), models characterized [129]
by smaller WAIC values are anticipated to exhibit superior mean predictive [130]
performance in the context of of model selection. Moreover, WAIC can be applied to [131]
complex hierarchical Bayesian models. [132]

We fitted both models to the data containing only *Q. serrata* and selected the [133]
optimal model after consideration of WAIC values and model simplicity. [134]

Model for analyzing all data [135]

We then constructed a model to analyze the dataset comprising all five species that was [136]
based on the previously selected optimal model. In this model, we incorporated the [137]
origin of each individual (i.e., coppice sprout or plantation) as an explanatory variable, [138]
as well as an additional random species effect. The model selection affected whether the [139]
standard deviation, $\sigma_{s[i]}$ was uniform or varied across different stands. [140]

$$\log_e Y_i \sim \text{Normal}(\mu_{s[i]}, \sigma_{s[i]}^2) \tag{2}$$

$$\mu_s = \bar{\mu} + \beta O + \epsilon_M + \epsilon_S \tag{3}$$

$$\epsilon_M \sim \text{Normal}(0, \sigma_M^2) \tag{4}$$

$$\epsilon_S \sim \text{Normal}(0, \sigma_S^2). \tag{5}$$

where the variable O denotes the origin of the individual (0: sprout, 1: plantation), and [141]

β is the coefficient of O. ϵ_{M} is a random stand effect, and σ_{M} is the standard deviation   142

thereof. Similarly, ϵ_{S} denotes a random species effect, and σ_{S} is its standard deviation.   143

Although *Z. serrata* had a small sample size, incorporating the random species effect   144

facilitates estimates of individual species by "borrowing strength" [28].   145

For the parameter estimation, the model was fitted to the whole dataset (S1 File),   146

i.e. the data for all five species (40 stands and 418 individuals), using Stan 2.32.2. The   147

code for the Stan model is given in the supporting information S4 File. Parameters used   148

in the models are summarized in Table 3.   149

Table 3. List of parameters

Name	Description
$\bar{\mu}$	Overall mean of log-transformed radiocesium activity concentration
μ_s	Mean of log-transformed radiocesium activity concentration at stand s
β	Coefficient of the trunk origin (0: sprout, 1: plantation)
$\epsilon_{\mathrm{M}[s]}$	Random stand effect on the mean of log-transformed radiocesium activity concentration at stand s
$\epsilon_{\mathrm{V}[s]}$	Random stand effect on the standard deviation of log-transformed radiocesium activity concentration at stand s
ϵ_{S}	Random species effect on the mean of log-transformed radiocesium activity concentration
$\bar{\sigma}$	Overall standard deviation of log-transformed radiocesium activity concentration
σ_s	Standard deviation of log-transformed radiocesium activity concentration at stand s
σ_{M}	Standard deviation of the random stand effect ϵ_{M}
σ_{V}	Standard deviation of the random effect ϵ_{V}
σ_{S}	Standard deviation of the random species effect ϵ_{S}

Prediction interval of the measurements of radiocesium activity   150
concentration for a new stand   151

Using the selected model with all data, we conducted a simple simulation to predict the   152

radiocesium activity concentration in growing shoots in a new stand. In this simulation,   153

we calculated the relationship between the sample size and the predicted value of the   154

radiocesium activity concentration of growing shoots in the study area.   155

In addition, we estimated the relationship between the sample size and the standard   156

deviation. Suppose a situation in which the radiocesium activity concentration is   157

measured in n individuals from a new stand.   158

Let Y_{new} be the radiocesium activity concentration of the growing shoots in the new   159

stand, and $y_{\mathrm{new}} = \log_{\mathrm{e}}(Y_{\mathrm{new}})$. Namely,

$$y_{\mathrm{new}} \sim \mathrm{Normal}(\mu_s, \sigma_s^2). \tag{6}$$

The mean concentration μ_s follows a normal distribution with an overall mean of $\bar{\mu}$ and a variance of σ_{M}^2,

$$\mu_s \sim \mathrm{Normal}(\bar{\mu}, \sigma_{\mathrm{M}}^2). \tag{7}$$

As described above (Table 2), the standard deviation σ_s depends on the model. If model 1 is selected, then σ_s is a constant:

$$\sigma_s = \bar{\sigma}$$

Otherwise, if model 2 is selected, then σ_s varies between stands:

$$\sigma_s = \bar{\sigma}\exp(\epsilon_{\mathrm{V}}[s]).$$

From Eqs 6 and 7,

$$y_{\mathrm{new}} \sim \mathrm{Normal}(\bar{\mu}, \sigma_s^2 + \sigma_{\mathrm{M}}^2).$$

Given the measurements $Y_1, Y_2, \ldots, Y_n$ (Bq kg^{-1}), we denote their logarithm values as $y_i = \log_{\mathrm{e}}(Y_i)$, and y_{n+1} should be distributed as follows [29]:

$$y_{n+1} \sim \mathrm{Normal}(\mu_n, \sigma_s^2 + \sigma_n^2) \tag{8}$$

where the expected value of μ_n after n individuals were measured is given by

$$\mu_n = \frac{\frac{1}{\sigma_{\mathrm{M}}^2}\bar{\mu} + \frac{1}{\sigma_{\mathrm{s}}^2}\sum_{i=1}^{n} y_i}{\frac{1}{\sigma_{\mathrm{M}}^2} + \frac{n}{\sigma_{\mathrm{s}}^2}},$$

and the expected value of $\bar{\mu}_n$ is

$$\bar{\mu}_n = \frac{\frac{1}{\sigma_{\mathrm{M}}^2}\bar{\mu} + \frac{n}{\sigma_{\mathrm{s}}^2}\mu_s}{\frac{1}{\sigma_{\mathrm{M}}^2} + \frac{n}{\sigma_{\mathrm{s}}^2}}, \tag{9}$$

In addition, σ_n can be calculated as follows [29],

$$\frac{1}{\sigma_n^2} = \frac{1}{\sigma_{\mathrm{M}}^2} + \frac{n}{\sigma_s^2}.$$

We calculated the mean μ_n for n = 1, …, 10, under the assumption that the true value of the stand radiocesium activity concentration is 10 Bq kg^{-1}. We also calculated the standard deviation in Eq 8, $\sqrt{\sigma_s^2 + \sigma_n^2}$.

Using the above relationships, we inferred how many individuals are required to adequately evaluate the expected value and error of the radiocesium activity concentration in a new stand.

Results

Radiocesium activity concentration measurements

The radiocesium activity concentration in the growing shoots ranged from 6 to 10265 Bq kg^{-1}, and the mean and median were 519 and 221 Bq kg^{-1}, respectively. The stand-level mean of the log-transformed values of the radiocesium activity concentrations ranged from 1.7 to 8.8, and their mean and median was 5.1 and 5.3, respectively (Fig 1.) The standard deviation in the stands ranged from 0.33 to 1.09 and the mean was 0.70. It should be noted that 7.7% of the total individuals surveyed were excluded from the analysis for the average value, as their samples could not be quantified due to low radioactivity levels; for the standard deviation, the value might

therefore be larger if the excluded samples were included.

Fig 1. Stand-level mean (left) and standard deviation (right) of log-transformed values of radiocesium activity concentration measurements in the growing shoots obtained for each species. Values larger than the third quartile + 1.5 × interquartile range (IQR) or smaller than the first quartile - 1.5 × IQR were treated as outliers. *Zelkova serrata* is not shown because the species occurred in only two stands.

Model selection

The results of the MCMC calculations for *Quercus serrata* showed that $\hat{R}$ values were smaller than 1.1 for all of the parameters, and that the Markov chains appeared to be well mixed. We therefore considered that the Markov chains converged successfully to stable distributions.

Table 4 shows the result of the model selection. Since there was only a small difference in the WAIC values between the two models (499 and 498), and because the values might vary slightly due to the sampling error, we adopted the simpler model, Model 1. Consequently, the standard deviation could be perceived as uniform across the stands, thereby dismissing the variations intrinsic to individual stands in predicting the radiocesium activity concentrations.

Table 4. Comparison of models (posterior means of the parameters and WAIC)

Model	$\bar{\mu}$	$\bar{\sigma}$	σ_{M}	σ_{V}	WAIC
Model 1	5.62	0.74	1.09		499
Model 2	5.64	0.70	1.08	0.26	498

Parameter estimates for the whole dataset

Since the results of the model selection allowed us to treat the standard deviation σ_s as being uniform among the stands, we let $\sigma_s = \bar{\sigma}$ and estimated the parameters of the model (Eq 2–5) using MCMC.

Table 5 shows a summary of the posterior distributions for the parameters. The posterior mean of $\bar{\sigma}$, which is the standard deviation of radiocesium activity concentration in the growing shoots, was 0.74 on a logarithmic scale, and the standard deviation was 0.03. The value of 0.74 on the logarithmic scale translates to a geometric standard deviation of 2.09. Using this value, we can expect that approximately 95% of

the measurement values are within an 18-fold ($\exp(2 \times 1.96 \times 0.74) = 18$) on the usual   209

linear scale, excluding the effects of the species and origin. The posterior mean of the   210

coefficient β, which represents the origin of the individuals, was estimated as -0.39, with   211

the entire 95% confidence interval exhibiting negative values. The negative value   212

indicates that the individuals from plantations would have a lower radiocesium activity   213

concentration than the individuals produced from sprouts, and that the log-transformed   214

mean concentration would decrease from 4.92 to 4.53. The posterior mean of σ_M, which   215

is the standard deviation of the random stand effect on the log-transformed mean   216

concentration, was 1.33, and larger than that of σ_S, 0.62, which is the standard   217

deviation of the random species effect on the log-transformed mean concentration.   218

Table 5. Summary of the posterior distribution of parameters for all species in the model. SD: standard deviation.

Parameter	Mean	SD	2.5% quantile	97.5% quantile
$\bar{\mu}$	4.92	0.41	4.06	5.65
β	-0.39	0.11	-0.61	-0.17
$\bar{\sigma}$	0.74	0.03	0.69	0.79
σ_M	1.32	0.17	1.04	1.71
σ_S	0.62	0.44	0.19	1.76

Fig 2 shows the posterior distributions of the random species effect. *Q. serrata* had a   219

relatively high value while that of *Zelkova serrata* was lower, corresponding the   220

log-transformed mean (Fig. 1). Values in the figure are shown using a logarithmic scale;   221

consequently, -1 on this scale is equivalent to a 0.37-fold decrease compared to the   222

overall mean radiocesium activity concentration of the growing shoots on a linear scale.   223

In the same manner, -0.5 on this scale is equivalent to a 0.61-fold decrease, while 0.25   224

and 0.5 are equivalent to increase of 1.6- and 2.7-fold, respectively.   225

Fig 2. Posterior distribution of the random species effect (on a logarithmic scale). Dots denote medians, thick lines denote 90% credible intervals, and thin lines denote 95% credible intervals.

Fig 3 shows the relationship between the number of measurement individuals (n)   226

and the expected value of the log-transformed radiocesium activity concentration   227

($\log_e(\mu_n)$), assuming a true mean of $10\,\mathrm{Bq\,kg^{-1}}$. This figure omits the random species   228

effect and assumes the origin of the stems to be sprouts. It shows that the expected   229

value progressively approximates the true value as the number of measurements   230

increases. Further, this relationship holds for other true values. ²³¹

Fig 3. Relationship between the number of measurement individuals and the expected value of the log-transformed radiocesium activity concentration when the true mean is $10\,\mathrm{Bq\,kg^{-1}}$ ($\log_\mathrm{e} 10 = 2.30$). The curve denotes the mean, and the dark region denotes the 95% prediction interval of the expected value.

As a result, the standard deviation of the predicted radiocesium activity ²³²
concentration on a logarithmic scale within a stand is given as ²³³

$$\sqrt{\sigma_s^2 + \sigma_n^2} = \sqrt{\sigma_s^2 + \frac{1}{\frac{1}{\sigma_\mathrm{M}^2} + \frac{n}{\sigma_s^2}}}$$
$$= \sqrt{\bar{\sigma}^2 + \frac{1}{\frac{1}{\sigma_\mathrm{M}^2} + \frac{n}{\bar{\sigma}^2}}}. \tag{10}$$

Discussion ²³⁴

In this study, we employed a Bayesian approach using MCMC methods to estimate ²³⁵
parameters in the mixed-effect models. Using WAIC values facilitated the selection of a ²³⁶
model for predicting radioactivity concentrations in the developing shoots of coppice ²³⁷
hardwoods. In addition, the Bayesian approach facilitated straightforward management ²³⁸
of uncertainties [29]. ²³⁹

Variation in radioactivity concentrations within a stand ²⁴⁰

Within the context of food safety, the public has a strong interest in radiocesium ²⁴¹
contamination of agricultural and forest products [7]. Local government bodies measure ²⁴²
the radioactivity concentrations of foods and items utilized for food production, such as ²⁴³
mushroom bed logs in the forestry sector, to prevent the distribution of foods and items ²⁴⁴
that have radioactivity concentrations that exceed standard limits. Consequently, ²⁴⁵
understanding the frequency distribution characteristics of the radioactivity ²⁴⁶
concentration in forest products obtained under certain conditions is very important for ²⁴⁷
the effective management and monitoring of these resources. In this study, we ²⁴⁸
investigated the variability in radiocesium activity concentrations in growing shoots of ²⁴⁹
hardwood species, which has been utilized to establish indices of radiocesium ²⁵⁰
absorption [10, 11], with a particular focus on the hardwood bed logs that are commonly ²⁵¹

used in mushroom cultivation. The radiocesium activity concentration in the forest ₂₅₂ products has been demonstrated to follow a log-normal distribution [15,19]. After the ₂₅₃ two largest nuclear power plant accidents to date, Chornobyl and Fukushima, ₂₅₄ radioactivity concentrations in many agricultural and forest products have been ₂₅₅ measured, and transfer factors (TF) and aggregated transfer factors (Tag) have been ₂₅₆ reported [30–32]. However, few studies have examined the variability — characterized ₂₅₇ by variance or standard deviation — in radioactivity concentrations of vegetation ₂₅₈ samples after log-transformation to attain normality. Khomutinin et al. [19,33] analyzed ₂₅₉ large systematic grid sampling data, with variations in the dimensions of sampling units ₂₆₀ and increments (distances between the grid nodes). In their research on vegetation, they ₂₆₁ found that sampling independence in the radiocesium activity concentration could be ₂₆₂ achieved by sampling over distance of 8–10 m in stands with uniform soil ₂₆₃ contamination [15]. Their findings were reflected in the sampling guidelines ₂₆₄ recommended and published by the International Atomic Energy Agency (IAEA) [16]. ₂₆₅ Khomutinin et al. [19] also reported that the log-transformed standard deviation of ₂₆₆ radioactivity concentration by ^{137}Cs sampled from 13 sites in Ukraine ranged from 0.22 ₂₆₇ to 0.62, and the mean and standard deviation was 0.38±0.11 . Their sampling sites ₂₆₈ covered a variety of different land use types including cultivated lands. ₂₆₉

Compared to the study in Ukraine, in this study, we estimated the standard ₂₇₀ deviation within 40 individual survey stands, focusing on the growing shoots (N=5) of ₂₇₁ five hardwood species of forests in Fukushima. The range in the radiocesium activity ₂₇₂ concentrations showed a marked discrepancy between soil and growing shoots; while the ₂₇₃ range in the soils was relatively stable with a 10-fold difference, that in the growing ₂₇₄ shoots showed fluctuations that ranged up to 1000-fold [9]. Based on the WAIC values ₂₇₅ (Table 4), we could select the simpler model with a constant standard deviation, instead ₂₇₆ of the model with a varying standard deviation among stands. Thus, we could treat the ₂₇₇ variability of radiocesium contamination of forest products as uniform irrespective of ₂₇₈ stands, as shown in crops and grasses [19]. The standard deviation of log-transformed ₂₇₉ concentration within each stand ($\bar{\sigma}$) was estimated to be 0.74 as the posterior mean, ₂₈₀ and its standard deviation was estimated to be 0.03 (Table 5). ₂₈₁

However, the within-stand standard deviation observed in this study and the study ₂₈₂ in Ukraine (0.38) [19] showed a two-fold difference. In Japan, woodlands are typically ₂₈₃

situated on steep undulating slopes, and the topographical characteristics of an area can change markedly within a relatively small area. We consider that these variations in the terrain may affect the soil properties. Furthermore, in terms of sampling methods, soil type, land usage, plant species, etc., numerous differences exist in the dataset used in the present study. Consequently, the factor that caused the two-fold difference between the present study and the preceding study in Ukraine was not specified. The standard deviations in the radioactivity concentration in vegetation samples are fundamental data that are essential for establishing radiation protection standards. Further studies are therefore required to clarify the principal factors that influence representative values and to comprehensively understand the prevalence of associated uncertainties.

In this study in Fukushima, we used sampling areas spanning several tens of square meters. To achieve statistical independence in sampling, Barnekow et al. [16] proposed that the center-to-center distance of sampling points for vegetation samples should not be less than 8–10 m. We consider that our sampling protocol satisfied this criterion in most cases. In contrast, in the survey of Khomutinin et al. [19], sampling grids ranged from 1 to 10 m in grid size, with 7 of the 13 surveyed sites having a grid size of 2.5 m or less. We considered the possibility that differences in sampling distance (increments) may have affected the differences observed in standard deviations, but the standard deviation in Khomutinin et al. [19] did not appear to be affected by the difference in grid size. All of the study sites in Khomutinin et al. [19] were located in crop fields and former agricultural fields and did not include forests; this is a notable difference between the datasets used in their study and this study. It is possible that forests show a greater variability of radiocesium activity concentrations in vegetation samples. For example, our Fukushima survey was conducted in mountainous areas with slopes of 6 to 30 degrees (11 to 73% gradient) [9], wheres all. 13 study sites in their Ukraine survey were located on flat terrain (confirmed on Google Maps from latitude and longitude). These differences may have accounted for the differences observed in the variability of radiocesium activity concentrations between these studies.

Factors affecting the radioactivity concentration

The species random effect was larger in *Q. serrata*, which is most commonly used for bed logs; the posterior mean was 0.44. The random effect for *Q. acutissima*, another

major hardwood species that is used for mushroom logs, was lower than that for *Q.* 315

serrata (Fig 2); the posterior mean was -0.14. The difference in the posterior mean of 316

the random effect was 0.57, which corresponds to a 1.8-fold difference when converted to 317

a linear scale. The lowest value was observed in *Z. serrata*; the posterior mean was -0.55 318

or 0.37-fold of that of *Q. serrata* when converted to a linear scale. The difference 319

between the maximum and minimum mean values for the five tree species was 2.7-fold. 320

Ipatyev et al. [34] reported a minimum 10-fold difference in Tag values ($m^2 \, kg^{-1}$) for 321

needles and leaves of five tree species in Belarus, with a range from 1.6×10^{-3} for birch 322

to 3.8×10^{-2} for pine. However, Calmon et al. [35] examined the Tag of trees after the 323

Chornobyl nuclear accident and found that the influence of tree species exerted only a 324

minor influence on these values, affecting them by a factor of 2 to 3. The present study 325

also suggested that the interspecific difference in radiocesium absorption from the soil 326

was small. 327

Furthermore, the data suggest a disparity in radiocesium contamination levels 328

contingent on the origin of the trunk, with stems that originated from sprouts 329

exhibiting a greater degree of contamination by radiocesium than stems that originated 330

from plantations. In this study, the ages of the measured stems ranged from one year to 331

a maximum of seven years. Therefore, the differences in radiocesium absorption among 332

species and the origin of the stems may differ from those of stems aged around 20 years 333

old, which are best suited for use as bed logs. Long-term monitoring is therefore 334

considered necessary in order to asses whether the observed relationships remain 335

consistent into the future. 336

The stand-level mean radiocesium concentrations observed in this study ranged from 337

1.7–8.8 in log-transformed values (Fig. 1). However, since 7.7% of the total number of 338

individuals were excluded due to their low radioactivity levels, it is plausible the true 339

distribution lies more towards the left tail. 340

Kanasashi et al. [9] pointed out that the amount of exchangeable potassium in the 341

surface soil largely influenced the radiocesium concentrations in the growing shoots. In 342

comparison, although species and origin differences also affect radiocesium 343

concentrations in the growing shoots, their impact is markedly diminished compaired to 344

the growing environment. These findings are consistent with a review [35], where 345

post-Chornobyl nuclear accident analysis of Tag values for trees showed a pronounced 346

influence of soil type, quantified as a level of 100, whereas the effect of tree species was 2. Given this large variability in the mean values, it is important to investigate the radiocesium activity concentrations across stand to determine if the forest owners can resume the bed log production for mushrooms.

Required sample size for appropriate estimation of radiocesium activity concentration

The objective of this study, evaluating the radiocesium activity concentrations of growing shoots in the coppices, is to protect communities from radiation exposure. This is particularly pertinent given the regulations that have been imposed on forest products used in food production, such as mushroom bed logs [7]. Assuming a normal distribution for the log-transformed radioactivity concentration estimates, the two parameters, i.e., the mean and standard deviation, are required to facilitate the accurate estimation of the proportion of the bed logs produced in a stand, which would adhere to the stipulated radioactivity regulation thresholds. Importantly, these parameter estimates have uncertainties. In this study, we showed that the within-stand standard deviation of the log-transformed radioactivity concentration of growing shoots of coppice woods could be regarded as uniform among stands and that the mean value was 0.74 (Table 5). The sample mean value of the log-transformed radioactivity concentration of growing shoots has been reported to vary markedly among stands [9]. Fig. 3 shows the relationship between the number of measurement individuals (n) in a stand and the expected value of the log-transformed radiocesium activity concentration for the case where true mean is $10\,\mathrm{Bq\,kg^{-1}}$ (2.30 in the log-transformed value). When the number of measurements is limited, the expected value approaches toward the overall mean of 4.92; however, with an increase in the data points, the curve progressively approximates the true value, and the slope becomes gentler. Thus, the expected value would be 2.92 in the case where $n = 1$, 2.46 in the case where $n = 5$, and 2.38 in the case where $n = 10$ (Eq 9, Fig. 3). Therefore, as the number of measurement individuals increases, the standard deviation of the predicted radiocesium activity concentration (log-transformed) within a stand diminishes; the standard deviation would be 0.98 in the case where $n = 1$, 0.80 in the case where $n = 5$, and 0.77 in the case where $n = 10$ (Eq 10). While the rate of decrease in the values slows down, the

measurement cost increases in proportion to the number of individuals measured. The ₃₇₈ IAEA recommends sampling at least five trees to investigate the radioactivity of the ₃₇₉ natural environments contaminated by nuclear accidents [36]. Khomutinin et al. [19] ₃₈₀ suggested that five-point samples taken from an area 1 m^2 were required in order to ₃₈₁ determine the radionuclide concentration in a combined vegetation sample. ₃₈₂ Consequently, we propose that sampling five individuals would be sufficient to estimate ₃₈₃ the radiocesium activity concentration in a coppice stand. ₃₈₄

Conclusion

In predicting radiocesium activity concentration in the growing shoots of hardwood species in forests exposed to radioactive fallout, the variance of the log-transformed measurement values within a stand can be assumed to be invariant irrespective of the stands. In this study, we found the mean standard deviation to be 0.74, a value that remained constant despite variations in the mean value of radiocesium activity concentration across stands and species. Furthermore, the findings showed that measurements from approximately five individuals would be sufficient for estimating the mean radiocesium activity concentration in a stand. The variance and mean of vegetation samples are fundamental parameters used in radiation protection measures, such as setting standard limits for radiation protection and determination surveys based on reference values. Numerous vegetation surveys have been conducted in forests, agricultural lands, and grasslands after the Fukushima and Chornobyl nuclear accidents. The statistical description and understanding of radioactive contamination in terrestrial plant ecosystems through integrated analysis of these data is an important topic that required further work.

Supporting information

S1 File   Data file of the radioactivity concentration activity in the growing shoots of five hardwood species in Miyakoji.

S2 File.   Stan model code file for model 1.

S3 File. Stan model code file for model 2. ₄₀₅

S4 File. Stan model code file for the model with all species. ₄₀₆

Acknowledgments ₄₀₇

The authors thank the Fukushima Chuo Forestry Co-operative for supporting this study. ₄₀₈
The authors thank Mr. Kentaro Matsuura (HOXO-M Inc.) for assistance with ₄₀₉
developing the Stan model incorporating the WAIC calculations. The authors also ₄₁₀
thank FORTE Science Communications (`https://www.forte-science.co.jp/`) for ₄₁₁
English language editing. This research was supported by the research program on ₄₁₂
development of innovative technology grants from the Project of the Bio-oriented ₄₁₃
Technology Research Advancement Institution (BRAIN) (Grant Number: JPJ007097) ₄₁₄
and grants from the Forestry and Forest Products Research Institute (201901, 202203). ₄₁₅

References

1. Washitani I. Satoyama landscapes and conservation ecology. In: Takeuchi K,
   Brown RD, Washitani I, Tsunekawa A, Yokohari M, editors. Satoyama — the
   traditional rural landscape of Japan —. Tokyo: Springer-Verlag Tokyo; 2003. p.
   16–23.

2. Fukamachi K, Oku H, Nakashizuka T. The change of a satoyama landscape and
   its causality in Kamiseya, Kyoto Prefecture, Japan between 1970 and 1995.
   Landsc Ecol. 2001;16:703–717. doi:10.1023/A:1014464909698.

3. Takeuchi K. Satoyama landscapes as managed nature. In: Takeuchi K, Brown
   RD, Washitani I, Tsunekawa A, Yokohari M, editors. Satoyama — the traditional
   rural landscape of Japan —. Tokyo: Springer-Verlag Tokyo; 2003. p. 9–16.

4. Itô H, Hino T, Sakuma D. Species abundance in floor vegetation of managed
   coppice and abandoned forest. For Ecol Manage. 2012;269:99–105.
   doi:10.1016/j.foreco.2011.12.017.

5. Morimoto Y. What is satoyama? Points for discussion on its future direction. Landsc Ecol Eng. 2011;7:163–171. doi:10.1007/s11355-010-0120-5.

6. Miura S. The effects of radioactive contamination on the forestry industry and commercial mushroom-log production in Fukushima, Japan. In: Nakanishi TM, Tanoi K, editors. Agricultural Implications of the Fukushima Nuclear Accident: The First Three Years. Tokyo: Springer Japan; 2016. p. 145–160.

7. Hashimoto S, Komatsu M, Miura S. Impacts of radioactive contamination of forest on life. In: Hashimoto S, Komatsu M, Miura S, editors. Forest Radioecology in Fukushima: Radiocesium Dynamics, Impact, and Future. Springer; 2022. p. 75–126.

8. Forestry Agency of Japan. Revision of the present index for mushroom bed logs; 2012 [cited 2024-07-04]. Available from: `https://www.rinya.maff.go.jp/j/tokuyou/attach/pdf/shihyouti-index-4.pdf` [in Japanese].

9. Kanasashi T, Miura S, Hirai K, Nagakura J, Itô H. Relationship between the activity concentration of ^{137}Cs in the growing shoots of *Quercus serrata* and soil ^{137}Cs, exchangeable cations, and pH in Fukushima, Japan. J Environ Radioact. 2020;220-221:106276. doi:10.1016/j.jenvrad.2020.106276.

10. Kobayashi NI, Ito R, Masumori M. Radiocesium contamination in forests and the current situation of growing oak trees for mushroom logs. In: Nakanishi TM, O'Brien M, Tanoi K, editors. Agricultural Implications of the Fukushima Nuclear Accident (III): After 7 Years. Singapore: Springer Singapore; 2019. p. 107–122.

11. Miura S. Studies on the revitalization of radioactive-contaminated mushroom log forests: Focus on shoots. In: Nakanishi TM, Tanoi K, editors. Agricultural Implications of the Fukushima Nuclear Accident (IV). Singapore: Springer Singapore; 2023. p. 203–220. doi:10.1007/978-981-19-9361-9_18.

12. Kato N, Kihou N, Fujimura S, Ikeba M, Miyazaki N, Saito Y, et al. Potassium fertilizer and other materials as countermeasures to reduce radiocesium levels in rice: results of urgent experiments in 2011 responding to the Fukushima Daiichi

Nuclear Power Plant accident. Soil Sci Plant Nutr. 2015;61(2):179–190. doi:10.1080/00380768.2014.995584.

13. Hiraide M, Sunagawa M, Neda H, Humaira'Lau bt Abdullah N, Yoshida S. Reducing radioactive cesium transfer from sawdust media to *Pleurotus ostreatus* fruiting bodies. J Wood Sci. 2015;61(4):420–430. doi:10.1007/s10086-015-1483-x

14. Hiraide M. Factors affecting the cesium transfer factor to shiitake (*Lentinula edodes*) cultivated in sawdust medium. J Wood Sci. 2021;67(1):17. doi:10.1186/s10086-021-01949-9

15. Khomutinin YV, Kashparov VA, Zhebrovska KI. Sampling optimization when radioecological monitoring. Kiev: Ukraine Institute for Agricultural Radiology; 2004.

16. Barnekow U, Fesenko S, Kashparov V, Kis-Benedek G, Matisoff G, Onda Y, et al. Guidelines on soil and vegetation sampling for radiological monitoring. Vienna: International Atomic Energy Agency; 2019.

17. Sakashita W, Miura S, Nagakura J, Kanasashi T, Shinomiya Y. Seasonal stability of ^{137}Cs in coppiced *Quercus serrata* current-year branches: toward the estimation of trunk ^{137}Cs activity concentrations without felling. Ecol Indic. 2021;133:108361. doi:10.1016/j.ecolind.2021.108361.

18. Yamada T, Takano N. Doublet peak area determination in NaI(Tl) scintillation spectrometry using maximum likelihood estimation. Appl Radiat Isot. 2014;87:407–409. doi:10.1016/j.apradiso.2013.11.069.

19. Khomutinin Y, Fesenko S, Levchuk S, Holiaka D, Kashparov V. Optimising sampling strategies for emergency response: vegetation sampling. J Environ Radioact. 2021;233:106605. doi:10.1016/j.jenvrad.2021.106605.

20. Gelman A, Jakulin A, Pittau MG, Su YS. A weakly informative default prior distribution for logistic and other regression models. Ann Appl Stat. 2008;2(4):1360–1383.

21. Gelman A, Simpson D, Betancourt M. The prior can often only be understood in the context of the likelihood. Entropy. 2017;19:555. doi:10.3390/e19100555.

22. Carpenter B, Gelman A, Hoffman M, Lee D, Goodrich B, Betancourt M, et al. Stan: a probabilistic programming language. J Stat Softw. 2017;76(1):1–32. doi:10.18637/jss.v076.i01.

23. Gabry J, Češnovar R. cmdstanr: R interface to 'CmdStan'; 2022 [cited 2023-07-04]. Available from: `https://mc-stan.org/cmdstanr/`.

24. R Core Team. R: a language and environment for statistical computing; 2022 [cited 2023-07-04]. Available from: `https://www.R-project.org/`.

25. Gelman A, Rubin DB. Inference from iterative simulation using multiple sequences. Stat Sci. 1992;7(4):457–472.

26. Brooks SP, Gelman A. General methods for monitoring convergence of iterative simulations. J Comput Graph Stat. 1998;7(4):434–455. doi:10.1080/10618600.1998.10474787.

27. Watanabe S. Asymptotic equivalence of Bayes cross validation and widely applicable information criterion in singular learning theory. J Mach Learn Res. 2010;11:3571–3594.

28. Kéry M, Schaub M. Bayesian population analysis using WinBUGS: a hierarchical perspective. Waltham: Academic Press; 2011.

29. Gelman A, Carlin JB, Stern HS, Dunson DB, Vehtari A, Rubin DB. Bayesian data analysis. 3rd ed. Boca Raton: CRC Press; 2013.

30. International Atomic Energy Agency. Quantification of radionuclide transfer in terrestrial and freshwater environments for radiological assessments. IAEA-TecDoc-1616. TECDOC Series. Vienna: International Atomic Energy Agency; 2009.

31. International Atomic Energy Agency. Handbook of parameter values for the prediction of radionuclide transfer in terrestrial and freshwater environments. No. 472 in Technical Reports Series. Vienna: International Atomic Energy Agency; 2010.

32. International Atomic Energy Agency. Environmental transfer of radionuclides in Japan following the accident at the Fukushima Daiichi Nuclear Power Plant. No. 1927 in TECDOC Series. Vienna: International Atomic Energy Agency; 2020.

33. Khomutinin Y, Fesenko S, Levchuk S, Zhebrovska K, Kashparov V. Optimising sampling strategies for emergency response: Soil sampling. J Environ Radioact. 2020;222:106344. doi:10.1016/j.jenvrad.2020.106344.

34. Ipatyev V, Bulavik I, Baginsky V, Goncharenko G, Dvornik A. Forest and Chernobyl: forest ecosystems after the Chernobyl nuclear power plant accident: 1986–1994. J Environ Radioact. 1999;42(1):9–38. doi:10.1016/S0265-931X(98)00042-3.

35. Calmon P, Thiry Y, Zibold G, Rantavaara A, Fesenko S. Transfer parameter values in temperate forest ecosystems: a review. J Environ Radioact. 2009;100(9):757–766. doi:10.1016/j.jenvrad.2008.11.005.

36. Barnekow U, Fesenko S, Kashparov V, Kis-Benedek G, Matisoff G, Onda Y, et al. Guidelines on soil and vegetation sampling for radiological monitoring. International Atomic Energy Agency; 2019.

Figure 1

Figure 2

Figure 3