SARS-CoV-2 antibodies cross-react and enhance dengue infection

Kamini Jakhar[†], Sudipta Sonar[†], Gagandeep Singh, Tania Sarkar, Mahima Tiwari, Jaskaran

Kaur, Deepak Kumar Rathore, Banwari Lal, Sandeep Kumar, Puneet Srivastav, Satendra

Kumar, Vikas Phagna, Lokesh Kumar, Vishal Gupta, Pallavi Kshetrapal, Savita Singh, Nitya

Wadhwa, Ramachandran Thiruvengadam, Sreevatsan Raghavan, Mudita Gosain, Tripti

Shrivastava, Sankar Bhattacharyya, Jayanta Bhattacharya, Shailendra Asthana and Shailendra

Mani[*]

† Contributed equally

*Corresponding author

Corresponding author: Dr Shailendra Mani PhD, Translational Health Science and Technology

Institute, Faridabad, NCR Delhi, India +91-129-2876478

Email: shailendramani@thsti.res.in

Author affiliations:

Translational Health Science and Technology Institute, Faridabad, Haryana, India-120001

Word Count

Abstract- 197

Main text- 3490

Abstract

Dengue disease is highly prevalent in tropical and subtropical regions worldwide. However, its

pathogenesis is still incompletely understood, particularly in comparison to other endemic

viruses. Antibody-dependent enhancement (ADE) is a well-known phenomenon for dengue

viruses. Given the recent surge in dengue cases and potential cross-reactivity with SARS-CoV-

2 antibodies, this study explores the impact of anti-SARS-CoV-2 antibodies on DENV-2

infection.

The study assessed the cross-reactivity of SARS-CoV-2 antibodies with the DENV-2 Virus.

Human convalescent plasma samples collected during different waves of COVID-19 and

monoclonal and polyclonal antibodies raised against SARS-CoV-2 were examined for their

potential to cause ADE of DENV-2 infection using cell-based assays. The study found that anti-

SARS-CoV-2 antibodies acquired from natural infection in humans or through experimental

immunization in animals were cross-reactive with DENV-2 and had the potential to enhance

DENV-2 infection in K562 and U937 cells. In-silico and in-vitro studies indicated a strong

interaction between SARS-CoV-2 antibodies and DENV-2 E-protein, providing a molecular

basis for these findings. This study is the first to demonstrate that anti-SARS-CoV-2 antibodies

can cross-react with DENV-2 and can enhance its infection through ADE. These findings have

implications for SARS-CoV-2 vaccine development and deployment strategies in regions

where dengue is endemic.

Summary:

Antibodies against SARS-CoV-2 (RBD and Spike) showed significant cross reactivity with

DENV-2 (E protein). Also, anti-SARS-CoV-2- commercial antibodies, immunised animal sera

and 46 human convalescent plasma samples (from different waves of pandemic) demonstrated

antibody-dependent enhancement (ADE) of DENV-2 infection.

Keywords; SARS-CoV-2; Cross-reactivity; Dengue; ADE; Vaccine-development.

Introduction

Dengue virus (DENV) is a positive-sense single-stranded RNA virus with a genome size of ~11kb belonging to the *Flaviviridae* family and causes the most prevalent mosquito-borne disease called dengue. Dengue is endemic to more than 120 countries with Asia contributing 70% of the global burden. The COVID-19 pandemic has had a substantial impact on Dengue cases globally with many countries reporting significant a surge since 2020. In 2021, prevalence of dengue has increased over three times in India (44,585 versus 193,245 cases), 19 times in Bangladesh (1,405 versus 28,429 cases), 2 times in Singapore (5,000 versus 11,000), over 1.8 times in Malaysia (87,000 versus 120,000 cases), and more than seven times (6,016 vs. 52,894 instances) in Pakistan. Taiwan, Sri Lanka, Mexico, and the Philippines have also reported notable increase, while Brazil documented a surge in severe cases and fatalities predominantly impacting younger population under 20 years of age during the period of 2021-2022 [[1], [2], [3], [4]]. DENV circulates as four serotypes (DENV 1-4) each containing multiple distinct genotypes. Infection with any serotype can lead to mild fever or severe forms like Dengue Haemorrhagic Fever (DHF) and Dengue Shock Syndrome (DSS). Primary infection provides lifelong immunity to homotypic secondary infection but gives partial immunity from heterotypic challenge. Such heterotypic secondary infections positively correlate with severe symptoms, as a consequence of higher viremia through Antibody-dependent Enhancement (ADE). ADE is caused by sub-neutralizing antibodies from previous infection binding to DENV virion and promoting FcR-mediated viral entry into the cells bearing Fc-receptor [5]. ADE occurs when the number of bound antibodies support stable attachment to Fc receptors, without reaching the neutralization threshold [6].

Presently there are multiple reports showing modest serological cross-reactivity between SARS-CoV-2 and DENV-2, significantly impacting dengue diagnostics [7] [8] [9]. However, it is yet unclear if this would have any clinical impact on dengue cases [10]. A study reported

IgG produced in rabbits upon immunization with purified SARS-CoV-2 S1-RBD, to cross-react

with DENV proteins, particularly the Envelope (E), precursor-Membrane (PrM), and Non-

structural protein 1 (NS1). However, these antibodies did not enhance DENV infection (ADE)

in THP-1 cells which express FcR[11]. Contrarily a study showed *in-vitro* inhibition

of dengue infection by sera from convalescent COVID-19 patients sera.

In this pretext, our current study aimed to examine if SARS-CoV-2 antibodies have any impact

on DENV-2 infection. To the best of our knowledge, this will be the first study to provide a

blueprint as to how SARS-CoV-2 antibodies cross-reactive with DENV-2 (particularly E

protein) may enhance dengue infection via ADE, using a combination of experimental and

computational studies.

Methods:

A detailed material and methods section is given as supplementary data. In brief, we selected a

group of convalescent plasma samples (n=48) before the Delta wave collected in India that is

15th of April 2021 (n=21), post the delta wave (n=17) and during Omicron wave (n=10). To

check if the SARS-CoV-2 antibodies enhance dengue virus infection in-vitro, human myeloid

cell lines K562 and U937 were used for the study. Along with the patient samples, a panel of

commercially available antibodies and animal sera raised against SARS-CoV-2

(Supplementary Table-2) were used. In ADE assay, the 4G2 antibody was used as a positive-

control while Rabishield®-100 was used as a Negative-control. ADE assay was performed to

obtain the percentage of DENV-2 positive cells by flowcytometry. The cross-reactivity of

SARS-CoV-2 antibodies with DENV protein E was tested using in-silico approach where

fourteen SARS-CoV-2 antibodies mainly targeting SARS-CoV-2 RBD were chosen. They were

docked onto the DENV-2 E-protein using multiple docking programs to choose the best pose.

The binding free energy of protein and antibody complexes was calculated in Prime MM-GBSA

by Prime module of Schrodinger suite. Biolayer Interferometry (BLI) was performed to study

the binding kinetics of two SARS-CoV-2 antibodies with dengue virus envelope and SARS-

CoV-2 spike protein. The cross-reactivity was also checked by immunofluorescence assay

where C6/36 cells were infected with DENV2 and stained with antibodies and animal sera

raised against SARS-CoV-2. Experimental values were represented as mean ± standard

deviation (SD). Statistical analysis was done using the ordinary one-way ANOVA followed by

Dunnett's multiple comparison test. Differences were considered significant at p value<0·05.

Results:

Antibodies against SARS-CoV-2 cross react with Dengue Virus

Previous studies have suggested considerable serological cross-reactivity between SARS-CoV-

2 and DENV. We employed multiple techniques to strengthen such observations. Using

confocal microscopy, we observed that the anti-spike polyclonal antibody (P1R) and anti-spike

mice serum (MSer) showed a significant cross-reactivity with an average MFI of 12·8 (p-

value<0·0001) and 13·4 (p-value<0·0001) (Figure-1A-B) with respect to the Negative-control,

Rabishield® whereas, the SARS-CoV-2 monoclonals (1A9 and MM41) also showed cross-

reactivity with DENV-2. The positive-control, 4G2 (pan-flavivirus-anti-envelope antibody)

showed maximum MFI of 18·28 whereas the Negative-control, anti-Rabies monoclonal

(Rabishield®) showed no reactivity with DENV-2. The binding kinetics of 1A9 (SARS-CoV-

2 spike antibody) in BLI showed high affinity with a dissociation constant (K_D) of 4·84 nM and

22·7 nM with DENV-2 Envelope and SARS-CoV-2 Spike respectively. (Figure-1C ii-iii). C8

(DENV-2 Envelope antibody) was used as a positive-control with K_D value of 3·24 nM (Figure-

1C (i)).

SARS-CoV-2 convalescent plasma enhances Dengue virus infection *in-vitro*

Probing the surface expression of FcγR-I, II and III in K562 and U937 cell lines showed both

cell lines to predominantly express FcγR-II (Supplementary Figure-1), as reported previously

[12]. When samples from convalescent COVID-19 patients (n=48, Supplementary Table-1)

were tested for their ability to cause ADE of DENV-infection in K562 cells, a significant

augmentation was observed (Figure-2). The percentage of DENV-2 infected K562 cells

(Supplementary Figure-2) was increased by 14 out of 21 samples from the pre-delta wave (p-

value<0·0001), 15 out 17 samples from the delta wave (p-value<0·0001) and 8 out of 10

samples from the Omicron wave (p-value=0·0001), in comparison with untreated virus control

(Figure-2A-D). Furthermore, using a subset of the convalescent plasma samples (n=15) similar

results were observed in U937 cells, where 9 out 15 serum samples enhanced DENV-2

infection. (p-value=0·0001) (Supplementary Figure-3A-B). A quantification of the neutralizing

antibody titers against SARS-CoV-2 Wuhan strain in these samples by live-virus neutralization

assay (Supplementary Table-2), did not show any association between the neutralizing antibody

titer and ADE efficiency (Supplementary Figure-4A-C). Similarly, we did not find any

association with severity of the COVID-19 patients (Supplementary Figure-4D).

A quantitative-PCR for intracellular DENV genomic RNA normalized to host cell GAPDH

transcripts (Supplementary Table-3), showed that 7 out 10 COVID-19 convalescent patient

samples caused higher viral replication in K562 cells when compared to control, probably

through increasing virus uptake in ADE (p-value<0·0001) (Supplementary Figure-5). This data

correlates with the increase in DENV-2 positive cells observed in flowcytometry.

Determination of the infectious titer of virus produced from infected cells by focus-forming

unit assay also showed a significant increase in the Foci-forming units/ml (FFU/ml) in 2 out of

3 COVID-19 convalescent plasma samples as compared to the virus control (p-value<0·0001)

(Supplementary figure-6).

Monoclonal and polyclonal antibodies raised against SARS-CoV-2 enhance dengue virus infection:

Since DENV is endemic to India it is possible that pre-existing anti-DENV antibodies are responsible for the observation of ADE as described above. In order to rule this possibility out, we tested commercially available monoclonal and polyclonal antibodies and sera raised against the SARS-CoV-2 spike and RBD for their potential to enhance the DENV-2 infection in ADE assay using K562 cells (Supplementary Table-4). Interestingly, all of these antibodies enhanced DENV-2 infection (p-value=0·0001) (Figure-3) confirming that antibodies binding to SARS-CoV-2 surface protein cross-react with DENV cause ADE. The RBD specific antibody (M5B) showed stronger enhancement compared to antibody against spike (MM41) with dose-dependent increase in DENV-2 infection (Figure-3A-B). Similarly, serum from Hamsters immunized with purified SARS-CoV-2 spike protein also showed ADE of dengue virus infection in K562 cells (Figure-3C).

As a control monoclonal antibody against Rabies virus, which do not cross react with DENV, (Negative-control), did not show any ADE of dengue infection in K562 cells suggesting a specific effect of anti-SARS-CoV-2 antibodies (Figure-3C). A similar trend of enhancement in viral infection was observed in U937 cells too using these antibodies (Supplementary figure-7). Three monoclonal antibodies namely 4G2, 1A9 and M4B showed significant ADE, as determined by infectious titer of virus released in the culture supernatant from K562 cells infected in the presence of these antibodies (p-value<0·0001) (Supplementary figure-6).

Having established that there is an enhancement of DENV-2 infection due to anti-SARS-CoV-2 antibodies, we next investigated whether it is time-dependent. We assessed the enhancement at 6 h, 18 h and 36 h post infection by virus and virus-antibody complexes, using FACS (Supplementary figure-8). We observed that the mice (Mser) and hamster (Hser) sera, showed a significant enhancement in percentage DENV-2 positive cells with respect to virus control

(adjusted p-values Mser=0·0071 and Hser=0·0001), even at 6 hpi, Although, we didn't observe

a significant enhancement due to 4G2 and anti-Spike monoclonals M3S and M2G at 6 hpi. At

18 hpi, there was a significant increase in the DENV-2 positive cells with respect to virus

control in 4G2 (Adjusted p-value=0·0024) as well as M2G (Adjusted p-value=0·0101), which

further enhanced significantly at 36 hpi (Adjusted p-values 4G2=0·0001 and M2G=0·0035).

The Hser showed highest increase in DENV-2 positive cells at 6 h which remained almost

constant till 36 h. (Adjusted p-value<0·0001).

Cross recognition of DENV-E protein by anti-Spike and anti-RBD SARS-CoV-2

antibodies *in-silico*

In parallel, in order to assess the cross-reactivity of SARS-CoV-2 antibodies with DENV-2 E-

protein dimer (Figure-4A) and to underpin the residues critically involved, we considered an

in-silico approach (Figure-4B). Since the structures for the antibodies used for *in-vitro* assays

weren't available, we selected a panel of reported SARS-CoV-2 antibodies, based on their

structural (extent of their characterization, availability of the structure information of their co-

crystals with low resolution, etc.) and biological (neutralization potential against SARS-CoV-

2, target epitopes, etc.) relevance (Supplementary Table-5).

For the identification of epitopes of DENV-2 E-protein, based on interaction fingerprinting, an

antibody against E-protein dimer i.e. C8 (Fab region, PDB: 4UTA) was used to characterize the

interaction site using its co-crystal. For the selected SARS-CoV-2 antibodies, in order to rule

out any biases, the protein-protein (PP) blind docking was carried out with the whole DENV-2

E dimer, for identification of the cross-reactive antibodies forming most promising complexes

with the E-protein. The consensus docked pose between Piper and HDOCK was used for post-

docking qualitative and quantitative analysis such as pose selection and pose-filtering criteria.

The top three SARS-CoV-2 antibodies which formed the best complexes with DENV-2

evaluated in terms of docking energy, interaction fingerprinting with co-crystal and MM-GBSA

scores are CR3022 (Fab region, PDB:6W7Y), S2E12 ab (Fab region, PDB:7K45) and DMAb

2196 (Fab region, PDB:8D8R) (Figure-4C). These complexes were picked for further analysis.

For the C8 antibody (4UTA) binding with DENV-2 E-protein, the residues in the interacting

epitopes were, T68, T70, E84, R99, G102, G104, K247, Q248, S274, K310 and E311 (formed

hydrogen bonds and/or salt-bridges); E71, S72, R73, C74, N103 and T155 (formed

hydrophobic contacts and Vander Waal interactions etc.), and were majorly localized in the

Domain-II (Figure-4D). The binding sites of CR3022, S2E12 ab and DMAb 2196 antibodies

were found to overlap nicely with the C8 binding sites, as it revealed well by the binding site

analysis (Figure-4D).

Furthermore, to quantitate their binding association at the residue level, the major contributing

amino acids were underscored in the form of residue-wise decomposition analysis using the

cut-off value of -1·5 kcal/mol. The contributing residues for CR3022 were E71, S72, R73, Q77,

S81, Q86 and N153 (Majorly in Domain-II); for S2E12ab, were S72, R73, Q77, S81, N83, R99,

N103, G152, D154 and T155 (Majorly in Domain-II including Fusion loop); and for DMAb

2196 the residues S72, R73, C74, Q77, E79, N83, E84, R99, and E311 (Majorly in Domain-II

including Fusion loop) were involved (Figure-4E, F). From residue-wise mapping it was

interesting to note that there were few overlapping residues in the interaction of E-protein with

C8 and the top three SARS-CoV-2 interacting antibodies [13]. The residue R73 appeared to be

the most common residue among all antibodies. Further in the pair-wise interaction of SARS-

CoV-2 and DENV-2 antibodies, the common residues between C8 and CR3022 were T70, E71,

S72, G102 and N103; between C8 and DMAb2196, were T70, S72, C74 and R99; and between

C8 and S2E12 ab were E71, R99, N103, T155 and K247 (Figure-4E and 4F). Furthermore, we

also performed the mining of paratope residues of SARS-CoV-2 antibodies which have shown

common interaction with DENV-2 E with equal and/or increased interaction energy. The

paratope residues involved in the interaction of DENV-2-E residues as well with CR3022 are

Y27(H), Q65(H), S100(H), G101(H) and E111 (L); with DMAb2196 residues are Y32(L),

V52(H), S54(H), Y91(L), S93(L), I100(H) and C102(H); and with S2E12, the residues are

V2(H), V3(L), L4 (H), L48(L), S53(L), I59(L), G65(L), V93(L), Q102(L) and C106(H)

(Supplementary Figure-9).

Overall, the interaction maps and per-residue energetic decompositions, underpinned that there

were several overlapping residues involved in the interaction of DENV-2-E with C8 and

CR3022, S2E12 ab and DMAb 2196 (Figure-4G and 4H). The epitope and paratope analysis

followed by thermodynamic energy (ΔG) calculations clearly indicated a high chance that

SARS-CoV-2 antibodies possibly interact with DENV-2 E-protein (Figure-4G and 4H). The

most cross-reactive residues involved in the interacting epitopes of C8, CR3022, S2E12 ab and

DMAb 2196 were also elucidated (Supplementary Table-6)

Cross reactivity and enhancement of DENV-2 infection due to CR3022 antibody

Based on the computational outcomes, one of the SARS-CoV-2 antibodies, CR3022 was further

investigated *in-vitro* for its cross-reactivity and ability to cause ADE. Confocal microscopy

images of CR3022 showed a statistically significant cross-reactivity with an average MFI of

7·68 (p-value<0·0001) w.r.t. the Negative-control, Rabishield® (Figure-5A). In BLI, it showed

high affinity with a dissociation constant (K_D) of <1·0pM with Spike protein (Figure-5B (i)),

and 2·83 nM with DENV-2 E-protein (Figure-5B (ii)). We observed a significant augmentation

in the percentage of infected cells at 2 µg and 4 µg concentrations of antibody (p-value=0·01)

in comparison with untreated virus control (Figure-5C).

Discussion:

In this study, we demonstrated that SARS-CoV-2 antibodies and plasma cross-react with

DENV-2. Also, SARS-CoV-2 spike, and RBD-directed antibodies have a very strong binding

affinity with stabilized DENV-2 E-protein dimer *in-vitro* with the dissociation constant in nM

range) as well as *in-silico*. We used the DENV-2 E-protein for BLI and computational studies

since it has been reported to have maximum cross-reactivity with SARS-CoV-2 S1 specific

antibodies [7]. Additionally, it is critical for viral entry into cells and is the major target not

only for neutralizing antibodies but also for infection-enhancing antibodies [14]. We did the

docking studies using dimeric DENV-2 E-protein since we intended to study the interaction

of DENV-2 with SARS-CoV-2 antibodies, which would happen before the viral entry,

extracellularly and it's known that the E-protein exists as a head-to-tail oriented homodimer

on the surface of mature-virions [15].

We further assessed whether the SARS-CoV-2 antibodies could enhance DENV-2 infection *in-*

vitro. We found that convalescent plasma samples of qRT-PCR confirmed SARS-CoV-2

infected individuals, from the pre-delta, delta and post-delta waves, significantly enhanced the

DENV-2 infection in cultured Fc-receptor positive K562 and U937 cells. Also, the

commercially available anti-Spike and anti-RBD SARS-CoV-2 antibodies and Spike-

immunized mice and hamster sera showed an increased DENV-2 infection in both the model

systems.

Our docking results indicated that there is a significant interaction of SARS-CoV-2 antibodies

with DENV-2 dimeric E-protein with the binding free energy of three of them, CR3022

(6W7Y), S2E12 ab (7K45) and DMAb2196 (8D8R), being comparable to the DENV-2 dimer

specific antibody, C8 (4UTA). These top three binders shared their interacting interface with

C8 antibody. The residues involved in the interaction of C8 antibody with DENV-2 E that were

also involved in the interaction of at-least one of the top three binders with E-protein were: R73,

E84, R99, G102 and E311 (Localized in Domain-II and Fusion-loop). The contributing residues

which were found to be majorly stabilizing the interaction of these top three SARS-CoV-2

antibodies with E-protein were S72, R73, Q77, S81, N83 and R99 (involved in interacting

epitopes of at least two out of the three). All the residues involved in the interaction of these

three SARS-CoV-2 antibodies with E-protein when mapped, were majorly found to lie in the

Domain-II and fusion-loop of E-protein. It is well known that the antibodies against two

epitopes of DENV are strong mediators of ADE: prM epitopes[16] and fusion loop epitopes

[17]. The fusion loop epitopes could be the ones, potentially involved in ADE. Multiple residues

have been identified from *in-silico* interaction mining, among them few residues such as R73

(common among all), T70, S72 (Domain-II), R99 and N103 (Fusion loop) and E311 (Domain-

III) seems critical as they are contributing with DENV-2 (C8:4UTA) and at least two antibodies

of SARS-CoV-2 CR3022:6W7Y; S2E12ab:7K45; and DMAb 2196:8D8R; as well. These

residues put forth the possible avenue to explore further for epitope-based therapeutic

interventions.

There are several limitations to our study. First, our study doesn't provide the mechanistic

insights into the pathway involved in ADE of DENV-2 infection due to SARS-CoV-2

antibodies. The extrinsic pathway, involves an enhanced FcR-mediated entry of the virus into

the host cells leading to an increase in the number of infected cells, whereas the intrinsic

pathway, involves a suppression of the host antiviral immune response causing increased viral

replication and would boost virus production per infected cell (burst size) [18] [19]. Although,

here we briefly did a time-course based kinetic study of DENV-2 infection upon treatment of

virus with SARS-CoV-2 antibodies and convalescent plasma. We observed an enhancement in

the percentage of DENV-2 infected cells, even at 6 h, indicating enhanced virus internalization,

since this is before E-protein from replicated new virions is expected [14] [15] which may

possibly indicate the presence of at least extrinsic pathway for ADE. However, it needs a lot of

further exploration for confirmation.

Secondly, the DENV-2 used for the assay was prepared in C6/36 cells and wasn't characterized

for the proportions of immature and mature-virions. It is well known that the virus prepared *in-*

vitro in mosquito cells has considerably a greater amount of immature virions as compared to

mammalian cells, probably due to differential furin-expression [20] [21] . Also, it is recognized

that laboratory adapted DENV strains are more dynamic than clinical strains and expose cryptic

epitopes more easily [22]. Thus our virus preparation might have major proportion of immature

virions but this probably won't change the fact that the ADE of DENV-infection would happen

since the viral heterogeneity in response to the natural DENV-infections is also well known

The antibody repertoire in the natural infections is known to have an abundance of poorly

neutralizing antibodies which are majorly generated in response to the immature conformations

[23], which is a result of incomplete furin processing, resulting in egress of immature-forms

along with and mature-forms of virions post-assembly [24] [25]. Also, the fully mature-virions

have been reported to be less susceptible to neutralization by cross-reactive and heterotypic

immune sera than partially mature variants [23]**.** Thus, the heterogeneity in the DENV-2

preparation used for the *in-vitro* assays can be probably closely related to the natural infection

scenario. Further, our results from the BLI and docking studies are based on the dimeric form

of E-protein (indicating mature-form). Thus, one may then ask that the cross-reactive epitopes

defined here, might not be all the interacting epitopes that would possibly be involved and

interactions with other conformations must also be considered. Although it can be challenging

to map the exact cross-reactive epitopes since DENV is known to have multiple transient states,

involving different arrangements and oligomeric states of E-proteins that which governs the

antigenic landscape or accessibility of DENV [26]. Still, a docking study with a heterotrimer of

E-protein (consisting of un-cleaved prM), being a crystallized transient state [27] may give a

better indication of more of the cross-reactive epitopes involved.

In summary, our findings demonstrated that the SARS-CoV-2 antibodies can enhance the

DENV-2 infection. A recent report by Nath *et. al.* demonstrated serological cross-reactivity of

SARS-CoV-2 positive serum samples with DENV1 and that the sera neutralized DENV1 [28]

but contrastingly, we didn't observe neutralization of DENV-2 at any antibody concentration

used. Also Panting *et. al.*, in a recent retrospective matched cohort study of pediatric patients,

diagnosed with SARS-CoV-2 and dengue coinfection, in Philippines, reported that the co-

infection predominantly presented with typical dengue symptoms more case fatality rate in

coinfection than mono-infection (6·7% versus 5%) [29], which could possibly be due to an

ADE of DENV-infection by SARS-CoV-2 antibodies. Further, it would be interesting to

investigate if the antibodies induced in response to various SARS-CoV-2 vaccines also possess

the potential to enhance DENV-infection. This finding can have significant implications

keeping in view the co-circulation of both DENV-2 and SARS-CoV-2 in several countries and

thus the vaccine development strategies against both the agents might need an inevitable

change. The findings suggest the need to consider DENV seasonality when deploying SARS-

CoV-2 vaccines to prevent severe infections, particularly in countries where DENV is endemic.

Finally, the study also identifies the cross-reactive epitopes and paratopes and lists some critical

residues involved in the interaction of SARS-CoV-2 antibodies with DENV-2 E-protein, which

can have prophylactic and therapeutic implications.

Acknowledgements:

The following reagent was obtained through BEI Resources, NIAID, NIH: SARS-Related

Coronavirus 2, Isolate USA-WA1/2020, NR-52281, Monoclonal Anti-SARS-Related

Coronavirus 2 Spike Glycoprotein, Clone 1-3D7 (produced *in-vitro*), NR-56488, Monoclonal

Anti-SARS-Related Coronavirus 2 Spike Glycoprotein Receptor Binding Domain-(RBD),

Clone 2TP1C3 (produced *in-vitro*), NR-55296, Polyclonal Anti-SARS-Related Coronavirus 2

Spike Glycoprotein (IgG, Rabbit), NR-52947." The RBD is proprietary reagents with IP No.

202011018845, we thank Dr. Barney Graham (VRC/NIAID/NIH) for providing us with spike

construct (SARS-2-CoV S 2P). We would like to thank Prof. Shinjini Bhatnagar, Prof. Pramod

Garg from THSTI for critically reviewing the manuscript, ABSL3 facility committee and staff

for their help and support. This research has been conducted with the significant contribution

and expertise of THSTI, NCR Biotech Science Cluster BIOREPOSITORY and DBT India

Consortium".

DBT India Consortium**, ** Name of all the collaborating Institutes/ Hospitals: Translational

Health Science Technology Institute, Clinical partners: Maulana Azad Medical College, Lok

Nayak Jai Prakash Hospital, and Lady Hardinge Medical College in Delhi, ESI Medical College

Hospital, Faridabad, Civil Hospital, Gurugram, Haryana; Civil Hospital, Palwal, Haryana; Al-

Falah School of Medical Science & Research Centre and Hospital, Dhauj, Haryana; Medanta

Hospital, Gurugram; Shaheed Hasan Khan Mewati Government Medical College, Nalhar,

Haryana.

Data availability:

All the data are presented in this manuscript. The virus strains used in this study have the

following accession numbers: original Wuhan (USA-WA1/2020, Genbank: MN985325),

B·1·617·2 (δ-variant) known as THSTI_287 (GenBank: MZ356566·1), and Omicron BA·1

(GISAID accession no.: EPI_ISL_8764350) Source data are provided with this paper.

Author Contributions:

SM conceptualized the idea. SM, SB, SA and TrS coordinated the research activity. KJ and SS

contributed to designing and execution of viral assays. KJ, SS, GS, TS, MT, JK, DKR, BL, SK,

PS, SK, VP, LK and VG contributed to data curation, investigation and methodology. KJ, SS,

GS, SB, SA, TrS and SM contributed to formal analysis. SS, SR, MG, PK, NW, RT, SB, JB,

TrS, SA and SM provided resources. KJ, SS, TrS, PK, NW, RT, SB, JB, SA and SM contributed

to writing, review & editing the draft. All authors had full access to all the data in the study and

had final responsibility for the decision to submit for publication.

Ethics statement

We obtained ethical approval from the Institutional Bio-Safety Committee of THSTI prior to

conducting all studies: (IBSC#495/2023) and RCGM (No.-BT/IBKP/137/2020).

Disclosure statement

The Author declares no conflict of interest.

Funding:

The current research is funded by the Translational Research Program Grant,

BT/PR30159/MED/15/188/2018 and the Bill and Melinda Gates Foundation GIISER-South

Asia (INV-030592). The computational work was supported by the national network project

(NNP-BT/PR40189/BTIS/137/50/2022), DBT. The work on dengue envelope proteins was

done with support of funding received from NBM (National Biopharma Mission) scheme of

BIRAC (Biotechnology Industry Research Assistance Council), India (BT/NBM099/02/18)

(2019-2023). SK acknowledge NBM BIRAC for his fellowship support.

Reference:

1. Khan S, Akbar SMF, Yahiro T, et al. Dengue Infections during COVID-19 Period: Reflection of Reality or Elusive Data Due to Effect of Pandemic. Int J Environ Res Public Health **2022**; 19.

2. Lu X, Bambrick H, Pongsumpun P, Dhewantara PW, Toan DTT, Hu W. Dengue outbreaks in the COVID-19 era: Alarm raised for Asia. PLoS Negl Trop Dis **2021**; 15:e0009778.

3. Sarkar S. Asia's "twindemic": dengue fever cases rocket as covid-19 rages on. BMJ **2022**; 378:o2090.

4. Manna S, Satapathy P, Bora I, Padhi BK. Dengue outbreaks in South Asia amid Covid-19: Epidemiology, transmission, and mitigation strategies. Front Public Health **2022**; 10:1060043.

5. Lee MF, Voon GZ, Lim HX, Chua ML, Poh CL. Innate and adaptive immune evasion by dengue virus. Front Cell Infect Microbiol **2022**; 12:1004608.

6. Pierson TC, Diamond MS. A game of numbers: the stoichiometry of antibody-mediated neutralization of flavivirus infection. Prog Mol Biol Transl Sci **2015**; 129:141-66.

7. Masyeni S, Santoso MS, Widyaningsih PD, et al. Serological cross-reaction and coinfection of dengue and COVID-19 in Asia: Experience from Indonesia. Int J Infect Dis **2021**; 102:152-4.

8. Alla D, Alla SSM, Vempati R, et al. Dengue & COVID-19: A Comparison and the Challenges at Hand. Cureus **2022**; 14:e31877.

9. Digwo DC, Elebe CP, Chigor VN, Madueke SN, Ezeh CK, Ike AC. Occurrence of False-Positive Tests and Cross-reactions Between COVID-19 and Dengue With Implications During Diagnosis: A Mixed Evidence Synthesis. Infectious Microbes & Diseases **2023**; 5:64-75.

10. Ulrich H, Pillat MM, Tarnok A. Dengue Fever, COVID-19 (SARS-CoV-2), and Antibody-Dependent Enhancement (ADE): A Perspective. Cytometry A **2020**; 97:662-7.

11. Cheng YL, Chao CH, Lai YC, et al. Antibodies against the SARS-CoV-2 S1-RBD cross-react

with dengue virus and hinder dengue pathogenesis. Front Immunol **2022**; 13:941923.

12. Littaua R, Kurane I, Ennis FA. Human IgG Fc receptor II mediates antibody-dependent

enhancement of dengue virus infection. J Immunol **1990**; 144:3183-6.

13. Srivastava M, Suri C, Singh M, Mathur R, Asthana S. Molecular dynamics simulation

reveals the possible druggable hot-spots of USP7. Oncotarget **2018**; 9:34289-305.

14. Wahala WM, Silva AM. The human antibody response to dengue virus infection. Viruses

2011; 3:2374-95.

15. Modis Y, Ogata S, Clements D, Harrison SC. A ligand-binding pocket in the dengue virus

envelope glycoprotein. Proc Natl Acad Sci U S A **2003**; 100:6986-91.

16. Rodenhuis-Zybert IA, van der Schaar HM, da Silva Voorham JM, et al. Immature dengue

virus: a veiled pathogen? PLoS Pathog **2010**; 6:e1000718.

17. de Alwis R, Williams KL, Schmid MA, et al. Dengue viruses are enhanced by distinct

populations of serotype cross-reactive antibodies in human immune sera. PLoS Pathog **2014**;

10:e1004386.

18. Modhiran N, Kalayanarooj S, Ubol S. Subversion of innate defenses by the interplay

between DENV and pre-existing enhancing antibodies: TLRs signaling collapse. PLoS Negl

Trop Dis **2010**; 4:e924.

19. Ubol S, Phuklia W, Kalayanarooj S, Modhiran N. Mechanisms of immune evasion induced

by a complex of dengue virus and preexisting enhancing antibodies. J Infect Dis **2010**;

201:923-35.

20. Murray JM, Aaskov JG, Wright PJ. Processing of the dengue virus type 2 proteins prM and

C-prM. J Gen Virol **1993**; 74 (Pt 2):175-82.

21. Davis CW, Nguyen HY, Hanna SL, Sanchez MD, Doms RW, Pierson TC. West Nile virus

discriminates between DC-SIGN and DC-SIGNR for cellular attachment and infection. J Virol

2006; 80:1290-301.

22. Chaichana P, Okabayashi T, Puiprom O, et al. Low levels of antibody-dependent

enhancement in vitro using viruses and plasma from dengue patients. PLoS One **2014**;

9:e92173.

23. Beltramello M, Williams KL, Simmons CP, et al. The human immune response to Dengue

virus is dominated by highly cross-reactive antibodies endowed with neutralizing and

enhancing activity. Cell Host Microbe **2010**; 8:271-83.

24. Pierson TC, Diamond MS. Degrees of maturity: the complex structure and biology of

flaviviruses. Curr Opin Virol **2012**; 2:168-75.

25. Cherrier MV, Kaufmann B, Nybakken GE, et al. Structural basis for the preferential

recognition of immature flaviviruses by a fusion-loop antibody. EMBO J **2009**; 28:3269-76.

26. Holzmann H, Stiasny K, York H, Dorner F, Kunz C, Heinz FX. Tick-borne encephalitis virus

envelope protein E-specific monoclonal antibodies for the study of low pH-induced

conformational changes and immature virions. Arch Virol **1995**; 140:213-21.

27. Rey FA, Stiasny K, Vaney MC, Dellarole M, Heinz FX. The bright and the dark side of

human antibody responses to flaviviruses: lessons for vaccine design. EMBO Rep **2018**;

19:206-24.

28. Nath H, Mallick A, Roy S, et al. COVID-19 serum can be cross-reactive and neutralizing

against the dengue virus, as observed by the dengue virus neutralization test. Int J Infect Dis

2022; 122:576-84.

29. Pantig FMT, Clemens SAC, Clemens R, Maramba-Lazarte CC, Madrid MAC. SARS-CoV-2

and Dengue Coinfection in Filipino Children: Epidemiology Profile, Clinical Presentation and

Outcomes. Pediatr Infect Dis J **2023**.

Figure 1: Cross-reactivity of antibodies and sera raised against SARS-CoV-2 Spike with DENV-2 (NGC) proteins.

(A) Confocal, digital images of DENV-2 (NGC) infected C6/36 cells reacted with 4G2, antibodies and sera raised against SARS-CoV-2 after 48 hpi. Cell nuclei are stained with DAPI while the other antibodies are detected using FITC-labeled respective secondary antibodies.

(B) Bar graph showing MFI of 4G2, MSer, 1A9, P1R (normalized w.r.t. the respective secondary antibody controls) with standard deviation. Analysis was performed using ImageJ and statistical significance was carried out using One-way ANOVA followed by Dunnett's test in GraphPad Prism 8·4·2. Asterisk (*) indicates statistically significant difference in MFI w.r.t. Rabishield® (Negative control).P value = 0·1234 (ns), 0·0332 (*), 0·0021 (**), 0·0002 (***), <0·0001 (****)

(C) Binding kinetics of C8 (DENV-E protein specific) and 1A9 (SARS-CoV-2 spike specific) antibodies with DENV-2 protein E and SARS-CoV-2 spike protein. Antibodies were loaded onto the sensors and serial dilutions of the proteins were used to study binding kinetics. The binding responses were calculated by subtracting data from reference and fitting globally with a 1:1 binding model using ForteBio's Data Analysis software 10·1. The data was considered validated if $\chi 2$ <0·5.

Figure 2: ADE of dengue virus infection by the SARS-CoV-2 positive patients' sera in K562 cells.
ADE due to convalescent plasma samples collected in the (A) pre-delta wave, n=21, (B) delta wave, n=17, and (C) Omicron wave, n=10. The bar graph represents an average percentage increase in DENV positive cells w.r.t. VC for each sample with standard deviation for duplicates. Statistical significance was carried out using One-way ANOVA followed by Dunnett's test in GraphPad Prism 8·4·2. Asterisk (*) indicates that the difference between the Virus control and the serum samples is statistically significant. P value = 0·1234 (ns), 0·0332 (*), 0·0021 (**), 0·0002 (***), <0·0001 (****)
(D) Representative dot plots of patient samples from Pre –delta wave (#143, #155), Delta wave (#96), and Omicron wave (#2, #7). Data was analyzed on FlowJo version 10·8·1.

Figure 3: ADE of dengue virus infection by the monoclonal and polyclonal antibodies and serum against SARS-CoV-2.

ADE due to (A) anti-spike monoclonal antibodies, M1B, 1A9, and different volumes of MM41 (from 2 µl to 6 µl) and anti-spike polyclonal antibody raised in rabbit, P1R, (B) anti-RBD monoclonal antibodies, M4B, and different volumes of M5B (from 2 µl to 6 µl), (C) hamster serum immunized with SARS-CoV-2 spike and negative control (Rabishield®). Statistical significance was calculated using One-way ANOVA followed by Dunnett's test in GraphPad Prism 8·4·2. Asterisk (*) indicates that the difference between the Virus control and the serum samples is statistically significant. P value = 0·1234 (ns), 0·0332 (*), 0·0021 (**), 0·0002 (***), <0·0001 (****).

(D) Representative dot plots. Data was analyzed using FlowJo version 10·8·1.

Figure 4. Computational insights exploring the cross-reactive SARS-CoV-2 antibodies at DENV-2 E protein: (A) Sequence and structural organization of the DENV-2 E dimer (PDB 1OAN). (B) Overview of the computational pipeline. (C) The binding free energy in kcal/mol of selected SARS-CoV-2 antibodies. The most likely complexes were generated using Protein-protein dockings. (D) The interaction map of top three antibodies (CR3022: 6W7Y: violet, S2E12 neutralizing antibody Fab fragment: 7K45: orange and DMAb 2196: 8D8R: magenta) along with the native antibody of DENV-2 E as control (C8:4UTA: green). The residue-wise zoom-out view is shown to highlight the critical residues mainly forming the interactions such as Hbonds and SBs. The DENV-2 antibody interactions are shown in green-colored residues and SARS-CoV-2 antibody interactions are highlighted in the pink-colored residues. (E) The matrix of critical residues involved in either establishing either H-bonds or in SBs along with their occurrence. (F) The per-residue decomposition of top three antibodies indicating the thermodynamic contributions of epitope residues. (G) The net binding energy between identified antibodies and DENV-2 E which claims as potential in both cases. (H) The 3D positioning of identified epitope residues at dimer of DENV. The residues which only interact with SARS-CoV-2 antibodies are shown in cyan, the residues interacting with only DENV-2 antibody are shown in green color. The residues which are interacting with both DENV-2 E as well as with SARS-CoV-2 RBD are shown in violet color. All the residues are rendered in licorice and atom-wise C:cyan/green/violet, N:blue, O:red and S:yellow.

Figure 5: Cross-reactivity and ADE of Dengue virus infection by the SARS-CoV-2 anti-RBD antibody CR3022.
(A) Confocal, digital images of DENV-2 (NGC) infected C6/36 cells stained with CR3022 along with the controls, 4G2 and Rabishield® (taken from Figure 1 to show the comparison) and the Mean Fluorescence Intensity (MFI).
(B) Binding kinetics of CR3022 with SARS-CoV-2 spike protein (i) and DENV-2 protein E (ii).
(C) ADE of DENV-2 infection by different concentrations of CR3022 in K562 cells by flowcytometry. Statistical significance was checked using One-way ANOVA followed by Dunnett's test. Asterisk (*) indicates a statistically significant difference between the control and treatment. P value = 0·1234 (ns), 0·0332 (*), 0·0021 (**), 0·0002 (***), <0·0001 (****).