Title: Foveolar drusen decrease fixation stability in pre-symptomatic AMD.

Authors: J. Murari[1], J. Gautier[2], J. Daout[1], L. Krafft[3], P. Senée[3,4], P. Mecê[3,5], K.Grieve[1,2], W. Seiple[6], *D.

Sheynikhovich[1], S. Meimon[3], M. Paques[2], A. Arleo[1]

Conflict of interest: No conflicting relationship relevant for this manuscript exists for any author.

Running title: Foveolar drusen decrease fixation stability

This article contains two video clips as additional online-only material: *Clip 1* and *Clip 2*.

This article contains additional online-only figures: *Supplemental Figure 1* and *Supplemental Figure 2*.

Word count (abstract and text): 6284

Abbreviations:

FEM: fixational eye movements

AMD: age-related macular degeneration

RPE: retinal pigment epithelium

AO-FIO: adaptive optics flood illumination ophthalmoscope

RMS-S2S: root mean square sample-to-sample

ISOA: isoline area

SD-OCT: spectral domain optical coherence tomography

Key words: AMD, drusen, fixational eye movements, fovea

[1]Sorbonne Université, INSERM, CNRS, Institut de la Vision, France ; [2]CHNO des Quinze-Vingts, INSERM-DGOS CIC, France ; [3]Office National d'Etudes et de Recherches Aérospatiales (ONERA), France ; [4]Quantel Medical SA, France ; [5]Institut Langevin, CNRS, ESPCI France ; [6]Lighthouse Guild, United States

J. Murari and J. Gautier contributed equally to this work.

S. Meimon, M. Paques and A. Arleo supervised equally this work.

*Corresponding author: Denis Sheynikhovich (denis.sheynikhovich@sorbonne-universite.fr)

Abstract

Purpose: This study aims at linking subtle changes of fixational eye movements (FEM) in controls and

in patients with foveal drusen using adaptive optics retinal imaging in order to find anatomo-functional

markers for pre-symptomatic Age-related Macular Degeneration (AMD).

Methods: We recruited 7 young controls, 4 older controls and 16 presymptomatic AMD patients with

foveal drusen from the Silversight Cohort. A high speed research-grade adaptive optics flood

illumination ophthalmoscope (AO-FIO) was used for monocular retinal tracking of fixational eye

movements. The system allows for sub-arcminute resolution, high-speed and distortion-free imaging of

the foveal area. Foveal drusen position and size were documented using gaze-dependent imaging on

a clinical-grade AO-FIO.

Results: FEM were measured with high precision (RMS-S2S=0.0015° on human eyes) and small foveal

drusen (median=60µm) were detected with high contrast imaging. Microsaccade amplitude, drift

diffusion coefficient and ISOline Area (ISOA) were significantly larger for patients with foveal drusen

compared with controls. Among the drusen participants, microsaccade amplitude was correlated to

drusen eccentricity from the center of the fovea.

Conclusions: A novel high-speed high-precision retinal tracking technique allowed for the

characterization of FEM at the microscopic level. Foveal drusen altered fixation stability, resulting in

compensatory FEM changes. Particularly, drusen at the foveolar level seemed to have a stronger

impact on microsaccade amplitudes and ISOA. The unexpected anatomo-functional link between small

foveal drusen and fixation stability opens up a new perspective of detecting oculomotor signatures of

eye diseases at the presymptomatic stage.

Introduction

Age-related macular degeneration (AMD) is the main cause of vision loss in developed

countries, affecting 180M people in 2020 with an expected prevalence of 288M by 2040[1]. AMD starts

with the accumulation of extracellular deposits, called drusen, in-between the Bruch's membrane and

the retinal pigment epithelium (RPE). Multiplication and expansion of drusen is associated with late

stages of AMD, which are classified into exudative (wet-AMD) and/or atrophic (dry-AMD)[2]. Wet-AMD is

caused by irregular blood vessel growth (i.e., neovascularization) at the macular level, and it affects

approximately 10-20% of AMD patients[3]. Although macular neovascularization can be successfully

treated using anti-VEGF therapies, these treatments do not prevent the development of atrophy, which

causes sight loss in the long term[4]. Geographic atrophy or dry-AMD is the irreversible loss of

photoreceptors caused either by drusen development or neovascular growth, and it accounts for about

80 to 90% of the cases[2]. No effective treatment currently exists for this form of AMD. The absence of

efficient treatment strategies for the late-stage AMD is exacerbated by the fact that patients frequently

remain unaware of their AMD condition[5], so that they typically receive a diagnosis only upon the

emergence of clinical indications, often at a stage too advanced for effective implementation of

preventive strategies. There is therefore an urgent need to characterize anatomo-functional

relationships associated with the earliest asymptomatic AMD stages aiming at complementing existing

medical imaging techniques with functional measures of retinal anomalies that can serve as biomarkers

of future disease onset. If indeed the pre-symptomatic structural anomalies are associated with

functional, e.g. oculomotor, changes, their timely detection may also incite the individuals with a risk of

developing AMD to engage in risk mitigation strategies to delay AMD onset and progression[6,7,8].

The objective of the present study is to test the novel hypothesis that the appearance of drusen,

an early structural anomaly in presymptomatic AMD, is associated with detectable changes in

microscopic eye movements during ocular fixation. Even though the development of early-to-late stages

of confirmed AMD has already been associated with changes in fixation stability, decreased visual

acuity, reduced visual field, slower reading speed, lower contrast sensitivity, and an overall deterioration

of functional vision useful for daily activities[9–11], these previous studies did not explore fixational eye

movements in presymptomatic stages, nor did they characterize the link between retinal structure

changes and fixational eye movements. In patients with central scotomas, fixation stability as assessed

by retinal motion compensation strategies has been shown to worsen with AMD progression[12].

Moreover, advanced geographic atrophies were reported to affect different components of fixational

eye movements, such as microsaccades (small saccades of amplitude below 1°) and ocular drift[13,14]

(random Brownian-walk-like eye movements between microsaccades). More specifically, these studies

report higher microsaccade amplitudes associated with macular lesions[13] and lower spectral whitening

of natural scenes by ocular drift in patients with long-lasting macular disease, compared to age-matched

controls[14]. Given the significant changes in fixational stability in advanced forms of AMD, it is important

to test whether even the first signs of potential future disease can induce changes in eye movement

statistics.

The hypothesis was tested by simultaneously performing *(i)* high-resolution retina imaging to

visualize and characterize small asymptomatic alterations of the foveal structure, *(ii)* high-speed retina

tracking to detect fixational eye movements. We studied the relationship between subtle structural

retinal anomalies and fixational eye movement characteristics by examining healthy controls and

participants with foveal drusen, but no scotoma or atrophy according to conventional measures. If the

appearance of first drusen indeed leads to fixation instability, as the current study shows, such a

structure-function relationship provides the possibility to use fine oculomotor measurements to predict

the appearance of retinal anomalies and therefore contribute to the development of early anatomo-

functional biomarkers for urgently needed screening approaches in presymptomatic AMD.

Methods

Participants

A cohort of 39 participants were recruited for the study as part of one of three groups: young

controls (n=7, [2 females], age=31±7 yo), older controls (n=20, [14 females], age=75±4 yo) and drusen

participants (n=12, [8 females], age=79±3 yo). The initial assignment of participants to the groups relied

on the detection of drusen by standard spectral-domain optical coherence tomography (SD-OCT)

imaging (Spectralis OCT, Heidelberg Engineering, Germany). All participants were recruited from the

SilverSight cohort[15], and they were fully screened and ascertained not to suffer from any age-related

sensory, cognitive, or motor pathology.

Prior to the beginning of the experiment, older controls and drusen participants were screened

using a novel imaging method, developed previously in our laboratory using Adaptive Optics Flood

Illumination Ophthalmoscope (AO-FIO, rtx1, Imagine Eyes), which is capable of detecting foveal drusen

not visible in standard SD-OCT or fundus imaging[16] (see the next section). As a consequence, 9

participants that had initially been recruited as older controls were actually found to have drusen in the

foveal region, and they were thus moved into the drusen group. In addition, a total of 12 participants

had to be excluded from study: 3 because of too blurry retina images on the rtx1; 3 due to poor signal-to-noise ratio during the fixation task; 2 because they were not fixating the right target; 3 because of excessive tiredness and discomfort during the experiment; and 1 because she exhibited fixation patterns typical of another disease. As a result, a total of 27 participants were kept for the analyses: 7 young controls ([2 females], age=31±7 yo), 4 older controls ([1 female], age=72±2 yo), and 16 drusen patients ([10 females], age=78±4 yo). Their demographic and ophthalmological characteristics are given in **Table 1**.

All clinical, experiment, and data management procedures were approved by the Ethical Committee "CPP Ile de France V" (ID RCB 2015-A01094-45, n. CPP: 16122 MSB), and they adhered to the tenets of the Declaration of Helsinki. A signed consent was obtained by each participant to be included in the study as well as for the inclusion in the Multimodal Ophthalmologic Imaging protocol (IMA-MODE, ID_RCB 2019-A00942-55 CPP: Sud Est III 2019-021-B) of the Quinze-Vingts National Ophthalmology Hospital.

Drusen detection by gaze-dependent imaging

In order to reliably detect the presence of even the smallest drusen, gaze-dependent fundus imaging of all older participants' central retina was performed using a commercial clinical-grade AO-FIO (rtx1, Imagine Eyes) following the method and protocol described by Rossi et al.[16]. Compared to direct single-shot retinal flood imaging, gaze dependent imaging computes the local contrast maxima across a stack of 5 images whose gaze direction has been sequentially shifted: one 4°x4° fovea-centered image of the retina, focused on the photoreceptors, and four 4 horizontally and vertically displaced images (±1° with respect to the center image). After image registration, the maximum of the standard deviation for each x, y pixel of the 5-image stack is computed to make gaze-varying structure visible. In the resulting contrast image, drusen and their delineation are clearly visible (**Fig. 2c**). This gaze-dependent retina imaging method allows for the detection of foveal drusen less than 25 μm in diameter that are not detected by standard imaging methods and it also visualizes much more reliably drusen 25-60 μm in diameter, providing more precise information about their size and position[16]. In a final step, drusen were manually annotated as ellipses by two experts in medical images, followed by a validation by two ophthalmologists (including M.P.). All the image filtering, registration, and processing were performed using custom software written in MATLAB (Mathworks, USA, 2023).

Ocular fixation task

Following drusen assessment, all participants were tested in an ocular fixation experiment in order to see whether the presence of drusen affects fixational eye movements. In order to precisely detect even the smallest eye movements, video recordings of the fovea during fixation were performed using the PARIS AO-FIO (ONERA)[17] located at the Quinze-Vingts National Ophthalmology Hospital in Paris, France. Both the PARIS AO-FIO retinal imaging setup and retinal motion tracking process are described in the following sections. Before the ocular fixation task, the dominant eye was dilated with a drop of 1% tropicamide (*Mydriaticum*, tropicamide 0.5%). The recordings were performed in a dark room with minimal stray light and with the participant's head fixed on a chinrest. The fixation task consisted of two phases. In the initial phase, participants were required to fixate a central cross for 5 seconds. This phase served to initialize the retinal motion tracking procedure. In the experimental phase, the participants were asked to fixate a circle (outer diameter: 17 arcmin (0.28°), inner diameter: 11 arcmin (0.18°)) for 30 seconds, across 5 trials. The instructions were *"Please, fixate the circle for 30 seconds. Do not speak and try not moving your head. Do not hesitate to blink if you need to.",* and a vocal indication about time progression was provided after 15 seconds.

Retina imaging setup

High-resolution structural imaging of the fovea was performed using the PARIS AO-FIO[17], through video recordings focused on the photoreceptor layer. Concurrently with retina imaging, the Paris AO-FIO allowed for the projection of NIR illumination patterns (i.e., stimuli) onto the retina via a deformable Digital Micromirror Device (DMD)[18] as well as for partial-field illumination[19]. The stimulus projection with the DMD was controlled using a custom-made Python code and the ALP4Lib python library wrapper. The PARIS AO-FIO system consists of two optical channels: the Wavefront correction channel, and the illumination and imaging channel, combined with the stimulation channel. A fibered SuperLuminescent Diode (Exalos, Switzerland) with nominal center wavelength of 750 nm measures optical aberrations, and an 850nm LED source (Thorlabs, USA) coupled with a liquid fiber homogeneously illuminate the retina over a 4° diameter field of view (**Fig. 1a**). The adaptive optics compensated for the spherical, cylindrical and some higher order aberrations of the eye's optics before the stimulus reached the retina. The backscattered light from the eye was reflected back on the

161 deformable mirror before reaching the imaging camera. The adaptive optics loop was controlled with a

162 LabVIEW program and image acquisition was monitored via the Holovibes software[20].

Retina motion tracking

During the initial phase of the ocular fixation task (5s), a full field sequence of images covering

a 4° diameter field of participants' retina was recorded at 100 Hz (2048 x 2048 pixel frames). The

recorded full field retinal images were first stabilized and then averaged into a unique squared full field

reference image of the center of the retina (**Fig. 1a**). Then, during the experimental phase of the task,

only a sub-band (2° x 0.5°, 1400 x 256 pixel, with the black circle target at the center) was recorded at

800 Hz to be used for retina motion tracking (**Fig. 1b**). Retinal motion was tracked using a phase-

correlation registration method[21,22] (**Fig. 1c**) after bandpass filtering the image sequences. The

algorithm consisted in finding the position that best fitted each frame of the high-speed video onto the

reference image. The best fit was considered as the peak value coordinates of the correlation map

computed in the Fourier domain. See **Clip 1**.

Even if the maximum sampling frequency of the camera was only limited by vertical pixel

resolution, we found that 1400-pixel horizontally offered a good tradeoff between field coverage and

video bandwidth. The resulting 0.684 MB/frame (16 bits) led to a very large 547 MB/s video bandwidth

that could only be recorded with solid state hard drives and ad-hoc recording software (Holovibes).

Videos were recorded on a separate computer than the one used for stimulus presentation.

Detection of retina motion related events

Processing of the raw pixel displacement of each frame relative to the reference images to

ocular movement in arcminutes with annotated events was made with a MATLAB custom program

(based on the work of Bowers and Gautier[23]). Microsaccades were detected using an acceleration

threshold inspired by Engbert[24]. An acceleration, rather than velocity, threshold (factor N=3 standard

deviation of the median) was used in order to differentiate fast ocular drift epochs from small and slower

microsaccades which can have similar velocities but distinct accelerations. The acceleration was

computed for the 30 s fixation, and smoothed using running average with a 70~ms window. This

adaptive algorithm allowed for thresholds that fitted the participants who had relatively small

microsaccades and stable ocular drift, as well as the participants with larger microsaccade and drift

amplitudes. In addition, to ensure that microsaccades were all entirely detected and that some parts of

ocular drift were not mislabeled as microsaccades, all traces were manually verified after the automatic

annotation. Blinks were detected automatically as sudden drops in mean frame pixel intensity, which

corresponded to the closed eye reflecting less light back to the camera. Rejected data were epochs of

ocular coordinates marked by the user as false data due to failures in the imaging or the tracking (to

the exclusion of blinks). They accounted for 0-3% of the fixation data for each participant. Everything

in-between the three types of events – microsaccades, blinks or rejected data – was automatically

marked as drift.

Root mean square sample to sample (RMS-S2S) deviation was calculated to measure the

precision of the retina tracking method. A given RMS-S2S of an eye-tracker quantifies the average

movement due to noise between successive data is its squared value[25,26]. Filtering ocular traces can

have a strong effect on RMS-S2S and skews the results[27]. However, in this tracking method, ocular

traces were used raw, and RMS-S2S appeared to be the right metric to measure precision.

Estimation of fixation stability

The Bivariate Contour Ellipse Area (BCEA) and, more recently, the ISOline Area (ISOA) metrics

have been used to clinically measure fixation stability of patients with central vision loss, including

AMD[28]. ISOA is more robust than BCEA as it can delineate non-ellipsoidal fixation patterns[29] and it can

be used to detect possible multiple retinal fixation loci (PRLs), such as in AMD patients. The choice of

the bandwidth and the extent (i.e., number of standard deviations) to compute the ISOA is relatively

arbitrary and open to criticism[29–31]. The choice of a 1° window for kernel density estimation, which

subtends ISOA calculation, has been shown to be robust and appropriate with respect to the classical

definition of the 1° foveola[30,31]. Concerning the ISOA contour extent, we used both the standard 68%

ISOA (i.e., encompassing 68% of the data points) and the 86% ISOA (i.e., computed over 2 standard

deviations in the 2D space defined from the center of the fixation distribution[32]. The rationale was that

the 86% ISOA contour would provide complementary insights on the global fixation pattern that might

better differentiate healthy controls from participants with foveal drusen.

Drift diffusion coefficient and amplitude spectrum computation

In the framework of Brownian motion[33,34], the eye movement during drift epochs can be modeled by a random walk, and motion statistics can be measured by the diffusion coefficient D. We computed the diffusion coefficient D by least square best fit to the measurements of the empirical standard deviation of the eye displacement as a function of time[35]. The motion amplitude spectra of the ocular drift segments were analyzed by using a multitaper spectral analysis. This analysis was carried separately on the horizontal and vertical components of eye motion[36,37], on overlapping segments of 1 s. The time half-bandwidth product was fixed at 2.5, and the last taper was dropped to maximize spectral concentration ratios in the Slepian sequences.

Statistical analysis

Linear mixed-effect model fit by maximum likelihood (*fitlme* function on MATLAB) was used to compare FEM components between groups with the formula (1). $FEM_{component}$ were microsaccade amplitudes, ISOAs or drift coefficients.

$$(1) FEM_{component} \sim groups + (1|trial)$$

Results

Detection and characterization of foveal drusen

Figure 2 shows foveal drusen detected in one participant by using gaze-dependent retinal fundus imaging protocol[16] (**Fig 2c)**, as well as the corresponding region in an SD-OCT cut across the fovea (**Fig 2a-b**). Overall, in 9 out of 20 participants (45%), initially deemed to have no drusen and therefore assigned to the control group based on SD-OCT images, the presence of foveal drusen was detected using the gaze-dependent detection procedure. These participants were therefore reassigned to the drusen group (see Methods). Across all participants of the drusen group, we quantified the number of foveal drusen, their eccentricity with respect to the center of the fovea, their areas, their diameters (i.e., computed as the average diameter of the ellipses), and the total area covered by all drusen (**Table 1**). The median drusen diameter in the study was 60 μm (0.2°) (**Fig 2e**).

Precision and robustness of retinal motion tracking

Retinal tracking during ocular fixation was performed using the Paris AO-FIO[17] as described in Methods. **Figure 3** shows the example traces recorded during 4s of the ocular fixation task in one control participant and one participant with ocular drusen. Our retinal tracking method allowed both microsaccades (shown in magenta) and ocular drift (shown in blue) to be easily visualized and detected in both control and drusen participants.

The precision of the tracking procedure was quantified by the average RMS-S2S and achieved 0.0015°, as measured on unfiltered data across all participants. This value is an order of magnitude lower than what is deemed sufficient to detect microsaccades[25] (0.03°). A low standard deviation of the tracking error (σ=0.0006°) suggests that the method is very robust from participant to participant. Finally, in order to assess possible noise sources stemming from either mechanical vibration of the Paris AO-FIO setup or defects of the tracking algorithm, as well as to get a standard RMS-S2S value, we quantified the precision of the tracking method on non-moving artificial eye (OEMI-7 Ocular Imaging Eye Model). We found an RMS-S2S of 0.00019° (σ=0.00009° over 4 recordings). The precision found on human eyes and the artificial eye is equal or better than any reported values in literature (0.0069° on a DPI Gen5.5 with an non-moving artificial eye[38] and 0.012° on a human eye[39]; 0.0012-0.0045 on the EWET1 prototypes on human eyes[39]) without being affected by lens-wobble artifacts such as in a DPI or distortion artifacts such as in an AO-SLO system (no RMS-S2S values found for AO-SLO in literature).

The sensitivity to noise was additionally assessed by power spectrum analysis. Power spectra were only weakly affected by noise across all subject groups, even at higher frequencies (above 100Hz, **Figs 5c,d**). The power spectrum of the still model eye was one to two orders of magnitude below human participants (**Supp. Fig. 1**), suggesting negligible setup-related noise.

An addition to be very precise, the method was also found to be resistant to variations in lens transparency (**Supp. Fig. 2**). More specifically, in 11 participants (41%), classified by orthoptists as having corticonuclear lens opacities, the mean RMS-S2S was 0.0018°, only slightly superior to the mean RMS-S2S of all other participants (0.0014°).

Effect of eccentricity of foveal drusen on microsaccades

To see whether the presence of foveal drusen is associated with changes in eye movements, we plotted individual distributions of microsaccade amplitudes sorted from lowest to highest mean amplitude (**Fig 4a).** The distributions of young and older participants from the control group were shifted toward the lowest amplitudes, whereas those from the drusen group towards the highest ones, suggesting a link between drusen properties and FEM characteristics. Strikingly, we also noted that the highest microsaccade amplitudes were associated with the presence of drusen mainly inside the foveola (the 0.4-0.6° foveal pit)[40], suggesting that drusen eccentricities may play an important role in oculomotor changes induced by the presence of drusen.

To quantify this effect, we computed Pearson's correlation coefficients between the measured drusen characteristics (i.e. their number, area and eccentricity) and microsaccade sizes. We found a significant negative correlation between the participants' drusen eccentricities and their microsaccade amplitudes (r=-0.74, p=0.001). Hence, the closer drusen were to the center of the fovea, the larger were microsaccades. Conversely, the more eccentric were the drusen in a particular participant, the more similar this participant was to the control subjects in terms of the microsaccade amplitude. No significant correlation was found between the number of drusen, the mean drusen area or the total drusen area and microsaccade amplitude (r=-0.003, p=0.99; r=0.16, p=0.53; r=0.13, p=0.61, respectively. The results in **Fig. 4b** show that microsaccade amplitudes start increasing exponentially when drusen are located closer than 0.8°-1° with respect to the center of the fovea.

To further characterize the role of drusen eccentricity, participants from the drusen group were subsequently classified into two subgroups. The classification took into account the approximate diameter of the foveola (0.4-0.6°[40]), of the stimulus (0.28°), and of the median foveal druse (0.2°). The first subgroup (named DIF, for "drusen inside the foveola") included the participants with drusen mainly located in the foveola (i.e., mean drusen eccentricity less than 1° with respect to the center of the fovea, n=8). The other subgroup (named DOF, for "drusen outside the foveola") consisted of participants with drusen located mainly outside their foveola (n=8).

Statistical analysis using linear mixed effects model revealed significant differences between the microsaccade amplitude in the DIF group and the three other groups (**Fig. 4c**; DIF vs young controls:

|t|=6.5, p=10⁻⁹; DIF vs older controls: |t|=6, p=10⁻⁹; DIF vs DOF: |t|=5, p=10⁻⁷). The group with drusen

outside the foveola (DOF) tended to have larger microsaccades than young controls (|t|=1.7, p=0.09)

and older controls (|t|=1.8 p=0.06). There was no difference between young and older controls (|t|=0.4,

p=0.66).

While no significant differences were found between groups in terms of *average* drusen area,

we hypothesized that large drusen closer to the center of the fovea should have an even stronger effect

on microsaccade amplitude. The rapid drop-off of both cone density and acuity in the fovea suggest

that both drusen area and eccentricity can have tremendous effect on foveal vision. We therefore

computed the correlation between drusen area divided by eccentricity (that we called drusen density)

and measured FEM characteristics. This new measure was indeed stronger correlated with

microsaccade amplitude (r=0.82, p=0.0001) than the eccentricity alone. There were no statistical

differences in the rate of microsaccades between groups and all participants produced approximately

1-3 microsaccades per second.

The above results show that drusen eccentricity, a simple property of retinal structure

associated with presymptomatic stage of AMD, is a crucial factor explaining the difference in

microsaccade size between control participants and subjects with a relatively elevated risk of

developing the disease. The quality of this simple structural measure in explaining functional changes

can be further improved by taking into account drusen area.

Impact of foveal drusen on ocular drift and eye motion spectrum

Apart from microsaccades, the ocular drift component of FEM (see **Fig. 3**) was also affected by

the presence of foveal drusen, but to a lesser extent. Indeed, the analysis of ocular drift in the four

participant groups defined in the previous section revealed significant differences in drift diffusion

coefficients between participants with drusen in the foveola and control subjects (**Figs. 5a,b**; DIF vs

young controls: |t|=2.5, p=0.01; DIF vs older controls: |t|=2, p=0.04). Statistical pairwise comparisons

between all other groups have shown that the DOF group was significantly different from young controls

(|t|=2, p=0.04), but not from older controls (|t|=1.7, p=0.1). No differences were found between young

and older controls (|t|=0.1, p=0.93, the absence of statistical differences here should be taken with

caution considering the low sample size of older controls). Individual diffusion coefficient revealed larger

variability within groups but less between groups (**Fig 5a,b**) when compared to mean microsaccade

amplitude (**Fig 4a**). No significant correlations were found between drift diffusion coefficient and drusen

properties across all subjects (eccentricity: r=0.09, p=0.72; drusen number: r=-0.003, p=0.99; mean

drusen area: r=-0.1, p=0.69; total drusen area: r=-0.05, p=0.85)

A different way to analyze FEM changes induced by drusen presence is look at the motion

spectrum across the subject groups. The spectral analysis of ocular drift revealed a progressive

increase in motion amplitude both with age and drusen centrality (**Fig. 5c**), marked in the middle

frequency range (10-40 Hz). No difference in motion amplitude was observed at high frequencies (100-

400 Hz) between the two drusen groups, in contrast to an age-related difference in control subjects

(**Fig. 5c**). When accounting for the spectral changes caused by microsaccades (**Fig. 5d**), the foveolar

drusen group clearly stood out across all motion frequencies, potentially impeding later-stage neural

processing of the visual input[35]. The differences between the two motion spectra (**Fig. 5c,d**) reflect the

statistical differences in drift diffusion coefficient between subjects groups presented above and

highlight the more important role of microsaccades, compared to ocular drift, in FEM changes induced

by the presence of central drusen.

Foveolar drusen impair fixation stability

Fixational movements of the eye, including microsaccades and drift, are the primary

determinants of fixational stability, a functional measure of ocular stability at the macroscopic level

usually evaluated by such measures as ISOline Area[28,29] (see Methods). The discovered relationship

between central drusen and microsaccade size, described in the previous section, suggests that a

similar relationship should exist between drusen eccentricity and ocular stability in the fixation task.

Indeed, we found a strong and significant negative correlation between ISOA (at 86% level) and drusen

eccentricity (r= -0.72, p=0.001) as well as highly significant positive correlation between ISOA and

drusen density (r= 0.72, p=0.0004 across all subjects. We checked that there were no significant

differences between participants in terms of microsaccade frequencies that could explain changes in

fixation stability. In order to better understand the effect of drusen on fixation stability and its topological

characteristics, we quantitatively compared average ISOA values across the four groups of subjects

using linear mixed modeling approach and visualized average ISOA contours. The participants with

drusen mainly in the foveola had the largest ISOA as compared to the other three groups (DIF vs young

controls: $|t|=5.5$, $p=10^{-7}$; DIF vs older controls: $|t|=4.9$, $p=10^{-6}$; DIF vs DOF: $|t|=5.4$, $p=10^{-7}$). ISOAs

remained similar between the DOF group and young controls ($|t|=0.24$, $p=0.81$) or older controls

($|t|=0.24$, $p=0.67$), and between young and older controls (($|t|=0.2$, $p=0.83$). These results extend our

conclusion from the previous section by showing that impaired fixation stability in subjects with drusen

in foveola is primarily mediated by the increase in microsaccade size and not by other FEM properties,

such as ocular drift.

Given the consistent and strong effect of foveal drusen on both fixation stability and

microsaccade amplitude, it is not surprising that there was a strong correlation between the ISOA 86%

values and microsaccade amplitudes across the entire study population (**Fig. 7a**; $r^2=0.85$, $p=0.001$).

Interestingly when looking at figure 6, in all groups except DIF, higher values for ISOA seemed to come

with both proportional increases in microsaccade amplitude and drift diffusion coefficient at the same

time. However, for the DIF group, drift diffusion coefficient appeared to lose its correlation with ISOA

compared to the rest of the study population.

Discussion

The objective of the present study was to see whether structural retinal changes in

presymptomatic AMD can have detectable consequences on functional aspects of vision. More

specifically, we tested the hypothesis that FEM abnormalities are associated with early structural retinal

anomalies, such as the appearance of small central drusen.

The primary finding of this study is that participants with foveal drusen exhibit significantly larger

microsaccades and higher fixation instability (measured by ISOA) as compared to healthy controls,

while maintaining equivalent microsaccade rates. Strikingly, drusen eccentricity with respect to the

center of the fovea appears to be the main determinant factor associated with larger microsaccades

and ISOAs. Indeed, we found that the more central the drusen were, the more unstable the fixation

was; conversely, the further from the center of the fovea the drusen were, the more the participants

tended to behave like controls, with minimal to no changes in fixation stability. We propose the following

quantification criterion to assess the point after which drusen eccentricity strongly affect fixation

characteristics: Microsaccade amplitude starts increasing exponentially when average drusen

eccentricity is less than 1° with respect to the center of the fovea. Considering the size of the typical

drusen and of the fixation target, the threshold point approximately corresponds to the size of the

foveola, i.e. the 350 µm foveal pit as defined by Kolb[40]. Our results further suggest that quantification

of microsaccade amplitude and ISOA in a simple ocular fixation test could in principle be sufficient to

discriminate healthy participants from those with retinal anomalies in the foveola, such as drusen.

Previous studies in patients with central scotomas suggest that these patients may develop one or more

eccentric areas of fixation (referred to as preferred retinal locus or PRL)[41], giving rise to an extended

overall fixation area. We therefore checked in our presymptomatic subjects from the DIF group whether

they express increased and possibly multipeaked fixation densities (as opposed to Gaussian-like

distributions expected for the other groups). This was indeed the case, as only in the DIF group the

ISOA contours exhibited multiple islands of fixation (see Fig. 6). In this study, we also evaluated the

impact of foveal drusen density and eccentricity on the ocular diffusion drift coefficient, which provides

a displacement estimator reflecting the randomness and stability of visual fixation[35]. The motion

spectrum of drift epochs without microsaccades appeared however impacted by the presence of drusen

and their centrality, especially in the middle frequency range. The dynamics of the complete eye

movements (i.e. microsaccades included) during fixation appears successively altered with age, with

the presence of foveal and further with foveolar drusen. This confirms the impact of drusen on

microsaccade dynamics, and the importance to finely characterize them. Further research is needed to

understand the role and implication of drusen on visual sampling and perception.

Another interesting observation stemming from this study concerns the definition of *healthy older adults*

from an ophthalmological point of view. Twenty participants of this study were initially recruited as

healthy controls based on the fact that (i) no visible foveal drusen were found using standard SD-OCT

imaging analyzed by ophthalmologists, and (ii) no visual symptoms (in terms of visual acuity, contrast

sensitivity, and so on) were observed during inclusion to the Silversight cohort. Subsequently, using

AO-FIO-based gaze-dependent retina imaging, 45% of these participants were found to actually have

small drusen in their fovea. This experimental observation, made with the use of more advanced

imaging technique, aligns with early studies suggesting that drusen are more prevalent than previously

assumed[42,43]. It also emphasizes the utmost importance of the development of sensitive measurement

techniques that can be applied at presymptomatic stages to detect the presence of small drusen. While

recent progress in retinal imaging resulted in retromode[44] or gaze-dependent[16] methods, which can

both detect drusen invisible in standard SD-OCT images, this study opens up a new avenue in drusen

detection research by showing that the appearance of small drusen can be detected based on simple

functional measures, i.e. by looking at the FEM. Moreover, the discovered link between central drusen

and fixation stability can explain the discrepancy present in the literature regarding the differences in

fixation stability between young and healthy older adults, with some studies reporting an age-related

decrease in stability[45–47] and other studies, including ours, reporting no effect of age[48–50]. If the presence

of foveal drusen is a common trait among older participants and these drusen do affect fixation stability,

as we observe in our experiment, then the putative effect of age on fixation stability in the above studies

can in fact be explained by the presence of participants with undetected central drusen in the sample

of older adults (since virtually all previous studies used standard retina fundus imaging or SD-OCT). In

conclusion, a combination of imaging methods and functional oculomotor measures, with a potential

addition of subjective patient-reported measures of retinal health[50] can be part of the general strategy

for multifactor screening for retinal diseases, including AMD, in the general population. DIF participants

were clearly clustered together around higher values of both ISOA and microsaccade amplitudes,

suggesting the possibility of automatic detection procedure for the existence of foveolar drusen that

could for instance use both ISOA and microsaccade amplitude values. While it is currently impossible

to know whether or not existing drusen will develop into severe forms of AMD[51], detection of drusen at

the earliest possible stage will allow to monitor their progression and take the necessary precautions

(e.g. apply anti-VEGF treatment[4]) before critical and irreversible damage occurs. Health style changes

might help slow down their growth, and new drugs are currently being clinically tested to prevent the

formation and evolution of drusen[7].

To the best of our knowledge, alterations of fine oculomotor control in relation to small,

asymptomatic changes of retinal structure have never been previously reported, whereas the

assessment of eye movements to identify non-invasive biomarkers of a number of diseases is not

new[52]. Research on eye movements has indeed existed for more than a century, and infrared video

eye trackers capabilities, accuracy, and accessibility have renewed the interest on eye movements to

clinicians and psychologists. FEM have been explored in neurological disorders such as amblyopia[53,54],

schizophrenia[55], Parkinson's disease[56], and others[57]. There have also been attempts to reveal new

FEM-based biomarkers to predict multiple sclerosis and follow its progression[58]. However, when it

comes to detecting subtle retinal structural anomalies by means of FEM analysis, it is key to be able to

record the whole spectrum of microsaccade amplitudes, down to 1 arcminute. The recent use of

adaptive optics retinal imaging instruments has made it possible to reach this resolution. The AO-FIO

used in this study allowed us to observe an average healthy microsaccade amplitude of 6.3 arcminutes,

whereas participants with drusen had an average microsaccade amplitude of 13.8 arcminutes. In the

past 20 years, FEM studies using pupil-based eye trackers have defined a microsaccade with

amplitudes up to 1-2°[25,59–61], and basic fixation tasks often showed average microsaccade amplitudes

of 20-60 arcminutes. It is thus possible that the differences between controls and drusen participants

found in our study would not have been detected using these less precise eye trackers and/or a fixation

target larger than ours, because microsaccade amplitude is known to be target-dependent[23]. Further

studies are being conducted in our team to confirm such hypotheses. This study, like many in the past

decade, shows that FEM reveal a lot of unique information about patients. The system we use is

probably too advanced and complex for immediate clinical use but the study is paving the road towards

future FEM analysis in ophthalmologic testing with future accessible tracking methods that are starting

to become available in the market.

While general functional significance of microsaccades is still a matter of debate, many

researchers agree that microsaccades play an important functional role in the presence of foveolar

stimuli, and the foveola size is even considered as a reference to define microsaccades[59,62]. As

discussed in Poletti's review on attention and eye movements at the foveal scale[59], any stimuli in the

foveola can be attention-grabbing and whether it is voluntary or not, it is possible to have microsaccade

directed towards different stimuli as close as 20 arcminutes from each other. It would not be correct to

consider foveal drusen as inputs to the visual system, but our data suggest that they might change

retinal processing in specific retinal locations. Drusen appears to distort foveal vision but the exact role

of or consequence on the oculomotor system remains to be determined. Fixation instability could be

the result of the oculomotor system adaptation to avoid drusen locally. The second hypothesis consists

in a more widespread RPE/photoreceptor dysfunction and degraded foveal sensory integration which

would in turn impair motor control. The local compensatory versus global oculomotor deficiency

hypotheses require further investigation. The relative impact of each eye's drusen on binocular vision

is another open subject of research to supplement these findings.

In conclusion, by combining state-of-the-art retinal imaging techniques and novel retinal motion tracking

approach, we report an association between the presence of foveal drusen and fixation instability, hence

linking structural and functional changes during aging. By demonstrating a strong association between

foveal drusen properties and oculomotor behavior in a simple ocular fixation task our results suggest

that FEM can be considered as a functional biomarker for innovative AMD screening approaches. More

generally, our results suggest a possibility that the presence of even small drusen may affect visual

perception. Thus, further studies are required to test potential effects of drusen on performance in tasks

requiring high visual acuity, contrast sensitivity and photoreceptor function.

References

1. Wong WL, Su X, Li X, et al. Global prevalence of age-related macular degeneration and disease burden projection for 2020 and 2040: a systematic review and meta-analysis. *The Lancet Global Health*. 2014;2(2):e106-e116. doi:10.1016/S2214-109X(13)70145-1

2. Danis RP, Lavine JA, Domalpally A. Geographic atrophy in patients with advanced dry age-related macular degeneration: current challenges and future prospects. *Clin Ophthalmol*. 2015;9:2159-2174. doi:10.2147/OPTH.S92359

3. Pugazhendhi A, Hubbell M, Jairam P, Ambati B. Neovascular Macular Degeneration: A Review of Etiology, Risk Factors, and Recent Advances in Research and Therapy. *Int J Mol Sci*. 2021;22(3):1170. doi:10.3390/ijms22031170

4. Rofagha S, Bhisitkul RB, Boyer DS, Sadda SR, Zhang K. Seven-Year Outcomes in Ranibizumab-Treated Patients in ANCHOR, MARINA, and HORIZON: A Multicenter Cohort Study (SEVEN-UP). *Ophthalmology*. 2013;120(11):2292-2299. doi:10.1016/j.ophtha.2013.03.046

5. Cervantes-Castañeda RA, Banin E, Hemo I, Shpigel M, Averbukh E, Chowers I. Lack of benefit of early awareness to age-related macular degeneration. *Eye*. 2008;22(6):777-781. doi:10.1038/sj.eye.6702691

6. Schultz NM, Bhardwaj S, Barclay C, Gaspar L, Schwartz J. Global Burden of Dry Age-Related Macular Degeneration: A Targeted Literature Review. *Clinical Therapeutics*. 2021;43(10):1792-1818. doi:10.1016/j.clinthera.2021.08.011

7. Fleckenstein M, Keenan TDL, Guymer RH, et al. Age-related macular degeneration. *Nat Rev Dis Primers*. 2021;7(1):1-25. doi:10.1038/s41572-021-00265-2

8. Seddon JM. Macular Degeneration Epidemiology: Nature-Nurture, Lifestyle Factors, Genetic Risk, and Gene-Environment Interactions – The Weisenfeld Award Lecture. *Investigative Ophthalmology & Visual Science*. 2017;58(14):6513-6528. doi:10.1167/iovs.17-23544

9. Verghese P, Vullings C, Shanidze N. Eye Movements in Macular Degeneration. *Annual Review of Vision Science*. 2021;7(1):773-791. doi:10.1146/annurev-vision-100119-125555

10. Pondorfer SG, Wintergerst MWM, Gorgi Zadeh S, et al. Association of Visual Function Measures with Drusen Volume in Early Stages of Age-Related Macular Degeneration. *Investigative Ophthalmology & Visual Science*. 2020;61(3):55. doi:10.1167/iovs.61.3.55

11. Calabrese A, Bernard JB, Hoffart L, et al. Wet versus dry age-related macular degeneration in patients with central field loss: different effects on maximum reading speed. *Investigative ophthalmology & visual science*. 2011;52(5):2417-2424.

12. Macedo AF, Crossland MD, Rubin GS. Investigating Unstable Fixation in Patients with Macular Disease. *Investigative Ophthalmology & Visual Science*. 2011;52(3):1275-1280. doi:10.1167/iovs.09-4334

13. Kumar G, Chung STL. Characteristics of Fixational Eye Movements in People With Macular Disease. *Investigative*
Ophthalmology & Visual Science. 2014;55(8):5125-5133. doi:10.1167/iovs.14-14608

14. Ağaoğlu MN, Chung STL. Exploration of the functional consequences of fixational eye movements in the absence of a
fovea. *Journal of Vision*. 2020;20(2):12. doi:10.1167/jov.20.2.12

15. LAGRENE K, Bécu M, Seiple WH, et al. Healthy and pathological visual aging in a French follow-up cohort study.
Investigative Ophthalmology & Visual Science. 2019;60(9):5915.

16. Rossi EA, Norberg N, Eandi C, et al. A New Method for Visualizing Drusen and Their Progression in Flood-Illumination
Adaptive Optics Ophthalmoscopy. *Translational Vision Science & Technology*. 2021;10(14):19. doi:10.1167/tvst.10.14.19

17. Gofas-Salas E, Mecê P, Petit C, et al. High loop rate adaptive optics flood illumination ophthalmoscope with structured
illumination capability. *Applied optics*. 2018;57(20):5635-5642.

18. Krafft L, Senée P, Gofas-Salas E, et al. Multimodal high-resolution retinal imaging using camera-based DMD-integrated
Adaptive Optics Flood-Illumination Ophthalmoscope. Published online May 10, 2023. doi:10.1364/opticaopen.22769519.v1

19. Krafft L, Gofas-Salas E, Lai-Tim Y, et al. Partial-Field Illumination Ophthalmoscope: improving the contrast of a camera-
based retinal imager. *Applied optics*. 2021;60(31):9951-9956.

20. Atlan M. Ultrahigh-throughput rendering of digital holograms. In: *Imaging and Applied Optics 2018 (3D, AO, AIO, COSI, DH,*
IS, LACSEA, LS&C, MATH, pcAOP) (2018), Paper DM5F.4. Optica Publishing Group; 2018:DM5F.4.
doi:10.1364/DH.2018.DM5F.4

21. Mecê P. *4D Exploration of the Retina for Adaptive Optics-Assisted Laser Photocoagulation*. PhD Thesis. These de doctorat,
Sorbonne Paris Cité; 2018.

22. Mece P, Chen A, Petit C, Mugnier L, Paques M, Meimon S. Real-time optical stabilization of retinal motion at micrometer
precision using Adaptive Optics Flood-Illumination Ophthalmsocope. Published online December 28, 2023.
doi:10.1364/opticaopen.24907464.v1

23. Bowers NR, Gautier J, Lin S, Roorda A. Fixational eye movements in passive versus active sustained fixation tasks. *Journal*
of Vision. 2021;21(11):16. doi:10.1167/jov.21.11.16

24. Engbert R. Microsaccades: a microcosm for research on oculomotor control, attention, and visual perception. In: Martinez-
Conde S, Macknik SL, Martinez LM, Alonso JM, Tse PU, eds. *Progress in Brain Research*. Vol 154. Visual Perception.
Elsevier; 2006:177-192. doi:10.1016/S0079-6123(06)54009-9

25. Holmqvist K, Nyström M, Andersson R, Dewhurst R, Jarodzka H, Van de Weijer J. *Eye Tracking: A Comprehensive Guide*
to Methods and Measures. OUP Oxford; 2011.

26. Holmqvist K, Örbom SL, Hooge ITC, et al. Eye tracking: empirical foundations for a minimal reporting guideline. *Behav Res*. 2023;55(1):364-416. doi:10.3758/s13428-021-01762-8

27. Niehorster DC, Zemblys R, Beelders T, Holmqvist K. Characterizing gaze position signals and synthesizing noise during fixations in eye-tracking data. *Behavior Research Methods*. 2020;52:2515-2534.

28. Whittaker SG, Budd J, Cummings RW. Eccentric fixation with macular scotoma. *Investigative Ophthalmology & Visual Science*. 1988;29(2):268-278.

29. Castet E, Crossland M. Quantifying eye stability during a fixation task: a review of definitions and methods. *Seeing Perceiving*. 2012;25(5):449-469. doi:10.1163/187847611X620955

30. Martinez-Conde S, Macknik SL, Troncoso XG, Dyar TA. Microsaccades Counteract Visual Fading during Fixation. *Neuron*. 2006;49(2):297-305. doi:10.1016/j.neuron.2005.11.033

31. Otero-Millan J, Troncoso XG, Macknik SL, Serrano-Pedraza I, Martinez-Conde S. Saccades and microsaccades during visual fixation, exploration, and search: Foundations for a common saccadic generator. *Journal of Vision*. 2008;8(14):21. doi:10.1167/8.14.21

32. Wang B, Shi W, Miao Z. Confidence Analysis of Standard Deviational Ellipse and Its Extension into Higher Dimensional Euclidean Space. *PLoS One*. 2015;10(3):e0118537. doi:10.1371/journal.pone.0118537

33. Intoy J, Rucci M. Finely tuned eye movements enhance visual acuity. *Nat Commun*. 2020;11(1):795. doi:10.1038/s41467-020-14616-2

34. Microsaccades Keep the Eyes' Balance During Fixation - Ralf Engbert, Reinhold Kliegl, 2004. Accessed July 18, 2023. https://journals.sagepub.com/doi/abs/10.1111/j.0956-7976.2004.00697.x

35. Kuang X, Poletti M, Victor JD, Rucci M. Temporal Encoding of Spatial Information during Active Visual Fixation. *Current Biology*. 2012;22(6):510-514. doi:10.1016/j.cub.2012.01.050

36. Thomson DJ. Spectrum estimation and harmonic analysis. *Proceedings of the IEEE*. 1982;70(9):1055-1096.

37. Babadi B, Brown EN. A review of multitaper spectral analysis. *IEEE Transactions on Biomedical Engineering*. 2014;61(5):1555-1564.

38. Holmqvist K, Blignaut P. Small eye movements cannot be reliably measured by video-based P-CR eye-trackers. *Behav Res*. 2020;52(5):2098-2121. doi:10.3758/s13428-020-01363-x

39. Holmqvist K, Orbom SL, Miller M, Kashchenevsky A, Shovman MM, Greenlee MW. Validation of a prototype hybrid eye-tracker against the DPI and the Tobii Spectrum. In: *ACM Symposium on Eye Tracking Research and Applications*. ACM; 2020:1-9. doi:10.1145/3379155.3391330

40.   Kolb H, Nelson RF, Ahnelt PK, Ortuño-Lizarán I, Cuenca N. The Architecture of the Human Fovea. In: Kolb H, Fernandez

E, Nelson R, eds. *Webvision: The Organization of the Retina and Visual System*. University of Utah Health Sciences Center;

1995. Accessed October 23, 2023. http://www.ncbi.nlm.nih.gov/books/NBK554706/

41.   Duret F, Issenhuth M, Safran AB. Combined use of several preferred retinal loci in patients with macular disorders when

reading single words. *Vision Research*. 1999;39(4):873-879. doi:10.1016/S0042-6989(98)00179-5

42.   Klein R, Klein BEK, Linton KLP. Prevalence of Age-related Maculopathy: The Beaver Dam Eye Study. *Ophthalmology*.

1992;99(6):933-943. doi:10.1016/S0161-6420(92)31871-8

43.   Bressler NM, Bressler SB, West SK, Fine SL, Taylor HR. The Grading and Prevalence of Macular Degeneration in

Chesapeake   Bay   Watermen.   *Archives   of   Ophthalmology*.   1989;107(6):847-852.

doi:10.1001/archopht.1989.01070010869032

44.   Cozzi M, Monteduro D, Airaldi M, et al. Retromode Imaging Technology for Detecting Drusen-Like Deposits in Healthy

Adults. *Ophthalmology Retina*. Published online July 2023:S2468653023003147. doi:10.1016/j.oret.2023.07.012

45.   Cutini A, Fragiotta S, Spadea L, et al. Normal values and repeatibility of bivariate contour ellipse area (BCEA) with

Microperimeter Mp-1. *Investigative Ophthalmology & Visual Science*. 2014;55(13):169.

46.   Agathos CP, Bernardin D, Huchet D, Scherlen AC, Assaiante C, Isableu B. Sensorimotor and cognitive factors associated

with the age-related increase of visual field dependence: a cross-sectional study. *AGE*. 2015;37(4):67. doi:10.1007/s11357-

015-9805-x

47.   Morales MU, Saker S, Wilde C, et al. Reference Clinical Database for Fixation Stability Metrics in Normal Subjects Measured

with the MAIA Microperimeter. *Translational Vision Science & Technology*. 2016;5(6):6. doi:10.1167/tvst.5.6.6

48.   Kosnik W, Fikre J, Sekulerf R. Visual Fixation Stability in Older Adults. 27(12).

49.   González EG, Mandelcorn MS, Mandelcorn ED, Tarita-Nistor L. Effect of Visual Feedback on the Eye Position Stability of

Patients with AMD. *Vision*. 2019;3(4):59. doi:10.3390/vision3040059

50.   Bécu M. *Impact of Healthy Aging on Spatial Cognition. Spatial Navigation and Gaze Dynamics in Ecological Conditions*.

phdthesis. Sorbonne Université; 2018. Accessed October 2, 2023. https://theses.hal.science/tel-02862784

51.   Ferris FL, Wilkinson CP, Bird A, et al. Clinical Classification of Age-related Macular Degeneration. *Ophthalmology*.

2013;120(4):844-851. doi:10.1016/j.ophtha.2012.10.036

52.   Marandi RZ, Gazerani P. Aging and eye tracking: in the quest for objective biomarkers. *Future Neurology*.

2019;14(4):FNL33. doi:10.2217/fnl-2019-0012

53.   Shaikh AG, Otero-Millan J, Kumar P, Ghasia FF. Abnormal Fixational Eye Movements in Amblyopia. Martinez-Conde S, ed.
591   *PLoS ONE*. 2016;11(3):e0149953. doi:10.1371/journal.pone.0149953

592   54.   Ghasia F, Wang J. Amblyopia and fixation eye movements. *Journal of the Neurological Sciences*. 2022;441:120373.
593   doi:10.1016/j.jns.2022.120373

594   55.   Egaña J, Devia C, Mayol R, et al. Small Saccades and Image Complexity during Free Viewing of Natural Images in
595   Schizophrenia.   *Frontiers   in   Psychiatry*.   2013;4.   Accessed   October   2,   2023.
596   https://www.frontiersin.org/articles/10.3389/fpsyt.2013.00037

597   56.   Kaski D, Bronstein AM. Ocular Tremor in Parkinson's Disease: Discussion, Debate, and Controversy. *Frontiers in Neurology*.
598   2017;8. Accessed May 24, 2022. https://www.frontiersin.org/article/10.3389/fneur.2017.00134

599   57.   Alexander RG, Macknik SL, Martinez-Conde S. Microsaccade Characteristics in Neurological and Ophthalmic Disease.
600   *Frontiers in Neurology*. 2018;9. Accessed July 18, 2023. https://www.frontiersin.org/articles/10.3389/fneur.2018.00144

601   58.   Sheehy CK, Bensinger ES, Romeo A, et al. Fixational microsaccades: A quantitative and objective measure of disability in
602   multiple sclerosis. *Mult Scler*. 2020;26(3):343-353. doi:10.1177/1352458519894712

603   59.   Poletti M. An eye for detail: Eye movements and attention at the foveal scale. *Vision Research*. 2023;211:108277.
604   doi:10.1016/j.visres.2023.108277

605   60.   Steinman RM, Cunitz RJ, Timberlake GT, Herman M. Voluntary Control of Microsaccades during Maintained Monocular
606   Fixation. *Science*. 1967;155(3769):1577-1579. doi:10.1126/science.155.3769.1577

607   61.   Collewijn H, Kowler E. The significance of microsaccades for vision and oculomotor control. *Journal of Vision*. 2008;8(14):20.
608   doi:10.1167/8.14.20

609   62.   Martinez-Conde S, Otero-Millan J, Macknik SL. The impact of microsaccades on vision: towards a unified theory of saccadic
610   function. *Nat Rev Neurosci*. 2013;14(2):83-96. doi:10.1038/nrn3405

Tag	Age (yo)	Tested eye	Visual acuity	Number of drusen	Mean drusen area ± std (arcmin²)	Total drusen area (arcmin²)	Mean drusen eccentricity ± std (arcmin)
DI1	79	OD	1.2	7	79 ± 14	556	27 ± 22
DI2	78	OD	1	7	169 ± 66	1182	28 ± 23
DI3	80	OD	1	5	75 ± 18	376	30 ± 25
DI4	84	OS	1.2	7	143 ± 32	998	37 ± 29
DI5	80	OD	1.2	32	339 ± 147	10861	42 ± 31
DI6	83	OS	1	12	176 ± 50	2121	42 ± 26
DI7	76	OS	1	9	131 ± 72	1184	45 ± 27
DI8	74	OD	1	2	106 ± 29	212	45 ± 14
DI9	75	OD	1.2	6	178 ± 40	1069	56 ± 32
DO1	73	OD	0.8	5	148 ± 78	741	62 ± 22
DO2	84	OS	1	13	177 ± 48	2313	63 ± 25
DO3	70	OS	1	15	180 ± 70	2710	62 ± 32
DO4	80	OD	0.7	6	128 ± 49	766	68 ± 19
DO5	81	OD	1.2	20	139 ± 39	2791	69 ± 42
DO6	74	OS	0.8	14	59 ± 15	826	76 ± 25
DO7	76	OD	1.2	28	189 ± 75	5311	92 ± 34
Mean for drusen group (n=16)	78 ± 4	/	1 ± 0.2	11 ± 8	133 ± 59	2126 ± 2000	53 ± 27
Mean for older controls (n=4)	72 ± 2	/	>1	/	/	/	/
Mean for young controls (n=7)	31 ± 7	/	>1	/	/	/	/

Figure legends

Figure 1. Principle of the AOFIO retinal tracking. (a) The 4° full field obtained after initial phase of the

fixation task (5 sec at 100 Hz, in red). Only the rectangular sub-part (in blue) was used as a reference

image for subsequent eye movement analysis, as it is enough to track and monitor eye movements

during central fixation (typically less than 2° in amplitude). During the experimental phase of the task,

the 2 x 0.5° subregion (in green) is tracked at very high speed (800 Hz, 16 bits). (b) An actual frame

recorded at this frequency. (c) Eye displacement is obtained by phase-correlation over the reference

image. (d) Custom eye tracking software processes large sequences of up to 5 Mbits/frame. (e) A 3x

zoom on an area 1° away from the fovea with clearly visible photoreceptor mosaic.

Figure 2. Drusen visualization and identification. (a) SD-OCT image of a patient. A 2x zoom on the 4° center area is shown in blue. The red and orange lines are the same size as the circle diameters in (c). **(b)** Fundus image with the OCT cut of (a) in green and the blue dotted area showing the 4° zone where gaze-dependent imaging is performed. **(c)** Gaze-dependent image of the 4°x4° zone around the fovea of the same patient. Pink arrows indicate the drusen: the white ring is the slope of bump cause by the druse the while the black center part is the tip. The red and orange circles are respectively of 0.5° and 1° of radius. The green line is the ISOA 86% On the bottom left corner the fixation target to scale. **(d-e)** Distribution of drusen eccentricities (c) and diameters (d) across all participants.

635 **Figure 3.** Retina motion traces of a healthy young subject (a-b) vs a subject with drusen (c-d) with the

636 corresponding movement in the visual field. The eye traces are shown for 3 seconds. Microsaccades

637 are annotated in pink and drift in light blue. Eye position is shown in degrees. The green line is the ISOA

638 86%.

Figure 4. Microsaccade characteristics across individuals and groups. **(a)** Violin plots of individual distributions of microsaccade amplitudes across all participants, color-coded by group and sorted with respect to the mean microsaccade amplitudes. Each dot is one microsaccade during the fixation task. **(b)** Microsaccade mean amplitude with respect to drusen eccentricity from the center of the fovea. Standard errors are shown in blue bars. **(c)** Microsaccade amplitudes across all four groups. Each dot is the average microsaccade amplitude for 30 seconds (one trial).

Figure 5. Drift dynamics and motion spectra across groups – horizontal motion. **(a)** Drift

diffusion coefficient distribution across all participants, sorted from lowest to highest. Each dot is one

trial. **(b)** Drift diffusion coefficient across groups **(c)** Motion spectrum of drifts only across the four

groups. **(d)** Whole motion spectrum (drifts + microsaccades) across the four groups. An increase in

dynamic across most frequencies is visible with age, and even further with the presence of foveal or

foveolar drusen. This progressive increase is particularly visible in the Mid-frequency (10-40 Hz, top

left inset) and very-high frequency range (100-400 Hz, top-right inset).

Figure 6. Impact of drusen on fixation stability. (a) The top row shows sample data respectively from a young adult, an older adult, a participant with foveal drusen and a participant with foveolar drusen. **(b)** Probability density of the young adult participant of (a) on the left and the foveolar participant of (a) on the right. ISOA contours encompassing 68% (purple) and 86% (green) of the eye position data.

Figure 7. Contribution of microsaccade amplitude and drift coefficient to fixation stability. (a) Microsaccade amplitude vs. 86% ISOA. **(b)** Drift coefficient vs. ISOA on the right. One symbol represents one trial of a participant from its colored group.