Discoveries

Saltational episodes of reticulate evolution in the *Drosophila saltans* species group

Carolina Prediger[1,2], Erina A. Ferreira[2], Samara Videira Zorzato[1,3], Aurélie Hua-Van[2], Lisa

Klasson[4], Wolfgang J. Miller[5], Amir Yassin[2,3*] and Lilian Madi-Ravazzi[1*]

[1]Department of Biology, UNESP - São Paulo State University, São José do Rio Preto, São

Paulo, Brazil.

[2]Laboratoire Évolution, Génomes, Comportement et Écologie, CNRS, IRD, Université Paris-

Saclay, Gif-sur-Yvette, France.

[3]Institut de Systématique, Évolution, Biodiversité (ISYEB), CNRS, MNHN, EPHE, Sorbonne

Université, Univ. des Antilles, Paris, France.

[4]Molecular evolution, Department of Cell and Molecular Biology, Science for Life

Laboratory, Uppsala University, Sweden.

[5]Center for Anatomy and Cell Biology, Department of Cell and Developmental Biology,

Medical University of Vienna, Austria.

[*] These authors contributed equally.

Correspondence: amir.yassin@universite-paris-saclay.fr; lilian.madi@unesp.br.

Running title: Phylogenomics of the *Drosophila saltans* group

Conflict of interest

The authors declare no conflict of interest.

Data availability statement

Genome sequences generated for this study are available on NCBI BioProject:

PRJNA1078835.

Acknowledgments

This paper is dedicated to Prof. Hermione E. M. C. Bicudo for her significant contributions to

the study of the evolutionary genetics of species of the *Drosophila saltans* species group. We

would like to thank David Ogereau for his assistance and insights in genome assembly for this

study. We thank Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) for the

financial support to L.M.R. (Number processes: 95/06165-1, 2014/14059-0 and 2016/ 11994-

5) enabling us to collect and sequence many strains used here. We extend our thanks to the

Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for their support,

specifically grant number 141545/2020-8, as well as the France Excellence Eiffel Scholarship

Program for funding C.P.'s PhD scholarship. *Illumina* sequencing of *D. neocordata* was

performed at the SNP&SEQ Technology Platform in Uppsala, Sweden, which is part of the

Swedish National Genomics Infrastructure and Science for Life Laboratory. SNP&SEQ is

supported by the Swedish Research Council and the Knut and Alice Wallenberg Foundation.

This project was partly supported by a grant from The Swedish research council VR (2014-

4353) to L.K. We want to thank the Nouragues research field station (managed by CNRS),

which benefits from "Investissement d'Avenir" grants managed by Agence Nationale de la

Recherche (AnaEE France ANR-11-INBS-0001; Labex CEBA ANR-10-LABX-25-01). The

project was partly funded by the Austrian Science Fund FWF grant P28255-B22 to W.J.M.

and by the Richard Lounsbery Foundation to A.Y.

Abstract

Phylogenomics reveals reticulate evolution to be widespread across taxa, but whether

reticulation is due to low statistical power or it is a true evolutionary pattern remains a field of

investigation. Here, we investigate the phylogeny and quantify reticulation in the *Drosophila*

saltans species group, a Neotropical clade of the subgenus *Sophophora* comprising 23 species

whose relationships have long been problematic. Phylogenetic analyses revealed conflicting

topologies between the X chromosome, autosomes and the mitochondria. We extended the

ABBA-BABA test of asymmetry in phylogenetic discordance to cases where no "true"

species tree could be inferred, and applied our new test (called 2A2B) to whole genome data

and to individual loci. We used four strategies, two of which consisted of windows from

pseudo-reference genomes aligned to either an outgroup or ingroup species, and two based on

our new assemblies using either conserved genes or ≥50 kb-long syntenic blocks with

conserved collinearity across Neotropical *Sophophora*. Evidence for reticulation varied

among the strategies, being lowest in the synteny-based approach, where it did not exceed

~7% of the blocks in the most conflicting species quartets. High incidences of low

phylogenetic resolution (polytomy) were restricted to three nodes on the tree, that coincided

with major paleogeographical events in South America. Our results identify possible technical

biases in quantifying reticulate evolution and indicate that episodic rapid radiations have

played a major role in the evolution of a largely understudied Neotropical clade.

Keywords: phylogenomic discordance; genome assembly; historical biogeography;

introgression; cyto-nuclear conflicts; Neotropical speciation; *Sophophora*.

Introduction

Knowledge of phylogenetic relationships among species is a requirement for many evolutionary studies. However, it is often difficult to reconstruct well-resolved bifurcating trees for some clades. This could either be due to the lack of signal in the evaluated data, a condition known as "soft polytomy", or due to persistent phylogenetic conflicts among datasets leading to "hard polytomies" and reticulate patterns of interspecific relationships. A plethora of biological processes could cause such conflicts, including incomplete lineage sorting (Maddison 1989; Maddison 1997; Walsh et al. 1999; Townsend et al. 2012), horizontal gene transfer, introgression and hybridization (Schrempf and Szöllősi 2020), and adaptive radiations (Glor 2010). Phylogenetic conflict may also be caused by technical errors, such as, sequencing error, contamination, wrong model selection and general lack of quality control (Philippe et al. 2011). Recent advances in genomic analyses have significantly reduced such errors and, in a wide range of taxa, increased the number of analyzed genes hence helping to resolve early conflicting topologies. However, in many other cases, whole genome analyses demonstrated persistent phylogenetic conflicts (e.g., in plants (Wickett et al. 2014; Gagnon et al. 2022), birds (Suh 2016), sponges and ctenophores (Philippe et al. 2009; Pick et al. 2010; Chang et al. 2015; Whelan et al. 2015; Simion et al. 2017), mammals (Morgan et al. 2013; Romiguier et al. 2013; Doronina et al. 2015), amphibians (Hime et al. 2021), and insects (Owen and Miller 2022)).

Of the different processes that can reduce phylogenetic resolution, introgression and hybridization have attracted much attention, and led to the development of a number of bioinformatic tools and tests that quantify their extent across the genome (Durand et al. 2011; Pease and Hahn 2015; Malinsky et al. 2021). Site-based methods usually count the number of bi-allelic sites supporting each of three possible topologies in a species triplet with an outgroup (Figure 1A). Comparisons between the proportions of the three topologies can yield

one of four possible outcomes (Figure 1B): (i) complete trifurcation, all topologies are equally

encountered; (ii) incomplete trifurcation, such as in the case of hybrid speciation wherein two

topologies significantly exceed the third one but do not significantly differ from each other;

(iii) incomplete bifurcation, such as in the case of asymmetric introgression wherein the

proportion of all topologies significantly differ; and (iv) complete bifurcation, one topology

significantly exceeds the two others, which in their turn have nearly equal proportions.

Categories ii and iii are often considered to evidence reticulate evolution. The earliest of

introgression tests, Patterson's D or the ABBA-BABA statistic (Green et al. 2010; Durand et

al. 2011), compared the two later cases (iii and iv), *i.e.* it presumed that a "true" species tree

exists. A later test, HyDe (Blischak et al. 2018), quantifies admixture (γ) from the ratio of

shared alleles with the test going from 0 (full isolation) to 0.5 (full hybridization) and

therefore it can also cover case ii. The two tests differ in how they measure significance, using

jackknifing or bootstrapping in Patterson's D and normal approximation in HyDe. Of late,

another site-based test was developed using χ^2 to test for deviation of parity between the three

topologies as in case i (Sayyari and Mirarab 2018). A unified test that can test the prevalence

of each of the four categories across the genome is still lacking.

Reminiscent to the problem of soft and hard polytomies, wherein the statistical power

of a phylogenetic analysis largely depends on the size of a locus, interpreting reticulation is

largely influenced by how a locus is defined. For example, at the species level, an analysis on

a whole concatenated alignment of a quartet should have a greater power (*i.e.* a total evidence

approach). However, sites are often physically linked and evolutionarily-dependent.

Consequently, whole genome analyses may be biased towards genomic regions with more

variable sites (Martin et al. 2015). Therefore, phylogenetic analyses are usually conducted at

individual loci under the multispecies coalescent model, which was shown to outperform

concatenation in multiple groups of animals (Jiang et al. 2020). In theory, a window size of

~100-kb long is appropriate to overcome the non-independence of sites under realistic

recombination rates (Pease and Hahn 2015), but the definition of such windows would largely

depend on the taxonomic scale of the analysis. For example, at shallow phylogenetic depths, it

is customary to use a pseudo-reference genome approach, whereas reads from multiple

species are aligned to a reference genome. Here, the distance from the reference genome

species can greatly influence the quality of mapping and subsequent analyses due to potential

gene duplications and rearrangements (XXX). Alternatively, single-copy orthologous genes,

such as those that have been popularized by the BUSCO program, are used (XXX). However,

such gene are presumably under strong selection, with the relaxation of selection in some

lineages may conflate introgression estimates (Frankel and Ané 2023). Proper estimation of

reticulation therefore requires a good understanding of how different locus-definition

strategies influences incongruence analyses.

Phylogenetic conflicts have been reported for the jumping fly *Drosophila saltans*

species group, a clade of the subgenus *Sophophora* with 23 Neotropical species (Magalhães

1962; Bächli 2024). The group retains its name from the peculiar "jumping" habit of its

larvae; *the larva seizes its posterior end with its mouthhooks, and stretches. The hooks pull*

loose suddenly, the larva straightens with considerable force, and as a result is thrown

several inches into the air" (Sturtevant 1942). The group was divided into five species

subgroups, namely, *saltans, parasaltans, cordata, elliptica* and *sturtevanti* subgroups, mostly

on the basis of male genitalia (Magalhães and Björnberg 1957). Although the monophyly of

the subgroups has been confirmed by different phylogenetic methods, the relationships among

and within them are not. Hypothesis for their evolutionary relationships have been proposed

using different methods and different morphological characters (Magalhães and Björnberg

1957; Throckmorton 1962; Throckmorton and Magalhães 1962; O'Grady et al. 1998; Yassin

2009; Souza et al. 2014; Roman et al. 2022), chromosome polymorphism (Bicudo 1973a),

reproductive isolation (Bicudo 1973b; Bicudo and Prioli 1978; Bicudo 1979), protein

polymorphism (Nascimento and Bicudo 2002) and gene sequences (Pélandakis and Solignac

1993; O'Grady et al. 1998; Rodríguez-Trelles et al. 1999; Castro and Carareto 2004; de Setta

et al. 2007; Roman et al. 2022). The evolutionary relationships proposed are summarized in

Supplementary Table S1.

Unlike other species groups in the subgenus *Sophophora*, such as the *melanogaster*,

obscura and *willistoni* groups, genomic resources and genetic investigations in the *saltans*

species group are scarce. Indeed, only four genomes have been sequenced and assembled to

date (Kim et al. 2021). To bridge this gap and to test for the extent of phylogenetic conflicts,

we sequenced and assembled genomes for 15 species with representatives from the five

subgroups. Phylogenetic analyses using well-conserved genes resolved the evolutionary

relationships among the subgroups but also highlighted conflicts between X-linked, autosomal

and mitochondrial loci. To test how each of the four incongruence categories prevails across

the genome, we devised a new χ^2-based test that we called 2A2B. The test uses pairwise

comparisons of the three topologies proportions shown in Figure 1. We applied this test to

sets of conserved orthologous genes and to long syntenic blocks with conserved collinearity

across the Neotropical *Sophophora*, as well as to 100-kb long windows from pseudo-reference

genomes aligned to either an ingroup (*D. sturtevanti*) or outgroup (*D. willistoni*) species. We

found reticulation levels to differ among the datasets and the subgroups, and to correlate with

rate of speciation and historical biogeography in the *saltans* group.

Results

Short-read assembly of 17 genomes recovered 90% of BUSCO genes

We sequenced using short-read Illumina approach 17 whole genomes from 15 species

collected across various locations in the Neotropical region. Genome size, estimated from 21-

kmer frequency spectrum using GenomeScope 2 (Ranallo-Benavidez et al. 2020), ranged

from 154.0 to 356.8 Mb. Our de novo assemblies using MaSuRCA (Zimin et al. 2013)

resulted in genome lengths ranging from 177.5 to 287.7 Mb, with N50 values ranging from 2

to 92 Kb (Supplementary Table S2). To evaluate the completeness of our assembled genomes,

we searched for single-copy genes (SCG) using BUSCO (Simão et al. 2015). We found that

over 90% of the searched genes were complete for all of the genomes (Supplementary Table

S2). Kim et al. (2021) assembled using both short Illumina and long Nanopore reads the

genomes of four *saltans* group species, all of which we have independently sequenced.

Whereas their assemblies' contigs were much longer, with N50 ranging from 2 to 6 Mb, the

BUSCO score for the same set of species did not largely differ (98% vs. 95-96% in our study;

Supplementary Table S2).

Muller elements analysis resolves relationships between the subgroups and unravels a minor

X-autosomal conflict in the sturtevanti subgroup

Phylogenomic analyses were performed using 2,159 SCG shared across all species.

Gene trees, inferred for each SCG using maximum-likelihood in IqTree2 (Minh et al. 2020)

produced 1,263 distinct topologies, with 206 of them found more than once. To test if SCG

chromosomal position may underlie the discrepancies in gene trees, we localized each SCG to

its corresponding Muller element according to the position of its *D. melanogaster* ortholog

identified by Blast (Camacho et al. 2009). As a result, we generated five independent datasets,

each corresponding to the Muller elements A, B, C, D, and E, comprising 337, 370, 425, 419,

and 568 SCGs, respectively. These datasets were then used to reconstruct the species trees

using the multi-species coalescent model, and the genes within them were concatenated for

Bayesian and Maximum Likelihood phylogenetic inferences.

The trees generated by the 5 data sets showed very similar topologies with well

supported nodes either in the Bayesian Inference implemented in BEAST (Bouckaert et al.

2019), the maximum-likelihood implemented in IqTree2, or for the multi-species coalescent

model analysis implemented in ASTRAL-III (Zhang et al. 2018) (Supplementary Figures S1,

S2 and S3). The *parasaltans* subgroup was placed as sister to all other subgroups, followed by

the emergence of the *sturtevanti* subgroup. The *cordata* and *elliptica* subgroups showed a

close relationship, and were sister to the *saltans* subgroup. The only discrepancy between the

topologies was the placement of *D. lehrmanae*, a newly discovered species in the *sturtevanti*

subgroup (Madi-Ravazzi et al. 2021). For *D. lehrmanae*, while maximum-likelihood and

multi-species coalescent analyses reported lack of branch support for multiple trees

(Supplementary Figures S2 and S3), Bayesian inference recovered well supported branches

and two topologies (Figure 2A and Supplementary Figure S1). The two distinct topologies

distinguished the Muller elements forming the X chromosome (elements A and D who are

fused in a fusion shared by the Neotropical *Sophophora*, *i.e.* the *saltans* and *willistoni* groups

(Sturtevant and Novitski 1941; Dobzhansky and Pavan 1943; Cavalcanti 1948), and the

Muller elements representing autosomal chromosomes (elements B, C and E).

The published genome of *D. prosaltans* (Kim et al. 2021) did not group with the

genome of this species sequenced by us, instead it grouped with *D. saltans*. The genome

previously published comes from a line collected in El Salvador in 1957. According to

Magalhães' (1962) detailed morphological revision of multiple geographical specimens of the

saltans group, the sampling site of this particular strain is outside the geographical range of *D.*

prosaltans, but within the expected range of *D. saltans*. Furthermore, the *D. saltans* and *D.*

prosaltans lines used in our study underwent thorough morphological analyzes (Souza et al.

2014; Roman and Madi-Ravazzi 2021), indicating that the lines we used were accurately

identified. Therefore, it is most likely that the previously sequenced *D. prosaltans* strain from

El Salvador was misidentified and we consider it here to belong to *D. saltans*.

Mitogenomes show cytonuclear conflicts in the sturtevanti *and* saltans *subgroups*

We assembled mitochondrial genomes for the 15 *saltans* group species using MitoZ

(Meng et al. 2019). We did not use the previously assembled four strains since several

mitochondrial scaffolds were likely removed in those assemblies (Kim et al. 2021). We

conducted phylogenetic analysis on the aligned mitogenomes genes using both IqTree2 and

MrBayes (Ronquist et al. 2012). Overall, the mitochondrial trees matched the topology of the

nuclear gene trees regarding the inter-subgroup relationships. However, three major

discrepancies were identified (Figure 2B). First, the position of *D. lehrmanae* within the

sturtevanti subgroup did not agree with either the X or autosomal SCG topologies, proposing

a topology wherein *D. lehrmanae* is a sister species of *D. sturtevanti* (a topology recovered

once in multi-species coalescent analysis (Muller element C, Supplementary Figure S3) and

maximum likelihood (Muller element B, Supplementary Figure S2)). Second, whereas the

mitochondrial tree recovered the monophyletic relationship between the *elliptica*, *cordata* and

saltans subgroups, the position of *D. neocordata* (*cordata* subgroup) differed, being sister to

the three species of the *elliptica* subgroup in the nuclear trees and to the six species of the

saltans subgroup in the mitochondrial tree. Third, whereas nuclear trees recovered three

lineages within the *saltans* subgroup, namely*, austrosaltans*, *nigrosaltans-pseudosaltans*, and

septentriosaltans-prosaltans-saltans, only two lineages are revealed by the mitochondrial tree.

Each of the mitochondrial clades involved one species from otherwise sister species in the

nuclear trees, *i.e. D. nigrosaltans* and *D. saltans* in one clade and their respective closely-

related species *D. pseudosaltans* and *D. prosaltans* in the other clade, suggesting that multiple

cytoplasmic introgression events might have occurred in this subgroup (Figure 2B). Because

D. saltans and *D. prosaltans* are reported as closely related species in the nuclear trees and are

separated in the two mitochondrial clades, the two mitoclades were called *S* and *P*,

respectively. Remarkably, the branching order in the *P* mitoclade is congruent with the

nuclear tree, indicating that the *S* mitoclade is most likely of reticulate origin, probably

reflecting an older introgression from *D. austrosaltans* to *D. nigrosaltans* and a more recent

introgression between *D. austrosaltans* and *D. saltans*.

The extent of reticulation differs between assembly- and pseudo-reference-based approaches

248   To quantify the extent of introgression at a genome-wide scale, we analyzed allele

distribution of bi-allelic phylogenetically informative sites in 28 species quartets using

Patterson's *D* estimate of the standard ABBA-BABA test. We used four distinct locus-

defining strategies to explore how strategy choice can influence estimates of *D*. Two

strategies relied on our *de novo* genome assemblies. The first strategy included the use of

single-copy genes (SCG) defined by BUSCO (2,159 loci). The second strategy identified

syntenic blocks equal to or greater than 50 kb-long with collinearity across the 15 *saltans*

assemblies and *D. willistoni*, *i.e.* across Neotropical *Sophophora* (1,797 loci). The third and

fourth strategies used a pseudo-reference approach, wherein the reads of each *saltans* group

species were mapped to an ingroup (*D. sturtevanti*, 2,328 loci) or an outgroup species (*D.*

willistoni, 1,863 loci) and a pseudo-reference was inferred by replacing variant sites in the

reference genome by the ones in the reads (as in Mai et al. 2020). In these pseudo-reference

strategies, a locus was defined as a 100 kb-long window of the reference genome. To avoid

the influence of linked loci from highly-variable regions on genome-wide estimates, we

randomly sampled 20 sites from each locus with ≥20 evaluated sites, and repeated resampling

for 100 reiterations for each locus-defining strategy.

For a bi-allelic site, the standard ABBA-BABA (Patterson's D) test considers a true

species tree exists, with site distribution AABB, and then evaluate the deviation from parity of

the two discordant configurations ABBA and BABA. We found that genome-wide absolute

Patterson's D estimates, $i.e.$ ($\sum$ABBA – $\sum$BABA)/($\sum$ABBA + $\sum$BABA), greatly differed

among the four strategies (Figure 3A). Synteny-based strategy had the lowest $|D|$ across the

28 quartets ($|D| = 0.025 \pm 0.004$) compared to BUSCO ($|D| = 0.066 \pm 0.014$, Student's t $P <$

0.01), $D.$ $sturtevanti$ pseudo-reference ($|D| = 0.176 \pm 0.025$, Student's t $P < 4.77$ x 10^{-6}), and

$D.$ $willistoni$ pseudo-reference ($|D| = 0.329 \pm 0.053$, Student's t $P < 5.97$ x 10^{-6}). So, of the

four strategies, the use of syntenic blocks is the most conservative.

Comparisons of locus-defining strategies identify sources of errors in whole-genome

Patterson's D *estimates of introgression in species quartets*

To gain further insights on the reasons of this discrepancy between the four locus-

defining strategies, we separately investigated assembly- and pseudoreference-based

approaches. The BUSCO approach mainly differs from the synteny-based approach in having,

on average, shorter sequence since many loci are ≤50-kb long. Short sequences usually shows

faster lineage-specific evolutionary rates (XXX), which can increase variance and also induce

biases on Patterson's D due to the violation of the molecular clock assumption (Frankel and

Ané 2023). Besides, the BUSCO analysis involves an additional step, which is the annotation

of the conserved protein-coding BUSCO genes. To account for these possible sources of

errors, we counted for each lineage the number of singletons distinguishing the outgroup from

the three ingroup species ($i.e.$ BBBA sites), as well as the singletons distinguishing between

the two most closely-related species in the quartet as defined from the phylogenetic analyses

above ($i.e.$ BAAA and ABAA). we called this test 3A1B, and it is reminiscent to the classical

Tajima's (1993) Relative Rate Test. Under the assumption of a constant molecular clock or

the absence of mis-annotation of a BUSCO gene in one of the two sister species both BBBA –

BAAA and BBBA – ABAA should be > 0. We focused on the *elliptica* subgroup with *D.*

saltans as an outgroup (*i.e.* (((*emarginata*,*neoelliptica*),*neosaltans*),*saltans*)) quartet, since

this quartet showed a strong discrepancy between the two assembly-based approaches, with

genome-wide *D* being 0.25 for BUSCO and -0.08 for synteny-based strategy. By applying the

3A1B test, we found that 37 BUSCO loci violated its assumption, all in the negative direction

for *D. emarginata*, *i.e.* BAAA > AAAB. By excluding these loci, Patterson's *D* dropped from

0.25 to 0.04. By contrast, only 5 loci violated the assumption in the synteny-based strategy.

Their exclusion did not change Patterson's *D* estimate which remained -0.08. We then applied

the 3A1B test for all locus-defining approaches, excluding loci that violated its basic

assumption in each quartet. The genome-wide absolute Patterson's *D* estimates dropped from

0.066 ± 0.014 to 0.030 ± 0.004 for BUSCO genes (Student's *t* $P < 0.05$) and from $0.025 \pm$

0.004 to only 0.023 ± 0.004 for synteny-based loci (Student's *t* $P = 0.76$). After correction,

genome-wide estimates from both BUSCO and synteny loci did not significantly differ

(Student's *t* $P = 0.28$), even if synteny estimates remained slightly lower (Figure 3B).

For pseudo-reference approaches, excluding rapidly evolving loci through the 3A1B

test may not be enough, since additional sources of errors related to the quality and depth of

mapping as well as reference genome biases constitute additional sources of errors. We

therefore reconducted the analyses using more stringent parameters, only retaining loci with a

mapping quality >20 and a depth >10 and <100 and excluding all within-species polymorphic

sites. These parameters dramatically reduced the number of evaluated loci to only 17% and

4% in the ingroup (*D. sturtevanti*) and outgroup (*D. willistoni*) reference genome approaches.

Estimation of genome-wide absolute Patterson's *D* to 0.101 ± 0.023 (Student's *t* $P < 0.05$) and

0.102 ± 0.017 (Student's *t* $P < 2.9 \times 10^{-4}$) for the ingroup and outgroup references,

respectively. This means that whereas ingroup and outgroup approaches were greatly different

before the application of the stringent criteria (Student's t $P < 0.05$), absolute Patterson's D

did not significantly differ between the two approaches under these criteria (Student's t $P =$

0.95). For the abovementioned *elliptica* subgroup example, whereas before the stringent

criteria Patterson's D were 0.02 and -0.71 for ingroup and outgroup approaches, these values

dropped to -0.005 and -0.27 in the two approaches, respectively. However, correlation across

quartets was strong (Pearson's $r = 0.56$, $P = 0.002$), indicating that distinct criteria may be

applied to improve estimate depending on the distance from the reference genome species,

with *D. willistoni* being more distant and having a more fragmented genome assembly.

The 2A2B test unravels lower signal for introgression when syntenic blocks are used

To consider cases where a true species tree cannot be inferred, we designed a test that

we call 2A2B. The test is an extension of the standard ABBA-BABA test in that it does not

only test the significance of deviation from parity of the ABBA and BABA patterns, but also

test the deviation from parity of each configuration to the BBAA pattern. Therefore, the test

allows classifying each species quartet into one of the four categories along the trifurcation-

bifurcation continuum given in Figure 1. We applied the test at both the genome-wide level

after corrections using the 3A1B test and mapping criteria (see above).

We found that for all locus-defining strategies, bifurcation categories (iii and iv)

predominated (Figure 4A). Only five exceptions were observed. One incidence of category ii

(incomplete trifurcation) was observed in the BUSCO strategy concerning the position of *D.*

lhermanae relative to both *D. sturtevanti* and *D. dacunhai*. Indeed, the position of *D.*

lhermanae was the only conflicting result in the phylogenetic analyses based on BUSCO

genes and it also showed a conflict with mitochondrial analysis (Figure 2). Four incidences of

category i (complete trifurcation) were observed in outgroup pseudo-reference strategy. Of

which, two concerned the position of *D. lehrmanae* to both *D. sturtevanti* on the one hand and

to each of the two sister species *D. dacunhai* and *D. milleri* on the other hand. The other two

complete trifurcation cases concerned the relationships between the three lineages of the

saltans subgroups, *i.e. austrosaltans*, *nigrosaltans-pseudosaltans*, *septentriosaltans-*

prosaltans-saltans, in cases where species of the different mitoclades where found analyzed

together, *i.e. D. pseudosaltans* and *D. saltans* or *D. nigrosaltans* and *D. septentriosaltans*

(Figure 2).

Category iv (complete bifurcation), which is the most consistent with a true species

tree, was the most frequent in synteny-based strategy (Figure 4). This was followed by both

the BUSCO and the outgroup pseudo-reference strategies but the difference in category iv

frequency was not significant (Fisher's exact test of category iv vs. all other categories).

However, with only four exceptions, ingroup pseudo-reference approach identified category

iii (incomplete bifurcation), which mostly indicates cases of introgression, in every possible

quartet on the tree (Fisher's exact test between synteny-based and ingroup pseudo-reference

strategy $P < 1 \times 10^{-4}$). The total number of bi-allelic sites (20 per locus) that were used in the

genome-wide analysis differed among the four strategies, averaging 22,893 (BUSCO) and

21,306 (synteny) assembly-based approaches, and 35,747 and 5,308 sites for the ingroup *D.*

sturtevanti and the outgroup *D. willistoni* pseudo-reference approaches, respectively. The low

number for the *D. willistoni* pseudo-reference is expected since increasing mapping quality

criteria on the genome of this distant species would only favor strongly conserved loci. For

the *D. sturtevanti* pseudo-reference, applying the same mapping quality criteria has likely

maintained less conserved loci. Although we have excluded heterozygous sites from every

species to reduce mapping biases of the reference allele, such biases may even extend to

homozygous sites linked to the excluded heterozygous site in the same read. Universal

mapping criteria that can reduce mapping biases independent from the reference genome do

not exist and mapping biases remain difficult to detect (XXX). However, these biases are the

most likely explanation of the extensive signal of introgression in the ingroup *D. sturtevanti*

pseudo-reference. To test the effect of mapping quality on introgression estimates, we applied

the 2A2B test at the locus level before and after applying mapping quality thresholds. In

agreement with our assumptions and the findings on Patterson's *D* (Figure 3), the proportion

of loci supporting categories ii and iii, *i.e.* categories that are the most associated with

reticulate evolution, did not exceed ~7% in assembly-based strategies, but they dropped after

application of mapping quality criteria from 46% to 11% and from 34% to 4% for *D.*

sturtevanti and *D. willistoni* pseudo-reference genomes strategies, respectively

(Supplementary Figure XXX). Ingroup pseudo-reference approaches, while providing more

data, may not reflect the true phylogenetic relationships between the species, hence inflating

the estimation of the extent of introgression.

Historical biogeographical events coincided with episodes of low phylogenetic resolution

Phylogenetic analyses and assembly-based 2A2B test identified three parts on the

species tree with evidence for incomplete bifurcation (*i.e.* category iii): at the inter-subgroup

level in any combination that involved representatives from at least two subgroups of the

cordata, *elliptica* and *saltans* subgroups, at the base of the *sturtevanti* subgroup, and in the

saltans subgroup in quartets involving species belonging to distinct mitoclades. Whereas

category iii is usually interpreted as evidence of introgression following secondary contact,

other biological processes such as ancestral population structure at the time of phyletic

radiation or asymmetry of evolutionary rates due to responses to distinct adaptive pressures

(XXX). Because all these processes can associate with major geographical changes affecting

species distribution, ecology and frequency of secondary contacts, a knowledge of the

historical biogeography of the clade under study is crucial for the understanding of the

processes that have likely led to phylogenetic conflicts.

Like phenotypic traits, ancestral geographical distribution can be inferred on

phylogenetic trees from present day geographical data (XXX). Therefore, we mapped current

distribution of the studied species on the Bayesian X tree that was dated in reference to recent

family-wide estimates (Suvorov et al . 2021; see Methods). The historical biogeography

supported an origin of the *saltans* group around 16 million years (myr) ago in the north-west

of South America, which currently corresponds to the Amazonian region (Figure 5). At that

time, this region was dominated by a vast wetland environment known as the 'Pebas System.'

The first radiation episode, *i.e.* the split between the ancestors of the *cordata*, *elliptica* and

saltans subgroups occurred circa 10 myr ago at this region, coinciding with the formation of

the Amazon River and its huge fluvial networks that led to the transformation of the Pebas

wetland into the dense and diverse Amazonian forests (Hoorn et al. 2022). The second

episode occurred nearly 3 myr ago, which correlates with the geological formation of the

isthmus of Panama (O'Dea et al. 2016), and the time of a connection between the Western

Amazonian and the Southern Atlantic rainforests during the Pliocene (Batalha-Filho et al.

2013; Ledo and Colli 2017). This episode involved the ancestors of the *nigrosaltans-*

pseudosaltans and *septentriosaltans-prosaltans-saltans* clades. Indeed, the first clade has a

northern to central representative (*D. nigrosaltans*) and a southern species (*D. pseudosaltans*),

whereas the second clade has a northern representative (*D. saltans*) and two central and

central to southern species (*D. septentriosaltans* and *D. prosaltans*, respectively). The third

radiation episode involves the ancestor of the *sturtevanti* subgroup and also occurred in the

late Pliocene, ca. 2.5 myr ago, at the likely time of a connection between the Western

Amazonian and the Southern Atlantic rainforests.

Discussion

Towards a comprehensive phylogeny of the saltans species group

A large number of *Drosophila* genomes have been sequenced and used in

phylogenetic analyses (Khallaf et al. 2021; Kim et al. 2021; Li et al. 2022; Suvorov et al.

2022), but studies with comprehensive sampling of nearly all species in a group remain

relatively uncommon (Mai et al. 2020; Conner et al. 2021; Yusuf et al. 2022; Moreyra et al.

2023). Despite minor inconsistencies, our phylogenomic analysis of 15 species of the

Drosophila saltans species group produced a consistent picture of the relationships between

the five subgroups of this clade. All X, autosomal and mitochondrial phylogenies showed the

parasaltans subgroup as the first to diverge, followed by the *sturtevanti* subgroup, and later

by a clade comprising the *cordata*, *elliptica* and *saltans* subgroups, in which the position of

the *cordata* subgroup differed between nuclear and mitochondrial trees. This general picture

has not been previously proposed despite the tremendous number of phylogenetic

investigations of this group (Magalhães 1962; Throckmorton 1962; Throckmorton and

Magalhães 1962; O'Grady et al. 1998; Rodríguez-Trelles et al. 1999; Castro and Carareto

2004; de Setta et al. 2007; Yassin 2009; Souza et al. 2014; Roman et al. 2022; see

Supplementary Table S1).

While our analysis has shed light on the intricate evolutionary dynamics within this

clade, further sampling holds the potential to provide a more comprehensive understanding

into this complex evolutionary history. For example, the inclusion of *D. subsaltans*, *D.

lusaltans*, *D. cordata*, and *D. rectangularis* through whole-genome sequencing promises to

provide insight onto unresolved phylogenetic questions raised from previously published

observations on reproductive isolation and morphology (Magalhães 1962; Bicudo and Prioli

1978). These questions include the monophyly and positioning of the *parasaltans* and *cordata*

subgroups. Additionally, the inclusion of the insular species *D. lusaltans* which presents low

reproductive isolation (Bicudo 1973b), can bring new insights into the reticulation evolution

in the *saltans* subgroup. The *saltans* subgroup showed the most dramatic signal of cyto-

nuclear discordance and reticulated evolution. Bicudo (1973b) investigated reproductive

isolation among the seven then described species of this subgroup, and in a remarkably partial

agreement with our nuclear phylogenomic trees, she concluded that *D. pseudosaltans*, *D.*

nigrosaltans and *D. austrosaltans* showed more basal relationships compared to *D. lusaltans*,

D. septentriosaltans, *D. prosaltans* and *D. saltans*, *i.e.* the latter species show less

reproductive isolation. Indeed, nearly all crosses among the last four species produce fertile

females with some even producing fertile females and males (Bicudo 1973b). This behavioral

porosity largely agrees with the high incidence of reticulate evolution we report here for this

subgroup.

Two widespread species of the *saltans* subgroup, *D. saltans* and *D. prosaltans*, show a

peculiar geographical disjunction, with the former species being restricted to Central America

whereas *D. prosaltans* is widespread in South America south of Costa Rica. The

discrimination between strains belonging to each species has long been erroneous

(Dobzhansky 1944; Mayr and Dobzhansky 1945; Spassky 1957; Magalhães 1962) and we

showed here that their misidentification persists even in the genomic era (Kim et al. 2021;

Suvorov et al. 2022). Interestingly, Bicudo (1973b) provided evidence for reproductive

reinforcement between these two sister species; sympatric populations in their junction zone

in Costa Rica demonstrated stronger reproductive isolation than allopatric populations of both

species. We have only included one to a few geographical lines from each species and a

broader sampling to investigate the extent of their reproductive isolation and genome porosity

is strongly needed.

Intra- and inter-genomic conflicts impact the inference of phylogenetic patterns

Concatenation helped recovering a sex chromosome versus autosome conflict, similar

to the one described by Mai et al. (2020) for the *Drosophila nasuta* subgroup. Like these

authors, this conflict was limited to a single part of the tree, *i.e.* the relationship of *D. pulau* to

D. sulfurigaster sulfuricaster and *D. s. bilimbata* in the *nasuta* group and the placement of *D.*

lehrmanae in the *sturtevanti* subgroup. The peculiarities of sexual chromosomes such as their

lower effective population size, different recombination and mutation rates, and greater

exposition to natural selection when found in hemizygosity, lead to higher rates of adaptive

evolution of sexual-linked genes compared with autosomal genes (*i.e.* faster-X evolution) and

also to the disproportional accumulation of genes related to reproductive isolation and

Dobzhansky-Muller hybrid incompatibilities (*i.e.* Haldane's rule) (Charlesworth et al. 2018).

Besides, low recombination rate underlies stronger linked selection effects against

introgressed ancestry. Altogether, those characteristics are thought to be responsible for the

resistance to hybridization in the sexual chromosomes (Ellegren 2009; Qvarnström and Bailey

2009; Sankararaman et al. 2016; Charlesworth et al. 2018; Seixas et al. 2018; Mai et al. 2020;

Matute et al. 2020; Moran et al. 2021; Reilly et al. 2022; Skov et al. 2023; but see David et al.

2022).

The second conflict regards a significant disagreement between mitochondrial

(mtDNA) and nuclear data. Discordance between nuclear and mitochondrial genomes is a

well-documented phenomenon in the tree of life as highlighted by Toews and Brelsford

(2012). Several characteristics of mtDNA, such as being haploid and uniparentally inherited,

resulting in a fourfold reduction in effective population size when compared with autosomal

chromosome loci, affect its evolution. Cytoplasmic introgression has long been recognized in

Drosophila (Solignac et al. 1986; Ballard 2000; Llopart et al. 2014). In a recent population

study within the *willistoni* group, multiple mitochondrial introgressions were observed in *D.*

paulistorum populations (Baião et al. 2023). These included an ancient introgression with a

highly divergent mitochondrial type, followed by more recent events. While nuclear-

mitochondrial incompatibilities likely posed challenges, the study also suggested two possible

alternatives to overcome these challenges: a selective advantage provided by the

mitochondrial type itself. Or a non-selective factor, such as *Wolbachia,* a bacteria known to

modify the reproduction of its host, could facilitate a mitochondrial type fixation (Baião et al.

2023). Although, interesting results have been reported from population approaches, conflicts

between nuclear and mitochondrial genomes have not been addressed in recent phylogenomic

analyses in the Drosophilidae (Mai et al. 2020; Khallaf et al. 2021; Suvorov et al. 2022; Yusuf

et al. 2022). The disagreement was particularly evident for the *saltans* subgroup, where it was

most likely of recent origins, separating species that have diverged only 0.7 myr ago, *i.e. D.*

nigrosaltans and *D. pseudosaltans*. Remarkably, the two mitotypes P and S do not correlate

with the degree of reproductive isolation inferred by Bicudo (1973b), contrary to nuclear tree,

indicating that cytoplasmic introgression in the *saltans* subgroup did not contribute to the

evolution of reproductive isolation in this clade.

Of the three subgroups for which multiple species were sequenced, the *saltans*

subgroup had the highest incidence of incongruences inferred by the 2A2B test. In this

subgroup, multiple large chromosomal inversions are known to be shared among closely-

related species (Dobzhansky and Pavan 1943; Cavalcanti 1948; Bicudo 1973a; Bicudo and

Prioli 1978) and evidence for balancing on ancestral inversion polymorphism has been

demonstrated in a number of cases (Bicudo 1973a). For example, one autosomal inversion is

polymorphic in *D. austrosaltans*, with one arrangement being homozygous in *D.*

septentriosaltans and *prosaltans*, and the alternative arrangement homozygous in the

remaining species including *D. saltans*. Such an inversion has likely been inherited from a

polymorphic ancestor of the *saltans* subgroup, and maintained polymorphic in the ancestor of

the *septentriosaltans-prosaltans-saltans* clade before different arrangements became fixed in

each of these species, with one arrangement being already fixed in the ancestor of the

nigrosaltans-pseudosaltans clade. Alternatively, the inversion might have originated in the

ancestor of the *septentriosaltans-prosaltans-saltans* clade, then it was introgressed in *D.*

austrosaltans, whereas the ancestral non-inverted configuration being introgressed on its own

in *D. saltans*. However, this introgression scenario is less parsimonious, especially given the

geographical isolation of *D. saltans* from the remaining species of the subgroup. Whether the

high degree of incongruences in the *saltans* subgroup is associated with large ancestral

inversions potentially absent in other bifurcating clades would require the future generation of

chromosome-level assemblies for multiple *saltans* group species.

The 2A2B test on syntenic blocks helps distinguishing "soft" and "hard" introgressions

Our analyses reveal that the choice of locus defining scheme can affect the degree of

phylogenomic uncertainty. Indeed, most studies that have provided strong evidence for

introgression using site-specific patterns comparisons have usually aligned reads from

multiple species to a single reference assembly (cf. studies compiled by Dagilis et al. 2021).

Whereas such an approach would increase the power (Martin et al. 2015; Pease and Hahn

2015), it also introduces biases due to paralogy, misalignments or absence of collinearity

among species, that are now known to bias phylogenetic inference itself (Valiente-Mullor et

al. 2021; Rick et al. 2023). Therefore, the most common practice now is to use sets of clade-

specific single-copy orthologs, such as the BUSCO genes, in inferring phylogenies (Manni et

al. 2021). Consequently, some recent introgression studies have focused on such genes. For

example, in a study of 155 genomes covering a wide range of drosophilid lineages, Suvorov et

al. (2022) analyzed these conserved genes. While finding evidence for widespread

introgression, they also showed that their discordance estimates were highly sensitive to the

length of the analyzed genes as well as by the slightest relaxation of selective pressures. We

attempted here to overcome this limitation by including larger syntenic blocks, which in

addition to conserved protein-coding sequences should also include presumably neutral

intronic and intergenic sites. Whereas, this approach showed the lowest $|D|$, indicating that

patterns predicted by introgression do not exceed 7% of the genome, it led to only 1.5-fold

increase in the number of analyzed sites, most likely because our assemblies were based on

only short reads. With the continuing progress in whole-chromosome sequencing techniques

and assembly-to-assembly alignment tools, a better understanding of the extent of probably

true or "hard" introgression events from artificial or "soft" ones may be attained.

Phyletic radiations occur when isolation barriers simultaneously and rapidly evolve

between multiple populations of a species with a broad geographical or ecological breadth.

Consequently, genetic lineages will be incompletely sorted in the emerging species leading to

a lack of phylogenetic signals. This situation usually occurs following a rapid environmental

change or expansion into a new range with new empty niches. Indeed, our historical

biogeography approach shows that such situation might have favored the episodic signals of

low phylogenetic signals in the *saltans* group since the episodes coincided with either the

transition from wetlands into Amazonia forests in the Late Miocene or the connection of the

Amazonian forest with either Central American or the Southern Atlantic forests in the

Pliocene. These events have been proposed to be responsible for a wide range of rapid

radiations in Neotropical taxa, such as butterflies, amphibians and birds (Batalha-Filho et al.

2013; O'Dea et al. 2016; Ledo and Colli 2017; Hoorn et al. 2022). The dating and the

analyses of the extent of phylogenetic discordance in the genomes of such clades are strongly

needed to identify the impact of these events on Neotropical diversification.

Materials and Methods

Sample collection, whole genome sequencing and assembly

We performed whole genome pool sequencing on female flies from 15 different

species from the *saltans* group, as well as three populations of *D. sturtevanti*. The specimens

used for sequencing were obtained from one or multiple strains, and detailed information

regarding the number of individuals and their collection locations can be found in

Supplementary Table S2. For all species, DNA extraction, PE library preparation using

NEBNext Protocol, and 75x coverage double-paired Illumina Novoseq sequencing were

conducted by End2End Genomics LLC, Davis, California. For all species except *D.*

neocordata, we assembled the genome using the Maryland Super Read Cabog Assembler

(MaSuRCA) (Zimin et al. 2013), which utilizes both the de Bruijn graph and overlap-layout-

consensus (OLC) methods to generate super-reads. K-mer based genome size was estimated

using KMC3 (Kokot et al. 2017) and GenomeScope v.2.0 (Ranallo-Benavidez et al. 2020). To

assess the assembly's completeness, we searched for single-copy-genes (SCG) using default

parameters in Busco5 (Simão et al. 2015) with the diptera_odb10 database (Kuznetsov et al.

2023). In addition to the sequenced flies, we also assembled Illumina reads the genomes of *D.*

saltans, *D. neocordata*, *D. prosaltans* and *D. sturtevanti* published by Kim et al. (2021)

(assembly numbers ASM1890357v1, ASM1890361v1, ASM1815127v1 and

ASM1815037v1, respectively), and we used the *D. neocordata* assembly published by Baião

et al. (2023) in all analyses.

Phylogenomics: Nuclear single-copy-genes

For phylogenomics analysis, SCG searches were carried out using 3,285 SCG from

diptera_odb10 database on Busco5 (Simão et al. 2015). SCG present in all species were kept

and aligned using the L-INS-i method implemented on MAFFT (Katoh and Standley 2013)

(mafft --localpair --maxiterate 1000 –adjustdirection).

Genomic data of different species of *Drosophila* support that genes tend to be situated

within the same Muller element across multiple species (see Schaeffer 2018). Taking this

gene linkage into account, we reconstructed five independent datasets (Muller elements A-E),

each comprising all SCG found in the respective Muller element. To achieve this, we

performed a tBlastn search against the *D. melanogaster*, and subsequently we concatenated

genes found within the same Muller element. Phylogenetic trees were then constructed using

maximum likelihood and Bayesian methods implemented in the softwares IQ-TREE (Nguyen

et al. 2015) and BEAST (Bouckaert et al. 2019), respectively. Additionally, maximum-

likelihood trees were generated for each gene, the output tree from each Muller element data-

set were used to reconstruct to species trees, using multi-species coalesce model implemented

in ASTRAL-III (Zhang et al. 2018). Branch support was assigned using 1,000 ultrafast

bootstrap in IQ-TREE (Hoang et al. 2018), posterior probability in BEAST and local posterior

probability in ASTRAL-III (Zhang et al. 2018).

Phylogenomics: Mitochondrial Genome

Mitochondrial genomes were assembled and annotated with MitoZ (Meng et al. 2019),

with the Megahit assembler (Li et al. 2015). In order to ensure the exclusion of nuclear-

embedded mitochondrial DNA sequences within the assembly, a strategic approach was

taken. Considering that mitochondrial reads are found in higher frequency than nuclear-

mitochondrial DNA sequences, the read subsampling were set to 0.5 gigabases (--

data_size_for_mt_assembly 0.5). The genes obtained from the mitochondrial genome were

aligned using the MAFFT alignment tool with the --auto parameter due to the close similarity

between sequences. Subsequently, the aligned genes were concatenated into a dataset for

phylogenetic analysis. The concatenated dataset served as the basis for reconstructing

phylogenetic trees using both Maximum Likelihood (ML) implemented in IQ-TREE (Nguyen

et al. 2015) and Bayesian Inference (BI) in BEAST (Bouckaert et al. 2019). Branch support

was assigned using 1,000 ultrafast bootstrap in IQ-TREE (Hoang et al. 2018) and posterior

probability in BEAST.

Identification of syntenic blocks

The 20 genomes of the *saltans* group, the 15 sequenced here and the 5 published by Kim et al.

(2021) and Baião et al. (2023) were preliminary annotated with Miniprot (miniprot -Iut16, Li

2023). The primary objective of this annotation was to accurately map proteins from the

robust and reliable genome annotation of *D. willistoni* (Clark et al. 2007). After the protein

mapping, the predicted gene loci were assessed to identify syntenic blocks present in the

Neotropical *Sophophora* (comprising *D. willistoni* and *saltans* group). The identification of

these blocks was based on gene order and orientation, achieved using an in-house Perl script.

First, this script compares the scaffolds' genes order and orientation between the reference

assemblies of *D. saltans* and *D. sturtevanti*, the synteny blocks were defined when all the

genes were found in the same order and orientation in both species. The identified collinear

blocks were then searched for the remaining genomes. Blocks with missing data, *i.e.* missing

genes for one or more species were subsequently removed, and the remaining blocks were

subjected to a size-based filtering with a threshold of ≥50kb. The selected blocks were aligned

using the Mafft software (Katoh and Standley 2013).

Pseudo-reference inference

Illumina reads were aligned on either the *D. willistoni* reference genome (Clark et al.

2007: 12) or the recently published *D. sturtevanti* assembly (Kim et al. 2021) using minimap2

(Li 2018). Minimap2 generated BAM files were assembled into a pseudo-reference for each

species using the BEDTools software (Quinlan and Hall 2010). Each scaffold was then split

into 100 kb-long windows, and phylogenetic analyses were conducted on each window

separately using IqTree. A supertree was then inferred for each pseudo-reference genome

approach using ASTRAL-III.

Quantifying reticulation using the 2A2B test

Combinations of 28 species quartets were evaluated to test reticulation, bi-allelic sited

shared by pairs were searched in every locus, whether using SCG, syntenic blocks or pseudo-

reference 100 kb-long windows. Loci with at least 20 informative sites were retained. For

each locus, the occurrences AABB, ABBA or BABA patterns were counted, and three

pairwise χ^2-based tests were conducted, denoted *D1*, *D2*, and *D3* as follows (D1 is

reminiscent to Patterson's D):

651                $D1 = (\sum ABBA - \sum BABA)/(\sum ABBA + \sum BABA),$

652                $D2 = (\sum AABB - \sum ABBA)/(\sum AABB + \sum ABBA),$ and

653                $D3 = (\sum AABB - \sum BABA)/(\sum AABB + \sum BABA).$

Each of these estimates is reminiscent to the classical Tajima's (1993) Relative Rate Test

(RTT), which tests the difference in branch lengths from two ingroup taxa to a third outgroup

taxon, with lengths estimated as counts of different sites. However, unlike RRT, branch

lengths in our analyses are counted in terms of shared differences, *i.e.* singletons are not

considered. Similar to RRT, we applied a chi-squared analysis to test for deviation from parity

for each parameter. According to the significance levels of the three estimates, a locus can be

classified under each of the four categories presented in Figure 1. The test is run by the 2A2B

perl script associated with this manuscript.

Historical biogeography

To determine the sampling locations of the evaluated species, we conducted searches

in TaxoDros (https://www.taxodros.uzh.ch/search/class.php). Additionally, we incorporated

sampling sites that we ourselves had conducted. The accuracy of our species identification

was confirmed through DNA barcoding. After inspection of the geographical points and

manual correction, we identified the most northern, southern, western, and eastern points for

each species. We used those points to reconstruct the ancestral geographical range by

conducting a BayesTraits (Meade and Pagel 2022) analysis for each cartesian point

separately. The analyses were carried out using the GEO model with the phylogenetic tree

generated reconstructed with the Muller element A (XL chromosome arm), 1,000,000 of

MCMC iterations and 25% burn-in. The divergence times were estimated using this tree

under Bayesian inference. The calibration point used was the split between *D. willistoni* and

D. sturtevanti at 17.5 myr, as estimated by Suvorov et al. (2022).

References

Bächli G. 2024. TaxoDros: the database on Taxonomy of Drosophilidae. Available from:

https://www.taxodros.uzh.ch/

Baião GC, Schneider DI, Miller WJ, Klasson L. 2023. Multiple introgressions shape

mitochondrial evolutionary history in *Drosophila paulistorum* and the *Drosophila*

willistoni group. *Molecular Phylogenetics and Evolution* 180:107683.

Ballard JWO. 2000. Comparative Genomics of Mitochondrial DNA in Members of the

Drosophila melanogaster Subgroup. *J Mol Evol* 51:48–63.

Batalha-Filho H, Fjeldså J, Fabre P-H, Miyaki CY. 2013. Connections between the Atlantic

and the Amazonian forest avifaunas represent distinct historical events. *J Ornithol*

154:41–50.

Bicudo HEMC. 1973a. Chromosomal polymorphism in the saltans group of Drosophila I. The

saltans subgroup. *Genetica* 44:520–552.

Bicudo HEMC. 1973b. Reproductive isolation in the saltans group of Drosophila. I. The

saltans subgroup. *Genetica* 44:313–329.

Bicudo HEMC. 1979. Reproductive isolation of the saltans group of Drosophila. IV. the

sturtevanti subgroup. *Revista Brasileira de Genética* 2:247–258.

Bicudo HEMC, Prioli AJ. 1978. Reproductive isolation in the saltans group of Drosophila II.

The parasaltans subgroup. *Genetica* 48:17–22.

Blischak PD, Chifman J, Wolfe AD, Kubatko LS. 2018. HyDe: A Python Package for

Genome-Scale Hybridization Detection. *Systematic Biology* 67:821–829.

Bouckaert R, Vaughan TG, Barido-Sottani J, Duchêne S, Fourment M, Gavryushkina A,

Heled J, Jones G, Kühnert D, Maio ND, et al. 2019. BEAST 2.5: An advanced

software platform for Bayesian evolutionary analysis. *PLOS Computational Biology*

15:e1006650.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.

BLAST+: architecture and applications. *BMC Bioinformatics* 10:421.

Castro JP de, Carareto CMA. 2004. *P* elements in the *saltans* group of *Drosophila*: a new

evaluation of their distribution and number of genomic insertion sites. *Molecular*

Phylogenetics and Evolution 32:383–387.

Cavalcanti AGL. 1948. Geographic Variation of Chromosome Structure in Drosophila

Prosaltans. *Genetics* 33:529–536.

Chang ES, Neuhof M, Rubinstein ND, Diamant A, Philippe H, Huchon D, Cartwright P.

2015. Genomic insights into the evolutionary origin of Myxozoa within Cnidaria.

Proceedings of the National Academy of Sciences 112:14912–14917.

Charlesworth B, Campos JL, Jackson BC. 2018. Faster-X evolution: Theory and evidence

from Drosophila. *Molecular Ecology* 27:3753–3771.

Clark AG, Eisen MB, Smith DR, Bergman CM, Oliver B, Markow TA, Kaufman TC, Kellis

M, Gelbart W, Iyer VN, et al. 2007. Evolution of genes and genomes on the

Drosophila phylogeny. *Nature* 450:203–218.

Conner WR, Delaney EK, Bronski MJ, Ginsberg PS, Wheeler TB, Richardson KM,

Peckenpaugh B, Kim KJ, Watada M, Hoffmann AA, et al. 2021. A phylogeny for the

Drosophila montium species group: A model clade for comparative analyses.

Molecular Phylogenetics and Evolution 158:107061.

Dagilis AJ, Peede D, Coughlan JM, Jofre GI, D'Agostino ERR, Mavengere H, Tate AD,

Matute DR. 2021. 15 years of introgression studies: quantifying gene flow across

Eukaryotes. :2021.06.15.448399. Available from:

https://www.biorxiv.org/content/10.1101/2021.06.15.448399v1

David JR, Ferreira EA, Jabaud L, Ogereau D, Bastide H, Yassin A. 2022. Evolution of

assortative mating following selective introgression of pigmentation genes between

two Drosophila species. *Ecology and Evolution* 12:e8821.

Dobzhansky TG, Pavan C. 1943. Studies on Brazilian species of Drosophila. *Boletim da*

Faculdade de Filosofia, Ciências e Letras da Universidade de São Paulo. Biologia

Geral. Separata 36:7–72.

Dobzhansky Th. 1944. Experiments on Sexual Isolation in Drosophila. *Proceedings of the*

National Academy of Sciences 30:335–339.

Doronina L, Churakov G, Shi J, Brosius J, Baertsch R, Clawson H, Schmitz J. 2015.

Exploring Massive Incomplete Lineage Sorting in Arctoids (Laurasiatheria,

Carnivora). *Molecular Biology and Evolution* 32:3194–3204.

Durand EY, Patterson N, Reich D, Slatkin M. 2011. Testing for Ancient Admixture between

Closely Related Populations. *Molecular Biology and Evolution* 28:2239–2252.

Ellegren H. 2009. The different levels of genetic diversity in sex chromosomes and

autosomes. *Trends in Genetics* 25:278–284.

Gagnon E, Hilgenhof R, Orejuela A, McDonnell A, Sablok G, Aubriot X, Giacomin L,

Gouvêa Y, Bragionis T, Stehmann JR, et al. 2022. Phylogenomic discordance suggests

polytomies along the backbone of the large genus Solanum. *American Journal of*

Botany 109:580–601.

Glor RE. 2010. Phylogenetic Insights on Adaptive Radiation. *Annual Review of Ecology,*

Evolution, and Systematics 41:251–270.

Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, Patterson N, Li H, Zhai

W, Fritz MH-Y, et al. 2010. A Draft Sequence of the Neandertal Genome. *Science*

328:710–722.

Hime PM, Lemmon AR, Lemmon ECM, Prendini E, Brown JM, Thomson RC, Kratovil JD,

754   Noonan BP, Pyron RA, Peloso PLV, et al. 2021. Phylogenomics Reveals Ancient

755   Gene Tree Discordance in the Amphibian Tree of Life. *Systematic Biology* 70:49–66.

756 Hoorn C, Boschman LM, Kukla T, Sciumbata M, Val P. 2022. The Miocene wetland of

757   western Amazonia and its role in Neotropical biogeography. *Botanical Journal of the*

758   *Linnean Society* 199:25–35.

759 Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version 7:

760   Improvements in Performance and Usability. *Molecular Biology and Evolution*

761   30:772–780.

762 Khallaf MA, Cui R, Weißflog J, Erdogmus M, Svatoš A, Dweck HKM, Valenzano DR,

763   Hansson BS, Knaden M. 2021. Large-scale characterization of sex pheromone

764   communication systems in Drosophila. *Nat Commun* 12:4165.

765 Kim BY, Wang JR, Miller DE, Barmina O, Delaney E, Thompson A, Comeault AA, Peede D,

766   D'Agostino ER, Pelaez J, et al. 2021. Highly contiguous assemblies of 101

767   drosophilid genomes.Coop G, Wittkopp PJ, Sackton TB, editors. *eLife* 10:e66405.

768 Kokot M, Długosz M, Deorowicz S. 2017. KMC 3: counting and manipulating k-mer

769   statistics. *Bioinformatics* 33:2759–2761.

770 Kuznetsov D, Tegenfeldt F, Manni M, Seppey M, Berkeley M, Kriventseva EV, Zdobnov

771   EM. 2023. OrthoDB v11: annotation of orthologs in the widest sampling of

772   organismal diversity. *Nucleic Acids Research* 51:D445–D451.

773 Ledo RMD, Colli GR. 2017. The historical connections between the Amazon and the Atlantic

774   Forest revisited. *Journal of Biogeography* 44:2551–2563.

775 Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. 2015. MEGAHIT: an ultra-fast single-node

776   solution for large and complex metagenomics assembly via succinct de Bruijn graph.

777   *Bioinformatics* 31:1674–1676.

Li F, Rane RV, Luria V, Xiong Z, Chen J, Li Z, Catullo RA, Griffin PC, Schiffer M, Pearce

S, et al. 2022. Phylogenomic analyses of the genus Drosophila reveals genomic signals

of climate adaptation. *Molecular Ecology Resources* 22:1559–1581.

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34:3094–

3100.

Li H. 2023. Protein-to-genome alignment with miniprot. *Bioinformatics* 39:btad014.

Llopart A, Herrig D, Brud E, Stecklein Z. 2014. Sequential adaptive introgression of the

mitochondrial genome in Drosophila yakuba and Drosophila santomea. *Molecular*

Ecology 23:1124–1136.

Maddison W. 1989. Reconstructing Character Evolution on Polytomous Cladograms.

Cladistics 5:365–377.

Maddison WP. 1997. Gene Trees in Species Trees. *Systematic Biology* 46:523–536.

Madi-Ravazzi L, Segala LF, Roman BE, Alevi KCC, Prediger C, Yassin A, Hua-VAN AL,

Miller WJ. 2021. Integrative taxonomy and a new species description in the sturtevanti

subgroup of the Drosophila saltans group (Diptera: Drosophilidae). *Zootaxa*

4980:269292.

Magalhães LE de. 1962. Notes on the taxonomy, morphology, and distribution of the saltans

group of Drosophila, with descriptions of four new species. *The University of Texas*

Publication 2:135–154.

Magalhães LE de, Björnberg AJS. 1957. Estudo da genitália masculina de "Drosophila" do

grupo "saltans" (Diptera). *Revista Brasileira de Biologia* 17:435–450.

Mai D, Nalley MJ, Bachtrog D. 2020. Patterns of Genomic Differentiation in the Drosophila

nasuta Species Complex. *Molecular Biology and Evolution* 37:208–220.

Malinsky M, Matschiner M, Svardal H. 2021. Dsuite - Fast D-statistics and related admixture

evidence from VCF files. *Molecular Ecology Resources* 21:584–595.

Manni M, Berkeley MR, Seppey M, Zdobnov EM. 2021. BUSCO: Assessing Genomic Data

Quality and Beyond. *Current Protocols* 1:e323.

Martin SH, Davey JW, Jiggins CD. 2015. Evaluating the Use of ABBA–BABA Statistics to

Locate Introgressed Loci. *Molecular Biology and Evolution* 32:244–257.

Matute DR, Comeault AA, Earley E, Serrato-Capuchina A, Peede D, Monroy-Eklund A,

Huang W, Jones CD, Mackay TFC, Coyne JA. 2020. Rapid and Predictable Evolution

of Admixed Populations Between Two Drosophila Species Pairs. *Genetics* 214:211–

230.

Mayr E, Dobzhansky Th. 1945. Experiments on Sexual Isolation in Drosophila. *Proceedings*

of the National Academy of Sciences 31:75–82.

Meade A, Pagel M. 2022. Ancestral State Reconstruction Using BayesTraits. In: Luo H,

editor. Environmental Microbial Evolution: Methods and Protocols. Methods in

Molecular Biology. New York, NY: Springer US. p. 255–266. Available from:

https://doi.org/10.1007/978-1-0716-2691-7_12

Meng G, Li Y, Yang C, Liu S. 2019. MitoZ: a toolkit for animal mitochondrial genome

assembly, annotation and visualization. *Nucleic Acids Research* 47:e63.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A,

Lanfear R. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic

Inference in the Genomic Era. *Molecular Biology and Evolution* 37:1530–1534.

Moran BM, Payne C, Langdon Q, Powell DL, Brandvain Y, Schumer M. 2021. The genomic

consequences of hybridization.Wittkopp PJ, editor. *eLife* 10:e69016.

Moreyra NN, Almeida FC, Allan C, Frankel N, Matzkin LM, Hasson E. 2023. Phylogenomics

provides insights into the evolution of cactophily and host plant shifts in *Drosophila*.

Molecular Phylogenetics and Evolution 178:107653.

Morgan CC, Foster PG, Webb AE, Pisani D, McInerney JO, O'Connell MJ. 2013.

Heterogeneous Models Place the Root of the Placental Mammal Phylogeny.

Molecular Biology and Evolution 30:2145–2156.

Nascimento AP, Bicudo HEMC. 2002. Esterase Patterns and Phylogenetic Relationships of

Drosophila Species in the Saltans Subgroup (Saltans Group). *Genetica* 114:41–51.

Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: A Fast and Effective

Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular*

Biology and Evolution 32:268–274.

O'Dea A, Lessios HA, Coates AG, Eytan RI, Restrepo-Moreno SA, Cione AL, Collins LS, de

Queiroz A, Farris DW, Norris RD, et al. 2016. Formation of the Isthmus of Panama.

Science Advances 2:e1600883.

O'Grady PM, Clark JB, Kidwell MG. 1998. Phylogeny of the Drosophila saltans species

group based on combined analysis of nuclear and mitochondrial DNA sequences.

Molecular Biology and Evolution 15:656–664.

Owen CL, Miller GL. 2022. Phylogenomics of the Aphididae: Deep relationships between

subfamilies clouded by gene tree discordance, introgression and the gene tree anomaly

zone. *Systematic Entomology* 47:470–486.

Pease JB, Hahn MW. 2015. Detection and Polarization of Introgression in a Five-Taxon

Phylogeny. *Systematic Biology* 64:651–662.

Pélandakis M, Solignac M. 1993. Molecular phylogeny of Drosophila based on ribosomal

RNA sequences. *J Mol Evol* 37:525–543.

Philippe H, Brinkmann H, Lavrov DV, Littlewood DTJ, Manuel M, Wörheide G, Baurain D.

2011. Resolving Difficult Phylogenetic Questions: Why More Sequences Are Not

Enough. *PLOS Biology* 9:e1000602.

Philippe H, Derelle R, Lopez P, Pick K, Borchiellini C, Boury-Esnault N, Vacelet J, Renard

E, Houliston E, Quéinnec E, et al. 2009. Phylogenomics Revives Traditional Views on

Deep Animal Relationships. *Current Biology* 19:706–712.

Pick KS, Philippe H, Schreiber F, Erpenbeck D, Jackson DJ, Wrede P, Wiens M, Alié A,

Morgenstern B, Manuel M, et al. 2010. Improved Phylogenomic Taxon Sampling

Noticeably Affects Nonbilaterian Relationships. *Molecular Biology and Evolution*

27:1983–1987.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* 26:841–842.

Qvarnström A, Bailey RI. 2009. Speciation through evolution of sex-linked genes. *Heredity*

102:4–15.

Ranallo-Benavidez TR, Jaron KS, Schatz MC. 2020. GenomeScope 2.0 and Smudgeplot for

reference-free profiling of polyploid genomes. *Nat Commun* 11:1432.

Reilly PF, Tjahjadi A, Miller SL, Akey JM, Tucci S. 2022. The contribution of Neanderthal

introgression to modern human traits. *Current Biology* 32:R970–R983.

Rick JA, Brock CD, Lewanski AL, Golcher-Benavides J, Wagner CE. 2023. Reference

Genome Choice and Filtering Thresholds Jointly Influence Phylogenomic Analyses.

Systematic Biology:syad065.

Rodríguez-Trelles F, Tarrío R, Ayala FJ. 1999. Molecular Evolution and Phylogeny of the

Drosophila saltans Species Group Inferred from the *Xdh* Gene. *Molecular*

Phylogenetics and Evolution 13:110–121.

Roman BE, Madi-Ravazzi L. 2021. <p>Male terminalia morphology of sixteen

species of the Drosophila saltans group

Sturtevant (Diptera, Drosophilidae)</p>. *Zootaxa*

5061:523–544.

Roman BE, Santana DJ, Prediger C, Madi-Ravazzi L. 2022. Phylogeny of Drosophila saltans

group (Diptera: Drosophilidae) based on morphological and molecular evidence.

PLOS ONE 17:e0266710.

Romiguier J, Ranwez V, Delsuc F, Galtier N, Douzery EJP. 2013. Less Is More in

Mammalian Phylogenomics: AT-Rich Genes Minimize Tree Conflicts and Unravel

the Root of Placental Mammals. *Molecular Biology and Evolution* 30:2134–2144.

Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L,

Suchard MA, Huelsenbeck JP. 2012. MrBayes 3.2: Efficient Bayesian Phylogenetic

Inference and Model Choice Across a Large Model Space. *Systematic Biology*

61:539–542.

Sankararaman S, Mallick S, Patterson N, Reich D. 2016. The Combined Landscape of

Denisovan and Neanderthal Ancestry in Present-Day Humans. *Current Biology*

26:1241–1247.

Sayyari E, Mirarab S. 2018. Testing for Polytomies in Phylogenetic Species Trees Using

Quartet Frequencies. *Genes* 9:132.

Schaeffer SW. 2018. Muller "Elements" in Drosophila: How the Search for the Genetic Basis

for Speciation Led to the Birth of Comparative Genomics. *Genetics* 210:3–13.

Schrempf D, Szöllősi G. 2020. The Sources of Phylogenetic Conflicts. In: No commercial

publisher | Authors open access book. p. 3.1:1. Available from: https://hal.science/hal-

02535482

Seixas FA, Boursot P, Melo-Ferreira J. 2018. The genomic impact of historical hybridization

with massive mitochondrial DNA introgression. *Genome Biology* 19:91.

de Setta N, Loreto ELS, Carareto CMA. 2007. Is the Evolutionary History of the O-Type P

Element in the saltans and willistoni Groups of Drosophila Similar to That of the

Canonical P Element? *J Mol Evol* 65:715–724.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO:

assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31:3210–3212.

Simion P, Philippe H, Baurain D, Jager M, Richter DJ, Di Franco A, Roure B, Satoh N, Quéinnec É, Ereskovsky A, et al. 2017. A Large and Consistent Phylogenomic Dataset Supports Sponges as the Sister Group to All Other Animals. *Current Biology* 27:958– 967.

Skov L, Coll Macià M, Lucotte EA, Cavassim MIA, Castellano D, Schierup MH, Munch K. 2023. Extraordinary selection on the human X chromosome associated with archaic admixture. *Cell Genomics* 3:100274.

Solignac M, Monnerot M, Mounolou J-C. 1986. Mitochondrial DNA evolution in themelanogaster species subgroup ofDrosophila. *J Mol Evol* 23:31–40.

Souza TAJ, Noll FB, Bicudo HEMC, Madi-Ravazzi L. 2014. Scanning Electron Microscopy of Male Terminalia and Its Application to Species Recognition and Phylogenetic Reconstruction in the Drosophila saltans Group. *PLOS ONE* 9:e97156.

Spassky B. 1957. Morphological differences between sibling species of Drosophila. *University of Texas Publication* 5721:48–61.

Sturtevant AH. 1942. The classification of the genus Drosophila, with descriptions of nine species. *University of Texas Publication* 4213:5–51.

Sturtevant AH, Novitski E. 1941. THE HOMOLOGIES OF THE CHROMOSOME ELEMENTS IN THE GENUS DROSOPHILA. *Genetics* 26:517–541.

Suh A. 2016. The phylogenomic forest of bird trees contains a hard polytomy at the root of Neoaves. *Zoologica Scripta* 45:50–62.

Suvorov A, Kim BY, Wang J, Armstrong EE, Peede D, D'Agostino ERR, Price DK, Waddell

PJ, Lang M, Courtier-Orgogozo V, et al. 2022. Widespread introgression across a

phylogeny of 155 *Drosophila* genomes. *Current Biology* 32:111-123.e5.

Tajima F. 1993. Simple methods for testing the molecular evolutionary clock hypothesis.

Genetics 135:599–607.

Throckmorton LH. 1962. The problem of phylogeny in the genus Drosophila.Studies in

Genetics.II. *University of Texas Publication* 6205:207–343.

Throckmorton LH, Magalhães LE de. 1962. Changes with evolution of pteridine

accumulations in species of the saltans group of the genus Drosophila. *University of*

Texas Publication 6205:489–505.

Toews DPL, Brelsford A. 2012. The biogeography of mitochondrial and nuclear discordance

in animals. *Molecular Ecology* 21:3907–3930.

Townsend JP, Su Z, Tekle YI. 2012. Phylogenetic Signal and Noise: Predicting the Power of

a Data Set to Resolve Phylogeny. *Systematic Biology* 61:835.

Valiente-Mullor C, Beamud B, Ansari I, Francés-Cuesta C, García-González N, Mejía L,

Ruiz-Hueso P, González-Candelas F. 2021. One is not enough: On the effects of

reference genome for the mapping and subsequent analyses of short-reads. *PLOS*

Computational Biology 17:e1008678.

Walsh HE, Kidd MG, Moum T, Friesen VL. 1999. Polytomies and the Power of Phylogenetic

Inference. *Evolution* 53:932–937.

Whelan NV, Kocot KM, Moroz LL, Halanych KM. 2015. Error, signal, and the placement of

Ctenophora sister to all other animals. *Proceedings of the National Academy of*

Sciences 112:5773–5778.

Wickett NJ, Mirarab S, Nguyen N, Warnow T, Carpenter E, Matasci N, Ayyampalayam S,

Barker MS, Burleigh JG, Gitzendanner MA, et al. 2014. Phylotranscriptomic analysis

of the origin and early diversification of land plants. *Proceedings of the National*

Academy of Sciences 111:E4859–E4868.

Yassin A. 2009. Phylogenetic relationships among species subgroups in the Drosophila

saltans group (Diptera: Drosophilidae): Can morphology solve a molecular conflict.

Zool Res 30:225–232.

Yusuf LH, Tyukmaeva V, Hoikkala A, Ritchie MG. 2022. Divergence and introgression

among the virilis group of Drosophila. *Evolution Letters* 6:537–551.

Zhang C, Rabiee M, Sayyari E, Mirarab S. 2018. ASTRAL-III: polynomial time species tree

reconstruction from partially resolved gene trees. *BMC Bioinformatics* 19:153.

Zimin AV, Marçais G, Puiu D, Roberts M, Salzberg SL, Yorke JA. 2013. The MaSuRCA

genome assembler. *Bioinformatics* 29:2669–2677.

 Figure legends

Figure 1 – The distribution of bi-allelic sites proportions among three ingroup species with an

outgroup and the rationale of the 2A2B test. A) Statistical pairwise comparisons of the three

possible bi-allelic patterns generate four categories. B) Each category corresponds to a

possible evolutionary scenario along the reticulation-bifurcation continuum. B) Based of the

frequency of these topologies in a genome fragment, this fragment can be categorized in (i)

complete reticulation, (ii) incomplete reticulation, (iii) incomplete bifurcation, and (iv)

complete bifurcation. The 2A2B test estimates significant deviation from parity in pairwise

comparisons of the topologies. *: significant deviation, ns = non-significant.

Figure 2 – Phylogenomic conflict between the X chromosome, autosomes, and the

mitochondria. A) Comparison of autosomal topology (left, represented by Muller element B)

and X chromosome topology (right, represented by Muller element A) demonstrates overall

agreement with minor incongruence. B) Mitochondrial-nuclear disagreement highlights

stronger incongruence between mitochondrial topology (left) and sexual chromosome

topology (right). All trees were inferred through Bayesian analysis. Divergence time

estimation (in million years ago, myr) for the sexual chromosome topology is provided (left).

Nodes are numbered for subsequent analyses. Maximum-likelihood and ASTRAL-III trees

are provided in Supplementary Figures S2 and S3, respectively. Posterior probabilities of all

nodes were equal to1.

Figure 3 – Extent of phylogenomic discordance across 28 species quartets as measured by A)

Patterson's *D* and B) the 2A2B test, using four locus-definition strategies (see text). For both

tests, assembly-based strategies have lower average *D* and provide stronger evidence for

reticulation at three episodes. Colors of bars in B) correspond to the four categories presented

in Figure 1, with grey bars corresponding to loci with less than 20 evaluated bi-allelic sites.

Figure 4 – Historical biogeography of the *saltans* species. Centroid geographical coordinates

for each species were inferred and ancestral coordinates were inferred using BayesTraits on

the Bayesian X chromosome chronogram given in Figure 2B. Internal nodes are projected on

the current map of South and Central America. Three episodes of rapid radiation were

identified by the 2A2B test (Figure 3B) and are identified here with their corresponding

possible paleogeographical events, with photos of adult males of four representative species

shown. Branch colors in the tree correspond to the subgroups colors given in Figure 2.

List of Supplementary Figures

Figure S1 – Phylogenetic relationships inferred using BEAST from BUSCO genes

concatenated according to their Muller's element.

1002

1003 **Figure S2 –** Phylogenetic relationships inferred using IqTree-2 from BUSCO genes

1004 concatenated according to their Muller's element.

1005

1006 **Figure S3 –** Phylogenetic relationships inferred using ASTRAL-III from BUSCO genes

1007 concatenated according to their Muller's element.

1008

1009 **Figure S4 –** Phylogenetic relationships inferred using ASTRAL-III from 100-kb long

1010 windows for pseudo-references produced by aligning short reads either to A) the ingroup *D.*

1011 *sturtevanti* or B) the outgroup *D. willistoni* reference assemblies.

1012

1013

1014

1015   **List of Supplementary Tables**

1016

1017   **Table S1 –** A synopsis of phylogenetic relationships previously published in the *Drosophila*

1018   *saltans* species group. Species subgroups are represented by the first two letters in capital,

1019   whereas species names are represented by the first three letter in small.

1020

1021   **Table S2 –** Source and genome assembly statistics including busco scores for the genomes

1022   used in this study.

1023

A

Frequency of patterns of bi-allelic sites

Sp. 1	B A B	B A B	B A B	B A B
Sp. 2	B B A	B B A	B B A	B B A
Sp. 3	A B B	A B B	A B B	A B B
O	A A A	A A A	A A A	A A A

B

(i)
Complete trifurcation (e.g., radiation)

(ii)
Incomplete trifurcation (e.g., hybrid speciation)

(iii)
Incomplete bifurcation (e.g., introgression)

(iv)
Complete bifurcation

sp.1 sp.2 sp.3 O

sp.1 sp.2 sp.3 O

sp.1 sp.2 sp.3 O

sp.1 sp.2 sp.3 O

The phylogenetic resolution continuum

~10 myr: Formation of the Amazonian River and the split of the *cordata-elliptica-saltans* subgroups

~3 myr: Formation of Panama Isthmus and radiation of the *saltans* subgroup

~2.5 myr: Connection between the Amazonian and Atlantic Forests and radiation of the *sturtevanti* subgroup

D. neocordata

D. emarginata

D. prosaltans

D. sturtevanti