Saltational episodes of reticulate evolution in the *Drosophila saltans* species group

Carolina Prediger[1,2], Erina A. Ferreira[2], Samara Videira Zorzato[1,3], Aurélie Hua-Van[2], Lisa

Klasson[4], Wolfgang J. Miller[5], Amir Yassin[2,3*] and Lilian Madi-Ravazzi[1*]

[1]Department of Biology, UNESP - São Paulo State University, São José do Rio Preto, São

Paulo, Brazil.

[2]Laboratoire Évolution, Génomes, Comportement et Écologie, CNRS, IRD, Université Paris-

Saclay, Gif-sur-Yvette, France.

[3]Institut de Systématique, Évolution, Biodiversité (ISYEB), CNRS, MNHN, EPHE, Sorbonne

Université, Univ. des Antilles, Paris, France.

[4]Molecular evolution, Department of Cell and Molecular Biology, Science for Life

Laboratory, Uppsala University, Sweden.

[5]Center for Anatomy and Cell Biology, Department of Cell and Developmental Biology,

Medical University of Vienna, Austria.

[*] These authors contributed equally.

Correspondence: amir.yassin@universite-paris-saclay.fr; lilian.madi@unesp.br.

Running title: Phylogenomics of the *Drosophila saltans* group

Conflict of interest

The authors declare no conflict of interest.

Data availability statement

Scripts used in this study, assemblies and intermediate files are publicly available on Figshare

(https://figshare.com/account/home#/collections/7100635; private link for review:

https://figshare.com/s/04b0b0f46808a8cdeefa) and will be made public upon acceptance.

1
2

25 Genome sequences generated for this study are available on NCBI BioProject:

26 PRJNA1078835.

3 2
4

Abstract

Phylogenomics revealed reticulate evolution to be widespread across taxa, but whether

reticulation is due to low statistical power (soft polytomy) or it is a true evolutionary pattern

(hard polytomy) remains a field of investigation. Here, we investigate the phylogeny and

quantify reticulation in the *Drosophila saltans* species group, a Neotropical clade of the

subgenus *Sophophora* comprising 23 species arranged in five subgroups, namely *cordata*,

elliptica, *parasaltans*, *saltans* and *sturtevanti.* The relationships within and between the

subgroups have long been problematic. We sequenced and assembled the genomes of 15

species. Phylogenetic analyses revealed conflicting topologies between the X chromosome,

autosomes and the mitochondria. We extended the ABBA-BABA test of asymmetry in

phylogenetic discordance to cases where no "true" species tree could be inferred, and applied

our new test (called 2A2B) to $\geq$50 kb-long 1,797 syntenic blocks with conserved collinearity

across Neotropical *Sophophora*. High incidences of reticulation (sometimes up to 90% of the

blocks) were restricted to three nodes on the tree, at the split between the *cordata-elliptica-*

saltans subgroups and at the origin of the *sturtevanti* and *saltans* subgroups. By contrast,

cases with asymmetric discordances, which are often interpreted as evidence for interspecific

introgression, did not exceed ~7% of the blocks. Historical biogeography analysis revealed

that short inter-speciation times and greater overlap of ancestral geographical ranges can

explain cases with predominant reticulation. Therefore, episodic rapid radiations have played

a major role in the evolution of this largely understudied Neotropical clade.

Keywords: phylogenomic discordance; genome assembly; historical biogeography;

introgression; cyto-nuclear conflicts; Neotropical speciation; *Sophophora*.

Introduction

Knowledge of phylogenetic relationships among species is a requirement for many evolutionary studies. However, it is often difficult to reconstruct well-resolved bifurcating trees for some clades. This could either be due to the lack of signal in the evaluated data, a condition known as "soft polytomy", or due to persistent phylogenetic conflicts among datasets leading to "hard polytomies" and reticulate patterns of interspecific relationships. A plethora of biological processes could cause such conflicts, including incomplete lineage sorting (Maddison 1989; Maddison 1997; Walsh et al. 1999; Townsend et al. 2012), horizontal gene transfer, introgression and hybridization (Schrempf and Szöllősi 2020), and adaptive radiations (Glor 2010). Phylogenetic conflict may also be caused by technical errors, such as, sequencing error, contamination, wrong model selection and general lack of quality control (Philippe et al. 2011). Recent advances in genomic analyses have significantly reduced such errors and, in a wide range of taxa, increased the number of analyzed genes hence helping to resolve early conflicting topologies. However, in many other cases, whole genome analyses demonstrated persistent phylogenetic conflicts (e.g., in plants (Wickett et al. 2014; Gagnon et al. 2022), birds (Suh 2016), sponges and ctenophores (Philippe et al. 2009; Pick et al. 2010; Chang et al. 2015; Whelan et al. 2015; Simion et al. 2017), mammals (Morgan et al. 2013; Romiguier et al. 2013; Doronina et al. 2015), amphibians (Hime et al. 2021), and insects (Owen and Miller 2022)).

Of the different processes that can lead to reticulated evolution, introgression and hybridization have attracted much attention, first because they challenged long-held concept of reproductive isolation between species, and second due to the development of a number of bioinformatic tools and tests that quantify phylogenetic discordance across the genome (Durand et al. 2011; Pease and Hahn 2015; Malinsky et al. 2021). Site-based methods usually count the number of bi-allelic sites supporting each of three possible topologies in a species

triplet with an outgroup. Comparisons between the proportions of the three topologies can

yield one of four possible outcomes (Figure 1B): (i) complete reticulation, all topologies are

equally encountered; (ii) incomplete reticulation, such as in the case of full hybridization

wherein two topologies significantly exceed the third one but do not significantly differ from

each other; (iii) incomplete bifurcation, such as in the case of asymmetric introgression

wherein the proportion of all topologies significantly differ; and (iv) complete bifurcation, one

topology significantly exceeds the two others, which in their turn have nearly equal

proportions. The earliest of introgression tests, Patterson's D or the ABBA-BABA statistic

(Green et al. 2010; Durand et al. 2011), compared the two later cases (iii and iv), *i.e.* it

presumed that a "true" species tree exists. A later test, HyDe (Blischak et al. 2018), quantifies

admixture (γ) from the ratio of shared alleles with the test going from 0 (full isolation) to 0.5

(full hybridization) and therefore it can also cover case ii. The two tests differ in how they

measure significance, using bootstrapping in Patterson's D and normal approximation in

HyDe. Of late, another site-based test was developed using χ^2 to test for deviation of parity

between the three topologies as in case i (Sayyari and Mirarab 2018). A unified test that can

test the prevalence of each of the four categories across the genome is still lacking.

Polytomies and incongruencies have been reported for the jumping fly *Drosophila*

saltans species group, a clade of the subgenus *Sophophora* with 23 Neotropical species

(Magalhães 1962; Bächli 2024). The group retains its name from the peculiar "jumping" habit

of its larvae; *"the larva seizes its posterior end with its mouthhooks, and stretches. The hooks*

pull loose suddenly, the larva straightens with considerable force, and as a result is thrown

several inches into the air" (Sturtevant 1942). The group was divided into five species

subgroups, namely, *saltans, parasaltans, cordata, elliptica* and *sturtevanti* subgroups, mostly

on the basis of male genitalia (Magalhães and Björnberg 1957). Although the monophyly of

the subgroups has been confirmed by different phylogenetic methods, the relationships among

9 5

and within them are not. Hypothesis for their evolutionary relationships have been proposed

using different methods and different morphological characters (Magalhães and Björnberg

1957; Throckmorton 1962; Throckmorton and Magalhães 1962; O'Grady et al. 1998; Yassin

2009; Souza et al. 2014; Roman et al. 2022), chromosome polymorphism (Bicudo 1973a),

reproductive isolation (Bicudo 1973b; Bicudo and Prioli 1978; Bicudo 1979), protein

polymorphism (Nascimento and Bicudo 2002) and gene sequences (Pélandakis and Solignac

1993; O'Grady et al. 1998; Rodríguez-Trelles et al. 1999; Castro and Carareto 2004; de Setta

et al. 2007; Roman et al. 2022). The evolutionary relationships proposed are summarized in

Supplementary Table S1.

Unlike other species groups in the subgenus *Sophophora*, such as the *melanogaster*,

obscura and *willistoni* groups, genomic resources and genetic investigations in the *saltans*

species group are scarce. Indeed, only four genomes have been sequenced and assembled to

date (Kim et al. 2021). To bridge this gap and to test for the extent of phylogenetic conflicts,

we sequenced and assembled genomes for 15 species with representatives from the five

subgroups. Phylogenetic analyses using well-conserved genes resolved the evolutionary

relationships among the subgroups but also highlighted conflicts between X-linked, autosomal

and mitochondrial loci. To test how each of the four incongruence categories prevails across

the genome, we devised a new χ^2-based test that we called 2A2B. The test uses pairwise

comparisons of the three topologies proportions shown in Figure 1. We applied this test to

sets of conserved orthologous genes and to long syntenic blocks with conserved collinearity

across the Neotropical *Sophophora*, as well as to 100-kb long windows from pseudo-reference

genomes aligned to either an ingroup (*D. sturtevanti*) or outgroup (*D. willistoni*) species. We

found reticulation levels to differ among the datasets and the subgroups, and to correlate with

rate of speciation and historical biogeography in the *saltans* group.

11 6
12

Results

Short-read assembly of 17 genomes recovered 90% of BUSCO genes

We sequenced using short-read Illumina approach 17 whole genomes from 15 species collected across various locations in the Neotropical region. Genome size, estimated from 21-kmer frequency spectrum using GenomeScope 2 (Ranallo-Benavidez et al. 2020), ranged from 154.0 to 356.8 Mb. Our de novo assemblies using MaSuRCA (Zimin et al. 2013) resulted in genome lengths ranging from 177.5 to 287.7 Mb, with N50 values ranging from 2 to 92 Kb (Supplementary Table S2). To evaluate the completeness of our assembled genomes, we searched for single-copy genes (SCG) using BUSCO (Simão et al. 2015). We found that over 90% of the searched genes were complete for all of the genomes (Supplementary Table S2). Kim et al. (2021) assembled using both short Illumina and long Nanopore reads the genomes of four *saltans* group species, all of which we have independently sequenced. Whereas their assemblies' contigs were much longer, with N50 ranging from 2 to 6 Mb, the BUSCO score for the same set of species did not largely differ (98% vs. 95-96% in our study; Supplementary Table S2).

Muller elements analysis resolves relationships between the subgroups and unravels a minor X-autosomal conflict in the sturtevanti subgroup

Phylogenomic analyses were performed using 2,159 SCG shared across all species. Gene trees, inferred for each SCG using maximum-likelihood in IqTree2 (Minh et al. 2020) produced 1,263 distinct topologies, with 206 of them found more than once. To test if SCG chromosomal position may underlie the discrepancies in gene trees, we localized each SCG to its corresponding Muller element according to the position of its *D. melanogaster* ortholog identified by Blast (Camacho et al. 2009). As a result, we generated five independent datasets, each corresponding to the Muller elements A, B, C, D, and E, comprising 337, 370, 425, 419,

and 568 SCG, respectively. These datasets were then used to reconstruct the species trees

using the multi-species coalescent model, and the genes within them were concatenated for

Bayesian and Maximum Likelihood phylogenetic inferences.

The trees generated by the 5 data sets showed very similar topologies with well supported

nodes either for the multi-species coalescent model analysis implemented in ASTRAL-III

(Zhang et al. 2018), the maximum-likelihood implemented in IqTree2, or Bayesian Inference

implemented in BEAST (Bouckaert et al. 2019) (Supplementary Figures S1, S2 and S3). The

parasaltans subgroup was placed as sister to all other subgroups, followed by the emergence

of the *sturtevanti* subgroup. The *cordata* and *elliptica* subgroups showed a close relationship,

and were sister to the *saltans* subgroup. The only discrepancy between the topologies was the

placement of *D. lehrmanae*, a newly discovered species in the *sturtevanti* subgroup (Madi-

Ravazzi et al. 2021). For *D. lehrmanae*, while maximum-likelihood and multi-species

coalescent analyses reported lack of branch support for multiple trees (Supplementary Figures

S2 and S3), Bayesian inference recovered well supported branches and two topologies (Figure

2A and Supplementary Figure S1). The two distinct topologies distinguished the Muller

elements forming the X chromosome (elements A and D who are fused in a fusion shared by

the Neotropical *Sophophora*, *i.e.* the *saltans* and *willistoni* groups (Sturtevant and Novitski

1941; Dobzhansky and Pavan 1943; Cavalcanti 1948), and the Muller elements representing

autosomal chromosomes (elements B, C and E).

The published genome of *D. prosaltans* (Kim et al. 2021) did not group with the

genome of this species sequenced by us, instead it grouped with *D. saltans*. The genome

previously published comes from a line collected in El Salvador in 1957. According to

Magalhães' (1962) detailed morphological revision of multiple geographical specimens of the

saltans group, the sampling site of this particular strain is outside the geographical range of *D.*

prosaltans, but within the expected range of *D. saltans*. Furthermore, the *D. saltans* and *D.*

prosaltans lines used in our study underwent thorough morphological analyzes (Souza et al.

2014; Roman and Madi-Ravazzi 2021), indicating that the lines we used were accurately

identified. Therefore, it is most likely that the previously sequenced *D. prosaltans* strain from

El Salvador was misidentified and we consider it here to belong to *D. saltans*.

Mitogenomes show cytonuclear conflicts in the sturtevanti *and* saltans *subgroups*

We assembled mitochondrial genomes for the 15 *saltans* group species using MitoZ

(Meng et al. 2019). We did not use the previously assembled four strains since several

mitochondrial scaffolds were likely removed in those assemblies (Kim et al. 2021). We

conducted phylogenetic analysis on the aligned mitogenomes genes using both IqTree2 and

MrBayes (Ronquist et al. 2012). Overall, the mitochondrial trees matched the topology of the

nuclear gene trees regarding the inter-subgroup relationships. However, three major

discrepancies were identified (Figure 2B). First, the position of *D. lehrmanae* within the

sturtevanti subgroup did not agree with either the X or autosomal SCG topologies, proposing

a topology wherein *D. lehrmanae* is a sister species of *D. sturtevanti* (a topology recovered

once in multi-species coalescent analysis (Muller element C, Supplementary Figure S3) and

maximum likelihood (Muller element B, Supplementary Figure S2)). Second, whereas the

mitochondrial tree recovered the monophyletic relationship between the *elliptica*, *cordata* and

saltans subgroups, the position of *D. neocordata* (*cordata* subgroup) differed, being sister to

the three species of the *elliptica* subgroup in the nuclear trees and to the six species of the

saltans subgroup in the mitochondrial tree. Third, whereas nuclear trees recovered three

lineages within the *saltans* subgroup, namely*, austrosaltans*, *nigrosaltans-pseudosaltans*, and

septentriosaltans-prosaltans-saltans, only two lineages are revealed by the mitochondrial tree.

Intriguingly, each of the mitochondrial clades involved one species from otherwise sister

species in the nuclear trees, *i.e. D. nigrosaltans* and *D. saltans* in one clade and their

17 9
18

respective closely-related species *D. pseudosaltans* and *D. prosaltans* in the other clade.

Because *D. saltans* and *D. prosaltans* are reported as closely related species in the nuclear

trees and are separated in the two mitochondrial clades, the two mitotypes were called S and

P, respectively. The distribution of closely-related species between distinct mitotypic groups

suggest that multiple cytoplasmic introgression events might have occurred in this subgroup

(Figure 2B).

The extent of reticulation differs between assembly- and pseudo-reference-based approaches

To quantify phylogenomic incongruences at a finer locus-scale, we analyzed allele

distribution of bi-allelic phylogenetically informative sites in 28 species quartets using four

distinct locus-defining strategies. Two strategies relied on our *de novo* genome assemblies.

The first strategy included the use of single-copy genes (SCG) defined by BUSCO (2,159

loci). The second strategy identified syntenic blocks equal to or greater than 50 kb-long with

collinearity across the 15 *saltans* assemblies and *D. willistoni*, *i.e.* across Neotropical

Sophophora (1,797 loci). The third and fourth strategies used a pseudo-reference approach,

wherein the reads of each *saltans* group species were mapped to an ingroup (*D. sturtevanti*,

2,328 loci) or an outgroup species (*D. willistoni*, 1,863 loci) and a pseudo-reference was

inferred by replacing variant sites in the reference genome by the ones in the reads (as in Mai

et al. 2020). In these strategies, a locus was defined as a 100 kb-long window of the reference

genome.

For a bi-allelic site, the standard ABBA-BABA (Patterson's *D*) test considers a true

species tree exists, with site distribution AABB, and then evaluate the deviation from parity of

the two discordant configurations ABBA and BABA. We found that genome-wide absolute

Patterson's *D* estimates, *i.e.* $(\sum ABBA - \sum BABA)/(\sum ABBA + \sum BABA)$, greatly differed

among the four strategies (Figure 3A). Synteny-based strategy had the lowest |*D*| across the

19 10
20

28 quartets ($|D| = 0.027 \pm 0.005$) compared to BUSCO ($|D| = 0.090 \pm 0.022$, Student's t $P <$

0.008), *D. sturtevanti* pseudo-reference ($|D| = 0.176 \pm 0.026$, Student's t $P < 4.77 \times 10^{-6}$), and

D. willistoni pseudo-reference ($|D| = 0.325 \pm 0.053$, Student's t $P < 5.97 \times 10^{-6}$). So, of the

four strategies, the use of syntenic blocks is the most conservative.

To consider cases where a true species tree cannot be inferred, we designed a test that

we call 2A2B. The test is an extension of the standard ABBA-BABA test in that it does not

only test the significance of deviation from parity of the ABBA and BABA patterns, but also

test the deviation from parity of each configuration to the AABB pattern. Therefore, the test

allows classifying each locus into one of the four categories along the reticulation-bifurcation

continuum given in Figure 1. Only loci with ≥20 evaluated sites were considered for this test.

We found assembly and pseudo-reference strategies significantly differed in multiple aspects

(Figure 3B). First, the extent of non-informative loci, *i.e.* loci with <20 evaluated sites, was

higher for assembly strategies. Similarly, the number of total evaluated sites ranged from

16,279-185,163 and 21,838-281,749 in BUSCO and syntenic strategies, respectively, to

364,472-1,177,118 and 153,238-452,562 in *D. sturtevanti* and *D. willistoni* pseudo-reference

genomes strategies, respectively. Second, assembly strategies had higher proportions of

complete reticulation (*i.e.* category i), averaging ~40 and 39% in BUSCO and syntenic

strategies, respectively, compared to pseudo-reference strategies, where it averaged ~9 and

20% in *D. sturtevanti* and *D. willistoni* pseudo-reference genomes strategies, respectively.

This effect was independent from the number of the evaluated sites ($r = -0.17$, $P = 0.40$, in the

synteny strategy), *i.e.* the high incidence of reticulation is not the result of low phylogenetic

signal leading to a soft polytomy. Third, partial reticulation and partial bifurcation (*i.e.*

categories ii and iii, respectively) predominated in the pseudo-reference strategies. Whereas

the proportion of these categories did not exceed ~7% in assembly-based strategies, it reached

~46% and 34% in *D. sturtevanti* and *D. willistoni* pseudo-reference genomes strategies,

21   11
22

respectively (Student's *t* for synteny vs. *sturtevanti* pseudo-reference strategies $P < 7.24 \times 10^{-9}$

). Because these two categories are related to hybridization and introgression, pseudo-

reference strategies therefore seem to provide stronger evidence for introgression, in

agreement to their higher |*D*| shown above. However, the phylogeny inferred from the

pseudo-reference alignments significantly differed from each other and from other analyses,

mainly in the position of the earliest branching subgroup, being the *sturtevanti* subgroup for

the *sturtevanti*-based pseudo-references and the *cordata* subgroup for the *willistoni*-based

pseudo-reference (Supplementary Figure S4). Pseudo-reference approaches, while providing

more data, may not reflect the true phylogenetic relationships between the species, hence

inflating the estimation of the extent of introgression.

Saltational episodes of reticulated evolution in the saltans *group*

Assembly-based 2A2B test identified three parts on the species tree with reticulation

indices (*i.e.* the proportion of blocks in categories iii and iv) exceeding 70% (Figure 3). At the

inter-subgroup level, high incidences of reticulation were observed in any combination that

involved representatives from at least two subgroups of the *cordata*, *elliptica* and *saltans*

subgroups. For the *sturtevanti* subgroup, ~90% of the blocks could not resolve the

relationships between *D. sturtevanti*, *D. lehrmanae* and the *dacunhai-milleri* clade, in

agreement with the conflicting topologies between the X, autosomes and mitochondrial loci

shown above. For the *saltans* subgroup, reticulation dominated (60-75%) in all comparisons

involving *D. austrosaltans*, and representatives of the *nigrosaltans-pseudosaltans* and the

septentriosaltans-prosaltans-saltans clades. However, not every subgroup with multiple

representatives showed excess reticulation, since for the *elliptica* subgroup, almost no

evidence for reticulate evolution was found whether *D. sturtevanti*, *D. neocordata* or any

species of the *saltans* subgroup were used as an outgroup.

23 12
24

To test if historical biogeography can explain the present incongruences, we mapped

current distribution of the studied species on the Bayesian X tree (see Methods). The

historical biogeography supported an origin of the *saltans* group around 16 million years

(myr) ago in the north-west of South America, which currently corresponds to the Amazonian

region (Figure 4A). At that time, this region was dominated by a vast wetland environment

known as the 'Pebas System.' The first radiation episode, *i.e.* the split between the ancestors

of the *cordata*, *elliptica* and *saltans* subgroups occurred circa 10 myr ago at this region,

coinciding with the formation of the Amazon River and its huge fluvial networks that led to

the transformation of the Pebas wetland into the dense and diverse Amazonian forests (Hoorn

et al. 2022). The second episode occurred nearly 3 myr ago, which correlates with the

geological formation of the isthmus of Panama (O'Dea et al. 2016), and the time of a

connection between the Western Amazonian and the Southern Atlantic rainforests during the

Pliocene (Batalha-Filho et al. 2013; Ledo and Colli 2017). This episode involved the

ancestors of the *nigrosaltans-pseudosaltans* and *septentriosaltans-prosaltans-saltans* clades.

Indeed, the first clade has a northern to central representative (*D. nigrosaltans*) and a southern

species (*D. pseudosaltans*), whereas the second clade has a northern representative (*D.*

saltans) and two central and central to southern species (*D. septentriosaltans* and *D.*

prosaltans, respectively). The third radiation episode involves the ancestor of the *sturtevanti*

subgroup and also occurred in the late Pliocene, ca. 2.5 myr ago, at the likely time of a

connection between the Western Amazonian and the Southern Atlantic rainforests.

Discussion

Towards a comprehensive phylogeny of the saltans species group

A large number of *Drosophila* genomes have been sequenced and used in

phylogenetic analyses (Khallaf et al. 2021; Kim et al. 2021; Li et al. 2022; Suvorov et al.

2022), but studies with comprehensive sampling of nearly all species in a group remain

relatively uncommon (Mai et al. 2020; Conner et al. 2021; Yusuf et al. 2022; Moreyra et al.

2023). Despite minor inconsistencies, our phylogenomic analysis of 15 species of the

Drosophila saltans species group produced a consistent picture of the relationships between

the five subgroups of this clade. All X, autosomal and mitochondrial phylogenies showed the

parasaltans subgroup as the first to diverge, followed by the *sturtevanti* subgroup, and later

by a clade comprising the *cordata*, *elliptica* and *saltans* subgroups, in which the position of

the *cordata* subgroup differed between nuclear and mitochondrial trees. This general picture

has not been previously proposed despite the tremendous number of phylogenetic

investigations of this group (Magalhães 1962; Throckmorton 1962; Throckmorton and

Magalhães 1962; O'Grady et al. 1998; Rodríguez-Trelles et al. 1999; Castro and Carareto

2004; de Setta et al. 2007; Yassin 2009; Souza et al. 2014; Roman et al. 2022; see

Supplementary Table S1).

While our analysis has shed light on the intricate evolutionary dynamics within this

clade, further sampling holds the potential to provide a more comprehensive understanding

into this complex evolutionary history. For example, the inclusion of *D. subsaltans*, *D.*

lusaltans, *D. cordata*, and *D. rectangularis* through whole-genome sequencing promises to

provide insight onto unresolved phylogenetic questions raised from previously published

observations on reproductive isolation and morphology (Magalhães 1962; Bicudo and Prioli

1978). These questions include the monophyly and positioning of the *parasaltans* and *cordata*

subgroups. Additionally, the inclusion of the insular species *D. lusaltans* which presents low

27   14

reproductive isolation (Bicudo 1973b), can bring new insights into the reticulation evolution

in the *saltans* subgroup. The *saltans* subgroup showed the most dramatic signal of cyto-

nuclear discordance and reticulated evolution. Bicudo (1973b) investigated reproductive

isolation among the seven then described species of this subgroup, and in a remarkably partial

agreement with our nuclear phylogenomic trees, she concluded that *D. pseudosaltans*, *D.*

nigrosaltans and *D. austrosaltans* showed more basal relationships compared to *D. lusaltans*,

D. septentriosaltans, *D. prosaltans* and *D. saltans*. Indeed, nearly all crosses among the last

four species produce fertile females with some even producing fertile females and males

(Bicudo 1973b). This behavioral porosity largely agrees with the high incidence of reticulate

evolution we report here for this subgroup.

Two widespread species of the *saltans* subgroup, *D. saltans* and *D. prosaltans*, show a

peculiar geographical disjunction. The discrimination between strains belonging to each

species has long been erroneous (Dobzhansky 1944; Mayr and Dobzhansky 1945; Spassky

1957; Magalhães 1962) and we showed here that their misidentification persists even in the

genomic era (Kim et al. 2021; Suvorov et al. 2022). Interestingly, Bicudo (1973b) provided

evidence for reproductive reinforcement between these two sister species; sympatric

populations in their junction zone in Costa Rica demonstrated stronger reproductive isolation

than allopatric populations of both species. We have only included one to a few geographical

lines from each species and a broader sampling to investigate the extent of their reproductive

isolation and genome porosity is strongly needed.

Intra- and inter-genomic conflicts impact the inference of phylogenetic patterns

Concatenation helped recovering a sex chromosome versus autosome conflict, similar

to the one described by Mai et al. (2020) for the *Drosophila nasuta* subgroup. Like these

authors, this conflict was limited to a single part of the tree, *i.e.* the relationship of *D. pulau* to

D. sulfurigaster sulfuricaster and *D. s. bilimbata* in the *nasuta* group and the placement of *D.*

lehrmanae in the *sturtevanti* subgroup. The peculiarities of sexual chromosomes such as their

lower effective number, different recombination and mutation rates, and greater exposition to

natural selection when found in hemizygosity, lead to higher rates of adaptive evolution of

sexual-linked genes compared with autosomal genes (*i.e.* faster-X evolution) and also to the

disproportional accumulation of genes related to reproductive isolation and Dobzhansky-

Muller hybrid incompatibilities (*i.e.* Haldane's rule) (Charlesworth et al. 2018). Altogether,

those characteristics are thought to be responsible for the resistance to hybridization in the

sexual chromosomes (Ellegren 2009; Qvarnström and Bailey 2009; Sankararaman et al. 2016;

Charlesworth et al. 2018; Seixas et al. 2018; Mai et al. 2020; Matute et al. 2020; Moran et al.

2021; Reilly et al. 2022; Skov et al. 2023; but see David et al. 2022).

The second conflict regards a significant disagreement between mitochondrial

(mtDNA) and nuclear data. Discordance between nuclear and mitochondrial genomes is a

well-documented phenomenon in the tree of life as highlighted by Toews and Brelsford

(2012). Several characteristics of mtDNA, such as being haploid and uniparentally inherited,

resulting in a fourfold reduction in effective population size when compared with autosomal

chromosome loci, affect its evolution. Cytoplasmic introgression has long been recognized in

Drosophila (Solignac et al. 1986; Ballard 2000; Llopart et al. 2014). In a recent population

study within the *willistoni* group, multiple mitochondrial introgressions were observed in *D.*

paulistorum populations (Baião et al. 2023). These included an ancient introgression with a

highly divergent mitochondrial type, followed by more recent events. While nuclear-

mitochondrial incompatibilities likely posed challenges, the study also suggested two possible

alternatives to overcome these challenges: a selective advantage provided by the

mitochondrial type itself. Or a non-selective factor, such as *Wolbachia,* a bacteria known to

modify the reproduction of its host, could facilitate a mitochondrial type fixation (Baião et al.

31 16
32

2023). Although, interesting results have been reported from population approaches, conflicts between nuclear and mitochondrial genomes have not been addressed in recent phylogenomic analyses in the Drosophilidae (Mai et al. 2020; Khallaf et al. 2021; Suvorov et al. 2022; Yusuf et al. 2022). The disagreement was particularly evident for the *saltans* subgroup, where it was most likely of recent origins, separating species that have diverged only 0.7 myr ago, *i.e. D. nigrosaltans* and *D. pseudosaltans*. Remarkably, the two mitotypes P and S do not correlate with the degree of reproductive isolation inferred by Bicudo (1973b), contrary to nuclear tree, indicating that cytoplasmic introgression in the *saltans* subgroup did not contribute to the evolution of reproductive isolation in this clade.

Of the three subgroups for which multiple species were sequenced, the *saltans* subgroup had the highest incidence of reticulation inferred by the 2A2B test. In this subgroup, multiple large chromosomal inversions are known to be shared among closely-related species (Dobzhansky and Pavan 1943; Cavalcanti 1948; Bicudo 1973a; Bicudo and Prioli 1978) and evidence for balancing on ancestral inversion has been demonstrated in a number of cases (Bicudo 1973a). Whether the high degree of reticulation in the *saltans* subgroup is associated with large ancestral inversions potentially absent in other bifurcating clades would require the future generation of chromosome-level assemblies for multiple *saltans* group species.

The 2A2B test on syntenic blocks helps distinguishing "soft" and "hard" introgressions

Our analyses test revealed that failure to account for a lack of phylogenetic signal in the tested loci and alignment on a reference genome both significantly inflate the estimation of the extent of interspecific introgression. Indeed, most studies that have provided strong evidence for introgression using site-specific patterns comparisons have usually aligned reads from multiple species to a single reference assembly (cf. studies compiled by Dagilis et al. 2021). Whereas such an approach would increase the power (Martin et al. 2015; Pease and

Hahn 2015), it also introduces biases due to paralogy, misalignments or absence of

collinearity among species, that are now known to bias phylogenetic inference itself

(Valiente-Mullor et al. 2021; Rick et al. 2023). Therefore, the most common practice now is

to use sets of clade-specific single-copy orthologs, such as the BUSCO genes, in inferring

phylogenies (Manni et al. 2021). Consequently, some recent introgression studies have

focused on such genes. For example, in a study of 155 genomes covering a wide range of

drosophilid lineages, Suvorov et al. (2022) analyzed these conserved genes. While finding

evidence for widespread introgression, they also showed that their discordance estimates were

highly sensitive to the length of the analyzed genes as well as by the slightest relaxation of

selective pressures. We attempted here to overcome this limitation by including larger

syntenic blocks, which in addition to conserved protein-coding sequences should also include

presumably neutral intronic and intergenic sites. Whereas, this approach showed the lowest D,

indicating that patterns predicted by introgression do not exceed 7% of the genome, it led to

only 1.5-fold increase in the number of analyzed sites, most likely because our assemblies

were based on only short reads. With the continuing progress in whole-chromosome

sequencing techniques and assembly-to-assembly alignment tools, a better understanding of

the extent of probably true or "hard" introgression events from artificial or "soft" ones may be

attained.

Phyletic radiations occur when isolation barriers simultaneously and rapidly evolve

between multiple populations of a species with a broad geographical or ecological breadth.

Consequently, genetic lineages will be incompletely sorted in the emerging species leading to

a lack of phylogenetic signals. This situation usually occurs following a rapid environmental

change or expansion into a new range with new empty niches. Indeed, our historical

biogeography approach shows that such situation might have favored the episodic signals of

low phylogenetic signals in the *saltans* group since the episodes coincided with either the

transition from wetlands into Amazonia forests in the Late Miocene or the connection of the

Amazonian forest with either Central American or the Southern Atlantic forests in the

Pliocene. These events have been proposed to be responsible for a wide range of rapid

radiations in Neotropical taxa, such as butterflies, amphibians and birds (Batalha-Filho et al.

2013; O'Dea et al. 2016; Ledo and Colli 2017; Hoorn et al. 2022). The dating and the

analyses of the extent of phylogenetic discordance in the genomes of such clades are strongly

needed to identify how much hard polytomies associate with these events may have occurred.

Materials and Methods

Sample collection, whole genome sequencing and assembly

We performed whole genome pool sequencing on female flies from 15 different

species from the *saltans* group, as well as three populations of *D. sturtevanti*. The specimens

used for sequencing were obtained from one or multiple strains, and detailed information

regarding the number of individuals and their collection locations can be found in

Supplementary Table S2. For all species, DNA extraction, PE library preparation using

NEBNext Protocol, and 75x coverage double-paired Illumina Novoseq sequencing were

conducted by End2End Genomics LLC, Davis, California. For all species except *D.*

neocordata, we assembled the genome using the Maryland Super Read Cabog Assembler

(MaSuRCA) (Zimin et al. 2013), which utilizes both the de Bruijn graph and overlap-layout-

consensus (OLC) methods to generate super-reads. K-mer based genome size was estimated

using KMC3 (Kokot et al. 2017) and GenomeScope v.2.0 (Ranallo-Benavidez et al. 2020). To

assess the assembly's completeness, we searched for single-copy-genes (SCG) using default

parameters in Busco5 (Simão et al. 2015) with the diptera_odb10 database (Kuznetsov et al.

2023). In addition to the sequenced flies, we also assembled Illumina reads the genomes of *D.*

37 19
38

448   *saltans*, *D. neocordata*, *D. prosaltans* and *D. sturtevanti* published by Kim et al. (2021)

449   (assembly numbers ASM1890357v1, ASM1890361v1, ASM1815127v1 and

450   ASM1815037v1, respectively), and we used the *D. neocordata* assembly published by Baião

451   et al. (2023) in all analyses.

453   *Phylogenomics: Nuclear single-copy-genes*

454   For phylogenomics analysis, SCG searches were carried out using 3,285 SCG from

455   diptera_odb10 database on Busco5 (Simão et al. 2015). SCG present in all species were kept

456   and aligned using the L-INS-i method implemented on MAFFT (Katoh and Standley 2013)

457   (mafft --localpair --maxiterate 1000 –adjustdirection).

458   Genomic data of different species of *Drosophila* support that genes tend to be situated

within the same Muller element across multiple species (see Schaeffer 2018). Taking this

gene linkage into account, we reconstructed five independent datasets (Muller elements A-E),

each comprising all SCG found in the respective Muller element. To achieve this, we

performed a tBlastn search against the *D. melanogaster*, and subsequently we concatenated

genes found within the same Muller element. Phylogenetic trees were then constructed using

maximum likelihood and Bayesian methods implemented in the softwares IQ-TREE (Nguyen

et al. 2015) and BEAST (Bouckaert et al. 2019), respectively. Additionally, maximum-

likelihood trees were generated for each gene, the output tree from each Muller element data-

set were used to reconstruct to species trees, using multi-species coalesce model implemented

in ASTRAL-III (Zhang et al. 2018).

Phylogenomics: Mitochondrial Genome

471   Mitochondrial genomes were assembled and annotated with MitoZ (Meng et al. 2019),

with the Megahit assembler (Li et al. 2015). In order to ensure the exclusion of nuclear-

embedded mitochondrial DNA sequences within the assembly, a strategic approach was

taken. Considering that mitochondrial reads are found in higher frequency than nuclear-

mitochondrial DNA sequences, the read subsampling were set to 0.5 gigabases (--

data_size_for_mt_assembly 0.5). The genes obtained from the mitochondrial genome were

aligned using the MAFFT alignment tool with the --auto parameter due to the close similarity

between sequences. Subsequently, the aligned genes were concatenated into a dataset for

phylogenetic analysis. The concatenated dataset served as the basis for reconstructing

phylogenetic trees using both Maximum Likelihood (ML) implemented in IQ-TREE (Nguyen

et al. 2015) and Bayesian Inference (BI) in BEAST (Bouckaert et al. 2019).

Identification of syntenic blocks

The 20 genomes of the *saltans* group, the 15 sequenced here and the 5 published by Kim et al.

(2021) and Baião et al. (2023) were preliminary annotated with Miniprot (miniprot -Iut16, Li

2023). The primary objective of this annotation was to accurately map proteins from the

robust and reliable genome annotation of *D. willistoni* (Clark et al. 2007). After the protein

mapping, the predicted gene loci were assessed to identify syntenic blocks present in the

Neotropical *Sophophora* (comprising *D. willistoni* and *saltans* group). The identification of

these blocks was based on gene order and orientation, achieved using an in-house Perl script.

First, this script compares the scaffolds' genes order and orientation between the reference

assemblies of *D. saltans* and *D. sturtevanti*, the synteny blocks were defined when all the

genes were found in the same order and orientation in both species. The identified collinear

blocks were then searched for the remaining genomes. Blocks with missing data, *i.e.* missing

genes for one or more species were subsequently removed, and the remaining blocks were

subjected to a size-based filtering with a threshold of ≥50kb. The selected blocks were aligned

using the Mafft software (Katoh and Standley 2013).

Pseudo-reference inference

Illumina reads were aligned on either the *D. willistoni* reference genome (Clark et al. 2007: 12) or the recently published *D. sturtevanti* assembly (Kim et al. 2021) using minimap2 (Li 2018). Minimap2 generated BAM files were assembled into a pseudo-reference for each species using the BEDTools software (Quinlan and Hall 2010). Each scaffold was then split into 100 kb-long windows, and phylogenetic analyses were conducted on each window separately using IqTree. A supertree was then inferred for each pseudo-reference genome approach using ASTRAL-III.

Quantifying reticulation using the 2A2B test

Combinations of 28 species quartets were evaluated to test reticulation, bi-allelic sited shared by pairs were searched in every locus, whether using SCG, syntenic blocks or pseudo-reference 100 kb-long windows. Loci with at least 20 informative sites were retained. For each locus, the occurrences AABB, ABBA or BABA patterns were counted, and three pairwise χ^2-based tests were conducted, denoted *D1*, *D2*, and *D3* as follows (D1 is reminiscent to Patterson's D):

$$D1 = (\textstyle\sum ABBA - \sum BABA)/(\sum ABBA + \sum BABA),$$

$$D2 = (\textstyle\sum AABB - \sum ABBA)/(\sum AABB + \sum ABBA), \text{ and}$$

$$D3 = (\textstyle\sum AABB - \sum BABA)/(\sum AABB + \sum BABA).$$

Each of these estimates is reminiscent to the classical Tajima's (1993) Relative Rate Test (RTT), which tests the difference in branch lengths from two ingroup taxa to a third outgroup taxon, with lengths estimated as counts of different sites. However, unlike RRT, branch lengths in our analyses are counted in terms of shared differences, *i.e.* singletons are not considered. Similar to RRT, we applied a chi-squared analysis to test for deviation from parity

for each parameter. According to the significance levels of the three estimates, a locus can be

classified under each of the four categories presented in Figure 1. The test is run by the 2A2B

perl script associated with this manuscript.

Historical biogeography

To determine the sampling locations of the evaluated species, we conducted searches

in TaxoDros (https://www.taxodros.uzh.ch/search/class.php). Additionally, we incorporated

sampling sites that we ourselves had conducted. The accuracy of our species identification

was confirmed through DNA barcoding. After inspection of the geographical points and

manual correction, we identified the most northern, southern, western, and eastern points for

each species. We used those points to reconstruct the ancestral geographical range by

conducting a BayesTraits (Meade and Pagel 2022) analysis for each cartesian point

separately. The analyses were carried out using the GEO model with the phylogenetic tree

generated reconstructed with the Muller element A (XL chromosome arm), 1,000,000 of

MCMC iterations and 25% burn-in. The divergence times were estimated using this tree

under Bayesian inference. The calibration point used was the split between *D. willistoni* and

D. sturtevanti at 17.5 myr, as estimated by Suvorov et al. (2022).

Acknowledgments

We would like to thank David Ogereau for his assistance and insights in genome assembly for

this study. We thank Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) for

the financial support to L.M.R. (Number processes: 95/06165-1, 2014/14059-0 and 2016/

11994-5) enabling us to collect and sequence many strains used here. We extend our thanks to

the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for their

45 23
46

support, specifically grant number 141545/2020-8, as well as the France Excellence Eiffel

Scholarship Program for funding C.P.'s PhD scholarship. *Illumina* sequencing of *D.*

neocordata was performed at the SNP&SEQ Technology Platform in Uppsala, Sweden,

which is part of the Swedish National Genomics Infrastructure and Science for Life

Laboratory. SNP&SEQ is supported by the Swedish Research Council and the Knut and Alice

Wallenberg Foundation. This project was partly supported by a grant from The Swedish

research council VR (2014-4353) to L.K. We want to thank the Nouragues research field

station (managed by CNRS), which benefits from "Investissement d'Avenir" grants managed

by Agence Nationale de la Recherche (AnaEE France ANR-11-INBS-0001; Labex CEBA

ANR-10-LABX-25-01). The project was partly funded by the Austrian Science Fund FWF

grant P28255-B22 to W.J.M. .

References

Bächli G. 2024. TaxoDros: the database on Taxonomy of Drosophilidae. Available from:

https://www.taxodros.uzh.ch/

Baião GC, Schneider DI, Miller WJ, Klasson L. 2023. Multiple introgressions shape

mitochondrial evolutionary history in *Drosophila paulistorum* and the *Drosophila*

willistoni group. *Molecular Phylogenetics and Evolution* 180:107683.

Ballard JWO. 2000. Comparative Genomics of Mitochondrial DNA in Members of the

Drosophila melanogaster Subgroup. *J Mol Evol* 51:48–63.

Batalha-Filho H, Fjeldså J, Fabre P-H, Miyaki CY. 2013. Connections between the Atlantic

and the Amazonian forest avifaunas represent distinct historical events. *J Ornithol*

154:41–50.

Bicudo HEMC. 1973a. Chromosomal polymorphism in the saltans group of Drosophila I. The

saltans subgroup. *Genetica* 44:520–552.

Bicudo HEMC. 1973b. Reproductive isolation in the saltans group of Drosophila. I. The

saltans subgroup. *Genetica* 44:313–329.

Bicudo HEMC. 1979. Reproductive isolation of the saltans group of Drosophila. IV. the

sturtevanti subgroup. *Revista Brasileira de Genética* 2:247–258.

Bicudo HEMC, Prioli AJ. 1978. Reproductive isolation in the saltans group of Drosophila II.

The parasaltans subgroup. *Genetica* 48:17–22.

Blischak PD, Chifman J, Wolfe AD, Kubatko LS. 2018. HyDe: A Python Package for

Genome-Scale Hybridization Detection. *Systematic Biology* 67:821–829.

Bouckaert R, Vaughan TG, Barido-Sottani J, Duchêne S, Fourment M, Gavryushkina A,

Heled J, Jones G, Kühnert D, Maio ND, et al. 2019. BEAST 2.5: An advanced

software platform for Bayesian evolutionary analysis. *PLOS Computational Biology*

15:e1006650.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.

BLAST+: architecture and applications. *BMC Bioinformatics* 10:421.

Castro JP de, Carareto CMA. 2004. *P* elements in the *saltans* group of *Drosophila*: a new

evaluation of their distribution and number of genomic insertion sites. *Molecular*

Phylogenetics and Evolution 32:383–387.

Cavalcanti AGL. 1948. Geographic Variation of Chromosome Structure in Drosophila

Prosaltans. *Genetics* 33:529–536.

Chang ES, Neuhof M, Rubinstein ND, Diamant A, Philippe H, Huchon D, Cartwright P.

2015. Genomic insights into the evolutionary origin of Myxozoa within Cnidaria.

Proceedings of the National Academy of Sciences 112:14912–14917.

Charlesworth B, Campos JL, Jackson BC. 2018. Faster-X evolution: Theory and evidence

from Drosophila. *Molecular Ecology* 27:3753–3771.

49   25

Clark AG, Eisen MB, Smith DR, Bergman CM, Oliver B, Markow TA, Kaufman TC, Kellis

M, Gelbart W, Iyer VN, et al. 2007. Evolution of genes and genomes on the

Drosophila phylogeny. *Nature* 450:203–218.

Conner WR, Delaney EK, Bronski MJ, Ginsberg PS, Wheeler TB, Richardson KM,

Peckenpaugh B, Kim KJ, Watada M, Hoffmann AA, et al. 2021. A phylogeny for the

Drosophila montium species group: A model clade for comparative analyses.

Molecular Phylogenetics and Evolution 158:107061.

Dagilis AJ, Peede D, Coughlan JM, Jofre GI, D'Agostino ERR, Mavengere H, Tate AD,

Matute DR. 2021. 15 years of introgression studies: quantifying gene flow across

Eukaryotes. :2021.06.15.448399. Available from:

https://www.biorxiv.org/content/10.1101/2021.06.15.448399v1

David JR, Ferreira EA, Jabaud L, Ogereau D, Bastide H, Yassin A. 2022. Evolution of

assortative mating following selective introgression of pigmentation genes between

two Drosophila species. *Ecology and Evolution* 12:e8821.

Dobzhansky TG, Pavan C. 1943. Studies on Brazilian species of Drosophila. *Boletim da*

Faculdade de Filosofia, Ciências e Letras da Universidade de São Paulo. Biologia

Geral. Separata 36:7–72.

Dobzhansky Th. 1944. Experiments on Sexual Isolation in Drosophila. *Proceedings of the*

National Academy of Sciences 30:335–339.

Doronina L, Churakov G, Shi J, Brosius J, Baertsch R, Clawson H, Schmitz J. 2015.

Exploring Massive Incomplete Lineage Sorting in Arctoids (Laurasiatheria,

Carnivora). *Molecular Biology and Evolution* 32:3194–3204.

Durand EY, Patterson N, Reich D, Slatkin M. 2011. Testing for Ancient Admixture between

Closely Related Populations. *Molecular Biology and Evolution* 28:2239–2252.

Ellegren H. 2009. The different levels of genetic diversity in sex chromosomes and

autosomes. *Trends in Genetics* 25:278–284.

Gagnon E, Hilgenhof R, Orejuela A, McDonnell A, Sablok G, Aubriot X, Giacomin L,

Gouvêa Y, Bragionis T, Stehmann JR, et al. 2022. Phylogenomic discordance suggests

polytomies along the backbone of the large genus Solanum. *American Journal of*

Botany 109:580–601.

Glor RE. 2010. Phylogenetic Insights on Adaptive Radiation. *Annual Review of Ecology,*

Evolution, and Systematics 41:251–270.

Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, Patterson N, Li H, Zhai

W, Fritz MH-Y, et al. 2010. A Draft Sequence of the Neandertal Genome. *Science*

328:710–722.

Hime PM, Lemmon AR, Lemmon ECM, Prendini E, Brown JM, Thomson RC, Kratovil JD,

Noonan BP, Pyron RA, Peloso PLV, et al. 2021. Phylogenomics Reveals Ancient

Gene Tree Discordance in the Amphibian Tree of Life. *Systematic Biology* 70:49–66.

Hoorn C, Boschman LM, Kukla T, Sciumbata M, Val P. 2022. The Miocene wetland of

western Amazonia and its role in Neotropical biogeography. *Botanical Journal of the*

Linnean Society 199:25–35.

Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. *Molecular Biology and Evolution*

30:772–780.

Khallaf MA, Cui R, Weißflog J, Erdogmus M, Svatoš A, Dweck HKM, Valenzano DR,

Hansson BS, Knaden M. 2021. Large-scale characterization of sex pheromone

communication systems in Drosophila. *Nat Commun* 12:4165.

Kim BY, Wang JR, Miller DE, Barmina O, Delaney E, Thompson A, Comeault AA, Peede D,

D'Agostino ER, Pelaez J, et al. 2021. Highly contiguous assemblies of 101

drosophilid genomes.Coop G, Wittkopp PJ, Sackton TB, editors. *eLife* 10:e66405.

Kokot M, Długosz M, Deorowicz S. 2017. KMC 3: counting and manipulating k-mer

statistics. *Bioinformatics* 33:2759–2761.

Kuznetsov D, Tegenfeldt F, Manni M, Seppey M, Berkeley M, Kriventseva EV, Zdobnov

EM. 2023. OrthoDB v11: annotation of orthologs in the widest sampling of

organismal diversity. *Nucleic Acids Research* 51:D445–D451.

Ledo RMD, Colli GR. 2017. The historical connections between the Amazon and the Atlantic

Forest revisited. *Journal of Biogeography* 44:2551–2563.

Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. 2015. MEGAHIT: an ultra-fast single-node

solution for large and complex metagenomics assembly via succinct de Bruijn graph.

Bioinformatics 31:1674–1676.

Li F, Rane RV, Luria V, Xiong Z, Chen J, Li Z, Catullo RA, Griffin PC, Schiffer M, Pearce

S, et al. 2022. Phylogenomic analyses of the genus Drosophila reveals genomic signals

of climate adaptation. *Molecular Ecology Resources* 22:1559–1581.

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34:3094–

3100.

Li H. 2023. Protein-to-genome alignment with miniprot. *Bioinformatics* 39:btad014.

Llopart A, Herrig D, Brud E, Stecklein Z. 2014. Sequential adaptive introgression of the

mitochondrial genome in Drosophila yakuba and Drosophila santomea. *Molecular*

Ecology 23:1124–1136.

Maddison W. 1989. Reconstructing Character Evolution on Polytomous Cladograms.

Cladistics 5:365–377.

Maddison WP. 1997. Gene Trees in Species Trees. *Systematic Biology* 46:523–536.

Madi-Ravazzi L, Segala LF, Roman BE, Alevi KCC, Prediger C, Yassin A, Hua-VAN AL,

Miller WJ. 2021. Integrative taxonomy and a new species description in the sturtevanti

subgroup of the Drosophila saltans group (Diptera: Drosophilidae). *Zootaxa*

4980:269292.

Magalhães LE de. 1962. Notes on the taxonomy, morphology, and distribution of the saltans

group of Drosophila, with descriptions of four new species. *The University of Texas*

Publication 2:135–154.

Magalhães LE de, Björnberg AJS. 1957. Estudo da genitália masculina de "Drosophila" do

grupo "saltans" (Diptera). *Revista Brasileira de Biologia* 17:435–450.

Mai D, Nalley MJ, Bachtrog D. 2020. Patterns of Genomic Differentiation in the Drosophila

nasuta Species Complex. *Molecular Biology and Evolution* 37:208–220.

Malinsky M, Matschiner M, Svardal H. 2021. Dsuite - Fast D-statistics and related admixture

evidence from VCF files. *Molecular Ecology Resources* 21:584–595.

Manni M, Berkeley MR, Seppey M, Zdobnov EM. 2021. BUSCO: Assessing Genomic Data

Quality and Beyond. *Current Protocols* 1:e323.

Martin SH, Davey JW, Jiggins CD. 2015. Evaluating the Use of ABBA–BABA Statistics to

Locate Introgressed Loci. *Molecular Biology and Evolution* 32:244–257.

Matute DR, Comeault AA, Earley E, Serrato-Capuchina A, Peede D, Monroy-Eklund A,

Huang W, Jones CD, Mackay TFC, Coyne JA. 2020. Rapid and Predictable Evolution

of Admixed Populations Between Two Drosophila Species Pairs. *Genetics* 214:211–

230.

Mayr E, Dobzhansky Th. 1945. Experiments on Sexual Isolation in Drosophila. *Proceedings*

of the National Academy of Sciences 31:75–82.

Meade A, Pagel M. 2022. Ancestral State Reconstruction Using BayesTraits. In: Luo H,

editor. Environmental Microbial Evolution: Methods and Protocols. Methods in

Molecular Biology. New York, NY: Springer US. p. 255–266. Available from:

https://doi.org/10.1007/978-1-0716-2691-7_12

Meng G, Li Y, Yang C, Liu S. 2019. MitoZ: a toolkit for animal mitochondrial genome

assembly, annotation and visualization. *Nucleic Acids Research* 47:e63.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A,

Lanfear R. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic

Inference in the Genomic Era. *Molecular Biology and Evolution* 37:1530–1534.

Moran BM, Payne C, Langdon Q, Powell DL, Brandvain Y, Schumer M. 2021. The genomic

consequences of hybridization.Wittkopp PJ, editor. *eLife* 10:e69016.

Moreyra NN, Almeida FC, Allan C, Frankel N, Matzkin LM, Hasson E. 2023. Phylogenomics

provides insights into the evolution of cactophily and host plant shifts in *Drosophila*.

Molecular Phylogenetics and Evolution 178:107653.

Morgan CC, Foster PG, Webb AE, Pisani D, McInerney JO, O'Connell MJ. 2013.

Heterogeneous Models Place the Root of the Placental Mammal Phylogeny.

Molecular Biology and Evolution 30:2145–2156.

Nascimento AP, Bicudo HEMC. 2002. Esterase Patterns and Phylogenetic Relationships of

Drosophila Species in the Saltans Subgroup (Saltans Group). *Genetica* 114:41–51.

Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: A Fast and Effective

Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular*

Biology and Evolution 32:268–274.

O'Dea A, Lessios HA, Coates AG, Eytan RI, Restrepo-Moreno SA, Cione AL, Collins LS, de

Queiroz A, Farris DW, Norris RD, et al. 2016. Formation of the Isthmus of Panama.

Science Advances 2:e1600883.

O'Grady PM, Clark JB, Kidwell MG. 1998. Phylogeny of the Drosophila saltans species

group based on combined analysis of nuclear and mitochondrial DNA sequences.

Molecular Biology and Evolution 15:656–664.

Owen CL, Miller GL. 2022. Phylogenomics of the Aphididae: Deep relationships between

subfamilies clouded by gene tree discordance, introgression and the gene tree anomaly

zone. *Systematic Entomology* 47:470–486.

Pease JB, Hahn MW. 2015. Detection and Polarization of Introgression in a Five-Taxon

Phylogeny. *Systematic Biology* 64:651–662.

Pélandakis M, Solignac M. 1993. Molecular phylogeny of Drosophila based on ribosomal

RNA sequences. *J Mol Evol* 37:525–543.

Philippe H, Brinkmann H, Lavrov DV, Littlewood DTJ, Manuel M, Wörheide G, Baurain D.

2011. Resolving Difficult Phylogenetic Questions: Why More Sequences Are Not

Enough. *PLOS Biology* 9:e1000602.

Philippe H, Derelle R, Lopez P, Pick K, Borchiellini C, Boury-Esnault N, Vacelet J, Renard

E, Houliston E, Quéinnec E, et al. 2009. Phylogenomics Revives Traditional Views on

Deep Animal Relationships. *Current Biology* 19:706–712.

Pick KS, Philippe H, Schreiber F, Erpenbeck D, Jackson DJ, Wrede P, Wiens M, Alié A,

Morgenstern B, Manuel M, et al. 2010. Improved Phylogenomic Taxon Sampling

Noticeably Affects Nonbilaterian Relationships. *Molecular Biology and Evolution*

27:1983–1987.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* 26:841–842.

Qvarnström A, Bailey RI. 2009. Speciation through evolution of sex-linked genes. *Heredity*

102:4–15.

61 31

Ranallo-Benavidez TR, Jaron KS, Schatz MC. 2020. GenomeScope 2.0 and Smudgeplot for

reference-free profiling of polyploid genomes. *Nat Commun* 11:1432.

Reilly PF, Tjahjadi A, Miller SL, Akey JM, Tucci S. 2022. The contribution of Neanderthal

introgression to modern human traits. *Current Biology* 32:R970–R983.

Rick JA, Brock CD, Lewanski AL, Golcher-Benavides J, Wagner CE. 2023. Reference

Genome Choice and Filtering Thresholds Jointly Influence Phylogenomic Analyses.

Systematic Biology:syad065.

Rodríguez-Trelles F, Tarrío R, Ayala FJ. 1999. Molecular Evolution and Phylogeny of the

Drosophila saltans Species Group Inferred from the *Xdh* Gene. *Molecular*

Phylogenetics and Evolution 13:110–121.

Roman BE, Madi-Ravazzi L. 2021. <p>Male terminalia morphology of sixteen

species of the Drosophila saltans group

Sturtevant (Diptera, Drosophilidae)</p>. *Zootaxa*

5061:523–544.

Roman BE, Santana DJ, Prediger C, Madi-Ravazzi L. 2022. Phylogeny of Drosophila saltans

group (Diptera: Drosophilidae) based on morphological and molecular evidence.

PLOS ONE 17:e0266710.

Romiguier J, Ranwez V, Delsuc F, Galtier N, Douzery EJP. 2013. Less Is More in

Mammalian Phylogenomics: AT-Rich Genes Minimize Tree Conflicts and Unravel

the Root of Placental Mammals. *Molecular Biology and Evolution* 30:2134–2144.

Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L,

Suchard MA, Huelsenbeck JP. 2012. MrBayes 3.2: Efficient Bayesian Phylogenetic

Inference and Model Choice Across a Large Model Space. *Systematic Biology*

61:539–542.

Sankararaman S, Mallick S, Patterson N, Reich D. 2016. The Combined Landscape of

Denisovan and Neanderthal Ancestry in Present-Day Humans. *Current Biology*

26:1241–1247.

Sayyari E, Mirarab S. 2018. Testing for Polytomies in Phylogenetic Species Trees Using

Quartet Frequencies. *Genes* 9:132.

Schaeffer SW. 2018. Muller "Elements" in Drosophila: How the Search for the Genetic Basis

for Speciation Led to the Birth of Comparative Genomics. *Genetics* 210:3–13.

Schrempf D, Szöllősi G. 2020. The Sources of Phylogenetic Conflicts. In: No commercial

publisher | Authors open access book. p. 3.1:1. Available from: https://hal.science/hal-

02535482

Seixas FA, Boursot P, Melo-Ferreira J. 2018. The genomic impact of historical hybridization

with massive mitochondrial DNA introgression. *Genome Biology* 19:91.

de Setta N, Loreto ELS, Carareto CMA. 2007. Is the Evolutionary History of the O-Type P

Element in the saltans and willistoni Groups of Drosophila Similar to That of the

Canonical P Element? *J Mol Evol* 65:715–724.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO:

assessing genome assembly and annotation completeness with single-copy orthologs.

Bioinformatics 31:3210–3212.

Simion P, Philippe H, Baurain D, Jager M, Richter DJ, Di Franco A, Roure B, Satoh N,

Quéinnec É, Ereskovsky A, et al. 2017. A Large and Consistent Phylogenomic Dataset

Supports Sponges as the Sister Group to All Other Animals. *Current Biology* 27:958–

967.

Skov L, Coll Macià M, Lucotte EA, Cavassim MIA, Castellano D, Schierup MH, Munch K.

2023. Extraordinary selection on the human X chromosome associated with archaic

admixture. *Cell Genomics* 3:100274.

Solignac M, Monnerot M, Mounolou J-C. 1986. Mitochondrial DNA evolution in

themelanogaster species subgroup ofDrosophila. *J Mol Evol* 23:31–40.

Souza TAJ, Noll FB, Bicudo HEMC, Madi-Ravazzi L. 2014. Scanning Electron Microscopy

of Male Terminalia and Its Application to Species Recognition and Phylogenetic

Reconstruction in the Drosophila saltans Group. *PLOS ONE* 9:e97156.

Spassky B. 1957. Morphological differences between sibling species of Drosophila.

University of Texas Publication 5721:48–61.

Sturtevant AH. 1942. The classification of the genus Drosophila, with descriptions of nine

species. *University of Texas Publication* 4213:5–51.

Sturtevant AH, Novitski E. 1941. THE HOMOLOGIES OF THE CHROMOSOME

ELEMENTS IN THE GENUS DROSOPHILA. *Genetics* 26:517–541.

Suh A. 2016. The phylogenomic forest of bird trees contains a hard polytomy at the root of

Neoaves. *Zoologica Scripta* 45:50–62.

Suvorov A, Kim BY, Wang J, Armstrong EE, Peede D, D'Agostino ERR, Price DK, Waddell

PJ, Lang M, Courtier-Orgogozo V, et al. 2022. Widespread introgression across a

phylogeny of 155 *Drosophila* genomes. *Current Biology* 32:111-123.e5.

Tajima F. 1993. Simple methods for testing the molecular evolutionary clock hypothesis.

Genetics 135:599–607.

Throckmorton LH. 1962. The problem of phylogeny in the genus Drosophila.Studies in

Genetics.II. *University of Texas Publication* 6205:207–343.

Throckmorton LH, Magalhães LE de. 1962. Changes with evolution of pteridine

accumulations in species of the saltans group of the genus Drosophila. *University of*

Texas Publication 6205:489–505.

Toews DPL, Brelsford A. 2012. The biogeography of mitochondrial and nuclear discordance

in animals. *Molecular Ecology* 21:3907–3930.

Townsend JP, Su Z, Tekle YI. 2012. Phylogenetic Signal and Noise: Predicting the Power of

a Data Set to Resolve Phylogeny. *Systematic Biology* 61:835.

Valiente-Mullor C, Beamud B, Ansari I, Francés-Cuesta C, García-González N, Mejía L,

Ruiz-Hueso P, González-Candelas F. 2021. One is not enough: On the effects of

reference genome for the mapping and subsequent analyses of short-reads. *PLOS*

Computational Biology 17:e1008678.

Walsh HE, Kidd MG, Moum T, Friesen VL. 1999. Polytomies and the Power of Phylogenetic

Inference. *Evolution* 53:932–937.

Whelan NV, Kocot KM, Moroz LL, Halanych KM. 2015. Error, signal, and the placement of

Ctenophora sister to all other animals. *Proceedings of the National Academy of*

Sciences 112:5773–5778.

Wickett NJ, Mirarab S, Nguyen N, Warnow T, Carpenter E, Matasci N, Ayyampalayam S,

Barker MS, Burleigh JG, Gitzendanner MA, et al. 2014. Phylotranscriptomic analysis

of the origin and early diversification of land plants. *Proceedings of the National*

Academy of Sciences 111:E4859–E4868.

Yassin A. 2009. Phylogenetic relationships among species subgroups in the Drosophila

saltans group (Diptera: Drosophilidae): Can morphology solve a molecular conflict.

Zool Res 30:225–232.

Yusuf LH, Tyukmaeva V, Hoikkala A, Ritchie MG. 2022. Divergence and introgression

among the virilis group of Drosophila. *Evolution Letters* 6:537–551.

Zhang C, Rabiee M, Sayyari E, Mirarab S. 2018. ASTRAL-III: polynomial time species tree

reconstruction from partially resolved gene trees. *BMC Bioinformatics* 19:153.

Zimin AV, Marçais G, Puiu D, Roberts M, Salzberg SL, Yorke JA. 2013. The MaSuRCA

genome assembler. *Bioinformatics* 29:2669–2677.

Figure legends

Figure 1 – Distribution of bi-allelic patterns along the reticulation-bifurcation continuum and the proposed 2A2B test. A) The distribution of bi-allelic sites of four species can generate three distinct topologies. B) Based of the frequency of these topologies in a genome fragment, this fragment can be categorized in (i) complete reticulation, (ii) incomplete reticulation, (iii) incomplete bifurcation, and (iv) complete bifurcation. The 2A2B test estimates significant deviation from parity in pairwise comparisons of the topologies. *: significant deviation, ns = non-significant.

Figure 2 – Phylogenomic conflict between the X chromosome, autosomes, and the mitochondria. A) Comparison of autosomal topology (left, represented by Muller element B) and X chromosome topology (right, represented by Muller element A) demonstrates overall agreement with minor incongruence. B) Mitochondrial-nuclear disagreement highlights stronger incongruence between mitochondrial topology (left) and sexual chromosome topology (right). All trees were inferred through Bayesian analysis. Divergence time estimation (in million years ago, myr) for the sexual chromosome topology is provided (left). Nodes are numbered for subsequent analyses. Maximum-likelihood and ASTRAL-III trees are provided in Supplementary Figures S2 and S3, respectively. Posterior probabilities of all nodes were equal to1.

Figure 3 – Extent of phylogenomic discordance across 28 species quartets as measured by A) Patterson's D and B) the 2A2B test, using four locus-definition strategies (see text). For both tests, assembly-based strategies have lower average D and provide stronger evidence for reticulation at three episodes. Colors of bars in B) correspond to the four categories presented in Figure 1, with grey bars corresponding to loci with less than 20 evaluated bi-allelic sites.

Figure 4 – Historical biogeography of the *saltans* species. Centroid geographical coordinates

for each species were inferred and ancestral coordinates were inferred using BayesTraits on

the Bayesian X chromosome chronogram given in Figure 2B. Internal nodes are projected on

the current map of South and Central America. Three episodes of rapid radiation were

identified by the 2A2B test (Figure 3B) and are identified here with their corresponding

possible paleogeographical events, with photos of adult males of four representative species

shown. Branch colors in the tree correspond to the subgroups colors given in Figure 2.

List of Supplementary Figures

Figure S1 – Phylogenetic relationships inferred using BEAST from BUSCO genes

concatenated according to their Muller's element.

Figure S2 – Phylogenetic relationships inferred using IqTree-2 from BUSCO genes

concatenated according to their Muller's element.

Figure S3 – Phylogenetic relationships inferred using ASTRAL-III from BUSCO genes

concatenated according to their Muller's element.

Figure S4 – Phylogenetic relationships inferred using ASTRAL-III from 100-kb long

windows for pseudo-references produced by aligning short reads either to A) the ingroup *D.*

sturtevanti or B) the outgroup *D. willistoni* reference assemblies.

List of Supplementary Tables

Table S1 – A synopsis of phylogenetic relationships previously published in the *Drosophila*

saltans species group. Species subgroups are represented by the first two letters in capital,

whereas species names are represented by the first three letter in small.

Table S2 – Source and genome assembly statistics including busco scores for the genomes

used in this study.

77 39
78

A

(i)	(ii)	(iii)	(iv)
Complete reticulation	Incomplete reticulation	Incomplete bifurcation	Complete bifurcation

B

Frequency of patterns of bi-allelic sites

Outgroup	A	A	B
Species 1	A	B	A
Species 2	B	B	B
Species 3	B	A	A

The reticulation-bifurcation continuum

A

Assembly-based Pseudo-reference-based

BUSCO Synteny *sturtevanti* *willistoni*

B

Unevaluated loci | Complete reticulation | Incomplete reticulation | Inomplete bifurcation | Complete bifurcation

~10 myr: Formation of the Amazonian River and the split of the *cordata-elliptica-saltans* subgroups

~3 myr: Formation of Panama Isthmus and radiation of the *saltans* subgroup

~2.5 myr: Connection between the Amazonian and Atlantic Forests and radiation of the *sturtevanti* subgroup

D. neocordata

D. emarginata

D. prosaltans

D. sturtevanti