Acute avoidance of hydrogen sulfide is modulated by external and internal states in *C.*

elegans

Longjun Pu[1, 2, 3], Lina Zhao[1, 2, 3], Jing Wang[1, 2, 3], Johan Henriksson[1, 4, 5], Patrick Laurent[6],

Changchun Chen[1, 2, 3*]

[1]Department of Molecular Biology, Umeå University, Umeå, Sweden

[2]Umeå Centre for Molecular Medicine, Umeå University, Umeå, Sweden

[3]Wallenberg Centre for Molecular Medicine, Umeå University, Umeå, Sweden

[4]The Laboratory for Molecular Infection Medicine Sweden (MIMS), Umeå University, Umeå,

Sweden

[5]Integrated Science Lab (Icelab), Umeå University, Umeå, Sweden

[6]Laboratory of Neurophysiology, ULB Neuroscience Institute (UNI), Université Libre de

Bruxelles (ULB), Belgium

*Corresponding author

E-mail: changchun.chen@umu.se

Abstract

Hydrogen sulfide (H_2S) can act as an energy source, a poison and a gasotransmitter in

organisms. We used the robust locomotory responses to H_2S in *Caenorhabditis elegans* to

delineate the molecular mechanisms governing sensory and adaptive responses to H_2S

exposure. We found that *C. elegans* exhibited transiently increased locomotory activity and

turning behavior as a strategy to escape the noxious H_2S stimulation. The behavioral

responses to H_2S were modulated by a complex network of signaling pathways, including

cyclic GMP signaling in ciliated sensory neurons, calcineurin, nuclear hormone receptors, to

the major starvation regulators such as insulin and TGF-β signaling. The response to H_2S was

substantially affected by the ambient O_2 levels and their prior experience in low

O_2 environments, suggesting an intricate interplay between O_2 and H_2S sensing mechanisms.

Prolonged exposure to H_2S robustly evoked H_2S detoxification coupled with reduced

locomotory response to the subsequent H_2S challenges. Intriguingly, the expression of genes

involved in iron homeostasis, including *ftn-1* and *smf-3*, was substantially modified in

exposure to H_2S, implying that labile iron levels are affected by H_2S. In support of this, iron

supplement significantly bolstered the behavioral response to H_2S. In addition, mitochondria,

one of the central hubs for H_2S metabolism, played a crucial role in adaptive responses to

H_2S. In summary, our study provides molecular insights into the mechanisms through

which *C. elegans* detects, modulates, and adapts its response to H_2S.

Introduction

The detection and response to chemical signals are crucial to interact effectively with the environment in various organisms, spanning from bacteria to humans. During the late Proterozoic era, the rise in oxygen (O_2) levels eliminated hydrogen sulfide (H_2S) from most of the habitats. Yet, low O_2 levels together with high concentrations of H_2S and carbon dioxide (CO_2) can still persist in enclosed environments where bacteria actively decompose organic matter (Olson & Straub, 2016). H_2S can easily penetrate biological membranes and interfere with various cellular processes, leading to toxicity. One of its detrimental effects is the disruption of cellular respiration by binding to cytochrome c oxidase (COX) and inhibiting the mitochondrial electron transport chain (Cooper & Brown, 2008; Khan et al., 1990; Nicholls & Kim, 1982b).

Animals inhabiting environments rich in H_2S have evolved mechanisms to either avoid or adapt to high concentrations of H_2S, providing them with survival advantages. Species that encounter periodic increases in H_2S often possess sensing abilities and exhibit avoidance behaviors. For example, benthic species undertake daily vertical migrations between sulfidic and non-sulfidic waters (Abel, Koenig, & Davis, 1987; Salvanes, Utne-Palm, Currie, & Braithwaite, 2011). Certain metazoan species have seized the ecological opportunities presented by H_2S-rich environments through the development of cellular adaptations, allowing them to occupy a unique ecological niche that is inaccessible to the other organisms. Livebearing fish species, for instance, have evolved COX subunits that are more resistant to H_2S and are more efficient in H_2S detoxification (Kelley et al., 2016; Pfenninger et al., 2014).

Free living nematodes thrive in decaying matter where they are exposed to significant fluctuations in O_2, CO_2, and H_2S levels over short distances (Rodriguez-Kabana, Jordan, & Hollis, 1965). Nematodes like *C. elegans* have evolved the ability to utilize these gasous stimuli to navigate their environment, locate bacterial food sources, or escape the dangers (Bretscher, Busch, & de Bono, 2008; Carrillo, Guillermin, Rengarajan, Okubo, & Hallem, 2013). Acute exposure to H_2S triggers transiently increased locomotion in *C. elegans* (Budde

& Roth, 2011). However, prolonged exposure to H_2S induces cellular adaptation to survive

the toxic environment. *C. elegans* can tolerate H_2S levels up to 50 ppm, but the concentrations

exceeding 150 ppm prove lethal (Budde & Roth, 2010, 2011; Fawcett, Hoyt, Johnson, &

Miller, 2015; Horsman, Heinis, & Miller, 2019; Miller, Budde, & Roth, 2011a; Miller &

Roth, 2007). Extended periods of low O_2 and/or high H_2S exposure stabilize the hypoxia-

inducible factor HIF-1, leading to transcriptional reprogramming (Budde & Roth, 2010).

Through HIF-1-dependent and independent mechanisms, acclimation of *C. elegans* to 50 ppm

H_2S extends lifespan and improves the response to hypoxia-induced protein aggregation

(Budde & Roth, 2010; Fawcett et al., 2015; Horsman et al., 2019; Miller & Roth, 2007;

Qabazard et al., 2014).

In this study, we investigated how the nematode *C. elegans* responds acutely to avoid

an increase of H_2S levels, as well as how it adapts at the molecular and physiological levels if

they fail to escape the H_2S exposure.

Results

cGMP signaling in ASJ cilia contributes to H_2S avoidance

When encountering aversive cues such as noxious gases, *C. elegans* initiates a pirouette

followed by an increase of its locomotory activity to escape the noxious stimuli. When

acutely exposed to 150 ppm H_2S, *C. elegans* exhibited an increased locomotory speed and

enhanced turning behavior, while decreasing the frequency of reversals, as a strategy to

escape the noxious stimuli (***Figure 1—figure supplement 1A–C***). The maximum speed was

reached after 6–8 minutes in H_2S, and returned to the baseline over a period of approximately

1 hour (***Figure 1—figure supplement 1D***). The reduced locomotory activity after 2 hours in

H_2S was fully reversible if these animals were immediately challenged by a different stimulus

such as hypoxia (***Figure 1—figure supplement 1E***). The magnitude of the avoidance

responses appeared to be concentration dependent. Under our assay conditions, acute

behavioral response to 50 ppm H_2S was minimal and lacked consistency, whereas the

response to 150 ppm H_2S was robust and reproducible (***Figure 1—figure supplement 1F***).

To gain molecular insight into H_2S evoked locomotory response, we performed a

candidate screen for mutants that failed to increase their locomotory speed when H_2S levels

acutely increased to 150 ppm (***Supplementary file 1***). We discovered that mutants defective

in ciliogenesis exhibited diminished H_2S responses (***Figure 1A–C***). These included *daf-19*

mutants lacking cilia, *dyf-3* mutants with reduced cilia length, and *dyf-7* mutants where

ciliogenesis occurs but the cilia are not anchored to their sensilla and are not exposed to the

external environment (Heiman & Shaham, 2009; Murayama, Toh, Ohshima, & Koga, 2005;

Starich et al., 1995; Swoboda, Adler, & Thomas, 2000). Our observations suggest that cilia

exposure to the external environment is necessary for H_2S detection. Cilia-mediated sensory

responses often involve the activation of guanylate cyclases and the opening of cGMP-gated

channels (Ferkey, Sengupta, & L'Etoile, 2021). We observed attenuated H_2S responses in *tax-*

4 or *tax-2* mutants, lacking the key subunits of cilia-enriched cGMP-gated channels (***Figure***

1D and E). Comparable observations have also been reported in acute response to CO_2,

suggesting that locomotory responses to CO_2 and H_2S are similar (Bretscher et al., 2008;

Hallem et al., 2011; Hallem & Sternberg, 2008). Consistent with this, mutations in genes

encoding the calcineurin subunits TAX-6 or CNB-1, or nuclear hormone receptor NHR-49,

impaired both H_2S and CO_2 responses (Hallem & Sternberg, 2008) (***Figure 1—figure***

supplement 2A–C). Additionally, locomotory response to H_2S was also repressed following a

24-hour period of starvation, as observed for CO_2 (***Figure 1—figure supplement 2D***)

(Bretscher et al., 2008; Hallem & Sternberg, 2008). In *C. elegans*, the regulation of

starvation-responses involves the reduced activity of two major pathways: insulin and TGF-β

signaling. Disrupting the genes encoding the insulin receptor DAF-2 or the TGF-β ligand

DAF-7 eliminated locomotory activity in response to acute H_2S (***Figure 1F and G, Figure***

1—figure supplement 2E). Insulin and TGF-β signaling involve the inhibition of

downstream transcription factors DAF-16 or DAF-3 respectively. H_2S avoidance defects of

daf-2 and *daf-7* mutants were fully suppressed in *daf-2; daf-16* or in *daf-7; daf-3* double

mutants (***Figure 1F and G***).

These observations prompted us to further investigate if the same molecular and

circuit mechanisms underlay the acute avoidance to CO_2 and H_2S. The response to CO_2

requires receptor guanylate cyclase GCY-9 (Hallem et al., 2011). Interestingly, GCY-9 was

dispensable for the locomotory responses to H_2S (***Figure 1H***). In addition, animals responded

more rapidly to CO_2 than to H_2S, but with a reduced magnitude (***Figure 1I***), implying that

distinct mechanisms likely drive these two responses. In a comprehensive survey of all

guanylate cyclase mutants, the *daf-11* strain emerged as the only mutant showing defects in

the avoidance of H_2S (***Figure 1D, Supplementary file 1***). Selective expression of *daf-11* in

ASJ neurons, but not in other neurons, partially restored the locomotory response to acute

H_2S (***Figures 1J, Figure 1—figure supplement 2F–I***). Consistently, the H_2S response defect

of *tax-4* mutants was also rescued by selective expression of its cDNA in ASJ neurons

(***Figure 1K***), suggesting that the receptor guanylate cyclase DAF-11 and the CNG channel

TAX-4 partially act in ASJ neurons to promote acute response to H_2S. This differed from the

site of action of the CO_2 sensor GCY-9, which primarily operates in BAG neurons (Hallem et

al., 2011). Taken together, these observations suggest that common signaling pathways

modulate the locomotory responses to both H_2S and CO_2, although the molecular and circuit

underpinning of H_2S and CO_2 sensing are different.

Activation of the O_2 sensing circuit antagonizes H_2S avoidance

The sensory responses of wild *C. elegans* are adapted to their living environment, which is

characterized by low O_2 levels and high concentrations of CO_2. In comparison to the

domesticated lab strain N2, wild isolates of *C. elegans* exhibit a strong response to high O_2

but are relatively indifferent to CO_2 (Beets et al., 2020; Bretscher et al., 2008; Hallem &

Sternberg, 2008). Local H_2S concentrations could also be significantly higher in decomposing

substances where wild *C. elegans* thrives. We suspected that wild strains could display a

modified response to H_2S in comparison to standard laboratory N2 animals. Indeed, wild

isolates CB4855, CB4856, and CB4858 exhibited attenuated responses to H_2S (***Figure 2A,***

Figure 2—figure supplement 1A). One predominant genetic trait in these strains is a

naturally occurring variation of neuropeptide receptor NPR-1 215F, which is less active than

NPR-1 215V in N2 animals (de Bono & Bargmann, 1998). This led us to investigate whether

the NPR-1 signaling plays a role in modulating acute avoidance of H_2S. Mutating *npr-1* in N2

strain background was sufficient to abolish locomotory responses to H_2S, as observed in the

wild isolates (***Figure 2B***). NPR-1 primarily acts within the RMG interneurons to modulate

responses to environmental stimuli by reducing signal output from these neurons (Laurent et

al., 2015; Macosko et al., 2009). The selective expression of the highly active *npr-1(215V)*

variant in RMG neurons effectively restored H_2S evoked locomotory response in *npr-1*

mutant worms (***Figure 2C***). Furthermore, optogenetic stimulation of RMG interneurons using

channelrhodopsin-2 (ChR2) significantly dampened locomotory response to H_2S of N2

animals, suggesting high RMG activity inhibits H_2S responses (***Figure 2—figure supplement***

1B).

The sensation of 21% O_2 involves the soluble guanylate cyclases GCY-35/GCY-36

and the downstream cyclic nucleotide gated (CNG) channels TAX-4/TAX-2 (Busch et al.,

2012; Cheung, Cohen, Rogers, Albayram, & de Bono, 2005; Couto, Oda, Nikolaev, Soltesz,

& de Bono, 2013; Gray et al., 2004; Laurent et al., 2015; Persson et al., 2009; Zimmer et al.,

2009). We examined if disrupting O_2 sensing machinery had the same effect as the inhibition

of RMG signaling by NPR-1(215V). We found that silencing O_2 sensing neurons by mutating

gcy-35 and *tax-4*, or ablating these neurons by cell-specific expression of the pro-apoptotic

gene *egl-1* restored H_2S evoked behavioral response in *npr-1* null mutants (***Figure 2D–F***).

Specific expression of *gcy-35* cDNA in URX, AQR, PQR was sufficient to repress acute

response to H_2S in *npr-1; gcy-35* double mutants (***Figure 2G***). Finally, enhancing the

presynaptic activities of URX, AQR, PQR neurons by the expression of a gain-of-function

PKC-1(A160E) (Dekker, Mcintyre, & Parker, 1993; Hiroki et al., 2022) significantly

dampened H_2S regulated locomotory activity (***Figure 2—figure supplement 1C***). Taken

together, these observations indicate that activation of O_2 sensing circuit is inhibitory to H_2S

evoked locomotory response.

H_2S exposure reprograms gene expression in *C. elegans*

Prolonged exposure to H_2S has a dramatic impact on the physiology of *C. elegans*: 50 ppm

H_2S exposure extends its lifespan, while 150 ppm H_2S is lethal (Budde & Roth, 2010, 2011;

Fawcett et al., 2015; Horsman et al., 2019; Miller & Roth, 2007 Miller, 2011 #5). We

conducted a comparative analysis of the transcriptome profiles in animals exposed to either

50 ppm or 150 ppm H_2S for the durations of 1 hour, 2 hours, or 12 hours. Our RNA-seq

analysis revealed that a one-hour exposure to either 50 ppm or 150 ppm H_2S was sufficient to

trigger significant changes in gene expression, with 518 or 304 genes showing differential

expression, respectively (***Figure 3A, Supplementary file 2, p<1e-10***). Genes induced by 1-

hour and 2-hour H_2S exposure displayed a considerable overlap (***Figure 3B, Figure 3—***

figure supplement 1A). Gene Ontology (GO) analysis revealed that biological processes such

as defense against bacteria and cysteine biosynthesis from L-serine were immediately

activated upon H_2S exposure (***Figure 3C and D, Figure 3—figure supplement 1B and C***).

As expected, we observed a robust induction of genes involved in H_2S detoxification (***Figure***

3E and F) (Miller, Budde, & Roth, 2011b; Niu et al., 2011; Vora et al., 2022). H_2S is

detoxified in mitochondria by sulfide:quinone oxidoreductase SQRD-1 to persulfide, which is

then metabolized by sulfur dioxygenase ETHE-1 to generate sulfite. Thiosulfate transferase

further oxidizes sulfite to produce thiosulfate (Hildebrandt & Grieshaber, 2008). Glutathione

S-transferases (GSTs) remove accumulated sulfur molecules during H_2S oxidation (Jackson,

Melideo, & Jorns, 2012). Notably, *gst-19* and *sqrd-1* were among the most significantly

upregulated genes after one or two hours of exposure to either 50 ppm or 150 ppm H_2S, while

ethe-1 showed a weak increase (***Figure 3E and F, Figure 3—figure supplement 1D and E,***

Supplementary file 2). H_2S activates the HIF-1 pathway through the interaction of

sulhydrylase/cysteine synthase CYSL-1 with the HIF-1 proline hydroxylase EGL-9 (Ma,

Vozdek, Bhatla, & Horvitz, 2012). We found that a proportion of genes with HIF-1 regulated

promoters displayed increased expression after 1 or 2 hours of H_2S exposure (***Figure 3—***

figure supplement 1F) (Miller et al., 2011b; Vora et al., 2022). These genes included the

above-mentioned detoxifying enzymes GST-19, SQRD-1, and ETHE-1, as well as enzymes

involved in cysteine metabolism, CYSL-1, CYSL-2, and CDO-1 (***Supplementary file 2***).

Mutations increasing SKN-1 activity bypass the requirement for HIF-1 to survive in high H_2S

(Horsman et al., 2019). Consistent with early observations (Miller et al., 2011a), we also

observed that a set of SKN-1 targets were differentially expressed in H_2S (***Figure 3—figure***

supplement 1G) (Niu et al., 2011). This included the heat shock protein encoding genes *hsp-*

16.2 and *hsp-16.41* (***Figure 3E and F, Figure 3—figure supplement 1D and E***).

Surprisingly, we also observed a rapid increase in the expression of genes associated with

intracellular H_2S production, such as *semo-1* and *mpst-3* genes (***Figure 3E and F, Figure***

3—figure supplement 1D and E) (Philipp et al., 2022; Qabazard et al., 2014). However,

these data aligned with earlier observations made in sulfide spring fish (Kelley et al., 2016;

Mathew, Schlipalius, & Ebert, 2011; Qabazard et al., 2014).

When exposed to the H_2S for 12 hours, the differentially expressed genes showed

reduced overlap with those identified at earlier time points (***Figure 3B, Figure 3—figure***

supplement 1A). This suggests that different defense strategies against H_2S might be

employed at various stages of exposure. Although similar sets of genes were induced after 1h

or 2h at 50 ppm and 150 ppm H_2S, the overlap between 50ppm and 150ppm decreased after

12 hours exposure (***Figure 3—figure supplement 1H–J***). This observation suggests that

defense mechanisms might also differ for prolonged exposure at low and high levels of H_2S.

In particular, we noted that 150 ppm H_2S specifically triggered the expression of the HIF-1

regulator *rhy-1* and the HIF-1 target *nhr-57* at all exposure time points, which was less

pronounced in response to 50 ppm H_2S (***Figures 3E and F, Figure 3—figure supplement***

1D, E, K and L).

Acute response to H₂S is modulated by HIF-1 signaling

Consistent with early observations (Budde & Roth, 2010), our RNA-seq data suggested that

H₂S exposure activates HIF-1 pathway (*Figure 3—figure supplement 1F*). Stabilization of

HIF-1 by H₂S is critical for animals to survive H₂S exposure by the induction of

detoxification genes as well as other adaptative responses (Ma et al., 2012). Such an

adaptation is usually accompanied with the reduction in sensory sensitivity to this stressor.

Therefore, we suspected that HIF-1 stabilization might dampen the acute response to H₂S.

Under normoxia conditions, the conserved proline-4-hydroxylase PHD/EGL-9 hydroxylates

HIF-1, which is subsequently recognized by the von Hippel-Lindau (VHL) tumor suppressor

protein for degradation (Kaelin & Ratcliffe, 2008; Semenza, 2010). We found that disrupting

hif-1 resulted in extremely transient behavioral response to H₂S characterized by an initial

bout of activity and a rapid decline of locomotory speed to the basal level (*Figure 4A*). While

eliminating PHD/EGL-9 or VHL activity, which stabilizes HIF-1, the locomotory response to

H₂S was completely inhibited (*Figure 4B*). *egl-9; hif-1* or *vhl-1; hif-1* double mutants had the

identical response as *hif-1* mutants (*Figure 4C and D*), confirming that mutating *egl-9* or *vhl-1* inhibits acute response to H₂S by activating HIF-1. Acclimating animals in 1% O₂ for more

than 12 hours also significantly decreased acute response to H₂S (*Figure 4E*). In addition,

animals carrying the non-degradable form of HIF-1 in all neurons were defective in H₂S

evoked locomotory responses (*Figure 4F*). Taken together, these observations suggest that

constitutive activation of HIF-1 promotes the adaption to H₂S and inhibits the behavioral

responses to H₂S stimulation.

The transient behavioral response to H₂S in *hif-1* mutants suggested that animals

deficient of HIF-1 can still detect the presence of H₂S. This observation together with the

importance of HIF-1 signaling in H₂S detoxification provoked us to explore whether

detoxification enzymes were also required for the sustained locomotory activity in H₂S.

Similar to *hif-1* mutants, both *sqrd-1* and *ethe-1* mutants displayed a transient response to H₂S

(*Figure 4G and H*). We next examined the contribution of other upregulated genes in

response to H₂S, including the genes encoding sulfhydrylase/cysteine synthases CYSL-1,

CYSL-2, and CYSL-3 as well as SEMO-1. Disrupting CYSLs and SEMO-1 encoding genes

phenocopied *hif-1* or *sqrd-1* mutants (***Figure 4I and J***). Our data suggest that while H_2S

detoxification system is not essential for H_2S sensing, it is required for preserving animals'

locomotory activity in the presence of H_2S.

Labile iron pool is critical to sustain the locomotory activity in H_2S

C. elegans tightly regulates iron homeostasis through the control of iron transporter SMF-3,

as well as the regulation of iron storage protein Ferritins FTN-1 and FTN-2 (Cha'on et al.,

2007; Gourley, Parker, Jones, Zumbrennen, & Leibold, 2003; Kim, Cho, Yoo, & Ahnn, 2004;

Pekec et al., 2022). Among the most differentially regulated genes, *ftn-1* appeared to be

consistently downregulated by H_2S (***Figure 5A and B, Figure 5—figure supplement 1A–D***).

The inhibited expression of *ftn-1* and *ftn-2* as well as the elevated expression of *smf-3* (***Figure***

5A and B, Figure 5—figure supplement 1A–D) prompted us to explore if labile iron levels

play a role in behavioral response to H_2S exposure. The labile iron pool is chelatable and

redox-active. In the presence of iron chelator 2,2'-dipyridyl (BP), H_2S evoked locomotory

response was fully inhibited (***Figure 5C***). This was not caused by the general sickness of BP-

exposed animals as they responded robustly to acute hypoxia (***Figure 5—figure supplement***

1E). By contrast, the supplementation of ferric ammonium citrate (FAC) significantly

prolonged animals' locomotory activity in H_2S (***Figure 5C***). In addition, disrupting *ftn-1*,

which was expected to increase labile iron levels (Anderson & Leibold, 2014; Romney,

Newman, Thacker, & Leibold, 2011), promoted the locomotory activity of animals in H_2S

(***Figure 5D***), Conversely, mutating *smf-3*, which impaired iron uptake (Anderson & Leibold,

2014; Romney et al., 2011), led to a faster decline of locomotory activity in H_2S (***Figure 5D***).

Furthermore, depleting labile iron pool by *ftn-1* overexpression resulted in an attenuated

response to H_2S (***Figure 5E***), while it did not affect acute response to hypoxia (***Figure 5—***

figure supplement 1F). These data suggests that high levels of free iron sustain locomotory

activity in H_2S, whereas low iron levels inhibit behavioral responses to H_2S. It has been

reported that hypoxia inhibits the expression of *ftn-1* and *ftn-2*, while upregulating *smf-3*

expression (Ackerman & Gems, 2012; Gourley et al., 2003; Romney et al., 2011).

Additionally, *hif-1* mutants have been reported to display reduced intracellular iron content

and increased susceptibility to iron starvation (Rajan et al., 2019; Romney et al., 2011). We

explored if the locomotory defect of *hif-1* mutants in H_2S could be explained by iron

deficiency (Rajan et al., 2019; Romney et al., 2011). The presence of FAC significantly

improved H_2S evoked locomotory activity of *hif-1* mutants (***Figure 5F***). In addition,

disrupting *ftn-1* partially suppressed the locomotory defect of *hif-1* mutants in H_2S (***Figure***

5G). These data confirm that labile iron pool is critical to maintain the locomotory activity of

animals in H_2S.

Iron deficiency causes mitochondrial dysfunction including decreased cytochrome

concentration and respiratory efficiency (Dallman, 1986; Masini, Salvioli, et al., 1994;

Masini, Trenti, et al., 1994; Walter et al., 2002). H_2S is known to exert both positive and

negative effects on the mitochondria, which are not only the primary targets of H_2S but also

represent one of the main cellular hubs of H_2S detoxification (Cooper & Brown, 2008; Khan

et al., 1990; Nicholls & Kim, 1982a). We probed the possibility that impairing mitochondrial

function might alter locomotory response to H_2S. To this end, we tested mutants with the

disruption of *gas-1*, *clk-1*, *mev-1,* or *isp-1* genes. Similar to iron depletion, these mutants

abrogated locomotory response to H_2S (***Figure 6A–D***). Additionally, exposing animals to the

mitochondrial complex I inhibitor rotenone fully repressed locomotory response to H_2S

(***Figure 6E***). While rotenone exposure inhibits H_2S response, it evoked increased locomotory

activity over 2 hours (***Figure 6E***). These observations confirm the critical role of

mitochondria in supporting locomotory response in exposure to H_2S in *C. elegans*.

Discussion

Animals' behavior and physiology are profoundly influenced by the surrounding

environment, in which they reside. Wild *C. elegans* isolates thrive in the decomposing

matters, where the local concentrations of O_2 are low while the levels of CO_2 and H_2S could

be high. These animals have adapted their behavior in such an environment, displaying

increased sensitivity to high O_2 exposure but dampened responses to CO_2. In contrast, the

standard laboratory strain N2, which is constantly exposed to high O_2 but low CO_2, is

relatively insensitive to elevated levels of O_2 but robustly avoids CO_2 exposure (Beets et al.,

2020; Bretscher et al., 2008; Carrillo et al., 2013; Hallem & Sternberg, 2008). As observed in

response to CO_2, N2 animals exhibit strong reorientation and acceleration to escape the H_2S

exposure, while wild isolates display a significantly diminished response to H_2S.

The responses to CO_2 and H_2S in N2 animals are modulated by common signaling

pathways, which include the ciliogenesis, the cGMP-gated channels TAX-2 and TAX-4,

calcineurin subunit TAX-6/CNB-1, and nuclear hormone receptor NHR-49. Additionally,

these two responses are affected by the nutritional state of the animals. In *C. elegans*,

starvation is signaled by reduced activity in two major signaling pathways: the insulin and

TGF-β pathways. H_2S and CO_2 responses are abolished in starved animals or mutants that

disrupt insulin or TGF-β signaling. Therefore, the modulation of behavioral responses to H_2S

and CO_2 by external and internal factors appears highly similar. However, CO_2 response is

mediated by the guanylate cyclase GCY-9 in BAG neurons (Hallem et al., 2011), which

appears to be dispensable for H_2S sensing. We observed a prominent role for the guanylate

cyclase DAF-11 expressed in the chemosensory neurons ASJ for H_2S responses. *daf-11*

signaling in ASJ also mediates chemosensory responses to volatile chemicals and to NO

(Birnby et al., 2000; Hao et al., 2018). However, DAF-11 controls the normal larval

development of *C. elegans* and may have pleiotropic indirect effects on sensory functions

(Murakami, Koga, & Ohshima, 2001). In absence of more direct evidence, its role as H_2S

sensor in ASJ remain uncertain.

Long-term exposure to H_2S can elicit adaptive responses including changes in gene

expression patterns to mitigate the toxic effects of H_2S. Within just one hour of H_2S exposure,

we observed the upregulation of genes associated with H_2S detoxification. This response

enables animals to detoxify H_2S and maintain cellular homeostasis under sulfidic conditions.

This adaptive strategy is accompanied by the reduced locomotory response to H_2S exposure.

The HIF-1 (hypoxia-inducible factor) signaling is a major pathway involved in the adaption

during prolonged H_2S exposure (Budde & Roth, 2011; Miller et al., 2011b). When the HIF-1

pathway is constitutively induced in the nervous system, the behavioral responses to H_2S are

eliminated. Similarly, reduced behavioral responses to H_2S are observed after 12 hours of

exposure to 1% O_2. It has been proposed that persulfides, polysulfides, and thiosulfate, which

are the by-products during sulfide detoxification, play a role in H_2S signaling (Mishanina,

Libiad, & Banerjee, 2015). However, in the case of locomotion responses to H_2S in *C.*

elegans, this is unlikely to be the mechanism for the initial response, as mutants lacking

detoxification enzymes still exhibit an acute and transient response to H_2S. Interestingly, the

sustained locomotory activity was abolished in these mutants, suggesting that either the by-

products of H_2S detoxification contribute to maintain the behavioral response or the lack of

detoxification enzyme results in rapid cellular toxicity, preventing sustained locomotory

activity in H_2S (Budde & Roth, 2011).

Interestingly, we provide evidence that labile iron pool is critical for H_2S responses.

Removal of labile iron pool by the chelator BP or *ftn-1* overexpression attenuated both acute

and sustained responses to H_2S, suggesting that intracellular iron is required for the sensation

of H_2S. While increasing labile iron levels sustains the locomotory activity in H_2S. In *C.*

elegans, labile iron homeostasis is predominantly regulated by the HIF-1 signaling, the

activation of which decreases the transcription of *ftn-1* and increases the expression of *smf-3*

(Ackerman & Gems, 2012; Romney et al., 2011). Consistent with this, we have observed a

downregulation of the ferritins FTN-1 and FTN-2, as well as an upregulation of the iron

transporter SMF-3 during exposure to H_2S. This suggests that H_2S may control the expression

of FTN-1 and SMF-3, and consequently the labile iron pool, through the HIF-1 pathway.

Although the exact mechanisms are not yet fully understood, several possibilities

shed light on the importance of labile iron for H_2S responses. Fe^{2+} acts as a critical cofactor

for the PHD/EGL-9 enzymes, which are involved in the hydroxylation-dependent

destabilization of HIF-1 (Myllyharju & Kivirikko, 1997). PHD inactivation by iron

sequestration by ferritin can increase HIF-1 signaling, suggesting a potential role of labile

iron as a negative feedback loop to limit HIF-1 activity (Siegert et al., 2015). In addition, iron

is necessary for the biosynthesis of iron-sulfur (Fe-S) clusters, critical cofactors involved in

redox reactions and mitochondrial respiration (Read, Bentley, Archer, & Dunham-Snary,

2021; Xu & Moller, 2011). Labile iron is beneficial for the biogenesis of Fe-S clusters despite

its oxidative properties under normal O_2 conditions (Liochev, 1996). Interestingly, hypoxia

has been observed to enhance Fe-S cluster biogenesis by increasing the availability of

bioactive iron (Ast et al., 2019). Therefore, low oxidative stress may alleviate the need to

maintain a low labile iron pool. In this context, the antioxidant properties of H_2S or the

interruption of mitochondrial respiration by H_2S might provide condition favorable to high

labile iron pool (Shefa, Kim, Jeong, & Jung, 2018; Sunda, Kieber, Kiene, & Huntsman,

2002).

The primary mechanism underlying the toxic effects of H_2S has been suggested to be

the direct inhibition of cytochrome c oxidase, a critical enzyme involved in mitochondrial

respiration (Cooper & Brown, 2008; Khan et al., 1990; Nicholls & Kim, 1982b). Intriguingly,

we have observed that genetic and pharmacological inhibition of mitochondrial respiration

prevents the locomotory response of *C. elegans* to H_2S. This effect may be mediated by the

induction of the HIF-1 pathway, which has been observed in mutants with impaired

mitochondrial respiration (Lee, Hwang, & Kenyon, 2010). Furthermore, acute inhibition of

mitochondrial respiration by rotenone also enhances locomotion while suppressing the

response to H_2S. Reducing mitochondrial respiration has been found to stimulate aversive

behavior in *C. elegans* (Melo & Ruvkun, 2012). Therefore, the acute inhibition of

mitochondrial respiration by H_2S could transiently promote aversive behavior in *C. elegans*,

while long-term inhibition of mitochondrial respiration may contribute to the induction of the

HIF-1 pathway and adaptation to H_2S.

In summary, this study unveils a novel behavior of *C. elegans* in their avoidance to

H_2S. The response to H_2S is intricately shaped by their environment, where it integrates with

other sensory signals, including food availability, CO_2 and O_2 levels. We emphasize the vital

role of H_2S detoxification system, particularly the involvement of mitochondria and labile

irons, in sustaining the locomotory activity in H_2S. In the future, it will be exciting to

elucidate the neuronal mechanisms driving H_2S avoidance and to explore how the changes of

labile iron pools modulate cellular and organismal behavior responses to noxious sensory

stimuli.

Materials and Methods

Strains

C. elegans were maintained using standard protocols (Brenner, 1974). Strains used in this

study are listed in ***Supplementary file 1*** and ***Supplementary file 3***.

Preparation of H_2S gas mixture

Different H_2S concentrations were created as previously described (Miller & Roth, 2007).

Briefly, the H_2S-containing gas mixture were prepared by diluting 5000 ppm H_2S in nitrogen

(N_2) with 7% O_2 balanced with N_2. The gas flow was tuned by Sierra Smart-Trak 100 mass

flow controllers. H_2S concentration in the mixture was measured using two H_2S detectors

(MSA ALTAIR 2X gas detector for H_2S and Clip Single Gas Detector, SDG, CROWCON).

The pre-defined gas mixtures of 7% O_2, 1% O_2 and 5% CO_2 with N_2 were purchased from Air

Liquide Gas AB. 5000 ppm H_2S stock in N_2 was obtained from Linde Gas AB. Gas mixtures

in all experiments were hydrated before use.

Molecular Biology

The Multisite Gateway system (Thermo Fisher Scientific, United States) was used to generate

expression vectors. Promoters, including *gcy-37* (2.7kb), *trx-1* (1 kb), *ftn-1*(2 kb), *daf-11*(3

kb), *odr-3*(2.7 kb), *gpa-11*(3 kb), *sra-6*(3 kb), *odr-1*(2.4 kb), *flp-21*(4.1kb), and *ocr-2*(2.4 kb)

were amplified from N2 genomic DNA and cloned into pDONR P4P1 using BP clonase. *gcy-*

35, *tax-4* and *pkc-1* cDNAs were amplified using the first strand cDNA library as the

template, while *daf-11* and *ftn-1* genomic sequences were amplified using genomic DNA as

the template. The entry clones of these genes pDONR 221 were generated using BP reaction.

To generate the gain of function mutation of *pkc-1* (A160E), Q5 Site-Directed Mutagenesis

Kit (NEB) was used according to manufacturer instructions. All expression vectors were

generated using LR reaction. The primer sequences were displayed in ***Supplementary file 4***.

To generate transgenic animals, the *daf-11* expression vectors were injected at the

concentration of 20 ng/μl supplemented with 50 ng/μl of a coelomocyte co-injection marker

(*unc-122p::GFP)* and 30 ng/μl of 1kb DNA ladder. For *tax-4* gene, the injection mixtures

were prepared using 40 ng/μl of *tax-4* expression vectors, 50 ng/μl of a coelomocyte co-

injection marker and 10 ng/μl of 1kb DNA ladder. The rest of expression constructs were

injected at 50 ng/μl together with 50 ng/μl of a coelomocyte marker.

CRISPR/Cas9 genome editing

Genes were disrupted using CRISPR/Cas-9 mediated genome editing as described (Dokshin,

Ghanta, Piscopo, & Mello, 2018; Ghanta & Mello, 2020). The strategy involved the

utilization of homology-directed insertion of custom designed single strand DNA template

(ssODN), which had two homology arms of 35bp flanking the targeted PAM site. Between

two homology arms, a short sequence containing a unique restriction enzyme cutting site and

in frame as well as out of frame stop codons was included. The insertion of ssODN template

would delete 16 bases of coding sequence, ensuring the proper gene disruption. To prepare

the injection cocktail, 0.5 μl of Cas-9 protein (IDT) was mixed with 5 μl of 0.4μg/μl

tracrRNA (IDT, United States) and 2.8 μl of 0.4 μg/μl crRNA (IDT). The mixture was

incubated at 37°C for at least 10 minutes before 2.2 μl of 1μg/μl ssODN (or 500 ng dsDNA)

and 2 μl of 0.6 μg/μl *rol-6* co-injection marker were added. Nuclease-free water was used to

bring the final volume to 20 μl. The injection mixture was centrifuged for 2 minutes before

use.

Behavioral assays

H_2S evoked locomotion activity was monitored as described previously (Laurent et al., 2015;

Zhao et al., 2022). Briefly, OP50 bacteria were seeded on the assay plates 16 hours before

use. The border of bacterial lawn was removed using a PDMS stamp. For each assay, 25 to 30

day-one adult animals were picked onto assay plates, allowed to settle down for 15 minutes,

and subsequently sealed within microfluidic chambers. A syringe pump (PHD 2000, Harvard

Apparatus) was employed to deliver gas mixtures into the microfluidic chamber at a constant

flow of 3 ml/min. The rapid switch between different gas mixtures were controlled by Telfon

valves coupled with ValveBank Perfusion Controller (AutoMate Scientific). The locomotory

activity at different gas mixtures were monitored using a high-resolution camera (FLIR)

mounted on a Zeiss Stemi 508 microscope. Videos were captured at a rate of 2 frames per

second, starting with a 2-minute recording in 7% O_2, followed by 11 minutes in H_2S, and

ending with a final 5-minute period in 7% O_2. For the long-term recording in H_2S, video

capture commenced with a 2-minute period in 7% O_2, followed by a duration of 148 minutes

in 150 ppm H_2S, and concluded with a final 2-minute interval in 7% O_2.

To assess the effects of rotenone on H_2S evoked locomotory response, day-one adult

animals were subjected to 10 μM rotenone for 10min, 30min, 1 hour and 2 hours on the drug

containing plates. Subsequently, the rotenone-treated animals were assayed on the standard

assay plates without the drug. To explore the impact of ferric ammonium citrate (FAC)

(F5879, Sigma-Aldrich) and 2,2′-Bipyridyl (BP) (D216305, Sigma-Aldrich) on behavioral

response to H_2S, L4 animals were exposed to 100 μM FAC or 5mg/ml BP for 16 hours in the

presence of bacterial food. The FAC or BP treated animals were assayed on FAC or BP

containing plates.

Optogenetic experiments were performed as previously outlined (Zhao et al., 2022).

Transgenic L4 animals were exposed to 100μM all-trans retinal (ATR) (R2500, Sigma) in the

dark for 16 hours. The ATR-fed animals were assayed with the illumination of 70 μW/mm^2

blue light during the whole assay period. Blue light was emitted from an ultra-high-power

LED lamp (UHP-MIC-LED-460, Prizmatix), and the light intensity was determined using a

PM50 Optical Power Meter (ThorLabs). To minimize the effect of transmitted light on the

transgenic animals, a long-pass optical filter was used to eliminate the lights with short

wavelengths during picking. All videos of behavioral analysis were analyzed using a home-

made MATLAB program Zentracker (https://github.com/wormtracker/zentracker). For all

behavior analyses, at least three assays were performed for each strain with more than 80

worms in total.

RNA extraction and sequencing

To obtain RNA samples for RNA-seq, thirty young adult animals were picked to each fresh

plate and allowed to lay eggs for 2 hours, after which the adult animals were removed and

eggs were allowed to develop into day one adults. Day one animals were subsequently

challenged with H_2S for three different time points (1 hour, 2 hours and 12 hours). For each

time point, five plates of day one animals were exposed to 50 ppm H_2S, 150 ppm H_2S or 7%

O_2 as a control group. Subsequently, the animals were immediately collected and rinsed three

times with M9 buffer. After washing, the worm pellet was frozen in liquid nitrogen. Animals

were homogenized using Bullet Blender (Next Advance) in the presence of Qiazol Lysis

Reagent and 0.5mm Zirconia beads at 4°C. RNA was prepared using RNeasy Plus Universal

Mini Kit (Qiagen). Six independent RNA samples were prepared for all different treatments.

The 2100 Bioanalyzer instrument (Agilent) was used for the RNA quality control with a

microfluidic chip specific for RNA (Agilent RNA 6000 Nano Kit). Then 1 μg of qualified

RNA samples in RNase-free ddH2O was used for library preparation. Library was

constructed and sequenced by Novogene.

RNA-seq analysis

The RNA-seq data was aligned using STAR v2.7.9a (Dobin et al., 2013) and gene expression

counts extracted by featureCounts v2.0.3 (Liao, Smyth, & Shi, 2014). Differential expression

was calculated using DESeq2 (Love, Huber, & Anders, 2014). GO analysis was performed

using the EnrichR R package (Kuleshov et al., 2016) on genes having p.adj < 1e-10. For

brevity, several highly similar GO categories were manually omitted (full EnrichR output

available at Github repository).

Data availability

Raw sequencing data has been deposited at ArrayExpress # E-MTAB-13296. The R code is

available at https://github.com/henriksson-lab/ce_h2s_tc.

Acknowledgments

We thank the Caenorhabditis Genetics Center (funded by NIH Office of Research

Infrastructure Programs P40 OD010440) and the National BioResources Project Japan for

strains.

Funding

The computations were enabled by resources provided by the National Academic

Infrastructure for Supercomputing in Sweden (NAISS) and the Swedish National

Infrastructure for Computing (SNIC) at UPPMAX partially funded by the Swedish Research

Council through grant agreements no. 2022-06725 and no. 2018-05973. This work is

supported by the Swedish VR Research Council grant MIMS (2021-06602) to J.H., the

Belgian National Fund for Scientific Research (FRS-FNRS) to P.L., the ERC starting grant

(802653 OXYGEN SENSING), the Swedish Research Council VR starting grant (2018-

02216), and the Wallenberg Centre for Molecular Medicine (Umeå) to C.C.

References

Abel, D. C., Koenig, C. C., & Davis, W. P. (1987). Emersion in the Mangrove Forest Fish

Rivulus-Marmoratus - a Unique Response to Hydrogen-Sulfide. *Environmental*

Biology of Fishes, 18(1), 67-72. doi:Doi 10.1007/Bf00002329

Ackerman, D., & Gems, D. (2012). Insulin/IGF-1 and Hypoxia Signaling Act in Concert to

Regulate Iron Homeostasis in *Caenorhabditis elegans. Plos Genetics, 8*(3).

doi:ARTN e1002498

10.1371/journal.pgen.1002498

Anderson, C. P., & Leibold, E. A. (2014). Mechanisms of iron metabolism in *Caenorhabditis*

elegans. *Frontiers in Pharmacology, 5*. doi:ARTN 113

10.3389/fphar.2014.00113

Ast, T., Meisel, J. D., Patra, S., Wang, H., Grange, R. M. H., Kim, S. H., . . . Mootha, V. K.

(2019). Hypoxia Rescues Frataxin Loss by Restoring Iron Sulfur Cluster Biogenesis.

Cell, 177(6), 1507-+. doi:10.1016/j.cell.2019.03.045

Beets, I., Zhang, G., Fenk, L. A., Chen, C., Nelson, G. M., Felix, M. A., & de Bono, M.

(2020). Natural Variation in a Dendritic Scaffold Protein Remodels Experience-

Dependent Plasticity by Altering Neuropeptide Expression. *Neuron, 105*(1), 106-121

e110. doi:10.1016/j.neuron.2019.10.001

Birnby, D. A., Link, E. M., Vowels, J. J., Tian, H., Colacurcio, P. L., & Thomas, J. H. (2000).

A transmembrane guanylyl cyclase (DAF-11) and Hsp90 (DAF-21) regulate a

common set of chemosensory behaviors in *Caenorhabditis elegans*. *Genetics, 155*(1),

85-104. Retrieved from <Go to ISI>://WOS:000086869200008

Brenner, S. (1974). The genetics of *Caenorhabditis elegans*. *Genetics, 77*(1), 71-94.

doi:10.1093/genetics/77.1.71

Bretscher, A. J., Busch, K. E., & de Bono, M. (2008). A carbon dioxide avoidance behavior is

integrated with responses to ambient oxygen and food in *Caenorhabditis elegans*.

Proc Natl Acad Sci U S A, 105(23), 8044-8049. doi:10.1073/pnas.0707607105

Budde, M. W., & Roth, M. B. (2010). Hydrogen sulfide increases hypoxia-inducible factor-1

activity independently of von Hippel-Lindau tumor suppressor-1 in *C. elegans*. *Mol*

Biol Cell, 21(1), 212-217. doi:10.1091/mbc.E09-03-0199

Budde, M. W., & Roth, M. B. (2011). The response of *Caenorhabditis elegans* to hydrogen

sulfide and hydrogen cyanide. *Genetics, 189*(2), 521-532.

doi:10.1534/genetics.111.129841

Busch, K. E., Laurent, P., Soltesz, Z., Murphy, R. J., Faivre, O., Hedwig, B., . . . de Bono, M.

(2012). Tonic signaling from O(2) sensors sets neural circuit activity and behavioral

state. *Nat Neurosci, 15*(4), 581-591. doi:10.1038/nn.3061

Carrillo, M. A., Guillermin, M. L., Rengarajan, S., Okubo, R. P., & Hallem, E. A. (2013). O-

2-Sensing Neurons Control CO2 Response in *C. elegans*. *Journal of Neuroscience,*

33(23), 9675-9683. doi:10.1523/Jneurosci.4541-12.2013

Cha'on, U., Valmas, N., Collins, P. J., Reilly, P. E., Hammock, B. D., & Ebert, P. R. (2007).

Disruption of iron homeostasis increases phosphine toxicity in *Caenorhabditis*

elegans. *Toxicol Sci, 96*(1), 194-201. doi:10.1093/toxsci/kfl187

Cheung, B. H., Cohen, M., Rogers, C., Albayram, O., & de Bono, M. (2005). Experience-

dependent modulation of *C. elegans* behavior by ambient oxygen. *Curr Biol, 15*(10),

905-917. doi:10.1016/j.cub.2005.04.017

Cooper, C. E., & Brown, G. C. (2008). The inhibition of mitochondrial cytochrome oxidase

by the gases carbon monoxide, nitric oxide, hydrogen cyanide and hydrogen sulfide:

chemical mechanism and physiological significance. *J Bioenerg Biomembr, 40*(5),

533-539. doi:10.1007/s10863-008-9166-6

Couto, A., Oda, S., Nikolaev, V. O., Soltesz, Z., & de Bono, M. (2013). In vivo genetic

dissection of O2-evoked cGMP dynamics in a *Caenorhabditis elegans* gas sensor.

Proc Natl Acad Sci U S A, 110(35), E3301-3310. doi:10.1073/pnas.1217428110

Dallman, P. R. (1986). Biochemical Basis for the Manifestations of Iron-Deficiency. *Annual*

Review of Nutrition, 6, 13-40. doi:DOI 10.1146/annurev.nu.06.070186.000305

de Bono, M., & Bargmann, C. I. (1998). Natural variation in a neuropeptide Y receptor

homolog modifies social behavior and food response in *C. elegans*. *Cell, 94*(5), 679-

689. doi:10.1016/s0092-8674(00)81609-8

Dekker, L. V., Mcintyre, P., & Parker, P. J. (1993). Mutagenesis of the Regulatory Domain of

Rat Protein Kinase-C-Eta - a Molecular-Basis for Restricted Histone Kinase-Activity.

Journal of Biological Chemistry, 268(26), 19498-19504. Retrieved from <Go to

ISI>://WOS:A1993LW81900055

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., . . . Gingeras, T. R.

(2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics, 29*(1), 15-21.

doi:10.1093/bioinformatics/bts635

Dokshin, G. A., Ghanta, K. S., Piscopo, K. M., & Mello, C. C. (2018). Robust Genome

Editing with Short Single-Stranded and Long, Partially Single-Stranded DNA Donors

in *Caenorhabditis elegans*. *Genetics, 210*(3), 781-787.

doi:10.1534/genetics.118.301532

Fawcett, E. M., Hoyt, J. M., Johnson, J. K., & Miller, D. L. (2015). Hypoxia disrupts

proteostasis in *Caenorhabditis elegans*. *Aging Cell, 14*(1), 92-101.

doi:10.1111/acel.12301

Ferkey, D. M., Sengupta, P., & L'Etoile, N. D. (2021). Chemosensory signal transduction in

Caenorhabditis elegans. *Genetics, 217*(3). doi:10.1093/genetics/iyab004

Ghanta, K. S., & Mello, C. C. (2020). Melting dsDNA Donor Molecules Greatly Improves

Precision Genome Editing in *Caenorhabditis elegans*. *Genetics, 216*(3), 643-650.

doi:10.1534/genetics.120.303564

Gourley, B. L., Parker, S. B., Jones, B. J., Zumbrennen, K. B., & Leibold, E. A. (2003).

Cytosolic aconitase and ferritin are regulated by iron in *Caenorhabditis elegans*. *J*

Biol Chem, 278(5), 3227-3234. doi:10.1074/jbc.M210333200

Gray, J. M., Karow, D. S., Lu, H., Chang, A. J., Chang, J. S., Ellis, R. E., . . . Bargmann, C. I.

(2004). Oxygen sensation and social feeding mediated by a *C. elegans* guanylate

cyclase homologue. *Nature, 430*(6997), 317-322. doi:10.1038/nature02714

Hallem, E. A., Spencer, W. C., McWhirter, R. D., Zeller, G., Henz, S. R., Ratsch, G., . . .

Ringstad, N. (2011). Receptor-type guanylate cyclase is required for carbon dioxide

sensation by *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A, 108*(1), 254-259.

doi:10.1073/pnas.1017354108

Hallem, E. A., & Sternberg, P. W. (2008). Acute carbon dioxide avoidance in *Caenorhabditis*

elegans. *Proc Natl Acad Sci U S A, 105*(23), 8038-8043.

doi:10.1073/pnas.0707469105

Hao, Y., Yang, W., Ren, J., Hall, Q., Zhang, Y., & Kaplan, J. M. (2018). Thioredoxin shapes

the *C. elegans* sensory response to Pseudomonas produced nitric oxide. *Elife, 7*.

doi:10.7554/eLife.36833

Heiman, M. G., & Shaham, S. (2009). DEX-1 and DYF-7 establish sensory dendrite length

by anchoring dendritic tips during cell migration. *Cell, 137*(2), 344-355.

doi:10.1016/j.cell.2009.01.057

Hildebrandt, T. M., & Grieshaber, M. K. (2008). Three enzymatic activities catalyze the

oxidation of sulfide to thiosulfate in mammalian and invertebrate mitochondria.

FEBS J, 275(13), 3352-3361. doi:10.1111/j.1742-4658.2008.06482.x

Hiroki, S., Yoshitane, H., Mitsui, H., Sato, H., Umatani, C., Kanda, S., . . . Iino, Y. (2022).

Molecular encoding and synaptic decoding of context during salt chemotaxis in *C.*

elegans. *Nature Communications, 13*(1). doi:ARTN 2928

10.1038/s41467-022-30279-7

Horsman, J. W., Heinis, F. I., & Miller, D. L. (2019). A Novel Mechanism To Prevent H2S

Toxicity in *Caenorhabditis elegans*. *Genetics, 213*(2), 481-490.

doi:10.1534/genetics.119.302326

Jackson, M. R., Melideo, S. L., & Jorns, M. S. (2012). Human sulfide:quinone oxidoreductase

catalyzes the first step in hydrogen sulfide metabolism and produces a sulfane sulfur

metabolite. *Biochemistry, 51*(34), 6804-6815. doi:10.1021/bi300778t

Kaelin, W. G., Jr., & Ratcliffe, P. J. (2008). Oxygen sensing by metazoans: the central role of

the HIF hydroxylase pathway. *Mol Cell, 30*(4), 393-402.

doi:10.1016/j.molcel.2008.04.009

Kelley, J. L., Arias-Rodriguez, L., Patacsil Martin, D., Yee, M. C., Bustamante, C. D., &

Tobler, M. (2016). Mechanisms Underlying Adaptation to Life in Hydrogen Sulfide-

Rich Environments. *Mol Biol Evol, 33*(6), 1419-1434. doi:10.1093/molbev/msw020

Khan, A. A., Schuler, M. M., Prior, M. G., Yong, S., Coppock, R. W., Florence, L. Z., &

Lillie, L. E. (1990). Effects of hydrogen sulfide exposure on lung mitochondrial

respiratory chain enzymes in rats. *Toxicol Appl Pharmacol, 103*(3), 482-490.

doi:10.1016/0041-008x(90)90321-k

Kim, Y. I., Cho, J. H., Yoo, O. J., & Ahnn, J. (2004). Transcriptional regulation and life-span

modulation of cytosolic aconitase and ferritin genes in *C.elegans*. *J Mol Biol, 342*(2),

421-433. doi:10.1016/j.jmb.2004.07.036

Kuleshov, M. V., Jones, M. R., Rouillard, A. D., Fernandez, N. F., Duan, Q. N., Wang, Z. C.,

. . . Ma'ayan, A. (2016). Enrichr: a comprehensive gene set enrichment analysis web

server 2016 update. *Nucleic Acids Research, 44*(W1), W90-W97.

doi:10.1093/nar/gkw377

Laurent, P., Soltesz, Z., Nelson, G. M., Chen, C., Arellano-Carbajal, F., Levy, E., & de Bono,

M. (2015). Decoding a neural circuit controlling global animal state in *C. elegans*.

Elife, 4. doi:10.7554/eLife.04241

Lee, S. J., Hwang, A. B., & Kenyon, C. (2010). Inhibition of Respiration Extends *C. elegans*

Life Span via Reactive Oxygen Species that Increase HIF-1 Activity. *Current*

Biology, 20(23), 2131-2136. doi:10.1016/j.cub.2010.10.057

Liao, Y., Smyth, G. K., & Shi, W. (2014). featureCounts: an efficient general purpose

program for assigning sequence reads to genomic features. *Bioinformatics, 30*(7),

923-930. doi:10.1093/bioinformatics/btt656

Liochev, S. I. (1996). The role of iron-sulfur clusters in in vivo hydroxyl radical production.

Free Radical Research, 25(5), 369-384. doi:Doi 10.3109/10715769609149059

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biology, 15*(12). doi:ARTN 550

10.1186/s13059-014-0550-8

Ma, D. K., Vozdek, R., Bhatla, N., & Horvitz, H. R. (2012). CYSL-1 interacts with the O2-

sensing hydroxylase EGL-9 to promote H2S-modulated hypoxia-induced behavioral

plasticity in *C. elegans*. *Neuron, 73*(5), 925-940. doi:10.1016/j.neuron.2011.12.037

Macosko, E. Z., Pokala, N., Feinberg, E. H., Chalasani, S. H., Butcher, R. A., Clardy, J., &

Bargmann, C. I. (2009). A hub-and-spoke circuit drives pheromone attraction and

social behaviour in *C. elegans*. *Nature, 458*(7242), 1171-1175.

doi:10.1038/nature07886

Masini, A., Salvioli, G., Cremonesi, P., Botti, B., Gallesi, D., & Ceccarelli, D. (1994). Dietary

Iron-Deficiency in the Rat .1. Abnormalities in Energy-Metabolism of the Hepatic

Tissue. *Biochimica Et Biophysica Acta-Bioenergetics, 1188*(1-2), 46-52. doi:Doi

10.1016/0005-2728(94)90020-5

Masini, A., Trenti, T., Caramazza, I., Predieri, G., Gallesi, D., & Ceccarelli, D. (1994).

Dietary Iron-Deficiency in the Rat .2. Recovery from Energy-Metabolism

Derangement of the Hepatic Tissue by Iron Therapy. *Biochimica Et Biophysica Acta-*

Bioenergetics, 1188(1-2), 53-57. doi:Doi 10.1016/0005-2728(94)90021-3

Mathew, N. D., Schlipalius, D. I., & Ebert, P. R. (2011). Sulfurous gases as biological

messengers and toxins: comparative genetics of their metabolism in model

organisms. *J Toxicol, 2011*, 394970. doi:10.1155/2011/394970

Melo, J. A., & Ruvkun, G. (2012). Inactivation of Conserved *C. elegans* Genes Engages

Pathogen- and Xenobiotic-Associated Defenses. *Cell, 149*(2), 452-466.

doi:10.1016/j.cell.2012.02.050

Miller, D. L., Budde, M. W., & Roth, M. B. (2011a). HIF-1 and SKN-1 Coordinate the

Transcriptional Response to Hydrogen Sulfide in *Caenorhabditis elegans*. *PLoS One,*

6(9). doi:ARTN e25476

10.1371/journal.pone.0025476

Miller, D. L., Budde, M. W., & Roth, M. B. (2011b). HIF-1 and SKN-1 coordinate the

transcriptional response to hydrogen sulfide in *Caenorhabditis elegans*. *PLoS One,*

6(9), e25476. doi:10.1371/journal.pone.0025476

Miller, D. L., & Roth, M. B. (2007). Hydrogen sulfide increases thermotolerance and lifespan

in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A, 104*(51), 20618-20622.

doi:10.1073/pnas.0710191104

Mishanina, A. V., Libiad, M., & Banerjee, R. (2015). Biogenesis of reactive sulfur species for

signaling by hydrogen sulfide oxidation pathways. *Nature Chemical Biology, 11*(7),

457-464. doi:10.1038/nchembio.1834

Murakami, M., Koga, M., & Ohshima, Y. (2001). DAF-7/TGF-beta expression required for

the normal larval development in C-elegans is controlled by a presumed guanylyl

cyclase DAF-11. *Mechanisms of Development, 109*(1), 27-35. doi:Doi

10.1016/S0925-4773(01)00507-X

Murayama, T., Toh, Y., Ohshima, Y., & Koga, M. (2005). The dyf-3 gene encodes a novel

protein required for sensory cilium formation in *Caenorhabditis elegans*. *J Mol Biol,*

346(3), 677-687. doi:10.1016/j.jmb.2004.12.005

Myllyharju, J., & Kivirikko, K. I. (1997). Characterization of the iron- and 2-oxoglutarate-

binding sites of human prolyl 4-hydroxylase. *Embo Journal, 16*(6), 1173-1180.

doi:DOI 10.1093/emboj/16.6.1173

Nicholls, P., & Kim, J. K. (1982a). Sulfide as an Inhibitor and Electron-Donor for the

Cytochrome-C Oxidase System. *Canadian Journal of Biochemistry, 60*(6), 613-623.

doi:DOI 10.1139/o82-076

Nicholls, P., & Kim, J. K. (1982b). Sulphide as an inhibitor and electron donor for the

cytochrome c oxidase system. *Can J Biochem, 60*(6), 613-623. doi:10.1139/o82-076

Niu, W., Lu, Z. J., Zhong, M., Sarov, M., Murray, J. I., Brdlik, C. M., . . . Reinke, V. (2011).

Diverse transcription factor binding features revealed by genome-wide ChIP-seq in

C. elegans. Genome Research, 21(2), 245-254. doi:10.1101/gr.114587.110

Olson, K. R., & Straub, K. D. (2016). The Role of Hydrogen Sulfide in Evolution and the

Evolution of Hydrogen Sulfide in Metabolism and Signaling. *Physiology (Bethesda),*

31(1), 60-72. doi:10.1152/physiol.00024.2015

Pekec, T., Lewandowski, J., Komur, A. A., Sobanska, D., Guo, Y., Switonska-Kurkowska,

K., . . . Ciosk, R. (2022). Ferritin-mediated iron detoxification promotes hypothermia

survival in *Caenorhabditis elegans* and murine neurons. *Nat Commun, 13*(1), 4883.

doi:10.1038/s41467-022-32500-z

Persson, A., Gross, E., Laurent, P., Busch, K. E., Bretes, H., & de Bono, M. (2009). Natural

variation in a neural globin tunes oxygen sensing in wild *Caenorhabditis elegans*.

Nature, 458(7241), 1030-1033. doi:10.1038/nature07820

Pfenninger, M., Lerp, H., Tobler, M., Passow, C., Kelley, J. L., Funke, E., . . . Plath, M.

(2014). Parallel evolution of cox genes in H2S-tolerant fish as key adaptation to a

toxic environment. *Nat Commun, 5*, 3873. doi:10.1038/ncomms4873

Philipp, T. M., Gong, W., Kohnlein, K., Ohse, V. A., Muller, F. I., Priebs, J., . . . Klotz, L. O.

(2022). SEMO-1, a novel methanethiol oxidase in *Caenorhabditis elegans*, is a pro-

aging factor conferring selective stress resistance. *Biofactors, 48*(3), 699-706.

doi:10.1002/biof.1836

Qabazard, B., Li, L., Gruber, J., Peh, M. T., Ng, L. F., Kumar, S. D., . . . Moore, P. K. (2014).

Hydrogen sulfide is an endogenous regulator of aging in *Caenorhabditis elegans*.

Antioxid Redox Signal, 20(16), 2621-2630. doi:10.1089/ars.2013.5448

Rajan, M., Anderson, C. P., Rindler, P. M., Romney, S. J., Ferreira Dos Santos, M. C., Gertz,

J., & Leibold, E. A. (2019). NHR-14 loss of function couples intestinal iron uptake

with innate immunity in *C. elegans* through PQM-1 signaling. *Elife, 8*.

doi:10.7554/eLife.44674

Read, A. D., Bentley, R. E. T., Archer, S. L., & Dunham-Snary, K. J. (2021). Mitochondrial

iron-sulfur clusters: Structure, function, and an emerging role in vascular biology.

Redox Biology, 47. doi:ARTN 102164

10.1016/j.redox.2021.102164

Rodriguez-Kabana, R., Jordan, J. W., & Hollis, J. P. (1965). Nematodes: Biological Control

in Rice Fields: Role of Hydrogen Sulfide. *Science, 148*(3669), 524-526.

doi:10.1126/science.148.3669.524

Romney, S. J., Newman, B. S., Thacker, C., & Leibold, E. A. (2011). HIF-1 regulates iron

homeostasis in *Caenorhabditis elegans* by activation and inhibition of genes involved

in iron uptake and storage. *PLoS Genet, 7*(12), e1002394.

doi:10.1371/journal.pgen.1002394

Salvanes, A. G. V., Utne-Palm, A. C., Currie, B., & Braithwaite, V. A. (2011). Behavioural

and physiological adaptations of the bearded goby, a key fish species of the extreme

environment of the northern Benguela upwelling. *Marine Ecology Progress Series,*

425, 193-202. doi:10.3354/meps08998

Semenza, G. L. (2010). Defining the role of hypoxia-inducible factor 1 in cancer biology and

therapeutics. *Oncogene, 29*(5), 625-634. doi:10.1038/onc.2009.441

Shefa, U., Kim, M. S., Jeong, N. Y., & Jung, J. (2018). Antioxidant and Cell-Signaling

Functions of Hydrogen Sulfide in the Central Nervous System. *Oxidative Medicine*

and Cellular Longevity, 2018. doi:Artn 1873962

10.1155/2018/1873962

Siegert, I., Schodel, J., Nairz, M., Schatz, V., Dettmer, K., Dick, C., . . . Jantsch, J. (2015).

Ferritin-Mediated Iron Sequestration Stabilizes Hypoxia-Inducible Factor-1 alpha

upon LPS Activation in the Presence of Ample Oxygen. *Cell Reports, 13*(10), 2048-

2055. doi:10.1016/j.celrep.2015.11.005

Starich, T. A., Herman, R. K., Kari, C. K., Yeh, W. H., Schackwitz, W. S., Schuyler, M. W., .

. . Riddle, D. L. (1995). Mutations affecting the chemosensory neurons of

Caenorhabditis elegans. *Genetics, 139*(1), 171-188. doi:10.1093/genetics/139.1.171

Sunda, W., Kieber, D. J., Kiene, R. P., & Huntsman, S. (2002). An antioxidant function for

DMSP and DMS in marine algae. *Nature, 418*(6895), 317-320. doi:DOI

10.1038/nature00851

Swoboda, P., Adler, H. T., & Thomas, J. H. (2000). The RFX-type transcription factor DAF-

19 regulates sensory neuron cilium formation in *C. elegans*. *Mol Cell, 5*(3), 411-421.

doi:10.1016/s1097-2765(00)80436-0

Vora, M., Pyonteck, S. M., Popovitchenko, T., Matlack, T. L., Prashar, A., Kane, N. S., . . .

Rongo, C. (2022). The hypoxia response pathway promotes PEP carboxykinase and

gluconeogenesis in *C. elegans*. *Nature Communications, 13*(1). doi:ARTN 6168

10.1038/s41467-022-33849-x

Walter, P. B., Knutson, M. D., Paler-Martinez, A., Lee, S., Xu, Y., Viteri, F. E., & Ames, B.

N. (2002). Iron deficiency and iron excess damage mitochondria and mitochondrial

DNA in rats. *Proceedings of the National Academy of Sciences of the United States of*

America, 99(4), 2264-2269. doi:DOI 10.1073/pnas.261708798

Xu, X. M., & Moller, S. G. (2011). Iron-Sulfur Clusters: Biogenesis, Molecular Mechanisms,

and Their Functional Significance. *Antioxidants & Redox Signaling, 15*(1), 271-307.

doi:10.1089/ars.2010.3259

Zhao, L., Fenk, L. A., Nilsson, L., Amin-Wetzel, N. P., Ramirez-Suarez, N. J., de Bono, M.,

& Chen, C. (2022). ROS and cGMP signaling modulate persistent escape from

hypoxia in *Caenorhabditis elegans. PLoS Biol, 20*(6), e3001684.

doi:10.1371/journal.pbio.3001684

Zimmer, M., Gray, J. M., Pokala, N., Chang, A. J., Karow, D. S., Marletta, M. A., . . .

Bargmann, C. I. (2009). Neurons detect increases and decreases in oxygen levels

using distinct guanylate cyclases. *Neuron, 61*(6), 865-879.

doi:10.1016/j.neuron.2009.02.013

Figure legends

Figure 1. cGMP signaling in ASJ neurons contributes to H_2S avoidance

(**A–H**) Locomotory responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2

for animals of the indicated genotype: WT (N2) and *daf-19(m86)* (**A**); WT and *dyf-3(m185)*

(**B**); WT and *dyf-7(m539)* (**C**); WT, *daf-11(m47)* and *tax-4(p678)* (**D**); WT and *tax-2(p691)*

(**E**); WT, *daf-2(e1370), daf-16(mgDf47)* and *daf-2(e1370); daf-16(mgDf47)* double mutants

(**F**); WT, *daf-7(e1372), daf-3(mgDf90)* and *daf-7(e1372); daf-3(mgDf90)* double mutants

(**G**); WT and *gcy-9(tm7632)* (**H**). In this and subsequent figures for H_2S assay, red bars on the

x-axis represent two intervals (4–5 minutes and 11–12 minutes, labeled as a and b

respectively in Figure 1D, F, G, J and K) used for statistical analysis unless otherwise

indicated. **** = $p < 0.0001$, *** = $p < 0.001$, ** = $p < 0.01$, * = $p < 0.05$, ns = not

significant, Mann–Whitney U test. (**I**) Comparison between the acute locomotory response of

WT animals to 150 ppm H_2S balanced with 7% O_2 and the acute locomotory response to 5%

CO_2 balanced with 7% O_2. Red bars on the x-axis indicate two intervals (3-4 minutes and 11-

12 minutes) used for statistical analysis. * = $p < 0.05$, ns = not significant, Mann–Whitney U

test. (**J**) Locomotory responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2

for animals of the indicated genotype: WT, *daf-11(m47)* and transgenic *daf-11(m47)*

expressing *daf-11* genomic DNA in ASJ neurons (two independent lines). *** = $p < 0.001$, **

= $p < 0.01$, * = $p < 0.05$, Mann–Whitney U test. (**K**) Locomotory responses to a switch from

7% O_2 to 150 ppm H_2S balanced with 7% O_2 for animals of the indicated genotype: WT, *tax-*

4(p678) and transgenic *tax-4(p678)* expressing *tax-4* cDNA in ASJ neurons (two independent

lines). *** = $p < 0.001$, ** = $p < 0.01$, * = $p < 0.05$, Mann–Whitney U test.

Figure 2. Activation of O_2 sensing circuit antagonizes H_2S avoidance

(**A**) Locomotory responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2 for

WT animals, wild isolates CB4858 and CB4855. Red bars on the x-axis represent two

intervals (4–5 minutes and 11–12 minutes, labeled as a and b respectively), used for statistical

analyses. ** = $p < 0.01$, * = $p < 0.05$, Mann–Whitney U test. (**B, C**) Locomotory responses to

a switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2 for animals of the indicated

genotype: WT and *npr-1(ad609)* (**B**); WT, *npr-1(ad609) and* transgenic *npr-1(ad609)*

expressing *npr-1* in RMG neurons using Cre-LoxP system (Macosko et al., 2009) (**C**). Red

bars on the x-axis indicate two intervals (4-5 minutes and 11-12 minutes) used for statistical

analysis. **** = $p < 0.0001$, *** = $p < 0.001$, ** = $p < 0.01$, Mann–Whitney U test. (**D–G**)

Locomotory responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2 for

animals of the indicated genotype: WT, *npr-1(ad609), gcy-35(ok769)* and *npr-1(ad609); gcy-*

35(ok769) double mutants (**D**); WT, *npr-1(ad609), tax-4(p678)* and *npr-1(ad609); tax-*

4(p678) double mutants (**E**); WT, *npr-1(ad609), qaIs2241*(genetic ablation of AQR, PQR and

URX neurons) and *npr-1(ad609); qaIs 2241* (**F**); WT, *npr-1(ad609), npr-1(ad609); gcy-*

35(ok769) double mutants and *transgenic npr-1(ad609); gcy-35(ok769)* double mutants

expressing *gcy-35* cDNA under *gcy-37* promoter, which drives expression in O_2 sensing

neurons (**G**). Red bars on the x-axis indicate two intervals (4-5 minutes and 11-12 minutes)

used for statistical analysis. *** = $p < 0.001$, ** = $p < 0.01$, * = $p < 0.05$, ns = not significant,

Mann–Whitney U test.

Figure 3. Prolonged H₂S exposure reprograms gene expression in *C. elegans*

(**A**) Venn diagram displaying the number of differentially expressed genes after 1-hour

exposure to 50 ppm or 150 ppm H_2S balanced with 7% O_2 in WT animals. $p<1e-10$. (**B**) Venn

diagram displaying the number of differentially expressed genes after 1-hour, 2-hour and 12-

hour exposure to 150 ppm H_2S balanced with 7% O_2 in WT animals. $p<1e-10$. (**C, D**)

Significantly enriched GO categories for differentially expressed genes with adjusted p value

$<1e-10$ in WT animals exposed to 50 ppm (**C**) or 150 ppm (**D**) H_2S balanced with 7% O_2 for 1

hour. (**E, F**) Volcano plots showing the differentially expressed genes with adjusted p value

$<1e-10$ in WT animals exposed to 50 ppm (**E**) or 150 ppm (**F**) H_2S balanced with 7% O_2 for 1

hour. A set of genes involved in H_2S detoxification, cysteine metabolism and stress response

were highlighted in red.

Figure 4. Acute response to H₂S is modulated by HIF-1 signaling

(**A–D**) Locomotory responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2

for animals of the indicated genotype: WT and *hif-1(ia4)* (**A**); WT, *egl-9(sa307) and vhl-*

1(ok161) (**B**); WT, *egl-9(sa307), hif-1(ia4) and egl-9(sa307); hif-1(ia4)* double mutants (**C**);

WT, *vhl-1(ok161), hif-1(ia4) and vhl-1(ok161); hif-1(ia4)* double mutants (**D**). Red bars on

the x-axis represent two intervals (4–5 minutes and 11–12 minutes, labeled as a and b

respectively), used for statistical analyses. *** = $p < 0.001$, ** = $p < 0.01$, * = $p < 0.05$, ns =

not significant, Mann–Whitney U test. (**E**) Locomotory responses of WT animals to a switch

from 7% O_2 to 150 ppm H_2S balanced with 7% O_2 after 0, 12, 24 or 36 hours in 1% O_2. Red

bars on the x-axis represent two intervals (4–5 minutes and 11–12 minutes, labeled as a and b

respectively), used for statistical analyses. *** = $p < 0.001$, ** = $p < 0.01$, ns = not

significant, Mann–Whitney U test. (**F**) Locomotory responses to a switch from 7% O_2 to 150

ppm H_2S balanced with 7% O_2 for animals carrying the non-degradable form of HIF-1. Non-

degradable form of HIF-1 (P621A and P621G) are expressed from a pan-neuronal promoter

unc-14 or form *hif-1* endogenous promoter, respectively. Red bars on the x-axis represent two

intervals (4–5 minutes and 11–12 minutes, labeled as a and b respectively), used for statistical

analyses. ** = $p < 0.01$, ns = not significant, Mann–Whitney U test. (**G–J**) Locomotory

responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2 for animals of the

indicated genotype: WT and *sqrd-1(tm3378)* (**G**); WT and *ethe-1(yum2895)* (**H**); WT, *cysl-*

1(ok762), cysl-2(ok3516) and *cysl-3(yum4)* (**I**); WT and *semo-1(yum2889)* (**J**). Red bars on

the x-axis represent two intervals (4–5 minutes and 11–12 minutes, labeled as a and b

respectively), used for statistical analyses. *** = $p < 0.001$, ** = $p < 0.01$, * = $p < 0.05$, ns =

not significant, Mann–Whitney U test.

Figure 5. Labile iron pool is critical to sustain the locomotory activity in H_2S

(**A, B**) Volcano plots showing the relative expression of the genes involved in the regulation

of iron homeostasis in WT animals after 2-hour exposure with 50 ppm (**A**) or 150 ppm (**B**)

H_2S balanced with 7% O_2. (**C**) Locomotory responses to a switch from 7% O_2 to 150 ppm

H_2S balanced with 7% O_2 for WT animals, WT animals pretreated with 100 μM 2,2′-

Bipyridyl (BP) and WT animals pretreated with 5 mg/ml ferric ammonium citrate (FAC) in

the presence of food for 16 hours. Red bars on the x-axis represent two intervals (4–5 minutes

and 14–15 minutes, labeled as a and b respectively), used for statistical analyses. ** = $p <$

0.01, * = $p < 0.05$, ns = not significant, Mann–Whitney U test. (**D**) Locomotory responses to a

switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2 for animals of the indicated

genotype: WT, *smf-3(ok1305)* and *ftn-1(ok3625)*. Red bars on the x-axis represent two

intervals (4–5 minutes and 14–15 minutes, labeled as a and b respectively), used for statistical

analysis. *** = $p < 0.001$, * = $p < 0.05$, ns = not significant, Mann–Whitney U test. (**E**)

Locomotory responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2 for WT

animals and animals overexpressing *ftn-1* genomic DNA under its own promoter (#1 and #2

indicate two independent lines). Red bars on the x-axis represent two intervals (4–5 minutes

and 11–12 minutes, labeled as a and b respectively), used for statistical analysis. **** = $p <$

0.0001, ** = $p < 0.01$, ns = not significant, Mann–Whitney U test. (**F, G**) Locomotory

responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2 for animals of the

indicated genotype: WT, *hif-1(ia4)* and *hif-1(ia4)* mutants pretreated with 5 mg/ml ferric

ammonium citrate (FAC) for 16 hours (**F**)*;* WT, *hif-1(ia4), ftn-1(ok3625)* and *hif-1(ia4); ftn-*

1(ok3625) double mutants (**G**). Red bars on the x-axis represent two intervals (4–5 minutes

and 8–9 minutes) used for statistical analysis. * = $p < 0.05$, ns = not significant, Mann–

Whitney U test.

Figure 6. Mitochondrial function is required for acute response to H_2S

(**A–D**) Locomotory responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2

for animals of the indicated genotype: WT and *gas-1(fc21)* (**A**); WT and *clk-1(qm30)* (**B**);

WT and *mev-1(kn1)* (**C**); WT and *isp-1(qm150)* (**D**). Red bars on the x-axis represent two

intervals (4–5 minutes and 11–12 minutes) used for statistical analysis. **** = $p < 0.0001$, **

= $p < 0.01$, * = $p < 0.05$, Mann–Whitney U test. (**E**) Locomotory responses to a switch from

7% O_2 to 150 ppm H_2S balanced with 7% O_2 for WT animals pretreated with 10 μM rotenone

for 0 minute, 10 minutes, 30 minutes, 60 minutes or 120 minutes. Red bars on the x-axis

represent two intervals (4–5 minutes and 11–12 minutes, labeled as a and b respectively),

used for statistical analysis. *** = $p < 0.001$, ** = $p < 0.01$, * = $p < 0.05$, ns = not significant,

Mann–Whitney U test.

Figure 1──figure supplement 1. Acute locomotory responses of *C. elegans* to hydrogen

sulfide.

(**A**) Changes in the locomotory activity of WT animals evoked by a switch from 7% O_2 to 150

ppm H_2S balanced with 7% O_2. (**B**) Reorientation movements (omega turns) of WT animals

evoked by a switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2. (**C**) Changes in

reversal frequency in WT animals experiencing a switch from 7% O_2 to 150 ppm H_2S

balanced with 7% O_2. (**D**) Locomotory activity of WT was recorded for 2 minutes at 7%

O_2 and 148 minutes at 150 ppm H_2S balanced with 7% O_2 followed by 2 additional minutes at

7% O_2. (**E**) Locomotory activity of WT animals evoked by acute hypoxia after preincubation

for 148 minutes at 150 ppm H_2S balanced with 7% O_2. Locomotory activity was recorded for

2 minutes at each gas interval. (**F**) Locomotory responses of WT animals to different

concentrations of H_2S balanced with 7% O_2. Red bars on the x-axis represent two intervals

(4–5 minutes and 11–12 minutes, labeled as a and b respectively) used for statistical analysis.

**** = $p < 0.0001$, ** = $p < 0.01$, * = $p < 0.05$, ns = not significant, Mann–Whitney U test.

Figure 1—figure supplement 2. Acute response to H_2S is regulated by multiple signaling

pathways

(**A–E**) Locomotory responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2

for animals of the indicated genotype: WT and *tax-6(ok2065)* (**A**); WT and *cnb-1(ok276)* (**B**);

WT and *nhr-49(nr2041)* (**C**); WT animals and WT animals starved for 24 hours (**D**); WT and

daf-2(e1370) (**E**). To assay *daf-2* animals, WT and *daf-2* mutants were kept in 15°C. L4

animals were picked, maintained at 25°C until day-one adults and assayed at room

temperature. Red bars on the x-axis represent two intervals (4–5 minutes and 11–12 minutes),

used for statistical analysis. **** = $p < 0.0001$, *** = $p < 0.001$, ** = $p < 0.01$, ns = not

significant, Mann–Whitney U test. (**F–I**) Locomotory responses to a switch from 7% O_2 to

150 ppm H_2S balanced with 7% O_2 for animals of the indicated genotype: WT, *daf-11(m47)*

and transgenic *daf-11* mutants expressing *daf-11* genomic DNA in different subsets of

neurons (**F–H**); The detailed expression pattern of each promoter is listed in (**I**). Red bars on

the x-axis represent two intervals (4–5 minutes and 11–12 minutes, labeled as a and b

respectively), used for statistical analysis. * = $p < 0.05$, ns = not significant, Mann–Whitney

U test.

Figure 2—figure supplement 1. NPR-1 signaling modulates locomotory response to H_2S

(**A**) Locomotory responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2 for

WT animals and wild isolate CB4856. Red bars on the x-axis represent two intervals (4–5

minutes and 11–12 minutes), used for statistical analysis. ** = $p < 0.01$, ns = not significant,

Mann–Whitney U test. (**B**) Optogenetic stimulation of RMG neurons using

channelrhodopsin-2 (ChR2). Expressing ChR2 in RMG neurons were achieved using Cre-

LoxP system (Macosko et al., 2009). L4 animals were picked, grown for 16 hours on the

seeded plates containing ATR, and assayed as day-one adults. Blue light was delivered

throughout the assay. Red bars on the x-axis represent two intervals (4–5 minutes and 11–12

minutes), used for statistical analysis. ** = $p < 0.01$, ns = not significant, Mann–Whitney U

test. (**C**) Activation of O_2 sensing neurons by expressing a gain-of-function allele of *pkc-1*

under *gcy-37* promoter. Red bars on the x-axis represent two intervals (4–5 minutes and 11–

12 minutes), used for statistical analysis. * = $p < 0.05$, ns = not significant, Mann–Whitney U

test.

Figure 3—figure supplement 1. Transcriptome reprogramming induced by H_2S

exposure

(**A**) Venn diagram displaying the number of differentially expressed genes after exposure in

50 ppm H_2S balanced with 7% O_2 in WT animals for 1 hour, 2 hours or 12 hours. Adjusted *p*

value <1e-10. (**B, C**) Significantly enriched GO categories for differentially expressed genes

with adjusted *p* value <1e-10 in WT animals exposed to 50 ppm (**B**) or 150 ppm (**C**) H_2S

balanced with 7% O_2 for 2 hours. (**D, E**) Volcano plot showing the differentially expressed

genes with adjusted *p* value <1e-10 in WT animals exposed to 50 ppm (**D**) or 150 ppm (**E**)

H_2S balanced with 7% O_2 for 2 hours. A set of genes involved in H_2S detoxification, cysteine

metabolism and stress response were highlighted in red. (**F, G**) The number of HIF-1(**F**) and

SKN-1(**G**) target genes induced by 50 ppm or 150 ppm H_2S exposure for 1hour, 2 hours or 12

hours in WT animals, with Adjusted *p* value < 1e-30. (**H–J**) Venn diagrams displaying the

number of differentially expressed genes (adjusted *p* value <1e-10) in WT animals of the

indicated conditions: animals were exposed to 50 ppm or 150 ppm H_2S for 1hour (**H**); for

2hours (**I**); and for 12 hours (**J**). (**K, L**) Volcano plots showing the differentially expressed

genes (adjusted p value <1e-10) in WT animals that were exposed to either 50 ppm (**K**) or

150 ppm (**L**) H_2S balanced with 7% O_2 for 12 hours. A set of genes involved in H_2S

detoxification, cysteine metabolism and stress response were highlighted in red.

Figure 5—figure supplement 1. *ftn-1* expression is consistently repressed in H_2S

(**A–D**) Volcano plots showing the relative expression of the genes involved in the regulation

of iron homeostasis (*ftn-1*, *ftn-2* and *smf-3*) in WT animals of the indicated conditions: 1-hour

exposure in 50 ppm H_2S balanced with 7% O_2 (**A**); 1-hour exposure in 150 ppm H_2S balanced

with 7% O_2 (**B**); 12-hour exposure in 50 ppm H_2S balanced with 7% O_2 (**C**); and 12-hour

exposure in 150 ppm H_2S balanced with 7% O_2 (**D**). (**E, F**) Locomotory responses to a switch

to 7% to 1% O_2 for animals of the indicated treatments or genotypes: WT animals, WT

animals treated with 100 μM 2,2′-Bipyridyl (BP), and WT animals treated with 5 mg/ml ferric

ammonium citrate (FAC) for 16 hours (**E**); WT and WT animals overexpressing *ftn-1*

genomic DNA under its own promoter (#1 and #2 indicate two independent lines) (**F**). Red

bar on the x-axis represents time interval (3–4 minutes) used for statistical analysis. * $= p <$

0.05, ns = not significant, Mann–Whitney U test.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 1-figure supplement 1

Figure 1-figure supplement 2

Promoter	neurons															
Pdaf-11	ASH	AWB	AWC	ASJ	ASK	ADL	ASI	ASE								
Podr-3	ASH	AWB	AWC						ADF	AWA						
Podr-1		AWB	AWC	ASJ	ASK		ASI									
Pflp-21	ASH					ADL	ASI	ASE			FLP	URA	RMG	MC	M2	M4
Pgpa-11	ASH					ADL										
Psra-6	ASH						ASI									PVQ

Figure 2-figure supplement 1

Figure 3-figure supplement 1

Figure 5-figure supplement 1