Avoidance of hydrogen sulfide is modulated by external and internal states in *C. elegans*

Longjun Pu[1, 2, 3], Lina Zhao[1, 2, 3], Jing Wang[1, 2, 3], Clementine Deleuze[6], Lars Nilsson[1, 2, 3], Johan

Henriksson[1, 4, 5], Patrick Laurent[6*], Changchun Chen[1, 2, 3*]

[1]Department of Molecular Biology, Umeå University, Umeå, Sweden

[2]Umeå Centre for Molecular Medicine, Umeå University, Umeå, Sweden

[3]Wallenberg Centre for Molecular Medicine, Umeå University, Umeå, Sweden

[4]The Laboratory for Molecular Infection Medicine Sweden (MIMS), Umeå University, Umeå,

Sweden

[5]Integrated Science Lab (Icelab), Umeå University, Umeå, Sweden

[6]Laboratory of Neurophysiology, ULB Neuroscience Institute (UNI), Université Libre de

Bruxelles (ULB), Belgium

*Corresponding author

E-mail: patrick.laurent.ulb@gmail.com; Changchun.chen@umu.se

Abstract

Hydrogen sulfide (H_2S) acts as an energy source, a toxin, and a gasotransmitter across diverse biological contexts. We use the robust locomotory responses of *Caenorhabditis elegans* to high levels of H_2S to elucidate the molecular mechanisms underlying its acute perception and adaptive responses. We find that the H_2S-evoked behavioral response is shaped by multiple environmental factors including oxygen (O_2) levels and nutritional state, and is modulated by various pathways such as insulin, TGF-β, and HIF-1 signaling, as well as by input from O_2-sensing neurons. Prolonged exposure to H_2S activates HIF-1 signaling, leading to the upregulation of stress-responsive genes, including those involved in H_2S detoxification. This promotes an adaptive state in which locomotory speed is reduced in H_2S, while responsiveness to other stimuli is preserved. In mutants deficient in HIF-1 signaling, iron storage, and detoxification mechanisms, animals display a robust initial response but rapidly enter a sleep-like behavior characterized by reduced mobility and diminished responsiveness to subsequent sensory stimuli. Furthermore, while acute production of mitochondria-derived reactive O_2 species (ROS) appears to initiate the avoidance response to H_2S, persistently high ROS promotes an adaptive state, likely by activating various stress-response pathways, without substantially compromising cellular H_2S detoxification capacity. Taken together, our study provides comprehensive molecular insights into the mechanisms through which *C. elegans* detects, modulates, and adapts its response to H_2S exposure.

Introduction

During the late Proterozoic era, the rise in oxygen (O_2) levels eliminated hydrogen sulfide (H_2S) from most habitats. Yet, low O_2 levels, along with high concentrations of hydrogen sulfide (H_2S) and carbon dioxide (CO_2), can persist in enclosed environments where bacteria actively decompose organic matter (Olson & Straub, 2016). H_2S can easily permeate biological membranes and interfere with various cellular processes. One of the most detrimental effects of H_2S is the disruption of cellular respiration by remodeling the mitochondrial electron transport chain and inhibiting cytochrome *c* oxidase (COX) (Cooper & Brown, 2008; Khan et al., 1990; Nicholls & Kim, 1982; Romanelli-Cedrez, Vairoletti, & Salinas, 2024).

The detection and response to chemical signals are crucial for various organisms to interact effectively with their environment. Animals living in H_2S-rich environments have evolved mechanisms to either avoid or adapt to these conditions, conferring them survival advantages. Species that encounter periodic increases in H_2S often possess sensing abilities and exhibit avoidance behaviors. For example, benthic species undertake daily vertical migrations between sulfidic and non-sulfidic waters (Abel, Koenig, & Davis, 1987; Salvanes, Utne-Palm, Currie, & Braithwaite, 2011). Certain metazoan species have seized the ecological opportunities presented by H_2S-rich environments through the development of cellular adaptations. Livebearing fish species, for instance, have evolved COX subunits that are more resistant to H_2S and more efficient at H_2S detoxification (Kelley et al., 2016; Pfenninger et al., 2014).

Free-living nematodes feed on bacteria that thrive in decaying organic matter, such as compost. This complex and dynamic environment, where species like *C. elegans* are commonly found, may exhibit significant fluctuations in H_2S levels over short distances (Adams, Farwell, Pack, & Bamesberger, 1979; Budde & Roth, 2011; Morra & Dick, 1991; Patange, Breen, Arsuffi, & Ruvkun, 2025; Rodriguez-Kabana, Jordan, & Hollis, 1965; Romanelli-Cedrez et al., 2024). Evidence suggests that *C. elegans* has evolved tolerance to prolonged H_2S exposure by reprogramming gene expression through H_2S-induced stabilization of the hypoxia-inducible factor HIF-1 (Budde & Roth, 2010; Powell-Coffman, 2010), and by developing an H_2S-

resistant electron transport chain (Romanelli-Cedrez et al., 2024). In particular, low

concentrations of H_2S are not only well tolerated but can also be beneficial. For instance,

exposure to 50 ppm H_2S has been shown to extend the lifespan and enhance thermotolerance

(Budde & Roth, 2010; Fawcett, Hoyt, Johnson, & Miller, 2015; Horsman, Heinis, & Miller,

2019; Miller & Roth, 2007; Qabazard et al., 2014). Moreover, endogenous H_2S production in

C. elegans modulates its interactions with actinobacteria (Patange et al., 2025). However,

exposure to concentrations exceeding 150 ppm proved lethal (Budde & Roth, 2010, 2011;

Fawcett et al., 2015; Horsman et al., 2019; Miller, Budde, & Roth, 2011a; Miller & Roth, 2007).

In addition to long-term adaptive mechanisms, *C. elegans* exhibits an acute behavioral

response to H_2S exposure (Budde & Roth, 2011). In this study, we investigated whether and

how the nematode *C. elegans* acutely avoids environments rich in H_2S, how it adapts at the

molecular and physiological levels to H_2S exposure, and how these adaptations alter the

avoidance response.

Results

***C. elegans* increases locomotory activity in response to acute H_2S exposure**

In response to aversive cues, *C. elegans* initiates a pirouette followed by an increase of its

locomotory speed to escape the noxious stimuli. When acutely exposed to 150 ppm H_2S, the

laboratory N2 strain exhibited an increased locomotory speed and enhanced turning behavior,

while generally decreasing the frequency of reversals, as a strategy to escape the noxious

stimuli (***Figure 1A–C, Figure 1—Video 1***). The maximum speed was reached after 6–8

minutes in H_2S and returned to the baseline over a period of approximately 1 hour (***Figure 1D***).

The reduced locomotory speed after 30 minutes in 150 ppm H_2S was reversible when these

animals were immediately challenged with another stimulus, such as near-UV light (***Figure***

1E), suggesting that neuromuscular function remains preserved. The magnitude and dynamics

of the H_2S-induced avoidance responses were concentration-dependent. Under our assay

conditions, 50 ppm H_2S did not elicit an acute speed increase, and responses to 75 ppm were

variable. In contrast, exposure to 150 ppm H_2S consistently triggered a robust and reproducible

escape response (***Figure 1F***), consistent with earlier reports that H_2S at 50 ppm exerts

beneficial effects, while H_2S at 150 ppm is toxic to *C. elegans* (Budde & Roth, 2010; Fawcett

et al., 2015; Horsman et al., 2019; Miller & Roth, 2007; Qabazard et al., 2014). To investigate

whether low levels of H_2S might be preferable or even attractive to the animals, we exposed

wild type animals to a linear gradient of H_2S (***Figure 1G***; See methods for details) (Bretscher,

Busch, & de Bono, 2008; Chang, Chronis, Karow, Marletta, & Bargmann, 2006; Gray et al.,

2004). Animals were allowed to move freely on a thin layer of bacteria within a microfluidic

chamber containing an H_2S gradient ranging from 150 ppm at one end to 0 ppm at the other.

As expected, *C. elegans* robustly avoids 150 ppm H_2S but does not accumulate at the extreme

end of 0 ppm *(Figure 1G, H).* The highest proportion of animals was found in the region with

approximately 40 ppm H_2S. These data suggest that H_2S acts as a potent repellent for *C. elegans*

at high concentrations but as an attractant at low levels.

Candidate gene screen identifies signaling pathways required for H_2S-evoked locomotion

To investigate the molecular mechanisms underlying the H_2S-evoked locomotory response in

C. elegans, we conducted a candidate gene screen for mutants that failed to increase their

locomotory speed upon exposure to an acute rise of H_2S level to 150 ppm. Our analysis

prioritized genes that are either known to be, or potentially involved in, sensory responses to

gaseous stimuli in both *C. elegans* and mammals (***Supplementary file 1***). We began by

examining mutants affecting globins, guanylate cyclases, and cyclic nucleotide-gated (CNG)

channels, which are involved in O_2 or CO_2 sensing in *C. elegans*, as well as wild isolates and

other previously characterized mutants with defective responses to gas stimuli (***Supplementary***

file 1) (Bretscher et al., 2008; Gray et al., 2004; Hallem & Sternberg, 2008; Persson et al., 2009).

Given the essential role of K^+ channels in acute hypoxia sensing in mammals, we also screened

all potassium (K^+) channel mutants (Gao, Gonzalez-Rodriguez, Ortega-Saenz, & Lopez-Barneo,

2017; Weir, Lopez-Barneo, Buckler, & Archer, 2005). To explore how the H_2S sensory signals

were transmitted within the neural circuit, we assayed mutants with defects in ciliogenesis or

cilia-mediated signaling, as well as animals deficient in neurotransmission involving classical

neurotransmitters, neuropeptides, or biogenic amines. Finally, considering that mitochondrial

function is central to H_2S detoxification, we included mitochondrial electron transport chain

(ETC) mutants and key factors implicated in H_2S and superoxide clearance. Together, these

functionally diverse categories allowed us to survey a wide spectrum of potential mechanisms

contributing to the behavioral response to H_2S.

In the screen, a collection of mutants displayed reduced speed response, whereas only

a few exhibited a lack of omega-turn and/or reversal responses to H_2S (***Supplementary file 1***).

The reversal inhibition upon H_2S exposure and its rebound following H_2S withdrawal appeared

to vary substantially across experiments for N2, suggesting that differences in population

density, food availability, and animals' physiological state prior to the assay may matter.

Nevertheless, our screen revealed that signaling pathways, including cGMP, insulin and TGF-

β signaling, contributed to H_2S-evoked avoidance (***Supplementary file 1***). A naturally

occurring variant of the neuropeptide receptor gene *npr-1*, previously implicated in the

modulation of O_2 and CO_2 sensing, also affected locomotory speed in H_2S (***Supplementary file***

1). In addition, mitochondrial components, including those involved in respiration and H_2S

detoxification, played critical roles in triggering and maintaining escape behavior in H_2S. In

contrast, we did not find evidence that globins, K^+ channels, or biogenic amine signaling

contributed significantly to the behavioral avoidance to H_2S (***Supplementary file 1***).

cGMP signaling in ASJ cilia contributes to H_2S avoidance

One group of mutants exhibiting reduced locomotory speed in H_2S had defects in ciliogenesis

(***Figure 2A–C***). This group included *daf-19* mutants lacking cilia, *dyf-3* mutants with reduced

cilia length, and *dyf-7* mutants in which ciliogenesis occurs but the cilia are not anchored to

their sensilla and are not exposed to the external environment (Heiman & Shaham, 2009;

Murayama, Toh, Ohshima, & Koga, 2005; Starich et al., 1995; Swoboda, Adler, & Thomas,

2000). These findings suggest that ciliogenesis and exposure of cilia to the external

environment are essential for animals to escape H_2S exposure. In addition, these mutants

exhibited consistently high reversal rate, likely associated with their reduced locomotory speed

(*Figure 2—figure supplement 1A–F*). Cilia-mediated sensory responses often involve the

activation of guanylate cyclases and the opening of cGMP-gated channels (Ferkey, Sengupta,

& L'Etoile, 2021). We observed significantly attenuated locomotory speed in response to H_2S

in *tax-4* or *tax-2* mutants, which lack key subunits of cilia-enriched cGMP-gated channels

(*Figure 2D, E*). In a comprehensive survey of all guanylate cyclase mutants, the *daf-11* strain

emerged as the only mutant showing speed response defects in the avoidance of H_2S (*Figure*

2D, supplementary file 1). Similar to the cilia mutants, animals deficient in *daf-11* had low

speed, attenuated omega turns, and constantly high reversal rate (*Figure 2D, Figure 2—figure*

supplement 1G, H). Selective expression of *daf-11* in ASJ neurons, but not in other neurons,

partially restored the speed response to acute H_2S (*Figures 2F, Figure 2—figure supplement*

2A–D). In addition, the H_2S response defect of *tax-4* mutants was also rescued by selective

expression of its cDNA in ASJ neurons (*Figure 2G*), suggesting that the receptor guanylate

cyclase DAF-11 and the CNG channel TAX-4 act at least partially in ASJ neurons to promote

acute avoidance to H_2S. To validate the role of ASJ neurons in H_2S responses, we blocked

neurotransmission from ASJ using the catalytic domain of *Tetanus Toxin (TeTx)*, which impairs

neurosecretion by specifically cleaving synaptobrevin (Schiavo et al., 1992). Cell-specific

expression of *TeTx* in ASJ significantly inhibited the speed response to H_2S *(Figure 2H)*.

Surprisingly, using the Ca^{2+} sensor GCaMP6s to monitor a direct response of ASJ neurons to

H_2S, we observed no H_2S-induced Ca^{2+} transients under any of the tested conditions, whereas

CO_2-evoked Ca^{2+} increases were readily detected *(Figure 2—figure supplement 2E, F)*. These

findings suggest that although ASJ neuronal activity and neurosecretion contribute to the H_2S

responses, ASJ neurons are unlikely to serve as the primary H_2S sensor.

Starvation modulates H_2S avoidance

In the candidate gene screen, another set of mutants with reduced speed responses to H_2S had

previously been shown to be defective in response to CO_2 (Hallem & Sternberg, 2008).

Specifically, the H_2S-evoked speed response was reduced in mutants deficient in nutrient-

sensitive signaling, including the insulin receptor DAF-2, the TGF-β ligand DAF-7, and the

nuclear hormone receptor NHR-49 (*Figure 3A–C, Figure 3—figure supplement 1A*). Similarly, mutants lacking other CO_2 response modulators, such as the calcineurin subunits TAX-6 and CNB-1, also failed to respond to H_2S (*Figure 3—figure supplement 1B, C*). In addition to the abolished speed response, we also observed that the H_2S-evoked omega-turn response was nearly abolished in *daf-2* mutants, and high proportion of animals exhibited persistently high reversal rate regardless of H_2S stimulation (*Figure 3—figure supplement 1D–F*). The H_2S avoidance defects observed in *daf-2* and *daf-7* single mutants were fully suppressed in *daf-2; daf-16* and *daf-7; daf-3* double mutants *(Figure 3A, B).* Therefore, it is likely that insulin and TGF-β signaling modulate H_2S responses by regulating the expression of relevant genes via DAF-16 and DAF-3 transcription factors, respectively. Given that H_2S-evoked speed response was modulated by the nutrient-sensitive pathways, we further explored whether the locomotory response to H_2S was sensitive to the nutrient state. Similar to the response to CO_2 (Bretscher et al., 2008; Hallem & Sternberg, 2008), we observed that a 24-hour period of starvation suppressed the speed response to H_2S *(Figure 3D).* However, the CO_2 sensor GCY-9 was dispensable for H_2S-evoked avoidance *(Figure 3—figure supplement 1G–I).* These findings suggest that even though acute avoidance of CO_2 and H_2S are modulated through shared mechanisms, distinct molecular sensors are used.

The O_2 sensing circuit antagonizes H_2S avoidance in wild isolates

Wild strains of *C. elegans* thrive in environmental niches characterized by variable O_2, CO_2 and H_2S levels (Adams et al., 1979; Bretscher et al., 2008; Budde & Roth, 2011; Gea, Barrena, Artola, & Sanchez, 2004; Hallem & Sternberg, 2008; Morra & Dick, 1991; Oshins, Michel, Louis, Richard, & Rynk, 2022; Patange et al., 2025; Rodriguez-Kabana et al., 1965). Compared to the laboratory reference strain N2, wild isolates display robust responses to high O_2 but reduced sensitivity to CO_2 stimulation (Beets et al., 2020; Bretscher et al., 2008; Carrillo, Guillermin, Rengarajan, Okubo, & Hallem, 2013; Hallem & Sternberg, 2008; Kodama-Namba et al., 2013; McGrath et al., 2009). It is thought that the unique O_2 and CO_2 responses of N2 strain evolved during its domestication on solid media at atmospheric gas concentrations

(Sterken, Snoek, Kammenga, & Andersen, 2015). We hypothesized that wild *C. elegans* strains

might also exhibit an attenuated response to H_2S, and therefore included a set of wild strains in

our candidate gene screen. Consistent with this hypothesis, wild isolates including CB4855,

CB4856, and CB4858 exhibited reduced locomotory speed in H_2S (***Figure 3E, Figure 3—***

figure supplement 1J).

A naturally occurring variation in the neuropeptide receptor NPR-1 predominantly

determines the differences in locomotory responses to O_2 and CO_2 between the laboratory N2

strain and wild isolates. The NPR-1(215F) variant found in wild isolates is less active than the

NPR-1(215V) found in N2 animals (de Bono & Bargmann, 1998). We wondered whether the

variations in the *npr-1* gene also contribute to differences in the acute response to H_2S. The

absence of *npr-1* was sufficient to reduce the speed response to H_2S, suggesting that active

NPR-1 signaling promotes H_2S avoidance (***Figure 3F***). NPR-1(215V) primarily acts within the

RMG interneurons to modulate responses to environmental stimuli by inhibiting signal output

from these neurons (***Figure 3G***) (Laurent et al., 2015; Macosko et al., 2009). Cell-specific

expression of the highly active *npr-1(215V)* variant in RMG neurons effectively restored the

H_2S-evoked locomotory speed in *npr-1* null mutant worms (***Figure 3H***). Furthermore,

optogenetic stimulation of RMG interneurons using channelrhodopsin-2 (ChR2) significantly

dampened the locomotory response to H_2S in N2 animals, suggesting that high RMG activity

inhibits H_2S responses (***Figure 3—figure supplement 1K***).

The perception of 21% O_2 involves the soluble guanylate cyclases GCY-35/GCY-36

and the downstream cyclic nucleotide-gated (CNG) channels TAX-4/TAX-2 in the URX, AQR,

and PQR O_2-sensing neurons, which relay the signals to RMG (Busch et al., 2012; Cheung,

Cohen, Rogers, Albayram, & de Bono, 2005; Couto, Oda, Nikolaev, Soltesz, & de Bono, 2013;

Gray et al., 2004; Laurent et al., 2015; Persson et al., 2009; Zimmer et al., 2009). We

investigated whether disrupting the O_2 sensing machinery would have the same effect as

inhibiting RMG signaling pathway by NPR-1(215V). Mutating *gcy-35* and *tax-4*, or genetically

ablating the O_2 sensing neurons URX, AQR, and PQR by cell-specific expression of the pro-

apoptotic gene *egl-1* restored H_2S-evoked speed response in *npr-1* null mutants (***Figure 3I–K***).

Specific expression of *gcy-35* cDNA in O_2 sensing neurons was sufficient to repress this

response to H_2S in *npr-1; gcy-35* double mutants (**Figure 3L**). Finally, enhancing the

presynaptic activity of URX, AQR, PQR neurons by the expression of a gain-of-function PKC-

1(A160E) (Dekker, Mcintyre, & Parker, 1993; Hiroki et al., 2022) significantly dampened H_2S-

induced locomotory speed (**Figure 3—figure supplement 1L**). Taken together, these

observations indicate that activation of the O_2-sensing circuit inhibits the H_2S-evoked

avoidance response.

H_2S exposure reprograms gene expression in *C. elegans*

Wild isolates that were relatively insensitive to H_2S exposure (**Figure 3E, Figure 3—figure**

supplement 1J) likely developed sensory and physiological adaptations to persist in

environments where H_2S may transiently increase (Budde & Roth, 2011; Patange et al., 2025;

Rodriguez-Kabana et al., 1965; Romanelli-Cedrez et al., 2024). These adaptations to H_2S are

mediated, at least in part, by the reprogramming of gene expression (Miller et al., 2011a). To

identify the genes whose expression was induced by prolonged H_2S exposure, we conducted a

comparative analysis of transcriptome profiles in animals exposed to 50 ppm or 150 ppm H_2S

for 1 hour, 2 hours, or 12 hours. RNA-seq analysis revealed that exposure to either 50 ppm or

150 ppm H_2S for one hour was sufficient to trigger significant changes in gene expression, with

518 or 304 genes showing differential expression, respectively (**Figure 4A, Supplementary file**

2). Genes induced by H_2S exposure for 1 or 2 hours displayed considerable overlap (**Figure 4B,**

Figure 4—figure supplement 1A). Gene Ontology (GO) analysis revealed that biological

processes such as defense against bacteria and cysteine biosynthesis from L-serine were

significantly enriched after H_2S exposure (**Figure 4C, D, Figure 4—figure supplement 1B, C**).

As expected, we observed a robust induction of genes involved in H_2S detoxification (**Figure**

4E, F) (Miller, Budde, & Roth, 2011b; Niu et al., 2011; Vora et al., 2022). H_2S is detoxified in

mitochondria by sulfide:quinone oxidoreductase SQRD-1 to produce persulfide, which is then

metabolized by sulfur dioxygenase ETHE-1 to generate sulfite. Thiosulfate transferase further

oxidizes sulfite to produce thiosulfate (Hildebrandt & Grieshaber, 2008). Glutathione S-

transferases (GSTs) remove accumulated sulfur molecules during H_2S oxidation (Jackson, Melideo, & Jorns, 2012). Notably, *gst-19* and *sqrd-1* were among the most significantly upregulated genes after one or two hours of exposure to either 50 ppm or 150 ppm H_2S, while *ethe-1* showed a weak increase (***Figure 4E, F, Figure 4—figure supplement 1D, E, Supplementary file 2***).

H_2S exposure is known to activate the HIF-1 pathway (Budde & Roth, 2010, 2011; Ma, Vozdek, Bhatla, & Horvitz, 2012; Miller et al., 2011a; Miller & Roth, 2007; Vora et al., 2022). In line with this, a proportion of genes with HIF-1 regulated promoters displayed increased expression after 1 or 2 hours of H_2S exposure (***Figure 4—figure supplement 1F***). Among these were the aforementioned detoxifying enzymes GST-19, SQRD-1, and ETHE-1, as well as CYSL-1, CYSL-2, and CDO-1, which are involved in cysteine metabolism (***Supplementary file 2***). Consistent with earlier studies, we also observed differential expression of a set of SKN-1 targets in response to chronic H_2S exposure (***Figure 4—figure supplement 1G***) (Horsman et al., 2019; Miller et al., 2011a; Niu et al., 2011). This included the heat shock protein genes *hsp-16.2* and *hsp-16.41* (***Figure 4E, F, Figure 4—figure supplement 1D, E***). Surprisingly, we detected a rapid increase in the expression of genes associated with intracellular H_2S production, including the methanethiol oxidase-encoding gene *semo-1* and the mercaptopyruvate sulfurtransferase-encoding gene *mpst-3* (***Figure 4E, F, Figure 4—figure supplement 1D, E***) (Philipp et al., 2022; Qabazard et al., 2014). Elevated expression of 3-mercaptopyruvate sulfurtransferase has similarly been reported in sulfide spring fish, suggesting a conserved response mechanism to H_2S (Kelley et al., 2016; Mathew, Schlipalius, & Ebert, 2011; Qabazard et al., 2014). However, the functional implications of increased expression of H_2S synthesis genes remains unclear.

When exposed to H_2S for 12 hours, the differentially expressed genes showed reduced overlap with those identified at earlier time points (***Figure 4B, Figure 4—figure supplement 1A***), suggesting that distinct defense strategies may be employed at various stages of H_2S exposure. Furthermore, substantial overlap was observed between the sets of genes induced by 50 ppm and 150 ppm H_2S after 1h or 2h exposure. However, this overlap decreased

significantly after 12 hours exposure (***Figure 4—figure supplement 1H–J***). Additionally, we

noticed that 150 ppm H_2S specifically triggered the expression of the HIF-1 regulator *rhy-1* and

the HIF-1 target *nhr-57* at all exposure time points, a response that was less pronounced at 50

ppm H_2S (***Figures 4E, F, Figure 4—figure supplement 1D, E, K and L***). Together, these

findings suggest that multiple defense mechanisms are induced to adapt to the stress imposed

by H_2S, which can diverge over time and at different H_2S concentrations.

Locomotory speed response to H_2S is modulated by HIF-1-induced detoxification

Given that HIF-1 signaling is robustly activated shortly after H_2S exposure (***Figure 4—figure***

supplement 1F) (Budde & Roth, 2010, 2011; Miller et al., 2011a), we next sought to determine

how HIF-1 signaling might contribute to H_2S-evoked avoidance response, and began by

examining the effects of HIF-1 stabilization. Acclimating animals in 1% O_2 for 12 hours or

longer significantly decreased the speed response to H_2S, indicating that HIF-1 stabilization

inhibits the H_2S-evoked avoidance behavior **(*Figure 5A*)**. To confirm this observation, we

examined several mutants with stabilized HIF-1. The conserved proline-4-hydroxylase

PHD/EGL-9 and the von Hippel-Lindau (VHL) tumor suppressor protein are required for HIF-

1 degradation (Kaelin & Ratcliffe, 2008; Semenza, 2010). Disruption of either *egl-9* or *vhl-1*

leads to HIF-1 stabilization under normoxic conditions. Similar to the effect of prolonged

hypoxia, both mutants exhibited a markedly reduced locomotory speed in H_2S **(*Figure 5B*)**.

Interestingly, the omega-turn and reversal responses to H_2S were also substantially inhibited in

egl-9 mutants **(*Figure 5—figure supplement 1A, B*)**. In contrast, the locomotory speed

response to 1% O_2 remained largely unaffected in *egl-9* deficient animals **(*Figure 5—figure***

***supplement 1C*),** suggesting a selective impairment of the H_2S response. In addition, expressing

the non-degradable forms of HIF-1, *hif-1*(P621A or P621G), under the pan-neuronal *unc-14*

promoter or under the endogenous *hif-1* promoter, was sufficient to inhibit H_2S evoked speed

responses (***Figure 5C***). Together, these findings indicate that the activation of HIF-1 signaling

during prolonged H_2S or hypoxia exposure mediates an adaptive response associated with

reduced behavioral responses to H_2S, and that HIF-1 stabilization in neurons alone is sufficient

to promote this behavioral adaptation.

On the other hand, animals lacking HIF-1 signaling have previously been shown to

display increased sensitivity to H_2S and become paralyzed rapidly upon exposure (Budde &

Roth, 2010; Horsman et al., 2019). Consistent with this, *hif-1* mutants exhibited a brief increase

in speed upon H_2S exposure, which rapidly declined to baseline levels (***Figure 5D***). They also

displayed a stronger initial omega-turn response and greater inhibition of the reversal rate than

WT (***Figure 5E, F***), confirming that animals deficient in HIF-1 signaling are sensitized to H_2S

exposure. This transient avoidance behavior was also observed in *egl-9; hif-1* and *vhl-1; hif-1*

double mutants (***Figure 5G, H***), suggesting that this phenotype is linked to the absence of HIF-

1 signaling. Importantly, *hif-1* mutants displayed a robust speed response to other stimuli, such

as near-UV light and 1% O_2 stimulation ***(Figure 5I, Figure5—figure supplement 1D)***,

suggesting that the brief avoidance response is specific to H_2S rather than reflecting a general

locomotory defect. Further supporting increased sensitivity to H_2S in *hif-1* mutants, the

locomotory speed response of *hif-1* mutants to subsequent near-UV light stimulation was nearly

abolished after 30 minutes of H_2S exposure, whereas *egl-9* mutants and WT animals remained

fully responsive ***(Figure 5I)***. The reduced movement and lack of responsiveness to other

stressors in *hif-1* mutants resemble the stress-induced sleep-like behavior previously observed

in response to noxious heat (Byrne Rodgers & Ryu, 2020). Prompted by this observation, we

next explored whether HIF-1-induced detoxification genes were also required to maintain H_2S-

evoked behavioral responses (***Figure 5—figure supplement 1E***). Similar to *hif-1* mutants,

sqrd-1 and *ethe-1* mutants displayed brief responses to H_2S, with locomotory activity rapidly

returning to baseline levels ***(Figure 5J, K, Figure 5—figure supplement 1F–I)***. However, both

mutants responded normally to 1% O_2, suggesting a specific hypersensitivity to H_2S toxicity

(Figure 5—figure supplement 1J). Furthermore, we examined the contribution of additional

genes whose expression was upregulated in response to H_2S, including the

sulfhydrylase/cysteine synthase-encoding genes *cysl-1*, *cysl-2*, and *cysl-3*, as well as *semo-1*.

Disrupting any of these genes phenocopied the responses observed in *hif-1, ethe-1*, or *sqrd-1*

mutants, while they displayed normal responses to 1% O_2 (**Figure 5L, Figure 5—figure**

supplement 1D, K–P). These observations suggest that while the HIF-1-induced H_2S

detoxification system is not essential for the initial sensing of H_2S, it is required to maintain

animals' locomotory activity in the presence of H_2S.

Labile iron pool sustains the locomotory activity in H_2S

Among the most differentially regulated genes upon exposure to H_2S, we observed consistent

downregulation of *ftn-1* and to a lesser extent *ftn-2*, whereas *smf-3* expression was increased

under specific H_2S conditions (**Figure 6A, B, Figure 6—figure supplement 1A–D**). The

ferritin-encoding genes *ftn-1* and *ftn-2,* along with the iron transporter *smf-3*, are critical for

maintaining intracellular labile iron homeostasis in *C. elegans* (Ackerman & Gems, 2012;

Gourley, Parker, Jones, Zumbrennen, & Leibold, 2003; Rajan et al., 2019; Romney, Newman,

Thacker, & Leibold, 2011), suggesting that H_2S exposure may interfere with cellular iron

storage. It has been shown that disrupting *ftn-1* increases labile iron pools in the cytoplasm,

while its overexpression reduces these levels (Anderson & Leibold, 2014; Romney et al., 2011).

We found that *ftn-1* mutants had enhanced speed response to H_2S, whereas *ftn-1* overexpression

attenuated H_2S-induced speed increase without damaging acute response to 1% O_2 (**Figure 6C,**

D, Figure 6—figure supplement 1E). In addition, *smf-3* mutants, which impair iron uptake,

exhibited a reduced speed response to H_2S (**Figure 6D**). These results support the idea that H_2S

exposure disrupts iron homeostasis.

To complement our genetic analysis, we pharmacologically depleted labile irons. In

the presence of the iron chelator 2,2'-dipyridyl (BP), the H_2S-evoked speed response was fully

lost (**Figure 6E**), whereas locomotory speed in response to hypoxia was maintained (**Figure**

6—figure supplement 1F). We noticed that, similar to *hif-1* mutants, both *smf-3* mutants and

BP-treated animals exhibited a strong initial omega-turn response to H_2S (**Figure 6—figure**

supplement 1G, I), suggesting that these animals are hypersensitive to H_2S toxicity. One

possibility is that low iron levels compromise H_2S detoxification capacity, as the detoxification

enzyme ETHE-1 requires iron for its activity (Kabil & Banerjee, 2012; Pettinati, Brem,

McDonough, & Schofield, 2015). By contrast, increasing iron levels by feeding ferric

ammonium citrate (FAC) or by disrupting *ftn-1* significantly sustained animals' locomotory

speed in H_2S (***Figure 6D, E***), likely by delaying H_2S-mediated iron depletion. However, the

reorientation and reversals were not significantly affected (***Figure 6—figure supplement 1G–***

J). This impact of FAC supplementation and *ftn-1* disruption on speed response was largely

lost in the absence of *hif-1* (***Figure 6F, G***), suggesting that the effects of iron require HIF-1-

dependent detoxification (***Figure 6H***). Yet, increased iron availability modestly improved the

locomotory activity of *hif-1* mutants in H_2S (***Figure 6F, G***), presumably by compensating for

the low iron levels in *hif-1* mutant animals (Rajan et al., 2019; Romney et al., 2011).

Mitochondrial function modulates H_2S-evoked locomotion responses

The mitochondrial electron transport chain (ETC) is not only the primary target of H_2S

toxicity, but is also an essential component of the H_2S detoxification pathways (Cooper &

Brown, 2008; Khan et al., 1990; Nicholls & Kim, 1982; Romanelli-Cedrez et al., 2024).

Based on this, we hypothesized that ETC plays a central role in the locomotory response to

H_2S. Indeed, mutating the genes critical for ETC function, including *gas-1*, *clk-1*, *mev-1,* and

isp-1, impaired the H_2S-evoked speed and reversal responses (***Figure 7A–D, Figure 7—***

figure supplement 1A–D). These data suggest that a functional ETC is required to trigger and

support H_2S-evoked avoidance behavior or that dysfunctional ETC induces an adaptive

response to cellular stressors. Consistently, exposing animals to rotenone, an ETC complex I

inhibitor, robustly inhibited the speed responses to H_2S at all time points we tested (***Figure***

7E). Interestingly, we noticed that transient rotenone exposure substantially increased

animals' basal locomotory speed (***Figure 7E***) (Onukwufor et al., 2022), suggesting that acute

ETC interruption evokes behavioral avoidance, in a manner similar to that observed with

other noxious stimuli. However, a 2-hour exposure to rotenone abolished the H_2S-evocked

speed response without increasing the basal locomotory speed (***Figure 7E***), suggesting that

prolonged ETC inhibition may either impair the cellular capacity to support H_2S-evoked

behavioral responses or drive animals' adaptation to cellular stress and consequently reduce

their locomotory response to H_2S.

Given that rotenone exposure and mutations in ETC genes increase mitochondrial

ROS generation (Ishii et al., 2013; Ochi et al., 2016; Ramsay & Singer, 1992; Yang &

Hekimi, 2010; Zorov, Juhaszova, & Sollott, 2014), and that high levels of H_2S also induce

ROS production (Jia et al., 2020), we sought to determine how mitochondrial ROS

contributes to speed response to high H_2S. We focused on superoxide dismutases (SODs), key

enzymes in superoxide detoxification. The *C. elegans* genome encodes five *sod* genes: *sod-1*

and *sod-5* encode cytosolic Cu/ZnSODs, *sod-2* and *sod-3* encode mitochondrial MnSODs,

and *sod-4* encodes the extracellular Cu/ZnSOD isoforms (Doonan et al., 2008; Zubovych,

Straud, & Roth, 2010). All *sod* single mutants exhibited a robust initial response to 150 ppm

H_2S *(Figure 7F, G).* Interestingly, the mitochondrial SOD mutants *sod-2* and *sod-3*, as well

as *sod-5*, showed an initial burst of locomotory activity followed by a rapid decline in speed

in the presence of H_2S *(Figure 7F, G)*, but maintained a high locomotory speed in response to

1% O_2 *(Figure 7—figure supplement 1E)*. In the quintuple *sod-1; sod-2; sod-3; sod-4; sod-5*

mutant, which lacks all superoxide dismutases and presumably accumulates higher ROS

levels, the H_2S-evoked speed and omega-turn responses were nearly abolished, and the

proportion of animals exhibiting reversals remained consistently high, a pattern resembling

that of *egl-9* and ETC mutants *(Figure 7G, Figure 7—figure supplement 1F, G)*. Similar to

sod-2 and *sod-3*, the quintuple *sod* mutant exhibited robust speed response to 1% O_2 (Zhao et

al., 2022). In addition, the quintuple mutants responded normally to near-UV light after 30

minutes of H_2S exposure *(Figure 7—figure supplement 1H)*, indicating that their

neuromuscular function remains largely preserved. Therefore, high ROS levels in *sod* mutants

do not substantially compromise the detoxification capacity required for survival in H_2S.

Instead, high ROS appears to promote adaptation to H_2S-induced stress, presumably by

activating several stress-responsive pathways such as HIF-1, NRF2/SKN-1, and DAF-

16/FOXO *(Figure 7H)* (Lennicke & Cocheme, 2021; Patten, Germain, Kelly, & Slack, 2010).

Discussion

Animals' behavior and physiology are profoundly influenced by the environments in which they evolved. The laboratory strain N2, which is adapted to low atmospheric CO_2 and H_2S concentrations, robustly avoids both gases (Beets et al., 2020; Bretscher et al., 2008; Carrillo et al., 2013; Hallem & Sternberg, 2008; Kodama-Namba et al., 2013; McGrath et al., 2009). Upon acute exposure to H_2S above 75 ppm, N2 exhibits an avoidance behavior characterized by reorientation and increased locomotion. This response is significantly reduced in wild isolates. These divergent responses between the laboratory strain and wild isolates are primarily driven by variations in the *npr-1* gene, which encodes a neuropeptide receptor, suggesting that *C. elegans* has undergone rapid evolutionary adaptation to its environment.

The behavioral response to a specific sensory stimulus is usually shaped by an interplay of multiple environmental and physiological cues. We observe that conditions dampening *C. elegans'* response to CO_2 also impair the response to H_2S. Specifically, the speed response to H_2S is significantly inhibited at high O_2 levels, and in mutants with defective insulin or TGF-β pathway that signal starvation. However, despite these shared modulation, CO_2 and H_2S sensing involve distinct molecular mechanisms. For example, CO_2 responses are mediated by the guanylate cyclase GCY-9 in BAG neurons (Hallem et al., 2011), which is dispensable for H_2S sensing. In addition, acute exposure to H_2S induces stronger reorientation but delayed acceleration compared to CO_2, suggesting different sensing and information processing mechanisms (Bretscher et al., 2008). Therefore, even though responses to CO_2 and H_2S are modulated by overlapping regulatory mechanisms, the two gases are detected by distinct sensors and processed differently.

In a candidate gene survey, we excluded the potential involvement of globins, K^+ channels, biogenic amines, and most guanylate cyclases in H_2S responses. However, we observed that H_2S avoidance requires the activity of guanylate cyclase DAF-11 in ASJ neurons, as well as neurosecretion from these neurons. Since the DAF-11 pathway and neurosecretion from ASJ neurons regulate developmental programs that modify sensory functions in *C. elegans* (Murakami, Koga, & Ohshima, 2001), it is not surprising that *daf-11* mutants display

pleiotropic phenotypes including impaired H_2S and CO_2 responses (Hallem & Sternberg, 2008).

In addition, we did not detect an H_2S-evoked calcium transient in ASJ neurons. Therefore,

although ASJ neurons and DAF-11 activity are clearly required, the exact sensory neurons,

circuits, and molecular triggers mediating acute H_2S avoidance remain to be elucidated.

Consistent with previous reports (Horsman et al., 2019; Miller et al., 2011a), H_2S

exposure substantially alters the gene expression in *C. elegans*, including those involved in H_2S

detoxification and iron homeostasis pathways. The repression of *ftn-1* and induction of *smf-3*

observed during H_2S exposure suggest that H_2S depletes intracellular labile iron. Iron is a

critical cofactor required for the activity of many enzymes and supports a wide range of cellular

processes (Dev & Babitt, 2017; Hentze, Muckenthaler, & Andrews, 2004). One important

enzyme that requires labile iron as cofactor is PHD/EGL-9, which targets HIF-1 for degradation.

Changes in iron availability are therefore expected to modulate EGL-9 activity, which in turn

alter the expression of HIF-1-regulated genes (Liochev, 1996; Myllyharju & Kivirikko, 1997;

Read, Bentley, Archer, & Dunham-Snary, 2021; Xu & Moller, 2011). However, ETHE-1, a

key H_2S detoxification enzyme upregulated upon HIF-1 stabilization, also requires iron for its

persulfide dioxygenase activity (Kabil & Banerjee, 2012; Pettinati et al., 2015). Therefore, H_2S-

evoked responses under varying iron availability are determined by the combined effects of

labile iron on HIF-1 signaling and on the H_2S detoxification enzymes, among other factors. For

instance, although reduced iron availability in *smf-3* or BP-treated animals may promote the

HIF-1 induced detoxification pathways by inactivating EGL-9, the actual enzymatic clearance

of H_2S is impaired due to low ETHE-1 activity. Supporting this, *smf-3* mutants and BP-treated

animals display increased sensitivity to H_2S, including enhanced initial omega-turn responses

and rapid inhibition of locomotion. In contrast, increased iron availability in *ftn-1* mutants or

FAC-treated animals likely delays iron depletion during H_2S exposure, preserving ETHE-1

detoxification capacity and postponing the onset of HIF-1-mediated adaptations that are

associated with reduced locomotion. This may explain the sustained high locomotory speed

observed in *ftn-1* mutants or FAC-treated animals. In the absence of HIF-1, iron

supplementation only partially improves the locomotory response to H_2S, likely because the

H_2S detoxification system including ETHE-1 cannot be transcriptionally induced. The modest

effect observed may instead reflect correction of iron deficiency in *hif-1* mutants.

At concentrations below 50 ppm, H_2S is well tolerated or even preferred, with

beneficial effects such as improved thermotolerance (Budde & Roth, 2010; Fawcett et al., 2015;

Horsman et al., 2019; Miller & Roth, 2007; Qabazard et al., 2014). Notably, behavioral

avoidance is absent below 50 ppm H_2S, suggesting that escape behavior is triggered only when

H_2S is not efficiently detoxified. This led us to hypothesize that animals with enhanced

detoxification capacity would show reduced avoidance of otherwise toxic H_2S levels. Indeed,

the speed response to H_2S is attenuated when the detoxification program is upregulated under

conditions of constitutive activation of HIF-1, such as in *egl-9* or *vhl-1* mutants, after prolonged

exposure to 1% O_2, or when HIF-1 signaling is specifically activated in neurons. These

observations support the idea that the initial avoidance response to H_2S is likely driven by acute

toxicity, while the subsequent decline in speed reflects a rapid activation of adaptive

mechanisms, particularly through stabilization of HIF-1. The fact that wild type animals remain

responsive to other stimuli after prolonged H_2S exposure suggests that reduced locomotory

speed in H_2S reflects active desensitization rather than general paralysis. However, persistently

high H_2S exposure is likely to exhaust cellular defense systems, leading to toxicity and paralysis.

By contrast, the rapid loss of locomotion observed in *hif-1* mutants and detoxification-defective

mutants is mediated by a mechanism distinct from adaptation. Their loss of responsiveness to

other stimuli after H_2S exposure suggests that these animals likely become sensitized and

rapidly intoxicated by H_2S due to impaired detoxification. Therefore, H_2S exposure promotes

a behavioral program that includes an initial reorientation and acceleration responses followed

by a progressive adaptation driven by cellular detoxification processes. A similar behavioral

program has been observed in *C. elegans* during noxious heat exposure, which induces short-

term heat avoidance followed by long-lasting cytoprotective adaptation and a gradual reduction

in avoidance (Byrne Rodgers & Ryu, 2020).

The interaction of H_2S with mitochondrial ETC is multifaceted, acting as an electron

donor at low concentrations and becoming a potent toxicant at high levels (Szabo et al., 2014),

in part by promoting superoxide generation through complex IV inhibition and reverse electron

transport at complex I (Cooper & Brown, 2008; Jia et al., 2020; Khan et al., 1990; Nicholls &

Kim, 1982; Romanelli-Cedrez et al., 2024). We propose that mitochondria play a dual role in

H_2S-evoked locomotory avoidance. On one hand, the mitochondrial ETC contributes to H_2S

detoxification and promotes adaptation. On the other hand, toxic levels of H_2S remodel the

ETC, leading to increased ROS production, which may serve as a trigger for the avoidance

response. Supporting the idea that acute mitochondrial ROS generation initiates avoidance of

high H_2S levels, short-term rotenone exposure, known to promote mitochondrial ROS

formation (Ochi et al., 2016; Ramsay & Singer, 1992; Zorov et al., 2014), substantially

increases locomotory speed (Onukwufor et al., 2022). Meanwhile, the speed response to high

H_2S is fully suppressed by rotenone. This inhibition could result either from excessive

mitochondrial ROS generated by rotenone, which may dampen the H_2S-triggered ROS spike,

or from direct complex I inhibition, which may disrupt other H_2S signaling processes required

to initiate avoidance. However, persistent mitochondrial ROS production appears to suppress

high locomotory speed and inhibit responsiveness to H_2S, as observed after 2-hour rotenone

exposure, in mitochondrial ETC mutants, and in animals lacking all superoxide dismutases.

One likely explanation is that mitochondrial ROS can activate a variety of stress-responsive

pathways, including HIF-1, NRF2/SKN-1, and DAF-16/ FOXO signaling (Lennicke &

Cocheme, 2021; Patten et al., 2010), priming animals' adaptation to prolonged stress rather

than causing toxicity. This is supported by the observation that even though SOD-deficient

animals do not display strong initial locomotory responses to H_2S, they remain responsive to

other stimuli after 30 minutes of H_2S exposure, suggesting that high ROS levels do not

compromise general viability or the H_2S detoxification capacity. Therefore, we favor a model

in which mitochondrial ROS exert a biphasic effect on H_2S-induced avoidance, facilitating H_2S

detection and avoidance under acute conditions, and contributing to locomotory inhibition

when it is chronically elevated.

In summary, this study unveils a novel behavior of *C. elegans* in their avoidance to

H_2S. The speed response to H_2S is intricately shaped by environmental context, integrating

inputs from other sensory cues such as food availability and O_2 levels. Although we did not

identify the specific neural circuit mediating H_2S signaling, our findings suggest that H_2S-

induced locomotion arises from acute remodeling of mitochondrial ETC and mitochondrial

ROS production. We emphasize the vital role of the HIF-1-induced detoxification pathways

and iron homeostasis in protecting against H_2S-induced mitochondrial toxicity. In future work,

it will be important to delineate the neural circuits underlying H_2S-evoked locomotory

avoidance and to clarify how H_2S-induced changes in mitochondrial respiration and labile iron

pools shape these responses.

Materials and Methods

Strains

C. elegans were maintained using standard protocols (Brenner, 1974). Strains used in this study

are listed in ***Supplementary file 1*** and ***Supplementary file 3***.

Preparation of H_2S gas mixture

Different H_2S concentrations were created as previously described (Miller & Roth, 2007).

Briefly, the H_2S-containing gas mixture were prepared by diluting 5000 ppm H_2S in nitrogen

(N_2) with 7% O_2 balanced with N_2. The gas flow was tuned by Sierra Smart-Trak 100 mass

flow controllers. H_2S concentration in the mixture was measured using two H_2S detectors (MSA

ALTAIR 2X gas detector for H_2S and Clip Single Gas Detector, SDG, CROWCON). The pre-

defined gas mixtures of 7% O_2, 1% O_2 and 5% CO_2 with N_2 were purchased from Air Liquide

Gas AB. 5000 ppm H_2S stock in N_2 was obtained from Linde Gas AB. Gas mixtures in all

experiments were hydrated before use.

Molecular Biology

The Multisite Gateway system (Thermo Fisher Scientific, United States) was used to generate

expression vectors. Promoters, including *gcy-37* (2.7kb), *trx-1* (1 kb), *ftn-1* (2 kb), *daf-11* (3 kb),

odr-3 (2.7 kb), *gpa-11* (3 kb), *sra-6* (3 kb), *odr-1* (2.4 kb), *flp-21* (4.1kb), and *ocr-2* (2.4 kb) were

amplified from N2 genomic DNA and cloned into pDONR P4P1 using BP clonase. g*cy-35*,

tax-4, and *pkc-1* cDNAs were amplified using the first strand cDNA library as the template,

while *daf-11* and *ftn-1* genomic sequences were amplified using genomic DNA as the template.

The entry clones of these genes pDONR 221 were generated using BP reaction. To generate

the gain of function mutation of *pkc-1* (A160E), Q5 Site-Directed Mutagenesis Kit (NEB) was

used according to manufacturer instructions. All expression vectors were generated using LR

reaction. The primer sequences were displayed in ***Supplementary file 4***.

To generate transgenic animals, the *daf-11* expression vectors were injected at the

concentration of 20 ng/μl supplemented with 50 ng/μl of a coelomocyte co-injection marker

(*unc-122p::GFP*) and 30 ng/μl of 1kb DNA ladder. For *tax-4* gene, the injection mixtures were

prepared using 40 ng/μl of *tax-4* expression vectors, 50 ng/μl of a coelomocyte co-injection

marker and 10 ng/μl of 1kb DNA ladder. The rest of expression constructs were injected at 50

ng/μl together with 50 ng/μl of a coelomocyte marker.

CRISPR/Cas9 genome editing

Genes were disrupted using CRISPR/Cas9 mediated genome editing as described (Dokshin,

Ghanta, Piscopo, & Mello, 2018; Ghanta & Mello, 2020). The strategy involved the utilization

of homology-directed insertion of custom designed single strand DNA template (ssODN),

which had two homology arms of 35bp flanking the targeted PAM site. Between two homology

arms, a short sequence containing a unique restriction enzyme cutting site as well as in-frame

and out-of-frame stop codons, was included. The insertion of ssODN template would delete 16

bases of coding sequence, ensuring the proper gene disruption. To prepare the injection cocktail,

0.5 μl of Cas9 protein (IDT) was mixed with 5 μl of 0.4μg/μl tracrRNA (IDT, United States)

and 2.8 μl of 0.4 μg/μl crRNA (IDT). The mixture was incubated at 37°C for at least 10 minutes

before 2.2 μl of 1μg/μl ssODN (or 500 ng dsDNA) and 2 μl of 0.6 μg/μl *rol-6* co-injection

marker were added. Nuclease-free water was used to bring the final volume to 20 μl. The

injection mixture was centrifuged for 2 minutes before use.

Behavioral assays

H$_2$S evoked locomotion activity was monitored as described previously (Laurent et al., 2015; Zhao et al., 2022). Briefly, OP50 bacteria were seeded on the assay plates 16 hours before use. The border of bacterial lawn was removed using a PDMS stamp. For each assay, 25 to 30 day-one adult animals were picked onto assay plates, allowed to settle down for 15 minutes, and subsequently sealed within microfluidic chambers. A syringe pump (PHD 2000, Harvard Apparatus) was employed to deliver gas mixtures into the microfluidic chamber at a constant flow of 3 ml/min. The rapid switch between different gas mixtures were controlled by Telfon valves coupled with ValveBank Perfusion Controller (AutoMate Scientific). The locomotory activity at different gas mixtures were monitored using a high-resolution camera (FLIR) mounted on a Zeiss Stemi 508 microscope. Videos were captured at a rate of 2 frames per second, starting with a 2-minute recording in 7% O$_2$, followed by 10 or 11 minutes in H$_2$S, and ending with 7% O$_2$. For the long-term recording in H$_2$S, video capture commenced with a 2-minute period in 7% O$_2$, followed by a duration of 148 minutes in 150 ppm H$_2$S, and concluded with a final 2-minute interval in 7% O$_2$. For each strain or condition, 3 to 4 replicates were performed.

Near-UV light experiments were conducted to assess whether animals remained responsive following a 30-minute incubation in 150 ppm H$_2$S balanced with 7% O$_2$. For each experiment, 6-minute videos were recorded. The first 2 minutes captured the baseline locomotion of the animals under white light, followed by 2 minutes of exposure to near-UV light (435 nm, 0.7 mW/mm^2), and concluded with 2 minutes of white light. 3 to 4 replicates were performed, with approximately 15 animals recorded per replicate.

The H$_2$S gradient experiment was performed in a PDMS chamber connected to a pump delivering gas at 0.5 ml/min. To establish a gradient ranging from 150 ppm to 0 ppm H$_2$S, we pumped 150 ppm H$_2$S balanced with 7% O$_2$ into one side and 7% O$_2$ into the opposite side. After 25 minutes, the number of animals in each of the 5 sections was counted (*Figure 1G*). Four replicates were performed, each using 30 day-one adult animals.

To assess the effects of rotenone on H_2S evoked locomotory response, day-one adult

animals were subjected to 10 μM rotenone for 10min, 30min, 1 hour and 2 hours on the drug

containing plates. Subsequently, the rotenone-treated animals were assayed on the standard

assay plates without the drug. To explore the impact of ferric ammonium citrate (FAC) (F5879,

Sigma-Aldrich) and 2,2′-Bipyridyl (BP) (D216305, Sigma-Aldrich) on behavioral response to

H_2S, L4 animals were exposed to 100 μM FAC or 5 mg/ml BP for 16 hours in the presence of

bacterial food. The FAC or BP treated animals were assayed on FAC or BP containing plates,

respectively.

Optogenetic experiments were performed as previously outlined (Zhao et al., 2022).

Transgenic L4 animals were exposed to 100 μM all-trans retinal (ATR) (R2500, Sigma) in the

dark for 16 hours. The ATR-fed animals were assayed with the illumination of 70 μW/mm^2

blue light during the whole assay period. Blue light was emitted from an ultra-high-power LED

lamp (UHP-MIC-LED-460, Prizmatix), and the light intensity was determined using a PM50

Optical Power Meter (ThorLabs). To minimize the effect of transmitted light on the transgenic

animals, a long-pass optical filter was used to eliminate the lights with short wavelengths during

picking. All videos of behavioral analysis were analyzed using a home-made MATLAB

program Zentracker (https://github.com/wormtracker/zentracker). For all behavior analyses, at

least three assays were performed for each strain with more than 80 worms in total.

Ca^{2+} Imaging

L4 animals expressing the GCaMP6s sensor in ASJ neurons were picked 24 hours before

imaging. The agarose pads were prepared 1 hour before the experiment. Animals were

incubated in 15 µl of 10 mM levamisole for 40 minutes prior to imaging. Between 15 and 20

animals were immobilized using 10 mM levamisole diluted in a concentrated OP50 bacterial

suspension and mounted on a 2% agarose pad in M9 buffer on a glass slide. The animals were

then covered with a microfluidic chamber. Imaging lasted for 16 minutes. O_2 (7%) was pumped

into the microfluidic chamber during the first and last 4 minutes. From minute 4 to 12, a mixture

of 7% O_2 and 150 ppm H_2S was pumped.

Imaging was performed on a Nikon AZ100 microscope equipped with a TwinCam

adapter (Cairn Research, UK), fitted with two DMK 33U monochrome cameras (The Imaging

Source, Germany). A 2x AZ Plan Fluor objective with 1x zoom was used, with an exposure

time of 300 ms. Excitation of GCaMP and RFP was provided by a CoolLED pE-300 (CoolLED,

UK). The TwinCam adapter was mounted with emission filters for green fluorescent protein

(510/20 nm) and red fluorescent protein (590/35 nm), along with a DC/T510LPXRXTUf2

dichroic mirror. Imaging data were analyzed using Neuron Analyzer, a custom-written Matlab

program for processing image stacks (code available at

https://github.com/neuronanalyser/neuronanalyser).

RNA extraction and sequencing

To obtain RNA samples for RNA-seq, thirty young adult animals were picked to each fresh

plate and allowed to lay eggs for 2 hours, after which the adult animals were removed and eggs

were allowed to develop into day-one adults. Day-one animals were subsequently challenged

with H_2S for three different time periods (1 hour, 2 hours and 12 hours). For each time period,

five plates of day-one animals were exposed to 50 ppm H_2S, 150 ppm H_2S or 7% O_2 as a control

group. Subsequently, the animals were immediately collected and rinsed three times with M9

buffer. After washing, the worm pellet was frozen in liquid nitrogen. Animals were

homogenized using Bullet Blender (Next Advance) in the presence of Qiazol Lysis Reagent

and 0.5mm Zirconia beads at 4°C. RNA was prepared using RNeasy Plus Universal Mini Kit

(Qiagen). Six independent RNA samples were prepared for each condition. The 2100

Bioanalyzer instrument (Agilent) was used for the RNA quality control with a microfluidic chip

specific for RNA (Agilent RNA 6000 Nano Kit). Then 1 μg of qualified RNA samples in

RNase-free ddH2O was used for library preparation. Library was constructed and sequenced

by Novogene.

RNA-seq analysis

The RNA-seq data were aligned using STAR v2.7.9a (Dobin et al., 2013) and gene expression

counts extracted by featureCounts v2.0.3 (Liao, Smyth, & Shi, 2014). Differential expression

was calculated using DESeq2 (Love, Huber, & Anders, 2014). GO analysis was performed

using the EnrichR R package (Kuleshov et al., 2016) on genes having p.adj < 1e-10. For brevity,

several highly similar GO categories were manually omitted (full EnrichR output available at

Github repository).

Data availability

Raw sequencing data has been deposited at ArrayExpress # E-MTAB-13296. The R code is

available at https://github.com/henriksson-lab/ce_h2s_tc.

Acknowledgments

We thank the Caenorhabditis Genetics Center (funded by NIH Office of Research Infrastructure

Programs P40 OD010440) and the National BioResources Project Japan for strains.

Funding

The computations were enabled by resources provided by the National Academic Infrastructure

for Supercomputing in Sweden (NAISS) and the Swedish National Infrastructure for

Computing (SNIC) at UPPMAX partially funded by the Swedish Research Council through

grant agreements no. 2022-06725 and no. 2018-05973. This work is supported by the Swedish

VR Research Council grant MIMS (2021-06602) to J.H., the Belgian National Fund for

Scientific Research (FRS-FNRS) to P.L., the ERC starting grant (802653 OXYGEN

SENSING), the Swedish Research Council VR starting grant (2018-02216), and the

Wallenberg Centre for Molecular Medicine (Umeå) to C.C.

References

Abel, D. C., Koenig, C. C., & Davis, W. P. (1987). Emersion in the Mangrove Forest Fish

Rivulus-Marmoratus - a Unique Response to Hydrogen-Sulfide. *Environmental*

Biology of Fishes, 18(1), 67-72. doi:Doi 10.1007/Bf00002329

Ackerman, D., & Gems, D. (2012). Insulin/IGF-1 and Hypoxia Signaling Act in Concert to

Regulate Iron Homeostasis in *Caenorhabditis elegans*. *Plos Genetics, 8*(3).

doi:ARTN e1002498

10.1371/journal.pgen.1002498

Adams, D. F., Farwell, S. O., Pack, M. R., & Bamesberger, W. L. (1979). Preliminary

Measurements of Biogenic Sulfur-Containing Gas Emissions from Soils. *Journal of*

the Air Pollution Control Association, 29(4), 380-383. doi:Doi

10.1080/00022470.1979.10470805

Anderson, C. P., & Leibold, E. A. (2014). Mechanisms of iron metabolism in *Caenorhabditis*

elegans. *Frontiers in Pharmacology, 5*. doi:ARTN 113

10.3389/fphar.2014.00113

Beets, I., Zhang, G., Fenk, L. A., Chen, C., Nelson, G. M., Felix, M. A., & de Bono, M.

(2020). Natural Variation in a Dendritic Scaffold Protein Remodels Experience-

Dependent Plasticity by Altering Neuropeptide Expression. *Neuron, 105*(1), 106-121

e110. doi:10.1016/j.neuron.2019.10.001

Brenner, S. (1974). The genetics of *Caenorhabditis elegans*. *Genetics, 77*(1), 71-94.

Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/4366476

Bretscher, A. J., Busch, K. E., & de Bono, M. (2008). A carbon dioxide avoidance behavior is

integrated with responses to ambient oxygen and food in *Caenorhabditis elegans*.

Proc Natl Acad Sci U S A, 105(23), 8044-8049. doi:10.1073/pnas.0707607105

Budde, M. W., & Roth, M. B. (2010). Hydrogen sulfide increases hypoxia-inducible factor-1

activity independently of von Hippel-Lindau tumor suppressor-1 in *C. elegans*. *Mol*

Biol Cell, 21(1), 212-217. doi:10.1091/mbc.E09-03-0199

Budde, M. W., & Roth, M. B. (2011). The response of *Caenorhabditis elegans* to hydrogen

sulfide and hydrogen cyanide. *Genetics, 189*(2), 521-532.

doi:10.1534/genetics.111.129841

Busch, K. E., Laurent, P., Soltesz, Z., Murphy, R. J., Faivre, O., Hedwig, B., . . . de Bono, M.

(2012). Tonic signaling from O(2) sensors sets neural circuit activity and behavioral

state. *Nat Neurosci, 15*(4), 581-591. doi:10.1038/nn.3061

Byrne Rodgers, J., & Ryu, W. S. (2020). Targeted thermal stimulation and high-content

phenotyping reveal that the *C. elegans* escape response integrates current behavioral

state and past experience. *PLoS One, 15*(3), e0229399.

doi:10.1371/journal.pone.0229399

Carrillo, M. A., Guillermin, M. L., Rengarajan, S., Okubo, R. P., & Hallem, E. A. (2013). O-

2-Sensing Neurons Control CO_2 Response in *C. elegans*. *Journal of Neuroscience,*

33(23), 9675-9683. doi:10.1523/Jneurosci.4541-12.2013

Chang, A. J., Chronis, N., Karow, D. S., Marletta, M. A., & Bargmann, C. I. (2006). A

distributed chemosensory circuit for oxygen preference in *C. elegans*. *PLoS Biol,*

4(9), e274. doi:10.1371/journal.pbio.0040274

Cheung, B. H., Cohen, M., Rogers, C., Albayram, O., & de Bono, M. (2005). Experience-

dependent modulation of *C. elegans* behavior by ambient oxygen. *Curr Biol, 15*(10),

905-917. doi:10.1016/j.cub.2005.04.017

Cooper, C. E., & Brown, G. C. (2008). The inhibition of mitochondrial cytochrome oxidase

by the gases carbon monoxide, nitric oxide, hydrogen cyanide and hydrogen sulfide:

chemical mechanism and physiological significance. *J Bioenerg Biomembr, 40*(5),

533-539. doi:10.1007/s10863-008-9166-6

Couto, A., Oda, S., Nikolaev, V. O., Soltesz, Z., & de Bono, M. (2013). In vivo genetic

dissection of O_2-evoked cGMP dynamics in a *Caenorhabditis elegans* gas sensor.

Proc Natl Acad Sci U S A, 110(35), E3301-3310. doi:10.1073/pnas.1217428110

de Bono, M., & Bargmann, C. I. (1998). Natural variation in a neuropeptide Y receptor

homolog modifies social behavior and food response in *C. elegans*. *Cell, 94*(5), 679-

689. doi:10.1016/s0092-8674(00)81609-8

Dekker, L. V., Mcintyre, P., & Parker, P. J. (1993). Mutagenesis of the Regulatory Domain of

Rat Protein Kinase-C-Eta - a Molecular-Basis for Restricted Histone Kinase-Activity.

Journal of Biological Chemistry, 268(26), 19498-19504. Retrieved from <Go to

ISI>://WOS:A1993LW81900055

Dev, S., & Babitt, J. L. (2017). Overview of iron metabolism in health and disease. *Hemodial

Int, 21 Suppl 1*(Suppl 1), S6-S20. doi:10.1111/hdi.12542

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., . . . Gingeras, T. R.

(2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics, 29*(1), 15-21.

doi:10.1093/bioinformatics/bts635

Dokshin, G. A., Ghanta, K. S., Piscopo, K. M., & Mello, C. C. (2018). Robust Genome

Editing with Short Single-Stranded and Long, Partially Single-Stranded DNA Donors

in *Caenorhabditis elegans*. *Genetics, 210*(3), 781-787.

doi:10.1534/genetics.118.301532

Doonan, R., McElwee, J. J., Matthijssens, F., Walker, G. A., Houthoofd, K., Back, P., . . .

Gems, D. (2008). Against the oxidative damage theory of aging: superoxide

dismutases protect against oxidative stress but have little or no effect on life span in

Caenorhabditis elegans. *Genes Dev, 22*(23), 3236-3241. doi:10.1101/gad.504808

Fawcett, E. M., Hoyt, J. M., Johnson, J. K., & Miller, D. L. (2015). Hypoxia disrupts

proteostasis in *Caenorhabditis elegans*. *Aging Cell, 14*(1), 92-101.

doi:10.1111/acel.12301

Ferkey, D. M., Sengupta, P., & L'Etoile, N. D. (2021). Chemosensory signal transduction in

Caenorhabditis elegans. *Genetics, 217*(3). doi:10.1093/genetics/iyab004

Gao, L., Gonzalez-Rodriguez, P., Ortega-Saenz, P., & Lopez-Barneo, J. (2017). Redox

signaling in acute oxygen sensing. *Redox Biol, 12*, 908-915.

doi:10.1016/j.redox.2017.04.033

Gea, T., Barrena, R., Artola, A., & Sanchez, A. (2004). Monitoring the biological activity of

the composting process: Oxygen uptake rate (OUR), respirometric index (RI), and

respiratory quotient (RQ). *Biotechnol Bioeng, 88*(4), 520-527. doi:10.1002/bit.20281

Ghanta, K. S., & Mello, C. C. (2020). Melting dsDNA Donor Molecules Greatly Improves

Precision Genome Editing in *Caenorhabditis elegans*. *Genetics, 216*(3), 643-650.

doi:10.1534/genetics.120.303564

Gourley, B. L., Parker, S. B., Jones, B. J., Zumbrennen, K. B., & Leibold, E. A. (2003).

Cytosolic aconitase and ferritin are regulated by iron in *Caenorhabditis elegans*. *J

Biol Chem, 278*(5), 3227-3234. doi:10.1074/jbc.M210333200

Gray, J. M., Karow, D. S., Lu, H., Chang, A. J., Chang, J. S., Ellis, R. E., . . . Bargmann, C. I.

(2004). Oxygen sensation and social feeding mediated by a *C. elegans* guanylate

cyclase homologue. *Nature, 430*(6997), 317-322. doi:10.1038/nature02714

Hallem, E. A., Spencer, W. C., McWhirter, R. D., Zeller, G., Henz, S. R., Ratsch, G., . . .

Ringstad, N. (2011). Receptor-type guanylate cyclase is required for carbon dioxide

sensation by *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A, 108*(1), 254-259.

doi:10.1073/pnas.1017354108

Hallem, E. A., & Sternberg, P. W. (2008). Acute carbon dioxide avoidance in *Caenorhabditis

elegans*. *Proc Natl Acad Sci U S A, 105*(23), 8038-8043.

doi:10.1073/pnas.0707469105

Heiman, M. G., & Shaham, S. (2009). DEX-1 and DYF-7 establish sensory dendrite length

by anchoring dendritic tips during cell migration. *Cell, 137*(2), 344-355.

doi:10.1016/j.cell.2009.01.057

Hentze, M. W., Muckenthaler, M. U., & Andrews, N. C. (2004). Balancing acts: molecular

control of mammalian iron metabolism. *Cell, 117*(3), 285-297. doi:10.1016/s0092-

8674(04)00343-5

Hildebrandt, T. M., & Grieshaber, M. K. (2008). Three enzymatic activities catalyze the

oxidation of sulfide to thiosulfate in mammalian and invertebrate mitochondria.

FEBS J, 275(13), 3352-3361. doi:10.1111/j.1742-4658.2008.06482.x

Hiroki, S., Yoshitane, H., Mitsui, H., Sato, H., Umatani, C., Kanda, S., . . . Iino, Y. (2022).

Molecular encoding and synaptic decoding of context during salt chemotaxis in *C.*

elegans. *Nature Communications, 13*(1). doi:ARTN 2928

10.1038/s41467-022-30279-7

Horsman, J. W., Heinis, F. I., & Miller, D. L. (2019). A Novel Mechanism To Prevent H2S

Toxicity in *Caenorhabditis elegans*. *Genetics, 213*(2), 481-490.

doi:10.1534/genetics.119.302326

Ishii, T., Miyazawa, M., Onouchi, H., Yasuda, K., Hartman, P. S., & Ishii, N. (2013). Model

animals for the study of oxidative stress from complex II. *Biochim Biophys Acta,*

1827(5), 588-597. doi:10.1016/j.bbabio.2012.10.016

Jackson, M. R., Melideo, S. L., & Jorns, M. S. (2012). Human sulfide:quinone oxidoreductase

catalyzes the first step in hydrogen sulfide metabolism and produces a sulfane sulfur

metabolite. *Biochemistry, 51*(34), 6804-6815. doi:10.1021/bi300778t

Jia, J., Wang, Z., Zhang, M., Huang, C., Song, Y., Xu, F., . . . Cheng, J. (2020). SQR

mediates therapeutic effects of H(2)S by targeting mitochondrial electron transport to

induce mitochondrial uncoupling. *Sci Adv, 6*(35), eaaz5752.

doi:10.1126/sciadv.aaz5752

Kabil, O., & Banerjee, R. (2012). Characterization of patient mutations in human persulfide

dioxygenase (ETHE1) involved in H_2S catabolism. *J Biol Chem, 287*(53), 44561-

44567. doi:10.1074/jbc.M112.407411

Kaelin, W. G., Jr., & Ratcliffe, P. J. (2008). Oxygen sensing by metazoans: the central role of

the HIF hydroxylase pathway. *Mol Cell, 30*(4), 393-402.

doi:10.1016/j.molcel.2008.04.009

Kelley, J. L., Arias-Rodriguez, L., Patacsil Martin, D., Yee, M. C., Bustamante, C. D., &

Tobler, M. (2016). Mechanisms Underlying Adaptation to Life in Hydrogen Sulfide-

Rich Environments. *Mol Biol Evol, 33*(6), 1419-1434. doi:10.1093/molbev/msw020

Khan, A. A., Schuler, M. M., Prior, M. G., Yong, S., Coppock, R. W., Florence, L. Z., &

Lillie, L. E. (1990). Effects of hydrogen sulfide exposure on lung mitochondrial

respiratory chain enzymes in rats. *Toxicol Appl Pharmacol, 103*(3), 482-490.

doi:10.1016/0041-008x(90)90321-k

Kodama-Namba, E., Fenk, L. A., Bretscher, A. J., Gross, E., Busch, K. E., & de Bono, M.

(2013). Cross-modulation of homeostatic responses to temperature, oxygen and

carbon dioxide in *C. elegans*. *PLoS Genet, 9*(12), e1004011.

doi:10.1371/journal.pgen.1004011

Kuleshov, M. V., Jones, M. R., Rouillard, A. D., Fernandez, N. F., Duan, Q. N., Wang, Z. C.,

. . . Ma'ayan, A. (2016). Enrichr: a comprehensive gene set enrichment analysis web

server 2016 update. *Nucleic Acids Research, 44*(W1), W90-W97.

doi:10.1093/nar/gkw377

Laurent, P., Soltesz, Z., Nelson, G. M., Chen, C., Arellano-Carbajal, F., Levy, E., & de Bono,

M. (2015). Decoding a neural circuit controlling global animal state in *C. elegans*.

Elife, 4. doi:10.7554/eLife.04241

Lennicke, C., & Cocheme, H. M. (2021). Redox metabolism: ROS as specific molecular

regulators of cell signaling and function. *Mol Cell, 81*(18), 3691-3707.

doi:10.1016/j.molcel.2021.08.018

Liao, Y., Smyth, G. K., & Shi, W. (2014). featureCounts: an efficient general purpose

program for assigning sequence reads to genomic features. *Bioinformatics, 30*(7),

923-930. doi:10.1093/bioinformatics/btt656

Liochev, S. L. (1996). The role of iron-sulfur clusters in in vivo hydroxyl radical production.

Free Radical Research, 25(5), 369-384. doi:Doi 10.3109/10715769609149059

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biology, 15*(12). doi:ARTN 550

10.1186/s13059-014-0550-8

Ma, D. K., Vozdek, R., Bhatla, N., & Horvitz, H. R. (2012). CYSL-1 interacts with the O_2-

sensing hydroxylase EGL-9 to promote H_2S-modulated hypoxia-induced behavioral

plasticity in *C. elegans*. *Neuron, 73*(5), 925-940. doi:10.1016/j.neuron.2011.12.037

Macosko, E. Z., Pokala, N., Feinberg, E. H., Chalasani, S. H., Butcher, R. A., Clardy, J., &

Bargmann, C. I. (2009). A hub-and-spoke circuit drives pheromone attraction and

social behaviour in *C. elegans*. *Nature, 458*(7242), 1171-1175.

doi:10.1038/nature07886

Mathew, N. D., Schlipalius, D. I., & Ebert, P. R. (2011). Sulfurous gases as biological

messengers and toxins: comparative genetics of their metabolism in model

organisms. *J Toxicol, 2011*, 394970. doi:10.1155/2011/394970

McGrath, P. T., Rockman, M. V., Zimmer, M., Jang, H., Macosko, E. Z., Kruglyak, L., &

Bargmann, C. I. (2009). Quantitative mapping of a digenic behavioral trait implicates

globin variation in *C. elegans* sensory behaviors. *Neuron, 61*(5), 692-699.

doi:10.1016/j.neuron.2009.02.012

Miller, D. L., Budde, M. W., & Roth, M. B. (2011a). HIF-1 and SKN-1 Coordinate the

Transcriptional Response to Hydrogen Sulfide in *Caenorhabditis elegans*. *PLoS One,*

6(9). doi:ARTN e25476

10.1371/journal.pone.0025476

Miller, D. L., Budde, M. W., & Roth, M. B. (2011b). HIF-1 and SKN-1 coordinate the

transcriptional response to hydrogen sulfide in *Caenorhabditis elegans*. *PLoS One,*

6(9), e25476. doi:10.1371/journal.pone.0025476

Miller, D. L., & Roth, M. B. (2007). Hydrogen sulfide increases thermotolerance and lifespan

in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A, 104*(51), 20618-20622.

doi:10.1073/pnas.0710191104

Morra, M. J., & Dick, W. A. (1991). Mechanisms of h(2)s production from cysteine and

cystine by microorganisms isolated from soil by selective enrichment. *Appl Environ*

Microbiol, 57(5), 1413-1417. doi:10.1128/aem.57.5.1413-1417.1991

Murakami, M., Koga, M., & Ohshima, Y. (2001). DAF-7/TGF-beta expression required for

the normal larval development in *C-elegans* is controlled by a presumed guanylyl

cyclase DAF-11. *Mechanisms of Development, 109*(1), 27-35. doi:Doi

10.1016/S0925-4773(01)00507-X

Murayama, T., Toh, Y., Ohshima, Y., & Koga, M. (2005). The dyf-3 gene encodes a novel

protein required for sensory cilium formation in *Caenorhabditis elegans*. *J Mol Biol,*

346(3), 677-687. doi:10.1016/j.jmb.2004.12.005

Myllyharju, J., & Kivirikko, K. I. (1997). Characterization of the iron- and 2-oxoglutarate-

binding sites of human prolyl 4-hydroxylase. *Embo Journal, 16*(6), 1173-1180.

doi:DOI 10.1093/emboj/16.6.1173

Nicholls, P., & Kim, J. K. (1982). Sulphide as an inhibitor and electron donor for the

cytochrome c oxidase system. *Can J Biochem, 60*(6), 613-623. doi:10.1139/o82-076

Niu, W., Lu, Z. J., Zhong, M., Sarov, M., Murray, J. I., Brdlik, C. M., . . . Reinke, V. (2011).

Diverse transcription factor binding features revealed by genome-wide ChIP-seq in

C. elegans. *Genome Research, 21*(2), 245-254. doi:10.1101/gr.114587.110

Ochi, R., Dhagia, V., Lakhkar, A., Patel, D., Wolin, M. S., & Gupte, S. A. (2016). Rotenone-

stimulated superoxide release from mitochondrial complex I acutely augments L-type

Ca^{2+} current in A7r5 aortic smooth muscle cells. *Am J Physiol Heart Circ Physiol,*

310(9), H1118-1128. doi:10.1152/ajpheart.00889.2015

Olson, K. R., & Straub, K. D. (2016). The Role of Hydrogen Sulfide in Evolution and the

Evolution of Hydrogen Sulfide in Metabolism and Signaling. *Physiology (Bethesda),*

31(1), 60-72. doi:10.1152/physiol.00024.2015

Onukwufor, J. O., Farooqi, M. A., Vodickova, A., Koren, S. A., Baldzizhar, A., Berry, B. J., .

. . Wojtovich, A. P. (2022). A reversible mitochondrial complex I thiol switch

mediates hypoxic avoidance behavior in *C. elegans*. *Nat Commun, 13*(1), 2403.

doi:10.1038/s41467-022-30169-y

Oshins, C., Michel, F., Louis, P., Richard, T. L., & Rynk, R. (2022). Chapter 3 - The

composting process. In R. Rynk (Ed.), *The Composting Handbook* (pp. 51-101):

Academic Press.

Patange, O., Breen, P., Arsuffi, G., & Ruvkun, G. (2025). Hydrogen sulfide mediates the

interaction between *C. elegans* and Actinobacteria from its natural microbial

environment. *Cell Reports, 44*(1), 115170. doi:10.1016/j.celrep.2024.115170

Patten, D. A., Germain, M., Kelly, M. A., & Slack, R. S. (2010). Reactive oxygen species:

stuck in the middle of neurodegeneration. *J Alzheimers Dis, 20 Suppl 2*, S357-367.

doi:10.3233/JAD-2010-100498

Persson, A., Gross, E., Laurent, P., Busch, K. E., Bretes, H., & de Bono, M. (2009). Natural

variation in a neural globin tunes oxygen sensing in wild *Caenorhabditis elegans*.

Nature, 458(7241), 1030-1033. doi:10.1038/nature07820

Pettinati, I., Brem, J., McDonough, M. A., & Schofield, C. J. (2015). Crystal structure of

human persulfide dioxygenase: structural basis of ethylmalonic encephalopathy. *Hum

Mol Genet, 24*(9), 2458-2469. doi:10.1093/hmg/ddv007

Pfenninger, M., Lerp, H., Tobler, M., Passow, C., Kelley, J. L., Funke, E., . . . Plath, M.

(2014). Parallel evolution of cox genes in H_2S-tolerant fish as key adaptation to a

toxic environment. *Nat Commun, 5*, 3873. doi:10.1038/ncomms4873

Philipp, T. M., Gong, W., Kohnlein, K., Ohse, V. A., Muller, F. I., Priebs, J., . . . Klotz, L. O.

(2022). SEMO-1, a novel methanethiol oxidase in *Caenorhabditis elegans*, is a pro-

aging factor conferring selective stress resistance. *Biofactors, 48*(3), 699-706.

doi:10.1002/biof.1836

Powell-Coffman, J. A. (2010). Hypoxia signaling and resistance in *C. elegans*. *Trends

Endocrinol Metab, 21*(7), 435-440. doi:10.1016/j.tem.2010.02.006

Qabazard, B., Li, L., Gruber, J., Peh, M. T., Ng, L. F., Kumar, S. D., . . . Moore, P. K. (2014).

Hydrogen sulfide is an endogenous regulator of aging in *Caenorhabditis elegans*.

Antioxid Redox Signal, 20(16), 2621-2630. doi:10.1089/ars.2013.5448

Rajan, M., Anderson, C. P., Rindler, P. M., Romney, S. J., Ferreira Dos Santos, M. C., Gertz,

J., & Leibold, E. A. (2019). NHR-14 loss of function couples intestinal iron uptake

with innate immunity in *C. elegans* through PQM-1 signaling. *Elife, 8*.

doi:10.7554/eLife.44674

Ramsay, R. R., & Singer, T. P. (1992). Relation of superoxide generation and lipid

peroxidation to the inhibition of NADH-Q oxidoreductase by rotenone, piericidin A,

and MPP+. *Biochem Biophys Res Commun, 189*(1), 47-52. doi:10.1016/0006-

291x(92)91523-s

Read, A. D., Bentley, R. E. T., Archer, S. L., & Dunham-Snary, K. J. (2021). Mitochondrial

iron-sulfur clusters: Structure, function, and an emerging role in vascular biology.

Redox Biology, 47. doi:ARTN 102164

10.1016/j.redox.2021.102164

Rodriguez-Kabana, R., Jordan, J. W., & Hollis, J. P. (1965). Nematodes: Biological Control

in Rice Fields: Role of Hydrogen Sulfide. *Science, 148*(3669), 524-526.

doi:10.1126/science.148.3669.524

Romanelli-Cedrez, L., Vairoletti, F., & Salinas, G. (2024). Rhodoquinone-dependent electron

transport chain is essential for *Caenorhabditis elegans* survival in hydrogen sulfide

environments. *J Biol Chem, 300*(9), 107708. doi:10.1016/j.jbc.2024.107708

Romney, S. J., Newman, B. S., Thacker, C., & Leibold, E. A. (2011). HIF-1 regulates iron

homeostasis in *Caenorhabditis elegans* by activation and inhibition of genes involved

in iron uptake and storage. *PLoS Genet, 7*(12), e1002394.

doi:10.1371/journal.pgen.1002394

Salvanes, A. G. V., Utne-Palm, A. C., Currie, B., & Braithwaite, V. A. (2011). Behavioural

and physiological adaptations of the bearded goby, a key fish species of the extreme

environment of the northern Benguela upwelling. *Marine Ecology Progress Series,*

425, 193-202. doi:10.3354/meps08998

Schiavo, G., Benfenati, F., Poulain, B., Rossetto, O., Polverino de Laureto, P., DasGupta, B.

R., & Montecucco, C. (1992). Tetanus and botulinum-B neurotoxins block

neurotransmitter release by proteolytic cleavage of synaptobrevin. *Nature, 359*(6398),

832-835. doi:10.1038/359832a0

Semenza, G. L. (2010). Defining the role of hypoxia-inducible factor 1 in cancer biology and

therapeutics. *Oncogene, 29*(5), 625-634. doi:10.1038/onc.2009.441

Starich, T. A., Herman, R. K., Kari, C. K., Yeh, W. H., Schackwitz, W. S., Schuyler, M. W., .

. . Riddle, D. L. (1995). Mutations affecting the chemosensory neurons of

Caenorhabditis elegans. *Genetics, 139*(1), 171-188. doi:10.1093/genetics/139.1.171

Sterken, M. G., Snoek, L. B., Kammenga, J. E., & Andersen, E. C. (2015). The laboratory

domestication of *Caenorhabditis elegans*. *Trends Genet, 31*(5), 224-231.

doi:10.1016/j.tig.2015.02.009

Swoboda, P., Adler, H. T., & Thomas, J. H. (2000). The RFX-type transcription factor DAF-

19 regulates sensory neuron cilium formation in *C. elegans*. *Mol Cell, 5*(3), 411-421.

doi:10.1016/s1097-2765(00)80436-0

Szabo, C., Ransy, C., Módis, K., Andriamihaja, M., Murghes, B., Coletta, C., . . . Bouillaud,

F. (2014). Regulation of mitochondrial bioenergetic function by hydrogen sulfide.

Part I. Biochemical and physiological mechanisms. *British Journal of Pharmacology,*

171(8), 2099-2122. doi:10.1111/bph.12369

Vora, M., Pyonteck, S. M., Popovitchenko, T., Matlack, T. L., Prashar, A., Kane, N. S., . . .

Rongo, C. (2022). The hypoxia response pathway promotes PEP carboxykinase and

gluconeogenesis in *C. elegans*. *Nature Communications, 13*(1). doi:ARTN 6168

10.1038/s41467-022-33849-x

Weir, E. K., Lopez-Barneo, J., Buckler, K. J., & Archer, S. L. (2005). Acute oxygen-sensing

mechanisms. *N Engl J Med, 353*(19), 2042-2055. doi:10.1056/NEJMra050002

Xu, X. M., & Moller, S. G. (2011). Iron-Sulfur Clusters: Biogenesis, Molecular Mechanisms,

and Their Functional Significance. *Antioxidants & Redox Signaling, 15*(1), 271-307.

doi:10.1089/ars.2010.3259

Yang, W., & Hekimi, S. (2010). A mitochondrial superoxide signal triggers increased

longevity in *Caenorhabditis elegans*. *PLoS Biol, 8*(12), e1000556.

doi:10.1371/journal.pbio.1000556

Zhao, L., Fenk, L. A., Nilsson, L., Amin-Wetzel, N. P., Ramirez-Suarez, N. J., de Bono, M.,

& Chen, C. (2022). ROS and cGMP signaling modulate persistent escape from

hypoxia in *Caenorhabditis elegans*. *PLoS Biol, 20*(6), e3001684.

doi:10.1371/journal.pbio.3001684

Zimmer, M., Gray, J. M., Pokala, N., Chang, A. J., Karow, D. S., Marletta, M. A., . . .

Bargmann, C. I. (2009). Neurons detect increases and decreases in oxygen levels

using distinct guanylate cyclases. *Neuron, 61*(6), 865-879.

doi:10.1016/j.neuron.2009.02.013

Zorov, D. B., Juhaszova, M., & Sollott, S. J. (2014). Mitochondrial reactive oxygen species

(ROS) and ROS-induced ROS release. *Physiol Rev, 94*(3), 909-950.

doi:10.1152/physrev.00026.2013

Zubovych, I. O., Straud, S., & Roth, M. G. (2010). Mitochondrial dysfunction confers

resistance to multiple drugs in *Caenorhabditis elegans*. *Mol Biol Cell, 21*(6), 956-

969. doi:10.1091/mbc.e09-08-0673

1014

Figure legends

Figure 1. Acute locomotory responses of *C. elegans* to hydrogen sulfide

(**A**) Changes in the locomotory speed of WT animals (N2 laboratory strain) evoked by a switch

from 7% O_2 to 150 ppm H_2S balanced with 7% O_2, followed by a return to 7% O_2. (**B**)

Proportion of WT animals undergoing reorientation movements (omega turns) during a switch

from 7% O_2 to 150 ppm H_2S balanced with 7% O_2, followed by a return to 7% O_2. (**C**) Fraction

of WT animals doing backward locomotion (reversals) during a switch from 7% O_2 to 150 ppm

H_2S balanced with 7% O_2, followed by a return to 7% O_2. (**D**) Locomotory speed of WT was

recorded for 2 minutes at 7% O_2, followed by 148 minutes at 150 ppm H_2S balanced with 7%

O_2, and then at 7% O_2. (**E**) Locomotory speed of WT animals in response to near-UV light (435

nm, 0.7 mW/mm^2) exposure, before and directly after 30-minute preincubation in 150 ppm H_2S

balanced with 7% O_2. Locomotory activity was recorded for 2 minutes before, during and after

UV light exposure. (**F**) Locomotory speed of WT animals to different concentrations of H_2S

balanced with 7% O_2. Red bars on the x-axis represent two intervals (4–5 minutes and 11–12

minutes, labeled a and b, respectively) used for statistical analysis. **** = $p < 0.0001$, ** = $p <$

0.01, * = $p < 0.05$, ns = not significant, Mann–Whitney U test. (**G**) Distribution of WT animals

in a microfluidic device after 25 minutes of aerotaxis. The gas inputs and the five chamber

sections used for scoring are indicated. (**H**) Distribution of WT animals in a gradient of H_2S

(150 ppm to 0 ppm). The bins correspond to different sections of the microfluidic chamber

described in (**G**). N= 4 aerotaxis assays.

1035

Figure 2. Neurosecretion from ASJ neurons contributes to H_2S avoidance

(**A–H**) Locomotory speed responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7%

O_2 for animals of the indicated genotype: WT and *daf-19(m86)* (**A**); WT and *dyf-3(m185)* (**B**);

WT and *dyf-7(m539)* (**C**); WT, *daf-11(m47)* and *tax-4(p678)* (**D**); WT and *tax-2(p691)* (**E**);

WT, *daf-11(m47)* and transgenic *daf-11(m47)* expressing *daf-11* genomic DNA in ASJ neurons

(two independent lines) (**F**); WT, *tax-4(p678)* and transgenic *tax-4(p678)* expressing *tax-4*

cDNA in ASJ neurons (two independent lines) (**G**); WT and transgenic WT expressing the

catalytic domain of the *tetanus* toxin (TeTx) in ASJ neurons (**H**). For the comparison of acute

locomotory speed responses between strains, red bars on the x-axis represent two intervals (4–

5 minutes and 11–12 minutes, labeled a and b, respectively, in panels D, F, and G) used for

statistical analysis. **** = $p < 0.0001$, *** = $p < 0.001$, ** = $p < 0.01$, * = $p < 0.05$, ns = not

significant, Mann–Whitney U test.

1048

Figure 3. Insulin, TGF-β, and high O_2 signaling pathways antagonizes H_2S avoidance

(**A, B**) Locomotory speed responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7%

O_2 for animals of the indicated genotype: WT, *daf-2(e1370), daf-16(mgDf47)*, and *daf-2(e1370);*

daf-16(mgDf47) double mutants (**A**); WT, *daf-7(e1372), daf-3(mgDf90),* and *daf-7(e1372);*

daf-3(mgDf90) double mutants (**B**). For the *daf-2* assays, WT and *daf-2* mutants were

maintained at 15°C. L4 animals were picked and shifted to 25°C until day-one adults, then

assayed at room temperature. (**C**) Hypothetical model for the regulation of H_2S avoidance by a

starvation program involving the insulin, TGF-β, and NHR-49 pathways. (**D**) Locomotory

speed response to a switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2 in fed and starved

1058 WT animals. (**E, F**) Locomotory speed responses to a switch from 7% O_2 to 150 ppm H_2S

1059 balanced with 7% O_2 for animals of the indicated genotype: WT and wild isolates (CB4858 and

1060 CB4855) (**E**); WT and *npr-1(ad609)* (**F**). (**G**) Hypothetical model showing how O_2 regulates

1061 H_2S avoidance via the inhibitory RMG circuit activity, which is modulated by O_2 sensory inputs

1062 and NPR-1 signaling. (**H–L**) Locomotory speed responses to a switch from 7% O_2 to 150 ppm

1063 H_2S balanced with 7% O_2 for animals of the indicated genotype: WT, *npr-1(ad609)*, and

1064 transgenic *npr-1(ad609)* expressing *npr-1* in RMG neurons using Cre-LoxP system (Macosko

1065 et al., 2009) (**H**); WT, *npr-1(ad609), gcy-35(ok769)*, and *npr-1(ad609); gcy-35(ok769)* double

1066 mutants (**I**); WT, *npr-1(ad609), tax-4(p678)* and *npr-1(ad609); tax-4(p678)* double mutants (**J**);

1067 WT, *npr-1(ad609), qaIs2241*(genetic ablation of AQR, PQR and URX neurons) and *npr-*

1068 *1(ad609); qaIs 2241* (**K**); WT, *npr-1(ad609), npr-1(ad609); gcy-35(ok769)* double mutants and

1069 *transgenic npr-1(ad609); gcy-35(ok769)* double mutants expressing *gcy-35* cDNA under *gcy-*

1070 *37* promoter, which drives *gcy-35* expression in O_2 sensing neurons (**L**). For the comparison of

1071 acute locomotory responses between strains, red bars on the x-axis represent two intervals (4–

1072 5 minutes and 11–12 minutes, labeled a and b, respectively, in panel E) used for statistical

1073 analysis. **** = $p < 0.0001$, *** = $p < 0.001$, ** = $p < 0.01$, * = $p < 0.05$, ns = not significant,

1074 Mann–Whitney U test.

1075

1076 **Figure 4. Prolonged H_2S exposure reprograms gene expression in *C. elegans***

1077 (**A**) Venn diagram displaying the number of differentially expressed genes after 1-hour

1078 exposure to 50 ppm or 150 ppm H_2S balanced with 7% O_2 in WT animals (adjusted $p < 1e-10$).

1079 (**B**) Venn diagram displaying the number of differentially expressed genes after 1-, 2-, and 12-

1080 hour exposure to 150 ppm H_2S balanced with 7% O_2 in WT animals (adjusted $p < 1e-10$). (**C,**

1081 **D**) Significantly enriched GO categories for differentially expressed genes with adjusted $p <$

1082 1e-10 in WT animals exposed to 50 ppm (**C**) or 150 ppm (**D**) H_2S balanced with 7% O_2 for 1

1083 hour. (**E, F**) Volcano plots showing the differentially expressed genes with adjusted $p < 1e-10$

1084 in WT animals exposed to 50 ppm (**E**) or 150 ppm (**F**) H_2S balanced with 7% O_2 for 1 hour. A

1085   set of genes involved in H_2S detoxification, cysteine metabolism and stress response were

1086   highlighted in red.

1087

Figure 5. Acute response to H_2S is modulated by HIF-1 signaling

(**A**) Locomotory speed responses of WT animals to a switch from 7% O_2 to 150 ppm H_2S

balanced with 7% O_2 after incubation of the animals for 0, 12, 24, or 36 hours in 1% O_2. (**B, C**)

Locomotory speed responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2 for

animals of the indicated genotype: WT, *egl-9(sa307)*, and *vhl-1(ok161)* (**B**); WT and transgenic

animals expressing the non-degradable form of HIF-1 (P621A or P621G) under the pan-

neuronal *unc-14* promoter or the endogenous *hif-1* promoter, respectively (**C**). (**D–F**)

Avoidance responses of WT and *hif-1(ia4)* to a switch from 7% O_2 to 150 ppm H_2S balanced

with 7% O_2, including locomotory speed (**D**), reorientation (omega-turn) (**E**), and reversal (**F**).

(**G, H**) Locomotory speed responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7%

O_2 for animals of the indicated genotype: WT, *egl-9(sa307), hif-1(ia4)*, and *egl-9(sa307); hif-*

1(ia4) double mutants (**G**); WT, *vhl-1(ok161), hif-1(ia4)*, and *vhl-1(ok161); hif-1(ia4)* double

mutants (**H**). (**I**) Average locomotory speed during 2 minutes of exposure to near-UV light for

WT, *hif-1(ia4)*, and *egl-9(sa307)* before (Ctrl = control) and after 30 minutes of exposure to

150 ppm H_2S balanced with 7% O_2. (**J–L**) Locomotory speed responses to a switch from 7%

O_2 to 150 ppm H_2S balanced with 7% O_2 for animals of the indicated genotype: WT and *sqrd-*

1(tm3378) (**J**); WT and *ethe-1(yum2895)* (**K**); WT, *cysl-1(ok762), cysl-2(ok3516)*, and *cysl-*

3(yum4) (**L**). For the comparison of acute locomotory responses between strains, red bars on

the x-axis represent two intervals (4–5 minutes and 11–12 minutes, labeled a and b, respectively,

in panels A, B, C, and L) used for statistical analysis. *** = $p < 0.001$, ** = $p < 0.01$, * = $p <$

0.05, ns = not significant, Mann–Whitney U test.

1109

Figure 6. Labile iron pool sustains the locomotory activity in H_2S

(**A, B**) Volcano plots showing the relative expression of the genes involved in the regulation of

iron homeostasis in WT animals after 2-hour exposure in 50 ppm (**A**) or 150 ppm (**B**) H_2S

balanced with 7% O_2. (**C–G**) Locomotory speed responses to a switch from 7% O_2 to 150 ppm

H_2S balanced with 7% O_2 for animals of indicated genotypes or treatments: WT and animals

overexpressing *ftn-1* genomic DNA under its own promoter (#1 and #2 indicate two

independent lines) (**C**); WT, *smf-3(ok1305)*, and *ftn-1(ok3625)* (**D**); WT, and WT pretreated

with 100 μM 2,2′-Bipyridyl (BP) or with 5 mg/ml ferric ammonium citrate (FAC) in the

presence of food for 16 hours (**E**); WT, *hif-1(ia4)*, and *hif-1(ia4)* mutants pretreated with 5

mg/ml ferric ammonium citrate (FAC) for 16 hours (**F**)*;* WT, *hif-1(ia4), ftn-1(ok3625)*, and *hif-*

1(ia4); ftn-1(ok3625) double mutants (**G**). (**H**) Hypothetical model of the interactions between

labile iron pool and HIF-1 signaling. For the comparison of acute locomotory speed responses

between strains or conditions, red bars on the x-axis represent two intervals (4–5 minutes and

11–12 minutes, labeled a and b, respectively, in panels C, D, and E) used for statistical analysis.

**** = $p < 0.0001$, *** = $p < 0.001$, ** = $p < 0.01$, * = $p < 0.05$, ns = not significant, Mann–

Whitney U test.

1126

Figure 7. Mitochondrial function is required for acute response to H_2S

(**A–D**) Locomotory speed responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7%

O_2 for animals of the indicated genotype: WT and *gas-1(fc21)* (**A**); WT and *clk-1(qm30)* (**B**);

WT and *mev-1(kn1)* (**C**); WT and *isp-1(qm150)* (**D**). (**E**) Locomotory speed responses to a

switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2 for WT animals pretreated with 10

μM rotenone for 0 minute, 10 minutes, 30 minutes, 60 minutes or 120 minutes. For the

comparison of acute locomotory speed responses between strains or conditions, red bars on the

x-axis represent two intervals (4–5 minutes and 11–12 minutes) used for statistical analysis.

*** = $p < 0.001$, ** = $p < 0.01$, * = $p < 0.05$, ns = not significant, Mann–Whitney U test. (**F–**

G) Locomotory speed responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2

for animals of the indicated genotype: WT, *sod-1(tm776)*, *sod-2(ok1030)*, and *sod-3(tm760)*

(**F**); WT, *sod-4(gk101)*, *sod-5(tm1146)*, and *sod-1(tm783); sod-2(ok1030); sod-3(tm760); sod-*

4(gk101); sod-5 (tm1146) (**G**). For the comparison of acute locomotory speed responses

between strains or conditions, red bars on the x-axis represent the time interval (6–10 minutes)

used for statistical analysis. *** = $p < 0.001$, ** = $p < 0.01$, * = $p < 0.05$, ns = not significant,

Mann–Whitney U test. (**H**) Hypothetical model of the role of mitochondria in response to toxic

levels of H_2S. In this model, mitochondria play a dual role in H_2S-evoked avoidance behavior.

A transient burst of mitochondrial ROS triggered by high H_2S levels initiates locomotory

avoidance, whereas sustained ROS elevation activates stress-responsive pathways, including

HIF-1, NRF2/SKN-1, and DAF-16/FOXO, promoting adaptation to prolonged H_2S exposure.

1147

Figure supplement legends

Figure 2—figure supplement 1. Acute avoidance response to H_2S is regulated by cGMP

signaling

(**A–H**) Fraction of animals undergoing reorientation movements (omega turns, left) or

backward locomotion (reversals, right) during a switch from 7% O_2 to 150 ppm H_2S balanced

with 7% O_2 for animals of the indicated genotype: WT and *daf-19(m86)* (**A, B**); WT and *dyf-*

3(m185) (**C, D**); WT and *dyf-7(m539)* (**E, F**); WT, *daf-11(m47)*, and *tax-4(p678)* (**G, H**).

1155

Figure 2—figure supplement 2. Acute response to H_2S is regulated by *daf-11* signaling in

ASJ neurons

(**A**) The detailed expression pattern of each promoter used to express *daf-11* genomic DNA.

Promoters shown in green rescued the *daf-11(m47)* mutant phenotype. (**B–D**) Locomotory

speed responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2 for animals of

the indicated genotype: WT, *daf-11(m47)*, and transgenic *daf-11* mutants expressing *daf-11*

genomic DNA in different subsets of neurons. Red bars on the x-axis represent two intervals

(4–5 minutes and 11–12 minutes, labeled a and b, respectively) used for statistical analysis. *

= $p < 0.05$, ns = not significant, Mann–Whitney U test. (**E**) Calcium transients evoked in ASJ

neurons of WT animals in response to a switch from 7% O_2 to 150 ppm H_2S balanced with 7%

O_2 for animals (N=42). The slow decline in GCaMP6s fluorescence is due to photobleaching.

(**F**) Calcium transients evoked in ASJ neurons of WT animals in response to a switch from 7%

O_2 to 3% CO_2 balanced with 7% O_2 for animals (N=16).

1169

Figure 3—figure supplement 1. Insulin, TGF-β, and O₂ signaling modulate locomotory

response to H₂S

(**A–C**) Locomotory speed responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7%

O_2 for animals of the indicated genotype: WT and *nhr-49(nr2041)* (**A**); WT and *tax-6(ok2065)*

(**B**); WT and *cnb-1(ok276)* (**C**). (**D–F**) Locomotory responses to a switch from 7% O_2 to 150

ppm H_2S balanced with 7% O_2 for WT and *daf-2(e1370)*, including locomotory speed (**D**),

reorientation (omega turns) (**E**), and reversal (**F**). WT and *daf-2* mutants were maintained at

15°C. L4 animals were picked and shifted to 25°C until day-one adults, then assayed at room

temperature. (**G–I**) Locomotory responses to a switch from 7% O_2 to 150 ppm H_2S balanced

with 7% O_2 for WT and *gcy-9(tm7632)*, including locomotory speed (**G**), reorientation (omega

turns) (**H**), and reversal (**I**). (**J**) Locomotory speed responses to a switch from 7% O_2 to 150

ppm H_2S balanced with 7% O_2 for WT animals and wild isolate CB4856. (**K**) Optogenetic

stimulation of RMG neurons using channelrhodopsin-2 (ChR2). ChR2 was expressed in RMG

neurons using the Cre-LoxP system (Macosko et al., 2009). Blue light was delivered throughout

the assay. Locomotory speed responses to a switch from 7% O_2 to 150 ppm H_2S balanced with

7% O_2 were recorded for day-one adults grown for 16 hours in the presence or absence of ATR.

(**L**) Locomotory speed responses to a switch from 7% O_2 to 150 ppm H_2S balanced with 7%

O_2, following activation of O_2 sensing neurons by expressing a gain-of-function allele of *pkc-*

1 in O_2 sensory neurons under *gcy-37* promoter. For the comparison of acute locomotory speed

responses between strains or conditions, red bars on the x-axis represent two intervals (4–5

minutes and 11–12 minutes, indicated with red bars on the x-axis) for statistical analysis. ****

$= p < 0.0001$, *** $= p < 0.001$, ** $= p < 0.01$, ns = not significant, Mann–Whitney U test.

1192

Figure 4—figure supplement 1. Transcriptome reprogramming induced by H₂S exposure

(**A**) Venn diagram displaying the number of differentially expressed genes with adjusted *p*

value <1e-10 in WT animals exposed to 50 ppm H_2S balanced with 7% O_2 for 1, 2, or 12 hours.

(**B, C**) Significantly enriched GO categories for differentially expressed genes with adjusted *p*

value <1e-10 in WT animals exposed to 50 ppm (**B**) or 150 ppm (**C**) H$_2$S balanced with 7% O$_2$

for 2 hours. (**D, E**) Volcano plots showing the differentially expressed genes with adjusted *p*

value <1e-10 in WT animals exposed to 50 ppm (**D**) or 150 ppm (**E**) H$_2$S balanced with 7% O$_2$

for 2 hours. A set of genes involved in H$_2$S detoxification, cysteine metabolism and stress

response are highlighted in red. (**F, G**) The number of HIF-1(**F**) and SKN-1(**G**) target genes

induced by 50 ppm or 150 ppm H$_2$S exposure for 1 hour, 2 hours or 12 hours in WT animals,

with adjusted *p* value < 1e-30. (**H–J**) Venn diagrams displaying the number of differentially

expressed genes (adjusted *p* value <1e-10) in WT animals under the indicated conditions:

animals exposed to 50 ppm or 150 ppm H$_2$S for 1hour (**H**); 2 hours (**I**); and 12 hours (**J**). (**K,**

L) Volcano plots showing the differentially expressed genes (adjusted *p* value <1e-10) in WT

animals that were exposed to either 50 ppm (**K**) or 150 ppm (**L**) H$_2$S balanced with 7% O$_2$ for

12 hours. A set of genes involved in H$_2$S detoxification, cysteine metabolism and stress

response are highlighted in red.

1210

Figure 5—figure supplement 1. Acute response to H$_2$S is modulated by HIF-1 signaling

(**A, B**) Fraction of WT and *egl-9(sa307)* animals undergoing reorientation movements (omega

turns) (**A**) or backward locomotion (reversals) (**B**) during a switch from 7% O$_2$ to 150 ppm H$_2$S

balanced with 7% O$_2$. (**C, D**) Locomotory speed responses to a switch from 7% O$_2$ to 1% O$_2$

for WT and *egl-9(sa307)* (**C**); WT, *hif-1(ia4)*, and *semo-1(yum2889)* (**D**). (**E**) HIF-1 signaling

regulates the H$_2$S mitochondrial detoxification pathway. The detoxification enzyme SQRD-1

(sulfide:quinone oxidoreductase) oxidizes H$_2$S, donating electrons to ubiquinone (uq), and

rhodoquinone (rq) of the mitochondrial electron transport chain (ETC). ETHE-1 (persulfide

dioxygenase) oxidizes sulfur compounds to sulfite. (**F–I**) Fraction of animals undergoing

reorientation movements (omega turns, F and H) or backward locomotion (reversals, G and I)

during a switch from 7% O$_2$ to 150 ppm H$_2$S balanced with 7% O$_2$ for the indicated genotypes:

WT and *sqrd-1(tm3378)* (**F, G**)*;* and WT and *ethe-1(yum2895)* (**H, I**). (**J**) Locomotory speed

responses to a switch from 7% O$_2$ to 1% O$_2$ for WT, *sqrd-1(tm3378)*, and *ethe-1(yum2895)*. (**K,**

L) Fraction of WT, *cysl-1(ok762)*, *cysl-2(ok3516)*, and *cysl-3(yum4)* animals undergoing

reorientation movements (omega turns, K) or backward locomotion (reversals, L) during a

switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2. (**M**) Locomotory speed responses to

a switch from 7% O_2 to 1% O_2 for WT, *cysl-1(ok762), cysl-2(ok3516)*, and *cysl-3(yum4)*

animals. (**N**) Locomotory speed responses to a switch from 7% O_2 to 150 ppm H_2S balanced

with 7% O_2 for WT and *semo-1(yum2889)*. (**O, P**) Fraction of animals undergoing reorientation

(omega turns, **O**) or backward locomotion (reversals, **P**) in response to a switch from 7% O_2 to

150 ppm H_2S balanced with 7% O_2. Comparisons of acute locomotory responses between

strains were performed using two intervals (4–5 minutes and 11–12 minutes, indicated with red

bars on the x-axis) for statistical analysis. *** = $p < 0.001$, ns = not significant, Mann–Whitney

U test.

1235

Figure 6—figure supplement 1. Labile iron pool sustains the locomotory activity in H_2S

(**A–D**) Volcano plots showing the relative expression of the genes involved in the regulation of

iron homeostasis (*ftn-1, ftn-2*, and *smf-3*) in WT animals under the following conditions: 1-hour

exposure in 50 ppm H_2S balanced with 7% O_2 (**A**); 1-hour exposure in 150 ppm H_2S balanced

with 7% O_2 (**B**); 12-hour exposure in 50 ppm H_2S balanced with 7% O_2 (**C**); and 12-hour

exposure in 150 ppm H_2S balanced with 7% O_2 (**D**). (**E, F**) Locomotory speed responses to a

switch from 7% O_2 to 1% O_2 for animals of the indicated genotype or treatment: WT, and two

independent transgenic lines overexpressing *ftn-1* genomic DNA under its own promoter (#1

and #2) (**E**); WT animals, WT animals treated with 100 μM 2,2′-Bipyridyl (BP), and WT

animals treated with 5 mg/ml ferric ammonium citrate (FAC) for 16 hours (**F**). Red bar on the

x-axis represents one-minute time interval (3–4 minutes) used for statistical analysis. * = $p <$

0.05, ns = not significant, Mann–Whitney U test. (**G, H**) Fraction of animals undergoing

reorientation movements (omega turns) (**G**) or backward locomotion (reversals) (**H**) during a

switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2 for animals of the indicated genotypes

WT, *smf-3(ok1305)*, and *ftn-1(ok3625)*. (**I, J**) Fraction of animals undergoing reorientation

movements (omega turns) (**I**) or backward locomotion (reversals) (**J**) during a switch from 7%

O_2 to 150 ppm H_2S balanced with 7% O_2 for WT animals, WT animals pretreated with either

100 μM 2,2′-Bipyridyl (BP) or with 5 mg/ml ferric ammonium citrate (FAC) in the presence of

food for 16 hours.

1255

Figure 7—figure supplement 1. Mitochondrial function is required for acute response to

H_2S

(**A–D**) Fraction of animals undergoing reorientation movements (omega turns) or backward

locomotion (reversals) during a switch from 7% O_2 to 150 ppm H_2S balanced with 7% O_2 for

the indicated genotype or treatment: WT and *clk-1(qm30)* (**A, B**); WT and *mev-1(kn1)* (**C, D**).

(**E**) Locomotory speed responses to a switch from 7% O_2 to 1% O_2 for WT, *sod-2(ok1030);* and

sod-3(tm760). (**F, G**) Fraction of animals undergoing reorientation movements (omega turns,

F) or backward locomotion (reversals, **G**) during a switch from 7% O_2 to 150 ppm H_2S balanced

with 7% O_2 for WT and *sod-1(tm783); sod-2(ok1030); sod-3(tm760); sod-4(gk101); sod-5*

(tm1146). (**H**) Locomotory response to acute near UV exposure for WT and *sod-1(tm783); sod-*

2(ok1030); sod-3(tm760); sod-4(gk101); sod-5 (tm1146) before and after 30 minutes of

exposure to 150ppm H_2S.

1268

1269   **Figure 1—Video 1. The locomotory response of WT animals to H_2S**

1270   An 8x accelerated video clip (original length: 16 minutes) showing WT (N2) animals exposed

1271   to 7% O_2 for 2 minutes, followed by 11 minutes in 150 ppm H_2S, and then returned to 7% O_2.

1272   The H_2S exposure segment is highlighted and begins at the 15-second mark, corresponding to

1273   2 minutes in the original recording.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 2—figure supplement 1

Figure 2—figure supplement 2

A

Promoter	neurons														
Pdaf-11	ASH	AWB	AWC	ASJ	ASK	ADL	ASI	ASE							
Podr-3	ASH	AWB	AWC						ADF	AWA					
Podr-1		AWB	AWC	ASJ	ASK		ASI								
Pflp-21	ASH				ADL	ASI	ASE			FLP	URA	RMG	MC	M2	M4
Pgpa-11	ASH				ADL										
Psra-6	ASH						ASI								PVQ
Ptrx-1				ASJ											

Figure 3—figure supplement 1

Figure 4—figure supplement 1

Figure 5—figure supplement 1

Figure 6—figure supplement 1

Figure 7—figure supplement 1