MSIsensor-RNA: microsatellite instability detection for bulk and single-cell gene expression data

Peng Jia[1,2,†], Xuanhao Yang[1,2, †], Xiaofei Yang[1,3], Tingjie Wang[1,3], Yu Xu[4], and Kai Ye[1,2,4,5,]*

[1]MOE Key Lab for Intelligent Networks & Networks Security, Faculty of Electronic and Information Engineering, Xi'an Jiaotong University, Xi'an, 710049, China

[2]School of Automation Science and Engineering, Faculty of Electronic and Information Engineering, Xi'an Jiaotong University, Xi'an, 710049, China

[3]School of Computer Science, Faculty of Electronic and Information Engineering, Xi'an Jiaotong University, Xi'an 710049, China

[4]School of Life Science and Technology, Xi'an Jiaotong University, Xi'an, 710049, China

[5]Genome Institute, the First Affiliated Hospital of Xi'an Jiaotong University, Xi'an 710061, China

*Correspondence: kaiye@xjtu.edu.cn (Ye K)

[†]These authors contributed equally to this work.

Abstract

Background: Microsatellite instability (MSI) is an indispensable biomarker in

cancer immunotherapy. Currently, MSI scoring methods by high-throughput omics

methods have gained popularity and demonstrated better performance than the gold

standard method for MSI detection. However, MSI detection method on expression

data, especially single-cell expression data is still lacking, limiting the scope of

clinical application and prohibiting investigation of MSI at single cell level.

Results: Herein, we developed MSIsensor-RNA, an accuracy, robust, adaptable, and

standalone software, to detect MSI status by its associated genes' expression values.

We demonstrated the favorable performance and promise of MSIsensor-RNA in both

bulk and single-cell gene expression data in multiplatform technologies including

RNA-seq, Microarray, and single-cell RNA-seq.

Conclusions: MSIsensor-RNA is a versatile, efficient, and robust method for MSI

status detection from both bulk and single-cell gene expression data in clinical

researches   and   applications.   MSIsensor-RNA   is   available   at

https://github.com/xjtu-omics/msisensor-rna.

Keywords: microsatellite instability, cancer, gene expression, multiplatform,

single-cell RNA-seq, RNA-seq, Microarray

Background

Microsatellite instability (MSI) refers to hypermutations of microsatellite sites due to inactivating alterations of mismatch repair (MMR) genes in malignancies [1, 2]. Currently, MSI is an indispensable pan-cancer biomarker in cancer immunotherapy therapy and prognosis, and it is routinely examined in multiple cancer types, particularly in colorectal cancer (CRC), stomach adenocarcinoma (STAD), and uterine corpus endometrial carcinoma (UCEC) [2-5]. For example, MSI positive patients are often resistant to 5-fluorouracil treatment but have a better outcome for immune checkpoint blockade treatment [4, 5].

In clinical settings, MSI detection mainly relies on the gold-standard experimental method, MSI-PCR [6], which is laborious and time-consuming. With the advancement of next-generation-sequencing technology, numerous features of genomics, epigenomics, transcriptomics, and histology are investigated, and novel MSI computational algorithms have been developed for a variety of scenarios [7-20]. Genomics-based methods quantify MSI according to genetic mutations at microsatellite sites, which achieve high accuracy and are becoming popular in clinical MSI detection. For example, MSIsensor [9] detects MSI with high concordance as 99.4% on MSK-IMPACT panel [21]. Epigenomics-based method MIRMMR [18] detects MSI using methylation levels in MMR pathway with 0.97

AUC. In addition, transcription levels of MSI-associated genes exhibit correlation

with MSI, hinting possibility of MSI detection using transcriptomics data [15-17].

Besides these high-throughput technologies, deep learning algorithms were also

applied to hematoxylin and eosin-stained slides to detect MSI [19, 20]. However, all

these MSI methods detected MSI at a sample level, lacking cell-level measuring of

MSI. Recently, single-cell RNA-seq (scRNA-seq) technology enables investigation

of cell specific transcriptome and sheds light on tumor heterogeneity and tumor

stages. In particular, the single-cell and spatial transcriptome enable the dynamic

analysis of MSI in the complex tumor microenvironment, such as in metastatic and

recurrent cancer [22]. However, current MSI detection methods designed for bulk

gene expression data do not perform well on scRNA-seq samples. For example, the

only software for gene expression data, PreMSIm [16], only provided fixed

signatures and a fixed model for all cancers, which limits the widely application of

the methods. Moreover, the normalized method in PreMSIm also leads to poor

performance with abnormal samples. Here, we developed MSIsensor-RNA, a robust

method for MSI-associated genes detection and MSI evaluation for both bulk gene

expression data and single-cell RNA-seq data.

Implementation

Dataset. We downloaded RNA-seq data of 1,428 TCGA samples across CRC, STAD,

and UCEC from TCGA Research Network (https://portal.gdc.cancer.gov) and

obtained their MSI status determined by gold standards (**Table S1**). We obtained 141

RNA-seq samples of ICGC from ICGC data portal (https://dcc.icgc.org), and their

MSI status reported by MIMcall [23]. Another 106 RNA-seq samples with the

matched MSI status were downloaded from public publication of Clinical Proteomic

Tumor Analysis Consortium (CPTAC) [24]. We also downloaded Microarray data

and their MSI status of 1,468 samples across CRC and STAD from GEO dataset

(https://www.ncbi.nlm.nih.gov/geo). For scRNA-seq data, we got the gene

expression data and their MSI status from 133 CRC samples in two recent

publications [25, 26].

Overall design. The pipeline of MSIsensor-RNA consists of data preprocessing,

informative genes selection, model training, and model testing (**Fig. 1 and Fig. S1**).

First, we preprocess the expression values of samples from Microarray, bulk

RNA-seq, and scRNA-seq. Next, we select an informative gene set for MSI

detection from 1,428 TCGA samples. Then we used these TCGA samples to train a

machine learning model for each cancer type for MSI scoring. Finally, we applied

the trained model to independent databases to test the performance of the

MSIsensor-RNA for each cancer type.

Data preprocessing. In MSIsensor-RNA, we accept Microarray expression value,

FPKM, TPM, and RESM read count as input. All values of expression matrix were

added 1 and followed by log2 transformed. Then, for each sample or cell, expression

values were normalized as a Gaussian distribution with 0 mean and 1 standard

deviation. For scRNA-seq sample, to obtain accurate MSI status, we only included

high-quality cells with at least 20% genes detected for MSI detection. If the number

of high-quality cells was less than 20, we sort all cells by the ratio of detected genes

in descending order, and the top 20 cells would be utilized for MSI detection. To

solve the dropout problem of scRNA-seq, we imputed zero values by the average of

the gene expression value in the given sample.

Selection of informative genes. We select informative genes for MSI classification

in terms of stability, discrimination, and generalization. Firstly, we remove

ribosomal genes, mitochondrial genes, and genes with low FPKM in TCGA dataset.

Secondly, we selected genes with discriminative gene expression signatures between

MSI samples and MSS samples. We perform rank-sum tests for expression values

between MSI samples and MSS samples for each gene, and only genes with *P* value

< 0.01 are included for the following analysis. Furthermore, we compute the fold of

*i*th gene by:

$$F^i = \left| \log2 \left(\frac{\frac{1}{n} \sum_{j=1}^{n} G_j^i}{\frac{1}{m-n} \sum_{k=n+1}^{m} G_k^i} \right) \right|$$

where *m* is the sample number for informative genes selection, *n* is the MSI sample

number, G_j^i is the gene expression value of ith for j sample. We only select genes

with fold > 0.5 for candidate informative genes. Finally, we keep genes with more

generalization ability for MSI detection. We calculate the area under the receiver

operating characteristic curve (AUC) of the gene expression value and only genes

with AUC > 0.65 are kept for next step. We also calculate the 10-fold cross

validation score of SVM and random forest, and only first quartet genes are included

the finial informative gene set (**Fig. S2**).

Machine learning model training and testing. We build a support vector machine

(SVM) model to classify the MSI status for CRC, STAD, and UCEC in TCGA

dataset. Firstly, we utilized SOMTE [27] to correct the imbalance between MSI and

MSS in each cancer type by amplifying the MSI samples. Then, we utilized the

expression values from correct data as input to train SVM model for MSI

classification. To evaluate the performance of MSIsensor-RNA, we tested the

trained model with 1,848 independent samples of multiplatform including 247

RNA-seq, 1,468 Microarray, and 133 scRNA-seq samples. For a scRNA-seq sample,

we calculated the MSI score with SVM model for each high-quality cell. Then the

average cell MSI score is used to evaluate the MSI status of a scRNA-seq sample.

PreMSIm running. To compare performance of MSIsensor-RNA with the only

standalone software PreMSIm, we also apply the data of Microarray, RNA-seq, and

131 scRNA-seq from 1,848 independent samples to PreMSIm. For Microarray and

132 RNA-seq samples, we test PreMSIm with two modes: PreMSIm-all and

133 PreMSIm-split. In PreMSIm-all, we integrate all input samples to PreMSIm

134 normalized module and predicted module. PreMSIm-split referred to input samples

135 one database for each run.

136 *Performance comparison of MSIsensor-RNA and PreMSIm.* In MSIsensor-RNA,

137 the predicted MSI probability by the SVM model was used to score the MSI status.

138 The probabilities were further transformed to MSI status by the Youden index [28].

139 We first compared the MSIsensor-RNA score between MSI and MSS samples to test

140 the performance of MSIsensor-RNA in multiplatform by rank sum test. To further

141 evaluate the performance of two MSI detected methods, we calculated AUC,

142 accuracy, F-score, precision, sensitivity, and specificity of MSIsensor-RNA and

143 PreMSIm in different sequencing technologies.

144 *Robustness testing of MSIsensor-RNA and PreMSIm.* To test the performance of

145 MSIsensor-RNA and PreMSIm at different normalized methods, we tested these two

146 methods with FPKM, TPM, and read counts format of TCGA samples and

147 calculated the AUC, F1-score, accuracy, precision, sensitivity, and specificity of

148 each normalized method. To overcome the bias of different normalized methods and

149 sequencing technology, we normalized the input data of each sample to a Gaussian

distribution with 0 mean and 1 standard deviation. However, in PreMSIm, the

normalization process was performed by genes, which means the normalized input

data of a sample would be influenced by other samples in the bulk. Here, we tested

the PreMSIm in two ways. Firstly, we input TCGA samples by three cancer types

and calculated the performance of predicted MSI result. Secondly, we input all

TCGA samples together to evaluate its performance. We further compared the MSI

result and performance of these two ways and found that the performance of

PreMSIm was affected by the way input was provided.

Results

The workflow of MSIsensor-RNA includes four modules (**Fig. 1 and Fig. S1**). First,

we preprocess the expression value of Microarray, bulk RNA-seq, and scRNA-seq

data. Then, we select a set of informative genes for MSI detection. Next, we train a

support vector machine (SVM) model to estimate MSI scores using gene expression

values of the selected informative genes. Finally, we apply the trained model to

predict MSI score for either one clinical sample or a single cell (**Table S1**). For a

given scRNA-seq sample, we also developed a model to report MSI status of this

sample by integrating MSI scores of cells within.

MSIsensor-RNA accepts a variety of expression data including FPKM, RESM

normalized read count, TPM, or microarray expression format as input. Input

expression values were added 1 and then log2 transformed following Z-score

normalization per sample or cell. In particular, for single cell module of

MSIsensor-RNA, we only included high-quality cell in following steps, and the

missing values of each gene in high-quality cells were imputed by the average of the

gene expression value in this sample.

The informative gene selection module consists of three key steps (**Fig. S2**): (i)

removing mitochondrial genes and ribosomal genes; (ii) filtering of genes, of which

expression values do not differ significantly between MSI and MSS samples; (iii)

keeping genes, of which expression values have high generalized scores for MSI

detection (online methods). We applied the gene selection module to 1,428 samples

based on the gene expressions (FPKM values) from three MSI-popular cancer types

(CRC, STAD, and UCEC) in TCGA dataset and finally obtained 109 informative

genes for MSI classification. We also performed this step for each type of CRC,

STAD, and UCEC, yielding 397, 206, and 86 informative genes, respectively (**Fig.**

S4 and Table S2-S5). We found that only eight informative genes are detected in all

three cancer types. Of which, we found that *MLH1* was the most important

informative gene for MSI detection, as confirmed by previous reports [15-17] (**Fig.**

S5).

To assess the performance of MSIsensor-RNA in bulk sample data, we first

trained tumor-specific models for CRC, STAD, and UCEC, as well as a model for

all three MSI-popular cancer types in the TCGA dataset. Then we compared the two

kinds of models (tumor-specific and MSI-popular) with the standalone software,

PreMSIm, in terms of the area under the curve (AUC) of the receiver operating

characteristic (ROC), accuracy, sensitivity, and specificity in 1,715 (1468

Microarray and 247 bulk RNA-seq samples) independent samples. Notably,

MSIsensor-RNA normalizes the expression value of informative genes for each

sample independently, while PreMSIm must normalize each gene for multiple

samples at the same time. Thus, we examined PreMSIm with all samples normalized

together (PreMSIm-all) or by database (PreMSIm-split).

For Microarray data, we computed MSI status by MSIsensor-RNA and PreMSIm

in 1,468 samples from 12 GEO accessions. The result showed that MSIsensor-RNA

predicted MSI with 0.952 AUC, while PreMSIm only performed 0.628 AUC in

PreMSIm-split and 0.912 AUC in PreMSIm-all mode (**Fig. 2A, S6, S7; Table S6**

and S7). Meanwhile, MSIsensor-RNA achieved much higher sensitivities than

PreMSIm-split, and preMSI-all (MSIsensor-RNA: 0.968, PreMSIm-split: 0.912,

PreMSIm-all: 0.384) and comparable specificities with PreMSIm-split, and

preMSI-all (MSIsensor-RNA: 0.843, PreMSIm-split: 0.912, PreMSIm-all: 0.873).

To evaluate the performance using bulk RNA-seq data, we compared

MSIsensor-RNA and two modes of PreMSIm on 247 independent samples from

ICGC and CPTAC. We noticed that MSIsensor-RNA achieved 0.997 AUC in

tumor-specific model and 0.985 AUC in MSI-popular model, which were

significantly greater than PreMSIm-all (0.5) and PreMSIm-split (0.870) (**Fig. 2B, S8,**

S9; Table S8 and S9). In addition, MSIsensor-RNA performed much better than

PreMSIm for both sensitivity (MSIsensor-RNA with tumor-specific model: 0.951,

MSIsensor-RNA with MSI-popular model: 0.973, PreMSIm-split: 0.834,

PreMSIm-all: 0.25) and specificity (MSIsensor-RNA with tumor-specific mode: 1,

MSIsensor-RNA with MSI-popular model: 0.923, PreMSIm-split: 0.906,

PreMSIm-all: 0.75). To further investigate the robustness of MSIsensor-RNA for

different input data types, we evaluated the performance of MSIsensor-RNA and

PreMSIm with FPKM, read count, and TPM normalized samples in TCGA as input.

We found that MSIsensor-RNA achieved 0.982 ± 0.040 AUC indicating the

robustness of MSIsensor-RNA regardless of the measurements of gene expression

(**Table S10**).

To assess the performance of MSIsensor-RNA and PreMSIm in scRNA-seq

samples, we applied the trained model of MSIsensor-RNA to 23,902 high-quality

cells from 133 samples to obtain sample specific MSI status and compared to the

ratio of cells labeled as MSI by PreMSIm. The result showed MSIsensor-RNA

detected MSI for scRNA-seq samples with 0.958 AUC, 0.9231 sensitivity, and

0.9362 specificity, while PreMSIm with 0.4969 AUC, 1 sensitivity, and 0.0319

specificity (**Fig. 2A**, **S10**; **Table S11 and S12**). The sample level MSI scores based

on scRNA-seq was significantly different between MSI and MSS samples by

MSIsensor-RNA (rank-sum test, $P = 1.01 \times 10^{-16}$) while no significant difference was

detected for PreMSIm (rank-sum test, $P = 0.9547$) (**Fig. 2B**). Having established the

effectiveness of MSIsensor-RNA on scRNA-seq sample, we investigated cell-level

MSI. We computed the MSI scores of 21,438 high-quality cells from 100 samples

(GSE178341) and found cell-type dependent MSI scores. For example, MSI scores

of epithelial and immune cells in MSI samples were greater than that in MSS

samples while no significant difference was detected between MSI and MSS for

stromal cells (**Fig. 2C, S11 and Table S13**). This indicated the potential of

MSIsensor-RNA to assess MSI at the single-cell level, providing a novel

measurement for the investigation of tumorigenic process.

Discussion

Microsatellite instability is important for the prognosis assessment of both 5-FU

chemotherapy [4] and immunotherapy [5]. In addition to gold-standard experimental

methods [6], MSI status is also evaluated according to genomic sequencing data

[7-14], gene expression data [15-17], methylation data [18], and H&E-stained slides

[19, 20]. Compared to variants in microsatellite regions, gene expression values are

more directly reflective of the features of MSI and easier to obtain. In this study, we

developed a robust method, MSIsensor-RNA, for MSI detection with gene

expression data. MSIsensor-RNA provided informative gene selections, model

training, and MSI detection modules. MSIsensor-RNA is able to process data from

multiple platforms, including Microarray, RNA-seq, and single cell RNA-seq.

Compared to the standalone method PreMSIm, MSIsensor-RNA also provided

modules for informative gene selection and model training so that users could apply

MSIsensor-RNA for different cancer types. MSIsensor-RNA also improved the

normalization method of the data, yielding a more robust result than PreMSIm (Fig.

2). In addition, MSIsensor-RNA facilitates the evaluation of MSI status at the single

cell level, which will be critical to better understanding the mechanism of MSI in

cancer immunotherapy in the future.

In most MSI detection methods, such as MSIsensor [10] and MSIsensor-pro [11],

MSI is quantified according to genetic mutations at microsatellite sites, the

consequence of MSI rather than the deficiency of the MMR system, the direct cause

of MSI. In this study, a set of MSI-associated genes was identified, and their

expression values were used for MSI evaluation. We found that *MLH1* is the most

important gene in all tested cancer types. In addition, unexpected expression of

MLH1 is commonly seen in Lynch syndrome [29]. Thus, we test the performance of

MSIsensor-RNA for samples with abnormal *MLH1* expression. We train a model

based on all informative genes and tested it by samples with simulated abnormal

MLH1 gene expression (**Table S14**). We found that the model achieved 0.974 and

0.972 AUCs when we set the *MLH1* expression value as the maximum and

minimum of all gene expression values, respectively. Furthermore, when *MLH1* was

excluded from the informative gene set, MSIsensor-RNA also achieved a 0.977

AUC, indicating the robustness of MSIsensor-RNA for MSI detection.

We demonstrate that MSIsensor-RNA achieved higher performance than other

methods based on gene expression and comparable performance compared to

DNA-based methods (**Table S15**). In our study design, MSIsensor-RNA detects MSI

according to the gene expression signature of genes on MSI associated pathways,

while MSIsensor evaluates MSI by computing the ratio of somatic microsatellite

mutations. Although MSIsensor achieved slightly higher performance than

MSIsensor-RNA, it cannot replace the applications of MSIsensor-RNA in gene

expression data. Currently, MSIsensor-RNA reports favorable performance in all

three MSI-popular cancers, including colorectal cancer, stomach adenocarcinoma,

and uterine corpus endometrial carcinoma. The MSI features are different in

different cancer types. Thus, the model obtained low performance when the testing

samples were inconsistent with training samples in cancer types (**Table S16-S18**).

Therefore, the performance of MSIsensor-RNA in other cancer types needs further

validation in the future.

Conclusions

MSIsensor-RNA is a cross-platform, efficient, and robust method for MSI status determination from both bulk and single-cell gene expression data. We demonstrated the effectiveness and robustness of MSIsensor-RNA across different platforms, hinting its potential in clinical research. Moreover, MSIsensor-RNA enables single-cell level MSI evaluation, providing a new tool to discover the role of MSI in tumorigenic process and to monitor cell-level dynamic changes during immunotherapy.

Availability and requirements

Project name: msisensor-rna

Project home page: https://github.com/xjtu-omics/msisensor-rna

Operating system(s): Unix System or Docker

Programming language: python

Other requirements: python packages including numpy, pandas and scikit-learn.

License: Custom License (see at homepage)

Any restrictions to use by non-academics: MSIsensor-RNA is free for

non-commercial use by academic, government, and non-profit/not-for-profit

institutions. A commercial version of the software is available and licensed through

Xi'an Jiaotong University. For more information, please contact with

pengjia@stu.xjtu.edu.cn or kaiye@xjtu.edu.cn.

Abbreviations

MSI: Microsatellite Instability

MMR: Mismatch Repair

CRC: Colorectal Cancer

STAD: Stomach Adenocarcinoma

UCEC: Uterine Corpus Endometrial Carcinoma

NGS: Next Generation Sequencing

ROC: Receiver Operating Characteristic

AUC: Area Under the Curve

scRNA-seq single-cell RNA sequencing

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

MSIsensor-RNA is a python program which is available at

https://github.com/xjtu-omics/msisensor-rna; Supplementary information:

Supplementary data are available at xxxx online.

Competing interests

The authors declare that they have no competing interests.

Funding

K.Y. and X.Y. are supported by the National Natural Science Foundation of China

(32125009, 32070663 and 62172325), by the National Key R&D Program of China

(2022YFC3400300), the Key Construction Program of the National "985" Project,

the Fundamental Research Funds for the Central Universities (xzy012020012), and

the Natural Science Basic Research Program of Shaanxi (2021GXLH-Z-098).

Authors' contributions

KY conceived and designed the study. PJ collected the data. PJ, XUY, XIY, TW, and YX

discussed and developed the method. PJ and XUY implemented the method. PJ and KY

wrote the manuscript. The authors read and approved the final manuscript.

Acknowledgments

We thank everyone in the Ye lab at Xi'an Jiaotong University for helpful comments

and discussion. We also thank the producers of data we used.

References

1.   Yamamoto H, Imai K: **Microsatellite instability: an update**. *Arch Toxicol*
2015, **89**(6):899-921.

2.   Hause RJ, Pritchard CC, Shendure J, Salipante SJ: **Classification and**
characterization of microsatellite instability across 18 cancer types. *Nat*
Med 2016, **22**(11):1342-1350.

3.   Rizvi NA, Hellmann MD, Snyder A, Kvistborg P, Makarov V, Havel JJ, Lee W,
Yuan J, Wong P, Ho TS *et al*: **Cancer immunology. Mutational landscape**
determines sensitivity to PD-1 blockade in non-small cell lung cancer.
Science 2015, **348**(6230):124-128.

4.   Ribic CM, Sargent DJ, Moore MJ, Thibodeau SN, French AJ, Goldberg RM,
Hamilton SR, Laurent-Puig P, Gryfe R, Shepherd LE *et al*: **Tumor**
microsatellite-instability status as a predictor of benefit from
fluorouracil-based adjuvant chemotherapy for colon cancer. *N Engl J*
Med 2003, **349**(3):247-257.

5.   Le DT, Uram JN, Wang H, Bartlett BR, Kemberling H, Eyring AD, Skora AD,
Luber BS, Azad NS, Laheru D *et al*: **PD-1 Blockade in Tumors with**
Mismatch-Repair Deficiency. *N Engl J Med* 2015, **372**(26):2509-2520.

6.   Baudrin LG, Deleuze JF, How-Kit A: **Molecular and Computational**
Methods for the Detection of Microsatellite Instability in Cancer. *Front*
Oncol 2018, **8**:621.

7.   Janavicius R, Matiukaite D, Jakubauskas A, Griskevicius L: **Microsatellite**
instability detection by high-resolution melting analysis. *Clin Chem* 2010,
56(11):1750-1757.

8.   Boland CR, Thibodeau SN, Hamilton SR, Sidransky D, Eshleman JR, Burt
RW, Meltzer SJ, Rodriguez-Bigas MA, Fodde R, Ranzani GN *et al*: **A**
National Cancer Institute Workshop on Microsatellite Instability for
cancer detection and familial predisposition: development of

international criteria for the determination of microsatellite instability in
colorectal cancer. *Cancer Res* 1998, **58**(22):5248-5257.

9.   Kautto EA, Bonneville R, Miya J, Yu L, Krook MA, Reeser JW,
Roychowdhury S: **Performance evaluation for rapid detection of**
pan-cancer microsatellite instability with MANTIS. *Oncotarget* 2016,
8(5).

10.   Huang MN, McPherson JR, Cutcutache I, Teh BT, Tan P, Rozen SG: **MSIseq:**
Software for Assessing Microsatellite Instability from Catalogs of
Somatic Mutations. *Sci Rep* 2015, **5**:13321.

11.   Niu B, Ye K, Zhang Q, Lu C, Xie M, McLellan MD, Wendl MC, Ding L:
MSIsensor: microsatellite   instability   detection   using   paired
tumor-normal sequence data. *Bioinformatics* 2014, **30**(7):1015-1016.

12.   Jia P, Yang X, Guo L, Liu B, Lin J, Liang H, Sun J, Zhang C, Ye K:
MSIsensor-pro: Fast, Accurate, and Matched-normal-sample-free
Detection   of   Microsatellite   Instability. *Genomics   Proteomics*
Bioinformatics 2020, **18**(1):65-71.

13.   Hempelmann JA, Scroggins SM, Pritchard CC, Salipante SJ: **MSIplus for**
Integrated Colorectal Cancer Molecular Testing by Next-Generation
Sequencing. *J Mol Diagn* 2015, **17**(6):705-714.

14.   Salipante SJ, Scroggins SM, Hampel HL, Turner EH, Pritchard CC:
Microsatellite instability detection by next generation sequencing. *Clin*
Chem 2014, **60**(9):1192-1199.

15.   Maruvka YE, Mouw KW, Karlic R, Parasuraman P, Kamburov A, Polak P,
Haradhvala NJ, Hess JM, Rheinbay E, Brody Y *et al*: **Analysis of somatic**
microsatellite indels identifies driver events in human tumors. *Nat*
Biotechnol 2017, **35**(10):951-959.

16.   Ratovomanana T, Cohen R, Svrcek M, Renaud F, Cervera P, Siret A,
Letourneur Q, Buhard O, Bourgoin P, Guillerm E *et al*: **Performance of**
Next-Generation   Sequencing   for   the   Detection   of   Microsatellite
Instability in Colorectal Cancer With Deficient DNA Mismatch Repair.
Gastroenterology 2021, **161**(3):814-826 e817.

17.   Pacinkova A, Popovici V: **Cross-platform Data Analysis Reveals a Generic**
Gene Expression Signature for Microsatellite Instability in Colorectal
Cancer. *Biomed Res Int* 2019, **2019**:6763596.

18.   Li L, Feng Q, Wang X: **PreMSIm: An R package for predicting**

microsatellite instability from the expression profiling of a gene panel in
cancer. *Comput Struct Biotechnol J* 2020, **18**:668-675.

19. Danaher P, Warren S, Ong S, Elliott N, Cesano A, Ferree S: **A gene**
expression assay for simultaneous measurement of microsatellite
instability and anti-tumor immune activity. *Journal for immunotherapy of*
cancer 2019, **7**(1):15.

20. Foltz SM, Liang WW, Xie M, Ding L: **MIRMMR: binary classification of**
microsatellite instability using methylation and mutations. *Bioinformatics*
2017, **33**(23):3799-3801.

21. Echle A, Grabsch HI, Quirke P, van den Brandt PA, West NP, Hutchins GGA,
Heij LR, Tan X, Richman SD, Krause J *et al*: **Clinical-Grade Detection of**
Microsatellite Instability in Colorectal Tumors by Deep Learning.
Gastroenterology 2020, **159**(4):1406-1416 e1411.

22. Kather JN, Pearson AT, Halama N, Jager D, Krause J, Loosen SH, Marx A,
Boor P, Tacke F, Neumann UP *et al*: **Deep learning can predict**
microsatellite instability directly from histology in gastrointestinal cancer.
Nat Med 2019, **25**(7):1054-1056.

23. Middha S, Zhang L, Nafa K, Jayakumaran G, Wong D, Kim HR, Sadowska J,
Berger MF, Delair DF, Shia J *et al*: **Reliable Pan-Cancer Microsatellite**
Instability Assessment by Using Targeted Next-Generation Sequencing
Data. *JCO Precis Oncol* 2017, **2017**(1):1-17.

24. Longo SK, Guo MG, Ji AL, Khavari PA: **Integrating single-cell and spatial**
transcriptomics to elucidate intercellular tissue dynamics. *Nat Rev Genet*
2021, **22**(10):627-644.

25. Fujimoto A, Fujita M, Hasegawa T, Wong JH, Maejima K, Oku-Sasaki A,
Nakano K, Shiraishi Y, Miyano S, Yamamoto G *et al*: **Comprehensive**
analysis of indels in whole-genome microsatellite regions and
microsatellite instability across 21 cancer types. *Genome Res* 2020.

26. Vasaikar S, Huang C, Wang X, Petyuk VA, Savage SR, Wen B, Dou Y, Zhang
Y, Shi Z, Arshad OA *et al*: **Proteogenomic Analysis of Human Colon**
Cancer Reveals New Therapeutic Opportunities. *Cell* 2019,
177(4):1035-1049 e1019.

27. Pelka K, Hofree M, Chen JH, Sarkizova S, Pirl JD, Jorgji V, Bejnood A,
Dionne D, Ge WH, Xu KH *et al*: **Spatially organized multicellular immune**
hubs in human colorectal cancer. *Cell* 2021, **184**(18):4734-4752 e4720.

28.   Lee HO, Hong Y, Etlioglu HE, Cho YB, Pomella V, Van den Bosch B,
Vanhecke J, Verbandt S, Hong H, Min JW *et al*: **Lineage-dependent gene**
expression programs influence the immune landscape of colorectal
cancer. *Nat Genet* 2020, **52**(6):594-603.

29.   Chawla NV, Bowyer KW, Hall LO, Kegelmeyer WP: **SMOTE: Synthetic**
Minority Over-sampling Technique. *Journal of Artificial Intelligence*
Research 2002, **16**:321-357.

30.   Schisterman EF, Perkins NJ, Liu A, Bondell H: **Optimal cut-point and its**
corresponding Youden Index to discriminate individuals using pooled
blood samples. *Epidemiology* 2005, **16**(1):73-81.

31.   Chen W, Hampel H, Pearlman R, Jones D, Zhao W, Alsomali M, Knight D,
Frankel WL: **Unexpected expression of mismatch repair protein is more**
commonly seen with pathogenic missense than with other mutations in
Lynch syndrome. *Hum Pathol* 2020, **103**:34-41.

Figure legends

Fig. 1. Workflow of MSIsensor-RNA. MSIsensor-RNA includes four modules:

data preprocessing, informative gene selection, SVM model training, and testing.

MSIsensor-RNA selects informative genes and trains SVM model by RNA-seq

samples from TCGA. MSI scores are predicted by the trained model for Microarray,

RNA-seq, and scRNA-seq samples.

Fig. 2. Performance of MSIsensor-pro. A-C. AUC of MSIsensor-RNA and

PreMSIm in Microarray (A), RNA-seq (B), and scRNA-seq (C) samples.

Tumor-specific: MSI results with tumor specific model; MSI-popular: MSI results

with three MSI-popular cancer types. **D.** Boxplot of MSIsensor-RNA score in

scRNA-seq samples. **E.** Violin plot of MSIsensor-RNA score of different cell types

in scRNA-seq samples. Epithelial, stromal, and immune cell types are defined in

Pelka et al.[25]

Supplementary tables

Table S1. Overview of samples in this study.

Table S2. Details of informative genes in CRC.

Table S3. Details of informative genes in STAD.

Table S4. Details of informative genes in UCEC.

Table S5. Details of informative genes in three MSI-popular cancers.

Table S6. MSI results of Microarray samples by MSIsensor-RNA and PreMSIm.

Table S7. MSI detection performance of MSIsensor-RNA and PreMSIm in

Microarray samples.

Table S8. MSI results of RNA-seq samples by MSIsensor-RNA and PreMSIm.

Table S9. MSI detection performance of MSIsensor-RNA and PreMSIm in RNA-seq

samples.

Table S10. MSI detection performance of MSIsensor-RNA and PreMSIm in

different normalized samples.

Table S11. MSI results of scRNA-seq samples by MSIsensor-RNA.

491   **Supplementary figures:**

492   See in supplementary materials.

494   **Figures:**

495   **Fig. 1**

496
497 **Fig. 2**