Regulatory Role of PKR in Systemic Inflammation-Triggered Neuroinflammation and

its Modulation of Glucose Metabolism and Cognitive Functions in Cholinergic Neurons

Abstract

Systemic inflammation may promote neuroinflammation and neurodegeneration. The double-

stranded RNA-dependent protein kinase (PKR) is a key signaling molecule that regulates

immune responses. This study aims to examine the role of PKR in regulating systemic

inflammation-induced neuroinflammation and cognitive dysfunctions using a laparotomy

mouse model. In the first part, wild-type C57BL/6J and C57BL/6-Tg(CD68-EGFP)1Drg/J

mice were assigned to undergo either laparotomy with sevoflurane anesthesia or sevoflurane

alone to examine effects of systemic inflammation on neuroinflammation and cognition. In

the second part, PKR$^{-/-}$ mice were used to study the role of PKR in modulating laparotomy-

induced systemic inflammation, neuroinflammation, and cognition. For the third part, PKR

was inhibited selectively in cholinergic neurons of ChAT-IRES-Cre-eGFP mice via

intracerebroventricular injection of rAAV-DIO-PKR-K296R. This examined the effects of

inhibiting PKR in cholinergic neurons on glucose metabolism and cognition in the

laparotomy model. Our study revealed that genetic deletion of PKR in mice potently

attenuated the laparotomy-induced peripheral and neural inflammation and cognitive deficits.

Furthermore, inhibiting PKR in the cholinergic neurons rescued the laparotomy-induced

brain glucose hypometabolism and cognitive impairment. Our results demonstrated the

critical role of PKR in regulating neuroinflammation and cognitive dysfunctions in a

peripheral inflammation model. PKR could be a pharmacological target for treating systemic

inflammation-induced neuroinflammation and cognitive dysfunctions.

Keywords

Alzheimer's disease; Laparotomy; Microglia; Neuroinflammation; Protein kinase R;

Systemic inflammation

List of abbreviations

AAV-CON: The AAV-treated group given only sevoflurane

AAV-LAP: The AAV-treated group subjected to laparotomy

AD: Alzheimer's disease

CA: Cornu ammonis

CD68-eGFP mice: C57BL/6-Tg(CD68-EGFP)1Drg/J mice

DAPI: 4', 6-Diamidino-2-phenylindole.

DG: Dentate gyrus

DIO: Double-floxed inverse orientation

dsRNA: Double-stranded RNA

eGFP: Enhanced green fluorescent protein

eIF2α: Eukaryotic initiation factor-2alpha

GC-MS/MS: Gas chromatography-tandem mass spectrometry

Iba1: Ionized calcium-binding adaptor molecule 1

JNK: c-Jun N-terminal kinase

LPS: Lipopolysaccharide

LTP: Long-term potentiation

MAPKs: Mitogen-activated protein kinases

MCP: Monocyte chemoattractant protein

NF-κB: Nuclear factor κB

NOR: Novel object recognition test

OFT: Open field test

PD: Parkinson's disease

PKR: Double-stranded RNA-dependent protein kinase

PND: Perioperative neurocognitive disorders

POD: Postoperative day

SAL-CON: The saline-treated control group given only sevoflurane

SAL-LAP: The saline-treated group subjected to laparotomy

SYM: Spontaneous alternation Y-maze test

TMEM119: Transmembrane protein 119

TRBP: TAR RNA-binding protein

Introduction

Systemic inflammation may promote neuroinflammation and neurodegeneration [1-3]. Neuroinflammation contributes to neurodegenerative disorders, including Alzheimer's disease (AD) [4], Parkinson's disease (PD) [5], and amyotrophic lateral sclerosis [6]. During systemic inflammation, the infiltration of circulating cytokines and macrophages into the brain can activate microglia [7, 8]. Activated microglia release pro-inflammatory cytokines, leading to neuronal injury and death [9]. These inflammatory mediators can further facilitate the recruitment of leukocyte into the brain, perpetuating neuroinflammation [10].

This study utilizes a nonbacterial endotoxin mouse model of laparotomy to address the systemic inflammation triggered by surgery. This laparotomy model is suitable for investigating perioperative neurocognitive disorder (PND) [11]. As a surgical model, laparotomy [12] outperforms other experimental models including tibial fracture surgery [13-15] and injection of bacteria [16] or lipopolysaccharide (LPS) [17] for studying peripheral inflammation because it allows for a longer period of observation up to 2 weeks [16, 17]. Therefore, the laparotomy model is appropriate for the investigation of whether and how neuroinflammation leads to cognitive dysfunctions upon systemic inflammation.

Double-stranded RNA-dependent protein kinase (PKR) is encoded by the *EIF2AK2* gene, regulates mRNA translation, transcription, proliferation, and apoptosis [18]. Beyond its established antiviral functions, PKR act as a critical signaling molecule modulating immune responses. It regulates macrophage activation [19], different inflammatory pathways, and inflammasome formation [20]. PKR can be activated by cytokines, viral double-stranded RNA (dsRNA), bacterial LPS, and

growth factors [20]. PKR activation can regulate activation of various critical inflammatory kinases, including mitogen-activated protein kinases (MAPKs) [21] and inhibitor of nuclear factor-κB (IκB) kinase [22]. Furthermore, PKR also activates and regulates the release of inflammasome-dependent cytokines [23, 24]. Through these myriad functions, PKR critically regulates inflammatory signaling cascades and responses.   In addition to regulating immune responses, PKR plays an essential role in cognition and age-related neurodegeneration. Our laboratory was among the first to reveal that PKR activation and eukaryotic initiation factor-2alpha (eIF2α) phosphorylation are implicated in neuronal apoptosis [25]. The overexpression of wild-type PKR in human neuroblastoma cells could enhance the apoptosis induced by β-amyloid peptides; while overexpression of dominant-negative PKR could attenuate the induced apoptosis. Primary cultured neurons from PKR knockout mice showed greater tolerance to β-amyloid neurotoxicity compared with their wild-type counterpart [25]. In addition, knockout of PKR could result in the enhanced late-phase long-term potentiation (LTP) in hippocampal sections, increased network excitability, as well as improved cognitive functions with the enhancement of learning and memory in mice [26]. All these studies suggested the role of PKR in modulating cognitive functions and neurodegeneration.

Considering the dual functions of PKR in modulating immune responses and cognition, we hypothesize that PKR regulates the transduction of systemic inflammation into neuroinflammation, and PKR in cholinergic neurons can modulate cognitive functions. This project aims at investigating whether PKR can be a pharmacological target for preventing systemic inflammation-triggered neuroinflammation and cognitive deficits.

Methods

Animal husbandry

Male wild-type C57BL/6J, C57BL/6-Tg(CD68-EGFP)1Drg/J (referred to herein as CD68-eGFP mice), PKR$^{-/-}$, and ChAT-IRES-Cre-eGFP mice (3 months) were obtained from the Centre for Comparative Medicine Research of the University of Hong Kong. This center is fully accredited by the Association for Assessment and Accreditation of Laboratory Animal Care. The mice were housed in cages under controlled temperature (20°C – 22°C), humidity (50% – 60%), with a 12 hour light/dark cycle and ad libitum access to food and water. Mice were allowed to acclimate for 1 week prior to experiment.

Ethics statement

All experiments were approved by the Department of Health of the Hong Kong SAR government and the Committee on the Use of Live Animals in Teaching and Research of the University of Hong Kong (#4911-18 and #5819-21).

Experimental protocols

Laparotomy was used as a model to examine neuroimmune responses induced by systemic inflammation. The experimental workflow is summarized in Fig. S1. In part 1, male wild-type C57BL/6J and CD68-eGFP mice were randomized into two groups: laparotomy with sevoflurane anesthesia and control group with sevoflurane anesthesia alone. In part 2, the PKR$^{-/-}$ mice were exposed to laparotomy with

sevoflurane anesthesia or sevoflurane anesthesia only to examine the role of PKR in regulating systemic inflammation-triggered neuroinflammation and cognitive deficits. In part 3, rAAV-DIO-PKR-K296R was intracerebroventricularly injected into the right lateral ventricle of ChAT-IRES-Cre-eGFP mice to suppress PKR activation specifically in cholinergic neurons. Mice were then subjected to laparotomy with sevoflurane or sevoflurane anesthesia alone. This allowed examination of the regulatory effects of blocking PKR in cholinergic neurons on glucose metabolism and cognitive function in the laparotomy model.

Laparotomy

For anesthetization, the animal was placed in a chamber with 5% sevoflurane (Sevorane™, Abbott, Switzerland) and transferred to a nose mask with anesthesia maintained at 3% sevoflurane with a gas flow of 800 ml/min using an inhalation anesthesia machine (Harvard, US). The laparotomy procedure followed previous descriptions with modifications [12]. Hair was removed from the surgical site, and the skin was disinfected. A 2 cm longitudinal midline cut was made at 0.5 cm below the lower right rib. A small intestinal segment of approximately 10 cm length was temporarily exteriorized and subjected to 2 minutes of gentle manipulation before being replaced in the abdomen. Abdominal muscle and skin suture was conducted using 5-0 vicryl and nylon suture respectively (Ethicon, USA).

Stereotaxic injection

Mice were anesthetized by intraperitoneal injection of a ketamine/xylazine mixture (80 mg/kg and 8 mg/kg, respectively). Prior to surgery, hair was removed

from the surgical site, and the skin was disinfected. The head was then fixed onto a stereotaxic frame (Narishige, Japan). A 1.5 cm straight midline incision was made to expose the skull. Either rAAV-DIO-PKR-K296R (packaged in AAV serotype 6) (2 µl; 1×10^{13} vg/mL) or 0.9% saline was injected intracerebroventricularly into right lateral ventricle (antero-posterior: -0.6 mm, medial-lateral: -1.3 mm, dorsal-ventral: -3.1 mm) at a flow rate of 0.5 µL/min using a Hamilton syringe, 33-gauge needle (7803-05, Hamilton, USA). Following the injection, the needle was retained in situ for 5 min allowing the solution to diffuse and avoiding solution backflow. The incision site was sealed with a nonabsorbable nylon suture (5-0, PS-2; Ethicon, USA).

Open field test

Open field test (OFT) is adopted to assess locomotor activity, exploratory behavior and anxiety in mice based on their innate propensity to avoid open spaces [27]. Mice were placed in a 40 cm square arena and allowed to freely explore for 10 min. Total distance traveled and time spent exploring central area were quantified as indices of locomotor activity and anxiety.

Novel object recognition test

Novel object recognition test (NOR) assesses recognition memory in mice based on their innate preference for novelty [28]. In the training session, mice were first allowed to explore an open field arena with two identical objects (A + A) for 10 min. After 24 hours, mice were allowed to explore the familiar arena with a novel object replacing one of the identical objects (A + B). Basically, driven by curiosity, the mice would spend more time exploring the novel object rather than the familiar

one. The discrimination index, calculated as a ratio of time exploring the novel verus familiar object, can be regarded as an index of recognition memory.

Spontaneous alternation Y-maze test

Spontaneous alternation Y-maze (SYM) test assesses short-term spatial working memory in rodents. Mice are allowed to explore the maze for 5 min [29]. A spontaneous alternation involves three successive entries into a new arm rather than returning to an earlier one. Driven by curiosity, the mice will tend to explore the new arm rather than the previously visited arm. The percentage of alternation was determined by dividing the number of spontaneous alternations by the total arm entries, then multiplying by 100. The percentage of alternation can be the indicator of short-term spatial working memory.

Puzzle box test

A puzzle box test was a five-day protocol modified by our laboratory [28] based on the previous published method [30]. It is designed to assess rodent's native problem-solving ability, short-term and long-term memory. The puzzle box was consisted of two main sections: an open-field area (600 × 280 mm) and an adjacent sheltered goal-box area (600 × 280 mm). Mice could enter the sheltered goal-box arena through a small underpass between these two compartments. Mice were required to complete escape tasks of increasing difficulty within a fixed period of time (Table S1). The first four days involved three trials per day, while the final day comprised a single trial. Each obstruction includes three trials; the initial two taking place on the same day, with the third trial conducted on the following day. Three trials

were performed: the first evaluated problem-solving skills; the second assessed short-term memory, and the third investigated long-term memory.

Arrangement of behavioral tests

A series of cognitive behavioral tests was conducted based on the following schedule. For minimized fatigue and interference, behavioral tests were divided across the study. A group of mice underwent OFT on postoperative day (POD) 10, followed by NOR on POD 11–12. A second group of mice underwent OFT on POD 10, followed by SYM on POD 11. The third cohort completed OFT on POD 9 and the puzzle box test on POD 10–14.

RNA extraction and quantitative real-time RT-PCR

Mice were sacrificed by carbon dioxide asphyxiation and decapitation, in line with the protocols recommended by American Veterinary Medical Association and then followed by transcardial perfusion with ice-cold saline. Frontal cortex, hippocampus, and liver were harvested, snap-frozen, and kept at −80°C. Total RNA was isolated from samples using the RNAiso Plus (TAKARA, Japan). Detailed protocols were described in Supplementary Methods. Quantitative real-time RT-PCR was performed with TB Green® Premix Ex Taq™ II (TAKARA, Japan) using a Bio-Rad CFX 96 real-time PCR detection system (Bio-Rad, USA) in triplicate. Primer sequences utilized for amplification are provided in Table S2. Relative expression was determined using the comparative CT method, normalizing the target gene mRNA to GAPDH.

228   *Immunohistochemical staining and confocal microscopy*

229   The processes for cryosection preparation described in the Supplementary

Methods were similar to those described previously for human brain sections [31].

Prior to confocal imaging, brain slices were pretreated with 10% goat serum blocking

solution, followed by overnight incubation at 4°C with primary antibodies listed in

Table S3, followed by 1h secondary antibody staining (Alexa Fluor, 1:400,

Invitrogen) at room temperature and co-staining with 4□, 6-diamidino-2-phenylindole

(DAPI). Confocal images were acquired using a LSM800 confocal microscope (Carl

Zeiss, Germany) with 10× or 20× objective or 40× oil immersion objective using Zen

Blue software. The frame size was set to 1024×1024 pixels, the bit depths to 16-bit,

averaging to 2×, and pinhole setting at 1 airy unit. Z-stack images were acquired,

followed by orthogonal projections from Z-stack images using Zen Blue software.

Image analysis was performed using ImageJ software (National Institute of Health,

USA).

Flow cytometry

244   The frontal cortex and hippocampus of CD68-eGFP mice were collected,

245   processed through homogenization and filtration to obtain single-cell suspensions,

246   which were then stained with DAPI. Detailed protocol was described in the

247   Supplementary Methods. Cellular fluorescence was measured using an Agilent

248   NovoCyte Quanteon analyzer (Agilent, USA).

250   *Targeted metabolomics by GC-MS/MS*

251    Samples were obtained from the collected frontal cortex and prepared using

252 the protocol described in the Supplementary Methods. Targeted metabolomics was

253 performed on central carbon metabolism using gas chromatography-tandem mass

254 spectrometry (GC-MS/MS) on Agilent 7890B GC-Agilent 7010 Triple Quadrupole

255 Mass Spectrometer system (Agilent, USA). The experimental parameters were stated

256 in the Supplementary Methods. Data were normalized by median and log-

257 transformed. Heat map and hierarchical clustering were conducted based on Euclidian

258 distance using Ward's algorithm. Further principal component analysis (PCA) and

259 partial least squares discriminant analysis (PLS-DA) were further performed using

260 Metaboanalyst 5.0.

262 *Statistical analysis*

263    Unless otherwise specified, all statistical analyses were performed on Prism

264 v8.0c (GraphPad Software, USA). Data were expressed as means ± SEM. The

265 normality of data were assessed using the D'Agostino-Pearson normality test.

266 Unpaired two-tailed Student's t-test at $p < 0.05$ was applied for comparing two group

267 to determine the statistical significance of treatment effects. Interactions between

268 mouse strain and treatment were assessed using two-way ANOVA with Bonferroni

269 post hoc test. Statistical significance was set at $p < 0.05$.

271 **Results**

272 *Laparotomy induced peripheral and neural inflammation in wild-type mice*

273    To investigate the systemic and neural immune responses following the

274 laparotomy challenge, we initially performed laparotomy in the male C57BL/6J mice.

Four hours following surgery, IL-1β and monocyte chemoattractant protein (MCP)-1 mRNA expression were enhanced in the liver of the laparotomy group when compared with that in the control counterpart ($p < 0.05$; Fig. 1a). IL-1β expression was also significantly upregulated in the hippocampus following the laparotomy challenge ($p < 0.05$; Fig. 1a). On POD 1, no significant differences in IL-1β, IL-6, MCP-1, and TNF-α mRNA expression in the liver were found between the laparotomy and control groups (Fig. 1b). This finding implied that the induced systemic inflammation is relatively transient. However, laparotomy significantly increased IL-1β and IL-6 expression in the frontal cortex and hippocampus versus controls ($p < 0.05$; Fig. 1b), indicating ongoing, persistent neuroimmune responses.

To further study the effects of laparotomy on neuroinflammation, we utilized transgenic CD68-eGFP mice for visualizing microglia/macrophage dynamics. The CD68-eGFP mice expressed high levels of enhanced green fluorescent protein (eGFP) in macrophages and activated microglia, which aided in identifying activated microglia and macrophages. Brain sections were immunostained with antibodies against eGFP, TMEM119 (transmembrane protein 119: a microglia-specific marker), and CD105 (endoglin: a vascular endothelial marker). On POD 14, more CD68-EGFP-positive cells were observed in the frontal cortex ($p < 0.01$) and hippocampus (including cornu ammonis (CA)1 and CA3 ($p < 0.05$), as well as dentate gyrus (DG) ($p < 0.01$)) of the laparotomy mice compared with those of the control counterpart (Fig. S2, S3). Furthermore, an increased number of vessel-associated macrophages (CD68-eGFP$^+$, TMEM119$^-$ cells associated with the blood vessels) in the frontal cortex was detected following laparotomy ($p < 0.05$; Fig. S2a, b). This result suggested the activation of microglia/potential infiltration of macrophages following laparotomy. Flow cytometry also demonstrated elevated numbers of CD68-eGFP-

positive cells in both frontal cortex and hippocampus on POD 14 when compared to controls ($p < 0.05$; Fig. 1c–f), suggesting laparotomy induced the activation of microglia/potential infiltration of macrophages in the brain.

Brain sections from the CD68-eGFP mice were immunostained with a microglial activation marker, ionized calcium-binding adaptor molecule 1 (Iba1), to further confirm whether laparotomy induces activation of microglia. More Iba1$^+$ cells were found in both frontal cortex and hippocampus of the laparotomy group when compared to controls (Frontal cortex: $p < 0.05$; CA1 and CA3: $p < 0.001$; DG: $p < 0.01$; Fig. 2). Increased fluorescence intensity was also observed in the hippocampus following laparotomy ($p < 0.05$; Fig. 2). Microglial morphology, an important indicator of neuroinflammation, was further quantified by skeleton analysis [32]. Reduced total endpoints or/and shorter process length per microglial cell were observed in the frontal cortex and the CA1 and DG of the hippocampus of the laparotomy group compared with those of the control counterpart ($p < 0.05$; Fig. 2b, d and h). This finding revealed that microglia were deramified following laparotomy.

Changes of body weight and cognitive performance in wild-type mice following laparotomy

Body weights were recorded throughout the experiment. Wild-type laparotomy mice exhibited notable weight loss in the first 6 days following surgery ($p < 0.05$; Fig. 3a) but no remarkable decreases thereafter.

A series of behavioral tests was performed to study whether laparotomy affects cognitive functions in the wild-type mice. During the OFT, total distance

traveled and central area exploration time were similar between laparotomy and

control groups (Fig. 3b, c). This finding implied that laparotomy did not affect the

locomotor activity, exploration, or anxiety. NOR then evaluated recognition memory

according to the innate preference of mice for novelty [28]. On the first day of NOR,

both groups of mice spent comparable time exploring the identical objects (Fig. 3d).

On day 2 of NOR, impaired recognition memory with lower discrimination index was

observed in the laparotomy group versus the controls ($p < 0.05$; Fig. 3e). As indicated

by the reduced percentage of alternation, impaired spatial working memory was

detected in the laparotomy group when compared to the control ($p < 0.05$; Fig. 3g)

during SYM with similar total number of entries (Fig. 3f). Puzzle box test was

adopted to examine the rodent's native problem-solving ability during trial (T)2, T5,

T8, and T11. Additionally, evaluations at the T3, T6, T9, and T12 provided insights

into their short-term memory recall, while the T4, T7, T10, and T13 were indicative

of their long-term memory function [28]. In the habituation task (T1), both groups

spent similar time to complete the escape task (Fig. 3h). Impaired problem-solving

abilities requiring a longer period of time to complete the escape task in T2, T5, and

T11 were detected in the laparotomy group compared to controls ($p < 0.05$; Fig. 3i).

Furthermore, the laparotomy group showed reduced short-term (with increased

latency in T12) ($p < 0.05$; Fig. 3j) and long-term (with increased latency in T7, T10,

and T13) memories when compared to controls (T7 and T13: $p < 0.05$; T10: $p < 0.01$;

Fig. 3k).

Knockout of PKR in mice potently downregulated the up-regulated pro-inflammatory cytokine expressions induced by laparotomy and ameliorated the microglial activation following laparotomy

Given that PKR regulates immune responses, we assessed whether genetic deletion of PKR could abrogate laparotomy-induced inflammation. IL-1β expression was significantly elevated in liver and hippocampus of the wild-type mice at four hours post-surgery ($p < 0.05$; Fig. 4a) but not in the $PKR^{-/-}$ mice after the surgery compared with that in the controls (Fig. 4a). Meanwhile, expression of IL-1β in liver and frontal cortex was significantly reduced in the $PKR^{-/-}$ laparotomy group when compared to the wild-type laparotomy group ($p < 0.05$; Fig. 4a). With regard to MCP-1 expression in the liver, a significant increase was observed in the wild-type mice after laparotomy ($p < 0.05$; Fig. 4a) while not in the $PKR^{-/-}$ mice (Fig. 4a). Hepatic TNF-α expression was significantly elevated in $PKR^{-/-}$ mice after laparotomy compared to $PKR^{-/-}$ controls ($p < 0.05$; Fig. 4a); wild-type mice showed no such difference.

On POD 1, the wild-type laparotomy group revealed a substantial increase in IL-1β and IL-6 mRNA expression in frontal cortex and hippocampus, compared to the wild-type controls ($p < 0.05$; Fig. 4b); no such increases were detected in the $PKR^{-/-}$ laparotomy group. Meanwhile, the $PKR^{-/-}$ laparotomy group showed significantly lower IL-6 expression in the frontal cortex versus the wild-type laparotomy group ($p < 0.01$; Fig. 4b). Between the surgical and control groups for both strains of mice, no remarkable differences in IL-1β, IL-6, MCP-1, and TNF-α mRNA expression were detected in the liver (Fig. 4b) and no remarkable alterations in MCP-1 and TNF-α expression were found in frontal cortex and hippocampus (Fig. 4b).

We then further investigated whether knockout of PKR could inhibit

activation of microglia in the laparotomy model. No observable difference in the

number and fluorescence intensity of Iba1$^+$ cells in the frontal cortex (Fig. 5a and b)

and hippocampus was found between the PKR$^{-/-}$ laparotomy group and its control

counterpart (Fig. 5c–h). Concurrently, laparotomy did not induce morphological

changes (total number of endpoints and summed branch lengths) in microglia in the

frontal cortex (Fig. 5a and b) and hippocampus (Fig. 5c–h) of PKR$^{-/-}$ mice compared

with those in the control. The PKR$^{-/-}$ mice showed ameliorated neuroimmune

responses compared to wild-type mice, indicating the critical role of PKR in

regulating neuroinflammation following laparotomy.

Knockout of PKR in mice ameliorated the cognitive deficits induced by laparotomy

Regardless of the mouse strain, laparotomy resulted in a considerable body

weight loss within 3 days ($p < 0.05$; Fig. 6a). PKR$^{-/-}$ mice restored weight by day 6,

earlier than that of the wild-type mice (on day 9). In the laparotomy group, neither

strain of mice lost significant weight after day 9 (Fig. 6a). Meanwhile, laparotomy did

not affect the locomotor activity (total distance travelled) and anxiety level (central

duration) of the PKR$^{-/-}$ mice (Fig. 6b and c). On NOR day 1, both strains spent

comparable time exploring the identical objects (Fig. 6d). On NOR day 2 with a novel

object, laparotomy impaired the recognition memory (reduced discrimination index)

of the wild-type mice ($p < 0.05$; Fig. 6e) but not that of the PKR$^{-/-}$ mice (Fig. 6e). In

the SYM, laparotomy impaired the spatial recognition memory (reduced percentage

of alternation) of the wild-type mice ($p < 0.05$; Fig. 6g) but not that of the PKR$^{-/-}$ mice

(Fig. 6g). No notable difference in the total entries was detected between the surgery and control groups (Fig. 6f).

The puzzle box test was conducted to further assess the rodent's problem-solving skill and short-term and long-term memories. Throughout habituation, no notable difference was detected between the laparotomy and control groups, regardless of the mouse strain (Fig. 6h). Between the mouse strains, the PKR$^{-/-}$ mice demonstrated greater latency to enter the goal zone than the wild-type mice ($p < 0.01$; Fig. 6h). In terms of problem-solving ability, the PKR$^{-/-}$ laparotomy mice performed similarly to their PKR$^{-/-}$ control counterparts (Fig. 6i). This finding differed from that observed in the wild-type laparotomy mice who took longer to solve task T2, T5, and T11 ($p < 0.05$; Fig. 6i). In terms of native problem-solving ability, significant differences were also detected between the PKR$^{-/-}$ and wild-type mice ($p < 0.01$; Fig. 6i). In T2, the PKR$^{-/-}$ laparotomy mice demonstrated greater latency when compared to the wild-type laparotomy group ($p < 0.05$; Fig. 6i); a similar trend was also detected in T11 between the PKR$^{-/-}$ controls and their wild-type control counterpart ($p < 0.05$; Fig. 6i). Furthermore, laparotomy did not impair the short- and long-term memories of the PKR$^{-/-}$ mice, and the wild-type laparotomy group showed significant impairment (T7, T12, and T13: $p < 0.05$; T10: $p < 0.01$; Fig. 6j, k). This result suggested that genetic deletion of PKR could ameliorate the cognitive deficits induced by laparotomy.

Inhibition of PKR in cholinergic neurons ameliorated laparotomy-induced cognitive dysfunctions

rAAV-DIO-PKR-K296R was intracerebroventricularly injected into the right lateral ventricle of the ChAT-IRES-Cre-eGFP mice to inhibit activation of PKR in cholinergic neurons. rAAV expresses a dominant negative form of PKR, that is, PKR-K296R, under the regulation of a double-floxed inverse orientation (DIO) switch (Fig. S4a). At 4 weeks after the stereotaxic injection of rAAV-DIO-PKR-K296R, the fluorescence intensity of p-PKR (T446) signal and the fluorescence ratio of p-PKR to the eGFP signal were reduced in the rAAV-DIO-PKR-K296R-treated group when compared to the saline-treated control ($p < 0.05$; Fig. S4b, c). Therefore, in ChAT-IRES-Cre-eGFP mice, the expression of PKR-K296R effectively suppressed the activation (phosphorylation) of PKR.

At 4 weeks following stereotaxic injection, mice were randomized into four groups: saline-treated control group given only sevoflurane (SAL-CON), saline-treated group subjected to laparotomy (SAL-LAP), AAV-treated group given only sevoflurane (AAV-CON), and AAV-treated group subjected to laparotomy (AAV-LAP). Within 3 days after laparotomy, the SAL-LAP and AAV-LAP groups lost significant weight ($p < 0.05$; Fig. 7a), similar to the wild-type and PKR$^{-/-}$ mice. On POD 6, their body weight returned to normal, and no remarkable difference was detected across all the groups (Fig. 7a).

Different behavioral tests were conducted to investigate the effects of inhibiting PKR in cholinergic neurons on cognition following laparotomy. rAAV-DIO-PKR-K296R injection and laparotomy did not affect locomotor activity (total distance travelled) and anxiety level (central duration) (Fig. 7b, c). In the SYM test, impaired spatial memory with lower percentage of alternation was detected in the SAL-LAP group when compared to the SAL-CON ($p < 0.05$; Fig. 7e); AAV-treated

mice showed no such difference. No remarkable difference in the total number of entries was detected among all groups (Fig. 7d). Our findings suggested that inhibiting PKR in cholinergic neurons could alleviate the spatial working memory deficits induced by laparotomy.

Considering the habituation task (T1) of the puzzle box test, all the groups of mice completed the escape task in about the same amount of time (Fig. 7f). In terms of problem-solving ability, contrasting behaviors were observed. In the T2 task, the SAL-LAP group demonstrated increased latency when compared to the SAL-CON ($p < 0.001$; Fig. 7g); no notable difference was found in the AAV-treated group (Fig. 7g). Meanwhile, the AAV-LAP group solved the T2 task faster than the SAL-LAP group ($p < 0.0001$; Fig. 7g). This result revealed that inhibiting PKR in cholinergic neurons could improve the cognitive functions of mice in the T2 task following laparotomy. In the T5 and T8 task, however, the AAV-treated and saline-treated mice showed increased latency time following laparotomy when compared to their control counterparts ($p < 0.05$; Fig. 7g). With respect to T11 task, the AAV-LAP group demonstrated a significant prolonged latency when compared to the AAV-CON ($p < 0.05$; Fig. 7g). Meanwhile, the AAV-CON group demonstrated better problem-solving skills than the SAL-CON ($p < 0.05$; Fig. 7g).

Regarding short-term memory, consistent results were observed in the T3 and T6 tasks (Fig. 7h). When compared with the SAL-CON group, the SAL-LAP mice showed impaired short-term memory with increased latency to solve the T3 ($p < 0.001$; Fig. 7h) and T6 tasks ($p < 0.05$; Fig. 7h). Such differences were not detected in the AAV-treated groups. In the T3 task, the AAV-LAP group escaped faster than the SAL-LAP group ($p < 0.05$; Fig. 7h). This finding indicated that inhibiting PKR in

cholinergic neurons helped alleviate the short-term memory deficits induced by laparotomy. However, for the T9 task, the AAV-treated and saline-treated mice showed increased latency following laparotomy when compared to the controls ($p < 0.05$; Fig. 7h). Moreover, the AAV-treated group demonstrated reduced latency for the T12 task compared with their saline-treated counterpart ($p < 0.05$; Fig. 7h). Therefore, inhibition of PKR in cholinergic neurons could improve short-term memory.

For long-term memory, contrasting behaviors were observed in the T4 and T7 tasks. In the T4 task, the SAL-LAP group showed impairment ($p < 0.05$; Fig. 7i); AAV-treated mice showed no such difference (Fig. 7i). In T7 task, a noticeable change was observed only in the AAV-treated groups ($p < 0.05$; Fig. 7i) but not in the saline-treated groups (Fig. 7i). No significant difference for all the tasks testing long-term memory (T4, T7, T10, and T13 tasks) was observed between the AAV-treated and SAL-treated groups. Therefore, no concrete conclusions concerning the influence of PKR inhibition in cholinergic neurons on long-term memory could be drawn.

Inhibition of PKR in cholinergic neurons alters glucose metabolism in the frontal cortex in a laparotomy model

MS-based targeted metabolomics was performed to examine the effects of PKR inhibition in cholinergic neurons on brain glucose metabolism. Distinct clustering was observed between the SAL-LAP and AAV-CON groups (Fig. 8a, b). The heatmap depicted the changes of quantified metabolites including tricarboxylic acids and glycolysis intermediates in the frontal cortex across all the groups (Fig. 8c).

Laparotomy significantly reduced the level of phosphoenolpyruvate in the SAL-LAP

group when compared to the control counterpart ($p < 0.05$; Fig. 8e). However, no

notable difference in phosphoenolpyruvate level was detected in the AAV-treated

group following laparotomy. Therefore, inhibiting PKR in cholinergic neurons could

ameliorate the laparotomy-induced changes in phosphoenolpyruvate level. Similarly,

laparotomy reduced the succinate level in the SAL-treated mice ($p < 0.05$; Fig. 8f) but

not in the AAV-treated group (Fig. 8f). No notable difference in glucose levels was

detected among all the groups (Fig. 8d). Apart from the laparotomy-induced changes

in glycolysis intermediates, inhibition of PKR in cholinergic neurons significantly

elevated the level of 6-phosphogluconate in the AAV-CON group when compared to

that in the SAL-CON group ($p < 0.01$; Fig. 8g), suggesting that PKR inhibition in

cholinergic neurons could alter brain energy metabolism.

Discussion

Neuroinflammation is a major cause of cognitive impairment in PND. In this

study, we employed a clinically relevant mouse model of laparotomy to induce

peripheral and neural inflammation and further demonstrate the regulatory role of

PKR in systemic inflammation-induced neuroinflammation and cognitive

dysfunctions.

Regarding the laparotomy model, systemic and neural immune responses

developed rapidly after the surgery. In theory, laparotomy results in peritoneal air

exposure [33, 34], tissue injury, and gut mucosa barrier damage, resulting in increased

inflammatory mediators and gut bacteria/endotoxin translocation to the circulation,

contributing to systemic inflammation [35]. Our study found a significant elevation in

the cytokine mRNA expression of liver (IL-1β and MCP-1) and brain tissues (IL-1β) at four hours after laparotomy (Fig. 1a). On POD 1, laparotomy triggered a significant upregulation of IL-1β and IL-6 expression in frontal cortex and hippocampus (Fig. 1b). IL-1β plays an important role in neuroinflammation and pathogenesis of AD [36] and PD [37]. Through activating nuclear factor κB (NF-κB), IL-1β promotes the upregulation of proinflammatory cytokines including IL-6 and TNF-α [38]. This finding may partially explain the increased IL-6 expression in the brain of the laparotomy group on POD 1 (Fig. 1b). However, the induced systemic inflammation is relatively transient, with proinflammatory cytokine expression levels in the liver returning to normal on POD 1 (Fig. 1b). Therefore, a transient systemic inflammation is already sufficient to induce neuroinflammation as evidenced by the persistent microgliosis revealed by flow cytometry (Figs. 1c–f) and confocal imaging (Fig. 2) on POD 14. Our findings are also consistent with a prior study that demonstrated significant microglial activation in frontal cortex and hippocampus following laparotomy [12]. Beyond the microglial activation, cognitive deficits (Fig. 3) were also detected in the wild-type laparotomy model mice. Therefore, the laparotomy model is a viable and relevant paradigm for studying systemic inflammation-triggered neuroimmune response and cognitive dysfunction.

Our study is the first to show that genetic deletion of PKR could alleviate laparotomy-induced peripheral and neural inflammation, as well as downregulate the increased expression of IL-1β and IL-6 in liver, frontal cortex, and hippocampus following laparotomy (Fig. 4a and b). Since PKR plays a critical role in IL-1β activation, PKR knockout can help prevent pro-IL-1β from cleaving into IL-1β. These findings were consistent with other previous studies showing that PKR could attenuate inflammation by regulating inflammasome activation [39] and various

inflammatory pathways including TNF [40], p38 MAPK [21], and NF-κB/IκB

signaling pathway [41]. PKR inhibition prevented IκB and NF-κB activation and

decreased IL-1β expression in primary murine mixed neuron–astrocyte–microglia co-

culture model [41] and RAW264.7 macrophages [42]. Subsequently, the reduced IL-

1β expression further downregulated IL-6 expression by preventing the activation of

NF-κB. Four hours following surgery, a significant elevation was found in TNF-α

expression in the liver of the PKR$^{-/-}$ laparotomy group but not in their wild-type

counterpart (Fig. 4a). However, no notable difference was detected between the PKR$^{-/-}$

and wild-type laparotomy groups. Similar findings were also observed in our earlier

investigation, which demonstrated a significant elevation in TNF-α expression in the

hippocampus of the PKR$^{-/-}$ mice after *Escherichia coli* challenge but not in their wild-

type control counterpart [16]. Indeed, PKR is involved in TNF-dependent activation

of c-Jun N-terminal kinase (JNK), NF-kB and protein kinase B. Meanwhile, knockout

of PKR could abrogate the TNF-induced activation of the above signaling pathways.

Therefore, the PKR$^{-/-}$ laparotomy mice may tend to highly express TNF-α to activate

the downstream pathways. Our study has taken a significant step beyond previous

research. Although previous works showed that genetic deletion of PKR in mice can

prevent activation of microglia following intraperitoneal injection of LPS [43], we

demonstrated that PKR also alleviates neuroinflammation induced by sterile systemic

inflammation as evidenced by the suppressed laparotomy-induced microglial

activation in frontal cortex and hippocampus of PKR$^{-/-}$ mice (Fig. 4, 5). Therefore,

PKR regulates the activation of microglia/macrophages. Since PKR is critical for

LPS-induced inducible nitric oxide synthase expression [44], it is responsible for the

macrophage inflammatory response by regulating the production of a

proinflammatory factor, nitric oxide [45]. Moreover, PKR is necessary for the

dsRNA-induced activation of macrophages [42]. All the aforementioned studies supported the role of PKR in activation of microglia/macrophages. Furthermore, knockout of PKR in mice could rescue the laparotomy-triggered impairment of problem-solving ability, spatial working memory, and short-term and long-term memories in mice (Figs. 6e, g, and i–k).

Another novel finding of this project is that inhibition of PKR in cholinergic neurons could rescue the laparotomy-induced cognitive deficits. The cholinergic system is selected for investigation because it is not only essential for cognition but also regulates inflammation through the cholinergic anti-inflammatory pathway [46], which connects systemic inflammation and neuroinflammation. This would elucidate the neuron-specific effects of PKR on cognition during inflammation beyond its immune regulatory roles. This study also demonstrated that inhibition of PKR in cholinergic neurons could alleviate the laparotomy-induced impairment of short-term and spatial working memory (Fig. 7e and h). A previous study showed that PKR inhibition could improve cortex-dependent memory consolidation in rats and mice [47]. Meanwhile, PKR$^{-/-}$ mice demonstrated enhanced conditioned taste aversion memory when compared with their wild-type control [47]; while PKR activation could impair hippocampal late-phase LTP and memory consolidation in PKR recombinant mice [48]. All the abovementioned studies supported the crucial role of PKR in regulating cognitive functions. Our research highlights PKR as a potential key target in treating inflammation-induced cognitive deficits, opening up promising possibilities for future therapeutic approaches focusing on PKR modulation.

We observed that laparotomy could lead to brain glucose hypometabolism with reduction in glycolysis and citric acid cycle intermediates (phosphoenolpyruvate and succinate) (Fig. 8), which is a hallmark of neurodegenerative diseases including

AD and PD in preclinical stages [49, 50]. Progressive reduction in brain glucose

metabolism positively correlates with amyloid load and cognitive decline in patients

with early AD or PD suffering from mild cognitive impairment [50-52]. AD patients

showed reduced concentrations of glycolytic intermediates including

phosphoenolpyruvate in cerebrospinal fluid [53]. By contrast, administering succinate

into the brain by retrodialysis could improve the brain reduced/oxidized nicotinamide

adenine dinucleotide redox state [54] and brain energy metabolism in patients

suffering from traumatic brain injury [55]. All of the abovementioned studies

established a close association between brain glucose metabolism and cognition.

Our study unraveled the novel role of PKR in regulating brain glucose

metabolism in the laparotomy model. Blocking PKR in cholinergic neurons could

rescue the laparotomy-induced alterations in glucose metabolism in the frontal cortex

on POD 14 (Fig. 8e and f). PKR regulates glucose homeostasis [56] and induces

insulin resistance [57, 58]. Increased eIF2α phosphorylation is a hallmark of insulin

resistance and obesity [57-59]. Under metabolic and obesity-induced stress, PKR

could form a complex with TAR RNA-binding protein (TRBP), activating JNK [56].

Inhibiting TRBP in the liver could improve insulin sensitivity, glucose metabolism,

and alleviate inflammation [56]. The interaction between PKR and TRBP links

metabolism to inflammatory signaling regulation. However, no prior study has

examined the effect of blocking PKR in brain on glucose metabolism. Our findings

supported that PKR in cholinergic neurons play a critical role in modulating glucose

metabolism. Further study to unveil the underlying regulatory networks is warranted.

Conclusions

The laparotomy model was a stable and viable model for studying

neuroimmune responses triggered by systemic inflammation as evidenced by

microgliosis and cognitive dysfunction. By genetic deletion of PKR, we demonstrated

the critical role of PKR in regulating peripheral and neural inflammation. Knockout of

PKR could help alleviate neuroimmune responses and cognitive impairment induced

by laparotomy. Furthermore, inhibition of PKR in the cholinergic neurons of the

laparotomy mice demonstrated a neuroprotective role and improved the glucose

metabolism and cognitive functions. On the basis of our combined findings, PKR

could be a pharmacological target for treating systemic inflammation-induced

neuroinflammation and cognitive deficits.

Conflict of Interest

Authors declare no competing interests.

Ethics approval and consent to participate

All experiments were approved by the Department of Health of the Hong

Kong SAR government and the CULATR of the University of Hong Kong (#4911-18

and #5819-21).

Availability of data and materials

The online version contains supplementary material. All raw data are available

at the HKU data repository.

Funding

This study is supported by General Research Fund (17123217).

References

1.   Hoogland I, Houbolt C, van Westerloo DJ, van Gool WA, van de Beek D. Systemic

inflammation and microglial activation: systematic review of animal experiments. J.

Neuroinflammation. 2015;12(1):1-13.

2.   Lopez‐Rodriguez AB, Hennessy E, Murray CL, Nazmi A, Delaney HJ, Healy D, et

al. Acute systemic inflammation exacerbates neuroinflammation in Alzheimer's disease:

IL‐1β drives amplified responses in primed astrocytes and neuronal network dysfunction.

Alzheimers. Dement. 2021;17(10):1735-55.

3.   Wang RPH, Ho YS, Leung WK, Goto T, Chang RCC. Systemic inflammation linking

chronic periodontitis to cognitive decline. Brain Behav. Immun. 2019;81:63-73.

4.   Heneka MT, Carson MJ, El Khoury J, Landreth GE, Brosseron F, Feinstein DL, et al.

Neuroinflammation in Alzheimer's disease. Lancet Neurol. 2015;14(4):388-405.

5.   Hirsch EC, Hunot S. Neuroinflammation in Parkinson's disease: a target for

neuroprotection? Lancet Neurol. 2009;8(4):382-97.

6.   Philips T, Robberecht W. Neuroinflammation in amyotrophic lateral sclerosis: role of

glial activation in motor neuron disease. Lancet Neurol. 2011;10(3):253-63.

7.   Fu R, Shen Q, Xu P, Luo JJ, Tang Y. Phagocytosis of microglia in the central nervous

system diseases. Mol. Neurobiol. 2014;49(3):1422-34.

8.   Kempuraj D, Thangavel R, Selvakumar GP, Zaheer S, Ahmed ME, Raikwar SP, et al.

Brain and Peripheral Atypical Inflammatory Mediators Potentiate Neuroinflammation and

Neurodegeneration. Front. Cell. Neurosci. 2017;11.

9.   Streit WJ, Mrak RE, Griffin WST. Microglia and neuroinflammation: a pathological

perspective. J. Neuroinflammation. 2004;1(1):14.

10.   Varvel NH, Neher JJ, Bosch A, Wang W, Ransohoff RM, Miller RJ, et al. Infiltrating

monocytes promote brain inflammation and exacerbate neuronal damage after status

epilepticus. Proc. Natl. Acad. Sci. U.S.A. 2016;113(38):E5665.

11.   Evered L, Silbert B, Knopman DS, Scott DA, DeKosky ST, Rasmussen LS, et al.

Recommendations for the Nomenclature of Cognitive Change Associated with Anaesthesia

and Surgery-2018. Anesthesiology. 2018;129(5):872-9.

12.   Huang C, Irwin MG, Wong GTC, Chang RCC. Evidence of the impact of systemic

inflammation on neuroinflammation from a non-bacterial endotoxin animal model. J.

Neuroinflammation. 2018;15(1):147.

13.   Lu Y, Xu X, Dong R, Sun L, Chen L, Zhang Z, et al. MicroRNA-181b-5p attenuates

early postoperative cognitive dysfunction by suppressing hippocampal neuroinflammation in

mice. Cytokine. 2019;120:41-53.

14.   Xu J, Dong H, Qian Q, Zhang X, Wang Y, Jin W, et al. Astrocyte-derived CCL2

participates in surgery-induced cognitive dysfunction and neuroinflammation via evoking

microglia activation. Behav. Brain Res. 2017;332:145-53.

15.   Degos V, Vacas S, Han Z, van Rooijen N, Gressens P, Su H, et al. Depletion of Bone

Marrow–derived Macrophages Perturbs the Innate Immune Response to Surgery and Reduces

Postoperative Memory Dysfunction. Anesthesiology. 2013;118(3):527-36.

16.   Poon DCH, Ho YS, You R, Tse HL, Chiu K, Chang RCC. PKR deficiency alters E.

coli-induced sickness behaviors but does not exacerbate neuroimmune responses or bacterial

load. J. Neuroinflammation. 2015;12(1):212.

17.   Carret-Rebillat AS, Pace C, Gourmaud S, Ravasi L, Montagne-Stora S, Longueville S,

et al. Neuroinflammation and Aβ Accumulation Linked To Systemic Inflammation Are

Decreased By Genetic PKR Down-Regulation. Sci. Rep. 2015;5(1):8489.

18.   Garcia M, Meurs E, Esteban M. The dsRNA protein kinase PKR: virus and cell

control. Biochimie. 2007;89(6-7):799-811.

19.   Maggi LB, Heitmeier MR, Scheuner D, Kaufman RJ, Buller RML, Corbett JA.

Potential role of PKR in double-stranded RNA-induced macrophage activation. EMBO J.

2000;19(14):3630-8.

20.   Williams BRG. Signal Integration via PKR. Sci STKE. 2001;2001(89):re2-re.

21.   Goh KC, deVeer MJ, Williams BRG. The protein kinase PKR is required for p38

MAPK activation and the innate immune response to bacterial endotoxin. EMBO J.

2000;19(16):4292-7.

22.   Zamanian-Daryoush M, Mogensen TH, DiDonato JA, Williams BRG. NF-κB

Activation by Double-Stranded-RNA-Activated Protein Kinase (PKR) Is Mediated through

NF-κB-Inducing Kinase and IκB Kinase. Mol. Cell. Biol. 2000;20(4):1278.

23.   Yim HCH, Williams BRG. Protein Kinase R and the Inflammasome. J. Interferon

Cytokine Res. 2014;34(6):447-54.

24.   Lu B, Nakamura T, Inouye K, Li J, Tang Y, Lundbäck P, et al. Novel role of PKR in

inflammasome activation and HMGB1 release. Nature. 2012;488(7413):670-4.

25.   Chang RCC, Suen KC, Ma CH, Elyaman W, Ng HK, Hugon J. Involvement of

double☐stranded RNA☐dependent protein kinase and phosphorylation of eukaryotic

initiation factor☐2α in neuronal degeneration. J. Neurochem. 2002;83(5):1215-25.

26.   Zhu Ping J, Huang W, Kalikulov D, Yoo Jong W, Placzek Andon N, Stoica L, et al.

Suppression of PKR Promotes Network Excitability and Enhanced Cognition by Interferon-γ-

Mediated Disinhibition. Cell. 2011;147(6):1384-96.

27.   You R, Liu Y, Chang RCC. A Behavioral Test Battery for the Repeated Assessment

of Motor Skills, Mood, and Cognition in Mice. J. Vis. Exp. 2019(145).

28. Ho YS, Lau CF, Lee K, Tan JY, Lee J, Yung S, et al. Impact of unilateral ureteral obstruction on cognition and neurodegeneration. Brain Res. Bull. 2021;169:112-27.

29. Wang RPH, Huang J, Chan KWY, Leung WK, Goto T, Ho YS, et al. IL-1β and TNF-α play an important role in modulating the risk of periodontitis and Alzheimer's disease. J. Neuroinflammation. 2023;20(1):71.

30. O'Connor AM, Burton TJ, Leamey CA, Sawatari A. The use of the puzzle box as a means of assessing the efficacy of environmental enrichment. J. Vis. Exp. 2014(94).

31. Chang RCC, Suen KC, Ma CH, Elyaman W, Ng HK, Hugon J. Involvement of double-stranded RNA-dependent protein kinase and phosphorylation of eukaryotic initiation factor-2alpha in neuronal degeneration. J. Neurochem. 2002;83(5):1215-25.

32. Young K, Morrison H. Quantifying microglia morphology from photomicrographs of immunohistochemistry prepared tissue using ImageJ. J. Vis. Exp. 2018(136):e57648.

33. Bao J, Tan S, Yu W, Lin Z, Dong Y, Chen Q, et al. The Effect of Peritoneal Air Exposure on Intestinal Mucosal Barrier. Gastroenterol Res Pract. 2014;2014:674875.

34. Tan S, Yu W, Lin Z, Chen Q, Shi J, Dong Y, et al. Peritoneal Air Exposure Elicits an Intestinal Inflammation Resulting in Postoperative Ileus. Mediators Inflamm. 2014;2014:924296.

35. Jacobi CA, Ordemann J, Zieren HU, Volk HD, Bauhofer A, Halle E, et al. Increased Systemic Inflammation After Laparotomy vs Laparoscopy in an Animal Model of Peritonitis. Arch. Surg. 1998;133(3):258-62.

36. Shaftel SS, Griffin WST, O'Banion MK. The role of interleukin-1 in neuroinflammation and Alzheimer disease: an evolving perspective. J. Neuroinflammation. 2008;5(1):7.

37. Saghazadeh A, Ferrari CC, Rezaei N. Deciphering variability in the role of interleukin-1β in Parkinson's disease. Rev Neurosci. 2016;27(6):635-50.

38.   Moynagh PN. The interleukin-1 signalling pathway in astrocytes: a key contributor to

inflammation in the brain. J. Anat. 2005;207(3):265-9.

39.   Lu B, Nakamura T, Inouye K, Li J, Tang Y, Lundbäck P, et al. Novel role of PKR in

inflammasome activation and HMGB1 release. Nature. 2012;488(7413):670-4.

40.   Takada Y, Ichikawa H, Pataer A, Swisher S, Aggarwal BB. Genetic deletion of PKR

abrogates TNF-induced activation of IkappaBalpha kinase, JNK, Akt and cell proliferation

but potentiates p44/p42 MAPK and p38 MAPK activation. Oncogene. 2007;26(8):1201-12.

41.   Couturier J, Paccalin M, Morel M, Terro F, Milin S, Pontcharraud R, et al. Prevention

of the β-amyloid peptide-induced inflammatory process by inhibition of double-stranded

RNA-dependent protein kinase in primary murine mixed co-cultures. J. Neuroinflammation.

2011;8(1):72.

42.   Maggi LB, Jr., Heitmeier MR, Scheuner D, Kaufman RJ, Buller RM, Corbett JA.

Potential role of PKR in double-stranded RNA-induced macrophage activation. EMBO J.

2000;19(14):3630-8.

43.   Carret-Rebillat AS, Pace C, Gourmaud S, Ravasi L, Montagne-Stora S, Longueville S,

et al. Neuroinflammation and Aβ accumulation linked to systemic inflammation are

decreased by genetic PKR down-regulation. Sci. Rep. 2015;5(1):1-8.

44.   He Y, Franchi L, Núñez G. The protein kinase PKR is critical for LPS-induced iNOS

production but dispensable for inflammasome activation in macrophages. Eur. J. Immunol.

2013;43(5):1147-52.

45.   McNeill E, Crabtree MJ, Sahgal N, Patel J, Chuaiphichai S, Iqbal AJ, et al.

Regulation of iNOS function and cellular redox state by macrophage Gch1 reveals specific

requirements for tetrahydrobiopterin in NRF2 activation. Free Radic. Biol. Med.

2015;79:206-16.

46. Rosas-Ballina M, Tracey KJ. Cholinergic control of inflammation. J. Intern. Med.

2009;265(6):663-79.

47. Stern E, Chinnakkaruppan A, David O, Sonenberg N, Rosenblum K. Blocking the

eIF2α Kinase (PKR) Enhances Positive and Negative Forms of Cortex-Dependent Taste

Memory. J. Neurosci. 2013;33(6):2517.

48. Jiang Z, Belforte JE, Lu Y, Yabe Y, Pickel J, Smith CB, et al. eIF2α Phosphorylation-

Dependent Translation in CA1 Pyramidal Cells Impairs Hippocampal Memory Consolidation

without Affecting General Translation. J. Neurosci. 2010;30(7):2582.

49. Malkov A, Popova I, Ivanov A, Jang SS, Yoon SY, Osypov A, et al. Aβ initiates brain

hypometabolism, network dysfunction and behavioral abnormalities via NOX2-induced

oxidative stress in mice. Commun. Biol. 2021;4(1):1054.

50. Firbank MJ, Yarnall AJ, Lawson RA, Duncan GW, Khoo TK, Petrides GS, et al.

Cerebral glucose metabolism and cognition in newly diagnosed Parkinson's disease: ICICLE-

PD study. J. Neurol. Neurosurg. Psychiatry. 2017;88(4):310.

51. Drzezga A, Lautenschlager N, Siebner H, Riemenschneider M, Willoch F, Minoshima

S, et al. Cerebral metabolic changes accompanying conversion of mild cognitive impairment

into Alzheimer's disease: a PET follow-up study. Eur. J. Nucl. Med. Mol. Imaging.

2003;30(8):1104-13.

52. Mosconi L, Pupi A, De Leon MJ. Brain glucose hypometabolism and oxidative stress

in preclinical Alzheimer's disease. Ann. N. Y. Acad. Sci. 2008;1147:180-95.

53. Bergau N, Maul S, Rujescu D, Simm A, Navarrete Santos A. Reduction of Glycolysis

Intermediate Concentrations in the Cerebrospinal Fluid of Alzheimer's Disease Patients.

Front. Neurosci. 2019;13:871-.

54.   Stovell MG, Mada MO, Helmy A, Carpenter TA, Thelin EP, Yan JL, et al. The effect

of succinate on brain NADH/NAD+ redox state and high energy phosphate metabolism in

acute traumatic brain injury. Sci. Rep. 2018;8(1):11140.

55.   Khellaf A, Garcia NM, Tajsic T, Alam A, Stovell MG, Killen MJ, et al. Focally

administered succinate improves cerebral metabolism in traumatic brain injury patients with

mitochondrial dysfunction. J. Cereb. Blood Flow. Metab. 2022;42(1):39-55.

56.   Nakamura T, Kunz RC, Zhang C, Kimura T, Yuan CL, Baccaro B, et al. A critical

role for PKR complexes with TRBP in Immunometabolic regulation and eIF2α

phosphorylation in obesity. Cell Rep. 2015;11(2):295-307.

57.   Nakamura T, Furuhashi M, Li P, Cao H, Tuncman G, Sonenberg N, et al. Double-

Stranded RNA-Dependent Protein Kinase Links Pathogen Sensing with Stress and Metabolic

Homeostasis. Cell. 2010;140(3):338-48.

58.   Carvalho BM, Oliveira AG, Ueno M, Araújo TG, Guadagnini D, Carvalho-Filho MA,

et al. Modulation of double-stranded RNA-activated protein kinase in insulin sensitive tissues

of obese humans. Obesity (Silver Spring). 2013;21(12):2452-7.

59.   Carvalho-Filho MA, Carvalho BM, Oliveira AG, Guadagnini D, Ueno M, Dias MM,

et al. Double-stranded RNA-activated protein kinase is a key modulator of insulin sensitivity

in physiological conditions and in obesity in mice. Endocrinology. 2012;153(11):5261-74.

Fig. 1. Laparotomy induced systemic inflammation and neuroinflammation. Relative

cytokine mRNA expression levels were measured in the liver, frontal cortex, and

hippocampus of C57BL/6J mice at (A) 4 h and (B) 1 day after laparotomy. Representative

flow cytometry plots illustrating the increase in CD68-eGFP[+] cells in the (C) frontal cortex

and (E) hippocampus of the CD68-eGFP mice on postoperative day 14. Morphological gating

strategy refers to dot plot SSC-H versus FSC-H to eliminate debris and aggregates in the (C)

frontal cortex and (E) hippocampus of the control and laparotomy groups. Doublets are

excluded by plotting FSC-H versus width and SSC-H versus width. Proportion of CD68-

eGFP$^+$ cells (labeled with eGFP) and cell viability (labeled with DAPI) were determined by

flow cytometry with FITC and pacific blue channel respectively. The data generated are

plotted in two-dimensional dot plots in which FITC-A versus Pacific-Blue-A. A four-

quadrant gate was established: Q1) live, CD68-eGFP$^+$ cells (FITC positive/Pacific blue

negative); Q2) dead, CD68-eGFP$^+$ cells (FITC positive/Pacific blue positive); Q3) live cells

without CD68 expression (FITC negative/Pacific blue negative); and Q4, dead cells without

CD68 expression (FITC negative/Pacific blue positive). Percentage of CD68-eGFP$^+$ cells in

the (D) frontal cortex and (F) hippocampus of the control and laparotomy groups. n = 6 per

group. Data are expressed as mean ± S.E.M. Differences were assessed by unpaired two-

tailed Student's t-test as follows: $^*p < 0.05$ and $^{**}p < 0.01$ compared with the control group.

Fig. 2. Laparotomy induced activation of microglia in the frontal cortex and hippocampus after laparotomy. Representative confocal images of the immunohistochemical staining of DAPI, and Iba1 in the (A) frontal cortex, (C) cornu ammonis (CA) 1, (E) CA3, and (G) dentate gyrus (DG) of the hippocampus sections from

CD68-eGFP mice of the control and laparotomy groups on postoperative day 14. Number of

Iba1 positive cells, relative fluorescence intensity of Iba1 positive cells, total number of

endpoints per Iba1 positive cell, and summed process length per Iba1 positive cell in the (B)

frontal cortex, (D) CA 1, (F) CA3, and (H) DG of the hippocampus sections were quantified.

n = 5 per group. Data are expressed as mean ± S.E.M. Differences were assessed by unpaired

two-tailed Student's t-test denoted as follows: $^{*}p < 0.05$, $^{**}p < 0.01$ and $^{***}p < 0.001$

compared with the control group.

Fig. 3. Cognitive impairment of C57BL/6J mice following laparotomy. (A) Percentage of the body weight change from baseline of the laparotomy and control groups throughout the postoperative period. (B) Total distance travelled and (C) central duration of mice in the open-field test. (D) Discrimination index, which is the ratio of exploration time of one object to two objects, was measured on the novel object recognition test day 1 (with two identical objects: A+A) and (E) day 2 (with a novel object replaced the familiar one: A+B). (F) Total number of arm entries and (G) percentage of alternation were measured in the spontaneous alternation Y-maze test. In the puzzle box test, the latency time for the mice entering the goal zone in different tasks was measured. All the tasks include (H) task for habituation (T1), (I) tasks for testing problem-solving skills (T2, T5, T8, and T11), and (J) tasks for evaluating short-term (T3, T6, T9, and T12) and (K) long-term memories (T4, T7, T10, and T13) in the puzzle box test. n = 9–11per group. Data are expressed as mean ± S.E.M. Differences were assessed by unpaired two-tailed Student's t-test denoted as follows: $^*p < 0.05$ and $^{**}p < 0.01$ compared with the control group.

Fig. 4. Knockout of PKR in mice ameliorated laparotomy-induced peripheral and neural inflammation after surgery. Relative cytokine mRNA expression levels were measured in the liver, frontal cortex, and hippocampus of the C57BL/6J mice and PKR-KO mice at (A) 4 h and (B) 1 day following laparotomy. n = 6 per group. Data are expressed as mean ± S.E.M. Differences were assessed by two-way ANOVA and Bonferroni's post hoc test denoted as follows: $^{*}p < 0.05$ and $^{**}p < 0.01$, compared with the control group.

Fig. 5. Knockout of PKR in mice ameliorated activation of microglia in the frontal

cortex and hippocampus following laparotomy. Representative confocal images of the

immunohistochemical staining of DAPI and Iba1 in the (A) frontal cortex, (C) cornu

ammonis (CA) 1, (E) CA3, and (G) dentate gyrus (DG) of the hippocampus sections from

PKR-KO mice of the control and laparotomy groups on postoperative day 14. Number of

Iba1 positive cells, relative fluorescence intensity of Iba1 positive cells, total number of

endpoints per Iba1 positive cell, and summed process length per Iba1 positive cell in the (B)

frontal cortex, (D) CA1, (F) CA3, and (H) DG of the hippocampus sections were quantified.

n = 5 per group. Data are expressed as mean ± S.E.M. Differences were assessed by unpaired

two-tailed Student's t-test.

Fig. 6. Genetic deletion of PKR in mice alleviated laparotomy-induced cognitive impairment. (A) Percentage of the body weight change from baseline of the C57BL/6J mice and PKR-KO mice throughout the postoperative period. (B) Total distance travelled and (C) central duration of mice in the open-field test. (D) Discrimination index, which is the ratio of exploration time of one object to two objects, was measured on the novel object recognition test day 1 (with two identical objects: A+A) and (E) day 2 (with a novel object replaced the familiar one: A+B). (F) Total number of arm entries and (G) percentage of alternation were

measured in the spontaneous alternation Y-maze test. In the puzzle box test, the latency time

for the mice entering the goal zone in different tasks was measured. All the tasks include (H)

task for habituation (T1), and (I) tasks for testing problem-solving skills (T2, T5, T8, and

T11), (J) tasks for evaluating short-term (T3, T6, T9, and T12) and (K) long-term memories

(T4, T7, T10, and T13). n = 8–11 per group. Data are expressed as mean ± S.E.M.

Differences were assessed by two-way ANOVA and Bonferroni's post hoc test denoted as

follows: $^{*}p < 0.05$ and $^{**}p < 0.01$ compared with the control group.

Fig. 7. Inhibition of PKR in cholinergic neurons ameliorated laparotomy-induced cognitive impairment. (A) Percentage of the body weight change from baseline of the ChAT-IRES-Cre-eGFP mice of the control group injected with saline (SAL) and the treatment group injected with the rAAV-DIO-PKR-K296R (AAV) throughout the postoperative period. (B) Total distance travelled and (C) central duration of mice in the open-field test. (D) Total number of arm entries and (E) percentage of alternation were measured in the spontaneous alternation Y-maze test. In the puzzle box test, the latency time

for the mice entering the goal zone in different tasks was measured. All the tasks include (F)

task for habituation (T1), (G) tasks for testing problem-solving skills (T2, T5, T8, and T11),

(H) tasks for evaluating short-term (T3, T6, T9, and T12) and (I) long-term memories (T4,

T7, T10, and T13). n = 6–8 per group. Data are expressed as mean ± S.E.M. Differences were

assessed by two-way ANOVA and Bonferroni's post hoc test denoted as follows: $^{*}p < 0.05$,

$^{**}p < 0.01$, $^{***}p < 0.001$, $^{****}p < 0.0001$ compared with the control group.

Fig. 8. Inhibition of PKR in cholinergic neurons ameliorated laparotomy-induced

changes in glucose metabolism in the frontal cortex. Two-dimensional score plots using

the (A) principal component analysis (PCA) and (B) partial least squares discriminant

analysis (PLS-DA). (C) Heatmap of the metabolites involved in central carbon metabolism in

the frontal cortex of the ChAT-IRES-Cre-eGFP mice injected with saline (SAL) or rAAV-

DIO-PKR-K296R (AAV). Both groups of mice were then randomly assigned into the

laparotomy and control groups. The effects of AAV treatment and laparotomy on glucose

metabolism were analyzed using gas chromatography-tandem mass spectrometry (GC-

MS/MS). The column represents the samples, and the row displays the glucose metabolites.

The heatmap scale ranges from −3 to 3. The brightness of each color corresponded to the

magnitude of the log2 fold change of the measured metabolites. Concentration of (D)

glucose, (E) phosphoenolpyruvate, (F) succinate, and (G) 6-phosphogluconate were

quantified in nmol/g. n = 4 per group. Data are expressed as mean ± S.E.M. Differences were

assessed by two-way ANOVA and Bonferroni's post hoc test denoted as follows: $^{*}p < 0.05$,

$^{**}p < 0.01$, compared with the control group.