Snapshots from Cryo-ET of active SARS-CoV-2 virions

Hideo Fukuhara[1,2,3,#], Hisham M. Dokainish[2,#,*], Shunsuke Kita[2], Koshiro Tabata[4,5], Akira Takasu[6],

Juha T. Huiskonen[7], Yuki Anraku[2], Toshiya Senda[6,8,9], David I. Stuart[10], Michihito Sasaki[4], Yasuko

Orba[4], Yasuhiko Suzuki[5,11,*], Hirofumi Sawa[4,5,12,13,*] & Katsumi Maenaka[1,2,3,5,12,14,*].

[1] Division of Pathogen Structure, International Institute for Zoonosis Control, Hokkaido University,

Kita-20, Nishi-10, Kita-ku, Sapporo 001-0020, Japan.

[2] Laboratory of Biomolecular Science and Center for Research and Education on Drug Discovery,

Faculty of Pharmaceutical Sciences, Hokkaido University, Kita-12, Nishi-6, Kita-ku, Sapporo 060-

0812, Japan.

[3] Global Station for Biosurfaces and Drug Discovery, Hokkaido University, Kita-12, Nishi-6, Kita-

ku, Sapporo 060-0812, Japan

[4] Division of Molecular Pathobiology, International Institute for Zoonosis Control, Hokkaido

University, Kita-20, Nishi-10, Kita-ku, Sapporo 001-0020, Japan

[5] Institute for Vaccine Research and Development (HU-IVReD), Hokkaido University, Kita-21,

Nishi-11, Kita-ku, Sapporo 001-0020, Japan

[6] Structural Biology Research Center, Institute of Materials Structure Science, High Energy

Accelerator Research Organization (KEK), Tokyo 135-0063, Japan

[7] Institute of Biotechnology, Helsinki Institute of Life Science HiLIFE, University of Helsinki, P.O.

Box 56, 00014 Helsinki, Finland.

[8] Department of Accelerator Science, School of High Energy Accelerator Science, SOKENDAI

(the Graduate University for Advanced Studies), 1-1 Oho, Tsukuba, Ibaraki 305-0801, Japan

[9] Faculty of Pure and Applied Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki

305-8572, Japan

[10] Division of Structural Biology, University of Oxford, The Wellcome Centre for Human Genetics,

Headington, Oxford OX3 7BN, UK; Chinese Academy of Medical Sciences Oxford Institute,

University of Oxford, Oxford OX3 7FZ, UK; Diamond Light Source Ltd, Harwell Science &

Innovation Campus, Didcot OX11 0DE, UK; Nuffield Department of Medicine, Wellcome Centre

for Human Genetics, University of Oxford, Oxford OX3 7BN, UK.

[11] Division of Bioresources, International Institute for Zoonosis Control, Hokkaido University,

Kita-20, Nishi-10, Kita-ku, Sapporo 001-0020, Japan

[12] One Health Research Center, Hokkaido University, Kita-19, Nishi-9, Kita-ku, Sapporo 001-

0020, Japan

[13] Global Virus Network, Baltimore, Maryland, USA

[14] Faculty of Pharmaceutical Sciences, Kyushu University, Fukuoka, 812-8582, Japan.

[#] Contributed equally.

*Correspondence authors:

Y.S. (email: suzuki@czc.hokudai.ac.jp)

H.D. (email: hisham.dokainish@pharm.hokudai.ac.jp)

H.S. (email: h-sawa@ivred.hokudai.ac.jp)

K.M. (email: maenaka@pharm.hokudai.ac.jp) (Lead Contact)

Abstract

Understanding the molecular properties of SARS-CoV-2 is crucial to tackle the continuing

variant emergence as well as future outbreaks. Current structural knowledge of the trimeric spike

protein is largely based on truncated recombinant proteins and inactivated full-length forms, which

may suffer from overstabilization through designed mutations and chemical fixation. Here, we

apply cryo-electron tomography (cryo-ET) at a Biosafety level 3 facility to study the virus structure

in its native, active state. The virus particles show variable shapes with diffusible spikes. While

the majority of spike proteins display typical prefusion trimer conformations, we identify atypical

open-trimer prefusion states, revealing a hidden level of flexibility. The sub-tomogram averaged

structure also suggests a loosely packed trimer structure. Spike dynamics reveal cryptic epitopes

in conserved regions among coronaviruses that can be targeted for broadly effective vaccines.

Therefore, the structural analysis of virus structures in their active forms will provide significant

implications on fusion mechanism as well as vaccine and antibody/drug design.

Main

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) continues to impact global health[1]. The ongoing emergence of new variants hinders the efficacy of developed vaccines and therapeutic antibodies[2,3]. The spike protein decorates the virus surface and plays central roles in the viral entry and immune shielding mechanisms[4,5]. The spike protein is a trimeric glycoprotein, wherein each protomer is formed of two subunits (S1 and S2) with a transmembrane region[6]. Numerous studies elucidated spike protein structures and dynamics, paving the way for mRNA vaccine developments[6-10]. Specifically, cryogenic electron microscopy (cryo-EM) has been used to determine structures of the recombinant spike protein in wild-type and emerging variants[3,6-12]. Moreover, cryo-ET has enabled the study of full-length spikes in the inactivated viruses, unraveling the flexibility of its stalk region and the protein distribution on the virus surface[7,13-18]. Structural studies provided fundamental information on the virus molecular structure, architecture, interactions, and dynamics[4].

Despite the enormous amount of structural information available on SARS-CoV-2 virus and its spike protein, our understanding of the virus in its active form is still lacking[19]. In fact, the majority of cryo-EM studies are performed using a truncated ectodomain of the spike protein, which includes several mutations (e.g. 2P and 6P proline mutations) designed to stabilize the prefusion state[20,21]. Accordingly, several approved vaccines include similar proline mutations, for improving production and stabilization[22]. Indeed, the proline mutations may induce local rigidity altering the nearby secondary structure[23]. Likewise, to reduce biological hazards, chemical fixation is usually applied in cryo-ET experiments[15,19]. Chemicals such as paraformaldehyde (PFA) are known to interact with charged residues and may cause morphological changes that alter protein structures[19,24]. In addition, electron (E)-beam inactivation was recently shown to be damaging to

spike protein structures[15]. According to previous structural studies, the spike protein has a

reasonably stable prefusion state with a number of conformational states that are governed by the

motion of its S1 receptor binding domains (RBDs) from Down to Up states[6]. In contrast, Hydrogen

Deuterium Exchange Mass Spectrometry (HDX-MS) study showed that the prefusion state of the

spike protein is quite dynamic, highlighting the existence of a form of the prefusion state in which

parts of S2 are exposed[23]. Furthermore, the binding of some antibodies to the S2 subunit in the

prefusion state suggests the formation of an unknown open trimer structure[25]. Collectively,

previous studies showed structural flexibility at several levels using artificially stabilized proteins

and inactivated viruses. However, true images of spike proteins are still missing.

Understanding the differences between the current structural knowledge of SARS-CoV-2 that

may suffer from overstabilization, and the active form is crucial for future vaccine and

antibody/drug development. Certainly, elucidating the dynamics of spike proteins in its native form

would shape the approach to targeting future variants. To address this point, we set up the high-

end cryo-electron microscope at a Biosafety level 3 (BSL3) area to employ cryo-ET and single-

particle tomography (SPT)[26] to unravel the structure of the spike protein using active viruses. In

this experiment, neither chemical fixation nor ultracentrifugation were applied, to avoid spike

structure alteration or detachment. To the best of our knowledge, this is the first structural study of

SARS-CoV-2 in its active form.

BSL3 High-end Cryo-electron Microscope.

We set up the latest generation of cryo-electron microscope (300 kV), Krios G4 (Thermo

Fisher Scientific) with a Falcon 4 direct electron detector and a Selectris energy filter in the

management area for BSL3 pathogens in the International Institute for Zoonosis Control,

Hokkaido University (**Fig. S1**). The detail of BSL3 cryo-EM system and procedure for handling

the grid preparation and data collection will be published elsewhere, but briefly, virus vitrification

using EM GP2 (Leica) and autogrid assembly were performed in the safety cabinet. The autogrid

was transferred to Krios G4 and cryo-ET data were collected (**Fig. S1**). This is one of the first

examples of operating a high-end cryo-electron microscope in the BSL3 area. The active SARS-

CoV-2 of the ancestral strain WK-521 in cell culture was directly frozen and observed. Thus, the

virus preparation did not include any ultracentrifugation or chemical fixation to avoid any kind of

artifacts as much as possible.

The Pleomorphic Virus Particles Morphology.

Early cryo-ET study of the inactivated SARS-CoV-2 demonstrated the formation of an

ellipsoidal or a spherical-shaped viral envelope, with an average of 96.6 nm for its longest axis[16].

In contrast, a more recent cryo-ET study suggested the formation of a dominant flat cylindrical

shape, in ancestral, alpha, beta and delta variants, with an average diameter of 102 nm[14]. Notably,

a later cryo-ET study of the delta variant has shown the formation of fused virus particles, where

one envelope encloses two sets of RNPs[15]. In summary, previous cryo-ET studies suggested some

level of pleomorphism.

In this study, 42 tomograms of the active ancestral virus were recorded and reconstructed,

including 46 virus particles (**e.g. Movie S1**). A tilt angular step of 1°, covering a range from −60°

to +60° was used, resulting in 121 tilt images per tomogram. **Fig. 1** shows a variety of envelope

shapes, including dominant spherical (L1: 99 ± 13.6 nm, L2: 94 ± 13.6 nm) and ellipsoidal shapes

(L1: 117 ± 19.7 nm, L2: 94 ± 15.9 nm), as well as oval (L1: 99 ± 21.5 nm, L2: 87 ± 10.6 nm),

cylindrical (L1: 93 ± 7.9 nm, L2: 57 ± 10.1 nm) and bean-like (L1: 100 ± 12.7 nm, L2: 65 ± 10.9

nm) envelops (summarized in **Table S1**). The measured diameters of the two longest axes (L1 and

L2) indicate a large variation in the virus size, wherein the maximum axis (L1) can vary from 72

to 150 nm (**Movie S2**). Remarkably, very large virus particles with L1:150, L2:125 (**Fig. 1c right**)

and L1:139, L2:102 nm (**Fig. 1c left**) were observed. Such large particles may indicate the fusion

of two virus particles in ancestral strain, as recently observed in delta variant[15]. **Fig. S2a and**

Movie S3 also show the observation of two virus particles apparently undergoing a fusion process.

In addition, **Fig. S2b** suggests that there is no correlation between the observed morphology and

ice thickness. It also shows that the third axis is correlated with the particle size. Collectively,

unlike previous results from inactivated viruses that exhibit a dominant morphology, our results

suggest a high level of flexibility, adaptability, and pleomorphism on virus shape in response to

surrounding environments.

Figure 1. Pleomorphology of active virus particles. a) Central sections (7.5 nm thick) through the cryo-

ET tomograms of the ancestral virions. b) Violin plot representation of the two longest diameters (L1 and

L2) in 46 virus particles. c) Tomograms slices of three virions with very large diameters (Left: L1:139,

L2:102; Middle: L1:125, L2:119; Right: L1:150, L2:125). d) Central slices of virions on carbon film.

Spike Distribution and Arrangements.

In the current dataset, one-third of the particles (14 out of 46) were observed on the carbon

film. These particles tend to include a higher density of spike proteins, in comparison to the ones

in the grid holes. On the carbon, spike proteins are pushed to the edge of the flattened particle,

reflecting high diffusibility within the native viral lipid membrane (**Fig. 1d**). Such high diffusibility

may allow the virus to relocate more spikes at the interface upon encountering natural host receptor,

Angiotensin-converting enzyme 2 (ACE2). On the other hand, within grid holes, spike proteins

are more dispersed on the surface, where fewer spikes can be identified in most cross sections. **Fig.**

2a shows that spike proteins can exist as a dimer of trimers or even a trimer of trimers (**Movie S1**

right), and similar complexes were previously observed in full-length spike proteins from the

Novavax vaccine[27]. **Fig. S3** indicates the formation of two spike trimers with shared

transmembrane regions, as reported previously[16]. Such arrangements raise a question regarding

the possibility of protomer swapping between trimeric spike structures. Both 247 prefusion and 44

postfusion conformations were observed across active virus particles. Specifically, the postfusion

state was found to occur in clusters (**Fig. 2b**). Such arrangements of the postfusion state may

facilitate membrane fusion by attaching several fusion peptides at the same time.

Figure 2. Distribution and atypical conformations of the spike protein. a) The 2D central slice of virus particle showing the formation of higher order structures including dimer of trimers and trimer of trimers. Isosurface is also shown in mesh representation, wherein PDB 6ZGE[10] is fitted and shown in surface representation. b) Tomograms slices (7.5 nm thick) of four virions with clusters of postfusion spike structures. Postfusion conformations are highlighted with white arrowheads. c) The isosurface representation of several spike proteins on the virus surface, wherein the closed/packed and open trimers of prefusion states can be identified as shown by dashed lines. d) The magnified view of 2D central slice tomograms of the typical spike protein densities including Down, 1Up, 2Up and 3Up conformations, as

well as atypical spike protein densities observed in active viruses. The detached S1 subunits are highlighted

by dashed lines. e) Proposed schematic representation of the spike opening mechanism at prefusion states,

in which one protomer S1 subunit undergoes a large conformational change.

Atypical Spike Conformations Indicate an Open Trimer.

Fig. 2c and d show all typical forms of the spike protein observed in our dataset. This

includes the prefusion Down, 1Up, 2Up and 3Up states, reflecting the inherent dynamical nature

of the spike protein, in the absence of ACE2 or neutralizing antibody (nAb). Notably, besides

typical conformations, atypical forms of the spike protein were also observed. Among these, spike

protomers may undergo large conformational changes, forming a novel open trimer state. Simple

fitting to a detached density indicates that the observed motion involves the S1 subunit of single

spike protomer. Accordingly, a range of open trimer structures can be seen (**Fig. 2d**), exposing the

S2 subunit. It is important to mention that most of spike protein conformations in our dataset were

typical prefusion states, where open trimer conformations occur in a small portion (~4%) of

observed prefusion states. These results suggest a hidden motion of the spike structure, in which

spike protomers are more flexible and can undergo conformational opening (**Fig. 2e**). Consistently,

previous HDX-MS studies using the recombinant ectodomain of the spike protein with a prefusion

stabilization (2P mutations) suggested the minor presence of an alternative conformation, in which

S2 residues are solvent exposed, which can be targeted by nAbs, such as 3A3 and RAY53[22,23,25].

Therefore, in active viruses, the typical closed/packed prefusion conformation is dominant but the

spike protein itself has ability to spontaneously transit to an open-trimer conformation.

Figure 3. The spike protein averaged map obtained by SPT analysis. a) The sub-tomogram averaged map with the superimposition of cryo-EM structures of the spike trimer (PDB 7M0J; Top)) and (PDB 6ZGE; Bottom). Purple, green and pink colors were used to represent spike protomers. The zoomed-in view of the trimeric S2 interface from 7M0J (pink and green ribbons) and 6ZGE (blue ribbon), fitted into the spike protein average map (Top center), indicating the S2 variation in fitted middle helices. The zoomed-in view of the terminal part of spike protein averaged map (Bottom center) with fitted PDB structures, 7M0J (pink, green and purple ribbons) and 6ZGE (blue ribbon). Arrows in central panels indicate structural shifts from 6ZGE to 7M0J. b) Top views of the sub-tomogram averaged maps of the C3 symmetry (top) and the C1 symmetry (bottom) fitted with prefusion-state spike protein (PDB 7M0J).

Spike Average Conformation Suggests a Loosely Packed Structure.

To gain further insights into the mechanism of the open-trimer formation, we performed single particle tomography (SPT) analysis of the prefusion packed conformation. 237 prefusion spike particles were manually picked (excluding open trimer and postfusion states). Around 28,000 tilt images (121 tilt per particle) were used for the 3D re-construction, with a voxel size of 1.884 Å. To avoid any bias, no previously published maps of the spike protein were used in the alignment and refinement processes. **Fig. 3a** and **Movie S4** show the sub-tomogram averaged map of the C3 symmetric spike protein, which was obtained with a global resolution of 14 Å (**Fig. S4**). In addition, a C1 symmetry map was also obtained at a slightly lower global resolution of 17.0 Å, and this was essentially similar to the C3 symmetric structure but with less density in one protomer (**Fig. 3b**). The general features of the maps obtained are similar to those of previous Cryo-EM studies. A 2.6 Å resolution structure of the ancestral spike protein (PDB 6ZGE)[10] fits adequately in the obtained low-resolution map (**Fig. 3a bottom**). However, a better fit was obtained using the asymmetrically refined structure (PDB 7M0J) of the engineered spike (u1S2q) with quadruple mutations (A570L, T572I, F855Y, N856I) [28,29] (**Fig. 3a top**), which was previously shown to exhibit intrinsic mobility, evidenced by high propensity of forming 1Up and 2Up conformations and sustained binding to the CR3022 antibody[29]. In detail, the SPT map shows missing density at the S2 middle helices interface (**Fig. 3a top center**). The helices were elongated and contributed to forming tightly packed S2 in many reported cryo-EM structures, however, there exists some variation of S2 middle helices interface in cryo-EM structures. Our cryo-ET map fits better to the 7MOJ structure harboring slightly bent helices (**Fig. 3a top center**). Notably, the S2 subunit shows multiple regions with charged residues (**Fig. S5**) at the trimer interface and, in the 7MOJ structure, the S2 subunit makes hydrophobic interactions at the interface to avoid charge repulsion. In addition, **Fig.**

3a bottom center highlights the fitting of 6ZGE and 7M0J to the proximal part of the stem helices,

wherein the 7M0J structure shows an upward shift that again better fit the averaged map. Previous

principal component (PC) analysis of 165 protomers from 52 spike Down PDB structures, shows

a similar behavior where PC1 and PC2 modes both show up and down motion in S2 with respect

to S1, indicating structural variation within the available PDB structures[30]. Therefore, these

structural features allow for more flexible movement of the protomers and the formation of the

above-mentioned open trimer structure as well as the transition to postfusion (**Fig. 4**). Notably, a

recent sub-tomogram averaged map of Env protein (9.1 Å resolution) in intact HIV particles, has

also suggested structural differences in middle helix region that suggest flexibility at the helical

bundle core[13].

Figure 4. Proposed model of spike structural changes for the membrane fusion. (top right panel)

Schematic representation of the spike structure with the domain organization. The color of each protomer

corresponds to Fig. 2e. (top left to center) New cryptic sites of the S2 subunit exposed in open-trimer

conformations of the spike protein, potential targets for vaccine and antibodies/drugs. (bottom) Schematic

presentation of the proposed fusion mechanism. Fixed or recombinant spikes protein take a closed or Up

RBD structure (gray box). On the active virus, spike proteins exist as a loosely packed trimer, with a

partially opened S1 subunit. When the ACE2 binding promotes the S2 subdomain exposure, the S2 subunit

irreversibly refolds to the postfusion state to fuse the viral envelope with the host cell membrane.

Implication on Fusion mechanism and Rational Vaccine and Antibody/Drug Design

Mutations of the spike protein are a major underlying cause for immune escape and

reduction of vaccine efficiency. The development of broadly neutralizing antibodies (bnAbs) is

important for preparedness for future pandemics. The recent review of Zhou et al. summarized

three major non-RBD sites targeted for bnAbs, (1) fusion peptide, (2) SD1 (the S1 subdomain 1)

and (3) the C-terminal stem helix[31]. Our cryo-ET map shows that the spike protein is flexible

enough for these sites to be to some extent exposed in open-trimer structures, possibly generating

bnAbs that exhibit neutralizing activity but with less potency. Either cocktails or bispecific forms

of bnAbs, which can recognize different epitopes of single spike protein spontaneously, may

improve the activity and tolerate mutations[31].

The development of long-term and broadly effective vaccines also requires new strategies

to target highly conserved epitope regions[32,33]. Unlike RBD which is a hot spot for mutations, the

S2 subunit shows higher conservation, wherein few mutations have emerged since the start of the

pandemic. In fact, the fusion peptide (FP) and heptad repeat regions (HR1 and HR2) of the S2

subunit are more conserved within all coronaviruses including SARS-CoV-1 and Middle East

respiratory syndrome coronavirus (MERS)-CoV[33]. This emphasizes the potential of developing

broadly effective vaccines that may cover current and future coronavirus outbreaks. However, in

current vaccine development strategies, the S2 subunit is considered as inaccessible, wherein

proline mutations are applied to further stabilize the prefusion state. On the contrary, our Cryo-ET

study here suggests the dynamic nature of the spike protein where parts of the S2 subunit are

accessible (**Fig. 2c and Fig. 4**). As described above, bnAbs may exhibit less potent activity but

vaccine-induced bnAbs would overcome this limitation, while the stabilization of the S2 subunit

may be necessary. In fact, the development of a multivalent S2-based vaccine was recently shown

to provide a wide effect against beta, delta and omicron variants as well as pangolin coronavirus[32].

Besides the S2 subunit, the currently observed dynamics of the active SARS-CoV-2 virus

(**Fig. 4**) indicates the accessibility of the interface of S1 and S2 subunits. Note that these regions

are conserved across SARS-CoV-2 variants. Indeed, the HDX-MS experiment of the recombinant

spike protein also shows the increased accessibility of the two subdomains (SD1 and SD2) in the

S1 subunit[23]. Collectively, accounting for spike dynamics and cryptic epitope formation may

facilitate the development of broadly effective vaccines. For instance, a vaccine design based on

the spike protomer may better account for spike protein dynamics, exposing cryptic epitopes (**Fig.**

4 and Fig. S6). To explore monomer dynamics, Molecular Dynamics (MD) simulations of the

spike monomer of the Omicron BA.2 variant were performed. **Fig. S7** shows that a large range of

motions in the S1 subunit can be observed in the absence of the trimeric complex, exposing cryptic

pockets located at the interface of the S2 subunit, RBD, and the SD1 subdomain, which may induce

a broader spectrum of immune effect. Indeed, monoclonal antibodies (mAbs) targeting cryptic

epitopes, such as CR3022, P008-60, and sd1.040, can break up spike trimeric structure[34-36]. It is

worth mentioning that an early study has reported the feasibility of using a monomeric spike

protein of SARS-CoV-1 as a vaccine candidate[37].

Unexpectedly, the active dynamics of the spike protein structure, especially open-trimer

conformations, reveal benefits for fusion efficiency. A small percentage of open-trimer

conformations definitively expose the S2' cleavage site for TMPRSS2 and cathepsins. In this sense,

the spike protein possesses the inherent potential to allow the S1-S2 dissociation, inducing the

exposure of the S2' cleavage site. This process could be further facilitated by the ACE2 binding

(**Fig. 4**). In retrospect, the effect of the D614G mutation would be anticipated to lead to a larger

exposure.

The spike protein in active virus is highly dynamic with multiple levels of flexibility

including, membrane diffusibility, stalk region knee-like motion[17], RBD Down to Up transition[38]

and open-trimer formation[23]. In addition, it is highly susceptible to change in temperature, pH and

even glycosylation[9,39]. Structural analysis using inactivated viruses or recombinant proteins is

essential, however, the effect of mutations, chemical fixations, and trimerization tags shall be

considered. Hence, it is crucial to study viral proteins in their native, active state. Cryo-ET at BSL3

facility is expected to be an indispensable method to tackle current and future pandemics.

Methods

Virus preparation

An ancestral SARS-CoV-2 strain, WK-521 (GISAID: EPI_ISL_408667), was kindly provided by Dr. Shimojima (National Institute of Infectious Diseases, Japan). The viral stock was prepared by the inoculation to Vero E6 cells constantly expressing human TMPRSS2 (Vero E6/TMPRSS2)[40]. At 2 days post-infection (dpi), the culturing supernatant was centrifuged at 440 × g for 2 min to remove the cell debris and stored at −80 °C as a virus stock. The titer of the stock was determined by plaque assay as previously described[41]. In brief, Vero E6/TMPRSS2 cells were inoculated with serial dilutions of virus stock for 1 h at 37 °C. The cells were overlaid with Dulbecco's Modified Eagle Medium (Nissui) containing 2% FBS and 0.5% Bacto Agar (Becton Dickinson). At 2 dpi, cells were fixed with 3.7% formaldehyde in phosphate-buffered saline (PBS) and stained with 1% crystal violet.

For the preparation of the cryo-ET sample, Vero E6/TMPRSS2 cells in a 6 well plate were inoculated with the virus at a multiplicity of infection (MOI) of 10 and incubated for 1 h at 37 °C. The media was then replaced with 400 μL of fresh Opti-MEM (Thermo Fisher Scientific). At 18 h post-infection, culture supernatant was collected and clarified with 0.45 μm filtration. The sample was kept at 37 °C and used immediately for grid preparation.

Data collection

For cryo-ET, 2 μL of sample was applied on each side of a freshly glow-discharged QUANTIFOIL R 1.2/1.3 Cu 200-mesh grid (Quantifoil Micro Tools). The grid was blotted at once for 15 s from the back side and plunged into liquid ethane using EM GP2 (Leica Microsystems) at 18 °C and 90% humidity chamber setting. The micrograph images were collected on a Krios G4

(Thermo Fisher Scientific) operating at a voltage of 300 kV with an energy filter Selectris (slit

width 10 eV) and a direct electron detector Falcon 4. Each micrograph was collected at a nominal

magnification of 64,000× (pixel size of 1.88Å) with an exposure time of 0.73 s (dose of 1.5 e/Å^2)

at a underfocus of 0.2 μm using a Volta phase plate (VPP). Tilt-series data collection was performed

with a dose-symmetric scheme tilting by 1° steps from −60° to 60° using Tomography software

(Thermo Fisher Scientific), **Table S2**.

Cryo-Electron Tomography Data Processing.

42 micrographs were processed using the IMOD package[42]. Patch tracking was used to

align tilt images. Ctfplotter was used to estimate defocus, phase shift, and astigmatism within the

context of the contrast transfer function[43]. Tomograms were binned 4x and reconstructed with

weighted back projection with a voxel size of 7.536 Å. SIRT-like filtered tomograms were

generated for visualization and particle picking. Reconstructed tomograms were also imported and

filtered in EMAN2[44]. Virus particles were manually segmented in EMAN2 and used for isosurface

visualization in Chimera-X software[45]. The two longest diameters of viruses and the thickness of

ice layer were calculated using 3dmod.

291 spike proteins, from 46 virions, were manually picked from the 4-fold binned SIRT

tomograms, including pre (247 particles) and postfusion (44 particles) states. The sub-tomogram

averaging of the prefusion state were processed in Relion 4.0[26]. The reference-free alignment was

performed, where the initial map was generated from the binned subtomograms and used

subsequently for other refinements. After obtaining an adequately representative map, the C3

symmetry was invoked. A series of refinements were performed starting from 4x binning to 2x to

unbinned. Finally, several cycles of the 3D refinement were performed at a voxel size of 1.884 Å

on unbinned subtomograms. After the 3D refinement, the CTF was corrected using ctffind-4.1.14[46]

and 2D images alignment was improved in each cycle. The cycles were repeated until no further

improvement in resolution was observed. The final average map of the C3 symmetry was estimated

to have 14.3 Å resolution at FSC = 0.143 cutoff in RELION4.0. To understand the effect of

symmetries, a final cycle of refinement was performed with C1 symmetry, providing a 17.0 Å

resolution map. The models of the spike protein, PDB 6ZGE and 7M0J, were rigid fitted in the C3

map using ChimeraX[45]. PyMOL software was used for visualizing the structures [47].

References

Hu, B., Guo, H., Zhou, P. & Shi, Z. L. Characteristics of SARS-CoV-2 and COVID-19. *Nat Rev Microbiol* **19**, 141-154 (2021). https://doi.org:10.1038/s41579-020-00459-7

Cao, Y. *et al.* Omicron escapes the majority of existing SARS-CoV-2 neutralizing antibodies. *Nature* **602**, 657-663 (2022). https://doi.org:10.1038/s41586-021-04385-3

Tamura, T. *et al.* Virological characteristics of the SARS-CoV-2 XBB variant derived from recombination of two Omicron subvariants. *Nat Commun* **14**, 2800 (2023). https://doi.org:10.1038/s41467-023-38435-3

Saville, J. W., Berezuk, A. M., Srivastava, S. S. & Subramaniam, S. Three-Dimensional Visualization of Viral Structure, Entry, and Replication Underlying the Spread of SARS-CoV-2. *Chem Rev* **122**, 14066-14084 (2022). https://doi.org:10.1021/acs.chemrev.1c01062

Wang, Q. *et al.* Structural and Functional Basis of SARS-CoV-2 Entry by Using Human ACE2. *Cell* **181**, 894-904 e899 (2020). https://doi.org:10.1016/j.cell.2020.03.045

Cai, Y. *et al.* Distinct conformational states of SARS-CoV-2 spike protein. *Science* **369**, 1586-1592 (2020). https://doi.org:10.1126/science.abd4251

Xu, C., Han, W. & Cong, Y. Cryo-EM and cryo-ET of the spike, virion, and antibody neutralization of SARS-CoV-2 and VOCs. *Curr Opin Struct Biol* **82**, 102664 (2023). https://doi.org:10.1016/j.sbi.2023.102664

Wrapp, D. *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* **367**, 1260-1263 (2020). https://doi.org:10.1126/science.abb2507

Zhou, T. *et al.* Cryo-EM Structures of SARS-CoV-2 Spike without and with ACE2 Reveal a pH-Dependent Switch to Mediate Endosomal Positioning of Receptor-Binding Domains. *Cell Host Microbe* **28**, 867-879 e865 (2020). https://doi.org:10.1016/j.chom.2020.11.004

Wrobel, A. G. *et al.* SARS-CoV-2 and bat RaTG13 spike glycoprotein structures inform on virus evolution and furin-cleavage effects. *Nat Struct Mol Biol* **27**, 763-767 (2020). https://doi.org:10.1038/s41594-020-0468-7

Fan, X., Cao, D., Kong, L. & Zhang, X. Cryo-EM analysis of the post-fusion structure of the SARS-CoV spike glycoprotein. *Nat Commun* **11**, 3618 (2020). https://doi.org:10.1038/s41467-020-17371-6

Saito, A. *et al.* Virological characteristics of the SARS-CoV-2 Omicron BA.2.75 variant. *Cell Host Microbe* **30**, 1540-1555 e1515 (2022). https://doi.org:10.1016/j.chom.2022.10.003

Mangala Prasad, V. *et al.* Cryo-ET of Env on intact HIV virions reveals structural variation and positioning on the Gag lattice. *Cell* **185**, 641-653 e617 (2022). https://doi.org:10.1016/j.cell.2022.01.013

Calder, L. J., Calcraft, T., Hussain, S., Harvey, R. & Rosenthal, P. B. Electron cryotomography of SARS-CoV-2 virions reveals cylinder-shaped particles with a double layer RNP assembly. *Commun Biol* **5**, 1210 (2022). https://doi.org:10.1038/s42003-022-04183-1

Song, Y. *et al.* In situ architecture and membrane fusion of SARS-CoV-2 Delta variant. *Proc Natl Acad Sci U S A* **120**, e2213332120 (2023). https://doi.org:10.1073/pnas.2213332120

Yao, H. *et al.* Molecular Architecture of the SARS-CoV-2 Virus. *Cell* **183**, 730-738 e713 (2020). https://doi.org:10.1016/j.cell.2020.09.018

Turonova, B. *et al.* In situ structural analysis of SARS-CoV-2 spike reveals flexibility mediated by three hinges. *Science* **370**, 203-208 (2020). https://doi.org:10.1126/science.abd5223

Mendonca, L. *et al.* Correlative multi-scale cryo-imaging unveils SARS-CoV-2 assembly and egress. *Nat Commun* **12**, 4629 (2021). https://doi.org:10.1038/s41467-021-24887-y

Zimmermann, L. & Chlanda, P. Cryo-electron tomography of viral infection - from applications to biosafety. *Curr Opin Virol* **61**, 101338 (2023). https://doi.org:10.1016/j.coviro.2023.101338

Hsieh, C. L. *et al.* Structure-based design of prefusion-stabilized SARS-CoV-2 spikes. *Science* **369**, 1501-1505 (2020). https://doi.org:10.1126/science.abd0826

21 Juraszek, J. *et al.* Stabilizing the closed SARS-CoV-2 spike trimer. *Nat Commun* **12**, 244 (2021).
https://doi.org:10.1038/s41467-020-20321-x

22 Claireaux, M. *et al.* A public antibody class recognizes an S2 epitope exposed on open
conformations of SARS-CoV-2 spike. *Nat Commun* **13**, 4539 (2022).
https://doi.org:10.1038/s41467-022-32232-0

23 Costello, S. M. *et al.* The SARS-CoV-2 spike reversibly samples an open-trimer conformation
exposing novel epitopes. *Nat Struct Mol Biol* **29**, 229-238 (2022). https://doi.org:10.1038/s41594-
022-00735-5

24 Ichikawa, T. *et al.* Chemical fixation creates nanoscale clusters on the cell surface by aggregating
membrane proteins. *Commun Biol* **5**, 487 (2022). https://doi.org:10.1038/s42003-022-03437-2

25 Silva, R. P. *et al.* Identification of a conserved S2 epitope present on spike proteins from all highly
pathogenic coronaviruses. *Elife* **12** (2023). https://doi.org:10.7554/eLife.83710

26 Zivanov, J. *et al.* A Bayesian approach to single-particle electron cryo-tomography in RELION-4.0.
Elife **11** (2022). https://doi.org:10.7554/eLife.83724

27 Bangaru, S. *et al.* Structural analysis of full-length SARS-CoV-2 spike protein from an advanced
vaccine candidate. *Science* **370**, 1089-1094 (2020). https://doi.org:10.1126/science.abe1502

28 Gobeil, S. M. *et al.* Effect of natural mutations of SARS-CoV-2 on spike structure, conformation,
and antigenicity. *Science* **373** (2021). https://doi.org:10.1126/science.abi6226

29 Henderson, R. *et al.* Controlling the SARS-CoV-2 spike glycoprotein conformation. *Nat Struct Mol*
Biol **27**, 925-933 (2020). https://doi.org:10.1038/s41594-020-0479-4

30 Dokainish, H. M. & Sugita, Y. Structural effects of spike protein D614G mutation in SARS-CoV-
2. *Biophys J* **122**, 2910-2920 (2023). https://doi.org:10.1016/j.bpj.2022.11.025

31 Zhou, D., Ren, J., Fry, E. E. & Stuart, D. I. Broadly neutralizing antibodies against COVID-19.
Curr Opin Virol **61**, 101332 (2023). https://doi.org:10.1016/j.coviro.2023.101332

32 Halfmann, P. J. *et al.* Multivalent S2-based vaccines provide broad protection against SARS-CoV-
2 variants of concern and pangolin coronaviruses. *EBioMedicine* **86**, 104341 (2022).
https://doi.org:10.1016/j.ebiom.2022.104341

33 Olukitibi, T. A. *et al.* Significance of Conserved Regions in Coronavirus Spike Protein for
Developing a Novel Vaccine against SARS-CoV-2 Infection. *Vaccines (Basel)* **11** (2023).
https://doi.org:10.3390/vaccines11030545

34 Huo, J. *et al.* Neutralization of SARS-CoV-2 by Destruction of the Prefusion Spike. *Cell Host*
Microbe **28**, 445-454 e446 (2020). https://doi.org:10.1016/j.chom.2020.06.010

35 Seow, J. *et al.* A neutralizing epitope on the SD1 domain of SARS-CoV-2 spike targeted following
infection and vaccination. *Cell Rep* **40**, 111276 (2022).
https://doi.org:10.1016/j.celrep.2022.111276

36 Bianchini, F. *et al.* Human neutralizing antibodies to cold linear epitopes and subdomain 1 of the
SARS-CoV-2 spike glycoprotein. *Sci Immunol* **8**, eade0958 (2023). https://doi.org:DOI:
10.1126/sciimmunol.ade0958

37 Li, J., Ulitzky, L., Silberstein, E., Taylor, D. R. & Viscidi, R. Immunogenicity and protection
efficacy of monomeric and trimeric recombinant SARS coronavirus spike protein subunit vaccine
candidates. *Viral Immunol* **26**, 126-132 (2013). https://doi.org:10.1089/vim.2012.0076

38 Dokainish, H. M. *et al.* The inherent flexibility of receptor binding domains in SARS-CoV-2 spike
protein. *Elife* **11** (2022). https://doi.org:10.7554/eLife.75720

39 Olia, A. S. *et al.* SARS-CoV-2 S2P spike ages through distinct states with altered immunogenicity.
J Biol Chem **297**, 101127 (2021). https://doi.org:https://doi.org/10.1016/j.jbc.2021.101127

40 Sasaki, M. *et al.* SARS-CoV-2 Bearing a Mutation at the S1/S2 Cleavage Site Exhibits Attenuated
Virulence and Confers Protective Immunity. *mBio* **12**, e0141521 (2021).
https://doi.org:10.1128/mBio.01415-21

41 Sasaki, M. *et al.* S-217622, a SARS-CoV-2 main protease inhibitor, decreases viral load and
ameliorates COVID-19 severity in hamsters. *Sci Transl Med* **15**, eabq4064 (2023).
https://doi.org:10.1126/scitranslmed.abq4064

42   Kremer, J. R., Mastronarde, D. N. & McIntosh, J. R. Computer visualization of three-dimensional
466   image data using IMOD. *J Struct Biol* **116**, 71-76 (1996). https://doi.org:10.1006/jsbi.1996.0013
467   43   Xiong, Q., Morphew, M. K., Schwartz, C. L., Hoenger, A. H. & Mastronarde, D. N. CTF
468   determination and correction for low dose tomographic tilt series. *J Struct Biol* **168**, 378-387 (2009).
469   https://doi.org:10.1016/j.jsb.2009.08.016
470   44   Tang, G. *et al.* EMAN2: an extensible image processing suite for electron microscopy. *J Struct Biol*
471   **157**, 38-46 (2007). https://doi.org:10.1016/j.jsb.2006.05.009
472   45   Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers, educators, and
473   developers. *Protein Sci* **30**, 70-82 (2021). https://doi.org:10.1002/pro.3943
474   46   Rohou, A. & Grigorieff, N. CTFFIND4: Fast and accurate defocus estimation from electron
475   micrographs. *J Struct Biol* **192**, 216-221 (2015). https://doi.org:10.1016/j.jsb.2015.08.008
476   47   Schrodinger, LLC. *The PyMOL Molecular Graphics System, Version 2.4*.
477
478

Acknowledgements

We thank Hiroshi Kida, Tatsuya Zenko, Atsushi Furukawa, Takashi Tadokoro, Robert

Parsons, Peijun Zhang, Fabian Eisenstein, Radostin Danev, Makoto Takeda for their support and

discussions. We also thank members of Facilities Division of Hokkaido University and

Thermofisher Scientific (Yoshihiro Tomatsu, Kazuhiro Aoyama, Yu Yong) for their technical

assistance. This work was supported in part by the Japan Agency for Medical Research and

Development (AMED) [JP17am0101093 (K.M.), JP20ae0101047 (K.M.), JP21fk0108463 (K.M.),

JP22ama121037 (K.M.), JP223fa627005 (K.M, K.T, M.S, Y.O, and H.S.), JP23ama121001 (T.S.)],

Hokkaido University Biosurface project (K.M.), Japan Science and Technology Agency (JST)

CREST grant number JPMJCR20H8 (H.F), JSPS Strategic Young Researcher Overseas Visits

Program for Accelerating Brain Circulation (K.M.), the Scientific Research on Innovative Areas

and International Group from the MEXT/JSPS KAKENHI [JP20H05873 (K.M)], 23HA2010 by

Ministry of Health, Labour and Welfare (MHLW) (H.S.), and Takeda Science Foundation (K.M.).

This research used computational resources of the supercomputer Fugaku (hp230052 and

hp220087) provided by the RIKEN Center for Computational Science (the evaluation environment

in the trial phase).

Author contributions

H.F., H.M.D., and K.M. designed the study; H.F., S.K., Y.S., H.S and K.M. set up BSL3

Cryo-EM facility; J.T.H. and D.I.S. assisted to set up and consult about BSL3 Cryo-EM facility;

H.F., K.T, M.S., Y.O., and H.S. prepared virus solution; H.F., S.K., K.T., Y.A., and H.S. performed

Cryo-ET data collection; H.M.D. performed Cryo-ET analysis; S.K., J.T.H., and K.M. advised

Cryo-ET analysis. A.S. T.S., and K.M. designed and prepared unique device of sample transfer for

Cryo-ET; H.M.D., H.F., and K.M. wrote the manuscript.

Competing interests

The authors declare no competing interests.

Data availability

All other data are available from the corresponding author upon request. Methods

describing structure determination are included in Supplementary Methods.

Snapshots from Cryo-ET of active SARS-CoV-2 virions

Hideo Fukuhara[1,2, #], Hisham M. Dokainish[2, #,*], Shunsuke Kita[2], Koshiro Tabata[3,4], Akira Takasu[5], Juha T. Huiskonen[6], Yuki Anraku[2], Toshiya Senda[5,7,8], David I. Stuart[9], Michihito Sasaki[3], Yasuko Orba[3], Yasuhiko Suzuki[4,10,*], Hirofumi Sawa[3,4,12,*] & Katsumi Maenaka[1,2,4,13,14,15,*].

Supplementary material

(Figure S1 to S7 and Table S1, S2)

Figure S1. BSL3 Cryo-EM facility at Hokkaido University and Cryo-ET analysis scheme.

Figure S2. Particle morphology and ice thickness. a) Tomogram slice of two virion particles undergoing fusion process. b) Side and central sections through the cryo-ET tomogram of the ancestral virions particles with different morphology. Ice thickness is highlighted by dotted line and reported in nm.

Spikes with shared Stem

Figure S3. 2D slice of full spike protein structures with shared stems.

Figure S4. Gold-standard Fourier shell correlation (FSC) curves for C3 and C1 symmetrized maps. Dotted line indicates the 0.143 FSC cutoff.

Figure S5. Charge distribution in Spike protomer. Electrostatic surface representation of the spike protein protomer from the high resolution structure (PDB 6ZGE)[1]. Highly charged regions are highlighted with dotted lines. Ribbon representation of the middle and top regions of S2 interface from two protomers with highlighted charged residues. Blue and red sticks represent positively and negatively charged amino acids, respectively.

Figure S6. SARS-CoV2 vaccine design approaches. Left: The cartoon representation of second generation of SARS-CoV-2 Vaccines that include 6 proline mutations (shown as orange spheres), inducing mainly RBD and NTD neutralizing Abs. Middle: S2-based vaccine approach that allow for targeting the S2 region. Right: Monomer-based vaccine approach to account for protein dynamic and target both S1 and S2 regions. Antibodies (blue and green) are schematically shown in all figures.

Figure S7. Conformational flexibility in the spike protein isolated monomer. a) Cartoon representation of different conformations of the head region of spike protomer obtained from three independent MD simulations. Two simulations form compact monomers conformations, while one simulation shows the opening of the S1 subunit (open monomer). b) Root mean square deviations (RMSD) of the Cα atoms upon fitting S2 region. All simulations show large deviation from a starting structure due to the motion of the S1 subunit.

Table S1. All particles diameter measurements (L1 and L2) in nm.

Ellipsoidal		Spherical		Oval		Bean-shape		Cylindrical	
L1	*L2*	*L1*	*L2*	*L1*	*L2*	*L1*	*L2*	*L1*	*L2*
101	88	90	81	122	89	86	51	105	61
125	100	105	104	107	101	109	65	94	49
125	119	107	102	97	83	119	66	91	50
150	125	102	93	71	76	89	79	92	73
147	88	105	101			94	55	83	52
99	83	87	86			103.0	75.0		
139	102	88	84						
108	86	108	102						
109	95	104	98						
97	81	85	82						
102	74	109	105						
103	87	83	78						
		94	94						
		106	94						
		142	138						
		90	86						
		98	98						
		94	91						
		86	82						

Table S2. Cryo-ET data acquisition and reconstruction statistics.

Data Collection		
Microscope	Titan Krios G4	
Magnification	64,000	
Voltage (kV)	300	
Detector	Falcon 4	
Energy filter	Selectris	
Pixel size (Å)	1.884	
Tilt Schemes	Dose-symmetric scheme	
Tilt angle increment (degree)	1	
Tilt angle range	-60 to +60	
Number of tilt Series	42	
Number of virions	46	
Defocus	0.2 using Volta Phase Plate	
Software	Tomography 4	
Total dose (e/Å^2)	181.5	
Reconstruction		
Software	IMOD, Relion	
Dataset	Prefusion spike	
Number of spikes	237	
Symmetry	C3	C1
Final resolution (Å)	14.3	17.0
Gold-standard	yes	yes
FSC threshold	0.143	0.143
Final pixelsize (Å)	1.884	1.884

Supplementary Method:

MD simulations of Spike monomer.

The monomeric spike protein model was based on the Omicron BA.2 Cryo-EM structure (PDB 7UB0)[2]. CHARMM-GUI was used to add missing residues, ions, water molecules for preparing final simulation model[3]. Glycan and disulfide bonds were added in a similar fashion to previous study[4]. Considering the potential large motion in monomer spike protein, a large box size was used with total number of atoms of 742586. All simulations were performed using GENESIS 2.0 Beta MD software on Fugaku supercomputer[5,6]. Three independent simulations for 100 ns each was performed using CHARMM 36M force field for protein residues and CHARMM TIP3P for ions and water molecules[7,8]. VMD software was used for trajectory visualization [9].

References

Wrobel, A. G. *et al.* SARS-CoV-2 and bat RaTG13 spike glycoprotein structures inform on virus evolution and furin-cleavage effects. *Nat Struct Mol Biol* **27**, 763-767 (2020). https://doi.org:10.1038/s41594-020-0468-7

Stalls, V. *et al.* Cryo-EM structures of SARS-CoV-2 Omicron BA.2 spike. *Cell Rep* **39**, 111009 (2022). https://doi.org:10.1016/j.celrep.2022.111009

Lee, J. *et al.* CHARMM-GUI Input Generator for NAMD, GROMACS, AMBER, OpenMM, and CHARMM/OpenMM Simulations Using the CHARMM36 Additive Force Field. *J. Chem. Theory Comput.* **12**, 405-413 (2016). https://doi.org:10.1021/acs.jctc.5b00935

Dokainish, H. M. *et al.* The inherent flexibility of receptor binding domains in SARS-CoV-2 spike protein. *Elife* **11** (2022). https://doi.org:10.7554/eLife.75720

Kobayashi, C. *et al.* GENESIS 1.1: A hybrid-parallel molecular dynamics simulator with enhanced sampling algorithms on multiple computational platforms. *J Comput Chem* **38**, 2193-2206 (2017). https://doi.org:10.1002/jcc.24874

Jung, J. *et al.* New parallel computing algorithm of molecular dynamics for extremely huge scale biological systems. *J Comput Chem* **42**, 231-241 (2021). https://doi.org:10.1002/jcc.26450

Huang, J. *et al.* CHARMM36m: an improved force field for folded and intrinsically disordered proteins. *Nat. Methods* **14**, 71-73 (2017). https://doi.org:10.1038/nmeth.4067

Guvench, O. *et al.* CHARMM additive all-atom force field for carbohydrate derivatives and its utility in polysaccharide and carbohydrate-protein modeling. *J. Chem. Theory Comput.* **7**, 3162-3180 (2011). https://doi.org:10.1021/ct200328p

Humphrey, W., Dalke, A. & Schulten, K. VMD: visual molecular dynamics. *J Mol Graph* **14**, 33-38, 27-38 (1996). https://doi.org:10.1016/0263-7855(96)00018-5