Complex genomic landscape of inversion polymorphism in Europe's most destructive forest

pest

Anastasiia Mykhailenko[1,2], Piotr Zieliński[1], Aleksandra Bednarz[1], Fredrik Schlyter[3,4], Martin N.

Andersson[5], Bernardo Antunes[1], Zbigniew Borowski[6], Paal Krokene[7], Markus Melin[8], Julia

Morales-García[1,2], Jörg Müller[9,10], Zuzanna Nowak[1], Martin Schebeck[11], Christian Stauffer[11],

Heli Viiri[12], Julia Zaborowska[1], Wiesław Babik[1], Krystyna Nadachowska-Brzyska[1]

Institute of Environmental Sciences, Faculty of Biology, Jagiellonian University, Gronostajowa 7, 30-387,

Kraków, Poland

²Doctoral School of Exact and Natural Sciences, Jagiellonian University, ul. Łojasiewicza 11, 30-348,

Kraków, Poland

³Chemical Ecology, Department of Plant Protection Biology, Swedish University of Agricultural Sciences

Alnarp, PO Box 190, 234 22 Lomma, Sweden

⁴ETM, Faculty of Forestry and Wood Sciences, Czech University of Life Sciences Prague, Kamýcká 129, 165

00, Praha, Czechia

⁵Department of Biology, Lund University, Sölvegatan 37, 223 62, Lund, Sweden

⁶Forest Research Institute, Sękocin Stary, Braci Leśnej 3, 05–090 Raszyn, Poland

⁷Division of Biotechnology and Plant Health, Norwegian Institute of Bioeconomy Research, Høgskoleveien 8,

Ås, 1433, Norway

⁸Natural Resources Institute Finland, Yliopistokatu 6b, 80100 Joensuu, Finland

⁹Field Station Fabrikschleichach, Animal Ecology and Tropical Biology, Biocenter, University of Würzburg,

Glashüttenstrasse 5, 96181 Rauhenebrach, Germany

¹⁰Bavarian Forest National Park, Freyungerstrasse 2, 94481 Grafenau, Germany

¹¹Institute of Forest Entomology, Forest Pathology and Forest Protection, Department of Forest and Soil

Sciences, University of Natural Resources and Life Sciences Vienna (BOKU), Peter-Jordan-Straße 82/I, 1190,

Vienna, Austria

¹²UPM-Kymmene, UPM Forest, BOX 85, 33100, Tampere, Finland

*Corresponding author: Krystyna Nadachowska-Brzyska. email: krystyna.nadachowska@uj.edu.pl

Abstract

In many species, polymorphic inversions underlie complex phenotypic polymorphisms and

facilitate local adaptation in the face of gene flow. Multiple polymorphic inversions can co-occur

in a genome, but the prevalence, evolutionary significance, and limits to complexity of genomic

inversion landscapes remain poorly understood. Here, we examine genome-wide variation in one of Europe's most destructive forest pests, the spruce bark beetle *Ips typographus*, scan for polymorphic inversions, and test whether inversions are involved in key adaptations in this species. We analyzed 240 individuals from 18 populations across the species' European range and, using a whole-genome resequencing approach, identified 27 polymorphic inversions covering approximately 28% of the genome. The inversions vary in size and in levels of intra-inversion recombination, are highly polymorphic across the species range, and often overlap, forming a complex genomic architecture. We test several mechanisms, including directional selection, overdominance and associative overdominance that can contribute to the maintenance of inversion polymorphisms in the genome. We show that the heterogeneous inversion landscape is likely maintained by the combined action of several evolutionary forces and that inversions are enriched in odorant receptor genes encoding key elements of recognition pathways for host plants, mates, and symbiotic fungi. Our results indicate that the genome of this major forest pest of growing social, political, and economic importance harbors one of the most complex inversion landscapes described to date posing a question about limits of genomic architecture complexity.

Introduction

Large structural variants (SVs), such as chromosomal inversions, translocations, insertions, and duplications, were among the earliest mutations described in natural populations (McClung 1905; Sturtevant 1921), but their systematic identification and in-depth analysis has only become possible with recent advances in sequencing technology (Wellenreuther et al. 2019). The increasing accumulation of genomic data has revealed not only the presence of structural variation between and within many species but also its important role in species adaptation and speciation (Lamichhaney et al. 2015; Hof et al. 2016; Cheng et al. 2018; Wellenreuther and Bernatchez 2018; Faria, Chaube, et al. 2019; Todesco et al. 2020; Mérot et al. 2021; Harringmeyer and Hoekstra 2022; Saitou et al. 2022). In particular, polymorphic chromosomal inversions have been at the center of recent debate about their potential to redirect the evolutionary process (Berdan et al. 2023).

Polymorphic inversions are chromosomal segments that occur in two orientations within populations: collinear and inverted haplotypes/arrangements. Inversions have been shown to be

involved in speciation, local adaptation, and maintenance of complex phenotypes (Lamichhaney

et al. 2015; Lohse et al. 2015; Wellenreuther and Bernatchez 2018; Fuller et al. 2019). This is

due to a key property of inversions: they suppress recombination within heterozygotes and

thereby prevent separation of coadapted variants. Because of their role as recombination

modifiers, inversions can act as supergenes, i.e., large elements of genomic architecture with

multiple linked functional elements (Thompson and Jiggins 2014). Supergenes keep coadapted

alleles together in the face of gene flow and prevent the formation of maladaptive recombinant

genotypes.

Classic examples of supergenes include inversions associated with different mating strategies in

ruffs (*Calidris pugnax*; Lamichhaney et al. 2015), mimicry phenotypes in *Heliconius* butterflies

(Joron et al. 2011), and social organization in fire ants (*Solenopsis* spp.; Purcell et al. 2014;

Gutiérrez-Valencia et al. 2021). In many other species, polymorphic inversions define locally

adapted ecotypes (Lowry and Willis 2010; Kirubakaran et al. 2016; Koch et al. 2021; Matschiner

et al. 2022; Reeve et al. 2023) or exhibit spatial frequency differences, e.g. by forming

geographic and climatic gradients (Ayala et al. 2014; Kapun et al. 2016; Ayala et al. 2017; Mérot

et al. 2021). While the vast majority of described cases are organisms with one or a few

inversions, several recent studies have reported species with many polymorphic inversions

(Faria, Chaube, et al. 2019; Todesco et al. 2020; Harringmeyer and Hoekstra 2022; Porubsky et

al. 2022; Reeve et al. 2023). These recent findings raise questions about the prevalence and

evolutionary significance of polymorphic inversions in natural populations. Are inversion-rich

genomes the exception or the rule? How much of the genome can be situated within polymorphic

inversions and, consequently, how large can the fraction of the genome with reduced

recombination be? The latter question is particularly important because, in addition to

suppressing recombination and keeping coadapted alleles together, inversion heterozygotes will

also prevent the formation of new allelic combinations and thus reduce the efficacy of natural

selection (Roesti et al. 2022).

Given the importance of recombination in purging deleterious mutations (Keightley and Otto

2006), a substantial fraction of the genome that is situated within polymorphic inversions raises

other questions. First, as the degree of recombination suppression depends on the number of

heterozygous genotypes, what is the probability of an individual being heterozygous at multiple

inversion regions? Second, are polymorphic inversions where both arrangements are equally

common across the species' range more common than inversions with one common and one rare

haplotype? Third, how common are mechanisms that potentially can mitigate negative

consequences of reduced recombination? Such mechanisms include double crossover-events and

gene conversion in heterozygotes, which can reshuffle allelic content between large parts of two

inversion arrangements and thereby reduce mutational load and create new haplotypes (Berdan et

al. 2021; Charlesworth 2023).

Long-term persistence of two inversion haplotypes can be facilitated by two main types of

selection: divergent and balancing (Faria, Johannesson, et al. 2019). Divergent selection can

favor different inversion genotypes in different environments and, when coupled with reduced

migration between divergent populations, can lead to speciation. Even when intraspecific gene

flow is high, divergent selection can lead to divergent ecotypes associated with locally

advantageous inversion genotypes. Alternatively, balancing selection may maintain balanced

inversion polymorphisms over time via several, not mutually exclusive, mechanisms, such as

overdominance, negative frequency dependence, antagonistic pleiotropy, and spatially or

temporally varying selection (Connallon and Clark 2014; Faria, Johannesson, et al. 2019).

Importantly, regardless of the selection mechanism, inversions will accumulate mutations

independently since recombination is suppressed in inversion heterozygotes. This will lead to

differentiated allelic content and increased differentiation between inversion haplotypes over

time (Faria, Johannesson, et al. 2019). Each inversion haplotype can thus be treated as a separate

"population" with a size that corresponds to the frequency of that arrangement within studied

population or species. Rare inversion arrangements will experience a high mutational load due to

reduced recombination and limited purging because of rarity of heterozygotes. However,

deleterious mutations can also accumulate on more frequent inversion haplotypes (Berdan et al.

2021) and lead to associative overdominance. This contributes to the maintenance of inversion

polymorphisms since independent accumulation of mutations continues over time but recessive

deleterious alleles private to an inversion arrangement are not visible to selection in inversion

heterozygotes.

The Eurasian spruce bark beetle (*Ips typographus* (L.): Curculionidae: Scolytinae; hereafter the

spruce bark beetle), plays a key role in Eurasian forest ecosystems. Under endemic conditions, it

attacks weakened Norway spruce (*Picea abies*) trees. However, if spruce resistance is

compromised by certain abiotic disturbances (e.g. snowbreaks, windfalls, high temperatures,

drought), an increased availability of stressed trees can trigger mass-propagation, leading to rapid

population increase and devastating outbreaks. The extent of recent outbreaks is unprecedented,

and impacts will likely increase in the coming decades in response to climate change

(Biedermann et al. 2019; Bentz et al. 2021; Müller et al. 2022). For example, during the first

decade of the 21st century the spruce bark beetle killed an estimated 14.5 million m^3 of timber

per year on average. The Czech Republic provides a particularly striking example of the beetles'

destructive potential. During the peak outbreak years 2017-2019 the beetles killed annually 3.1-

5.4% of the country's growing stock of Norway spruce, which in 2019 translated to 23 million

m^3. Historically, spruce forests in Central Europe have been most heavily affected by bark beetle

outbreaks, while in northern Europe outbreaks have been less frequent and destructive (Hlásny et

al. 2019). However, this may change with climate warming that probably will make the boreal

forests of northern Europe more vulnerable to bark beetle outbreaks (Lange et al. 2006; Jönsson

et al. 2007; Müller 2011; Bentz et al. 2021; Müller et al. 2022). As an example, heatwaves and

severe summer drought in Sweden in 2018 initiated a bark beetle outbreak killing over 30 million

m^3 Norway spruce in the next years (Öhrn et al. 2021; Wulff and Roberge 2022).

Increasing bark beetle attacks and other forest disturbances have already triggered social and

political conflicts in parts of Europe and have highlighted the urgent need for improved

management strategies (Müller 2011; Vega and Hofstetter 2015; Tomáš Hlásny et al. 2021).

Indeed, a rapidly growing body of research focuses on the species' ecology, and the causes and

consequences of outbreaks (reviewed in: Wermelinger 2004; Vega and Hofstetter 2015;

Biedermann et al. 2019; Wermelinger and Jakoby 2022). Despite this enormous interest, one

aspect of the species' biology remains largely unexplored: we know almost nothing about the

species' genome-wide variation and the evolutionary mechanisms that shape this variation. The

lack of population genomics studies restricts our understanding of the genomic basis of

adaptation and adaptive potential in the spruce bark beetle. Such information could provide a

critical missing link between applied and basic research and serve as a foundation for effective

management. This is particularly important because, as revealed in this study, the spruce bark

beetle genome harbors a complex inversion polymorphism landscape that may play a critical role

in many evolutionary processes, including key species adaptations (Kirkpatrick and Barton 2006;

Wellenreuther and Bernatchez 2018; Berdan et al. 2021).

Here, we investigated genome-wide variation across spruce bark beetle populations with a
special focus on chromosomal inversion polymorphisms. We found one of the most complex
polymorphic inversion architectures described to date and investigated several evolutionary
mechanisms, including directional selection, overdominance and associative overdominance, that
can maintain inversion polymorphisms in the genome. We also tested associations between
inversion polymorphisms and two key fitness-related traits - diapause and olfaction. Our results
suggest that inversions are involved in key species adaptations and raise questions about the
prevalence and role of inversion polymorphisms in species characterized by large effective
population sizes, little geographic subdivision, and no clear phenotypic variation across the
species range.

Results and Discussion
We resequenced whole genomes of the spruce bark beetles from 18 European populations
(Figure 1, Table S1). After initial raw data processing from 253 individuals (including data
quality check, trimming and GATK guided variant calling, genotyping, recalibration and
filtering; for details see Supplementary Information), we used 240 individuals in downstream
analyses. The mean per individual sequencing depth was 23.2x (range: 5 – 53x). Sequencing
coverage on IpsContig9 was consistently lower in individuals sexed as males (on average 0.57
individual coverage). Thus, we considered IpsContig 9 to be a sex (X) chromosome (females
carry XX and males XYp chromosomes). After quality filtering, we retained 5.245 million SNPs
covering the entire length of the genome assembly (236.8 Mb) but analyzed a subset of 5.067
million SNPs located on the 36 longest contigs and that represented 78% of the assembly.
Alignment of the *Ips typographus* and *Ips nitidus* genome assemblies indicated no major
misassembly problems (*Ips typographus* contigs align to single *Ips nitidus* chromosomes) and
high synteny between karyotypes of both species (Figure S1). Given these results, contig sizes,
genome assembly quality scores and species karyotype, it is likely that many spruce bark beetle
contigs represent entire chromosome arms. Such genome quality is sufficient for all downstream
analyses, in particular SNP-based polymorphic inversion identification (Mérot et al. 2021; Reeve
et al. 2023).

Complex genomic inversion landscape

Twenty-nine candidate inversions ("inversions" henceforth) were identified following the criteria

described in the Methods section (Table 1, Figure 2, Figure S2, and Figure S3). Briefly, we

considered a genomic region to harbor a polymorphic inversion if local PCA analysis identified

the region as an outlier, and/or the region exhibited high linkage disequilibrium (LD) and PCA

performed on SNPs from this region separated individuals not by geography but into three

distinct groups (with a group classified as putative heterozygous individuals characterized by

higher heterozygosity compared to the other groups). Two putative inversions (Inv16.1 and

Inv23.1) were most likely part of the same inversion: they are in strong LD (Figure S4) and the

same beetles were genotyped as homo- and heterozygotes in both, which is unlikely to occur for

two unlinked inversions. The same situation was found for Inv16.2 and Inv23.2. No other

inversions were in strong LD with each other, which would suggest co-segregation (Figure S4).

Thus, overall, we found 27 inversions in 17 contigs, including one located on the X chromosome

(Inv9). Approximate inversion sizes varied from 0.1 to 10.8 Mb (Table 1) and inversions

constituted approximately 28% of the analyzed part of the genome and at least 18.6% of the

entire assembly. Estimated inversion ages ranged from 0.5 to 2.6 My (Table 1, assuming a

mutation rate of 2.9×10^{-9}; see Table S1 for results for different mutation rates) and were

strongly negatively correlated with the major inversion haplotype frequency (Figure S5).

Inversion regions exhibited a reduced population recombination rate in heterozygous individuals

(Figure S2) and moderate to high genetic differentiation between inversion arrangements (F_{ST}

between homozygous individuals was 0.15-0.64; Figure 1; Figure S2).

While 12 contigs contained single inversions, five contigs showed patterns consistent with

multiple adjacent or overlapping inversions (Figure 2, Figure S2, Figure S3). IpsContig14 and

IpsContig22 contained complexes of multiple adjacent and sometimes overlapping inversions

(Figure 1 J-O), and three other contigs contained two overlapping inversions each (IpsContigs: 7,

16 and 23, Figure S2 and Figure S3). Additionally, four putative double-crossover events were

identified within four inversions (Inv5, Inv18, Inv22.3, Inv22.4; Figure 1, Figure S2 and Figure

S3).

While the variation patterns observed on contigs with single inversions were clear and leave no

doubt about the presence of polymorphic inversions, the complex patterns (overlapping

inversions and inversions with double crossover events) were more difficult to interpret and

deserve additional explanation and caution. Putative overlapping inversions were identified on

the basis of overlapping clusters of high LD, PCA clustering into three distinct groups along PC2

or PC3 (in addition to the three distinct groups along PC1), and moderate to high F_{ST} between

individuals classified as alternatively homozygous (Figure 2 and Figure S3). In the case of

putative double-crossover events, the observed LD clusters were separated by regions of lower

LD and intermediate groups of individuals were visible between the main clusters along the PC1

(Figure S3). The F_{ST} between individuals classified as alternatively homozygotes was

significantly reduced in the region of the putative double-crossover (Figure 2 and Figure S2, S3).

The most difficult to disentangle patterns were associated with Inv2, Inv5, Inv7.1 and Inv7.2.

Putative Inv2 showed a signature of overlapping LD clusters (overlapping inversions) but

ambiguous PCA clustering (Figure S3). The patterns were clear and consistent with inversion

polymorphisms when only populations from the southern region were analyzed (Figure S6; thus,

Inv2 was genotyped only in this part of the species range), suggesting other structural differences

between the southern and northern regions or unidentified genome assembly problems. Principal

component analysis of Inv5 identified three clusters that most likely correspond to three

inversion genotypes, but also several smaller clusters that may indicate multiple independent

recombination events (double-crossover events that occurred at different locations within Inv5,

Figure S2 and Figure S3) or some assembly problems within Inv5 (however, IpsContig5 aligns

perfectly with *Ips nitidus* chromosome 9, Figure S1). Thus, Inv5 was only genotyped for three

major genotype groups for which multiple lines of evidence (LD, heterozygosity, F_{ST}) suggested

an inversion polymorphism. Inv7.1 and Inv7.2 are characterized by overlapping LD clusters, and

PCA and F_{ST} patterns consistent with overlapping inversions. However, Inv7.2 showed low LD

in the middle of the inversion and unexpected clustering of individuals in one of the clusters

along PC1 – the cluster was split into two distinct groups with individuals clustered according to

their geographic location within each of these groups (Figure S2 and Figure S3). This suggests

possible misassembly and it is likely that Inv7.2 is shorter than predicted from the size of the LD

boundaries. Genotyping of Inv16.2 and Inv23.2 was only possible in part of the species range, as

PCA clustering into three genotype groups was only visible when analyzing the northern group

only (Figure S6).

Even though we are not able to rule out that some of the patterns were created or disturbed by

potential misassembly or additional structural rearrangements (e.g., duplications; Kim et al.

2022), the evidence for the vast majority of polymorphic inversions in the spruce bark beetle is

strong. Extensive collinearity with *Ips nitidus* suggested no major misassembly problems and

further supports our identification of polymorphic inversions in the spruce bark beetle genome

(Figure S1; Wang et al. 2023). Long-read sequencing and Hi-C data is needed to shed light on

the problematic/complex cases as well as to precisely identify the inversion boundaries. It was

not surprising that we detected polymorphic inversions in the spruce bark beetle genome, as

there are many well-known examples of polymorphic inversions in natural populations

(Wellenreuther and Bernatchez 2018). What was striking, however, was the complex genomic

landscape of polymorphic inversions we found in this species. The spruce bark beetle has at least

27 large inversions covering a substantial part of the genome (28% of the analyzed part of the

genome). Numerous (a dozen or more) polymorphic inversions have been described for several

species (e.g. Faria, Chaube, et al. 2019; Tigano et al. 2021; Harringmeyer and Hoekstra 2022)

but it remains an open question whether many polymorphic inversions within species is the

exception or the rule.

The exceptionally complex inversion architecture we found in the spruce bark beetle, with

multiple adjacent and often overlapping inversions, resembles well known examples from

Heliconius butterflies (Jay et al. 2021) or fire ants (Wang et al. 2013). In these insects, multiple

adjacent inversions are the basis for mimicry phenotypes and complex social organization,

respectively. The presence of clusters of adjacent inversions and inversion overlaps are

consistent with theoretical expectations of stepwise extension of recombination suppression on

supergenes (Jay et al. 2022) and with a highly polygenic architecture of adaptation (Schaal et al.

2022).

Genome-wide variation and its geographic structuring – collinear vs. inversion regions

Analyses based on both the whole genome and collinear parts only revealed a clear latitudinal

structuring of genetic variation in the spruce bark beetle (Figure 1). Both PCA and NGSadmix

supported the presence of two distinct genetic groups corresponding to southern and northern

populations, with Polish populations showing varying degrees of admixture between the two

clusters. Based on these results, we divided the 18 studied populations into a northern, southern,

and Polish group. Despite unambiguous NGSadmix division into two genetic clusters, the

genome-wide genetic differentiation between the northern and southern group was very low (F_{ST}

$= 0.021$). Similarly, F_{ST} between all population pairs showed low levels of differentiation,

ranging from 0.000 to 0.035 (calculated excluding inversions; Table S2). Mean genome-wide

nucleotide diversity was moderate ($\pi = 0.0062$) and per population π ranged from 0.0055 to

0.0066 (Figure S7). There was a weak negative correlation between nucleotide diversity and

latitude ($r^2 = 0.32$, p = 0.033, Figure S8), as northern populations had slightly lower genetic

variation than southern populations ($\pi_{southern} = 0.0065$; $\pi_{northern} = 0.0061$; $\pi_{Polish} = 0.0066$).

Southern populations had an excess of rare alleles and, consequently, had more negative

Tajima's D values along the genome than northern populations (mean Tajima's D was -0.458

and -0.062 in the southern and northern group, respectively; Figure S7). All these results are

consistent with previous phylogeographic studies of the spruce bark beetle that analyzed a much

smaller number of genetic markers. The data suggests high levels of connectivity among

populations and a very recent differentiation into two genetic clusters (Stauffer et al. 1999; Sallé

et al. 2007; Bertheau et al. 2013; Mayer et al. 2015). More recent RADseq data confirms a very

weak genetic structure in the spruce bark beetle across much of Sweden (Ellerstrand et al. 2022),

as is expected in a species with high dispersal (Nilssen 1984) and/or recent divergence. Tajima's

D values indicate different demographic history of southern and northern group (e.g., recent

expansion of southern populations and bottleneck in northern populations, Figure S7).

Inversion regions in the spruce bark beetle did not structure in the same way as the collinear part

of the genome and in most of the cases do not show any clustering into a southern and northern

group (Figure S2). Frequencies of five inversions show significant correlation with latitude

(Figure 3; r^2 ranged from 0.34 to 0.68; see more discussion below). Almost all identified

inversions were polymorphic across the European species range, except for one inversion (Inv9)

that was polymorphic only in northern populations. For three inversions (Inv2, Inv5,

Inv16.2+Inv23.2) unambiguous genotyping was only possible across part of the species range

(see above). The differentiation between inversion haplotypes was moderate or high (mean F_{ST}

range 0.15-0.64), (Figure 2; Figure S2) and, according to our age estimates, the origin of the

inversions in the spruce bark beetle predates the Last Glacial Period. Many inversions may be

several million years old, likely also predating the within-species differentiation into a southern

and northern group. This was not unexpected, as many known inversion polymorphisms have

been segregating within species for hundreds of thousands or millions of years (see table in

(Wellenreuther and Bernatchez 2018)), sometimes even persisting through multiple speciation

events (Brelsford et al. 2020).

Inversions and key fitness-related traits

Inversion polymorphism is often associated with the maintenance of complex polymorphic

phenotypes (Schwander et al. 2014; Lamichhaney et al. 2015). Although the spruce bark beetle

does not exhibit easily identifiable phenotypes, such as distinct color patterns or mating types,

we were able to test for associations between inversions and two complex traits of key

importance for many insects: diapause and olfaction. Diapause allows species to suspend

development during unfavorable conditions. While multiple environmental factors can influence

this complex process, many aspects of diapause, such as its induction and termination, are

heritable (Roff 1996) and may be controlled by a small number of loci or be highly polygenic

(Paolucci et al. 2016; Pruisscher et al. 2018; Kozak et al. 2019; Denlinger 2022). The spruce bark

beetle exhibits two diapause strategies that could be associated with polymorphic inversions: a

facultative photoperiod-regulated diapause and an obligate photoperiod-independent diapause

(Schebeck et al. 2022). However, we found no association between diapause phenotypes and

inversion genotypes (exact G test, Table S3, Figure S9), nor did we find any highly differentiated

genomic regions between facultatively and obligately diapausing individuals (Figure S10). This

suggests a polygenic nature of diapause phenotypes in the spruce bark beetle.

Olfaction is another key fitness-related trait in many insects, including bark beetles. Insect

odorant receptors (ORs) are encoded by a large and dynamically evolving gene family. Some of

the receptors are evolutionarily conserved across species within insect orders, however, many are

species- or genus-specific. In the spruce bark beetle, detection of odorants is essential for host

and mate finding, as well as recognition and maintenance of symbiosis with specific fungi

(Hansson and Stensmyr 2011; Kandasamy et al. 2019). We examined 73 antennally expressed

ORs (Yuvaraj et al. 2021) and found that 46% of these were located within inverted regions

(Table 1), even though inverted regions constituted only 28% of the analyzed part of the genome.

A permutation test confirmed this over-representation of OR genes within inversions (p = 0.03).

In addition, several *Ips*-specific ORs (5 out of 7 ORs from an *Ips*-specific OR clade (Hou et al.

2021)) were located in inverted regions, specifically on IpsContig13 (4 out of 7 ORs). These 5

ORs (ItypOR23, ItypOR27, ItypOR28, ItypOR29, and ItypOR49) have been functionally
characterized, responding to compounds primarily produced by beetles (pheromones), the host
tree, or fungal symbionts (Hou et al. 2021; Powell et al. 2021). We found no difference in the OR
number (no OR deletions) between inversion haplotypes. The majority of ORs located in
inversions harbor multiple nonsynonymous variants segregating within the spruce bark beetle
populations (Table S4) and nine ORs (Table S5) had at least one fixed or nearly fixed (d_{xy}>0.9)
nonsynonymous variant between inversion haplotypes. These diverged ORs were located at Inv5
(ItypOR43JOI, ItypOR29), Inv13 (ItypOR23, ItypOR27), Inv14.1 (ItypOR20NTE, ItypOR36)
and Inv16.1 (ItypOR30, ItypOR31, ItypOR16). Among these ORs only ItypOR30 had dN/dS >1
(2 nonsynonymous substitutions, McDonald-Kreitman test p=0.55).

Two inversions harboring multiple OR genes showed significant latitudinal variation (Figure 3;
Inv13 and Inv16.1+23.1, Table 1). Olfactory receptor activity and odorant binding were also
among the enriched gene ontology categories in inversions that showed a correlation between
inversion haplotype frequency and latitude (Figure 3, Table S6; $0.01 < p < 0.05$). In *Drosophila
pseudoobscura* odorant receptors were also associated with inversions (Fuller et al. 2016; Fuller
et al. 2017). Interestingly, one of the inversions, Inv13, includes a gene encoding ItypOR23,
receptor that has been previously shown to primarily respond to an odor from fungi (Hou et al.
2021). This OR was also among the most divergent between alternative inversion haplotypes
(Table S5). We hypothesize that different OR alleles associated with these inversions may be
involved in spruce bark beetle interactions with fungal associates present in different parts of
Europe. As many other bark beetle species, *I. typographus* is associated with several, spatially
differentiated, symbiotic fungal species that can help beetles to exhaust and overcome tree's
defense system (Lieutier et al. 2009). Interestingly, recent studies suggested that beetles can
recognize volatile organic compounds of specific beneficial fungi that can be located and picked
up and that individual beetles may also have preferences for different fungal species (Kandasamy
et al. 2019; Zhao et al. 2019; Kandasamy et al. 2023).German spruce bark beetles have, for
example, been shown to be more attracted to fungal species that are common in Germany
(*Grosmannia penicillata, Endoconidiophora polonica*) than to rarer species (*Leptographium
europhioides*). Unpublished data from Swedish beetles suggests that they are more attracted to *L.
europhioides*, which is common in Sweden (personal communication D. Kandasamy). Although
preliminary, these observations suggest that beetle preferences may be tuned to the local fungal

community, and we speculate that inversions may be involved in recognition of region-specific

fungal species.

Several other interesting behavioral strategies are polymorphic among spruce bark beetle

individuals, including the existence of pioneer individuals that are the first to infest host trees,

and re-emergence of females after egg laying to establish so-called sister broods in new trees.

Other less obvious/visible phenotypes could also be associated with inversion polymorphisms.

Gene ontology enrichment analysis provided a list of categories of interest (Table S7) but

comprehensive research, both bottom-up and top-down, is needed to understand the relationship

between spruce bark beetle phenotypes and the inversion polymorphism landscape.

Evolutionary mechanisms maintaining inversion polymorphism in the spruce bark beetle

Several not mutually exclusive evolutionary mechanisms can maintain polymorphic inversions

within species, in particular divergent and balancing selection (Wellenreuther and Bernatchez

2018; Faria, Johannesson, et al. 2019). The importance of divergent selection has been postulated

based on allele frequency patterns and associations of polymorphic inversions with local

adaptations that persist despite extensive intraspecific gene flow (Tigano and Friesen 2016). For

example, a recent study of deer mice (*Peromyscus maniculatus*) (Harringmeyer and Hoekstra

2022) identified multiple polymorphic inversions with clinal variation across environmental

gradients in two distinct habitats. Such frequency changes have also been reported across hybrid

zones (Faria, Chaube, et al. 2019) or latitudinal gradients (Mérot et al. 2021). Although the

spruce bark beetle does not occupy distinct environmental niches, it inhabits forests across a very

wide latitudinal gradient (spanning at least 16 degrees). It is also a species with high dispersal

capacity and extensive gene flow, as indicated by low F_{ST} across its range. We found a

significant correlation between the frequency of inversion haplotypes in populations and

geographic location (latitude) for five inversions (Figure 3; r^2 ranged from 0.34 to 0.68). There

were no significant correlations between haplotype frequencies and longitude (Figure S11). In

most cases, differences in inversion haplotype frequencies were small (Figure 3), except for the

two inversions with the strongest correlations ($r^2 > 0.6$; Inv12 and Inv13). Climate and land

cover variation (summarized via PCA) was associated with allele frequency changes of multiple

SNPs (59), however, no such association was found for inversion genotypes (Figure 4). The

identified SNPs were in the proximity of 14 genes, five of which had assigned gene ontology

categories (Table S8). All the above results indicated that probably there is selection across

environmental gradients in the spruce bark beetle but that this is not the only, or even a major,

force maintaining inversion polymorphism within the species.

While the 'local adaptation' is a major hypothesis proposed to explain inversion polymorphism

(Christmas et al. 2019; Akopyan et al. 2022; Harringmeyer and Hoekstra 2022), balancing

selection and related mechanisms may also be important in maintaining polymorphic

arrangements (Faria, Johannesson, et al. 2019; Mérot et al. 2020; Berdan et al. 2021). Such

mechanisms include overdominance, associative overdominance, frequency-dependent selection,

and spatially and temporally varying selection. We found no support for overdominance playing

a role in the spruce bark beetle, as no excess of inversion heterozygotes compared to Hardy-

Weinberg expectations was detected in any of the populations. The only significant deviations

from Hardy-Weinberg expectations were detected in Inv12 and Inv22.4 in geographic regions

and/or across the whole species range (Table S9). We therefore looked more closely at mutation

load, which can say something about the role of associative overdominance in the maintenance

of inversion polymorphism. Theory predicts that recessive deleterious mutations will accumulate

on both inversion arrangements but that most of these mutations will be private to only one

arrangement (Navarro et al. 2000; Faria, Johannesson, et al. 2019; Berdan et al. 2021). This

would lead to associative overdominance, as in heterozygotes the effects of deleterious recessive

alleles on one arrangement would be masked by the wild-type alleles on the other arrangement.

The result would be long-term maintenance of the inversion polymorphism, resulting in strong

divergence between inversion haplotypes (Navarro et al. 2000; Guerrero et al. 2012; Berdan et

al. 2021).

Interestingly, several stable evolutionary scenarios that maintain polymorphic inversions are

possible (for details see Figure 4 in (Berdan et al. 2021)). These scenarios differ in the expected

mutation load, fitness, and frequency of the corresponding genotypes. Given the haplotype

frequencies we observed in spruce bark beetle inversions, two scenarios are likely. First, that

minority arrangements experience higher mutation load (due to reduced recombination and lower

population size) but are maintained in the population at low frequency due to, e.g., associative

overdominance. Such a mechanism would favor balanced inversion polymorphisms of

intermediate to large sizes (harboring hundreds of genes; (Ohta 1971; Connallon and Olito 2022)

and has been shown to play a role in maintaining polymorphic inversions in several insect

species (Yang et al. 2002; Jay et al. 2021). Second, mutation load may accumulate on one or

both inversion arrangements but be mitigated by the haplotype structuring process, i.e., the

existence of multiple diverged sub-haplotypes among inversion homozygotes that reduces the

mutation load within homozygotes. If this process operates within one or both inversion

arrangements it may result in more equal frequencies of alternative inversion haplotypes.

However, such a mechanism is only possible when genetic variation and mutation load is high

(Berdan et al. 2021).

In contrast to these theoretical expectations, we observed no sign of increased mutation load

(measured by the π_N/π_S ratio) in inversion regions compared to the collinear part of the spruce

bark beetle genome (Table S10; Figure 5). We also found no sign of haplotype structuring

capable of reducing mutation load within inversion homozygotes (Figure S12). The only

significant within-homozygote clustering we observed was in a few inversion haplotypes.

However, the clustering divided individuals into southern and northern clades suggesting either

neutral differentiation or divergent selection acting on one of the inversion arrangements, rather

than haplotype structuring being associated with mutation load (Figure S12). These results are

consistent with observations of no significant mutation load in other species where inversion

haplotypes are subject to divergent selection that results in geographic structuring (Harringmeyer

and Hoekstra 2022; Huang et al. 2022). In such cases, inversions facilitate adaptive divergence

but do not tend to accumulate a mutation load. However, geographic clustering of inversion

polymorphisms is weak in the spruce bark beetle, and many inversions appear to have a slightly

lower mutation load associated with the more common inversion haplotype (Figure 5, Table

S11). It is possible that accumulation of a mutation load in the spruce bark beetle is mitigated by

high effective population size in this species and/or gene conversion and double crossover

events, which despite their apparent rarity have been detected in several spruce bark beetle

inversions.

Overall, the absence of heterozygote excess does not support a role of overdominance in the

maintenance of inversion polymorphism in the spruce bark beetle. Likewise, the absence of an

elevated mutation load in inverted regions does not support a role of associative overdominance

either. However, since we only have genomic data available, we cannot conclusively rule out

that other potential mechanisms, such as negative frequency-dependent selection or antagonistic

pleiotropy, could maintain balanced inversion polymorphisms. Additional temporal data are

needed to test whether temporally varying selection has affected the frequencies of inversion

haplotypes in the spruce bark beetle and further research is essential to determine the role of

different mechanisms maintaining inversion polymorphisms within the species.

Far-reaching consequences of having an inversion-rich genome

The presence of multiple polymorphic inversions can have significant consequences for the

evolution of a species, as well as for evolutionary inferences based on genome-wide

polymorphism data. Importantly, polymorphic inversions are a reservoir of genetic variation that

can facilitate adaptation to rapidly changing environments. Indeed, several studies have shown

that polymorphic inversions support rapid adaptation to changing climatic conditions (Rane et al.

2015; Kapun and Flatt 2019; McCulloch and Waters 2023) or adaptive tracking of fluctuating

environments (Nunez et al. 2023). Spruce bark beetle populations are subject to seasonal weather

changes and a rapidly changing environment due to strong anthropogenic pressures (Hofmann et

al. 2024). Warmer weather and drought periods have been associated with a predicted

intensification of bark beetle outbreaks (Biedermann et al. 2019; Bentz et al. 2021; Hlásny et al.

2021), which may act as a strong selection factor within bark beetle populations. Whether

inversions are involved in rapid adaptations in the spruce bark beetle is an open question that

requires further investigation.

Abundant polymorphic inversions within the genome can have far-reaching consequences for

inferences about e.g. demographic history and selection. It is well known that non-equilibrium

demography and selection can leave similar genomic signatures. Traditionally, demographic

analyses have used non-coding parts of the genome, based on the assumption that directional

selection mostly affects protein-coding regions. However, growing evidence for the importance

of background selection in shaping genome-wide diversity is moving the field towards

incorporating linked selection into inferences of demographic history (Li et al. 2012; Johri et al.

2021). We believe that new approaches should also consider the potential influence of

polymorphic inversion landscapes, because variation patterns of inverted regions can be shaped

by different types of balancing selection. In addition, genomics scans for selection in inversion-

rich genomes may be biased due to reduced recombination within inversion regions. Importantly,

the effect of reduced recombination may extend outside inversions (Adrion et al. 2020; Koury

2023; Li et al. 2023). The influence of inversions and inversion-rich genomes on various

evolutionary inference methods may become a standard scenario to test their robustness (Novo et

al. 2023).

Materials and methods

Species genome and its quality

The spruce bark beetle karyotype is n = 14 + XYp with male being a heterogametic sex. The

species genome assembly consists of 272 contigs, and 36 largest (>1Mb) contigs constitute the

majority of the assembly (78% of 236.8 Mb long assembly, N50 = 6.65 Mb; Powell et al. 2021).

Telomeric motifs (sequences present at the end of chromosomes) were identified at the end of

eight contigs (including five largest contigs). BUSCO analysis indicated that 99.5 % genes

present in the insect database (insects_odb9) were also present among the predicted *Ips*

typographus genes. To further investigate the quality of the *Ips typographus* assembly we

aligned it to *Ips nitidius* reference genome (chromosome-level assembly, 16 chromosomes; 2.3

Mya divergence from *Ips typographus*; (Wang et al. 2023). The alignment was conducted with

minimap2 (Li 2018; *asm20* preset recommended for sequences expected to differ in up to 20%).

We used *Ips typographus* contigs as either an alignment target or as a query.

Sampling

Adult spruce bark beetles were collected with pheromone-baited traps in the spring and summer

of 2020. In total, we sampled 18 populations throughout Europe with 13-14 individuals per

locality (253 individuals in total) (Figure 2; Table S5). Throughout the text, we use the term

'population' to refer to a particular site or a collection of closely situated sites (within about 50

km). In Austria, we pooled individuals from three localities that were up to 120 km apart because

of small sample sizes (five beetles or less per site). Populations from Fennoscandia are referred

to as northern populations (or the northern group) and populations from central Europe will be

referred to as southern populations (or the southern group). Polish populations are considered

separately from other central European populations (due to high admixture proportions from

northern group identified in downstream analysis). Beetles were brought alive to the laboratory,

kept on a paper diet for several days, dissected, sexed based on genitalia morphology, and

subjected to DNA extraction (described below).

DNA extraction and genome re-sequencing

DNA was extracted from the whole body of dissected beetles using the Wizard Genomic DNA

Purification Kit (Promega). The concentration of extracted DNA was estimated using a Qubit

fluorimeter (Thermo Fisher Scientific). Genomic libraries were prepared with NEBNext Ultra II

FS DNA Library Prep with Beads (New England Biolabs), with single indexes. Individual

libraries were combined into three pools and 2×150 bp paired-end sequenced in three lanes of a

S4 flowcell using the NovaSeq 6000 instrument and v1 sequencing chemistry (Illumina Inc.).

Sequencing was done by the National Genomics Infrastructure, SNP&SEQ Technology Platform

(Uppsala, Sweden). To assess the overall genotyping error, we prepared and sequenced duplicate

libraries for nine individuals.

Data preparation and filtering

Details of raw data processing and filtering are described in the Supplementary Files. Shortly,

raw reads were mapped to the reference genome (Powell et al. 2021) using Bowtie 2 version

2.4.2 (Langmead and Salzberg 2012). Duplicated reads were removed using Picard

MarkDuplicates version 2.24.1 (Broad Institute 2019). To detect and correct systematic errors in

base quality scores, recalibration was done using the Genome Analysis Toolkit (GATK) version

4.1.9.0, BaseRecalibrator, and ApplyBQSR (McKenna et al. 2010; Depristo et al. 2011). Variant

calling and genotyping was done using GATK HaplotypeCaller, CombineGVCFs, and

GenotypeGVCFs. GATK VariantRecalibrator and ApplyVQSR were used to calculate and filter

(by variant) quality score log-odds (VQSLOD). Bcftools version 1.11 (Danecek et al. 2021) was

used to remove insertions and deletions (indels) as well polymorphisms five bases up- and

downstream. GATK VariantFiltration was applied to mask all genotypes with low sequencing

depth or low genotype quality (McKenna et al. 2010; Depristo et al. 2011). Variants which were

not biallelic single nucleotide polymorphisms or did not meet the recommended hard filtering

thresholds (GATK Team, see Supplementary materials) were filtered out. To filter out

polymorphisms that could come from duplicated regions we removed variants located within

repeat-masked regions of the genome (Powell et al. 2021), variants with excessive overall

coverage (mean + 1 SD), and variants with heterozygote excess. Variants for which genotypes

could be detected in less than half of the individuals were also removed. We used PLINK version

1.90b6.18 (Purcell et al. 2007) to detect sample contaminations, swaps and duplications, and

familial relationships (e.g. sibling pairs present in the data) which might bias downstream

analyses. Individuals with excessive coverage were removed, as these could be a result of human

errors during library preparation or pooling. We focused on contigs longer than 1 Mb that

together constituted 78% of the genome assembly, i.e. a total of 186 Mb. Since part of

IpsContig33 had high similarity to mtDNA this contig was not included in the downstream

analysis. Genotyping error was assessed using GATK Genotype Concordance.

Genome-wide genetic variation and its geographic structuring

Genome-wide genetic structuring was explored by PCA using PLINK. The most likely number

of genetic clusters and admixture proportions was estimated using NGSadmix (Skotte et al.

2013). The analysis was run for five different K-values (1-5; 10 replicates per K-value), using a

minor allele frequency (MAF) filter of 0.05 and 10,000 iterations, and a SNP dataset that was

pruned for linkage disequilibrium using PLINK (–indep-pairwise 50 10 0.1 option). To choose

the most likely number of genetic clusters, the results were examined using CLUMPAK

(http://clumpak.tau.ac.il/index.html). To examine the influence of inversions on genetic

clustering and to facilitate inversion genotyping, NGSadmix was run separately for 1) all

autosomal contigs without potential inversions and 2) each potential inversion.

To assess genetic differentiation among different spruce bark beetle populations Weir and

Cockerham's (Weir and Cockerham 1984) F_{ST} was estimated using VCFtools version 0.1.16

(Danecek et al. 2011). F_{ST} was calculated between population groups identified by NGSadmix,

as well as among all population pairs. Additionally, we summarized F_{ST} values in 100 kb

overlapping windows (using 20 kb steps) along the contigs. Window-based analyses were done

for each contig and each population pair. In addition, absolute sequence divergence (d_{xy}),

nucleotide diversity, and Tajima's D statistic were estimated and summarized for 100 kb non-

overlapping windows using ANGSD version 0.935-44-g02a07fc (Korneliussen et al. 2014).

These statistics were calculated for each population separately and were based on a maximum

likelihood estimate of the folded site frequency spectrum (SFS). We excluded sites with mapping

quality below 30 phred, base quality score below 20, and coverage less than three times the

population sample size and more than three times the average coverage, following the approach

used in Delmore et al. (Delmore et al. 2018). The ANGSD calculations were based on allele

frequencies estimated from genotype likelihoods (Li 2011) and ngsPopGen scripts

(https://github.com/mfumagalli/ngsPopGen).

Identification of inversions, their geographic distribution, and variation patterns

To identify inversions, we followed standard population genetic approach (see e.g. Huang et al.

2020; Mérot et al. 2021; Reeve et al. 2023). Potential chromosomal inversion regions were

identified based on: i) per contig PCAs, ii) local PCAs, iii) patterns of heterozygosity, and iv) LD

clustering. Per contig PCAs were performed as a first screen for inversion existence using

PLINK (Purcell et al. 2007). Local PCAs were performed to identify inversions approximate

positions using the *lostruct* R package (Li and Ralph 2019). Analysis was performed using two

independent approaches: localPCA analysis using all analyzed contigs and localPCA analysis for

each contig independently. The first analysis was run following the approach described in

Huang et al. (2020)Windows were considered outliers if their MDS scores were greater than 2SD

from the mean across all windows and the maximum number of windows between what was

considered separate inversions was set to 20. In the second analysis separate contigs were subject

to k-means clustering algorithm (R) to define groups of windows that formed a single inversion.

K-means clustering algorithm was performed on the first MDS scores. For each contig different

k (number of window clusters) were tested so that each structurally different part of the contig

was isolated into one cluster. For both approaches, different window sizes (1, 10 and 100kb)

were tested.

Linkage disequilibrium among SNPs (thinned by selecting one SNP every 10 kb; MAF > 5%)

was calculated for each contig using PLINK. We considered a genomic region to be an inversion

region if 1) local PCA analysis identified the region as an outlier, and/or 2) the region exhibited

high LD (most SNPs having $r^2 > 0.4$), and 3) PCA performed on SNPs from this region separated

individuals into three distinct groups with heterozygosity patterns matching the expectation of

the middle group of the PCA presenting higher heterozygosity.

Genotyping of individual beetles with respect to the inversion haplotypes they carried was done

based on NGSadmix clustering (with K = 2 inversion heterozygotes having mixed ancestry in

approximate 50/50 proportions; Figure S13). In a few more complex cases (putative overlapping

inversions and double-crossover events) genotyping was done by clustering individuals based on

the PCA groups visible along PC1 and PC2 (for more details see Figure S3 and related text).

Contigs with less than 10,000 SNPs provided ambiguous results and were excluded from

genotyping. Approximate inversion boundaries were defined based on local PCAs and sharp

borders detected in LD clusters. The population genetic approach used in this study is a standard

approach to detect putative large polymorphic inversions but is not able to provide details on

inversion breakpoints.

To further support inversion evidence, we estimated population recombination rates (rho) for

each putative inversion region using individuals with a particular inversion genotype (separately

for each homozygous and heterozygous individuals). We followed the approach used in Jones et

al. (2019) and sampled 20 individuals from each genotype group in cases when more individuals

were available. Rho was not calculated if there were less than 10 individuals of a particular

genotype. Watterson theta estimates were used to create a custom likelihood lookup table using

the Ldhat 2.2 program *complete* (McVean et al. 2002). The *interval* program was used to

estimate the population recombination rate across investigated inversion (and adjacent regions in

the same contig). The *interval* algorithm was run for 2 million iterations and the chain was

sampled every 10,000 iterations with a burn-in of 100,000 generations (*stat* program in Ldhat

package, block penalty = 5). Population recombination rates were summarized in non-

overlapping 100 kb windows using a custom perl script.

Inversion genotype and haplotype frequencies were calculated using an in-house R-script.

Frequencies were calculated 1) within each population, 2) within the southern and northern

group, and 3) for all sampled populations combined. Deviations from Hardy-Weinberg

equilibrium were estimated for all three datasets. Inversions with only two haplotypes were

tested using Fisher's exact tests. Inversions with more than two haplotypes (including

recombinant haplotypes between two inversion arrangements) were tested using permutation

test, and sex chromosome inversions were tested as described in Graffelman & Weir (Graffelman

and Weir 2018). All tests were done using the R package *HardyWeinberg* (Graffelman 2015). To

investigate if inversion haplotypes differed in frequency along environmental gradients, Pearson

correlation between inversion haplotype frequencies and latitude/longitude was calculated.

To assess levels of genetic differentiation between inversion haplotypes, F_{ST} and d_{xy} between

alternative inversion haplotypes (AA, BB) were estimated following the approach described

above. Deletions present in alternative inversion haplotypes were not included in d_{xy} calculations

but were summarized separately using custom scripts.

Testing for enrichment of gene ontology terms in inversions

We used genes annotations from Powell et. al (2021) and tested for overrepresentation of gene

ontology (GO) terms among genes located within inversions using R package topGO (Alexa and

Rahnenführer 2022), applying Fisher's exact test and "weight01" algorithm (Alexa et al. 2006)

to deal with the GO graph structure; only GO categories with at least 10 members among the

SNP associated genes were considered.

Inversion age estimation

Absolute sequence divergence between alternative inversion haplotypes was used to calculate the

approximate time of divergence of inverted and non-inverted haplotypes. We used the equation T

$= d_{xy}/2\mu$, where T is the divergence time in generations, μ is the mutation rate per site per

generation, and d_{xy} is a mean d_{xy} calculated based on per SNP values estimated in ANGSD. Since

mutation rates of the spruce bark beetle are unknown, we used a range of mutation rate estimates

available for some diploid, sexually reproducing insects including *Drosophila melanogaster*

(Krasovec 2021)*, Heliconius melpomene* (Keightley et al. 2015), and *Chironomus riparius*

(Oppold and Pfenninger 2017). Per generation per bp mutation rate estimates varied from 2.1 to

11.7×10^{-9}. This approach could only give us rough inversion age estimates due to the

uncertainty of the mutation rate estimates, probable intraspecific variation in mutation rate

(Krasovec 2021), and a (likely) substantial influence of selection and gene flux (Charlesworth

2023).

Mutation load estimation

To estimate mutation load we calculated the ratio of nucleotide diversity at non-synonymous

sites (π_N) vs. synonymous sites (π_S). Mutation load (π_N/π_S) was calculated separately for each

inversion homozygote and for the collinear part of the genome (all individuals). We computed

nucleotide diversity for each site using SNPGenie (Nelson et al. 2015). The π_N/π_S ratio was

estimated in 200 kb windows using an in-house R script. To account for the fact that inversions

can greatly suppress recombination in surrounding parts of the genome (Koury 2023) the

collinear part of the genome was divided into two groups: 1) a group including all collinear 200

kb windows outside inversions and 2) a group including all collinear windows outside inversions

but excluding windows that came from contigs with inversions (so called strict filtering). Both

collinear datasets were used to test for overall differences in mutation load between inversions

and the collinear part of the genome (using two-sided t-test). One-sided t-tests were used to test

whether minor (less frequent) homokaryotypes had higher mutation loads than major (more

frequent) homokaryotypes. Homokaryotypes that contained fewer than four 200 kb windows and

were present in few individuals (two thresholds were tested: < 4 and < 10 individuals) were

excluded from the analysis. Additionally, windows with a small number of genes were excluded

(two thresholds were tested: < 5 and < 10 genes).

Haplotype structuring, i.e., the presence of two or more distinct sub-haplotypes among inversion

homozygote haplotypes, can prevent fitness degeneration on one or both inversion haplotypes by

carrying partially complementary sets of deleterious recessive alleles (Charlesworth and

Charlesworth 1997; Berdan et al. 2021). To check if any inversion homozygotes exhibited

haplotype structuring we first phased the data using Beagle 5.2 (default settings (Browning and

Browning 2007)). Next, in each inverted region and homokaryotype we filtered out all variants

with MAF < 0.1 and used PGDSpider 2.1.1.0 (Lischer and Excoffier 2012) to convert variant

call formats (VCF) to full length sequences. Finally, we constructed neighbor-joining trees for

alleles within haplotypes using MEGA7 and investigated if topologies showed the presence of

distinct clusters within homokaryotype groups (Kumar et al. 2016).

Phenotype-genotype associations

To test whether inversion polymorphisms were associated with diapause phenotypes we

resequenced whole genomes of 18 individuals from a spruce bark beetle diapause study by

Schebeck et al. (Schebeck et al. 2022) (10 beetles expressing facultative diapause and 8 beetles

expressing obligatory diapause). DNA sequencing was performed using the DNBseq platform

(BGI Tech Solutions, Poland) to a mean coverage of 20×. The data was processed in the same

way as described above and combined with other sequenced individuals before performing PCA

in PLINK. To test for differentiation in inversion haplotype frequencies between diapause

phenotypes an exact G test was run using Genepop 4.1.2 (Raymond and Rousset 1995). F_{ST}

between individuals expressing facultative and obligate diapause was estimated using VCFtools

version 0.1.16 (Danecek et al. 2011) and summarized in 100 kb overlapping windows (20 kb

steps).

To check if spruce bark beetle odorant receptors (ORs) genes were associated with inversions we

examined 73 OR genes recently annotated by Yuvaraj et al. (Yuvaraj et al. 2021). OR sequences

were mapped to the bark beetle reference genome using minimap2 (Li 2018), and 71 out of 73

ORs were located in the 186 Mb covered by the 36 contigs we analyzed. Three ORs mapped to

more than one contig. ItypOR9 and ItypOR58 mapped to the end of IpsContig16 and 23,

suggesting possible assembly error and duplication of end-of-contig sequences, which are

difficult to assemble. ItypOR59NTE mapped to three nearby locations on IpsContig6, suggesting

either assembly error or recent duplication. For these three ORs we used one randomly chosen

location in downstream analyses. To test if inversions were enriched in OR genes, we ran

permutation tests (10,000 iterations; permutating inversion locations). To check if alternative

inversion arrangements harbored different numbers of OR genes (e.g. that one arrangement

carried a deletion) we compared sequence coverage within ORs in individual beetles identified

as inversion homozygotes. Additionally, we counted nonsynonymous segregating variants

present in ORs situated within inversions and checked if any of these SNPs are highly diverged

between inversion haplotypes (by calculating d_{xy}). Finally, to investigate whether natural

selection was involved in shaping ORs nonsynonymous variation patterns between inversion

haplotypes we calculated d_N/d_S ratio for ORs with multiple highly diverged nonsynonymous

variants and performed MacDonald-Kreitman test for ORs with $d_N/d_S > 1$. D_{xy} was calculated

using GATK VariantsToTable outputs and custom perl scripts and dN/dS was calculated using

pairwise distances option in MEGA11 (Syn-Nonsynonymous substitution model, Kumar

method). MacDonald-Kreitman test was performed in DnaSP v6 (Rozas et al. 2017).

Genotype-environment association

Genotype-environment association (GEA) analyses were done to test if allele frequency changes

in SNPs were associated with the beetle populations' local environment. The analysis was

performed using 1) all SNPs present in the analyzed part of the genome and 2) excluding SNPs

present within inversion regions and coding inversions as single SNPs. Many different

environmental variables were summarized along two principal components (see below). To

control for the confounding effect of the overall genetic differentiation, we used Latent Factors

Mixed Models (LFMM (Caye et al. 2019)) as implemented in the lfmm2() function from the R

package LEA (Gain and François 2021). We used only SNPs with < 20% missing data and MAF

>= 0.1, and that occurred in individuals with < 30% missing data. Because LFMM cannot handle

missing data, we imputed missing genotypes with impute() from LEA. We ran LFMM for 1) all

SNPs that passed the filters described above and 2) for a dataset where each inversion was

represented as a single "SNP" inversion genotype. We used five latent factors (K = 5) in

lfmm2(). P-values were calculated using lfmm2.test() from LEA and false discovery rate (FDR)-

corrected using the p.adjust() R function with method = "fdr".

Each beetle population's local environment was characterized according to climate and land-

cover data. We used all 19 bioclimatic variables from WorldClim version 2.1. (Fick and Hijmans

2017) with a resolution of ~1 km^2. Environmental variables included annual mean temperature,

mean diurnal range, isothermality, temperature seasonality, maximum temperature of warmest

month, minimum temperature of coldest month, temperature annual range, mean temperature of

wettest quarter, mean temperature of driest quarter, mean temperature of warmest quarter, mean

temperature of coldest quarter, annual precipitation, precipitation of wettest month, precipitation

of driest month, precipitation seasonality, precipitation of wettest quarter, precipitation of driest

quarter, precipitation of warmest quarter, and precipitation of coldest quarter. These variables

are averaged over the years 1970-2000. Proportions of forest, cropland, and built-up areas were

downloaded from https://lcviewer.vito.be/ for 2015 with a spatial resolution of ~100 m^2

(Buchhorn et al. 2019). These global land-cover maps are part of the Copernicus Land Service,

derived from PROBA-V satellite observations, and have an accuracy of 80% as measured by the

CEOS land product validation subgroup. We also included the proportion of land area covered by

spruce trees (genus *Picea*) at a resolution of ~1 km^2, as obtained by Brus et al. (2012) using a

statistical mapping approach. All environmental variables were reprojected to a final resolution

of ~1 km^2 using the Lambert azimuthal equal area method. We then extracted mean values for all

environmental variables (Figure S14) within a 50 km radius from each population location using

the R package terra (Hijmans 2023). Finally, PCA was used to summarize the multi-scale

environmental variation among populations. The first two PCA components (PC1 and PC2)

explained 25% and 23% of the environmental variation, respectively, and were used as the final

input for the GEA analyses. PC1 represented environmental variation mainly related to latitude,

with northern populations showing higher values indicative of higher temperature variation and

lower temperatures during the coldest months. PC2 represented environmental variation mostly

related to temperature and amount of cropland, with higher values representing localities with

higher temperatures during the warmest months and a higher proportion of cropland (and

conversely less forest cover and spruce) (Figure S15). Additionally, we identified genes closest

to SNPs that showed significant association with PC.

Data Availability

All DNA sequences have been deposited to the National Center for Biotechnology Information

Sequence Read Archive under the BioProject ID PRJNA1013983.

Acknowledgments

We thank A. Hietala, E. Stengel, K. Zub, Å. Lindelöw, O. Langvall, M. Holmlund, E.

Kristensen, U. Johansson, R. Modlinger, J. Reisenberger, K. Szreder, W. Skowroński L.

Stanecki, and M. Ahlström for help in sampling spruce bark beetles across Europe. T. Mokrzycki

helped with beetle sexing and identification. We thank members of the Genomics and

Experimental Evolution Group at Jagiellonian University for their help in improving this

manuscript. We thank Tomasz Gaczorek for the help in writing scripts and optimizing data

analysis. This work was funded by a Polish National Science Center 2018/30/E/NZ8/00105 grant

to K.N.B, and the Foundation in Memory of Oscar and Lili Lamm to M.N.A.

Author contributions

A.M., P.Z. and K.N.B. conceived the study, performed the main analyses, and wrote the manuscript. A.B. managed sample shipment, sexed beetles, isolated DNA, and prepared samples for sequencing. F.S., P.K. and M.N.A. provided an unpublished version of the spruce bark beetle genome and valuable insights on the species' ecology and sensory biology. M.M., J.M., Z.B., M.S., P.K., and C.S. organized sampling, provided beetles for analysis and insights on sampled populations. B.A. and W.B. performed GEA analysis. Z.N. analyzed diapause data. J.M. phased the data. M.S. provided diapause samples. J.Z performed *Ips nitidus* analysis. W.B. helped in data interpretation and provided feedback on all manuscript versions. All authors read and approved the final manuscript.

References

Adrion JR, Galloway JG, Kern AD. 2020. Predicting the landscape of recombination using deep learning. *Mol. Biol. Evol.* 37:1790–1808.

Akopyan M, Tigano A, Jacobs A, Wilder AP, Baumann H, Therkildsen NO. 2022. Comparative linkage mapping uncovers recombination suppression across massive chromosomal inversions associated with local adaptation in Atlantic silversides. *Mol. Ecol.* 31:3323–3341.

Alexa A, Rahnenführer J. 2022. topGO: Enrichment analysis for gene ontology (2.46. 0)[computer software]. Bioconductor version: Release (3.14)e.

Alexa A, Rahnenführer J, Lengauer T. 2006. Improved scoring of functional groups from gene expression data by decorrelating GO graph structure. *Bioinformatics* 22:1600–1607.

Ayala D, Acevedo P, Pombi M, Dia I, Boccolini D, Costantini C, Fontenille D. 2017. Chromosome inversions and ecological plasticity in the main African malaria mosquitoes. *Evolution* 71:686–701.

Ayala D, Ullastres A, González J, Chain FJJ, Planck M. 2014. Adaptation through chromosomal inversions in Anopheles. *Front. Genet.* 5:1–10.

Bentz BJ, Hansen EM, Davenport M, Soderberg D. 2021. Complexities in predicting mountain pine beetle and spruce beetle response to climate change. In: Bark beetle management, ecology and climate change. Elsevier.

Berdan EL, Barton NH, Butlin R, Charlesworth B, Faria R, Fragata I, Gilbert KJ, Jay P, Kapun

M, Lotterhos KE, et al. 2023. How chromosomal inversions reorient the evolutionary

process. *J. Evol. Biol.* 36:1761–1782.

Berdan EL, Blanckaert A, Butlin RK, Bank C. 2021. Deleterious mutation accumulation and the

long-term fate of chromosomal inversions. *PLoS Genet.* 17: e1009411.

Bertheau C, Schuler H, Arthofer W, Avtzis DN, Mayer F, Krumböck S, Moodley Y, Stauffer C.

2013. Divergent evolutionary histories of two sympatric spruce bark beetle species. *Mol.*

Ecol. 22:3318–3332.

Biedermann PHW, Müller J, Grégoire JC, Gruppe A, Hagge J, Hammerbacher A, Hofstetter

RW, Kandasamy D, Kolarik M, Kostovcik M, et al. 2019. Bark Beetle Population

Dynamics in the Anthropocene: Challenges and Solutions. *Trends Ecol. Evol.* 34:914–924.

Brelsford A, Purcell J, Avril A, Tran Van P, Zhang J, Brütsch T, Sundström L, Helanterä H,

Chapuisat M. 2020. An Ancient and Eroded Social Supergene Is Widespread across

Formica Ants. *Curr. Biol.* 30:304-311.e4.

Browning SR, Browning BL. 2007. Rapid and accurate haplotype phasing and missing-data

inference for whole-genome association studies by use of localized haplotype clustering.

Am. J. Hum. Genet. 81:1084–1097.

Brus DJ, Hengeveld GM, Walvoort DJJ, Goedhart PW, Heidema AH, Nabuurs GJ, Gunia K.

2012. Statistical mapping of tree species over Europe. *Eur. J. For. Res.* 131:145–157.

Buchhorn M, Smets B, Bertels L, Lesiv M, Tsendbazar, N. E., Herold M, Fritz S. 2019.

Copernicus global land service: Land cover 100m: Epoch 2015: Globe. Version V2. 0.2.

Caye K, Jumentier B, Lepeule J, François O. 2019. LFMM 2: Fast and accurate inference of

gene-environment associations in genome-wide studies. *Mol. Biol. Evol.* 36:852–860.

Charlesworth B. 2023. The effects of inversion polymorphisms on patterns of neutral genetic

diversity. *Genetics* 224:1–21.

Charlesworth B, Charlesworth D. 1997. Rapid fixation of deleterious alleles can be caused by

Muller's ratchet. *Genet. Res.* 70:63–73.

Cheng C, Tan JC, Hahn MW, Besansky NJ. 2018. Systems genetic analysis of inversion

polymorphisms in the malaria mosquito Anopheles gambiae. *Proc. Natl. Acad. Sci. U. S. A.*

115:E7005–E7014.

Christmas MJ, Wallberg A, Bunikis I, Olsson A, Wallerman O, Webster MT. 2019.

Chromosomal inversions associated with environmental adaptation in honeybees. *Mol.*

Ecol. 28:1358–1374.

Connallon T, Clark AG. 2014. Balancing selection in species with separate sexes: Insights from

fisher's geometric model. *Genetics* 197:991–1006.

Connallon T, Olito C. 2022. Natural selection and the distribution of chromosomal inversion

lengths. *Mol. Ecol.* 31:3627–3641.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G,

Marth GT, Sherry ST, et al. 2011. The variant call format and VCFtools. *Bioinformatics*

27:2156–2158.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T,

McCarthy SA, Davies RM. 2021. Twelve years of SAMtools and BCFtools. *Gigascience*

10:1–4.

Delmore KE, Lugo Ramos JS, Van Doren BM, Lundberg M, Bensch S, Irwin DE, Liedvogel M.

2018. Comparative analysis examining patterns of genomic differentiation across multiple

episodes of population divergence in birds. *Evol. Lett.* 2:76–87.

Denlinger D. 2022. Insect diapause. Cambridge University Press.

Depristo MA, Banks E, Poplin R, Garimella K V., Maguire JR, Hartl C, Philippakis AA, Del

Angel G, Rivas MA, Hanna M, et al. 2011. A framework for variation discovery and

genotyping using next-generation DNA sequencing data. *Nat. Genet.* 43:491–501.

Ellerstrand SJ, Choudhury S, Svensson K, Andersson MN, Kirkeby C, Powell D, Schlyter F,

Jönsson AM, Brydegaard M, Hansson B, et al. 2022. Weak population genetic structure in

Eurasian spruce bark beetle over large regional scales in Sweden. *Ecol. Evol.* 12:1–12.

Faria R, Chaube P, Morales HE, Larsson T, Lemmon AR, Lemmon EM, Rafajlović M, Panova

M, Ravinet M, Johannesson K, et al. 2019. Multiple chromosomal rearrangements in a

hybrid zone between Littorina saxatilis ecotypes. *Mol. Ecol.* 28:1375–1393.

Faria R, Johannesson K, Butlin RK, Westram AM. 2019. Evolving Inversions. *Trends Ecol.*

Evol. 34:239–248.

Fick SE, Hijmans RJ. 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for

global land areas. *Int. J. Climatol.* 37:4302–4315.

Fuller ZL, Haynes GD, Richards S, Schaeffer SW. 2016. Genomics of natural populations: How

differentially expressed genes shape the evolution of chromosomal inversions in Drosophila

pseudoobscura. *Genetics* 204:287–301.

Fuller ZL, Haynes GD, Richards S, Schaeffer SW. 2017. Genomics of natural populations:
Evolutionary forces that establish and maintain gene arrangements in Drosophila
pseudoobscura. *Mol. Ecol.* 26:6539–6562.

Fuller ZL, Koury SA, Phadnis N, Schaeffer SW. 2019. How chromosomal rearrangements shape
adaptation and speciation: Case studies in Drosophila pseudoobscura and its sibling species
Drosophila persimilis. *Mol. Ecol.* 28:1283–1301.

Gain C, François O. 2021. LEA 3: Factor models in population genetics and ecological genomics
with R. *Mol. Ecol. Resour.* 21:2738–2748.

Graffelman J. 2015. Exploring diallelic genetic markers: The HardyWeinberg package. *J. Stat.*
Softw. 64:1–23.

Graffelman J, Weir BS. 2018. Multi-allelic exact tests for Hardy–Weinberg equilibrium that
account for gender. *Mol. Ecol. Resour.* 18:461–473.

Guerrero RF, Rousset F, Kirkpatrick M. 2012. Coalescent patterns for chromosomal inversions
in divergent populations. *Philos. Trans. R. Soc. B Biol. Sci.* 367:430–438.

Gutiérrez-Valencia J, Hughes PW, Berdan EL, Slotte T. 2021. The Genomic Architecture and
Evolutionary Fates of Supergenes. *Genome Biol. Evol.* 13:1–19.

Hansson BS, Stensmyr MC. 2011. Evolution of insect olfaction. *Neuron* 72:698–711.

Harringmeyer OS, Hoekstra HE. 2022. Chromosomal inversion polymorphisms shape the
genomic landscape of deer mice. *Nat. Ecol. Evol.* 6:1965–1979.

Hijmans R. 2023. terra: Spatial Data Analysis. R package version 1.7-18. :https://CRAN.R-
project.org/package=terra.

Hlásny T, König L, Krokene P, Lindner M, Montagné-Huck C, Müller J, Qin H, Raffa KF,
Schelhaas MJ, Svoboda M, et al. 2021. Bark Beetle Outbreaks in Europe: State of
Knowledge and Ways Forward for Management. *Curr. For. Reports* 7:138–165.

Hof AEV t., Campagne P, Rigden DJ, Yung CJ, Lingley J, Quail MA, Hall N, Darby AC,
Saccheri IJ. 2016. The industrial melanism mutation in British peppered moths is a
transposable element. *Nature* 534:102–105.

Hofmann S, Schebeck M, Kautz M. 2024. Diurnal temperature fluctuations improve predictions
of developmental rates in the spruce bark beetle Ips typographus. *J. Pest Sci.* 2024.

Hou XQ, Yuvaraj JK, Roberts RE, Zhang DD, Unelius CR, Löfstedt C, Andersson MN. 2021.
Functional Evolution of a Bark Beetle Odorant Receptor Clade Detecting Monoterpenoids

of Different Ecological Origins. *Mol. Biol. Evol.* 38:4934–4947.

Huang K, Andrew RL, Owens GL, Ostevik KL, Rieseberg LH. 2020. Multiple chromosomal

inversions contribute to adaptive divergence of a dune sunflower ecotype. *Mol. Ecol.*

29:2535–2549.

Huang K, Ostevik KL, Elphinstone C, Todesco M, Bercovich N, Owens GL, Rieseberg LH.

2022. Mutation Load in Sunflower Inversions Is Negatively Correlated with Inversion

Heterozygosity. *Mol. Biol. Evol.* 39:1–17.

Jay P, Chouteau M, Whibley A, Bastide H, Parrinello H, Llaurens V, Joron M. 2021. Mutation

load at a mimicry supergene sheds new light on the evolution of inversion polymorphisms.

Nat. Genet. 53:288–293.

Jay P, Tezenas E, Véber A, Giraud T. 2022. Sheltering of deleterious mutations explains the

stepwise extension of recombination suppression on sex chromosomes and other

supergenes.

Johri P, Riall K, Becher H, Excoffier L, Charlesworth B, Jensen JD. 2021. The Impact of

Purifying and Background Selection on the Inference of Population History: Problems and

Prospects. *Mol. Biol. Evol.* 38:2986–3003.

Jones JC, Wallberg A, Christmas MJ, Kapheim KM, Webster MT, Singh N. 2019. Extreme

Differences in Recombination Rate between the Genomes of a Solitary and a Social Bee.

Mol. Biol. Evol. 36:2277–2291.

Joron M, Frezal L, Jones RT, Chamberlain NL, Lee SF, Haag CR, Whibley A, Becuwe M,

Baxter SW, Ferguson L, et al. 2011. Chromosomal rearrangements maintain a polymorphic

supergene controlling butterfly mimicry. *Nature* 477:203–206.

Kandasamy D, Gershenzon J, Andersson MN, Hammerbacher A. 2019. Volatile organic

compounds influence the interaction of the Eurasian spruce bark beetle (*Ips typographus*)

with its fungal symbionts. *ISME J.* 13:1788–1800.

Kandasamy D, Zaman R, Nakamura Y, Zhao T, Hartmann H, Andersson MN, Hammerbacher A,

Gershenzon J. 2023. Conifer-killing bark beetles locate fungal symbionts by detecting

volatile fungal metabolites of host tree resin monoterpenes.

Kapun M, Fabian DK, Goudet J, Flatt T. 2016. Genomic Evidence for Adaptive Inversion Clines

in Drosophila melanogaster. *Mol. Biol. Evol.* 33:1317–1336.

Kapun M, Flatt T. 2019. The adaptive significance of chromosomal inversion polymorphisms in

Drosophila melanogaster. *Mol. Ecol.* 28:1263–1282.

Keightley PD, Otto SP. 2006. Interference among deleterious mutations favours sex and

recombination in finite populations. *Nature* 443:89–92.

Keightley PD, Pinharanda A, Ness RW, Simpson F, Dasmahapatra KK, Mallet J, Davey JW,

Jiggins CD. 2015. Estimation of the Spontaneous Mutation Rate in Heliconius melpomene.

Mol. Biol. Evol. 32:239–243.

Kim KW, De-Kayne R, Gordon IJ, Omufwoko KS, Martins DJ, Ffrench-Constant R, Martin SH.

2022. Stepwise evolution of a butterfly supergene via duplication and inversion. *Philos.*

Trans. R. Soc. B Biol. Sci. 377.

Kirubakaran TG, Grove H, Kent MP, Sandve SR, Baranski M, Nome T, De Rosa MC, Righino

B, Johansen T, Otterå H, et al. 2016. Two adjacent inversions maintain genomic

differentiation between migratory and stationary ecotypes of Atlantic cod. *Mol. Ecol.*

25:2130–2143.

Koch EL, Morales HE, Larsson J, Westram AM, Faria R, Lemmon AR, Lemmon EM,

Johannesson K. 2021. Genetic variation for adaptive traits is associated with polymorphic

inversions in *Littorina saxatilis*. *Evol. Letters* 5:196–213.

Korneliussen TS, Albrechtsen A, Nielsen R. 2014. ANGSD: Analysis of Next Generation

Sequencing Data. *BMC Bioinformatics* 15:1–13.

Koury SA. 2023. Predicting recombination suppression outside chromosomal inversions in

Drosophila melanogaster using crossover interference theory. *Heridity* 130:196–208.

Kozak GM, Wadsworth CB, Kahne SC, Bogdanowicz SM, Harrison RG, Coates BS, Dopman

EB. 2019. Genomic Basis of Circannual Rhythm in the European Corn Borer Moth. *Curr.*

Biol. 29:3501-3509.e5.

Krasovec M. 2021. The spontaneous mutation rate of Drosophila pseudoobscura. *G3 Genes,*

Genomes, Genet. 11: jkab151.

Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics Analysis

Version 7.0 for Bigger Datasets. *Mol. Biol. Evol.* 33:1870–1874.

Lamichhaney S, Fan G, Widemo F, Gunnarsson U, Thalmann DS, Hoeppner MP, Kerje S,

Gustafson U, Shi C, Zhang H, et al. 2015. Structural genomic changes underlie alternative

reproductive strategies in the ruff (*Philomachus pugnax*). *Nat. Genet.* 48:84–88.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat. Methods*

9:357–359.

Li H. 2011. A statistical framework for SNP calling, mutation discovery, association mapping

and population genetical parameter estimation from sequencing data. *Bioinformatics*

27:2987–2993.

Li H. 2018. Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics* 34:3094–

3100.

Li H, Berent E, Hadjipanteli S, Galey M, Muhammad-Lahbabi N, Miller DE, Crown KN. 2023.

Heterozygous inversion breakpoints suppress meiotic crossovers by altering recombination

repair outcomes. *PLoS Genet.* 19:1–20.

Li H, Ralph P. 2019. Local PCA shows how the effect of population structure differs along the

genome. *Genetics* 211:289–304.

Li J, Li H, Jakobsson M, Li S, Sjödin P, Lascoux M. 2012. Joint analysis of demography and

selection in population genetics: Where do we stand and where could we go? *Mol. Ecol.*

21:28–44.

Lieutier F, Yart A, Salle A. 2009. Stimulation of tree defenses by Ophiostomatoid fungi can

explain attack success of bark beetles on conifers. *Ann. For. Sci.* 66:801.

Lischer HEL, Excoffier L. 2012. PGDSpider: An automated data conversion tool for connecting

population genetics and genomics programs. *Bioinformatics* 28:298–299.

Lohse K, Clarke M, Ritchie MG, Etges WJ. 2015. Genome-wide tests for introgression between

cactophilic Drosophila implicate a role of inversions during speciation. *Evolution* 69:1178–

1190.

Lowry DB, Willis JH. 2010. A widespread chromosomal inversion polymorphism contributes to

a major life-history transition, local adaptation, and reproductive isolation. *PLoS Biol.*

8:e1000500.

Matschiner M, Maria J, Barth I, Tørresen OK, Star B, Baalsrud HT, Servane M, Brieuc O,

Pampoulie C, Bradbury I, et al. 2022. Supergene origin and maintenance in Atlantic cod.

Nat. Ecol. Evol. 6:469–481.

Mayer F, Piel FB, Cassel-Lundhagen A, Kirichenko N, Grumiau L, Økland B, Bertheau C,

Grégoire JC, Mardulyn P. 2015. Comparative multilocus phylogeography of two Palaearctic

spruce bark beetles: Influence of contrasting ecological strategies on genetic variation. *Mol.*

Ecol. 24:1292–1310.

McClung CE. 1905. The chromosome complex of orthopteran spermatocytes. *Biol. Bull.* 9:304–340.

McCulloch GA, Waters JM. 2023. Rapid adaptation in a fast-changing world: Emerging insights from insect genomics. *Glob. Chang. Biol.* 29:943–954.

McKenna A, Hanna M, Banks E, Andrey Sivachenko KC, Kernytsky A, Garimella K, Altshuler D, Gabriel S, Daly M, DePristo MA. 2010. The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* 20:1297–1303.

McVean G, Awadalla P, Fearnhead P. 2002. A coalescent-based method for detecting and estimating recombination from gene sequences. *Genetics* 160:1231–1241.

Mérot C, Berdan EL, Cayuela H, Djambazian H, Ferchaud AL, Laporte M, Normandeau E, Ragoussis J, Wellenreuther M, Bernatchez L. 2021. Locally Adaptive Inversions Modulate Genetic Variation at Different Geographic Scales in a Seaweed Fly. *Mol. Biol. Evol.* 38:3953–3971.

Mérot C, Llaurens V, Normandeau E, Bernatchez L, Wellenreuther M. 2020. Balancing selection via life-history trade-offs maintains an inversion polymorphism in a seaweed fly. *Nat. Commun.* 11:1-11.

Navarro A, Barbadilla A, Ruiz A. 2000. Effect of inversion polymorphism on the neutral nucleotide variability of linked chromosomal regions in drosophila. *Genetics* 155:685–698.

Nelson CW, Moncla LH, Hughes AL. 2015. SNPGenie: Estimating evolutionary parameters to detect natural selection using pooled next-generation sequencing data. *Bioinformatics* 31:3709–3711.

Nilssen AC. 1984. Long-range aerial dispersal of bark beetles and bark weevils (Coleoptera, Scolytidae and Curculionidae) in northern Finland. *Ann. Entomol. Fenn.* 50:37–42.

Novo I, Ordás P, Moraga N, Santiago E, Quesada H, Caballero A. 2023. Impact of population structure in the estimation of recent historical effective population size by the software GONE. *Genetics Selection Evolution* 55:1–15.

Nunez JCB, Lenhart BA, Bangerter A, Murray CS, Mazzeo GR, Yu Y, Nystrom TL, Tern C, Erickson PA, Bergland AO. 2023. A cosmopolitan inversion facilitates seasonal adaptation in overwintering Drosophila. *Genetics* 226: iyad207.

Ohta T. 1971. Associative overdominance caused by linked detrimental mutations. *Genet. Res.*

18:277–286.

Oppold AM, Pfenninger M. 2017. Direct estimation of the spontaneous mutation rate by short-
term mutation accumulation lines in Chironomus riparius. *Evol. Lett.* 1:86–92.

Paolucci S, Salis L, Vermeulen CJ, Beukeboom LW, van de Zande L. 2016. QTL analysis of the
photoperiodic response and clinal distribution of period alleles in Nasonia vitripennis. *Mol.*
Ecol. 25:4805–4817.

Porubsky D, Höps W, Ashraf H, Hsieh PH, Rodriguez-Martin B, Yilmaz F, Ebler J, Hallast P,
Maria Maggiolini FA, Harvey WT, et al. 2022. Recurrent inversion polymorphisms in
humans associate with genetic instability and genomic disorders. *Cell* 185:1986-2005.e26.

Powell D, Groβe-Wilde E, Krokene P, Roy A, Chakraborty A, Löfstedt C, Vogel H, Andersson
MN, Schlyter F. 2021. A highly-contiguous genome assembly of the Eurasian spruce bark
beetle, Ips typographus, provides insight into a major forest pest. *Commun. Biol.* 4:1–9.

Pruisscher P, Nylin S, Gotthard K, Wheat CW. 2018. Genetic variation underlying local
adaptation of diapause induction along a cline in a butterfly. *Mol. Ecol.* 27:3613–3626.

Purcell J, Brelsford A, Wurm Y, Perrin N, Chapuisat M. 2014. Convergent genetic architecture
underlies social organization in ants. *Curr. Biol.* 24:2728–2732.

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, De
Bakker PIW, Daly MJ, et al. 2007. PLINK: A tool set for whole-genome association and
population-based linkage analyses. *Am. J. Hum. Genet.* 81:559–575.

Rane R V., Rako L, Kapun M, Lee SF, Hoffmann AA. 2015. Genomic evidence for role of
inversion 3RP of Drosophila melanogaster in facilitating climate change adaptation. *Mol.*
Ecol. 24:2423–2432.

Raymond M, Rousset F. 1995. An Exact Test for Population Differentiation. *Evolution* 49:1280.

Reeve J, Butlin RK, Koch EL, Stankowski S, Faria R. 2023. Chromosomal inversion
polymorphisms are widespread across the species ranges of rough periwinkles (*Littorina*
saxatilis and *L . arcana*). Early view.

Roesti M, Gilbert KJ, Samuk K. 2022. Chromosomal inversions can limit adaptation to new
environments. 31:1–16.

Roff D a. 1996. The evolution of threshold traits in animals. *Q. Rev. Biol.* 71:3–35.

Rozas J, Ferrer-Mata A, Sanchez-DelBarrio JC, Guirao-Rico S, Librado P, Ramos-Onsins SE,
Sanchez-Gracia A. 2017. DnaSP 6: DNA sequence polymorphism analysis of large data

sets. *Mol. Biol. Evol.* 34:3299–3302.

Saitou M, Masuda N, Gokcumen O. 2022. Similarity-based analysis of allele frequency

distribution among multiple populations identifies adaptive genomic structural variants.

Mol. Biol. Evol. 39:1–19.

Sallé A, Arthofer W, Lieutier F, Stauffer C, Kerdelhué C. 2007. Phylogeography of a host-

specific insect: Genetic structure of Ips typographus in Europe does not reflect past

fragmentation of its host. *Biol. J. Linn. Soc.* 90:239–246.

Schaal SM, Haller BC, Lotterhos KE. 2022. Inversion invasions: When the genetic basis of local

adaptation is concentrated within inversions in the face of gene flow. *Philos. Trans. R. Soc.*

B Biol. Sci. 377.

Schebeck M, Dobart N, Ragland GJ, Schopf A, Stauffer C. 2022. Facultative and obligate

diapause phenotypes in populations of the European spruce bark beetle Ips typographus. *J.*

Pest Sci. 95:889–899.

Schwander T, Libbrecht R, Keller L. 2014. Supergenes and complex phenotypes. *Curr. Biol.*

24:R288–R294.

Skotte L, Korneliussen TS, Albrechtsen A. 2013. Estimating individual admixture proportions

from next generation sequencing data. *Genetics* 195:693–702.

Stauffer C, Lakatos F, Hewitt GM. 1999. Phylogeography and postglacial colonization routes of

Ips typographus L. (Coleoptera, Scolytidae). *Mol. Ecol.* 8:763–773.

Sturtevant AH. 1921. Linkage variation and chromosome maps. *Proc. Natl. Acad. Sci. USA*

7:181–183.

Thompson MJ, Jiggins CD. 2014. Supergenes and their role in evolution. *Heredity (Edinb).*

113:1–8.

Tigano A, Friesen VL. 2016. Genomics of local adaptation with gene flow. *Mol. Ecol.* 25:2144–

2164.

Tigano A, Jacobs A, Wilder AP, Nand A, Zhan Y, Dekker J, Therkildsen NO, Lohmueller K.

2021. Chromosome-Level Assembly of the Atlantic Silverside Genome Reveals Extreme

Levels of Sequence Diversity and Structural Genetic Variation. *Genome Biol. Evol.* 13:1–

17.

Todesco M, Owens GL, Bercovich N, Légaré JS, Soudi S, Burge DO, Huang K, Ostevik KL,

Drummond EBM, Imerovski I, et al. 2020. Massive haplotypes underlie ecotypic

differentiation in sunflowers. *Nature* 584:602–607.

Wang J, Wurm Y, Nipitwattanaphon M, Riba-Grognuz O, Huang YC, Shoemaker D, Keller L.

2013. A Y-like social chromosome causes alternative colony organization in fire ants.

Nature 493:664–668.

Wang Z, Liu Y, Wang H, Roy A, Liu H, Han F, Zhang X, Lu Q. 2023. Genome and

transcriptome of Ips nitidus provide insights into high-altitude hypoxia adaptation and

symbiosis. *iScience* 26:107793.

Weir BS, Cockerham CC. 1984. Estimating F-statistics for the analysis of population structure.

Evolution 38:1358–1370.

Wellenreuther M, Bernatchez L. 2018. Eco-Evolutionary Genomics of Chromosomal Inversions.

Trends Ecol. Evol. 33:427–440.

Wellenreuther M, Mérot C, Berdan E, Bernatchez L. 2019. Going beyond SNPs: The role of

structural genomic variants in adaptive evolution and species diversification. *Mol. Ecol.*

28:1203–1209.

Yang YY, Lin FJ, Chang HY. 2002. Comparison of recessive lethal accumulation in inversion-

bearing and inversion-free chromosomes in Drosophila. *Zool. Stud.* 41:271–282.

Yuvaraj JK, Roberts RE, Sonntag Y, Hou XQ, Grosse-Wilde E, Machara A, Zhang DD, Hansson

BS, Johanson U, Löfstedt C, et al. 2021. Putative ligand binding sites of two functionally

characterized bark beetle odorant receptors. *BMC Biol.* 19:1–21.

Zhao T, Kandasamy D, Krokene P, Chen J, Gershenzon J, Hammerbacher A. 2019. Fungal

associates of the tree-killing bark beetle , Ips typographus , vary in virulence , ability to

degrade conifer phenolics and in fl uence bark beetle tunneling behavior. *Fungal Ecol.*

38:71–79.

1133

1134

1135   **Tables**

1136

Table 1 List of identified chromosomal inversions in the *Ips typographus* genome. ID: inversion

name; Contig: contig name; Size: size of the inversions (Mb); Start and End: coordinates of the

inversion (Mb); Age: approximate age of the inversion in Myr; Odorant receptors: odorant

receptors present within inversion (in the order they appear along the contig sequence). Note that

Inv16.1 and Inv23.1 are parts of the same inversion and Inv16.2 and Inv23.2 are a part of another

single inversion.

ID	Contig	Size	Start	End	Age	Odorant receptors
Inv2	IpsContig2	4.04	12.67	16.71	0.5	
Inv3	IpsContig3	0.14	1.11	1.25	1.1	
Inv5	IpsContig5	10.84	0.00	10.84	1.3	ItypOR33, ItypOR41, ItypOR40, ItypOR10, ItypOR47, ItypOR50, ItypOR29, ItypOR43JOI, ItypOR34, ItypOR52NTE, ItypOR4, ItypOR3, ItypOR53, ItypOR2, ItypOR19
Inv6	IpsContig6	0.34	8.93	9.27	1.0	
Inv7.1	IpsContig7	0.67	0.00	0.67	1.7	
Inv7.2	IpsContig7	6.92	0.00	6.92	1.1	ItypOR1, ItypOR17
Inv9	IpsContig9	3.30	1.71	5.01	0.6	
Inv10	IpsContig10	0.08	6.05	6.13	1.8	
Inv12	IpsContig12	0.07	3.63	3.70	1.5	
Inv13	IpsContig13	4.50	0.00	4.50	1.7	ItypOR28, ItypOR23, ItypOR49, ItypOR27
Inv14.1	IpsContig14	2.08	0.00	2.08	1.9	ItypOR36, ItypOR44, ItypOR18JF, ItypOR20NTE
Inv14.2	IpsContig14	0.67	2.08	2.75	1.0	
Inv14.3	IpsContig14	0.76	2.78	3.54	2.1	
Inv14.4	IpsContig14	0.11	3.73	3.84	2.1	
Inv14.5	IpsContig14	0.57	4.23	4.80	2.3	
Inv14.6	IpsContig14	2.48	0.00	2.48	0.6	ItypOR36, ItypOR44, ItypOR18JF, ItypOR20NTE
Inv15	IpsContig15	1.92	0.86	2.78	1.9	
Inv16.1	IpsContig16	4.83	0.00	4.83	1.6	ItypOR58, ItypOR9, ItypOR11, ItypOR31, ItypOR30, ItypOR16, ItypOR35JF
Inv16.2	IpsContig16	4.83	0.00	4.83	0.7	ItypOR58, ItypOR9, ItypOR11, ItypOR31, ItypOR30, ItypOR16, ItypOR35JF
Inv17	IpsContig17	0.21	2.48	2.69	2.2	
Inv18	IpsContig18	2.32	0.00	2.32	2.1	
Inv22.1	IpsContig22	0.32	0.00	0.32	2.1	
Inv22.2	IpsContig22	0.23	0.42	0.65	2.1	
Inv22.3	IpsContig22	1.23	0.68	1.91	2.0	ItypOR22CTE

Inv22.4	IpsContig22	0.20	1.92	2.12	2.1	
Inv22.5	IpsContig22	2.24	0.00	2.24	0.6	ItypOR22CTE
Inv23.1	IpsContig23	2.10	0.00	2.10	1.4	ItypOR58, ItypOR9
Inv23.2	IpsContig23	1.84	0.26	2.10	0.6	ItypOR58, ItypOR9
Inv26	IpsContig26	0.10	0.12	0.22	2.6	

1143

Figures

Figure 1 Genomic structure and differentiation in *Ips typographus*. (A) Whole-genome PCA,

where colors correspond to what population an individual belongs to. Legend indicates

geographical grouping of populations: S-South, P-Poland, N-North. (B, C) Geographical

distribution and genetic differentiation of the 18 beetle populations analyzed. Blue and red are

the colors of the main genetic groups: southern (red), northern (blue). (D) Genome-wide genetic

differentiation (F_{ST}) calculated between northern and southern groups. Vertical lines separate

different contigs shown in grey and black.

1152

1153

1154

1155

Figure 2 Identification of chromosomal inversions in *Ips typographus*. Each row of the figure panels shows the results of per contig PCA, per contig linkage disequilibrium analysis, and genetic differentiation (F_{ST}) analysis for a selected contig. Panels A-C show results for a contig with no inversions (IpsContig8) and little differentiation between southern and northern populations. Panels D-O show different contigs with increasingly complex inversion patterns: a single inversion (D-F, Inv15); a single inversion with a double-crossover signal (G-I, Inv18); multiple adjacent inversions with one inversion overlapping with the first two inversions on the contig (J-L, Inv14.1, Inv14.2, Inv14.3, Inv14.4, Inv14.5, and Inv.14.6); multiple adjacent inversions with one large inversion overlapping with several smaller ones (M-O, Inv22.1, Inv22.2, Inv22.3, Inv22.4, and Inv22.5). The large overlapping inversion is visible as a yellow background in the LD plot (N). The PCA grouping shown in (M) corresponds to genotypes of the largest inversion on IpsContig22 (Inv22.3), including genotypes that include haplotypes produced with the double crossover event. The last panel (P) shows the 36 largest contigs of the spruce bark beetle genome with inversions indicated in red. Colored dots indicate the inversions that are presented in detail in panels A-O. Colored bars indicate the position of each inversion. Colors of the bars in plots E, K and N correspond to adjacent F_{ST} plots (panels F, L and O), where F_{ST} is calculated between haplotypes of indicated inversion along the contig in question. In the plot I F_{ST} is calculated between three haplotypes: homozygotes for two arrangements and the homozygotes for recombinant haplotype created after putative double crossover event. Dots in the PCA plots represent individual beetles and are colored according to the heterozygosity of the individual (darker color represents low heterozygosity). Both axes in LD plots (B, E, H, K, N) represent positions along the contig in megabases (Mb); low levels of linkage are shown in blue and higher levels in yellow to dark red. Sex chromosome is indicated with a grey star in P panel.

1184   **Figure 3** Correlation between inversion haplotype frequency and population latitude for
1185   chromosomal inversions detected in European *Ips typographus* populations. Significant
1186   correlations are indicated with asterisks (*: p < 0.05; **: p < 0.01; ***: p < 0.001). Inversions
1187   harboring odorant receptors are indicated in green.
1188

1189
1190
1191

1192

Figure 4 Genotype-environment associations across 36 contigs in the *Ips typographus* genome.

Different colors represent different contigs. Each dot represents a single SNP. The upper three

panels show results for all SNPs and the lower three panels show results using data where each

inversion was coded as a single locus (shown in a blue circle). MAF: Minor Allele Frequency.

PC1, PC2, and PC3 represent the first three of PCA used to summarize the environmental

variation among populations.

1199

1200
1201
1202

1203

Figure 5 Mutation load analysis in *Ips typographus*.

(A) The ratio of nonsynonymous to synonymous nucleotide diversity π_N/π_S computed for 200 kb
windows along collinear parts of the genome (COL) and different inversion haplotypes (blue:
major haplotype; yellow: minor haplotype; green: haplotype produced by double crossover
events between minor and major haplotypes). n = 4 to 55 windows per inversion haplotype and n
= 531 for the collinear genome. Inversion haplotypes that were present in less than four
individuals, did not include four or more 200 kb windows or included fewer than five genes were
not included in the analysis. For better visibility, π_N/π_S outliers above 0.5 are not shown (a total
of 58 π_N/π_S values, 45 of them in COL). The lower and upper box hinges correspond to the first
and third quartiles, whiskers show 1.5 * the inter-quartile range. (B) Distribution of π_N/π_S values
per 200 kb window (C) Correlation between mean π_N/π_S for all inversion haplotypes and their
frequency across spruce bark beetle populations.

1216

1217