Complex genomic landscape of inversion polymorphism in Europe's most destructive forest

pest

Anastasiia Mykhailenko[1,2], Piotr Zieliński[1], Aleksandra Bednarz[1], Fredrik Schlyter[3,4], Martin N.

Andersson[5], Bernardo Antunes[1], Zbigniew Borowski[6], Paal Krokene[7], Markus Melin[8], Julia

Morales-García[1,2], Jörg Müller[9,10], Zuzanna Nowak[1], Martin Schebeck[11], Christian Stauffer[11],

Heli Viiri[12], Wiesław Babik[1], Krystyna Nadachowska-Brzyska[1]

[1]Institute of Environmental Sciences, Faculty of Biology, Jagiellonian University, Gronostajowa

7, 30-387, Kraków, Poland

[2]Doctoral School of Exact and Natural Sciences, Jagiellonian University, ul. Łojasiewicza 11,

30-348, Kraków, Poland

[3]Chemical Ecology, Department of Plant Protection Biology, Swedish University of Agricultural

Sciences Alnarp, PO Box 190, 234 22 Lomma, Sweden

[4] ETM, Faculty of Forestry and Wood Sciences, Czech University of Life Sciences Prague,

Kamýcká 129, 165 00, Praha, Czechia

[5]Department of Biology, Lund University, Sölvegatan 37, 223 62, Lund, Sweden

[6] Forest Research Institute, Sękocin Stary, Braci Leśnej 3, 05–090 Raszyn, Poland

[7] Division of Biotechnology and Plant Health, Norwegian Institute of Bioeconomy Research,

Høgskoleveien 8, Ås, 1433, Norway

[8]Natural Resources Institute Finland, Yliopistokatu 6b, 80100 Joensuu, Finland

[9]Field Station Fabrikschleichach, Animal Ecology and Tropical Biology, Biocenter, University

of Würzburg, Glashüttenstrasse 5, 96181 Rauhenebrach, Germany

[10]Bavarian Forest National Park, Freyungerstrasse 2, 94481 Grafenau, Germany

[11]Institute of Forest Entomology, Forest Pathology and Forest Protection, Department of Forest

and Soil Sciences, University of Natural Resources and Life Sciences Vienna (BOKU), Peter-

Jordan-Straße 82/I, 1190, Vienna, Austria

[12]UPM-Kymmene, UPM Forest, BOX 85, 33100, Tampere, Finland

Corresponding author: Krystyna Nadachowska-Brzyska

Abstract

In many species, polymorphic inversions underlie complex phenotypic polymorphisms and appear to facilitate local adaptation in the face of gene flow. Multiple polymorphic inversions can co-occur in a genome, but the prevalence, evolutionary significance, and limits to complexity of genomic inversion landscapes remain poorly understood. Here, we examine genome-wide variation in one of Europe's most destructive forest pests, the spruce bark beetle *Ips typographus*, scan for polymorphic inversions, and test whether inversions are involved in key adaptations in this species. We sampled 244 individuals from 18 populations across the species' European range and, using a whole-genome resequencing approach, identified 27 polymorphic inversions covering at least 28% of the genome. The inversions vary in size and in levels of intra-inversion recombination, are highly polymorphic across the species range, and often overlap, forming an extremely complex genomic architecture. We show that the heterogeneous inversion landscape is likely maintained by the combined action of several evolutionary forces and that inversions are enriched in odorant receptor genes encoding key elements of recognition pathways for host plants, mates, and symbiotic fungi. Our results indicate that the genome of this major forest pest of growing social, political, and economic importance harbors one of the most complex inversion landscapes described to date and is pushing the limits of genomic architecture complexity.

Introduction

The eight spined spruce bark beetle (*Ips typographus*; hereafter spruce bark beetle) is one of the most destructive insect pests in Europe, causing mass mortality in spruce-dominated forests[1,2]. The extent of recent outbreaks is unprecedented and impacts will likely increase in the coming decades in response to climate change[3–5]. Increasing bark beetle attacks and other forest disturbances have already triggered social and political conflicts in parts of Europe and have highlighted the urgent need for improved management strategies[2,6,7]. Indeed, a rapidly growing body of research focuses on the species' ecology, the causes and consequences of outbreaks, and their social aspects[2,4,8,9]. Despite this enormous interest, one aspect of the species' biology remains largely unexplored: we know almost nothing about the species' genome-wide variation and the evolutionary mechanisms that shape this variation. The lack of population genomics studies restricts our understanding of the genomic basis of adaptation and adaptive potential in

the spruce bark beetle. Such information could provide a critical missing link between applied

and basic research and serve as a foundation for effective management. This is particularly

important because, as discovered and described in this study, the spruce bark beetle genome

harbors an extremely complex inversion polymorphism landscape that may play a critical role in

many evolutionary processes, including key species adaptations[10–12].

Polymorphic chromosomal inversions are chromosomal segments that occur in two orientations

within populations: collinear and inverted haplotypes/arrangements. Inversions have been shown

to be involved in speciation, local adaptation, and/or maintenance of complex phenotypes[12–15].

This is due to a key property of inversions: they suppress recombination within heterozygotes

and thereby prevent separation of coadapted variants. Because of their role as recombination

modifiers, inversions can act as supergenes, i.e., large elements of genomic architecture with

multiple linked functional elements[16]. Supergenes keep coadapted alleles together in the face of

gene flow and prevent the formation of maladaptive recombinant genotypes.

Classic examples of supergenes include inversions associated with different mating strategies in

ruffs (*Calidris pugnax*)[15], mimicry phenotypes in Heliconius butterflies[17], and social organization

in fire ants (*Solenopsis* spp.)[18,19]. In many other species, polymorphic inversions define locally

adapted ecotypes[20,21] or exhibit spatial frequency differences, e.g. by forming geographic and

climatic gradients[22,23]. While the vast majority of described cases are organisms with one or a

few inversions, several recent studies have reported species with many polymorphic inversions[24–

26]. These recent findings raise questions about the prevalence and evolutionary significance of

polymorphic inversions in natural populations. Are inversion-rich genomes the exception or the

rule? How much of the genome can be situated within polymorphic inversions and,

consequently, how large can the fraction of the genome with reduced recombination be? The

latter question is particularly important because, in addition to suppressing recombination and

keeping coadapted alleles together, inversion heterozygotes will also prevent the formation of

new allelic combinations and thus reduce the efficacy of natural selection[27].

Given the importance of recombination in purging deleterious mutations[28], the presence of

multiple inversions that cover a substantial fraction of the genome raises other questions. First, as

the degree of recombination suppression depends on the number of heterozygous genotypes, how

frequent are inversion haplotypes in inversion-rich species? Second, are balanced

polymorphisms (where both inversion arrangements are equally common across the species'

range) more common than inversions with one common and one rare haplotype? Third, how

common are mechanisms that mitigate negative consequences of reduced recombination? Such

mechanisms include double crossover-events and gene conversion in heterozygotes, which can

reshuffle allelic content between large parts of two inversion arrangements and thereby reduce

mutational load and create new haplotypes[10,29].

Long-term persistence of two inversion haplotypes can be facilitated by two main types of

selection: divergent and balancing[30]. Divergent selection can favor different inversion genotypes

in different environments and, when coupled with reduced migration between divergent

populations, can lead to speciation. Even when intraspecific gene flow is high, divergent

selection can lead to divergent ecotypes associated with locally advantageous inversion

genotypes. Alternatively, balancing selection may maintain balanced inversion polymorphisms

over time via several, not mutually exclusive, mechanisms, such as overdominance, negative

frequency dependence, antagonistic pleiotropy, and spatially or temporally varying selection[30,31].

Importantly, regardless of the selection mechanism, inversions will accumulate mutations

independently since recombination is suppressed in inversion heterozygotes. This will lead to

differentiated allelic content and increased differentiation between inversion haplotypes over

time[30]. Each inversion haplotype can be treated as a separate "population" with a size that

corresponds to the frequency of that arrangement within studied population or species. Rare

inversion arrangements will suffer from a high mutational load due to reduced recombination

and limited purging since homozygotes are rare. However, deleterious mutations can also

accumulate on more frequent inversion haplotype[10] and lead to associative overdominance. This

contributes to the maintenance of inversion polymorphisms since independent accumulation of

mutations continues over the time but recessive deleterious alleles private to an inversion

arrangement are not visible to selection in inversion heterozygotes.

Here, we investigated genome-wide variation across spruce bark beetle populations with a

special focus on chromosomal inversion polymorphisms. We found one of the most complex

polymorphic inversion architectures described to date and tested several evolutionary

mechanisms, including directional and balancing selection, that can maintain inversion

polymorphisms in the genome. We also tested associations between inversion polymorphisms

and the two key fitness-related traits diapause and olfaction. Our results suggest that inversions

are involved in key species adaptations and raise questions about the prevalence and role of

inversion polymorphisms in species characterized by large effective population sizes, little

geographic subdivision, and no clear phenotypic variation across the species range.

Results and Discussion

After careful filtering of the whole-genome re-sequencing data, we used 240 individuals in

downstream analyses. The mean per individual sequencing depth was $23.2\times$ (range: 5 - $53\times$).

Sequencing coverage on IpsContig9 was consistently lower in individuals sexed as males (on

average 0.57 individual coverage). Thus, we considered IpsContig 9 to be a sex (X)

chromosome. After quality filtering, we retained 5.245 million SNPs covering the entire genome

assembly but analyzed a subset of 5.067 million SNPs located on the 35 longest contigs and

representing 75% of the genome assembly.

Complex genomic inversion landscape

Twenty-nine candidate inversions ("inversions" henceforth) were identified following the criteria

described in the Methods section (Table 1, Figure 1, Figure S1, and Figure S2). Two possible

inversions (Inv16.1 and Inv23.1) were most likely part of the same inversion: the same beetle

individuals were genotyped as both homo- and heterozygotes, which is unlikely to occur for two

unlinked inversions. The same situation was found for Inv16.2 and Inv23.2. No other inversions

were in strong linkage disequilibrium (LD) with each other, which would suggest co-segregation

(Figure S3). Thus, overall, we found 27 inversions in 17 contigs, including one located on the X

chromosome (Inv9). Inversion size varied from 0.1 to 10.8 Mb (Table 1) and inversions

constituted 28% of the analyzed part of the genome. Estimated inversion age ranged from 0.5 to

2.6 My (Table 1, assuming a mutation rate of 2.9×10^{-9}; see Table S1 for results for different

mutation rates). Inversion regions exhibited a reduced population recombination rate (rho, Figure

S4) and moderate to high genetic differentiation between inversion arrangements (F_{ST} between

homozygous individuals was 0.15-0.64; Figure 1; Figure S1).

While 12 contigs contained single inversions, five contigs showed patterns consistent with

multiple adjacent or overlapping inversions (Figure 1, Figure S1, Figure S2). IpsContig14 and

IpsContig22 contained complexes of multiple adjacent and sometimes overlapping inversions

(Figure 1 j-o), and three other contigs contained two overlapping inversions each (IpsContigs: 7,

16 and 23, Figure S1). Additionally, four putative double crossover-events were identified within

four inversions (Inv5, Inv18, Inv22.3, Inv22.4; Figure 1, Figure S1, Figure S2). In these cases,

LD clusters were separated by regions of lower LD and intermediate groups of individuals were

visible between the main clusters along the first principal component.

It was not surprising that we detected polymorphic inversions in the spruce bark beetle genome,

as there are many well-known examples of polymorphic inversions in natural populations[12].

What was striking, however, was the extremely complex genomic landscape of polymorphic

inversions we found in this species. The spruce bark beetle has at least 27 large inversions

covering a substantial part of the genome (28% of the analyzed part of the genome, but this is

probably an underestimate since we focused on ca. 75% of the genome and on large inversions >

0.1 Mb only). Numerous (a dozen or more) polymorphic inversions have so far only been

described for a few species (e.g.[26,32,33]). It is still an open question whether many polymorphic

inversions within species is the exception or the rule.

The exceptionally complex inversion architecture we found in the spruce bark beetle, with

multiple adjacent and often overlapping inversions, resembles well known examples from

Heliconius butterflies[34] or fire ants[35]. In these insects multiple adjacent inversions are the basis

for mimicry phenotypes and complex social organization, respectively. The presence of clusters

of adjacent inversions and inversion overlaps are consistent with theoretical expectations of

stepwise extension of recombination suppression on supergenes[36] and with a highly polygenic

architecture of adaptation[37].

Genome-wide variation and its geographic structuring – collinear vs. inversion regions

Analyses based on both the whole genome and collinear parts only revealed a clear latitudinal

structuring of genetic variation in the spruce bark beetle (Figure 2). PCA and NGSadmix

supported the presence of two distinct genetic groups corresponding to southern and northern

populations, with Polish populations showing varying degrees of admixture between the two

clusters. Based on these results, we divided the 18 studied populations into a northern, southern,

and Polish group. Despite unambiguous NGSadmix-division into two genetic clusters, the

genome-wide genetic differentiation between the northern and southern group was extremely

low ($F_{ST} = 0.021$). Similarly, F_{ST} between all population pairs showed low levels of

differentiation, ranging from 0.000 to 0.035 (Table S2). Mean genome-wide nucleotide diversity

was moderate ($\pi = 0.0062$) and per population π ranged from 0.0055 to 0.0066 (Figure S5).

There was a weak negative correlation between nucleotide diversity and latitude ($r^2 = 0.32$, p =

0.033, Figure S6), as northern populations had slightly lower genetic variation than southern

populations ($\pi_{southern} = 0.0065$; $\pi_{northern} = 0.0061$; $\pi_{Polish} = 0.0066$). Southern populations had an

excess of rare alleles and, consequently, had more negative Tajima's D values along the genome

than northern populations (mean Tajima's D was -0.458 and -0.062 in the southern and northern

group, respectively; Figure S7). All these results are consistent with previous phylogeographic

studies of the spruce bark beetle that analyzed a much smaller number of genetic markers. The

data suggests high levels of connectivity among spruce bark beetle populations and a very recent

differentiation into two genetic clusters[38–41]. More recent RADseq data confirms a very weak

genetic structure in the spruce bark beetle across much of Sweden[42], as is expected in a species

with high dispersal[43] and recent divergence.

Inversion regions in the spruce bark beetle did not structure geographically into a southern and

northern group. Almost all identified inversions were polymorphic across the European species

range, except for one inversion (Inv9) that was polymorphic only within northern populations.

For three inversions (Inv2, Inv5, Inv16.2+Inv23.2) unambiguous genotyping was only possible

across part of the species range. The differentiation between inversion haplotypes was high,

suggesting long-term persistence of inversion polymorphisms within this species (Figure 1;

Figure S1). According to our age estimates, the origin of the inversions in the spruce bark beetle

predates the Last Glacial Period and many inversions may be several million years old, likely

also predating the within-species differentiation into a southern and northern group. This was not

unexpected, as many known inversion polymorphisms have been segregating within species for

hundreds of thousands or millions of years (see table in[12]), sometimes even persisting through

multiple speciation events[44].

Inversions and key fitness-related traits

Inversion polymorphism is often associated with the maintenance of complex polymorphic

phenotypes[15,45]. Although the spruce bark beetle does not exhibit easily identifiable phenotypes,

such as distinct color patterns or mating types, we were able to test for associations between

inversions and two complex traits of key importance for many insects: diapause and olfaction.

Diapause allows species to suspend development during unfavorable conditions. While multiple

environmental factors can influence this complex process, many aspects of diapause, such as its

induction and termination, are heritable[46] and may be controlled by a small number of loci or be

highly polygenic[47–50]. The spruce bark beetle exhibits two diapause strategies that could be

associated with polymorphic inversions: a facultative photoperiod-regulated diapause and an

obligate photoperiod-independent diapause[51]. However, we found no association between

diapause phenotypes and inversion genotypes (exact G test, Table S3, Figure S8), nor did we

find any highly differentiated genomic regions between facultatively and obligately diapausing

individuals (Figure S9). This suggests a polygenic nature of diapause phenotypes in the spruce

bark beetle.

Olfaction is another key fitness-related trait in many insects, including bark beetles. Insect

odorant receptors (ORs) are encoded by a large and dynamically evolving gene family. Some of

the receptors are evolutionarily conserved across species within insect orders, however, many are

species- or genus-specific. In the spruce bark beetle, detection of odorants is essential for host

and mate finding, as well as recognition and maintenance of symbiosis with specific fungi[52,53].

We examined 73 antennally expressed ORs[54] and found that 46% of these were located within

inverted regions (Table 1), even though inverted regions constituted only 28% of the analyzed

part of the genome. A permutation test confirmed this over-representation of OR genes within

inversions (p = 0.03). In addition, several *Ips*-specific ORs (5 out of 7 ORs from an *Ips*-specific

OR clade[55]) were located in inverted regions, specifically on IpsContig13 (4 out of 7 ORs).

These 5 ORs (ItypOR23, ItypOR27, ItypOR28, ItypOR29, and ItypOR49) have been

functionally characterized, responding to selectively compounds primarily produced by beetles

(pheromones), the host tree, or fungal symbionts, respectively (Powell et al., 2021; Hou et al.,

2021). We found no difference in the OR composition (no OR deletions) between inversion

haplotypes.

Two inversions harboring multiple OR genes showed significant latitudinal variation (Figure 3;

Inv13 and Inv16.1+23.1, Table 1). Interestingly, one of the inversion, Inv13, include a gene

encoding ItypOR23, receptor that has been previously shown to primarily respond to an odor

from fungi[55]. We hypothesize that different OR alleles associated with these inversions are

involved in spruce bark beetle interactions with fungal associates present in different parts of

Europe. Not only are spruce bark beetle populations exposed to different fungal species

throughout the beetle's range, but individual beetles may also have preferences for different

fungal species[53,56]. German spruce bark beetles have, for example, been shown to be more

attracted to fungal species that are common in Germany (*Grosmannia penicillata,*

Endoconidiophora polonica) than to rarer species (*Leptographium europhioides*). Unpublished

data from Swedish beetles suggests that they are more attracted to *L. europhioides*, which is

common in Sweden (personal communication D. Kandasamy). Although preliminary, these

observations suggest that beetle preferences may be tuned to the local fungal flora and we

speculate that inversions may be involved in recognition of region-specific fungal species.

Several other interesting behavioral strategies are polymorphic among spruce bark beetle

individuals, including the existence of pioneer individuals that are the first to infest host trees,

and re-emergence of females after egg laying to establish so-called sister broods in new trees.

Other less obvious/visible phenotypes could also be associated with inversion polymorphisms.

Comprehensive research, both bottom-up and top-down, is needed to understand the relationship

between spruce bark beetle phenotypes and the inversion polymorphism landscape.

Evolutionary mechanisms maintaining inversion polymorphism in the spruce bark beetle

Several non-mutually exclusive evolutionary processes can maintain polymorphic inversions

within species, particularly divergent and balancing selection[12,30]. The importance of divergent

selection has been postulated based on allele frequency patterns and associations of polymorphic

inversions with local adaptations that persist despite extensive intraspecific gene flow[57]. For

example, a recent study of deer mice (*Peromyscus maniculatus*)[26] identified multiple

polymorphic inversions with clinal variation across environmental gradients in two distinct

habitats. Such frequency changes have also been reported across hybrid zones[33] or latitudinal

gradients[23]. Although the spruce bark beetle does not occupy distinct environmental niches, it

inhabits forests across a very wide latitudinal gradient (spanning at least 16 degrees). It is also a

species with high dispersal capacity and extensive gene flow, as indicated by low F_{ST} across its

range. We found a significant correlation between the frequency of inversion haplotypes in

populations and geographic location (latitude) for five inversions (Figure 3; r^2 ranged from 0.34

to 0.68). There were no significant correlations between haplotype frequencies and longitude

(Figure S10). In most cases, differences in inversion haplotype frequencies were small (Figure

3), except for the two inversions with the strongest correlations ($r^2 > 0.6$; Inv12 and Inv13). We

found no association between inversion genotypes and climate or land cover variables (Figure 4),

but several SNPs showed significant correlation with the first principal component in a PCA of

many environmental variables. These results indicated that there probably is selection across

environmental gradients in the spruce bark beetle but that this is not the only, or even a major,

force maintaining inversion polymorphism within the species.

While the 'local adaptation' hypothesis is a major hypothesis proposed to explain inversion

polymorphism[26,58,59], balancing selection and related mechanisms may also be important in

maintaining polymorphic arrangements[10,30,60]. Such mechanisms include overdominance,

associative overdominance, frequency-dependent selection, and spatially and temporally varying

selection. We found no support for overdominance playing a role in the spruce bark beetle, as no

excess of inversion heterozygotes was detected in any of the populations, geographic regions or

across the whole species range. We therefore looked more closely at mutation load, which can

say something about the role of associative overdominance in the maintenance of inversion

polymorphism. Theory predicts that recessive deleterious mutations will accumulate on both

inversion arrangements but that most of these mutations will be private to only one

arrangement[10,30,61]. This would lead to associative overdominance, as in heterozygotes the effects

of deleterious recessive alleles on one arrangement would be masked by the wild-type alleles on

the other arrangement. The result would be long-term maintenance of the inversion

polymorphism, resulting in strong divergence between inversion haplotypes[10,61,62].

Interestingly, several stable evolutionary scenarios that maintain polymorphic inversions are

possible (for details see Figure 4 in[10]). These scenarios differ in the expected mutation load,

fitness, and frequency of the corresponding genotypes. Given the haplotype frequencies we

observed in spruce bark beetle inversions, two scenarios are likely. First, that minority

arrangements suffer from higher mutation load (due to reduced recombination and lower

population size) but are maintained in the population at low frequency due to, e.g., associative

overdominance. Such a mechanism would favor balanced inversion polymorphisms of

intermediate to large sizes[63,64] and has been shown to play a role in maintaining polymorphic

inversions in several insect species[34,65]. Second, mutation load may accumulate on one or both

inversion arrangements but be mitigated by the haplotype structuring process, i.e., the existence

of multiple diverged sub-haplotypes among inversion homozygotes that reduces the mutation

load within homozygotes. If this process operates within one or both inversion haplotypes it may

result in more equal frequencies of alternative inversion haplotypes. However, such a mechanism

is only possible when genetic variation and mutation load is high[10].

In contrast to these theoretical expectations, we observed no sign of increased mutation load

(measured by the π_N/π_S ratio) in inversion regions compared to the collinear part of the spruce

bark beetle genome (Table S4; Figure 5). We also found no sign of haplotype structuring capable

of reducing mutation load within inversion homozygotes (Figure S11). The only significant

within-homozygote clustering we observed was in a few inversion haplotypes, divided

individuals into southern and northern clades and suggests that divergent selection has been

acting on one of the inversion arrangements, rather than haplotype structuring being associated

with mutation load (Figure S11). These results are consistent with observations of no significant

mutation load in other species where inversion haplotypes are subject to divergent selection that

results in geographic structuring[26,66]. In such cases, inversions facilitate adaptive divergence but

do not tend to accumulate a mutation load. However, geographic clustering of inversion

polymorphisms is weak in the spruce bark beetle, and many inversions appear to have a slightly

lower mutation load associated with the more common inversion haplotype (Figure 5, Table S4).

It is possible that accumulation of a mutation load in the spruce bark beetle is mitigated by high

effective population size in this species and/or gene conversion and double crossover events,

which despite their apparent rarity have been detected in several spruce bark beetle inversions.

Overall, the absence of heterozygote excess observed in the spruce bark beetle does not support a
role of overdominance in the maintenance of inversion polymorphism in this species. Likewise,
the absence of an elevated mutation load in inverted regions does not support a role of
associative overdominance either. However, since we only have genomic data available, we
cannot conclusively rule out that other potential mechanisms, such as negative frequency-
dependent selection or antagonistic pleiotropy, could maintain balanced inversion
polymorphisms. Additional temporal data are needed to test whether temporally varying
selection has affected the frequencies of inversion haplotypes in the spruce bark beetle. Our
results indicate that inversions in this species are maintained as polymorphic by a complex
interaction of different, not mutually exclusive mechanisms. Further research is essential to
determine the role of different mechanisms.

Far-reaching consequences of having an inversion-rich genome

The presence of multiple polymorphic inversions can have significant consequences for the
evolution of a species, as well as for evolutionary inferences based on genome-wide
polymorphism data. Importantly, polymorphic inversions are a reservoir of genetic variation that
can facilitate adaptation to rapidly changing environments. Indeed, several studies have shown
that polymorphic inversions support rapid adaptation to changing climatic conditions[67–69] or
adaptive tracking of fluctuating environments[70]. Spruce bark beetle populations are subject to
seasonal weather changes and a rapidly changing environment due to strong anthropogenic
pressures. Warmer weather and drought periods have been associated with a predicted
intensification of bark beetle outbreaks[3,4,6], which may act as a strong selection factor within
bark beetle populations. Whether inversions are involved in rapid adaptations in the spruce bark
beetle is an open question that requires further investigation.

Abundant polymorphic inversions within the genome can have far-reaching consequences for
inferences about demographic history and selection. It is well known that non-equilibrium
demography and selection can leave similar genomic signatures. Traditionally, demographic
analyses have used non-coding parts of the genome, based on the assumption that directional
selection mostly affects protein-coding regions. However, growing evidence for the importance

of background selection in shaping genome-wide diversity is moving the field towards

incorporating linked selection into inferences of demographic history[71,72]. We believe that new

approaches should also consider the potential influence of polymorphic inversion landscapes,

because variation patterns of inverted regions can be shaped by different types of balancing

selection. In addition, genomics scans for selection in inversion-rich genomes may be biased due

to reduced recombination within inversion regions. Importantly, the effect of reduced

recombination may extend outside inversions[73,74].

Materials and methods

Study system

The spruce bark beetle, *Ips typographus* (Coleoptera: Curculionidae: Scolytinae), plays a key

role in Eurasian forest ecosystems. Under normal (endemic) conditions, this forest pest attacks

mainly weakened Norway spruce (*Picea abies*) trees. However, if tree resistance is compromised

by certain abiotic disturbances (e.g. snowbreaks, windfalls, high temperatures, drought), an

increased availability of stressed trees can trigger mass-propagation, leading to rapid population

increase and devastating outbreaks. During outbreaks, the beetles attack healthy trees and cause

massive tree mortality with hundreds or thousands of hectares of dead spruce stands.

During the first decade of the 21st century the spruce bark beetle killed an estimated 14.5 million

m^3 of timber per year on average, and this number is expected to increase due to climate

change[75]. The Czech Republic provides a particularly striking example of the beetles' destructive

potential. During the peak outbreak years 2017-2019 the beetles killed annually 3.1-5.4% of the

country's growing stock of Norway spruce, which in 2019 translated to 23 million m^3. In some

regions there was an almost total depletion of Norway spruce[76]. From a socio-economic

perspective, bark beetle outbreaks and subsequent intensive salvage cuttings negatively affects

the quality of life for people living in outbreak areas and cause serious disturbances to the wood

market. In the Czech Republic, decreasing timber prices caused severe revenue losses for forest

owners and required state interventions in the amount of 260 million EUR in 2018-2019[76].

Historically, spruce forests in Central Europe have been most heavily affected by bark beetle

outbreaks, while in northern Europe outbreaks have been less frequent and destructive[1].

However, this may change with climate warming that probably will make the boreal forests of

northern Europe more vulnerable to bark beetle outbreaks[3,5,7,77,78]. As an example, heatwaves and

severe summer drought in Sweden in 2018 initiated a bark beetle outbreak killing over 30

million m^3 Norway spruce the next years[79,80].

Sampling

Adult spruce bark beetles were collected with pheromone-baited traps in the spring and summer

of 2020. In total, we sampled 18 populations throughout Europe with 13-14 individuals per

locality (244 individuals in total) (Figure 2; Table S5). Throughout the text, we use the term

'population' to refer to a particular site or a collection of closely situated sites (within about 50

km). In Austria, we pooled individuals from three localities that were up to 120 km apart because

of small sample sizes (five beetles or less per site). Populations from Scandinavia will be referred

to as northern populations (or the northern group) and populations from central Europe will be

referred to as southern populations (or the southern group). Polish populations are considered

separately from other central European populations (due to high admixture proportions from

northern group identified in downstream analysis). Beetles were brought alive to the laboratory,

kept on a paper diet for several days, dissected, sexed based on genitalia morphology, and

subjected to DNA extraction (described below).

DNA extraction and genome re-sequencing

DNA was extracted from the whole body of dissected beetles using the Wizard Genomic DNA

Purification Kit (Promega). The concentration of extracted DNA was estimated using a Qubit

fluorometer (Thermo Fisher Scientific). Genomic libraries were prepared with NEBNext Ultra II

FS DNA Library Prep with Beads (New England Biolabs), with single indexes. Individual

libraries were combined into three pools and 2×150 bp paired-end sequenced in three lanes of a

S4 flowcell using the NovaSeq 6000 instrument and v1 sequencing chemistry (Illumina Inc.).

Sequencing was done by the National Genomics Infrastructure, SNP&SEQ Technology Platform

(Uppsala, Sweden). To assess the overall genotyping error, we prepared and sequenced duplicate

libraries for nine individuals.

Data preparation and filtering

Details of raw data processing and filtering are described in the Supplementary Files. Shortly, raw reads were mapped to the reference genome[81] using Bowtie 2[82]. Duplicated reads were removed using Picard MarkDuplicates (Broad Institute 2019). To detect and correct systematic errors in base quality scores, recalibration was done using the Genome Analysis Toolkit (GATK), BaseRecalibrator, and ApplyBQSR[83,84]. Variant calling and genotyping was done using GATK HaplotypeCaller, CombineGVCFs, and GenotypeGVCFs. GATK VariantRecalibrator and ApplyVQSR were used to calculate and filter (by variant) quality score log-odds (VQSLOD). Bcftools[85] was used to remove insertions and deletions (indels) as well polymorphisms five bases up- and downstream. GATK VariantFiltration was applied to mask all genotypes with low sequencing depth or low genotype quality[83,84]. Variants which were not biallelic single nucleotide polymorphisms or did not meet the recommended hard filtering thresholds (GATK Team, see Supplementary materials) were filtered out. To filter out polymorphisms that could come from duplicated regions we removed variants located within reapeat-masked regions of the genome[81], variants with excessive overall coverage, and variants with heterozygote excess. Variants for which genotypes could be detected in less than half of the individuals were also removed. We used PLINK[86] to detect sample contaminations, swaps and duplications, and unknown familial relationships (e.g. sibling pairs present in the data) which might bias downstream analyses. Individuals with excessive coverage were removed, as these could be a result of human errors during library preparation or pooling. We focused on contigs longer than 1 Mb that together constituted 78% of the genome assembly, i.e. a total of 186 Mb. Since part of IpsContig33 had high similarity to mtDNA this contig was not included in the downstream analysis. Genotyping error was assessed using GATK Genotype Concordance.

Genome-wide genetic variation and its geographic structuring

Genome-wide genetic structuring was explored by PCA using PLINK. The most likely number of genetic clusters and admixture proportions was estimated using NGSadmix[87]. The analysis was run for five different K-values (1-5; 10 replicates per K-value), using a minor allele frequency (MAF) filter of 0.05 and 10,000 iterations, and a SNP dataset that was pruned for linkage disequilibrium using PLINK (–indep-pairwise 50 10 0.1 option). To choose the most likely number of genetic clusters, the results were examined using CLUMPAK

(http://clumpak.tau.ac.il/index.html). To examine the influence of inversions on genetic

clustering and to facilitate inversion genotyping, NGSadmix was run separately for (1) all

autosomal contigs without potential inversions and (2) each potential inversion.

To assess genetic differentiation among different spruce bark beetle populations Weir and

Cockerham's[88] F_{ST} was estimated using VCFtools[89]. F_{ST} was calculated between population

groups identified by NGSadmix, as well as among all population pairs. Additionally, we

summarized F_{ST} values in 100 kb overlapping windows (using 20 kb steps) along the contigs.

Window-based analyses were done for each contig and each population pair. In addition,

absolute sequence divergence (d_{xy}), nucleotide diversity, and Tajima's D statistic were estimated

and summarized for 100 kb non-overlapping windows using ANGSD[90]. These statistics were

calculated for each population separately and were based on a maximum likelihood estimate of

the folded site frequency spectrum (SFS). We excluded sites with mapping quality below 30

phred, quality score below 20, and coverage less than three times the population sample size and

more than three times the average coverage, following the approach used in Delmore et al.[91]. The

ANGSD calculations were based on allele frequencies estimated from genotype likelihoods[92] and

ngsPopGen scripts (https://github.com/mfumagalli/ngsPopGen).

Population recombination rates were estimated following the approach used in Jones et al.[93],

using 20 individuals from the southern and northern group (10 individuals per group). Watterson

theta estimates were used to create a custom likelihood lookup table using the Ldhat program

complete[94]. The *Interval* program was used to estimate the population recombination rate across

investigated contigs in 1 Mb segments. The *interval* algorithm was run for 2 million iterations

and the chain was sampled every 10,000 iterations with a burn-in of 100,000 generations (*stat*

program in Ldhat package, block penalty = 5). Population recombination rates were estimated

for each contig, and the results were summarized in non-overlapping 100 kb windows using a

custom perl script.

Identification of inversions, their geographic distribution, and variation patterns

Potential chromosomal inversion regions were identified based on the results of per contig PCAs,

local PCAs, patterns of heterozygosity, and LD clustering. Per contig PCAs were performed

using PLINK[86]. Local PCAs were performed using the *lostruct* R package[95], following the

approach described in Huang et al.[96]. Linkage disequilibrium among SNPs (thinned by selecting

one SNP every 10 kb; MAF > 5%) was calculated for each contig using PLINK. We considered

a genomic region to be an inversion region if (1) local PCA analysis identified the region as an

outlier, and/or (2) the region exhibited high LD (most SNPs having $r^2 > 0.4$), and (3) PCA

performed on SNPs from this region separated individuals into two (or more) distinct groups

with heterozygosity patterns matching the expectation of at least one group having lower levels

of heterozygosity (i.e., the group consisting of individuals with homozygous inversion).

Genotyping of individual beetles with respect to the inversion haplotypes they carried was done

based on inversion region PCA1 loading scores and/or inversion-specific NGSadmix clustering

(with K = 2 inversion heterozygotes having mixed ancestry in approximate 50/50 proportions;

Figure S12). In a few special cases (such as overlapping inversions or complex double crossover

patterns) genotyping was done based on inversion region PCA2 loading scores or was limited to

either the southern or northern group. Contigs with less than 10,000 variants provided ambiguous

results and were excluded from genotyping. Putative inversion boundaries were defined based on

local PCAs and sharp borders detected in LD clusters.

Inversion genotype and haplotype frequencies were calculated using an in-house R-script.

Frequencies were calculated (1) within each population, (2) within the southern and northern

group, and (3) for all sampled populations combined. Deviations from Hardy-Weinberg

equilibrium were estimated for all three datasets. Inversions with only two haplotypes were

tested using exact Fisher tests. Inversions with more than two haplotypes (including recombinant

haplotypes between two inversion arrangements) were tested using permutation test, and sex

chromosome inversions were tested as described in Graffelman & Weir[97]. All tests were done

using the R package *HardyWeinberg*[98]. To investigate if inversion haplotypes differed in

frequency along environmental gradients, Pearson correlation coefficients between inversion

haplotype frequencies and latitude/longitude was calculated.

To assess levels of genetic differentiation between inversion haplotypes, F_{ST} and d_{xy} between

alternative inversion haplotypes (AA, BB) were estimated following the approach described

above. Deletions present in alternative inversion haplotypes were not included in d_{xy} calculations

but were summarized separately using custom scripts.

Inversion age estimates

Absolute sequence divergence between alternative inversion haplotypes was used to calculate the

approximate time of divergence of inverted and non-inverted haplotypes. We used the equation T

$= d_{xy}/2\mu$, where T is the divergence time in generations, μ is the mutation rate per site per

generation, and d_{xy} is a mean d_{xy} calculated based on per SNP values estimated in ANGSD.

Since mutation rates of the spruce bark beetle are unknown, we used a selection of mutation rate

estimates available for some diploid, sexually reproducing insects including *Drosophila*

melanogaster[99]*, Heliconius melpomene*[100], and *Chironomus riparius*[101]. Per generation mutation

rate estimates varied from 2.1 to 11.7×10^{-9}. This approach could only give us rough inversion

age estimates due to the uncertainty of the mutation rate estimates, probable intraspecific

variation in mutation rate[99], and a (likely) substantial influence of gene flux[29].

Mutation load estimation

To estimate mutation load we calculated the ratio of nucleotide diversity at non-synonymous

sites (π_N) vs. synonymous sites (π_S). Mutation load (π_N/π_S) was calculated separately for each

inversion homozygote and for the collinear part of the genome. We computed nucleotide

diversity for each site using SNPGenie[102]. The π_N/π_S ratio was estimated in windows of 200 kb

using an in-house R script. To account for the fact that inversions can greatly suppress

recombination in surrounding parts of the genome[74] the collinear part of the genome was divided

into two groups: (1) a group including all collinear 200 kb windows outside inversions and (2) a

group including all collinear windows outside inversions but excluding windows that came from

contigs with inversions (so called strict filtering). Both collinear datasets were used to test for

overall differences in mutation load between inversions and the collinear part of the genome

(using two-sided t-test). One-sided t-tests were used to test whether minor (less frequent)

homokaryotypes had higher mutation loads than major (more frequent) homokaryotypes.

Homokaryotypes that contained fewer than four 200 kb windows and were present in few

individuals (two thresholds were tested: < 4 and < 10 individuals) were excluded from the

analysis. Additionally, windows with a small number of genes were excluded (two thresholds

were tested: < 5 and < 10 genes).

Haplotype structuring, i.e., the existence of two or more distinct sub-haplotypes among inversion

homozygote haplotypes, can halt fitness degeneration on one or both inversion haplotypes by

carrying partially complementary sets of deleterious recessive alleles[10,103]. To check if any

inversion homozygotes exhibited haplotype structuring we first phased the data using Beagle 5.2

(default settings[104]). Next, in each inverted region and homokaryotype we filtered out all variants

with MAF < 0.1 and used PGDSpider 2.1.1.0[105] to convert variant call formats (VCF) to full

length sequences. Finally, we constructed neighbor-joining trees for alleles within haplotypes

using MEGA7[106].

Phenotype-genotype associations

To test whether inversion polymorphisms were associated with diapause phenotypes we genome

sequenced 18 individuals from a spruce bark beetle diapause study by Schebeck et al.[51] (10

beetles expressing facultative diapause and 8 beetles expressing obligatory diapause). DNA

sequencing was performed using the DNBseq platform (BGI Tech Solutions, Poland) to a mean

coverage of 20×. The data was processed in the same way as described above and combined with

other sequenced individuals before performing PCA in PLINK. To test for differentiation in

inversion haplotype frequencies between diapause phenotypes an exact G test was run using

Genepop 4.1.2[107]. F_{ST} between individuals expressing facultative and obligate diapause was

estimated using VCFtools[89] and summarized in 100 kb overlapping windows (20 kb steps).

To check if spruce bark beetle odorant receptors (ORs) genes were associated with inversions we

examined 73 OR genes recently annotated by Yuvaraj et al.[54]. OR sequences were mapped to the

bark beetle reference genome using minimap2[108], and 71 out of 73 ORs were located in the 186

Mb covered by the 36 contigs we analyzed. Three ORs mapped to more than one contig.

ItypOR9 and ItypOR58 mapped to the end of IpsContig16 and 23, suggesting possible assembly

error and duplication of end-of-contig sequences, which are difficult to assemble. ItypOR59NTE

mapped to three nearby locations on IpsContig6, suggesting either assembly error or recent

duplication. For these three ORs we used one randomly chosen location in downstream analyses.

To test if inversions were enriched in OR genes, we ran permutation tests (10,000 iterations;

permutating inversions locations). To check if alternative inversion arrangements harbored

different numbers of OR genes (e.g. that one arrangement carried a deletion) we compared

sequence coverage within ORs in individual beetles identified as inversion homozygotes.

Genotype-environment association

Genotype-environment association (GEA) analyses were done to test if allele frequency changes

in SNPs were associated with the beetle populations' local environmental. Many different

environmental variables were summarized along two principal components (see below). To

control for confounders due to the overall genetic differentiation, we used Latent Factors Mixed

Models (LFMM[109]) as implemented in the lfmm2() function from the R package LEA[110]. We

used only SNPs with < 20% missing data and MAF >= 0.1, and that occurred in individuals with

< 30% missing data. Because LFMM cannot handle missing data, we imputed missing genotypes

with impute() from LEA. We ran LFMM for (1) all SNPs that passed the filters described above

and (2) for a dataset where each inversion was represented as a single "SNP" inversion genotype.

We used five latent factors (K = 5) in lfmm2(). P-values were calculated using lfmm2.test() from

LEA and false discovery rate (FDR)-corrected using the p.adjust() R function with method =

"fdr".

Each beetle population's local environment was characterized according to climate and land-

cover data. We used all 19 bioclimatic variables from WorldClim version 2.1.[111] with a resolution

of ~1 km^2. These variables are averaged over the years 1970-2000. Proportions of forest,

cropland, and built-up areas were downloaded from https://lcviewer.vito.be/ for 2015 with a

spatial resolution of ~100 m^2[112]. These global land-cover maps are part of the Copernicus Land

Service, derived from PROBA-V satellite observations, and have an accuracy of 80% as

measured by the CEOS land product validation subgroup. We also included the proportion of

land area covered by spruce trees (genus *Picea*) at a resolution of ~1 km^2, as obtained by Brus et

al.[113] using a statistical mapping approach. All environmental variables were reprojected to a

final resolution of ~1 km^2 using the Lambert azimuthal equal area method. We then extracted

mean values for all environmental variables (Figure S13) within a 50 km radius from each

population location using the R package terra[114]. Finally, PCA was used to summarize the multi-

scale environmental variation among populations. The first two PCA components (PC1 and PC2)
explained 25% and 23% of the environmental variation, respectively, and were used as the final
input for the GEA analyses. PC1 represented environmental variation mainly related to latitude,
with northern populations showing higher values indicative of higher temperature seasonality
and lower temperatures during the coldest months. PC2 represented environmental variation
mostly related to temperature and amount of cropland, with higher values representing localities
with higher temperatures during the warmest months and a higher proportion of cropland (and
conversely less forest cover and spruce) (Figure S14).

Data Availability

Acknowledgments

We thank A. Hietala, E. Stengel, K. Zub, Å. Lindelöw, O. Langvall, M. Holmlund, E.
Kristensen, U. Johansson, R. Modlinger, J. Reisenberger, K. Szreder, W. Skowroński L.
Stanecki, and M. Ahlström for help in sampling spruce bark beetles across Europe. T. Mokrzycki
helped with beetle sexing and identification. We thank members of the Genomics and
Experimental Evolution Group at Jagiellonian University for their help in improving this
manuscript. We thank Tomasz Gaczorek for the help in writing scripts and optimizing data
analysis. This work was funded by a Polish National Science Center 2018/30/E/NZ8/00105 grant
to K.N.B, and the Foundation in Memory of Oscar and Lili Lamm to M.N.A.

Author contributions

A.M., P.Z. and K.N.B. conceived the study, performed the main analyses, and wrote the
manuscript. A.B. managed sample shipment, sexed beetles, isolated DNA, and prepared samples
for sequencing. F.S., P.K. and M.N.A. provided an unpublished version of the spruce bark beetle
genome and valuable insights on the species' ecology and sensory biology. M.M., J.M., Z.B.,
M.S., P.K., and C.S. organized sampling, provided beetles for analysis and insights on sampled
populations. B.A. and W.B. performed GEA analysis. Z.N. analyzed diapause data. J.M. phased

the data. M.S. provided diapause samples. W.B. helped in data interpretation and provided

feedback on all manuscript versions. All authors read and approved the final manuscript.

References

1. Hlásny, T. *et al.* Living with bark beetles: impacts , outlook and management options.

From Science to Policy vol. 8 (2019).

2. Vega, F. & Hofstetter, R. Bark Beetles: Biology and Ecology of Native and Invasive

Species. Elsevier (2015).

3. Bentz, B. J., Hansen, E. M., Davenport, M. & Soderberg, D. Complexities in predicting

mountain pine beetle and spruce beetle response to climate change. In *Bark Beetle*

Management, Ecology, and Climate Change. Elsevier (2021).

4. Biedermann, P. H. W. *et al.* Bark Beetle Population Dynamics in the Anthropocene:

Challenges and Solutions. *Trends Ecol. Evol.* **34**, 914–924 (2019).

5. Müller, M., Olsson, P. O., Eklundh, L., Jamali, S. & Ardö, J. Features predisposing forest

to bark beetle outbreaks and their dynamics during drought. *For. Ecol. Manage.* **523**,

120480 (2022).

6. Hlásny, T. *et al.* Bark Beetle Outbreaks in Europe: State of Knowledge and Ways Forward

for Management. *Curr. For. Reports* **7**, 138–165 (2021).

7. Müller, M. How natural disturbance triggers political conflict: Bark beetles and the

meaning of landscape in the Bavarian Forest. *Glob. Environ. Chang.* **21**, 935–946 (2011).

8. Wermelinger, B. & Jakoby, O. Bark beetles. in *Disturbance Ecology* (ed. Thomas

Wohlgemuth, Anke Jentsch, R. S.) 271–293 (2022).

9. Wermelinger, B. Ecology and management of the spruce bark beetle Ips typographus - A

review of recent research. *For. Ecol. Manage.* **202**, 67–82 (2004).

10. Berdan, E. L., Blanckaert, A., Butlin, R. K. & Bank, C. Deleterious mutation

accumulation and the long-term fate of chromosomal inversions. *PLoS Genet.* **17**, (2021).

11. Kirkpatrick, M. & Barton, N. Chromosome inversions, local adaptation and speciation.

Genetics **173**, 419–434 (2006).

12. Wellenreuther, M. & Bernatchez, L. Eco-Evolutionary Genomics of Chromosomal

Inversions. *Trends Ecol. Evol.* **33**, 427–440 (2018).

13. Lohse, K., Clarke, M., Ritchie, M. G. & Etges, W. J. Genome-wide tests for introgression

between cactophilic Drosophila implicate a role of inversions during speciation. *Evolution*

69, 1178–1190 (2015).

14.   Fuller, Z. L., Koury, S. A., Phadnis, N. & Schaeffer, S. W. How chromosomal

rearrangements shape adaptation and speciation: Case studies in Drosophila

pseudoobscura and its sibling species Drosophila persimilis. *Mol. Ecol.* **28**, 1283–1301

(2019).

15.   Lamichhaney, S. *et al.* Structural genomic changes underlie alternative reproductive

strategies in the ruff (*Philomachus pugnax*). *Nat. Genet.* **48**, 84–88 (2015).

16.   Thompson, M. J. & Jiggins, C. D. Supergenes and their role in evolution. *Heredity* **113**, 1–

8 (2014).

17.   Joron, M. *et al.* Chromosomal rearrangements maintain a polymorphic supergene

controlling butterfly mimicry. *Nature* **477**, 203–206 (2011).

18.   Purcell, J., Brelsford, A., Wurm, Y., Perrin, N. & Chapuisat, M. Convergent genetic

architecture underlies social organization in ants. *Curr. Biol.* **24**, 2728–2732 (2014).

19.   Gutiérrez-Valencia, J., Hughes, P. W., Berdan, E. L. & Slotte, T. The Genomic

Architecture and Evolutionary Fates of Supergenes. *Genome Biol. Evol.* **13**, 1–19 (2021).

20.   Lowry, D. B. & Willis, J. H. A widespread chromosomal inversion polymorphism

contributes to a major life-history transition, local adaptation, and reproductive isolation.

PLoS Biol. **8**, e1000500 (2010).

21.   Kirubakaran, T. G. *et al.* Two adjacent inversions maintain genomic differentiation

between migratory and stationary ecotypes of Atlantic cod. *Mol. Ecol.* **25**, 2130–2143

(2016).

22.   Kapun, M., Fabian, D. K., Goudet, J. & Flatt, T. Genomic Evidence for Adaptive

Inversion Clines in Drosophila melanogaster. *Mol. Biol. Evol.* **33**, 1317–1336 (2016).

23.   Mérot, C. *et al.* Locally Adaptive Inversions Modulate Genetic Variation at Different

Geographic Scales in a Seaweed Fly. *Mol. Biol. Evol.* **38**, 3953–3971 (2021).

24.   Porubsky, D. *et al.* Recurrent inversion polymorphisms in humans associate with genetic

instability and genomic disorders. *Cell* **185**, 1986-2005.e26 (2022).

25.   Todesco, M. *et al.* Massive haplotypes underlie ecotypic differentiation in sunflowers.

Nature **584**, 602–607 (2020).

26.   Harringmeyer, O. S. & Hoekstra, H. E. Chromosomal inversion polymorphisms shape the

genomic landscape of deer mice. *Nat. Ecol. Evol.* **6**, 1965–1979 (2022).

27.   Roesti, M., Gilbert, K. J. & Samuk, K. Chromosomal inversions can limit adaptation to

705   new environments. 1–16 (2022).

706   28.   Keightley, P. D. & Otto, S. P. Interference among deleterious mutations favours sex and

707   recombination in finite populations. *Nature* **443**, 89–92 (2006).

708   29.   Charlesworth, B. The effects of inversion polymorphisms on patterns of neutral genetic

709   diversity. *Genetics* **224**, 1–21 (2023).

710   30.   Faria, R., Johannesson, K., Butlin, R. K. & Westram, A. M. Evolving Inversions. *Trends*

711   *Ecol. Evol.* **34**, 239–248 (2019).

712   31.   Connallon, T. & Clark, A. G. Balancing selection in species with separate sexes: Insights

713   from fisher's geometric model. *Genetics* **197**, 991–1006 (2014).

714   32.   Tigano, A. *et al.* Chromosome-Level Assembly of the Atlantic Silverside Genome

715   Reveals Extreme Levels of Sequence Diversity and Structural Genetic Variation. *Genome*

716   *Biol. Evol.* **13**, 1–17 (2021).

717   33.   Faria, R. *et al.* Multiple chromosomal rearrangements in a hybrid zone between Littorina

718   saxatilis ecotypes. *Mol. Ecol.* **28**, 1375–1393 (2019).

719   34.   Jay, P. *et al.* Mutation load at a mimicry supergene sheds new light on the evolution of

720   inversion polymorphisms. *Nat. Genet.* **53**, 288–293 (2021).

721   35.   Wang, J. *et al.* A Y-like social chromosome causes alternative colony organization in fire

722   ants. *Nature* **493**, 664–668 (2013).

723   36.   Jay, P., Tezenas, E., Véber, A. & Giraud, T. Sheltering of deleterious mutations explains

724   the stepwise extension of recombination suppression on sex chromosomes and other

725   supergenes. *PLoS Biology* **20**, e3001698 (2022).

726   37.   Schaal, S. M., Haller, B. C. & Lotterhos, K. E. Inversion invasions: When the genetic

727   basis of local adaptation is concentrated within inversions in the face of gene flow. *Philos.*

728   *Trans. R. Soc. B Biol. Sci.* **377**, (2022).

729   38.   Stauffer, C., Lakatos, F. & Hewitt, G. M. Phylogeography and postglacial colonization

730   routes of Ips typographus L. (Coleoptera, Scolytidae). *Mol. Ecol.* **8**, 763–773 (1999).

731   39.   Sallé, A., Arthofer, W., Lieutier, F., Stauffer, C. & Kerdelhué, C. Phylogeography of a

732   host-specific insect: Genetic structure of Ips typographus in Europe does not reflect past

733   fragmentation of its host. *Biol. J. Linn. Soc.* **90**, 239–246 (2007).

40.   Bertheau, C. *et al.* Divergent evolutionary histories of two sympatric spruce bark beetle
species. *Mol. Ecol.* **22**, 3318–3332 (2013).

41.   Mayer, F. *et al.* Comparative multilocus phylogeography of two Palaearctic spruce bark
beetles: Influence of contrasting ecological strategies on genetic variation. *Mol. Ecol.* **24**,
1292–1310 (2015).

42.   Ellerstrand, S. J. *et al.* Weak population genetic structure in Eurasian spruce bark beetle
over large regional scales in Sweden. *Ecol. Evol.* **12**, 1–12 (2022).

43.   Nilssen, A. C. Long-range aerial dispersal of bark beetles and bark weevils (Coleoptera,
Scolytidae and Curculionidae) in northern Finland. *Ann. Entomol. Fenn.* **50**, 37–42
(1984).

44.   Brelsford, A. *et al.* An Ancient and Eroded Social Supergene Is Widespread across
Formica Ants. *Curr. Biol.* **30**, 304-311.e4 (2020).

45.   Schwander, T., Libbrecht, R. & Keller, L. Supergenes and complex phenotypes. *Curr.*
Biol. **24**, 288–294 (2014).

46.   Roff, D. a. The Evolution of Threshold Traits in Animals. *Q. Rev. Biol.* **71**, 3–35 (1996).

47.   Pruisscher, P., Nylin, S., Gotthard, K. & Wheat, C. W. Genetic variation underlying local
adaptation of diapause induction along a cline in a butterfly. *Mol. Ecol.* **27**, 3613–3626
(2018).

48.   Kozak, G. M. *et al.* Genomic Basis of Circannual Rhythm in the European Corn Borer
Moth. *Curr. Biol.* **29**, 3501-3509.e5 (2019).

49.   Paolucci, S., Salis, L., Vermeulen, C. J., Beukeboom, L. W. & van de Zande, L. QTL
analysis of the photoperiodic response and clinal distribution of period alleles in Nasonia
vitripennis. *Mol. Ecol.* **25**, 4805–4817 (2016).

50.   Denlinger, D. *Insect diapause*. Cambridge University Press (2022).

51.   Schebeck, M., Dobart, N., Ragland, G. J., Schopf, A. & Stauffer, C. Facultative and
obligate diapause phenotypes in populations of the European spruce bark beetle Ips
typographus. *J. Pest Sci. (2004).* **95**, 889–899 (2022).

52.   Hansson, B. S. & Stensmyr, M. C. Evolution of insect olfaction. *Neuron* **72**, 698–711
(2011).

53.   Kandasamy, D., Gershenzon, J., Andersson, M. N. & Hammerbacher, A. Volatile organic
compounds influence the interaction of the Eurasian spruce bark beetle (*Ips typographus*)

with its fungal symbionts. *ISME J.* **13**, 1788–1800 (2019).

54. Yuvaraj, J. K. *et al.* Putative ligand binding sites of two functionally characterized bark

beetle odorant receptors. *BMC Biol.* **19**, 1–21 (2021).

55. Hou, X. Q. *et al.* Functional Evolution of a Bark Beetle Odorant Receptor Clade Detecting

Monoterpenoids of Different Ecological Origins. *Mol. Biol. Evol.* **38**, 4934–4947 (2021).

56. Kandasamy, D. *et al.* Conifer-killing bark beetles locate fungal symbionts by detecting

volatile fungal metabolites of host tree resin monoterpenes. *PLoS Biology* **21**, e3001887

(2023).

57. Tigano, A. & Friesen, V. L. Genomics of local adaptation with gene flow. *Mol. Ecol.* **25**,

2144–2164 (2016).

58. Akopyan, M. *et al.* Comparative linkage mapping uncovers recombination suppression

across massive chromosomal inversions associated with local adaptation in Atlantic

silversides. *Mol. Ecol.* **31**, 3323–3341 (2022).

59. Christmas, M. J. *et al.* Chromosomal inversions associated with environmental adaptation

in honeybees. *Mol. Ecol.* **28**, 1358–1374 (2019).

60. Mérot, C., Llaurens, V., Normandeau, E., Bernatchez, L. & Wellenreuther, M. Balancing

selection via life-history trade-offs maintains an inversion polymorphism in a seaweed fly.

Nat. Commun. **11**, (2020).

61. Navarro, A., Barbadilla, A. & Ruiz, A. Effect of inversion polymorphism on the neutral

nucleotide variability of linked chromosomal regions in drosophila. *Genetics* **155**, 685–

698 (2000).

62. Guerrero, R. F., Rousset, F. & Kirkpatrick, M. Coalescent patterns for chromosomal

inversions in divergent populations. *Philos. Trans. R. Soc. B Biol. Sci.* **367**, 430–438

(2012).

63. Ohta, T. Associative overdominance caused by linked detrimental mutations. *Genet. Res.*

18, 277–286 (1971).

64. Connallon, T. & Olito, C. Natural selection and the distribution of chromosomal inversion

lengths. *Mol. Ecol.* **31**, 3627–3641 (2022).

65. Yang, Y. Y., Lin, F. J. & Chang, H. Y. Comparison of recessive lethal accumulation in

inversion-bearing and inversion-free chromosomes in Drosophila. *Zool. Stud.* **41**, 271–282

(2002).

66.   Huang, K. *et al.* Mutation Load in Sunflower Inversions Is Negatively Correlated with
Inversion Heterozygosity. *Mol. Biol. Evol.* **39**, 1–17 (2022).

67.   McCulloch, G. A. & Waters, J. M. Rapid adaptation in a fast-changing world: Emerging
insights from insect genomics. *Glob. Chang. Biol.* **29**, 943–954 (2023).

68.   Rane, R. V., Rako, L., Kapun, M., Lee, S. F. & Hoffmann, A. A. Genomic evidence for
role of inversion 3RP of *Drosophila melanogaster* in facilitating climate change
adaptation. *Mol. Ecol.* **24**, 2423–2432 (2015).

69.   Kapun, M. & Flatt, T. The adaptive significance of chromosomal inversion
polymorphisms in *Drosophila melanogaster*. *Mol. Ecol.* **28**, 1263–1282 (2019).

70.   Nunez, J. C. B. *et al.* A cosmopolitan inversion drives seasonal adaptation in
overwintering Drosophila. *bioRxiv* 2022.12.09.519676 (2022).

71.   Li, J. *et al.* Joint analysis of demography and selection in population genetics: Where do
we stand and where could we go? *Mol. Ecol.* **21**, 28–44 (2012).

72.   Johri, P. *et al.* The Impact of Purifying and Background Selection on the Inference of
Population History: Problems and Prospects. *Mol. Biol. Evol.* **38**, 2986–3003 (2021).

73.   Li, H. *et al.* Heterozygous inversion breakpoints suppress meiotic crossovers by altering
recombination repair outcomes. *PLoS Genet.* **19**, 1–20 (2023).

74.   Koury, S. A. Predicting recombination suppression outside chromosomal inversions in
Drosophila melanogaster using crossover interference theory. *Heridity* **130**, 196–208
(2023).

75.   Seidl, R., Schelhaas, M., Rammer, W. & Verkerk, P. J. Increasing forest disturbances in
Europe and their impact on carbon storage. *Nat. Clim. Chang.* **4**, 806–810 (2014).

76.   Hlásny, T. *et al.* Devastating outbreak of bark beetles in the Czech Republic: Drivers,
impacts, and management implications. *For. Ecol. Manage.* **490**, 119075 (2021).

77.   Lange, H., Økland, B. & Krokene, P. Thresholds in the life cycle of the spruce bark beetle
under climate change. *Interjournal Complex Syst.* **1648**, 1–10 (2006).

78.   Jönsson, A. M., Harding, S., Bärring, L. & Ravn, H. P. Impact of climate change on the
population dynamics of Ips typographus in southern Sweden. *Agric. For. Meteorol.* **146**,
70–81 (2007).

79.   Wulff, S. & Roberge, C. Inventering av granbarkborreangrepp i Götaland och Svealand.
(2022).

80. Öhrn, P., Berlin, M., Elfstrand, M., Krokene, P. & Jönsson, A. M. Seasonal variation in
Norway spruce response to inoculation with bark beetle-associated bluestain fungi one
year after a severe drought. *For. Ecol. Manage.* **496**, 119443 (2021).

81. Powell, D. *et al.* A highly-contiguous genome assembly of the Eurasian spruce bark
beetle, Ips typographus, provides insight into a major forest pest. *Commun. Biol.* **4**, 1–9
(2021).

82. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods*
9, 357–359 (2012).

83. Depristo, M. A. *et al.* A framework for variation discovery and genotyping using next-
generation DNA sequencing data. *Nat. Genet.* **43**, 491–501 (2011).

84. McKenna, A. *et al.* The Genome Analysis Toolkit: A MapReduce framework for
analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

85. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, 1–4 (2021).

86. Purcell, S. *et al.* PLINK: A tool set for whole-genome association and population-based
linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

87. Skotte, L., Korneliussen, T. S. & Albrechtsen, A. Estimating individual admixture
proportions from next generation sequencing data. *Genetics* **195**, 693–702 (2013).

88. Weir, B. S. & Cockerham, C. C. Estimating F-statistics for the analysis of population
structure. *Evolution* **38**, 1358–1370 (1984).

89. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158
(2011).

90. Korneliussen, T. S., Albrechtsen, A. & Nielsen, R. ANGSD: Analysis of Next Generation
Sequencing Data. *BMC Bioinformatics* **15**, 1–13 (2014).

91. Delmore, K. E. *et al.* Comparative analysis examining patterns of genomic differentiation
across multiple episodes of population divergence in birds. *Evol. Lett.* **2**, 76–87 (2018).

92. Li, H. A statistical framework for SNP calling, mutation discovery, association mapping
and population genetical parameter estimation from sequencing data. *Bioinformatics* **27**,
2987–2993 (2011).

93. Jones, J. C. *et al.* Extreme Differences in Recombination Rate between the Genomes of a
Solitary and a Social Bee. *Mol. Biol. Evol.* **36**, 2277–2291 (2019).

94. McVean, G., Awadalla, P. & Fearnhead, P. A coalescent-based method for detecting and

estimating recombination from gene sequences. *Genetics* **160**, 1231–1241 (2002).

95. Li, H. & Ralph, P. Local PCA shows how the effect of population structure differs along the genome. *Genetics* **211**, 289–304 (2019).

96. Huang, K., Andrew, R. L., Owens, G. L., Ostevik, K. L. & Rieseberg, L. H. Multiple chromosomal inversions contribute to adaptive divergence of a dune sunflower ecotype. *Mol. Ecol.* **29**, 2535–2549 (2020).

97. Graffelman, J. & Weir, B. S. Multi-allelic exact tests for Hardy–Weinberg equilibrium that account for gender. *Mol. Ecol. Resour.* **18**, 461–473 (2018).

98. Graffelman, J. Exploring diallelic genetic markers: The HardyWeinberg package. *J. Stat. Softw.* **64**, 1–23 (2015).

99. Krasovec, M. The spontaneous mutation rate of Drosophila pseudoobscura. *G3 Genes, Genomes, Genet.* **11**, (2021).

100. Keightley, P. D. *et al.* Estimation of the Spontaneous Mutation Rate in Heliconius melpomene. *Mol. Biol. Evol.* **32**, 239–243 (2015).

101. Oppold, A. M. & Pfenninger, M. Direct estimation of the spontaneous mutation rate by short-term mutation accumulation lines in Chironomus riparius. *Evol. Lett.* **1**, 86–92 (2017).

102. Nelson, C. W., Moncla, L. H. & Hughes, A. L. SNPGenie: Estimating evolutionary parameters to detect natural selection using pooled next-generation sequencing data. *Bioinformatics* **31**, 3709–3711 (2015).

103. Charlesworth, B. & Charlesworth, D. Rapid fixation of deleterious alleles can be caused by Muller's ratchet. *Genet. Res.* **70**, 63–73 (1997).

104. Browning, S. R. & Browning, B. L. Rapid and accurate haplotype phasing and missing-data inference for whole-genome association studies by use of localized haplotype clustering. *Am. J. Hum. Genet.* **81**, 1084–1097 (2007).

105. Lischer, H. E. L. & Excoffier, L. PGDSpider: An automated data conversion tool for connecting population genetics and genomics programs. *Bioinformatics* **28**, 298–299 (2012).

106. Kumar, S., Stecher, G. & Tamura, K. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol. Biol. Evol.* **33**, 1870–1874 (2016).

107. Raymond, M. & Rousset, F. An Exact Test for Population Differentiation. *Evolution* **49**,

1280 (1995).

108.   Li, H. Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–
3100 (2018).

109.   Caye, K., Jumentier, B., Lepeule, J. & François, O. LFMM 2: Fast and accurate inference
of gene-environment associations in genome-wide studies. *Mol. Biol. Evol.* **36**, 852–860
(2019).

110.   Gain, C. & François, O. LEA 3: Factor models in population genetics and ecological
genomics with R. *Mol. Ecol. Resour.* **21**, 2738–2748 (2021).

111.   Fick, S. E. & Hijmans, R. J. WorldClim 2: new 1-km spatial resolution climate surfaces
for global land areas. *Int. J. Climatol.* **37**, 4302–4315 (2017).

112.   Buchhorn, M. *et al.* Copernicus global land service: Land cover 100m: Epoch 2015:
Globe. Version V2. 0.2. (2019).

113.   Brus, D. J. *et al.* Statistical mapping of tree species over Europe. *Eur. J. For. Res.* **131**,
145–157 (2012).

114.   Hijmans, R. terra: Spatial Data Analysis. R package version 1.7-18. https://CRAN.R-
project.org/package=terra. (2023).

Tables

910 **Table 1** List of identified chromosomal inversions in the *Ips typographus* genome. ID: inversion

911 name; Contig: contig name; Size: size of the inversions (Mb); Start and End: coordinates of the

912 inversion (Mb); Age: approximate age of the inversion in Myr; Odorant receptors: odorant

913 receptors present within inversion. Note that Inv16.1 and Inv23.1 are parts of the same inversion

914 and Inv16.2 and Inv23.2 are a part of another single inversion.

ID	Contig	Size	Start	End	Age	Odorant receptors
Inv2	IpsContig2	4.04	12.67	16.71	0.5	
Inv3	IpsContig3	0.14	1.11	1.25	1.1	
Inv5	IpsContig5	10.84	0.00	10.84	1.3	ItypOR33, ItypOR41, ItypOR40, ItypOR10, ItypOR47, ItypOR50, ItypOR29, ItypOR43JOI, ItypOR34, ItypOR52NTE, ItypOR4, ItypOR3, ItypOR53, ItypOR2, ItypOR19
Inv6	IpsContig6	0.34	8.93	9.27	1.0	
Inv7.1	IpsContig7	0.67	0.00	0.67	1.7	
Inv7.2	IpsContig7	6.92	0.00	6.92	1.1	ItypOR1, ItypOR17
Inv9	IpsContig9	3.30	1.71	5.01	0.6	
Inv10	IpsContig10	0.08	6.05	6.13	1.8	
Inv12	IpsContig12	0.07	3.63	3.70	1.5	
Inv13	IpsContig13	4.50	0.00	4.50	1.7	ItypOR28, ItypOR23, ItypOR49, ItypOR27
Inv14.1	IpsContig14	2.08	0.00	2.08	1.9	ItypOR36, ItypOR44, ItypOR18JF, ItypOR20NTE
Inv14.2	IpsContig14	0.67	2.08	2.75	1.0	
Inv14.3	IpsContig14	0.76	2.78	3.54	2.1	
Inv14.4	IpsContig14	0.11	3.73	3.84	2.1	
Inv14.5	IpsContig14	0.57	4.23	4.80	2.3	
Inv14.6	IpsContig14	2.48	0.00	2.48	0.6	ItypOR36, ItypOR44, ItypOR18JF, ItypOR20NTE
Inv15	IpsContig15	1.92	0.86	2.78	1.9	
Inv16.1	IpsContig16	4.83	0.00	4.83	1.6	ItypOR58, ItypOR9, ItypOR11, ItypOR31, ItypOR30, ItypOR16, ItypOR35JF
Inv16.2	IpsContig16	4.83	0.00	4.83	0.7	ItypOR58, ItypOR9, ItypOR11, ItypOR31, ItypOR30, ItypOR16,

						ItypOR35JF
Inv17	IpsContig17	0.21	2.48	2.69	2.2	
Inv18	IpsContig18	2.32	0.00	2.32	2.1	
Inv22.1	IpsContig22	0.32	0.00	0.32	2.1	
Inv22.2	IpsContig22	0.23	0.42	0.65	2.1	
Inv22.3	IpsContig22	1.23	0.68	1.91	2.0	ItypOR22CTE
Inv22.4	IpsContig22	0.20	1.92	2.12	2.1	
Inv22.5	IpsContig22	2.24	0.00	2.24	0.6	ItypOR22CTE
Inv23.1	IpsContig23	2.10	0.00	2.10	1.4	ItypOR58, ItypOR9
Inv23.2	IpsContig23	1.84	0.26	2.10	0.6	ItypOR58, ItypOR9
Inv26	IpsContig26	0.10	0.12	0.22	2.6	

Figures

Figure 1 Identification of chromosomal inversions in *Ips typographus*. Each row of figure panels

shows the results of per contig PCA, per contig linkage disequilibrium analysis, and genetic

differentiation (F_{ST}) analysis for a selected contig. Panels a-c show results for a contig with no

inversions (IpsContig8) and little differentiation between southern and northern populations.

Panels d-o show different contigs with increasingly complex inversion patterns: a single

inversion (d-f, Inv15); a single inversion with a double-crossover signal (g-i, Inv18); multiple

adjacent inversions with one inversion overlapping with the first two inversions on the contig (j-

l, Inv14.1, Inv14.2, Inv14.3, Inv14.4, Inv14.5, and Inv.14.6); multiple adjacent inversions with

one large inversion overlapping with several smaller ones (m-o, Inv22.1, Inv22.2, Inv22.3,

Inv22.4, and Inv22.5). The large overlapping inversion is visible as a yellow background in the

LD plot (n). The PCA grouping shown in (m) corresponds to genotypes of the largest inversion

on IpsContig22 (Inv22.3), including genotypes that include haplotypes produced with the double

crossover event. The last panel (p) shows the 36 largest contigs of the spruce bark beetle genome

with inversions indicated in red. Dots in the PCA plots represent individual beetles and are

colored according to the heterozygosity of the individual (darker color represents low

heterozygosity). Both axis on LD plots (b,e,h,k,n) represent conting's positions in megabases

(Mb); low levels of linkage are shown in blue and higher levels in yellow to dark red. F_{ST} values

in panels f,i,l,o show genetic differentiation between inversion haplotypes along the contig in

question.

Figure 2 Genomic structure and differentiation in *Ips typographus*. (a) Whole-genome PCA,
where colors correspond to genetic clustering of beetle individuals revealed by NGSadmix
analysis. (b,c) Geographical distribution and genetic differentiation of the 18 beetle populations
analyzed. Blue dots and bars: northern populations (N); red dots and bars: southern populations
(S); violet dots: Polish populations (P). (d) Genome-wide genetic differentiation (F_{ST}) calculated
between northern and southern populations. Vertical lines separate different contigs shown in
grey and black.
946

947
948

949 **Figure 3** Correlation between inversion haplotype frequency and population origin (latitude) for
950 chromosomal inversions detected in European *Ips typographus* populations. Significant
951 correlations are indicated with asterisks (*: p < 0.05; **: p < 0.01; ***: p < 0.001). Inversions
952 harboring odorant receptors are indicated in green.

953
954

 Figure 4 Genotype-environment associations across 36 contigs in the *Ips typographus* genome.
Different colors represent different contigs. The upper three panels show results for all SNPs and
the lower three panels show results using data where each inversion was coded as a single locus
(shown in a blue circle). MAF: Minor Allele Frequency. PC1, PC2, and PC3 represent the first
three of PCA used to summarize the environmental variation among populations.

Figure 5 Mutation load analysis in *Ips typographus*. (a)
The ratio of nonsynonymous to synonymous nucleotide diversity π_N/π_S computed for 200 kb
windows along collinear parts of the genome (COL) and different inversion haplotypes (blue:
major haplotype; yellow: minor haplotype; green: haplotype produced by double crossover
events between minor and major haplotypes). n = 4 to 55 windows per inversion haplotype and n
= 531 for the collinear genome. Inversion haplotypes that were present in less than four
individuals, did not include four or more 200 kb windows or include less than five genes were
not included in the analysis. For better visibility, π_N/π_S outliers above 0.5 are not shown (a total
of 58 π_N/π_S values, 45 of them belonging to COL). The lower and upper box hinges correspond
to the first and third quartiles, whiskers show 1.5 * the inter-quartile range. (b) Distribution of
π_N/π_S values per 200 kb window (c) Correlation between mean π_N/π_S for all inversion haplotypes
and their frequency across spruce bark beetle populations.
974

975