Premature endocycling of *Drosophila* follicle cells causes pleiotropic defects in

oogenesis

Hunter C. Herriage[1] and Brian R. Calvi[1,2]

1: Department of Biology, Indiana University, Bloomington, IN 47405

2: Corresponding author, bcalvi@indiana.edu

Keywords: endocycling, border cells, polyploidy, oogenesis, epithelium, *Drosophila*

Summary Statement

A premature switch to polyploid endocycles in *Drosophila* ovarian follicle cells caused

pleiotropic defects in oogenesis and compromised female fertility, revealing new ways in which

unscheduled endocycles cause developmental defects.

Abstract

Endocycling cells grow and repeatedly duplicate their genome without dividing. Cells switch

from mitotic cycles to endocycles in response to developmental signals during the growth of

specific tissues in a wide range of organisms. The purpose of switching to endocycles,

however, remains unclear in many tissues. Additionally, cells can switch to endocycles in

response to conditional signals, which can have beneficial or pathological effects on tissues.

However, the impact of these unscheduled endocycles on development is underexplored.

Here, we use *Drosophila* ovarian somatic follicle cells as a model to examine the impact of

unscheduled endocycles on tissue growth and function. Follicle cells normally switch to

endocycles at mid-oogenesis. Inducing follicle cells to prematurely switch to endocycles

resulted in lethality of the resulting embryos. Analysis of ovaries with premature follicle cell

endocycles revealed aberrant follicular epithelial structure and pleiotropic defects in oocyte

growth, developmental gene amplification, and the migration of a special set of follicle cells

known as border cells. Overall, these findings reveal how unscheduled endocycles can disrupt

tissue growth and function to cause aberrant development.

Introduction

Many tissues grow via a variant cell cycle known as the endocycle. Endocycles are characterized by alternating G and S phases without mitotic division, resulting in the growth of large polyploid cells (Calvi, 2013). A regulated switch from mitotic cycles to endocycles is part of normal developmental growth programs of specific tissues across *Eukarya* (Fox and Duronio, 2013; Ovrebo and Edgar, 2018; Shu et al., 2018). In contrast, cells can also switch to unscheduled endocycles in response to stress and other conditional signals (Bailey et al., 2021; Hassel et al., 2014; Qi and Calvi, 2016; Rotelli et al., 2019). These induced endocycling cells can be beneficial for wound healing and regeneration, but they can also have pathological effects, including the promotion of cancer (Bailey et al., 2021; Herriage et al., 2023; Pienta et al., 2022; Shu et al., 2018). Much remains unknown, however, about how unscheduled endocycling affects tissue structure and function.

In this study we use the somatic follicle cells of the *Drosophila* ovary as a model to evaluate the effects of unscheduled endocycles on tissue function. A *Drosophila* ovary consists of ~16 ovarioles, each composed of one germarium stem cell compartment followed by an "assembly line" of progressively developing egg chambers in 14 stages of oogenesis (Fig. 1A) (Duhart et al., 2017; King, 1970). These developing egg chambers contain the germline (15 endocycling nurse cells and one oocyte) ensheathed by an epithelium of somatic follicle cells (Fig. 1A) (Jia et al., 2015; King, 1970). Follicle cells originate from a population of two to four somatic stem cells in the germarium and proliferate mitotically over the course of three days until stage 6 of oogenesis (Fadiga and Nystul, 2019; Margolis and Spradling, 1995; Sun and Deng, 2007). In response to Notch signaling at stages 6 / 7, follicle cells switch to the endocycle and subsequently complete three genome duplications over the next ~24 hours (Fig. 1A) (Calvi et al., 1998; Deng et al., 2001; Lopez-Schier and St Johnston, 2001; Spradling, 1993). After achieving a DNA copy number (C) of 16C, follicle cells arrest endocycles at different times (asynchronously) during late stage 9. Follicle cells then synchronously initiate a specialized DNA rereplication program in stage 10B that selectively amplifies the DNA copy number of eggshell (chorion) protein genes, a process known as chorion gene amplification (Fig. 1A) (Calvi et al., 1998; Spradling and Mahowald, 1980). Disruption of this process causes thin eggshells and embryonic lethality (Calvi, 2006; Spradling and Mahowald, 1980). Follicle cells also support egg growth by mediating transport of yolk proteins into the oocyte and

participate in patterning of the egg and embryo (Brennan et al., 1982; Duhart et al., 2017;

Merkle et al., 2020; Raikhel and Dhadialla, 1992). While it is known that follicle cells have

these important roles in oogenesis, it remains unclear what biological function is served by the

endocycle switch occurring specifically at stage 6. In this study, we find that unscheduled

endocycles can have severe negative consequences, with broader impacts on understanding

how they may contribute to human disease.

Results and Discussion

<u>Induction of premature endocycles in follicle cells</u>

To address whether a switch to follicle cell endocycles before stage 6 would affect

oogenesis, we induced premature endocycling by expressing the protein Roughex (Rux), a

repressor of Cyclin A / CDK1 activity and mitotic entry (Avedisov et al., 2000; Sprenger et al.,

1997). A *tj-GAL4* transgene induced expression of a *UAS-Rux* transgene in all ovarian somatic

cells*,* and a temperature-sensitive GAL80^{ts} repressor of GAL4 permitted control over timing of

expression (Brand and Perrimon, 1993; McGuire et al., 2004; Weaver et al., 2020). The *tj-*

GAL4 tub-GAL80^{ts} / UAS-Rux experimental and *tj-GAL4 tub-GAL80^{ts} / +* control animals were

raised at 18°C to repress GAL4 activity. Adult flies were then shifted to 29°C to induce *UAS-*

Rux expression and ovaries were examined one and two days thereafter (Fig. 1A). Within one

day after *UAS-Rux* expression, stage 1-6 follicle cells ceased labeling for the mitotic marker

phosphorylated Histone H3 (pH3) but continued to incorporate the nucleotide analog 5-ethynyl

2′-deoxyuridine (EdU), suggesting that they had switched to G / S endocycles (Fig. S1A-D′).

We next asked what effect premature endocycle induction would have on follicle cells as

they matured to later stages of oogenesis (Fig. 1A). Two days after a shift to 29°C, stage 10B

egg chambers from control females had a normal ~641 follicle cells, and measurement of DAPI

fluorescence indicated that they had the normal DNA content of ~16C (Fig. 1A-B′, D, E) (Calvi

et al., 1998; Sun and Deng, 2007). In contrast, induction of premature endocycles two days

earlier resulted in stage 10B egg chambers with four-fold fewer follicle cells (~179) and nearly

a four-fold increase in average DNA content (mean ploidy ~56 C) (Fig. 1C-E). Immunolabeling

for E-cadherin also indicated that these stage 10B follicle cells were larger than normal,

consistent with the known scaling of cell size with DNA ploidy (Fig. 1B′, C′) (Cadart and Heald,

2022; Tepass et al., 1996; Uemura et al., 1996). These results suggest that Rux-expressing

follicle cells had skipped two mitotic divisions and instead underwent two extra endocycle

genome doublings. These results are consistent with the known rate of oogenesis at 29°C

(Gandara and Drummond-Barbosa, 2022). These stage 10B egg chambers would have been

in approximately stage 4 / 5 two days earlier at the time of temperature shift (Gandara and

Drummond-Barbosa, 2022). Normally, follicle cells in a stage 4 / 5 egg chamber undergo an

average of two mitotic divisions over one day as it matures to stage 6 / 7, after which follicle

cells endocycle over the next day from stage 7 to stage 10A (Calvi et al., 1998; Margolis and

Spradling, 1995). Given that there were two extra genome doublings instead of the normal two

mitotic divisions during the ~24 hours from stage 4 / 5 to stage 6 / 7, these data also suggest

that the duration of each induced endocycle is approximately equal to that of the mitotic cycles

that they supplanted (Margolis and Spradling, 1995). Moreover, the increased DNA content of

these follicle cells at stage 10 suggests that starting endocycles early can override the

undefined cell-autonomous mechanisms that arrest endocycles to limit follicle cell ploidy to

16C.

<u>Premature endocycling has deleterious effects on oogenesis and female fertility</u>

We next examined whether premature follicle cell endocycles affected the morphology

of late-stage egg chambers. Stage 10B egg chambers from Rux-expressing females appeared

similar to those from controls (Fig 1B-C'). The main body follicle cells underwent normal

migration posteriorly during stage 9 to over the growing oocyte (Figs. 1B-C', 2A-D).

Measurements indicated that one day after temperature shift, stage 10B egg chambers from

Rux-expressing females had oocytes of a similar size to controls (Fig. 2A- E). These egg

chambers would have been in approximately stage 6 / 7 one day earlier at the time of

temperature shift, a stage when follicle cells normally enter endocycles, and, therefore, would

have begun endocycling no more than a few hours early. Conversely, two days after

temperature shift, the Rux-expressing stage 10B egg chambers that had begun endocycles in

stage 4 / 5 had oocytes that were smaller than controls (Fig. 2C-E). Correspondingly, the

length of the nurse cell compartment in these stage 10B egg chambers was larger than

controls after two days of Rux expression (Fig. 2F). These results suggest that the two extra

follicle cell endocycles had non-autonomous effects on germline growth, which may include the

function of follicle cells to mediate oocyte yolk deposition (Brennan et al., 1982; Horne-

Badovinac and Bilder, 2005; Isasti-Sanchez et al., 2021; Raikhel and Dhadialla, 1992).

Another important function of follicle cells is to synthesize the eggshell late in

oogenesis, a process that is supported by chorion gene amplification (Fig. 1A) (Calvi et al.,

1998; Spradling and Mahowald, 1980). This selective gene amplification can be detected as

distinct foci of EdU incorporation beginning in stage 10B (Calvi et al., 1998). EdU labeling of *tj-*

GAL4 GAL80^{ts} / + control ovaries at 29°C showed that all follicle cells over the oocyte had

nuclear foci of EdU at the onset of stage 10B, indicating that the onset of amplification was

synchronized among follicle cells, as previously described (Fig. 2G, I-I′, K) (Calvi et al., 1998).

Follicle cells that had been expressing Rux for 24 hours also had the normal transition to

synchronous amplification in stage 10B (Fig. 2H). In contrast, induction of premature

endocycles two days earlier resulted in some stage 10B egg chambers with a mixture of follicle

cells that had either the normal amplification foci or pan-nuclear EdU incorporation indicative of

continued genomic replication (Fig. 2J-K). These results suggest that a premature start to

endocycles in earlier oogenesis disrupts the normal cessation of endocycles and the

synchronous initiation of amplification in later oogenesis. Further analyses of these induced

endocycles may provide insight into the mechanisms that coordinate the transition from

endocycles to amplification during stage 10, which may be similar to the mechanisms that

govern endocycle arrest and determine final DNA content in other polyploid cell types.

Defects in chorion gene amplification are known to result in thin eggshells and reduced

embryo viability (Calvi, 2006). We therefore examined whether premature endocycles impaired

eggshell synthesis. Females expressing *UAS-Rux* began laying eggs with thin shells one day

after induction, a phenotype that greatly increased in frequency after two days (Fig. 2L-P, Fig.

S2A, B). Expression of Rux in females also resulted in reduced viability of their embryos, as

evidenced by a reduction in egg hatching beginning on day one, which was further reduced on

days two and three (Fig. 2Q). Three or more days of Rux expression resulted in ovarioles with

degenerating stage 8-9 egg chambers, diagnostic of an ovarian stress response known as the

vitellogenic checkpoint (Fig. S2 C, D) (Peterson et al., 2015; Serizier and McCall, 2017). This

result suggests that very early entry into endocycles in stage 1 / 2 triggers a general stress

response three days later in stage 8. All together, these data indicate that induction of

premature follicle cell endocycles has deleterious effects on egg quality and female fertility.

<u>Premature endocycles cause structural defects in the follicular epithelium</u>

After Rux expression, there was a one to several day delay in later oogenesis defects

and reduced embryo viability. This delay suggested that these defects were the manifestation

of premature endocycle entry at earlier stages. To examine the structure of the follicular

epithelium in early stage egg chambers, we immunolabeled for the cell-cell junction protein E-

cadherin (Godt and Tepass, 1998; Oda et al., 1997; Tepass et al., 1996; Uemura et al., 1996).

Although EdU and pH3 labeling had indicated that follicle cells switch to endocycles within one

day of Rux expression, this duration of induced endocycling did not cause major defects in

epithelial structure (Figs. S1A-B′, 3A-B″). These follicle cells had a uniform, cuboidal structure

and E-cadherin labeling on their apical surface that is adjacent to the germline (Fig. 3A-B″)

(Godt and Tepass, 1998; Gonzalez-Reyes and St Johnston, 1998). In contrast, after two days

of Rux expression, follicle cells were all larger with wide variations in their size and shape

within an egg chamber (Fig. 3C-D″). Some follicle cells had highly irregular shapes, with

different parts of the cell having unequal thicknesses in the apical / basal axis, and many with

highly irregular E-cadherin labeling (Fig. 3D-D″). In confocal sections through the middle of egg

chambers, the interface between the follicular epithelium and germline appeared wavy instead

of the normal smooth circular interface, suggesting global deformations of tissue morphology

(Fig. 3C′, D′). Labeling for the membrane-associated protein Discs large 1 (Dlg1) also revealed

irregular cell sizes and shapes, but Dlg1 had an overall normal basolateral localization,

suggesting that there was not a severe disruption of cell polarity (Fig. S3A-D″) (Khoury and

Bilder, 2020; Woods et al., 1996). All follicle cells in these aberrant egg chambers had an

increased DNA content consistent with having proceeded through premature endocycles (Fig.

3C-E).

These disruptions of epithelial structure in early stages may contribute to the observed

defects in later stages of oogenesis over subsequent days. This includes the reduction in

oocyte growth in stage 10 egg chambers, which is mediated by a paracelluar transport of yolk

protein across the follicular epithelium beginning in stage 8 (Isasti-Sanchez et al., 2021;

Riechmann, 2021). This regulated permeability of the epithelium is known as patency, and is

mediated by remodeling of cell-cell junctions through a mechanism that includes E-cadherin

(Isasti-Sanchez et al., 2021). Thus, the aberrant E-cadherin labeling and cell and epithelial

morphology we observe in induced endocycling cells may disturb the remodeling of cell-cell

contacts that is required for patency and oocyte growth. More broadly, these results suggest

that disrupted epithelial structure and permeability in other tissues may be another way for

unscheduled endocycles to contribute to disease.

<u>Premature endocycling inhibits border cell migration</u>

Unlike the main body follicle cells that migrate on the surface of the egg chamber,

another group of follicle cells delaminate from the anterior epithelium and migrate posteriorly

between the nurse cells during stage 9 (Mishra et al., 2019; Montell et al., 1992; Montell et al.,

2012; Spradling, 1993). This collective migration event is initiated in early stage 9 when two

specialized follicle cells, known as the polar cells, recruit 4-6 neighboring follicle cells to

become border cells via JAK-STAT signaling (Montell et al., 1992; Silver and Montell, 2001).

After arriving at their final destination at the anterior of the growing oocyte in stage 10A, this

migrating border cell cluster forms the sperm entry port known as the micropyle (Bishop and

King, 1984; King, 1970). Because these border cells are a powerful model for mechanisms of

collective cell migration during metastasis, we wished to investigate the effect of premature

endocycling on their migration.

We identified border cells based on their elevated E-cadherin levels in control and Rux-

expressing ovaries two days after induction at 29°C and measured the distance they had

migrated relative to main body follicle cells in the same egg chamber during stage 9-10A

(Niewiadomska et al., 1999; Oda et al., 1997). In controls, the migration of the border cell

cluster between the nurse cells stayed on-pace with the anterior of the main body follicle cells

that are migrating on the surface of the egg chamber during stage 9, with both populations of

follicle cells reaching the oocyte at the onset of stage 10A (Fig. 4A, A', C). In contrast, Rux

expression resulted in border cells lagging behind the main body follicle cells to varying extents

(Fig. 4B-C). In some egg chambers, border cells had failed to even delaminate from the

epithelium at the anterior pole of the egg chamber by stage 10 (Fig. 4C). Having abnormal

numbers of polar cells or border cells can inhibit border cell cluster delamination and migration

(Silver et al., 2005; Silver and Montell, 2001). Labeling for the polar cell marker protein

Fasciclin III (Fas III) showed that after two days of Rux expression egg chambers had the

normal two polar cells, including those in which border cells migrated slowly or failed to

delaminate from the epithelium (Fig. 4D-E''') (Ruohola et al., 1991). Co-labeling against E-

cadherin indicated that these migrating clusters had normal structure and cell polarity (Fig. 4D-

E'''). While some Rux-expressing egg chambers had a reduced number of cells per cluster,

even clusters composed of the normal number of cells were delayed in their migration (Fig. 4B,

B′, E-E‴). Our observation of impaired collective border cell migration contrasts with recent

evidence that polyploidy enhances migration of cancer cells *in vitro,* and suggestions that it

promotes collective cell migration during metastasis (Godinho et al., 2014; Mallin et al., 2023;

Wang et al., 2019). Our results raise the possibility that the effects of unscheduled endocycles

on cell migration are dependent on cell type and tissue context (Lamb et al., 2021; Penfield

and Montell, 2023). In addition, oncogenic mutations may synergize with the polyploid state to

give rise to emergent migratory properties (Almeida Machado Costa et al., 2022).

<u>Conclusions</u>

Overall, our findings show that unscheduled endocycles can have severe pleiotropic

effects on epithelial structure and function. Further characterization of these effects will lead to

a greater understanding of the different ways in which unscheduled endocycling can impact

development and contribute to disease.

Materials and Methods

Drosophila genetics: Crosses of *tj-GAL4 tub-GAL80^ts^ / CyO* to *w^1118^* (control) or *UAS-Rux / SM6a* (experimental) were conducted, and their offspring were reared, at 18°C during larval and early adult development. The resulting adult female offspring were fed wet yeast with males for 2 days at 18°C before being transferred to new wet yeast and shifted to 29°C for 1, 2, or 3 days (as noted in the text). The females and males were transferred to new wet yeast every 2 days during the experiments. The *tj-GAL4, tub-GAL80^ts^ / CyO* strain was a generous gift from L. Weaver. The *UAS-Rux / SM6a* strain was obtained from the Bloomington Drosophila Stock Center (BDSC #9166).

Immunofluorescence microscopy: Ovaries were fixed and labeled as previously described (Schwed et al., 2002). For E-cadherin labeling, fixation buffers were supplemented with 1 mM $CaCl_2$. The following antibodies and dilutions were used: rat anti-shg (fly E-cadherin), 1:20 (DSHB DCAD2); rabbit anti-phosphorylated histone H3 (pH3), 1:1000 (Millipore 06-570); mouse anti-Dlg1, 1:200 (DSHB 4F3); mouse anti-FasIII, 1:50 (DSHB 7G10). EdU incorporation *in vitro* was for 1 hour as previously described (Calvi and Lilly, 2004). Asynchronous chorion gene amplification was defined by stage 10B egg chambers with both punctate and pan-nuclear EdU labeling of follicle cell nuclei. Egg chambers with more than one pan-nuclear EdU labeled nucleus were scored as "asynchronous." Fluorescence micrographs were taken on a Leica SP8 confocal. DIC images of stage 10B egg chambers were taken using a Leica DM5500 B. Brightfield images of laid eggs were taken using a Zeiss SteREO Discovery.V12. Nuclear DNA content was quantified using ImageJ/FIJI by measuring mean DAPI fluorescence per pixel in a nucleus, subtracting local mean background DAPI fluorescence per pixel, and then multiplying the difference by the number of pixels per nucleus to obtain the corrected total DAPI fluorescence intensity. The number of follicle cells, oocyte area, nurse cell compartment length, and border cell migration were quantified using ImageJ/FIJI.

Eggshell scoring and egg hatch rate: Eggs from well-fed and mated females were collected on grape plates with a thin layer of yeast paste for 2-4 hours for each respective day of 29°C transgene induction. After counting the number of eggs with thin eggshells, eggs were lined up and incubated at 18°C to ensure no GAL4-driven transgene expression affected their viability. They were then scored for hatching over four subsequent days.

<u>Statistical Analysis:</u> Data graphing and statistical analyses were carried out in GraphPad

Prism. Data were tested for normality and homoscedasticity to determine correct analyses.

Sample size (n) is defined as the total of independent observations / measurements used for

the chosen analysis across multiple biological replicates. All averages shown are mean ± s.d.,

except for Figure 2K, which is mean ± s.e.m. Two-tailed Mann-Whitney tests were used to test

for significance in Figs. 1D, 2E, and 3E. Fig. 2E was corrected for multiple comparisons. A two-

tailed unpaired Welch's t-test was used to test for significance in Fig. 1E. A Mixed Effects Two-

Way ANOVA was used to test for significance in Fig. 2F. A two-tailed unpaired t-test was used

to test for significance in Fig. 2K.

Acknowledgements

We thank S. Brown and Y.T. Huang for experimental assistance. We are appreciative of the

thoughtful feedback on the research and manuscript from S. Brown, L. Hesting, Y.T. Huang, P.

Rangarajan, and C. Walczak. We thank A. Kun and J. Powers of the IU Light Microscopy

Imaging Center, supported in part by the Office of the Vice Provost of Research. Special

thanks to L. Weaver for her generous gift of the *tj-GAL4 tub-GAL80^{ts} / CyO* strain.

Competing Interests

The authors declare no competing interests.

Author Contributions

H.C.H. performed experiments and statistical analyses. Both authors wrote and edited the

manuscript. B.R.C. secured funding and provided mentorship and supervision.

Funding

This work was supported by National Institutes of Health Grant NIH R01GM113107 to B.R.C.

References

Almeida Machado Costa, C., Wang, X. F., Ellsworth, C. and Deng, W. M. (2022). Polyploidy in development and tumor models in Drosophila. *Semin Cancer Biol* **81**, 106-118.

Avedisov, S. N., Krasnoselskaya, I., Mortin, M. and Thomas, B. J. (2000). Roughex mediates G(1) arrest through a physical association with cyclin A. *Mol Cell Biol* **20**, 8220-8229.

Bailey, E. C., Kobielski, S., Park, J. and Losick, V. P. (2021). Polyploidy in Tissue Repair and Regeneration. *Cold Spring Harb Perspect Biol* **13**.

Bishop, D. L. and King, R. C. (1984). An ultrastructural study of ovarian development in the otu-7 mutant of Drosophila melanogaster. *J. Cell Sci.* **67**, 87-119.

Brand, A. and Perrimon, N. (1993). Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development* **118**, 401-415.

Brennan, M. D., Weiner, A. J., Goralski, T. J. and Mahowald, A. P. (1982). The follicle cells are a major site of vitellogenin synthesis in Drosophila melanogaster. *Dev Biol* **89**, 225-236.

Cadart, C. and Heald, R. (2022). Scaling of biosynthesis and metabolism with cell size. *Molecular Biology of the Cell* **33**, pe5.

Calvi, B. R. (2006). Developmental DNA amplification. In *DNA replication and human disease* (ed. M. L. DePamphilis), pp. 233-255. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press.

Calvi, B. R. (2013). Making big cells: one size does not fit all. *Proc Natl Acad Sci U S A* **110**, 9621-9622.

Calvi, B. R., Lilly, M. A. and Spradling, A. C. (1998). Cell cycle control of chorion gene amplification. *Genes Dev* **12**, 734-744.

Deng, W. M., Althauser, C. and Ruohola-Baker, H. (2001). Notch-Delta signaling induces a transition from mitotic cell cycle to endocycle in Drosophila follicle cells. *Development* **128**, 4737-4746.

Duhart, J. C., Parsons, T. T. and Raftery, L. A. (2017). The repertoire of epithelial morphogenesis on display: Progressive elaboration of Drosophila egg structure. *Mech Dev* **148**, 18-39.

Fadiga, J. and Nystul, T. G. (2019). The follicle epithelium in the Drosophila ovary is maintained by a small number of stem cells. *Elife* **8**.

Fox, D. T. and Duronio, R. J. (2013). Endoreplication and polyploidy: insights into development and disease. *Development* **140**, 3-12.

Gandara, A. C. P. and Drummond-Barbosa, D. (2022). Warm and cold temperatures have distinct germline stem cell lineage effects during Drosophila oogenesis. *Development* **149**.

Godinho, S. A., Picone, R., Burute, M., Dagher, R., Su, Y., Leung, C. T., Polyak, K., Brugge, J. S., Thery, M. and Pellman, D. (2014). Oncogene-like induction of cellular invasion from centrosome amplification. *Nature* **510**, 167-171.

Godt, D. and Tepass, U. (1998). Drosophila oocyte localization is mediated by differential cadherin-based adhesion. *Nature* **395**, 387-391.

Gonzalez-Reyes, A. and St Johnston, D. (1998). The Drosophila AP axis is polarised by the cadherin-mediated positioning of the oocyte. *Development* **125**, 3635-3644.

Hassel, C., Zhang, B., Dixon, M. and Calvi, B. R. (2014). Induction of endocycles represses
apoptosis independently of differentiation and predisposes cells to genome instability.
Development **141**, 112-123.
Herriage, H. C., Huang, Y. T. and Calvi, B. R. (2023). The antagonistic relationship between
apoptosis and polyploidy in development and cancer. *Semin Cell Dev Biol*.
Horne-Badovinac, S. and Bilder, D. (2005). Mass transit: epithelial morphogenesis in the
Drosophila egg chamber. *Dev Dyn* **232**, 559-574.
Isasti-Sanchez, J., Munz-Zeise, F., Lancino, M. and Luschnig, S. (2021). Transient opening
of tricellular vertices controls paracellular transport through the follicle epithelium during
Drosophila oogenesis. *Dev Cell* **56**, 1083-1099 e1085.
Jia, D., Huang, Y. C. and Deng, W. M. (2015). Analysis of Cell Cycle Switches in Drosophila
Oogenesis. *Methods Mol Biol* **1328**, 207-216.
Khoury, M. J. and Bilder, D. (2020). Distinct activities of Scrib module proteins organize
epithelial polarity. *Proc Natl Acad Sci U S A* **117**, 11531-11540.
King, R. C. (1970). *Ovarian Development in Drosophila melanogaster*. New York: Academic
Press.
Lamb, M. C., Kaluarachchi, C. P., Lansakara, T. I., Mellentine, S. Q., Lan, Y., Tivanski, A. V.
and Tootle, T. L. (2021). Fascin limits Myosin activity within Drosophila border cells to
control substrate stiffness and promote migration. *Elife* **10**.
Lopez-Schier, H. and St Johnston, D. (2001). Delta signaling from the germ line controls the
proliferation and differentiation of the somatic follicle cells during Drosophila oogenesis.
Genes Dev **15**, 1393-1405.
Mallin, M. M., Kim, N., Choudhury, M. I., Lee, S. J., An, S. S., Sun, S. X., Konstantopoulos,
K., Pienta, K. J. and Amend, S. R. (2023). Cells in the polyaneuploid cancer cell
(PACC) state have increased metastatic potential. *Clin Exp Metastasis* **40**, 321-338.
Margolis, J. and Spradling, A. (1995). Identification and behavior of epithelial stem cells in
the Drosophila ovary. *Development* **121**, 3797-3807.
McGuire, S. E., Mao, Z. and Davis, R. L. (2004). Spatiotemporal gene expression targeting
with the TARGET and gene-switch systems in Drosophila. *Sci STKE* **2004**, pl6.
Merkle, J. A., Wittes, J. and Schupbach, T. (2020). Signaling between somatic follicle cells
and the germline patterns the egg and embryo of Drosophila. *Curr Top Dev Biol* **140**,
55-86.
Mishra, A. K., Campanale, J. P., Mondo, J. A. and Montell, D. J. (2019). Cell interactions in
collective cell migration. *Development* **146**.
Montell, D. J., Rorth, P. and Spradling, A. C. (1992). slow border cells, a locus required for a
developmentally regulated cell migration during oogenesis, encodes Drosophila C/EBP.
Cell **71**, 51-62.
Montell, D. J., Yoon, W. H. and Starz-Gaiano, M. (2012). Group choreography: mechanisms
orchestrating the collective movement of border cells. *Nat Rev Mol Cell Biol* **13**, 631-
645.
Niewiadomska, P., Godt, D. and Tepass, U. (1999). DE-Cadherin is required for intercellular
motility during Drosophila oogenesis. *J Cell Biol* **144**, 533-547.
Oda, H., Uemura, T. and Takeichi, M. (1997). Phenotypic analysis of null mutants for DE-
cadherin and Armadillo in Drosophila ovaries reveals distinct aspects of their functions
in cell adhesion and cytoskeletal organization. *Genes Cells* **2**, 29-40.
Ovrebo, J. I. and Edgar, B. A. (2018). Polyploidy in tissue homeostasis and regeneration.
Development **145**.

Penfield, L. and Montell, D. J. (2023). Nuclear lamin facilitates collective border cell invasion into confined spaces in vivo. *J Cell Biol* **222**.

Peterson, J. S., Timmons, A. K., Mondragon, A. A. and McCall, K. (2015). The End of the Beginning: Cell Death in the Germline. *Curr Top Dev Biol* **114**, 93-119.

Pienta, K. J., Hammarlund, E. U., Austin, R. H., Axelrod, R., Brown, J. S. and Amend, S. R. (2022). Cancer cells employ an evolutionarily conserved polyploidization program to resist therapy. *Semin Cancer Biol* **81**, 145-159.

Qi, S. and Calvi, B. R. (2016). Different cell cycle modifications repress apoptosis at different steps independent of developmental signaling in Drosophila. *Mol Biol Cell* **27**, 1885-1897.

Raikhel, A. S. and Dhadialla, T. S. (1992). Accumulation of yolk proteins in insect oocytes. *Annu Rev Entomol* **37**, 217-251.

Riechmann, V. (2021). Paracellular transport: Opening the gates for growth. *Dev Cell* **56**, 1075-1077.

Rotelli, M. D., Policastro, R. A., Bolling, A. M., Killion, A. W., Weinberg, A. J., Dixon, M. J., Zentner, G. E., Walczak, C. E., Lilly, M. A. and Calvi, B. R. (2019). A Cyclin A-Myb-MuvB-Aurora B network regulates the choice between mitotic cycles and polyploid endoreplication cycles. *PLoS Genet* **15**, e1008253.

Ruohola, H., Bremer, K., Baker, D., Swedlow, J., Jan, L. and Jan, Y. (1991). Role of neurogenic genes in establishment of follicle cell fate and oocyte polarity during oogenesis in Drosophila. *Cell* **66**, 433-449.

Schwed, G., May, N., Pechersky, Y. and Calvi, B. R. (2002). Drosophila minichromosome maintenance 6 is required for chorion gene amplification and genomic replication. *Mol Biol Cell* **13**, 607-620.

Serizier, S. B. and McCall, K. (2017). Scrambled Eggs: Apoptotic Cell Clearance by Non-Professional Phagocytes in the Drosophila Ovary. *Front Immunol* **8**, 1642.

Shu, Z., Row, S. and Deng, W. M. (2018). Endoreplication: The Good, the Bad, and the Ugly. *Trends Cell Biol* **28**, 465-474.

Silver, D. L., Geisbrecht, E. R. and Montell, D. J. (2005). Requirement for JAK/STAT signaling throughout border cell migration in Drosophila. *Development* **132**, 3483-3492.

Silver, D. L. and Montell, D. J. (2001). Paracrine signaling through the JAK/STAT pathway activates invasive behavior of ovarian epithelial cells in Drosophila. *Cell* **107**, 831-841.

Spradling, A. and Mahowald, A. (1980). Amplification of genes for chorion proteins during oogenesis in Drosophila melanogaster. *Proc Natl Acad Sci U S A* **77**, 1096-1100.

Spradling, A. C. (1993). Developmental Genetics of Oogenesis. In *The Development of Drosophila melanogaster* (ed. M. B. a. A. Martinez-Arias). Cold Spring Harbor, New York: Cold Spring Harbor Press.

Sprenger, F., Yakubovich, N. and O'Farrell, P. H. (1997). S-phase function of Drosophila cyclin A and its downregulation in G1 phase. *Curr Biol* **7**, 488-499.

Sun, J. and Deng, W. M. (2007). Hindsight mediates the role of notch in suppressing hedgehog signaling and cell proliferation. *Dev Cell* **12**, 431-442.

Tepass, U., Gruszynski-DeFeo, E., Haag, T. A., Omatyar, L., Torok, T. and Hartenstein, V. (1996). shotgun encodes Drosophila E-cadherin and is preferentially required during cell rearrangement in the neurectoderm and other morphogenetically active epithelia. *Genes Dev* **10**, 672-685.

Uemura, T., Oda, H., Kraut, R., Hayashi, S., Kotaoka, Y. and Takeichi, M. (1996). Zygotic
Drosophila E-cadherin expression is required for processes of dynamic epithelial cell
rearrangement in the Drosophila embryo. *Genes Dev* **10**, 659-671.
Wang, X., Zheng, M., Fei, F., Li, C., Du, J., Liu, K., Li, Y. and Zhang, S. (2019). EMT-related
protein expression in polyploid giant cancer cells and their daughter cells with different
passages after triptolide treatment. *Med Oncol* **36**, 82.
Weaver, L. N., Ma, T. and Drummond-Barbosa, D. (2020). Analysis of Gal4 Expression
Patterns in Adult Drosophila Females. *G3 (Bethesda)* **10**, 4147-4158.
Woods, D. F., Hough, C., Peel, D., Callaini, G. and Bryant, P. J. (1996). Dlg protein is
required for junction structure, cell polarity, and proliferation control in Drosophila
epithelia. *J Cell Biol* **134**, 1469-1482.

Figure Legends

Figure 1: Induction of premature follicle cell endocycles. (A) Drawing of an ovariole with

select stages of oogenesis and follicle cell cycle transitions indicated below. Follicle cells are

pink. Above is the experimental timeline in relationship to the timing of egg chamber

maturation. (B-C′) DAPI staining and immunolabeling against E-cadherin in control (B, B′) and

Rux-expressing stage 10B egg chambers (C, C′); scale bar = 30 μm. (D) Quantification of DNA

Content (C) in control and Rux-expressing stage 10B egg chambers.; ****$p < 0.0001$; n = 85

nuclei in control and n = 104 nuclei in *UAS-Rux* across four biological replicates. (E)

Quantification of number of DAPI-labeled nuclei in control and Rux-expressing stage 10B egg

chambers; ****$p < 0.0001$; n = 12 egg chambers in both across four biological replicates.

Figure 2: Premature endocycling compromises egg quality and female fertility. (A-D)

Differential interference contrast images of control (A, C) and Rux-expressing (B, D) stage 10B

egg chambers after one (A, B) or two (C, D) days of incubation at 29°C; scale bar = 100 μm.

(E) Quantification of oocyte area in control and Rux-expressing stage 10 egg chambers.; *$p <$

0.05; for day one, n = 13 oocytes for control and n = 15 oocytes for UAS-*Rux* across three

biological replicates; for day 2, n = 20 oocytes for control and n = 24 oocytes for UAS-*Rux*

across five biological replicates. (F) Quantification of nurse cell compartment length; **$p =$

0.0021, ***$p = 0.0002$, ****$p < 0.0001$. For day 1, n = 20 egg chambers for control and UAS-

Rux across three biological replicates; for day 2, n = 27 egg chambers for control and n = 30

egg chambers for *UAS-Rux* across five biological replicates. (G-J′) DAPI staining and EdU

labeling of DNA replication in control (G, I, I′) and Rux-expressing (H, J, J′) stage 10B egg

chambers after one (G, H) or two (I-J′) days of incubation at 29°C; scale bar = 25 μm (G-I, J) or

10 μm (I′, J′). (K) Quantification of percent stage 10B egg chambers that had follicle cells

exhibiting pan-nuclear EdU labeling in control and Rux-expressing ovaries. ***$p = 0.0003$; n =

26 egg chambers for control and n = 19 egg chambers for *UAS-Rux* across three biological

replicates. (L-O) Brightfield images of eggs laid by control (L, N) and Rux-expressing females

(M, O) after one (L, M) or two (N, O) days of incubation at 29°C; scale bar = 200 μm.

Quantification of percent of eggs with thin shells (P) and hatch rate (Q) across three biological

replicates.

Figure 3: Premature endocycling disrupts follicular epithelial structure. (A-D″) DAPI
staining and immunolabeling against E-cadherin in control (A-A″, C-C″) or Rux-expressing (B-
B″, D-D″) early-stage egg chambers after one (A-B″) or two (C-D″) days of incubation at 29°C;
scale bar = 15 µm (A-D″). Stages (S) of oogenesis and germarium (G) are indicated. (E)
Quantification of DNA Content (C) in control and Rux-expressing stage 4 / 5 egg chambers
after two days of incubation at 29°C.; ****$p < 0.0001$; n = 25 nuclei for control and n = 15 nuclei
for *UAS-Rux* across three biological replicates.

Figure 4: Premature endocycling inhibits border cell migration. (A-B′) DAPI staining and
immunolabeling against E-cadherin in control (A, A′) and Rux-expressing (B, B′) egg chambers
after two days of incubation at 29°C; scale bar = 30 µm (A, B) or 15 µm (A′, B′). White arrows
indicate border cell clusters. White lines indicate extent of migration of border cell clusters, and
red lines indicate extent of migration of main body cells. (C) Quantification of border cell
migration during stage 9 from control and Rux-expressing females after two days of incubation
at 29°C across five biological replicates. Each dot represents one egg chamber with the
distance migrated by its main body and border cells plotted on the x- and y-axis, respectively.
(D- E‴) DAPI staining and immunolabeling against FasIII and E-cadherin in control (D-D‴) and
Rux-expressing (E-E‴) border cell clusters after two days of incubation at 29°C; scale bar = 10
µm.

Figure 1

A

~Stage at temperature shift   S4/5   ← 2 days →   Assay stage 10

Stem cells   S6   S7   S9   S10   S14

10 mitotic cycles   **3 endocycles (16C)**   **post-endocycle / amplification**
~3 days   **~24h**   **~15hr**

B *tj*-GAL4 Control — DAPI — 30 μm

C *tj*-GAL4 UAS-*Rux* — DAPI — 30 μm

B′ *tj*-GAL4 Control — DAPI / E-cadherin — 30 μm

C′ *tj*-GAL4 UAS-*Rux* — DAPI / E-cadherin — 30 μm

D DNA Content (C) Stage 10 — **** — + / UAS-*Rux* / *tj*-GAL4

E # of follicle cells Stage 10 — **** — + / UAS-*Rux* / *tj*-GAL4

FIGURE 2

FIGURE 3

1 Day at 29°C

A | *tj*-GAL4 Control
S2
S3/4
S1 | S5/6
G
15 µm | DAPI

B | *tj*-GAL4 UAS-*Rux*
S2
S3/4
15 µm | DAPI

2 Days at 29°C

C | *tj*-GAL4 Control
S2 | S3 | S4
15 µm | DAPI

D | *tj*-GAL4 UAS-*Rux*
S3/4
S5
15 µm | DAPI

A′ | *tj*-GAL4 Control
S2
S3/4
S1 | S5/6
G
15 µm | E-cadherin

B′ | *tj*-GAL4 UAS-*Rux*
S2
S3/4
15 µm | E-cadherin

C′ | *tj*-GAL4 Control
S2 | S3 | S4
15 µm | E-cadherin

D′ | *tj*-GAL4 UAS-*Rux*
S3/4
S5
15 µm | E-cadherin

A″ | *tj*-GAL4 Control
S2
S3/4
S1 | S5/6
G
15 µm | DAPI
E-cadherin

B″ | *tj*-GAL4 UAS-*Rux*
S2
S3/4
15 µm | DAPI
E-cadherin

C″ | *tj*-GAL4 Control
S2 | S3 | S4
15 µm | DAPI
E-cadherin

D″ | *tj*-GAL4 UAS-*Rux*
S3/4
S5
15 µm | DAPI
E-cadherin

E

2 Days at 29°C

****

DNA Content (C) Stage4/5

+   UAS-*Rux*

tj-GAL4

FIGURE 4