Title:

Multiomics-assisted characterization of rice-Yellow Stem Borer interaction provides genomic and

mechanistic insights into stem borer tolerance in rice

Running title: Characterization of rice-Yellow Stem Borer interaction

Authors:

Gokulan C. G.[1], Umakanth Bangale[2], Vishalakshi Balija[2], Suneel Ballichatla[2], Gopi Potupureddi[2],

Deepti Rao[1], Prashanth Varma[2], Nakul Magar[2], Karteek J.[2], Sravan M.[2], Padmakumari A. P.[2], Gouri

S Laha[2], Subba Rao LV[2], Kalyani M. Barbadikar[2], Meenakshi Sundaram Raman[2], Hitendra K.

Patel[1,3], M Sheshu Madhav[2,4, *], Ramesh V. Sonti[1,5,*]

Affiliation:

[1]CSIR-Centre for Cellular and Molecular Biology, Telangana, India - 500007.

[2]ICAR-Indian Institute of Rice Research, Telangana, India - 500030.

[3]Academy of Scientific and Innovative Research, Uttar Pradesh, India - 201002.

[4]Current address: ICAR-Central Tobacco Research Institute, Andhra Pradesh, India - 533105.

[5]Current address: International Centre for Genetic Engineering and Biotechnology, New Delhi,

India - 110067

[*]Addresses for correspondence: sheshu24@gmail.com; sonti@ccmb.res.in

ORCiD:

Gokulan C. G.: 0000-0002-4305-2818

AP Padmakumari: 0000-0003-2318-2540

Hitendra K. Patel: 0009-0003-4016-4739

Sheshu Madhav: 0000-0001-6100-5751

Ramesh V. Sonti: 0000-0003-4845-0601

SUMMARY:

Yellow stem borer (YSB), *Scirpophaga incertulas* (Walker) (Lepidoptera: Crambidae), is a major

pest of rice in India, that can lead to 20-60% loss in rice production. Effective management of YSB

infestation is challenged by the non-availability of adequate source of resistance and poor

understanding of resistance mechanisms, thus necessitating studies for generating resources to breed

YSB resistant rice and to understand rice-YSB interaction. In this study, by using bulk-segregant

analysis in combination with next-generation sequencing, Quantitative Trait Loci (QTL) intervals in

five rice chromosomes were mapped that could be associated with YSB tolerance at vegetative

phase in a highly tolerant rice line named SM92. Further, multiple SNP markers that showed

significant association with YSB tolerance in rice chromosomes 1, 5, 10, and 12 were developed.

RNA-sequencing of the susceptible and tolerant lines revealed multiple genes present in the

candidate QTL intervals to be differentially regulated upon YSB infestation. Comparative

transcriptome analysis revealed a putative candidate gene that was predicted to encode an alpha-

amylase inhibitor. Analysis of the transcriptome and metabolite profiles further revealed a possible

link between phenylpropanoid metabolism and YSB tolerance. Taken together, our study provides

insights on rice-YSB interaction at genomic, transcriptomic and metabolomic level, thereby

facilitating the understanding of tolerance mechanism. Importantly, a promising breeding line and

markers for YSB tolerance have been developed that can potentially aid in marker-assisted breeding

of YSB resistance among elite rice cultivars.

Keywords: Rice, Yellow Stem Borer, QTL mapping, Insect resistance, Transcriptomics,

Metabolomics, Markers, Plant breeding.

SIGNIFICANCE STATEMENT:

Global rice production is threatened by various pests, among which stem borers pose serious

challenges. Hence, understanding the molecular intricacies of rice-stem borer interaction is

necessary for effective pest management. Here, we used a multi-omics approach to unravel the

mechanisms that might help rice combat yellow stem borer infestation, thus providing insights and

scope for developing YSB tolerant rice varieties. To facilitate the latter, we developed markers that

co-segregate with tolerance.

INTRODUCTION

Several pests affect rice cultivation globally and pose serious threats to grain production and quality. Over a hundred pests have been reported to infest rice, of which 15-20 cause severe agro-economic losses (Pathak, 1968; Pathak, 1977). Yield losses caused by insects can vary depending on the insect biotype, severity of the infestation, environmental factors, growth stage and genetic background of the host (Pathak, 1977). Among the major pests of rice, stem borers pose a serious threat as they cause devastating effects on rice production in both temperate and tropical regions. Commonly found stem borers in Asia include yellow stem borer (YSB; *Scirpophaga incertulas*), striped stem borer (SSB; *Chilo suppressalis*), white stem borer, dark-headed stem borer, and pink stem borer. YSB and SSB were the cause of a steady yield loss of an estimated 5%-10% in Asia (M.D. Pathak and Z.R.Khan, 1994). YSB is a predominant pest among other stem borers in the Indian sub-continent (M.D. Pathak and Z.R.Khan, 1994; Gururaj Katti, Chitra Shanker, Padmakumari A.P., 2011). A study in a field condition revealed that YSB infestation explained about 70% variation in the yield (Gangwar *et al.*, 1986). The female moths of YSB lay eggs on the leaf tips. After hatching, the larvae disperse through wind or move downwards and bore into the stem just above the water level. YSB infestation in rice is manifested as different symptoms based on the growth stage of the plants. The larvae of YSB borer into the rice stem at the base and feed on the growing tissue leading to formation of "dead hearts" in vegetative phase and white ears/ white heads in reproductive phase. Both dead hearts and white ears lead to a considerable yield loss as they both directly affect the production of rice by interfering with the number of productive tillers and panicles, respectively (Pathak, 1968; Muralidharan, K. and Pasalu, 2006). However dead heart damage at vegetative phase can be compensated up to 10% (Rubia *et al.*, 1996).

The control of stem borers in the field is still challenging and is majorly dependent on timely chemical intervention. The usage of chemicals affects human health, pollutes the ecosystem, and destroys the natural predators of insects and leads to development of resistance to insecticides. Using resistant or tolerant rice cultivars to counter the attack by pests and pathogens is one of the highly recommended practices in pest management. Resistance or tolerance among different cultivars to different species of stem borers has been reported (Heinrichs, Elvis & Medrano, F.G. and Rapusas, 1985). It was observed that the resistance of these varieties was primarily due to antibiosis, that is, an antagonistic reaction of rice towards stem borer which is detrimental to the latter (Pathak, 1977). It was found that rice plants can compensate for the damage through the reallocation of metabolites and photosynthates to uninjured stems/tillers and by initiating the

production of new tillers (Rubia *et al.*, 1996). Compensation was observed to be an effective

strategy to tolerate YSB infestation during the vegetative growth stage and not during the

reproductive stage. It was proposed that breeding rice lines for tolerance to YSB based on the

ability of the varieties to compensate for the YSB-caused injury could be a better approach (Rubia

et al., 1996).

Taking note of the extent of spread and severity of the insect (Padmakumari AP and Katti G,

2018), the necessity for efficient pest management is imperative. Hence, the development of rice

lines with YSB tolerance and using such lines for breeding purposes is of primary importance. This

study reports the identification of a promising rice line that shows enhanced tolerance to YSB

during vegetative phase of the crop. To further characterize the line and identify it as a potential

source of tolerance, bulk segregant analysis and QTLseq analysis of the SM92 mapping population

was performed. Putative QTL intervals linked to YSB tolerance were identified in five rice

chromosomes. Using SNP genotyping of the segregating population, several markers that showed

significant association with YSB tolerance were identified. In order to elucidate the mechanism of

tolerance, RNA-sequencing of the tolerant and susceptible lines upon YSB infestation was

performed. The analysis revealed modulation of phenyl propanoid pathway, lipid transport, and

alpha-amylase inhibition could be key pathways involved in YSB tolerance in rice. Metabolite

profiling further indicated accumulation of various phenylpropanoid metabolites upon YSB

infestation. Overall, this work identified, characterized, and developed resources for YSB tolerance

that could be of potential use in rice breeding.

RESULTS

SM92 was identified to be highly tolerant to YSB

A rice line named SM92 (described in Padmakumari A.P et al 2017; Potupureddi et al. 2021) was found to show tolerance to YSB during vegetative stage of growth (Figure 1A). A cut-stem assay was performed to reveal the status of YSB larvae inside the stems of SM and SM92. The results showed a significantly reduced number of live larvae inside the stems of SM92 as compared to that in SM (Figure 1B, C, D). This suggests the possibility of antibiosis mechanism in SM92.

QTL-seq revealed putative genomic intervals linked to YSB tolerance at vegetative phase

Bulk segregant Analysis (BSA) was performed to identify the genomic regions that are associated with YSB tolerance. For this, the mutant line in the M_6 generation (SM92) was crossed with the wildtype line. A segregating population containing 314 F_2 progenies was screened for YSB tolerance in field conditions for dead heart symptoms (Data S1). The phenotypic segregation followed normal distribution suggesting that the trait is governed by Quantitative Trait Loci (QTL; Figure 2A). The tolerant and susceptible DNA bulks as well as the parental DNA were subjected to a high coverage (~40X) of next-generation whole genome sequencing. The obtained raw reads were processed, aligned to the reference genome, and the variants were called. The variants were then used for the QTL-seq analysis to identify the QTL associated with YSB tolerance. QTL-seq analysis was performed using two different parameters, namely delta-SNP-index and G-prime (G'). The analyses overall revealed regions in 5 of the 12 rice chromosomes, namely chromosomes 1, 5, 7, 10, and 12, to be putatively linked to YSB tolerance (Figure 2B, C). In total, 10 QTL intervals were predicted in these 5 chromosomes (Table 1).

A comparison between the YSB tolerance QTL intervals and previously reported insect resistance QTL was performed. The QTL intervals identified in this study overlapped with the predicted QTL intervals linked to resistance against rice leaf folder (Chr1; Selvaraju et al., 2007) and BPH (Chr10 and Chr12; Sun et al., 2005; Jena et al., 2006) (Table S3). Although these QTL intervals are mapped for insect resistance, there is no further information on the possible genes or their alleles that might be involved in resistance. Also, no markers have been developed from these intervals for application in breeding programmes.

A large number of genes in the QTL intervals exhibit variations

The genes that harbor SNPs and that are present in the QTL interval were annotated using Variant Effect Predictor (VEP). There were 25,065 SNPs present in the QTL intervals in all five chromosomes (Figure S1A; Data S2). About 94% of these SNPs were in non-coding regions of the genome (Figure S1B). A total of 1101 genes harbor SNPs which include SNPs in coding (UTRs and exonic) and/or non-coding (upstream, downstream, and intronic) regions. Genes harboring SNPs in their coding regions were 459 in number and 34 among them harbored SNPs that were predicted to be deleterious (Data S3). Over-representation analysis of the genes that harbor mutations in the coding sequences revealed a significant enrichment of laccase-encoding genes, that are involved in phenylpropanoid/lignin metabolism (Figure S1C).

Marker-Trait Association Analysis revealed markers exhibiting significant association with YSB tolerance at vegetative phase

A set of fifty SNPs were selected as markers for genotyping the F_2 individuals using Kompetitive Allele Specific PCR (KASP^{TM}, LGC Genomics, UK; Data S4, S5). These SNPs were selected based on their effect on the genes that harbor them, the genomic distance between the preceding and the following SNP in such a way that the distance between the two markers is less than 250-300 kbp. KASP assay was performed from the F_2 plants showing extreme as well as intermediate phenotypes (Data S6). Further, a marker-trait association analysis was carried out to identify the markers that are significantly associated with the phenotype (Data S7). The KASP-based genotyping revealed a significant ($P<0.05$) association of fourteen alleles from chromosomes 1, 5, 10, and 12 with YSB tolerance in the F_2 individuals (Table 2). Each of the regions was calculated to explain about 5% to 9% phenotype variation observed in the F_2 progenies.

YSB infestation causes an extensive transcriptional reprogramming

Transcriptomics-based characterization of rice-YSB interaction was performed to understand the possible mechanism of tolerance to YSB. For this, RNA sequencing (RNA-seq) was performed on RNA isolated from the stem tissues of SM and SM92, with and without YSB infestation (3 days post infestation). The analysis of RNA-seq data showed several differentially expressed genes (DEGs) with overlapping and unique DEGs between SM and SM92 (Figure 3A, B; Data S8, S9). The number of DEGs was slightly more in SM (n=1950) when compared to SM92 (n=1857). Also, SM92 had a lesser number of downregulated genes than SM. Overall, only 13.7% of the DEGs were common between SM and SM92, suggesting a contrasting transcriptional reprogramming occurring in tolerant and susceptible lines upon YSB infestation.

Gene ontology (GO) and pathway enrichment analysis of the DEGs revealed the similarities and
differences between SM and SM92 in terms of their transcriptomics response to YSB infestation
(Figure 3C, D). Specifically, YSB-induced genes in SM92 were enriched in phenylpropanoid
metabolism, alpha-amylase inhibitors/lipid transfer protein and glycoside hydrolases (Figure 3D).
An in-depth analysis using MapMan revealed genes belonging to the GO term ribosome biogenesis
were exclusively downregulated in SM (Figure S2A). The categories consisting of genes involved
in the metabolism of simple phenols, lignin and lignans showed a striking difference between SM
and SM92, wherein these genes were majorly upregulated in SM92 (Figure S2B).

Phenylpropanoid pathway genes show remarkably contrasting expression pattern at
vegetative phase infestation in tolerant and susceptible lines

Genes involved in the biosynthesis of phenylpropanoids showed a remarkable upregulation in
SM92 while some of the genes were downregulated in SM (Figure 3D; Figure S3). The
Phenylpropanoid Pathway (PPP) is an important secondary metabolism pathway that is essential for
normal plant growth, development and defense against various pests and pathogens (Dong and Lin,
2021). PPP leads to the production of various types of lignin and lignin derivatives. In line with this,
overall, thirteen *laccase*-encoding genes, that are predicted to be involved in lignin metabolism,
were upregulated in SM92 upon YSB infestation. Markedly, five of the thirteen *laccase* genes are
present within the chromosome 1 QTL interval (Table S4). Further analysis revealed that
cumulatively 118 genes that are in the QTL intervals are differentially expressed in SM and SM92
upon YSB infestation (Data S10). A deeper look into these genes indicated lignin metabolism-
related genes, *laccases* (n=5), to be enriched among the genes upregulated exclusively in SM92
(Figure S4A). Promoter analysis of the five *laccase* genes revealed the presence of tandem CATG
palindromic motif (CATGCATG) in the promoters of all five genes (Fig S4B). Further analysis
indicated a significant overlap between the observed motif and the binding element of a B3-domain
containing transcription factor (Figure S4C). It could hence be speculated that a common mode of
regulation might be acting on these *laccase* genes in SM92, probably mediated by a B3-domain
containing transcription factor.

Comparative analysis suggests putative candidate genes for stem borer tolerance in rice
A comparative transcriptomics analysis was performed between the data generated in this study and
a previously published data where rice lines with varying degree of resistance to striped stem borer
(SSB) were infested with SSB (Wang *et al.*, 2018). It was observed that 527 genes were commonly
upregulated by SSB and YSB in the respective tolerant lines (Figure 4A; Data S11). A gene set

enrichment analysis revealed that genes annotated as alpha-amylase inhibitors/lipid transfer protein,

carbohydrate metabolism, and cell wall-related genes were significantly enriched (Figure 4B). A

striking observation was that the highest upregulated gene in SM92 upon YSB infestation –

OsLTPL146, a Lipid Transfer Protein-like (LTPL) gene – is present within the mapped QTL

interval in Chromosome 10. In addition, the same gene was significantly downregulated in SM.

OsLTPL146 exhibited a similar expression pattern upon SSB infestation as well. That is,

OsLTPL146 was upregulated in a resistant cultivar and downregulated in a susceptible cultivar upon

SSB infestation (Figure 4C). Interestingly, it was also observed that twenty-five lipid-

metabolism/transport related genes majorly including *LTPL* class of genes were upregulated in

SM92 upon YSB infestation, while only five genes showed differential expression in SM (Figure

S5A). Other lipid/fatty-acid metabolism related genes that are involved in cutin, suberine, and wax

synthesis were also exclusively upregulated in SM92 (Figure S6). Another unique expression

pattern observed was that of the glutathione metabolism genes consisting of the Phi and Tau classes

of *glutathione transferases* (*GSTs*). These genes were exclusively upregulated in SM92 post YSB

attack (Figure S6).

As mentioned above, twenty-five enriched genes belonged to the lipid metabolism/transport

category. Of the twenty-five genes, twenty-two genes were annotated as "Protease inhibitor/seed

storage/LTP family protein precursor". To elucidate the actual biochemical properties of these gene

products, the structures of all the twenty-five protein sequences were predicted using AlphaFold2

and their structural homologues were identified. The results suggest that fifteen genes likely encode

lipid transfer proteins as their structures were homologous to characterized lipid transfer proteins

from various organisms, five gene products were homologous to alpha-amylase inhibitors and of the

remaining five genes, one coded for a lyase, one for a phospholipase D, one for annexin and the

structures could not be predicted for the other two gene products (Figure S5B).

Notably, structure prediction analysis suggests that the predicted structure of *OsLTPL146* is

homologous to an alpha-amylase inhibitor whose structure is experimentally solved. This result

indicates that *OsLTPL146* – the highest upregulated gene in SM92 upon YSB infestation – likely

encodes an alpha-amylase inhibitor (Figure 4D). Further, owing to the difference in the expression

pattern of *OsLTPL146* in SM and SM92, the promoter sequence of *OsLTPL146* in SM and SM92

were compared. The alignment revealed considerable variations in the promoter and 5' untranslated

region (UTR) of the gene between SM and SM92in certain known cis-elements. The variations

included single nucleotide polymorphisms, small insertions, and deletions (Figure S7). Whether the

variations in the promoter/regulatory sequence of *OsLTPL146* are involved in the YSB-mediated

induction of the transcript and/or YSB tolerance is another avenue for further research. Taken

together, detailed investigation of OsLTPL146 promoter variation and protein biochemistry could

reveal its role in stem borer resistance.

Levels of a few phenylpropanoid pathway metabolites are altered upon YSB infestation

An untargeted metabolite profiling of sheath samples that were either uninfested or infested with

YSB larvae was performed to gain insights into the mechanism of tolerance to YSB. Over 1000

metabolites were identified in all the tested conditions (Data S12, and S13). The overall profile

indicated a difference in the metabolite levels between uninfested SM and SM92 plants. However,

YSB infestation seems to result in highly comparable metabolite profile in both SM and SM92

(Figure 5A). Further, the levels of phenylpropanoid pathway (PPP) associated metabolites were

analysed specifically, as the RNA-seq data indicated an upregulation of genes associated with the

pathway. The results showed the presence of 27, and 25 PPP metabolites in SM and SM92 samples,

respectively (Figure S8). Four of the PPP metabolites including 5-O-Caffeoylshikimic acid,

Coniferyl acetate, Caffeate, and L-Phenylalanine exhibited significant differential accumulation in

SM and/or SM92 upon YSB infestation when compared to uninfested samples (Figure 5B). YSB

infestation resulted in increased levels of 5-O-Caffeoylshikimic acid and Coniferyl acetate in both

SM and SM92. On the other hand, Caffeate and L-Phenylalanine exhibited significantly reduced

accumulation in SM92 upon YSB infestation. Owing to the variation among the samples, difference

in the levels of other metabolites could not be reliably identified. Targeted analysis of PPP

metabolites could provide a better understanding of the regulation of the pathway with respect to

YSB infestation.

DISCUSSION

Stem borers cause maximum yield loss among all the insects that attack rice in India (Savary *et al.*, 2019). Complete resistance to stem borers in nature is scarce and unidentified yet. Being a monophagous pest, it feeds only on rice plants and perpetuates. As a borer, by nature of its feeding habit it must feed on the plant for its survival. The damage occurs in the process of entering the plant and therefore some extent of damage is inevitable. Hence, the dependance on chemical insecticides prevail despite their negative impact on health and the ecosystem (Su *et al.*, 2014). Most stem borer tolerant varieties of rice exhibit compensation and antibiosis traits (Rubia *et al.*, 1996). However, the exact mechanism of tolerance and the molecular players involved are unknown.

The advent of bulk segregant analysis and next-generation sequencing methods have accelerated QTL identification and gene mapping for the traits of interest. This study identified a rice line, named SM92, that is highly tolerant to YSB. Screening the F_2 population, generated through bi-parental mating, at vegetative phase revealed that YSB tolerance to be a quantitative trait. Therefore, the segregants with extreme phenotypes were used for preparing tolerant and susceptible DNA bulks and were subjected to high-coverage sequencing. Analysis of the sequence data using QTL-seq pipeline indicated the presence of QTL intervals in five rice chromosomes that are putatively linked to YSB tolerance, thus validating the observed segregation pattern. Notably, three of the identified genomic regions were found to be overlapping with known insect resistance QTL intervals (Table S3). That is the Chr1 QTL was partly overlapping with a previously identified rice leaf folder resistance QTL whereas the Chr10 and Chr12 QTL intervals partly overlap with BPH resistance QTL. This observation suggests that the identified genomic intervals are of potential importance in insect resistance/tolerance and are likely to contain genes/alleles that are part of the insect resistance/tolerance mechanism of rice. The QTL mapping data indicated that SM alleles contribute to tolerance arising from Chromosome 1 QTL interval whereas SM92 alleles contribute to tolerance arising from other intervals (Figure 2B). Further analysis indicated that *laccase* genes, involved in lignin metabolism, are overrepresented in the QTL intervals.

Laccases and plant defense

Laccase (LAC; EC1.10.3.2) belongs to a multi-copper oxidase family of enzymes that play a crucial role in plant development and stress responses (Yu *et al.*, 2021). Various studies have identified the multi-faceted roles of laccases in plants that include cell wall elongation, secondary cell wall formation, lignification, pigmentation, flavonoid biosynthesis, metal ion stress, and seed setting.

Laccases are involved in catalyzing the polymerization of lignin monomers in the apoplast (Zhao *et*

al., 2013). Lignin, being a component of the secondary cell wall, plays important roles in protecting

cells from pathogens. Previous studies have established a positive correlation between lignin

accumulation and resistance to bacterial and fungal infections (Lee *et al.*, 2019; Peltier *et al.*, 2009).

Previous studies have also associated the levels of lignin pathway precursors including

hydroxycinnamic acid positively with resistance to stem borers (Santiago *et al.*, 2013; Malvar *et al.*,

2008). A study in maize showed a positive correlation between levels of phenylpropanoids

including *p*-coumarate and syringyl lignin to Mediterranean Corn Borer resistance (Gesteiro *et al.*,

2021). Our transcriptome profiling indicated thirteen *laccase* genes to be significantly upregulated

in SM92 upon YSB infestation. Additionally, some of the genes involved in the phenylpropanoid

metabolism pathway exhibited a strikingly contrast expression pattern in SM (downregulated) and

SM92 (upregulated) upon YSB infestation. Notably, five of the thirteen *laccase* genes are in the

identified QTL intervals in Chromosomes 1. It was also noted that some of the phenylpropanoid

pathway genes were downregulated in SM, the susceptible line (Figure S3). Together, the data

suggest a likely involvement of lignin metabolism in conferring tolerance to YSB. Although

previous studies have pointed out a generic role of laccases in pathogen/pest resistance, our study

has provided promising candidate genes that can be studied further to understand the mechanism of

tolerance at molecular level.

Ribosomal proteins as potential candidates to understand plant defense mechanisms

On examining the gene enrichment in the susceptible interaction, a remarkable number of genes that

code for ribosomal subunit proteins (RPs – Ribosomal proteins) were found to be significantly

downregulated. Both the large and small subunit protein-encoding genes were highly

downregulated. Previous studies have pointed out that RPs possess moonlighting functions. Such

extra-ribosomal functions of RPs include RNA-binding and promoting/preventing translation, role

in apoptosis, cell-cycle regulation, and stress resilience in prokaryotes as well as eukaryotes (Wool,

1996; Weisberg, 2008; Warner and McIntosh, 2009; Trautmann and Ramsey, 2022). In plants, two

RPs viz. RPL12 (Ribosomal Protein Large subunit 12) and RPL19 in *Nicotiana benthamiana* and

Arabidopsis thaliana were shown to be important for non-host resistance as the mutants exhibited

delayed hypersensitive response to non-host pathogens (Nagaraj *et al.*, 2016; Ramu *et al.*, 2020). A

targeted transcriptome profiling of thirty-four large subunit genes indicated that they are highly

stress responsive. The genes were either upregulated or downregulated depending on the nature of

the stress. Infection of rice plants with the bacterial blight pathogen caused upregulation of the

majority of the tested RPs (Moin *et al.*, 2016). Downy mildew infection in grapevine resulted in

accumulation of RPs during the early timepoints post-infection (Santos *et al.*, 2020). However, our

data revealed a significant downregulation of ribosomal protein-coding genes. This could be an

energy-conserving strategy that the host deploys to enhance defense against insect attack. But it is

also possible that reduced expression of RPs is a consequence of the mechanisms that the pest uses

to suppress defense, and this ultimately results in the host succumbing to the attack. It is noteworthy

that the tolerant line did not show any alteration in the expression of RP-encoding genes (Figure

S2A), thus suggesting susceptibility being a consequence of RP gene downregulation. Further

studies that follow the expression pattern of RP-encoding genes at different time points post

infestation can provide a clearer picture of the role of RPs in rice-stem borer interaction.

Insights gained through structure prediction analysis

Another striking observation from the transcriptomics data was the differential expression of

approximately twenty genes that were annotated as Lipid transfer-like proteins (LTPLs)/Protease

inhibitors (PIs). One of the genes *OsLTPL146* showed the highest upregulation ($\log_2$FC>10) upon

YSB infestation in the tolerant line while it was downregulated in the susceptible line. Importantly,

OsLTPL146 is located in the Chromosome 10 QTL interval, suggesting a possible link between the

expression of the gene and YSB tolerance at vegetative phase. A similar pattern of expression was

observed for *OsLTPL146* in an independent transcriptomics study performed in rice upon striped

stem borer infestation (Wang *et al.*, 2018). Owing to the high sequence similarity between LTPs

and PIs, based on the annotation and sequence alone it is hard to predict the biochemical function of

their gene products. Hence, the recently developed, artificial intelligence-based AlphaFold2 was

used to predict the structure of the differentially expressed LTPLs (Jumper *et al.*, 2021). The results

indicated that fifteen out of the twenty LTPLs exhibit structural homology to lipid transfer proteins

while the rest five were homologous to alpha-amylase inhibitors. The presence of proteins of

different classes suggest that multiple modes of tolerance are likely in action including protease

inhibition, lipid transfer, and alpha-amylase inhibition.

Lipid transfer proteins (LTP and LTP-like (LTPL) proteins) have been associated with pathogen

resistance, plant development, abiotic stress development, and cuticular wax metabolism among

others (Gao *et al.*, 2022). LTPs are small proteins that bind to lipids and aid in their transport across

cellular compartments (Salminen *et al.*, 2016). Multiple rice LTPs/LTPLs have been associated

with pollen development, tolerance to abiotic stress such as drought, and plant development (Gao *et*

al., 2022). *OsLTP5* was found to be induced by cutin monomers, abscisic acid, and salicylic acid

(Kim *et al.*, 2008). Cutin monomers possess a defense elicitation property against a potential

invasion (Schweizer *et al.*, 1996; Fauth *et al.*, 1998; Kolattukudy *et al.*, 1995). Class I LTPs were

identified to be induced upon infection by the blast fungus, *Magnaporthe oryzae* (Tae *et al.*, 2006).

The damage caused by fungus and chewing insects results in cuticle damage which in turn leads to

the generation of cutin monomers that subsequently activate defense signaling such as accumulation

of H_2O_2 (Fauth *et al.*, 1998). Moreover, our transcriptome analysis also revealed enrichment of

genes involved in cutin/wax biosynthesis to be upregulated in SM92 upon YSB infestation,

suggesting a possible production of cutin and/or reorganization of the cuticle being associated with

YSB tolerance.

Alpha-amylase inhibitors (AAIs) belong to multi-family proteins that interact with and inhibit

alpha-amylase enzymes (Franco *et al.*, 2002; Oneda *et al.*, 2004). Alpha amylases are important

digestive enzymes that help in the digestion of plant-derived starch/amylose that are ingested by the

insects (Lage, 2018). The evolutionary arms race between plants and pests has resulted in the

evolution of AAIs that inhibit alpha-amylase activity in the insect gut and thus negatively impacts

insect survival (Franco *et al.*, 2002).

Multi-layered mechanisms to counter stem borer infestation in rice

Studies so far have identified such tolerance-associated mechanisms individually in different

plants/varieties. But our observations in combination with other reports in the literature suggest a

multimodal mechanism of tolerance to YSB in SM92. Our results suggest that the striking

upregulation of multiple Lipid Transfer Protein like genes and alpha-amylase inhibitor genes in

SM92 is potentially linked to YSB tolerance in a manner that is either dependent on or independent

of wax/cuticle damage. Upregulation of a considerable number of lipid transfer protein-like (LTPL)

genes indicates a possibility of major alteration in the lipid composition of the cells. The

upregulation of cutin/wax synthesis genes in combination with *LTPLs* provides a reasonable

indication of the involvement of lipid translocation during insect attack as a plant defense strategy.

More studies on this can provide novel insights into plant defense signaling and mechanisms

involving lipid moieties. Also, future studies directed towards OsLTPL146, and its amylase

inhibition activity could provide a native source of alpha-amylase inhibition thereby eliminating the

need for a source of foreign origin.

Additionally, the upregulation of phenylpropanoid metabolism genes and laccases adds one more

layer to the complex mechanism of YSB tolerance in SM92. The presence of signature motif in the

promoter sequence of similarly regulated *laccases* suggests a common regulatory network that

might control the expression of lignin metabolism genes in SM92 upon YSB infestation. Therefore,
SM92 is a very promising line for gaining insights into fundamental resistance mechanisms as well
as for breeding programmes to introgress YSB tolerance into breeding lines. The former can be
accomplished by studying the candidate genes from this study based on QTL-seq and
transcriptomics results, while the latter has been facilitated by the markers that were identified to be
significantly associated with tolerance.

Breeding for YSB tolerance
Marker-assisted breeding is a viable option for introgressing traits from one crop variety to another.
In this regard, several SNP markers have been identified in this study that span the QTL intervals in
four different chromosomes. Notably, the Chr10 QTL interval was fine mapped to a ~180kb region
with five linked SNP markers. On the other hand, a few markers have been identified to be linked to
YSB tolerance in Chr1, Chr5, and Chr12 QTL intervals. Further mapping studies and marker-trait
association analyses using high density markers could help in fine mapping the QTL intervals
thereby providing additional markers with tight linkage/association to YSB tolerance.

Overall, to the best of our knowledge, the study presented here is the first to provide important
insights at genomic, transcriptomic, and metabolomic levels to uncover the genomic loci and
mechanisms of tolerance to yellow stem borer in rice. Stem borers, one of the major group of pests
affecting global rice production, in combination with changing and erratic climatic conditions, pose
a serious threat to food security. Knowledge from this study would aid in orienting future research
to better understand the mechanisms as well as provide additional gene targets to breed stem borer
resistance   in   rice   and   possibly   in   other   related   crops.

EXPERIMENTAL PROCEDURES

Plant materials and Insect

Plant materials used for this study include the cultivar Samba Mahsuri (SM) and the EMS mutant line named SM92. SM92 is registered as a germplasm at ICAR-National Bureau of Plant Genetic Resources (National Identity: IC646828). Yellow Stem Borer (YSB; *S. incertulas*) moths were collected from the rice field and were released for egg laying on rice plants. The neonate larvae emerging from the egg masses were used for infestation of the mutant lines, F_2, and $F_{2:3}$ progeny under field conditions by releasing 2 neonate larvae per tiller at the maximum tillering phase (Bentur JS *et al*, 2011; Padmakumari AP and Katti G, 2018). The phenotype was scored when the dead heart symptoms were apparent on the susceptible cultivar (IRRI, 2014). All the YSB screening activities were conducted in the greenhouse and/or rice field facility of ICAR – Indian Institute of Rice Research, Hyderabad, India (17.32 N, 78.39 E).

Cut-stem assay

Cut stem insect bioassays were carried out at the ICAR- IIRR. At vegetative phase, stem cuttings were collected from healthy tillers of both SM and SM92. The stems were cut into 4-5 cm long pieces and placed on Petri plates containing Whatman paper moistened with benzimidazole (75 mg/L). Freshly hatched neonate YSB larvae were then released onto each cut stem at 5 larvae/stem. The plates were kept in an incubator at 27±2°C. After 4 days, the plates were opened, and the stems were dissected carefully under a microscope. Before dissection, the stem pieces were observed for the presence of feeding holes. Upon dissection, the stems were observed for feeding injury and for the number of dead/surviving larvae were counted.

Bulk segregant Analysis and sequencing

For mapping the genomic region(s) associated with YSB tolerance, bulk segregant analysis (BSA) followed by next-generation sequencing and QTL-seq analysis was performed as described earlier (Takagi *et al.*, 2013; Mansfeld and Grumet, 2018). A cross was made between SM (WT; recipient) and SM92 (Mutant; donor). The F_1 plants were confirmed for hybridity and advanced to F_2 generation comprising 314 plants. The F_2 progenies and the parental lines were screened during the vegetative growth phase. The plants were scored based on the percentage of dead heart as recommended by the Standard Evaluation System of Rice (IRRI, 2014). F_2 plants showing the extreme phenotypes i.e., tolerance (n=21) and susceptibility (n=23) to YSB were selected to constitute the tolerant and susceptible bulks, respectively, after confirming the phenotypes of the $F_{2:3}$ lines. DNA from the bulk constituents and parents were isolated using the CTAB method and

quantified using Qubit^{TM} High-Sensitivity dsDNA Assay in a Qubit^{TM} 4 fluorometer (Thermo

Fisher Scientific, USA). The DNA quality was assessed on 0.8% agarose gel through

electrophoresis. Equal concentrations of DNA from the tolerant and susceptible plants were bulked

separately. 1 µg DNA from all the samples was used to prepare sequencing libraries with TruSeq

Nano DNA HT Sample Preparation Kit (Illumina, USA) and qualified libraries were sequenced on

the Illumina HiSeq2500 platform to obtain 2x150 bp reads at approximately 40x coverage. The

library preparation and sequencing were outsourced to Nucleome Informatics Private Limited,

Hyderabad, India.

Sequencing analysis and gene mapping

Raw reads were obtained from the service provider and processed using *FastQC* to assess the

quality. The reads were then used for aligning against the reference genome (Nipponbare) using

BWA mem. The initial alignment (*.bam*) files were sorted, and duplicates were removed using

samtools. The analysis-ready bam files were used for calling variants using *freebayes* and the

variants were filtered and processed using *VCFtools* and *bcftools*. The variants were filtered to

retain only Single Nucleotide Polymorphisms (SNPs) with a mapping quality greater than 30, a

minimum and the maximum depth of 10 and 200, respectively. The filtered SNPs were then

provided as input to the R language-based QTL mapping tool called *QTLseqr* with default filtering

criteria and 0.5 Mb window size. The *QTLseqr* identifies QTL intervals by calculating two

parameters namely, Delta-SNP-index and the G' statistics (Mansfeld and Grumet, 2018; Magwene

et al., 2011). The DNA sequence and analysis statistics are provided in Table S1.

SNP annotation and analyses

Information of the putative QTL intervals obtained from the QTL-seq analysis was used to obtain

the variations in the intervals. *Bedtools* (Quinlan and Hall, 2010) was used to extract the SNPs in

the QTL interval from the whole genome variant file. The QTL SNPs were then annotated using

Variant Effect Predictor (VEP) to identify the effect of the SNPs on the genome. Microsoft Excel

was used for further analyses like segregating the SNPs based on their effects.

Marker development and validation

SNPs having a high impact on the associated genes were chosen based on the effect (non-sense

mutations) and SIFT score (missense mutations causing intolerable amino acid changes) provided

by VEP. The presence and absence of selected SNPs in SM92 (mutant line) and SM (the wildtype

line), respectively, were confirmed through Sanger's sequencing. A total of 50 SNPs (42 from the

QTL intervals and 8 from the rest of the genome) were finally selected for developing Kompetitive

Allele Specific PCR (KASP[TM]) assays (LGC Genomics, UK). The F_2 individuals derived from

SM92 X SM cross were genotyped using KASP assays for all 50 SNPs in the ABI ViiA7 RT-PCR

System or Bio-Rad CFX384 system. The genotyping was performed in a 5 µl reaction per assay

under conditions recommended by the manufacturer. KASP reaction condition was as follows: 94 □

- 15 minutes; 10 cycles of 94□ for 20 seconds and 61-55□ for 1 minute (touch-down of

0.6□/cycle); 26 cycles of 94□ for 20 seconds and 55□ for 1 minute. TASSEL was used to

statistically assess the association between the genotypes and YSB tolerance.

RNA sequencing and analysis

SM and SM92 plants were infested with YSB larvae during the tillering stage. Stem samples were

collected from the infested stems an area 1 cm on either side of the larval entry hole 3 days after

release of larvae. Also, uninfested stem samples were collected as a control. All the samples were

collected as duplicates. The samples were flash-frozen immediately after collection and stored at -

80□ until further processing. RNA was isolated from the samples using RNeasy Plant RNA

Isolation Kit (Qiagen, Germany) with on-column DNA digestion using RNase-Free DNase Set

(Qiagen, Germany). Purified RNA was quantified using Qubit[TM] RNA HS Assay Kit (Thermo

Fisher Scientific, USA). RNA integrity was assessed using agarose gel electrophoresis and

BioAnalyzer 2100 (Agilent Technologies, USA). 1 µg of total RNA samples was used for library

preparation for each sample using the Illumina TruSeq Stranded Total RNA with Ribo-Zero Plant

(Illumina, USA). Sequencing was performed on Illumina's NovaSeq6000 platform with cycling

conditions to obtain 2x100bp reads on an S2 flowcell to obtain ~70 million reads per sample. The

RNA sequence statistics are provided in Table S2. Library preparation and sequencing were

performed in the NGS facility of CSIR-CCMB, Hyderabad.

The raw data was obtained, and the quality was assessed using *FastQC*. Adapters and low-quality

reads were trimmed using *Trim Galore!* (Krueger). *RNA-STAR* was used to align the trimmed reads

to the rice genome (MSU Rice Genome Annotation Project - version 7) (Dobin *et al.*, 2013). Reads

that mapped to the genes were quantified using *featureCounts* (Liao *et al.*, 2014). Differential gene

expression was obtained using the R library *DEseq2* (Love *et al.*, 2014). Genes with a false

discovery rate ≤ 0.05 and $\log_2$ fold change ≤ -2 or ≥ 2 were considered differentially expressed

genes (DEGs). For gene ontology, and pathway analyses of the DEGs, tools including gProfiler,

ShinyGO, and MapMan were used (Thimm *et al.*, 2004). *Venny* was used for generating Venn

diagrams (Oliveros, 2007).

Structure prediction and homology search

Amino acid sequences of the DEGs classified under the Lipid Transport/Metabolism category were retrieved from RAPDB (RAP-DB; Sakai et al., 2013; Kawahara et al., 2013). The sequences were subjected to structure prediction using ColabFold (AlphaFold2.ipynb - Colaboratory) which uses the AlphaFold2 algorithm to predict 3D structure models of proteins (Mirdita *et al.*, 2022). In order to identify the structural homologues of the predicted structures, a structural homology search was performed using the PDB search option of DALI server (Dali server; Holm, 2022). Multiple sequence alignment of protein sequences and the phylogenetic tree were generated using Clustal Omega (Sievers *et al.*, 2011; Goujon *et al.*, 2010). The phylogenetic tree visualization and annotation were performed using iTOL (Letunic and Bork, 2021).

Metabolite profiling and analysis

SM and SM92 plants were infested with YSB larvae during the tillering stage. Stem samples were collected from the infested stems an area 1 cm on either side of the larval entry hole 3 days after release of larvae. Also, uninfested stem samples were collected as a control. All the samples were collected in triplicates. The samples were flash-frozen immediately after collection and stored at -80□ until further processing. Further, the frozen samples were lyophilized thoroughly, weighed, and 30 mg of the lyophilized samples were powdered. Metabolites were extracted in 1ml of 80% methanol at 30°C for 30 minutes in shaking condition before centrifuging the samples at 14000rpm for 5 minutes. The supernatant was filtered through 0.22μm membrane. The extracts (10μl each) were analysed by the Shimadzu Prominence HPLC coupled with Shimadzu triple quadrupole LCMS-8045 mass spectrometer (Shimadzu Corporation, Kyoto, Japan). HPLC was performed with a CLC0181 column (Shimpack Gist C18 75*4.0mm, 3μm) with a runtime of 50 minutes at a flow rate of 0.8mL/minute and column temperature of 30°C. Mobile phases used for HPLC include 0.04% (v/v) acetic acid in water and 100% acetonitrile. The electrospray ionization (ESI)–MS analysis was performed in both positive and negative ion modes. Full-scan mass spectra were acquired over a mass range of m/z 50–2000. The sample extraction and metabolite profiling were outsourced to Novelgene Technologies Pvt. Ltd., Hyderabad, India.

The raw intensity data of the detected metabolites was analysed using the one factor statistical analysis module of MetaboAnalyst 5.0 using KEGG compound identifiers (Pang *et al.*, 2021). The data was normalized using quantile normalization method, log-transformed, and scaled using the auto-scaling option. The results were downloaded and further processed in Microsoft Excel.

DATA AVAILABILITY

SM and SM92 parent whole genome sequences are deposited in NCBI-SRA under the accession PRJNA658718. The F_2 bulk DNA data are deposited under the accession PRJNA972670. The transcriptome data are deposited in NCBI-Gene Expression Omnibus (GEO) under the accession GSE245213.

AUTHOR CONTRIBUTIONS

Conceptualisation - RVS, MSM, HKP, and PAP. Investigation - GCG (QTL-seq, RNA-seq, metabolomics analyses and KASP genotyping); UB, VB, SB, PAP (YSB larvae rearing, F_2 population generation, and screening of rice plants); DR (QTLseq data analysis); PV, NM, KJ, SM, PAP (screening of plants in various seasons). Assessment and supervision of the work - GSL, SRLV, KMB, MSR, PAP, HKP, MSM, and RVS. Writing of original draft – GCG. Manuscript review and editing - PAP, KMB, HKP, MSM, and RVS. Funding acquisition – MSM, HKP, and RVS.

ACKNOWLEDGEMENTS

This study was supported by grants to MSM, HKP, and RVS from the Council of Scientific and Industrial Research (CSIR), Government of India (MLP0121 Phase-I and Phase-II) and the JC Bose Fellowship of RVS granted by the Science and Engineering Research Board (SERB), Department of Science and Technology, Government of India (GAP0444). We thank Dr. Rajkanwar Nathawat (Current address: Yale University) for her guidance and suggestions on protein structure prediction analysis. We thank the skilled workers at ICAR-IIRR and CSIR-CCMB for their support in field activities and maintenance.

CONFLICT OF INTEREST

The authors declare that no conflict of interest exists.

552 **Supporting Information:**

553 **Table S1: S**tatistics of the whole genome sequence data generated in this study.

554 **Table S2: S**tatistics of the RNA sequence data generated in this study.

555 **Table S3:** Overlapping insect resistance QTL intervals

556 **Table S4:** Laccase-encoding genes that are situated in Chromosome 1 QTL interval.
557

Figure S1: Functional profiling of genes present in the QTL intervals and carrying SNPs.

Figure S2: Differential expression of genes in SM and SM92.

Figure S3: Differential regulation of phenylpropanoid pathway genes.

Figure S4: Enrichment of lignin metabolism related genes in the QTL intervals.

Figure S5: Differentially expressed lipid-related genes and their phylogeny.

Figure S6: Other significantly enriched pathways and corresponding differentially expressed genes.

Figure S7: Variation in the regulatory region of *OsLTPL146* in SM and SM92.

Figure S8: Multiple phenylpropanoid pathway metabolites are accumulated in SM and SM92.
566

Data S1: Phenotype scores of F2 progenies

Data S2: List of genes present in all the identified QTL intervals, the SNPs and their effects.

Data S3: List of genes predicted to harbor high effect causing SNPs in the QTL regions.

Data S4: List of SNPs selected as markers for trait association study using KASP assays.

Data S5: KASP assay information

Data S6: Genotype-phenotype information of screened F2 progenies

Data S7: Marker-Trait Association Analysis of F2 progenies

Data S8: Differentially expressed genes in SM upon YSB infestation at 72hpi (infested vs
uninfested)

Data S9: Differentially expressed genes in SM92 upon YSB infestation at 72hpi (infested vs
uninfested)

Data S10: List of differentially expressed genes that are located within the QTL intervals and their
log2 fold change values.

Data S11: Log2 fold change values of genes upregulated upon striped stem borer and yellow stem
borer in respective tolerant and susceptible lines

Data S12: Raw peak intensity data of metabolites

Data S13: Normalized peak intensity data of metabolites

References:

Bentur JS, Padmakumari AP, Jhansilakshmi V, Padmavathi Ch, Kondala Rao Y, A.S. and P.I. (2011) *Insect resistance in Rice (2000-2009).*,.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M. and Gingeras, T.R. (2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*, **29**, 15–21. Available at: https://academic.oup.com/bioinformatics/article/29/1/15/272537 [Accessed April 23, 2023].

Dong, N.Q. and Lin, H.X. (2021) Contribution of phenylpropanoid metabolism to plant development and plant–environment interactions. *J. Integr. Plant Biol.*, **63**, 180–209. Available at: https://onlinelibrary.wiley.com/doi/full/10.1111/jipb.13054 [Accessed April 23, 2023].

Fauth, M., Schweizer, P., Buchala, A., Markstädter, C., Riederer, M., Kato, T. and Kauss, H. (1998) Cutin Monomers and Surface Wax Constituents Elicit H2O2 in Conditioned Cucumber Hypocotyl Segments and Enhance the Activity of Other H2O2 Elicitors. *Plant Physiol.*, **117**, 1373. Available at: /pmc/articles/PMC34901/ [Accessed October 14, 2022].

Franco, O.L., Rigden, D.J., Melo, F.R. and Grossi-de-Sá, M.F. (2002) Plant α-amylase inhibitors and their interaction with insect α-amylases. *Eur. J. Biochem.*, **269**, 397–412. Available at: https://onlinelibrary.wiley.com/doi/full/10.1046/j.0014-2956.2001.02656.x [Accessed October 14, 2022].

Gangwar, S.K., Chakraborty, S., Dasgupta, M.K. and Huda, A.K.S. (1986) MODELLING YIELD LOSS IN INDICA RICE IN FARMERS' FIELDS DUE TO MULTIPLE PESTS. *Ecosyst. Environ.*, **17**, 165–171.

Gao, H., Ma, K., Ji, G., Pan, L. and Zhou, Q. (2022) Lipid transfer proteins involved in plant–pathogen interactions and their molecular mechanisms. *Mol. Plant Pathol.* Available at: https://onlinelibrary.wiley.com/doi/full/10.1111/mpp.13264 [Accessed October 13, 2022].

Gesteiro, N., Butrón, A., Estévez, S. and Santiago, R. (2021) Unraveling the role of maize (Zea mays L.) cell-wall phenylpropanoids in stem-borer resistance. *Phytochemistry*, **185**, 112683.

Goujon, M., McWilliam, H., Li, W., Valentin, F., Squizzato, S., Paern, J. and Lopez, R. (2010) A new bioinformatics analysis tools framework at EMBL–EBI. *Nucleic Acids Res.*, **38**, W695–W699. Available at: https://academic.oup.com/nar/article/38/suppl_2/W695/1097251

[Accessed August 12, 2022].

Gururaj Katti, Chitra Shanker, Padmakumari A.P., P.I.C. (2011) Rice stem borers in India-

species composition and distribution. *Tech. Bull. No.59.*, 89.

Heinrichs, Elvis & Medrano, F.G. and Rapusas, H.R. (1985) *Genetic Evaluation for Insect*

Resistance in Rice., Available at: http://books.irri.org/9711041103_ content. Pdf.

Holm, L. (2022) Dali server: structural unification of protein families. *Nucleic Acids Res.*, **50**,

W210–W215.   Available   at: https://academic.oup.com/nar/article/50/W1/W210/6591528

[Accessed October 13, 2022].

IRRI (2014) *Standard evaluation system for rice. International Rice Research Institute*, Available

at: http://www.clrri.org/ver2/uploads/SES_5th_edition.pdf [Accessed July 31, 2020].

Jena, K.K., Jeung, J.U., Lee, J.H., Choi, H.C. and Brar, D.S. (2006) High-resolution mapping of

a new brown planthopper (BPH) resistance gene, Bph18(t), and marker-assisted selection for

BPH resistance in rice (Oryza sativa L.). *Theor. Appl. Genet.*, **112**, 288–297. Available at:

https://link.springer.com/article/10.1007/s00122-005-0127-8 [Accessed July 1, 2022].

Jumper, J., Evans, R., Pritzel, A., et al. (2021) Highly accurate protein structure prediction with

AlphaFold.   *Nat.*   *2021*   *5967873*,   **596**,   583–589.   Available   at:

https://www.nature.com/articles/s41586-021-03819-2 [Accessed October 14, 2022].

Kawahara, Y., la Bastide, M. de, Hamilton, J.P., et al. (2013) Improvement of the oryza sativa

nipponbare reference genome using next generation sequence and optical map data. *Rice*, **6**, 3–

10.   Available   at: https://thericejournal.springeropen.com/articles/10.1186/1939-8433-6-4

[Accessed August 12, 2022].

Kim, T.H., Park, J.H., Kim, M.C. and Cho, S.H. (2008) Cutin monomer induces expression of

the rice OsLTP5 lipid transfer protein gene. *J. Plant Physiol.*, **165**, 345–349. Available at:

https://linkinghub.elsevier.com/retrieve/pii/S0176161707001162 [Accessed August 11, 2020].

Kolattukudy, P.E., Rogers, L.M., Li, D., Hwang, C.S. and Flaishman, M.A. (1995) Surface

signaling in pathogenesis. *Proc. Natl. Acad. Sci. U. S. A.*, **92**, 4080. Available at:

/pmc/articles/PMC41890/?report=abstract [Accessed October 14, 2022].

Krueger, F. TrimGalore: A wrapper around Cutadapt and FastQC to consistently apply adapter and

quality trimming to FastQ files, with extra functionality for RRBS data. Available at:

https://github.com/FelixKrueger/TrimGalore [Accessed April 23, 2023].

Lage, J.-L. Da (2018) The Amylases of Insects. *Int. J. Insect Sci.*, **10**, 117954331880478. Available
at: /pmc/articles/PMC6176531/ [Accessed October 14, 2022].

Lee, M.-H., Jeon, H.S., Kim, S.H., et al. (2019) Lignin-based barrier restricts pathogens to the
infection site and confers resistance in plants. *EMBO J.*, **38**, e101948. Available at:
https://onlinelibrary.wiley.com/doi/full/10.15252/embj.2019101948 [Accessed October 13,
2022].

Letunic, I. and Bork, P. (2021) Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic
tree display and annotation. *Nucleic Acids Res.*, **49**, W293–W296. Available at:
https://academic.oup.com/nar/article/49/W1/W293/6246398 [Accessed August 12, 2022].

Liao, Y., Smyth, G.K. and Shi, W. (2014) featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics*, **30**, 923–930. Available at:
https://academic.oup.com/bioinformatics/article/30/7/923/232889 [Accessed April 23, 2023].

Love, M.I., Huber, W. and Anders, S. (2014) Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol.*, **15**, 1–21. Available at:
https://genomebiology.biomedcentral.com/articles/10.1186/s13059-014-0550-8 [Accessed
April 23, 2023].

M.D. Pathak and Z.R.Khan (1994) *Insect Pests of Rice*, International Rice Research Institute.
Available at: http://books.irri.org/9712200280_content.pdf [Accessed December 6, 2021].

Magwene, P.M., Willis, J.H. and Kelly, J.K. (2011) The Statistics of Bulk Segregant Analysis
Using Next Generation Sequencing. *PLOS Comput. Biol.*, **7**, e1002255. Available at:
https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1002255 [Accessed
October 23, 2022].

Malvar, R.A., Butrón, A., Ordás, B. and Santiago, R. (2008) Causes of natural resistance to stem
borers in maize. *Crop Prot. Res. Adv.*, **313**.

Mansfeld, B.N. and Grumet, R. (2018) QTLseqr: An R Package for Bulk Segregant Analysis with
Next□Generation Sequencing. *Plant Genome*, **11**, 180006. Available at:
https://onlinelibrary.wiley.com/doi/10.3835/plantgenome2018.01.0006 [Accessed December
25, 2020].

Miller, G.L. (1959) Use of Dinitrosalicylic Acid Reagent for Determination of Reducing Sugar.
Anal. Chem., **31**, 426–428. Available at: https://pubs.acs.org/doi/abs/10.1021/ac60147a030

[Accessed August 29, 2023].

Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S. and Steinegger, M. (2022)
ColabFold: making protein folding accessible to all. *Nat. Methods 2022 196*, **19**, 679–682.
Available at: https://www.nature.com/articles/s41592-022-01488-1 [Accessed August 12,
2022].

Moin, M., Bakshi, A., Saha, A., Dutta, M., Madhav, S.M. and Kirti, P.B. (2016) Rice ribosomal
protein large subunit genes and their spatio-temporal and stress regulation. *Front. Plant Sci.*, **7**,
1284.

Muralidharan, K. and Pasalu, I.C. (2006) Assessments of crop losses in rice ecosystems due to
stem borer damage (Lepidoptera: Pyralidae). *Crop Prot.*, **25**, 409417.

Nagaraj, S., Senthil-Kumar, M., Ramu, V.S., Wang, K. and Mysore, K.S. (2016) Plant
ribosomal proteins, RPL12 and RPL19, play a role in nonhost disease resistance against
bacterial pathogens. *Front. Plant Sci.*, **6**, 1192.

Oliveros, J.C. (2007) Venny. An interactive tool for comparing lists with Venn's diagrams.
Available at: https://bioinfogp.cnb.csic.es/tools/venny/index.html.

Oneda, H., Lee, S. and Inouye, K. (2004) Inhibitory Effect of 0.19 α-Amylase Inhibitor from
Wheat Kernel on the Activity of Porcine Pancreas α-Amylase and Its Thermal Stability. *J.*
Biochem., **135**, 421–427. Available at: https://academic.oup.com/jb/article/135/3/421/1072114
[Accessed October 14, 2022].

Padmakumari A.P, Seshu Madhav M, Uma Kanth B, Sunil B, Gopi, P, Karteek J, Laha GS,
Sundaram, RM , Subba Rao, LV, H.K.P. and R.S. (2017) *EMS mutants as a novel source of*
tolerance to yellow stem borer., Hyderabad.

Padmakumari AP and Katti G (2018) Designing insect bioassays for studying components of
insect pest management-rice yellow stem borer – as a case study. In *National Symposium*
Entomology2018:Advances and Challenges. Hyderabad, p. 117.

Pang, Z., Chong, J., Zhou, G., et al. (2021) MetaboAnalyst 5.0: narrowing the gap between raw
spectra and functional insights. *Nucleic Acids Res.*, **49**, W388–W396. Available at:
https://academic.oup.com/nar/article/49/W1/W388/6279832 [Accessed April 23, 2023].

Pathak, M.D. (1977) DEFENSE OF THE RICE CROP AGAINST INSECT PESTS. *Ann. N. Y.*
Acad.   Sci.,   **287**,   287–295.   Available   at:

https://onlinelibrary.wiley.com/doi/full/10.1111/j.1749-6632.1977.tb34247.x   [Accessed
December 6, 2021].

Pathak, M.D. (1968) Ecology of Common Insect Pests of Rice.
http://dx.doi.org/10.1146/annurev.en.13.010168.001353, **13**, 257–294. Available at:
https://www.annualreviews.org/doi/abs/10.1146/annurev.en.13.010168.001353   [Accessed
December 6, 2021].

Peltier, A.J., Hatfield, R.D. and Grau, C.R. (2009) Soybean Stem Lignin Concentration Relates
to Resistance to Sclerotinia sclerotiorum. *Plant Dis.*, **93**, 149–154. Available at:
https://pubmed.ncbi.nlm.nih.gov/30764097/ [Accessed October 13, 2022].

Potupureddi, G., Balija, V., Ballichatla, S., et al. (2021) Mutation resource of Samba Mahsuri
revealed the presence of high extent of variations among key traits for rice improvement. *PLoS*
One, **16**, e0258816. Available at:
https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0258816 [Accessed May 24,
2022].

Quinlan, A.R. and Hall, I.M. (2010) BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics*, **26**, 841–842. Available at:
https://academic.oup.com/bioinformatics/article/26/6/841/244688 [Accessed May 8, 2023].

Ramu, V.S., Dawane, A., Lee, S., Oh, S., Lee, H.K., Sun, L., Senthil-Kumar, M. and Mysore,
K.S. (2020) Ribosomal protein QM/RPL10 positively regulates defence and protein translation
mechanisms during nonhost disease resistance. *Mol. Plant Pathol.*, **21**, 1481–1494. Available
at: https://onlinelibrary.wiley.com/doi/full/10.1111/mpp.12991 [Accessed October 14, 2022].

Rubia, E.G., Heong, K.L., Zalucki, M., Gonzales, B. and Norton, G.A. (1996) Mechanisms of
compensation of rice plants to yellow stem borer Scirpophaga incertulas (Walker) injury. *Crop*
Prot., **15**, 335–340.

Sakai, H., Lee, S.S., Tanaka, T., et al. (2013) Rice Annotation Project Database (RAP-DB): an
integrative and interactive database for rice genomics. *Plant Cell Physiol.*, **54**. Available at:
https://pubmed.ncbi.nlm.nih.gov/23299411/ [Accessed August 12, 2022].

Salminen, T.A., Blomqvist, K. and Edqvist, J. (2016) Lipid transfer proteins: classification,
nomenclature, structure, and function. *Planta*, **244**, 971–997. Available at:
https://pubmed.ncbi.nlm.nih.gov/27562524/ [Accessed October 14, 2022].

Santiago, R., Barros-Rios, J. and Malvar, R.A. (2013) Impact of Cell Wall Composition on
Maize Resistance to Pests and Diseases. *Int. J. Mol. Sci. 2013, Vol. 14, Pages 6960-6980*, **14**,
6960–6980. Available at: https://www.mdpi.com/1422-0067/14/4/6960/htm [Accessed
October 13, 2022].

Santos, R.B., Nascimento, R., Coelho, A. V. and Figueiredo, A. (2020) Grapevine–Downy
Mildew Rendezvous: Proteome Analysis of the First Hours of an Incompatible Interaction.
Plants 2020, Vol. 9, Page 1498, **9**, 1498. Available at: https://www.mdpi.com/2223-
7747/9/11/1498/htm [Accessed October 14, 2022].

Savary, S., Willocquet, L., Pethybridge, S.J., Esker, P., McRoberts, N. and Nelson, A. (2019)
The global burden of pathogens and pests on major food crops. *Nat. Ecol. Evol.*, **3**, 430–439.
Available at: https://www.nature.com/articles/s41559-018-0793-y.

Schweizer, P., Jeanguenat, A., Whitacre, D., Métraux, J.P. and Mösinger, E. (1996) Induction
of resistance in barley againstErysiphe graminisf.sp.hordeiby free cutin monomers. *Physiol.*
Mol. Plant Pathol., **49**, 103–120.

Selvaraju, K., Shanmugasundaram, P., Mohankumar, S., Asaithambi, M. and
Balasaraswathi, R. (2007) Detection of quantitative trait locus for leaffolder (Cnaphalocrocis
medinalis (Guenée)) resistance in rice on linkage group 1 based on damage score and flag leaf
width. *Euphytica*, **157**, 35–43. Available at: https://link.springer.com/article/10.1007/s10681-
007-9394-6 [Accessed July 1, 2022].

Sievers, F., Wilm, A., Dineen, D., et al. (2011) Fast, scalable generation of high-quality protein
multiple sequence alignments using Clustal Omega. *Mol. Syst. Biol.*, **7**, 539. Available at:
https://onlinelibrary.wiley.com/doi/full/10.1038/msb.2011.75 [Accessed August 12, 2022].

Su, J., Zhang, Z., Wu, M. and Gao, C. (2014) Geographic susceptibility of Chilo suppressalis
Walker (Lepidoptera: Crambidae), to chlorantraniliprole in China. *Pest Manag. Sci.*, **70**, 989–
995. Available at: https://onlinelibrary.wiley.com/doi/full/10.1002/ps.3640 [Accessed October
13, 2022].

Sun, L., Su, C., Wang, C., Zhai, H. and Wan, J. (2005) Mapping of a Major Resistance Gene to
the Brown Planthopper in the Rice Cultivar Rathu Heenati. *Breed. Sci.*, **55**, 391–396. Available
at: http://www. [Accessed July 1, 2022].

Tae, H.K., Moon, C.K., Jong, H.P., Seong, S.H., Byung, R.K., Byoung, Y.M., Mi, C.S. and

Sung, H.C. (2006) Differential expression of rice lipid transfer protein gene(LTP) classes in
765   response to abscisic acid, salt, salicylic acid, and the fungal pathogenMagnaporthe grisea. *J.*
766   *Plant Biol. 2006 495*, **49**, 371–375. Available at:
767   https://link.springer.com/article/10.1007/BF03178814 [Accessed October 14, 2022].

768   **Takagi, H., Abe, A., Yoshida, K., et al.** (2013) QTL-seq: Rapid mapping of quantitative trait loci
769   in rice by whole genome resequencing of DNA from two bulked populations. *Plant J.*, **74**,
770   174–183. Available at: https://onlinelibrary.wiley.com/doi/full/10.1111/tpj.12105 [Accessed
771   July 29, 2020].

772   **Thimm, O., Bläsing, O., Gibon, Y., et al.** (2004) MAPMAN: a user-driven tool to display
773   genomics data sets onto diagrams of metabolic pathways and other biological processes. *Plant*
774   *J.*, **37**, 914–939. Available at: https://pubmed.ncbi.nlm.nih.gov/14996223/ [Accessed April 23,
775   2023].

776   **Trautmann, H.S. and Ramsey, K.M.** (2022) A Ribosomal Protein Homolog Governs Gene
777   Expression and Virulence in a Bacterial Pathogen. *J. Bacteriol.* Available at:
778   https://journals.asm.org/doi/10.1128/jb.00268-22 [Accessed October 14, 2022].

779   **Wang, Y., Ju, D., Yang, X., Ma, D. and Wang, X.** (2018) Comparative Transcriptome Analysis
780   Between Resistant and Susceptible Rice Cultivars Responding to Striped Stem Borer (SSB),
781   Chilo suppressalis (Walker) Infestation. *Front. Physiol.*, **9**, 1717.

782   **Warner, J.R. and McIntosh, K.B.** (2009) How Common Are Extraribosomal Functions of
783   Ribosomal Proteins? *Mol. Cell*, **34**, 3–11.

784   **Weisberg, R.A.** (2008) Transcription by Moonlight: Structural Basis of an Extraribosomal Activity
785   of Ribosomal Protein S10. *Mol. Cell*, **32**, 747–748.

786   **Wool, I.G.** (1996) Extraribosomal functions of ribosomal proteins. *Trends Biochem. Sci.*, **21**, 164–
787   165.

788   **Yu, Y., Xing, Y., Liu, F., Zhang, X., Li, X., Zhang, J. and Sun, X.** (2021) The Laccase Gene
789   Family Mediate Multi-Perspective Trade-Offs during Tea Plant (Camellia sinensis)
790   Development and Defense Processes. *Int. J. Mol. Sci. 2021, Vol. 22, Page 12554*, **22**, 12554.
791   Available at: https://www.mdpi.com/1422-0067/22/22/12554/htm [Accessed October 13,
792   2022].

793   **Zhao, Q., Nakashima, J., Chen, F., et al.** (2013) LACCASE Is Necessary and Nonredundant with

794 PEROXIDASE for Lignin Polymerization during Vascular Development in Arabidopsis. *Plant*

S.No.	Chromosome	Start[a]	End[b]	Length[c]	Number of SNPs[d]	Mean Q-val
1	1	33465932	36594124	3128192	741	0.025
2		37253341	37887039	633698	166	0.033
3	5	1931441	2666236	734795	268	0.030
4		3360636	3839012	478376	134	0.042
5	7	2039475	2340432	300957	94	0.035
6	10	21348600	22786612	1438012	719	0.013

795 *Cell*, **25**, 3976. Available at: /pmc/articles/PMC3877815/ [Accessed
812 October 13, 2022].

Table 1: List of the putative QTL predicted to be associated with YSB tolerance. SNPs - Single Nucleotide Polymorphisms; Q-val - False Discovery Rate.

7		7703041	10065030	2361989	1979	0.034
8	12	16700754	18048717	1347963	1026	0.030
9		19115124	19867810	752686	825	0.043
10		22145804	23226149	1080345	529	0.031

[a] Beginning of the mapped QTL interval
[b] End of the mapped QTL interval
[c] Length of the mapped QTL interval in base pairs
[d] Number of SNPs situated in the mapped QTL interval

845 **Table 2:** Markers that are significantly associated with YSB tolerance.

Marker	Chromosome	Position	REF[a]	ALT[b]	SNP-index Tol[c]	SNP-index Sus[d]	Delta SNP-index	P	%PVE	Gene ID	Description
S2	1	33886610	C	G	0	0.61	-0.61	0.034	5.7	Os01g0800266	Similar to Phosphoglycerate kinase. (Os01t0800266-00)
S3		33957364	G	A	0.29	0.62	-0.33	0.022	6.7	Os01g0801700	Ribosomal protein S35, mitochondrial domain containing protein. (Os01t0801700-01)
S5		34710185	G	A	0	0.63	-0.63	0.007	9.2	Os01g0816400	Similar to microtubule organization protein. (Os01t0816400-01)
S6		35088627	C	T	0	0.69	-0.69	0.019	7.1	Non-exonic[†]	-
S17	5	2366109	G	A	0.65	0	0.65	0.047	5.0	Os05g0141200	Hypothetical conserved gene. (Os05t0141200-00)
S18		3157367	C	T	0.59	0	0.59	0.033	5.8	Non-exonic	-
S19		3743040	G	T	0.91	0	0.91	0.022	6.6	Non-exonic	-
S20		4321188	C	T	0.66	0	0.66	0.047	5.1	Os05g0172300	Domain of unknown function DUF632 domain containing protein. (Os05t0172300-01)
S28	10	21786052	C	T	0.73	0	0.73	0.017	7.2	Non-exonic	-
S29		21853916	T	A	0.82	0	0.82	0.017	7.2	Non-exonic	-
S30		21955122	C	T	0.68	0	0.68	0.017	7.2	Os10g0557800	Conserved hypothetical protein. (Os10t0557800-01)
S31		21955776	T	A	0.76	0.45	0.31	0.038	5.5	Os10g0557900	Peptidase M10A and M12B, matrixin and adamalysin family protein. (Os10t0557900-00)
S32		21965877	C	G	0.69	0.38	0.31	0.017	7.2	Os10g0558125	Similar to Matrixin family protein, expressed. (Os10t0558125-00)
S45	12	18013596	A	T	0.63	0	0.63	0.034	5.8	Non-exonic	-

846
[a] Allele present in the wildtype line
[b] Allele present in the mutant line
[c] SNP-index of the tolerant bulk
[d] SNP-index of the susceptible bulk
[*] Control – SNPs from genomic regions outside QTL intervals
[†] Non-exonic – SNPs from the QTL intervals that are not exonic
%PVE - % Phenotypic Variation Explained
854

Figure 1

SM92 exhibited a high level of tolerance to YSB infestation. (A) Field screening with YSB

larvae revealed an increased incidence of dead hearts in SM as compared to SM92. (B) Violin plot

showing the number of live and dead YSB larvae in the infested stems of SM and SM92. The

number of live larvae were significantly less in the stems of SM92 as compared to SM. Asterisk

indicates *P*-value of less than 0.05 as calculated using a 2-way ANOVA test. (C) and (D) Images

indicating the presence of a live larvae in SM and a dead larva in SM92, respectively. Black

arrowheads denote the larvae.

Figure 2

Mapping genomic loci associated with YSB tolerance in F2 population using BSA-QTL-seq.

(A) Phenotype scores of the F_2 population consisting of 314 progenies followed a continuous

distribution suggesting the quantitative nature of YSB tolerance. The green bar indicates progenies

that constituted tolerant bulk (n=21) and the red bars indicate the progenies that constituted

susceptible bulk (n=24). (B) Delta SNP-index plot generated by the QTL-seq pipeline shows two

genomic loci (red arrowheads) that could be linked to YSB tolerance. The red, green, and blue lines

represent the confidence intervals at 90%, 95%, and 99%, respectively. (C) G' plot generated by the

QTL-seq pipeline shows additional genomic loci (blue arrowheads) that might be associated with

YSB tolerance. The red line represents the q-value cut-off of 0.05. The black line in (B) and (C)

represents the moving average line calculated using a window size of 500kb.

Figure 3

Transcriptome profiling revealed putative rice functions associated with YSB tolerance. (A)

Bar chart depicting the number of differentially expressed genes in SM and SM92 upon YSB

infestation in comparison to uninfested samples, as obtained through RNA-sequencing based

transcriptome profiling of stem samples at 72 hours post infestation. (B) Venn diagram comparing

the differentially expressed genes in SM and SM92. (C) and (D) Gene enrichment analysis showing

the upregulated pathways in SM and SM92 post YSB infestation.

Figure 4

Comparative transcriptome indicated a possible candidate gene for YSB tolerance. (A) Venn

diagram showing the comparison of genes upregulated in rice lines that are tolerant to striped stem

borer (SSB) from Wang et al (2018) and yellow stem borer (YSB). The intersection indicates 527

genes that are commonly upregulated. (B) Gene enrichment analysis of the stem borers-induced
genes revealed the enrichment of functions involved in alpha-amylase inhibition, glycoproteins, and
secreted proteins. (C) Expression of the gene *OsLTPL16* was observed to be upregulated by SSB
and YSB in tolerant rice lines and downregulated in susceptible rice lines. (D) Structure prediction
and homology search suggests that *OsLTPL146* might encode an alpha-amylase inhibitor. Structure
model of OsLTPL146 is colored green and the solved structure of wheat alpha-amylase inhibitor
(PDB:1hss) is colored brown.

Figure 5

YSB infestation causes considerable changes in the plant metabolite composition. (A) Principal
Component Analysis (PCA) plot showing the clustering of SM and SM92 samples infested with
YSB (pink- and purple-shaded oval) indicating an overlapping response in both SM and SM92. The
uninfested samples, however, showed a loose clustering, probably indicating differences in the
basal metabolite composition. (B) Box plot showing the normalized intensity of certain
phenylpropanoid pathway metabolites in SM and SM92 upon YSB infestation and uninfested
conditions. *P*-values were calculated using unpaired t-test with multiple comparison correction
using Holm-Sidak method. P<0.05 is considered to be statistically significant.

919   34

A

A

B