Robust estimation of cancer and immune cell-type proportions from bulk tumor ATAC-Seq data.

Aurélie AG Gabriel[1,2,3,4], Julien Racle[1,2,3,4], Maryline Falquet[3,5,6,7], Camilla Jandus[3,5,6,7], David Gfeller[1,2,3,4,*]

Affiliations:

[1] Department of Oncology, Ludwig Institute for Cancer Research, University of Lausanne, Lausanne, Switzerland

[2] Agora Cancer Research Centre, Lausanne, Switzerland

[3] Swiss Cancer Center Leman (SCCL), Switzerland

[4] Swiss Institute of Bioinformatics (SIB), Lausanne, Switzerland.

[5] Ludwig Institute for Cancer Research, Lausanne Branch, Lausanne, Switzerland

[6] Department of Pathology and Immunology Faculty of Medicine, University of Geneva, Geneva, Switzerland

[7] Geneva Center for Inflammation Research, Geneva, Switzerland

* Corresponding author: david.gfeller@unil.ch

Abstract

Assay for Transposase-Accessible Chromatin sequencing (ATAC-Seq) is a widely used technique to explore gene regulatory mechanisms. For most ATAC-Seq data from healthy and diseased tissues such as tumors, chromatin accessibility measurement represents a mixed signal from multiple cell types. In this work, we derive reliable chromatin accessibility marker peaks and reference profiles for all major cancer-relevant cell types. We then integrate these data into the EPIC deconvolution framework to quantify cell-type heterogeneity in bulk ATAC-Seq data. Our EPIC-ATAC tool accurately predicts non-malignant and malignant cell fractions in tumor samples. When applied to a breast

25 cancer cohort, EPIC-ATAC accurately infers the immune contexture of the main breast cancer

26 subtypes.

Introduction

Gene regulation is a dynamic process largely determined by the physical access of chromatin-binding

factors such as transcription factors (TFs) to regulatory regions of the DNA (*e.g.,* enhancers and

promoters) (1). The genome-wide landscape of chromatin accessibility is essential in the control of

cellular identity and cell fate and thus varies in different cell types (1,2). Over the last decade, Assay

for Transposase-Accessible Chromatin (ATAC-Seq) (3) has become a reference epigenomic technique

to profile chromatin accessibility and the activity of gene regulatory elements in diverse biological

contexts including cancer (4) and across large cohorts (5). Several optimized ATAC-seq protocols have

been developed to improve the quality of ATAC-Seq data and expand its usage to different tissue

types. These include the OMNI-ATAC protocol, which leads to cleaner signal and is applicable to

frozen samples (6,7), as well as the formalin-fixed paraffin-embedded (FFPE)-ATAC protocol adapted

to FFPE samples. The reasonable cost and technical advantages of these protocols foreshadow an

increased usage of ATAC-Seq in cancer studies.

Most biological tissues are composed of multiple cell types. For instance, tumors are complex

ecosystems including malignant and stromal cells as well as a large diversity of immune cells. This

cellular heterogeneity, in particular the presence of specific immune cell types, impacts tumor

progression as well as response to immunotherapy (8–10). Most existing ATAC-Seq data from tumors

were performed on bulk samples, thereby including information from both cancer and non-

malignant cells. Precisely quantifying the proportions of different cell types in such samples

represents therefore a promising way to explore the immune contexture and the composition of the

tumor micro-environment (TME) across large cohorts. Carefully assessing cell-type heterogeneity is

also important to handle confounding factors in genomic analyses in which samples with different

cellular compositions are compared. Recently, single-cell ATAC-Seq (scATAC-Seq) has been developed

to explore cellular heterogeneity with high resolution in complex biological systems (11–13).

However, the resulting data are sensitive to technical noise and such experiments require important

resources, which so far limits the use of scATAC-Seq in contrast to bulk sequencing in the context of

large cohorts.

In the past decade, computational deconvolution tools have been developed to predict the

proportion of diverse cell types from bulk genomic data obtained from tumor samples (14–22). A

large number of these tools model bulk data as a mixture of reference profiles identified in purified

cell populations for each cell type. The accuracy of the predictions of cell-type proportions relies on

the quality of these reference profiles as well as on the use of cell-type specific markers (14). A

limitation of most deconvolution algorithms is that they do not predict the proportion of cell types

that are not present in the reference profiles (here referred to as 'uncharacterized' cells). In the

context of cancer samples, these uncharacterized cell populations include malignant cells whose

molecular profiles differ not only from one cancer type to another, but also from one patient to

another even within the same tumor type (5). A few tools consider uncharacterized cells in their

deconvolution framework by using cell-type specific markers not expressed in the uncharacterized

cells (17,21,23,24). These tools include EPIC (Estimating the Proportion of Immune and Cancer cells)

which simultaneously quantifies immune, stromal, vascular as well as uncharacterized cells from bulk

tumor samples (17,25).

Most deconvolution algorithms have been initially developed for transcriptomic data (RNA-Seq data)

(17–19,21,26–29). More recently they have been adapted for other omics layers such as methylation

(30–33) and proteomics (34) or chromatin accessibility. For the latter, a specific framework called

DeconPeaker (20) was developed to estimate cell-type proportions from bulk samples.

Deconvolution tools developed initially for other omics modalities, such as RNA-Seq, can also be

applied on ATAC-Seq if appropriate ATAC-Seq profiles are provided to the tool. For example, the

popular deconvolution tool, CIBERSORT (26), was used to deconvolve leukemic ATAC-Seq samples

(35). Other methods have been proposed to decompose ATAC-Seq bulk profiles into subpopulation-

specific profiles (36,37) or compartments (38). However, these methods have more requisites: (i) the

integration of the ATAC-Seq data with single-cell or bulk RNA-Seq (36,37) and HIChIP data (36) or, (ii)

subsequent feature annotation to associate compartments with cell types or biological processes

(38).

The application of existing bulk ATAC-Seq data deconvolution tools to solid tumors is limited. First,

current computational frameworks do not quantify populations of uncharacterized cell types.

Second, ATAC-Seq based markers (*i.e.,* chromatin accessible regions called peaks) and reference

profiles generated so far have been derived in the context of hematopoietic cell mixtures (20,35).

Markers and profiles for major populations of the TME (*e.g.,* stromal and vascular cells) are thus

missing. While cell-type specific markers have been identified from scATAC-Seq data (2), not all TME-

relevant cell types are covered (*e.g.*, lack of scATAC-Seq data from neutrophils due to extracellular

traps formation). Also, these markers have not been curated to fulfill the requirements of tools such

as EPIC to quantify uncharacterized cells (*i.e.*, markers of a cell-type should not be accessible in other

human tissues).

In this study, we collected ATAC-Seq data from pure cell types to identify cell-type specific

marker peaks and to build reference profiles from most major non-malignant cell types typically

observed in tumors. These data were integrated in the EPIC (17) framework to perform bulk ATAC-

Seq samples deconvolution (Figure 1). Applied on peripheral blood mononuclear cells (PBMCs) and

tumor samples, the EPIC-ATAC framework showed accurate predictions of the proportions of non-

malignant and malignant cells with similar or higher performances than other existing tools.

Results

ATAC-Seq data from sorted cell populations reveal cell-type specific marker peaks and reference profiles

A key determinant for accurate predictions of cell-type proportions by most deconvolution tools is the availability of reliable cell-type specific markers and reference profiles. To identify robust chromatin accessibility marker peaks of cancer relevant cell types, we collected 564 samples of sorted cell populations from twelve studies including eight immune cell types (B cells (35,39,40), CD4+ T cells (35,40–43), CD8+ T cells (35,39–41,43), natural killer (NK) cells (35,39), dendritic cells (DCs) (39,41,44), macrophages (40,41), monocytes (35,39,40,44,45) and neutrophils (46,47), as well as fibroblasts (41,48) and endothelial (41,49) cells (Figure 1 box 1, Figure 2A, Supplementary Table 1). To limit batch effects, the collected samples were homogeneously processed from read alignment to peak calling. For each cell type, we derived a set of stable peaks, *i.e.,* peaks observed across samples and studies (see Materials and Methods).

These peaks were then used to perform pairwise differential analysis to identify marker peaks for each cell type (Figure 1, box 2). To ensure that the cell-type specific marker peaks are not accessible in other human tissues, we included in the differential analysis ATAC-Seq samples from diverse human tissues from the ENCODE data (50,51) (Supplementary Figure 1). To select a sufficient number of peaks prior to peak filtering, the top 200 peaks recurrently differentially accessible across all cell-type pairs were selected as cell-type specific markers (see Materials and Methods). Using the human atlas study (2), markers with potential residual accessibility in human tissues were then filtered out (Figure 1, box 3, see Materials and Methods). The resulting marker peaks specific to the immune cell types were considered for the deconvolution of PBMC samples (PBMC markers). For tumor bulk sample deconvolution, the list of markers was further refined based on the correlation patterns of the markers in tumor bulk samples from diverse cancer types from The Cancer Genome Atlas (TCGA) (5) (Figure 1, box 4, see the Material and methods). The latter filtering ensures the relevance of the markers in the TME context since cell-type specific TME markers are expected to be correlated in

tumor bulk ATAC-Seq measurements (52). 716 markers of immune, fibroblasts and endothelial cell

types remained after the later filtering and were considered for the deconvolution of bulk tumor

samples (TME markers).

To assess the quality and reproducibility of these markers, we performed principal component

analysis (PCA) based on each set of marker peaks. Computing silhouette coefficients based on the

cell-type classification and on the study of origin showed that samples clustered by cell type and not

by study of origin (averaged silhouette coefficients above 0.45 for cell type and around 0 for study of

origin). Two-dimensional UMAP representations of the samples confirmed this observation (Figure

2B). These results indicate limited remaining batch effects after data processing and marker

selection.

We then used the collected samples to generate chromatin accessibility profiles by computing the

average of the normalized counts for each peak in each cell type as well as peak variability in each

cell type (17) (see Material and methods). Figure 2C represents the average chromatin accessibility of

each marker peak in each cell type of the reference dataset and highlights, as expected, the cell-type

specificity of the selected markers (see also Supplementary Tables 2 and 3), which was confirmed in

independent ATAC-Seq data from sorted cells and single-cell ATAC-Seq samples from blood and

diverse human tissues (Figure 2D and 2E, see Materials and methods).

Annotations of the marker peaks highlight their biological relevance

To characterize the different marker peaks, we annotated them using ChiPSeeker (53). We observed

that most of the markers are in distal and intergenic regions (Figure 2F), which is expected

considering the large proportion of distal regions in the human genome and the fact that such

regions have been previously described as highly cell-type specific (35). We also noticed that 7% of

the PBMC and TME marker peaks are in promoter regions in contrast to 4% when considering

matched genomic regions randomly selected in the set of peaks identified prior to the differential

analysis (see Material and methods), which suggest enrichment in our marker peaks for important

regulatory regions.

To assess the biological relevance of the marker peaks, we associated each marker peak to its

nearest gene using ChIP-Enrich based on the "nearest transcription start site (TSS)" locus definition

(54) (Supplementary Tables 4 and 5). Nearest genes reported as known marker genes in public

databases of gene markers (*i.e.,* PanglaoDB (55) and CellMarker (56)) are listed in Table 1.

In each set of cell-type specific peaks, we observed an overrepresentation of chromatin binding

proteins (CBPs) reported in the JASPAR2022 database (57) (using Signac (58) and MonaLisa (59) for

assessing the overrepresentation) and the ReMap catalog (60) (using RemapEnrich, see Material and

Methods). Overrepresented CBPs also reported as known marker genes in the PanglaoDB and

CellMarker databases are listed in Table 1. Detailed peaks annotations are summarized in

Supplementary Tables 4 and 5.

Based on the "nearest TSS" annotation, we tested, using ChIP-Enrich (54), whether each set of cell-

type specific marker peaks was enriched for regions linked to specific biological pathways (GO

pathways). Figure 2G highlights a subset of the enriched pathways that are consistent with prior

knowledge on each cell type. Some of these pathways are known to be characteristic of immune

responses to inflammatory or tumoral environments. The complete list of enriched pathways is listed

in the Supplementary Tables 6 and 7. Overall, these analyses demonstrate that the proposed cell-

type specific marker peaks capture some of the known biological properties associated to each cell

type.

Cell type	Nearest genes	Enriched CPBs
Bcells	DHTKD1 LHPP WDFY4 ARID5B HHEX SIDT2 CD82 MS4A1 FCHSD2 USP8 RHCG ATF7IP2 CIITA GGA2 SNX29P2 C16orf74 CBFA2T3 CD79B BCL2 GNG7 CD22 FCER2 FCRL1 LY9 PTPRC LAPTM5 IGLL5 VPREB3 CENPM AFF3 SP100 INPP5D DTNB CD86 RFTN1 ST6GAL1 NGLY1 OSBPL10 TLR9 CD38 SMIM14 ARHGAP24 ADAM19 EBF1 BASP1 CD83 PLEKHG1 CCR6 CCND3 HDAC9 CDCA7L BLK MTSS1 LYN PLEKHF2 MOB3B PAX5	SPIB POU2F2 TCF4 EBF1 TCF3 NFKB1 STAT1 NFKB2 IKZF1 FOXO1 FOXP1 BCL6 POU2AF1 STAT3 BACH2 IKZF3 FLI1 TBX21 JUNB MITF NKX6-2 RBPJ
CD4_Tcells	IL2RA CD6 CD5 CD4 RORA PTPRC CTLA4 ICOS SLC9A9 FHIT TCF7 FYB1 ATXN1 CD40LG	TCF7 RUNX3 SOHLH2 IRF9 GATA3 TBX21 MAF STAT3 RORA BATF CREM

CD8_Tcells	MKI67 JAML MAML2 KLRD1 NELL2 LAG3 PPP1R13B PTPRC LYST CASP8 CD8A CD8B CD96 BTLA GZMA THEMIS ETV1	ETV1 FOXP3 TBX21 FOXP1 EOMES CREM IRF4 ZEB1 ARNT JUNB TCF7
NK	PRF1 ZBTB16 KLRD1 SPN CD226 SH2D1B CD247 IL2RB CXCR4 NMUR1 GNLY ZAP70 TXK	EOMES TBX21 NFIL3 FOS JUN
DCs	C12orf75 LYZ APP CD8A RIOX2 NFKB1 QDPR ABCG2 PRELID2 DST CD36 IDO2 PCMTD1	SPIB IRF8 MYB NR4A1 REL CUX2 FOXO1 ETV6 IRF5 BATF3 RUNX2
Neutrophils	TLE3 CA4 CYP4F3 CEACAM8 PGLYRP1 FPR1 CTSS ALPL PI3 MMP9 CXCR1 DRC1 ASPRV1 LTF MGAM SLC25A37	FOS
Monocytes	VENTX GLT1D1 CLEC4E CARS2 SLC24A4 C16orf74 FFAR2 STXBP2 NLRP3 CYRIA CMTM7 TGFBI DIAPH1 VCAN MCTP1 IFNGR1 STX11 CAPZA2 CD36 MTSS1 DENND3 ASAH1 TNFRSF10B BNIP3L NACC2 MAMDC2 FBP1	CEBPA CEBPD CEBPB CEBPE SPI1 VENTX JUND RXRA TCF7L2
Macrophages	CXCL12 PSAP P2RY6 SLCO2B1 CMKLR1 MMP19 LGMN CLEC10A C5AR1 FPR3 LILRB4 RGL1 SIGLEC1 MMP9 CD80	STAT1 SPI1 FOSL2 FOS SPIC
Endothelial	FAM107B ROBO4 FLI1 ACVRL1 FLT1 DOCK9 ABCC1 S1PR1 ELOVL1 PLPP3 ASAP2 SNRK ECSCR ARAP3 LAMA4 BMP6 SERPINE1 LAMB1 DOCK4 NOS3	ETV2 ELF1 FLI1 ELK3 FOSB ETS1 ERG GATA2 ZEB1 ETS2 FOXC1 SOX18
Fibroblasts	LOX CAV1 COL15A1	FOSL2 FOSB FLI1 HIF1A PBX1

Table 1: List of nearest genes and enriched CBPs reported in the PanglaoDB or CellMarker databases.

EPIC-ATAC accurately estimates immune cell fractions in PBMC ATAC-Seq samples

The cell-type specific marker peaks and profiles derived from the reference samples were integrated

to the EPIC deconvolution tool (17,25). We will refer to this ATAC-Seq deconvolution framework as

EPIC-ATAC.

To test the accuracy of EPIC-ATAC predictions, we first collected PBMCs from five healthy donors. In

each donor, half of the cells was used to generate a bulk ATAC-Seq dataset and the other half was

used to determine the cellular composition of each sample, *i.e.,* the proportions of monocytes, B

cells, CD4+ T cells, CD8+ T cells, NK cells and dendritic cells, by multiparametric flow cytometry

(Figure 3A, see Materials and methods). We then applied EPIC-ATAC to the bulk ATAC-Seq data. The

predicted cell fractions are consistent with the cell fractions obtained by flow cytometry (Figure 3B,

Pearson correlation coefficient of 0.78 and root mean squared error (RMSE) of 0.10).

As a second validation, we applied EPIC-ATAC to pseudo-bulk PBMC samples (referred to as the

PBMC pseudobulk dataset, generated using three publicly available PBMC scATAC-Seq datasets

(13,61,62), see Material and methods). A high correlation (0.91) between EPIC-ATAC predictions and

true cell-type proportions and a low RMSE (0.05) were observed for this dataset (Figure 3C).

The accuracy of the predictions obtained with EPIC-ATAC was then compared with the accuracy of

other deconvolution approaches which could be used with our reference profiles and marker peaks

(Figure 3D-E). To this end, we considered both the DeconPeaker method (20) originally developed for

bulk ATAC-Seq as well as several algorithms developed for bulk RNA-Seq (CIBERSORTx (19),

QuanTIseq (21), ABIS (18), and MCPcounter (22)). To enable meaningful comparison across the cell

types considered in this work and use the method initially developed for bulk RNA-Seq

deconvolution, the marker peaks and profiles derived in this work were used in each of these

methods. DeconPeaker and CIBERSORTx include the option to define cell-type specific markers and

profiles from a set of reference samples. We thus fed our ATAC-Seq samples collection to both

algorithms and used the resulting profiles and marker peaks to perform bulk ATAC-Seq

deconvolution. The resulting predictions are referred to as DeconPeaker-Custom and CIBERSORTx-

Custom.

Many tools displayed high correlation and low RMSE values, similar to those of EPIC-ATAC, and no

single tool consistently outperformed the others (Figure 3D-E, Supplementary Figure 2A-C). The fact

that our marker peaks and reference profiles could be used with EPIC-ATAC and other existing tools

demonstrates their broad applicability.

Predictions accuracies were also evaluated in each cell type separately. Since the number of samples

was low in each dataset, samples from both datasets were combined for this analysis. EPIC-ATAC

demonstrated good accuracies across cell types with RMSE values ranging from 0.02 for B cells to

0.13 for NK cells (Supplementary Figure 3). As expected, predictions with all tools were more

accurate for frequent cell types with well-characterized markers (*e.g.,* CD8/CD4 T cells, B cells)

compared to less frequent cell types (*e.g.,* NK cells, dendritic cells) (Supplementary Figure 2 and 3).

Note that MCPcounter is a marker-based method that derives cell-type specific scores which cannot

215 be compared between cell types. This method was thus only included in the benchmark considering

216 each cell type separately.

218 **EPIC-ATAC accurately predicts fractions of cancer and non-malignant cells in tumor**

219 **samples**

220 We evaluated the ability of the EPIC-ATAC framework to predict not only immune and stromal cells

221 proportions but also the proportion of cells for which reference profiles are not available (*i.e.,*

222 uncharacterized cells). For this purpose, we considered two previously published scATAC-Seq

223 datasets containing basal cell carcinoma and gynecological cancer samples (13,63). We generated

224 two pseudobulk datasets by averaging the chromatin accessibility signal across all cells of each

225 sample (see Material and methods). Applying EPIC-ATAC to both datasets shows that this framework

226 is able to simultaneously predict the proportions of both uncharacterized cells and immune, stromal

227 and vascular cells (Figure 4A). In these cancer samples, the proportion of uncharacterized cells can be

228 seen as a proxy of the proportion of cancer cells.

229 As for the PBMC datasets, we compared EPIC-ATAC performances to other existing deconvolution

230 tools. For both datasets, EPIC-ATAC led to the highest performances and was the only method to

231 accurately predict the proportion of uncharacterized cells (Figure 4B, Supplementary Figure 4 and 5).

232 Although quanTIseq also allows users to perform such predictions, the method resulted in lower

233 correlation and higher RMSE values when comparing the estimated and true proportions of the

234 uncharacterized cells (Figure 4B, Supplementary Figure 4).

235 In the EPIC-ATAC and quanTIseq frameworks, predictions correspond to absolute cell-type fraction,

236 *i.e.,* proportions of all cells present in the bulk, while the estimations obtained from the other tools

237 correspond to relative cell fractions, *i.e.,* proportions of cells present in the reference profiles

238 (CIBERSORTx, DeconPeaker) or to scores with arbitrary units (ABIS, MCPcounter). We thus conducted

239 a second benchmark excluding the predictions of uncharacterized cell fractions and rescaling both

240 estimations and true proportions to sum to 1 (see Material and methods). EPIC-ATAC outperformed

most of the other methods also when excluding the uncharacterized cells (Figure 4C, Supplementary Figure 4 and 5).

Supplementary Figure 6 reports the performances of each tool when considering each cell type separately. Overall, EPIC-ATAC showed comparable or higher correlation and lower RMSE values when compared to the other deconvolution tools.

T cell subtypes quantification reveals the ATAC-Seq deconvolution limits for closely related cell types.

To explore the limitations of ATAC-Seq deconvolution, we next evaluated whether EPIC-ATAC could predict the proportions of T-cell subtypes. To this end, we considered naive and non-naive CD8+ as well as naïve, helper/memory and T regulatory CD4+ T cells. We redefined our list of cell-type specific marker peaks and reference profiles including also these five T-cell subtypes (Supplementary Tables 8-9, Supplementary Figure 7A) and observed that the markers were conserved in external data (Supplementary Figure 7B). The annotations of the markers associated to the T-cell subtypes are available in Supplementary Tables 10-13.

We capitalized on the more detailed cell-type annotation of the PBMC datasets as well as the basal cell carcinoma dataset to evaluate the EPIC-ATAC prediction of cell-subtype fractions using these updated markers and profiles. Overall, the correlations observed between the predictions and true proportions of T cells decreased when considering T-cell subtypes rather than CD4+ and CD8+ cell types only (Figure 5A). In particular, low accuracies were obtained for helper/memory CD4+ and naïve T-cell subtypes (Figure 5B). Similar results were obtained using other deconvolution tools (Supplementary Figure 8).

EPIC-ATAC accurately infers the immune contexture in a bulk ATAC-Seq breast cancer cohort

We applied EPIC-ATAC to a breast cancer cohort of 42 breast ATAC-Seq samples including samples

from two breast cancer subtypes, *i.e.,* 35 oestrogen receptor (ER)-positive human epidermal growth

factor receptor 2 (HER2)-negative (ER+/HER2-) samples and 7 triple negative (TN) tumors (64). No cell

sorting was performed in parallel to the chromatin accessibility sequencing. We thus used EPIC-ATAC

to estimate cell-type proportions. We observed a higher proportion of T cells, B cells, NK cells and

macrophages in the TN samples in comparison to ER+/HER2- samples (Figure 6A). We then compared

the cellular composition of ER+/HER2- subgroups identified in the original study (clusters CA-A, CA-B

and CA-C). A higher infiltration of T and B cells was observed in cluster CA-C and higher proportions

of endothelial cells and fibroblasts were observed in cluster CA-B (Figure 6B). These predictions are

consistent with the infiltration level estimations reported in the original publication, although no

differences in macrophages infiltration was observed between the ER+/HER2- subgroups in our case

(64).

EPIC-ATAC performs similarly to EPIC RNA-seq based deconvolution and better than gene

activity based deconvolution

We finally compared the accuracy of EPIC when applied on ATAC-Seq data and on RNA-Seq data. For

this purpose, we used the 10X multiome PBMC dataset (62) which provides for each cell both its

chromatin accessibility profile and its gene expression profile and simulated 100 pseudobulks with

diverse cellular compositions (see Material and methods). We used EPIC-ATAC to perform ATAC-Seq

based deconvolution on the chromatin accessibility levels of the peaks and the original EPIC tool to

perform standard RNA-seq deconvolution on the gene expression levels. ATAC-Seq peaks can also be

aggregated, based on peak distances to each gene, into gene activity (GA) variables as proxy for gene

expression. We thus applied the GA transformation to the 10x multiome PBMC dataset and

performed GA-based RNA deconvolution using the original EPIC tool (See Material and methods).

Figure 7 shows that EPIC-ATAC performs similarly to the EPIC RNA-seq based deconvolution and

outperforms the GA-based RNA deconvolution. The lower performances of GA based RNA

deconvolution could be explained by the fact that GA features, by construction, do not perfectly

match the transcriptomic data.

Discussion

Bulk chromatin accessibility profiling of biological tissues like tumors represents a reliable and

affordable technology to map the activity of gene regulatory elements across multiple samples in

different conditions. Here, we collected ATAC-Seq data from pure cell populations covering major

immune and non-immune cancer-relevant cell types from diseased, stimulated and healthy samples.

This enabled us to identify reliable cell-type specific marker peaks and chromatin accessibility profiles

for both PBMC and solid tumor sample deconvolution. We integrated these data in the EPIC

deconvolution framework to accurately predict the fraction of both malignant and non-malignant cell

types from bulk tumor ATAC-Seq samples.

In cases where specific cell types are expected in a sample but are not part of our list of reference

profiles (*e.g.,* neuronal cells in brain tumors), custom marker peaks and reference profiles can be

provided to EPIC-ATAC to perform cell-type deconvolution and we provide the code to generate such

markers and profiles based on ATAC-Seq data from sorted cells, following the approach developed in

this work (Figure 1, see Code availability).

Solid tumors contain large and heterogeneous fractions of cancer cells for which it is

challenging to build reference profiles. To our knowledge this work provides the first benchmark of

deconvolution tools adapted to ATAC-Seq data in the context of solid tumor samples. We show that

the EPIC-ATAC framework, in contrast to other existing tools, allows users to accurately predict the

proportion of cells not included in the reference profiles (Figure 4 and Supplementary Figure 4).

These uncharacterized cells can include cancer cells but also other non-malignant cells. Since the

major cell types composing TMEs were included in our reference profiles, the proportion of

uncharacterized cells approximates the proportion of the cancer cells in most cases.

The pseudobulk approach provides unique opportunities to design benchmarks with known

cell-type proportions but also comes with some limitations. Indeed, pseudobulks are generated from

single-cell data which are noisy and whose cell-type annotation is challenging in particular for closely

related cell types. These limitations might lead to chromatin accessibility profiles that deviates from

true bulk data and errors in the true cell-type proportions. For this reason, we anticipate that the

newly generated benchmarking PBMCs dataset with ground truth cell proportions obtained by flow

cytometry will nicely complement pseudobulk from scATAC-Seq data in future benchmarks of ATAC-

Seq deconvolution. The qualitative evaluation of our method on true bulk ATAC-Seq samples from

breast cancer patients and the observation of similar immune compositions in TN and ER+/HER2-

samples as the ones identified in the original paper (Figure 6) further support the accuracy of EPIC-

ATAC to deconvolve bulk ATAC-Seq data, without requiring additional scATAC-Seq data which are not

always available for all cancer types.

Overall the evaluation of the EPIC-ATAC deconvolution resulted in an average absolute error

of 7% across cell types. This number is consistent with previous observations in RNA-Seq data

deconvolution (17). Considering this uncertainty, the quantification of low frequency populations

remains challenging (65). While the estimated proportions of these populations by EPIC-ATAC are

low (*e.g.,* dendritic cells), comparing such estimations across samples should be performed with care

due to the uncertainty of the predictions.

Another limitation of cell-type deconvolution is often reached when closely related cell types

are considered. In the reference-based methods used in this study, this limit was reached when

considering T-cell subtypes in the reference profiles (Figure 5 and Supplementary figure 8). We thus

recommend to use the EPIC-ATAC framework using the markers and reference profiles based on the

major cell populations. We additionally provide the marker peaks of the T-cell subtypes which could

be used to build cell-type specific chromatin accessibility signatures or perform "peak set enrichment

analysis" similarly to gene set enrichment analysis (GSEA, (66)). Such application could be useful for

the annotation of scATAC-Seq data, which often relies on matched RNA-Seq data and for which there

is a lack of markers at the peak level (67).

Another possible application of our marker peaks relies on their annotation (Figure 2G,

Supplementary Tables 4-5), which could be used to expand the list of genes and CBPs associated to

each cell type or subtype. For example, the neutrophils marker peaks were enriched for motifs of TFs

such as SPI1 (Supplementary Table 4), which was not listed in the neutrophil genes in the databases

used for annotation but has been reported in previous studies as involved in neutrophils

development (68). The annotations related to the set of major cell types and T-cell subtypes are

provided in Supplementary Tables 4-5 and 10-11. Finally, the annotation of marker peaks highlighted

pathways involved in immune responses to tumoral environments (Figure 2G). Examples of these

pathways are the toll-like receptor signaling pathway involved in pathogen-associated and

recognition of damage-associated molecular patterns in diverse cell types including B and T cells

(69,70), glucan metabolic processes which are known to be related to trained immunity which can

lead to anti-tumor phenotype in neutrophils (71) or the Fc-receptor signaling observed in NK cells

(72,73). These observations suggest that our marker peaks contain regulatory regions not only

specific to cell types but also adapted to the biological context of solid tumors.

Conclusion

In this work, we identified biologically relevant cell-type specific chromatin accessibility markers and

profiles for all major cancer-relevant cell types. We capitalized on these markers and profiles to

predict cell-type proportions from bulk PBMC and solid tumor ATAC-Seq data

(https://github.com/GfellerLab/EPIC-ATAC). Evaluated on diverse tissues, EPIC-ATAC shows reliable

predictions of immune, stromal, vascular and cancer cell proportions. With the expected increase of

ATAC-Seq studies in cancer, the EPIC-ATAC framework will enable researchers to deconvolve bulk

ATAC-Seq data from tumor samples to support the analysis of regulatory processes underlying tumor

development, and correlate the TME composition with clinical variables.

Materials and methods

Generation of an ATAC-Seq reference dataset of cancer relevant cell types.

Pre-processing of the sorted ATAC-Seq datasets

We collected pure ATAC-Seq samples from 12 studies. The data include samples from (i) ten major immune, stromal and vascular cell types (B (35,39,40), CD4+ (35,40–43), CD8+ (35,39–41,43), natural killer (NK) (35,39), dendritic (DCs) cells (39,41,44), macrophages (40,41), monocytes (35,39,40,44,45) and neutrophils (46,47) as well as fibroblasts (41,48) and endothelial (41,49) cells (See Figure 2A), and (ii) eight tissues from distinct organs (*i.e* bladder, breast, colon, liver, lung, ovary, pancreas and thyroid) from the ENCODE data (50,51). The list of the samples and their associated metadata (including cell types and accession number of the study of origin) is provided in Supplementary Table 1. To limit batch effects, the samples were reprocessed homogeneously from the raw data (fastq files) processing to the peak calling. For that purpose, raw fastq files were collected from GEO using the SRA toolkit and the PEPATAC framework (74) was used to process the raw fastq files based on the following tool: trimmomatic for adapter trimming, bowtie2 (with the PEPATAC default parameters) for reads pre-alignment on human repeats and human mitochondrial reference genome, bowtie2 (with the default PEPATAC parameters: *--very-sensitive -X 2000*) for alignment on the human genome (hg38), samtools (PEPATAC default parameters: *-q 10*) for duplicates removal and MACS2 (75) (PEPATAC default parameters: *--shift -75 --extsize 150 --nomodel --call-summits --nolambda --keep-dup all -p 0.01*) for peak calling in each sample. After alignment, reads mapping on chromosome M were excluded. TSS enrichment scores were computed for each sample and used to filter out samples with low quality (criteria of exclusion: TSS score < 5) (See Supplementary Table 1 containing the TSS score of each sample). 789 samples (including 564 from our ten reference cell-types) had a TSS score > 5.

Generation of a consensus set of peaks

Peak calling was performed in each sample individually. Peaks were then iteratively collapsed to

generate a set of reproducible peaks. For each cell type, peaks collapse was performed adapting the

iterative overlap peak merging approach proposed in the PEPATAC framework. A first peaks collapse

was performed at the level of each study of origin, *i.e.,* if peaks identified in distinct samples

overlapped (minimum overlap of 1bp between peaks), only the peak with the highest peak calling

score was kept. Also, only peaks detected in at least half of the samples of each study were

considered for the next step. If a study had only two samples, only peaks detected in both samples

were considered. After this first selection, a second round of peaks collapse was performed at the

cell-type level to limit batch effects in downstream analyses. For each cell type, only peaks detected

in all the studies of origin were considered. The final list of peaks was then generated by merging

each set of reproducible peaks. Peaks located on chromosome Y were excluded from the rest of the

analyses. ATAC-Seq counts were retrieved for each sample and each peak using featureCounts (76).

Identification of cell-type specific markers

Differential accessibility analysis

To identify cell-type specific markers, we split the samples collection in ten folds (created with the

create_folds function from the R package splitTools (77)). For each fold, we performed pairwise

differential accessibility analysis across the ten cell types considered in the reference samples as well

as the ENCODE samples from diverse organs. The differential analysis was performed using limma

((78), version 3.56.2). Effective library sizes were computed using the method of trimmed mean of

M-values (TMM) from the edgeR package in R ((79), version 3.42.4). Due to differences of library size

across all samples collected, we used voom from the limma package (80) to transform the data and

model the mean-variance relationship. Finally, a linear model was fitted to the data to assess the

differential accessibility of each peak across each pair of cell types. To identify our marker peaks, all

peaks with log2 fold change higher than 0.2 were selected and ranked by their maximum adjusted *p-*

value across all pairwise comparisons. The top 200 features (with the lowest maximum adjusted *p-*

value) were considered as cell-type specific marker peaks. The marker peaks identified in at least

three folds were considered in the final list of marker peaks.

Marker peaks filtering

Modules of open chromatin regions accessible in all (universal modules) or in specific human tissues

have been identified in the study Zhang *et al.* (2). These regions were used to refine the set of marker

peaks and exclude peaks with residual accessibility in other cell types than those considered for

deconvolution. More precisely, for immune, endothelial and fibroblasts specific peaks, we filtered

out the peaks overlapping the universal modules as well as the tissue specific modules except the

immune (modules 8 to 25), endothelial (modules 26 to 35) and stromal related modules (modules 41

to 49 and 139-150) respectively. As a second filtering step, we retained markers exhibiting the

highest correlation patterns in tumor bulk samples from different cancer types, *i.e.,* The Cancer

Genome Atlas (TCGA) samples (5). We used the Cancer Genomics Cloud (CGC) (81) to retrieve the

ATAC-Seq counts for each marker peaks in each TCGA sample (using *featureCounts*). For each set of

cell-type specific peaks, we identified the most correlated peaks using the *findCorrelation* function of

the caret R package ((82), version 6.0-94) with a correlation cutoff value corresponding to the 90^{th}

percentile of pairwise Pearson correlation values.

Evaluation of the study of origin batch effect

To identify potential batch effect issues, we run principal component analysis (PCA) based on the

cell-type specific peaks after normalizing ATAC-Seq counts using full quantile normalization (FQ-FQ)

implemented in the EDASeq R package (83) to correct for depth and GC biases. These data were used

to visualize the data in two-dimensional space running Uniform Manifold Approximation (UMAP)

based on the PBMC and TME markers (Figure 2B). We also run PCA and used the ten first principal

components to evaluate distances between samples and compute silhouette coefficients based on

the cell type and study of origin classifications.

Building the reference profiles

It has been previously demonstrated in the context of RNA-Seq based deconvolution approaches (16,17) that the transcripts per million (TPM) transformation is appropriate to estimate cell fractions from bulk mixtures. We thus normalized the ATAC-Seq counts of the reference samples using a TPM-like transformation, *i.e.,* dividing counts by peak length, correcting samples counts for depth and rescaling counts so that the counts of each sample sum to 10^6. We then computed for each peak the median of the TPM-like counts across all samples from each cell type to build the reference profiles of the ten cell types considered in the EPIC-ATAC framework (Figure 2C). In the EPIC algorithm, weights reflecting the variability of each feature of the reference profile can be considered in the constrained least square optimization. We thus also computed the inter-quartile range of the TPM-like counts for each feature in each cell type. Two ATAC-Seq reference profiles are available in the EPIC-ATAC framework: (i) a reference profile containing profiles for B cells, CD4+ T cells, CD8+ T cells, NK, monocytes, dendritic cells and neutrophils to deconvolve PBMC samples, and (ii) a reference profile containing profiles for B cells, CD4+ T cells, CD8+ T cells, NK, dendritic cells, macrophages, neutrophils, fibroblasts and endothelial cells to deconvolve tumor samples. The reference profiles are available in the EPICATAC R package and the reference profiles restricted to our cell-type specific marker peaks are available in the Supplementary Tables 2 and 3.

Assessing the reproducibility of the marker peaks signal in independent samples

We evaluated the chromatin accessibility level of the marker peaks in samples that were not included in the peak calling step. Firstly, we considered samples from two independent studies (84,85) providing pure ATAC-Seq data for five immune cell types (*i.e.,* B, CD4+ T cells, CD8+ T cells, Monocytes, Macrophages) (Figure 2D). To consider the other cell types, samples that were excluded from the reference dataset due to a low TSS enrichment score were also considered in this validation dataset (Supplementary Table 1). Secondly, we collected the data from a single-cell atlas chromatin

472 accessibility from human tissues and considered the cell types included in our reference data (2)

473 (Figure 2E). We used the cell-type annotations provided in the original study (GEO accession number:

474 GSE184462). The Signac R package ((58), 1.9.0) was used to extract fragments counts for each cell

475 and each marker peak and the ATAC-Seq signal of each marker peak was averaged across all cells of

476 each cell type.

478 **Annotation of the marker peaks**

479 The cell-type specific markers were annotated using ChIPseeker R package ((53), version 1.34.1) and

480 the annotation from *TxDb.Hsapiens.UCSC.hg38.knownGene* in R to identify the regions in which the

481 marker peaks are (*i.e.,* promoter, intronic regions, etc.) and ChipEnrich to associate each peak to the

482 nearest gene TSS (54). The nearest genes identified were then compared to cell-type marker genes

483 listed in the PanglaoDB (55) and CellMarker databases (56). PanglaoDB provides an online interface

484 to explore a large collection on single-cell RNA-Seq data as well as a community-curated list of cell-

485 type marker genes. CellMarker is a database providing a large set of curated cell-type markers for

486 more than 400 cell types in human tissues retrieved from a large collection of single-cell studies and

487 flow cytometry, immunostaining or experimental studies. ChipEnrich was also used to perform gene

488 set enrichment and identify for each set of cell-type specific peaks potential biological pathways

489 regulated by the marker peaks. The enrichment analysis was performed using the *chipenrich* function

490 (*genesets = "GOBP" , locusdef = "nearest_tss"*) from the chipenrich R package (v2.22.0).

491 Chromatin accessibility peaks can also be annotated for chromatin binding proteins (CBPs) such as

492 transcription factors (TFs), whose potential binding in the peak region is reported in databases. In our

493 study we chose the JASPAR2022 (57) database and the ReMap database (60).

494 Using the JASPAR2022 database, we assessed, for each cell type, whether the cell-type specific

495 marker peaks were enriched in specific TFs motifs using two TFs enrichment analysis frameworks:

496 Signac (58) and MonaLisa (59). For the MonaLisa analysis, the cell-types specific markers peaks were

497 categorized in bins of sequences, one bin per cell type (use of the *calcBinnedMotifEnrR* function). To

test for an enrichment of motifs, the sequences of each bin were compared to a set of background

peaks with similar average size and GC composition obtained by randomly sampling regions in all the

peaks identified from the reference dataset. The enrichment test was based on a binomial test. For

the Signac analysis, we used the *FindMotif* function to identify over-represented TF motifs in each set

of cell-type specific marker peaks (query). This function used a hypergeometric test to compare the

number of query peaks containing the motif with the total number of peaks containing the motif in

the background regions (matched to have similar GC content, region length and dinucleotide

frequencies as the query regions), corresponding in our case to the peaks called in the reference

dataset.

The ReMap database associates chromatin binding proteins (CBPs), including TFs, transcriptional

coactivators and chromatin-remodeling factors, to their DNA binding regions based on DNA-binding

experiments such as chromatin immunoprecipitation followed by sequencing (ChIP-seq). For each

association of a CBP to its binding region, the cell type in which the binding has been observed is

reported in the ReMap database (biotype). We used the ReMapEnrich R package (version 0.99) to

test if the cell-type specific marker peaks are significantly enriched in CBPs-binding regions listed in

the Remap 2022 catalog. We considered the non-redundant peaks catalog from Remap 2022,

containing non-redundant binding regions for each CBP in each biotype. Similarly to the previously

mentioned enrichment methods, we chose the consensus peaks called in the reference samples as

universe for the enrichment test. Note that, for each cell type, an enrichment was retained only if the

biotype in which the CBP-regions were identified matched the correct cell-type.

Running EPIC-ATAC on bulk ATAC-Seq data

The samples used to generate the reference profiles were aligned using the hg38 reference genome.

To assure the compatibility of any input bulk ATAC-Seq dataset with the EPIC-ATAC marker peaks and

reference profiles, we provide an option to lift over hg19 datasets to hg38 (use of the liftOver R

package). Subsequently, the features of the input bulk matrix are matched to our reference profiles

features. To match both sets of features, we determine for each peak of the input bulk matrix the

distance to the nearest peak in the reference profiles peaks. Overlapping regions are retained and

the feature IDs are matched to their associated nearest peaks. If multiple features are matched to

the same reference peak, the counts are summed. In RNA-Seq based deconvolution, EPIC uses an

estimation of the amount of mRNA in each reference cell type to derive cell proportions. For the

ATAC-Seq based deconvolution these values were set to 1 to give similar weights to all cell-types

quantifications.

<u>Datasets used for the evaluation of ATAC-Seq deconvolution</u>

PBMCs ATAC-Seq data from healthy donors

Peripheral blood mononuclear cell (PBMC) isolation

Venous blood from five healthy donors was collected at the local blood transfusion center of Geneva

in Switzerland, under the approval of the Geneva University Hospital's Institute Review Board, upon

written informed consent and in accordance with the Declaration of Helsinki. PBMCs were freshly

isolated by Lymphoprep (Promega) centrifugation (1800 rpm, 20 minutes, without break, room

temperature). Red blood cell lysis was performed using red blood lysis buffer (Qiagen) and platelets

were removed by centrifugation (1000 rpm, 10 minutes without break, room temperature). Cells

were counted and immediately used.

Flow cytometry

Immune cell populations were identified using multiparameter flow cytometry and the following

antibodies: FITC anti-human CD45RA (HI100, Biolegend), PerCP-Cyanine5.5 anti-human CD19 (H1B19,

Biolegend), PE anti-human CD3 (SK7, Biolegend), PE-Dazzle anti-human CD14 (M0P9, BD

Biosciences), PE-Cyanine7 anti-human CD56 (HCD56, Biolegend), APC anti-human CD4 (RPA-T4,

Biolgend), APC-Cyanine7 anti-human CCR7 (G043H7, Biolegend), Brilliant Violet 421 anti-human CD8

(RPA-T8, Biolegend), Brilliant Violet 510 anti-human CD25 (BC96, Biolegend), Brilliant Violet 711 anti-

human CD16 (3G8, Biolegend), Brilliant Violet 786 anti-human CD127 (A019D5, Biolegend), Ultra-

Brilliant Violet anti-human CD45 (HI30, BD Biosciences), FITC anti-human Celc9a (8F9, Miltenyi) , PE

anti-human XCR1 (S15046E, Biolegend), PE-Dazzle anti-human BDCA-2 (201A, Biolegend), APC anti-

human BDCA-3 (AD5-14H12, Miltenyi), Brilliant Violet 421 anti-human CD3 (UCHT1, Biolegend),

Brilliant Violet 421 anti-human CD14 (M5E2, BD Pharmingen), Brilliant Violet 421 anti-human CD19

(SJ25C1, Biolegend), Brilliant Violet 510 anti-human BDCA-1 (L161, Biolegend), Brilliant Violet 650

anti-human CD11c (3.9, Biolegend), Brilliant Violet 711 anti-human CD11c (N418, Biolegend) and

Brilliant Violet 711 anti-human HLA-DR (L243, Biolegend). Dead cells were excluded using the Zombie

UV™ Fixable Viability Kit (Biolegend). Intracellular staining was performed after fixation and

permeabilization of the cells with the FoxP3 Transcription Factor Staining Buffer Set (00-5523-00,

Invitrogen) using Alexa 700 anti-human FoxP3 antibody (259D/C7, BD Biosciences). Data were

acquired on LSRFortessa flow cytometer and analysed using FlowJo software (v10.7.1).

Cell preparation for ATAC-Sequencing

50000 CD45+ cells were sorted from total PBMCs using anti-human Ultra-Brilliant Violet (BUV395)

CD45 (HI30, BD Biosciences) with a FACSAria II (Becton Dickinson) and were collected in PBS with

10% Foetal Bovine Serum (FBS). Cell pellets were resuspended in cold lysis buffer (10mM Tris-Cl pH

7.4, 10mM NaCl, 3mM MgCl2, 0,1% NP40 and water) and immediately centrifuged at 600g for 30min

at 4°C. Transposition reaction was performed using the Illumina Tagment DNA Enzyme and Buffer kit

(20034210, Illumina) and transposed DNA was eluted using the MinElute PCR Purification Kit

(Qiagen). Libraries were generated by PCR amplification using indexing primers and NEBNext High-

Fidelity Master Mix (New England BioLabs) and were purified using AMPure XP beads (A63880,

Beckman Coulter). Libraries were quantified by a fluorometric method (QubIT, Life Technologies) and

their quality assessed on a Fragment Analyzer (Agilent Technologies). Sequencing was performed as a

paired end 50 cycles run on an Illumina NovaSeq 6000 (v1.5 reagents) at the Genomic Technologies

Facility (GTF) in Lausanne, Switzerland. Raw sequencing data were demultiplexed using the

bcl2fastq2 Conversion Software (version 2.20, Illumina).

Data processing

The same steps as for the processing of the reference ATAC-Seq samples were followed. (See Pre-

processing of the ATAC-Seq datasets).

ATAC-Seq pseudobulk data from PBMCs and cancer samples

To evaluate the accuracy of our ATAC-Seq deconvolution framework, we generated pseudo-bulk

datasets from 5 single-cell datasets:

• **PBMC pseudobulk dataset:** combination of three single-cell datasets for PBMCs.

o Dataset 1 corresponds to a scATAC-Seq dataset obtained from Satpathy et al. (13)

(GEO accession number: GSE129785). This dataset contains FACS-sorted populations

of PBMCs. Since the cells of some cell types came from a unique donor, all the cells

of this dataset were aggregated to form one pseudobulk. Ground truth cell fractions

were obtained by dividing the number of cells in each cell type by the total number

of cells.

o Dataset 2 (included in the PBMC pseudobulk dataset) was retrieved from Granja *et*

al. (61) (GEO accession number GSE139369). B cells, monocytes, dendritic, CD8+,

CD4+ T, NK cells, neutrophils from healthy donors were considered. The neutrophil

cells came from a single donor. As for dataset 1, we thus aggregated all the cells to

generate one pseudobulk. Ground truth cell fractions were obtained by dividing the

number of cells in each cell type by the total number of cells.

o Dataset 3 (included in the PBMC pseudobulk dataset) corresponds to the 10X

multiome dataset of PBMC cells (62). Since these data come from one donor, one

pseudobulk sample was generated for this dataset. The pseudobulk was generated

by averaging the ATAC-Seq signal from all cells from the following cell types: B cells,

CD4+ T cells , CD8+ T cells, NK cells, Dendritic cells and monocytes.

• **Basal cell carcinoma dataset:** obtained from the study of Satpathy *et al.* (13). This dataset is

a scATAC-Seq dataset composed of 13 basal cell carcinoma samples composed of immune (B

cells, plasma cells, CD4+ T cells, CD8+ T cells, NK cells, myeloid cells), stromal (endothelial

and fibroblasts) and cancer cells. Plasma cells and cancer cells were both considered as

uncharacterized cells (*i.e.,* cell types not included in the reference profiles). Cell annotations

were retrieved from the original study.

• **Gynecological cancer dataset:** obtained from the study of Regner *et al.* (63) (GEO accession

number GSE173682). In this study, the authors performed scATAC-Seq on 11 gynecological

cancer samples from two tumor sites (*i.e* endometrium and ovary) and composed of immune

(B cells, NK and T cells grouped under the same cell-type annotation, macrophages, mast

cells), stromal (fibroblast, endothelial, smooth muscle) and cancer cells. Mast cells, smooth

muscle and cancer cells were considered as uncharacterized cells. Cell annotations were

retrieved from the original study.

For Basal cell carcinoma and Gynecological cancer datasets, one pseudobulk per sample was

generated and ground truth cell fractions were obtained for each sample by dividing the number of

cells in each cell type by the total number of cells in the sample.

For each dataset, raw fragments files were downloaded from the respective GEO accession numbers

and data were preprocessed using ArchR ((86), ArchR R package 1.0.2). Cells with TSS score below

four were removed. Doublets removal was performed using the *doubletsRemoval* function from

ArchR. To match as much as possible real bulk ATAC-seq data processing, peak calling was not

performed on each cell type or cell cluster as usually done in scATAC-Seq studies but using all cells

for each dataset from the PBMC pseudobulk data or grouping cells by sample for the Basal cell

carcinoma and Gynecological cancer datasets. Peak calling was performed using MACS2 within the

ArchR framework. Fragments counts were extracted using ArchR for each peak called to generate

single-cell peak counts matrices. These matrices were normalized using a TPM-like transformation,

i.e., dividing counts by peak length and correcting samples counts for depth. Finally, for each peak,

the average of the normalized counts was computed across all the cells for each dataset from the

PBMC pseudobulk data and across all the cells of each sample for the Basal cell carcinoma and

Gynecological cancer datasets. Averaged data were then rescaled so that the sum of counts of each

sample sum to 10^6.

Bulk ATAC-Seq data from a breast cancer cohort

Bulk ATAC-Seq samples from a breast cancer cohort was obtained from Kumegawa *et al.* (64). These

data include 42 breast cancer samples which can be classified based on two features: (i) the breast

cancer subtype ER+/HER2- or triple negative, and (ii) the molecular classification provided by the

original study (CA-A, CA-B and CA-C). The ATAC-Seq raw counts and the samples metadata were

retrieved from figshare (87). As for the previously mentioned datasets, raw counts were normalized

using the TPM-like transformation prior to bulk deconvolution.

<u>**Benchmarking of the EPIC-ATAC framework against other existing deconvolution tools**</u>

The performances of the EPIC-ATAC framework were benchmarked against the following

deconvolution tools:

• quanTIseq (21) is a deconvolution tool using constrained least square regression to

deconvolve RNA-Seq bulk samples. No reference profiles are available in this framework to

perform ATAC-Seq deconvolution and quanTIseq does not provide the option to

automatically build reference profiles from pure bulk samples. quanTIseq was thus run using

the reference profiles derived in this work for the EPIC-ATAC framework and the *quanTIseq*

function from the quantiseqr R package (parameters: scaling set to 1 for all cell types and

method set to "lsei").

- DeconPeaker [20] relies on SIMPLS, a variant of partial least square regression to perform bulk RNA-Seq and bulk ATAC-Seq deconvolution. ATAC-Seq reference profiles are available in this deconvolution framework however not all cell types considered in the EPIC-ATAC framework are included in the DeconPeaker reference profiles. This tool was thus run using different reference profiles: (i) the reference profiles derived in this work for the EPIC-ATAC framework (corresponds to "DeconPeaker" or "DeconPeaker_ourmarkers" in our analyses), and (ii) reference profiles automatically generated by DeconPeaker from the sorted reference samples collected in this work (corresponds to "DeconPeaker_cust." in our analyses). The results of DeconPeaker obtained using its original markers and profiles are also provided for the cell types in common with the cell types considered in this work in Supplementary Figures 3 and 5. Deconvolution was run using the deconvolution module deconPeaker (using findctsps with the following parameter: --lib-strategy=ATAC-Seq). DeconPeaker outputs cell-type proportions relative to the total amount of cells from the reference cell types.

- CIBERSORTx [19] is a deconvolution algorithm based on linear support vector regression. CIBERSORTx does not provide ATAC-Seq reference profiles, however it is possible to automatically generate new profiles from a set of pure bulk samples. This tool was thus run using different reference profiles: i) the reference profiles derived in this work for the EPIC-ATAC framework (corresponds to "CIBERSORTx" or "CIBERSORTx_ourMarkers" in our analyses), and ii) reference profiles automatically generated by CIBERSORTx from the sorted reference samples collected in this work (corresponds to "CIBERSORTx_cust." in our analyses). To run CIBERSORTx, we used the docker container provided by the authors of CIBERSORTx on their website. The algorithm was run using the default options (i.e --absolute FALSE, --rmbatchBmode FALSE and –rmbatchSmode FALSE), which results in cell-type proportions relative to the total amount of cells from the reference cell types.

- ABIS (18) uses robust linear modeling to estimate cell-type proportions in bulk RNA-Seq samples. No ATAC-Seq reference profiles are available in the deconvolution framework. ABIS was run using the EPIC-ATAC reference profiles by using the *rlm* function from the MASS R package (as performed in the deconvolute_abis function from the immunedeconv R package (16) was used to quantify each cell type from the reference profiles. The cell-types quantifications returned by this approach are in arbitrary units. To compare the estimations and the true cell proportions, we scaled the estimations of each sample between 0 and 1 to obtained relative proportions.

- MCPcounter (22): MCPcounter returns scores instead of cell type proportions. The scores were obtained using the *appendSignatures* function from the MCPcounter R package by providing the list of marker peaks specific to each cell type. The cell-type scores are not comparable between cell type, MCPcounter was thus included only in the evaluation of the performances in each cell type separately.

For all the tools, TPM-like data were used as input bulk samples for the deconvolution.

Since CIBERSORTx, ABIS and DeconPeaker do not predict proportions of uncharacterized cells, we performed two benchmarking analyses: (i) including all cell types and (ii) excluding the cell types that are absent from the reference profiles (uncharacterized cells) and rescaling the estimated and true proportions of the immune cells, endothelial cells and fibroblasts so that their sum equals 1.

Comparing deconvolution based on RNA-Seq, gene activity or peaks features.

100 pseudobulks were generated from the 10X PBMC multiome dataset (62) based on 3000 cells for each pseudobulk. Cell fractions were defined using the *rdirichlet* function from the *gtools* R package. Three sets of features were extracted from the data, *i.e.,* gene expression features extracted from the RNA-Seq layer, ATAC-Seq peaks and gene activity derived from the ATAC-Seq layer. The same cells sampling was considered for each modality.

Gene activity features were extracted from the single-cell data using ArchR (1.0.2), which considers

distal elements and adjusts for large differences in gene size in the gene activity score calculation.

Gene activity pseudobulks were built by averaging the gene activity scores across all cells belonging

to the pseudobulk. For ATAC-Seq pseudobulk, peaks called using ArchR on all cells form the 10X

dataset were considered (see the method section "ATAC-Seq pseudobulk data from PBMCs and

cancer samples") and counts were averaged across all cells of each pseudobulk. For RNA-Seq

pseudobulks, counts were also averaged across all cells of each pseudobulk. All aggregated data were

depth normalized across each features to 10^6. Cell-type deconvolution was performed on each

pseudobulk using EPIC-ATAC on the peak matrix using our ATAC-Seq marker peaks and reference

profiles. The RNA-Seq and gene activity pseudobulks were deconvolved with EPIC.

Code availability

The code to download and preprocess publicly available ATAC-Seq samples as well as the code used

to identify our cell-type specific marker peaks and generate the reference profiles is available on

GitHub (https://github.com/GfellerLab/EPIC-ATAC_Manuscript). A README file is provided on the

GitHub repository with more details on how to use the code.

The code to perform ATAC-Seq deconvolution using the EPIC-ATAC framework is available as an R

package called EPICATAC and is available on GitHub (https://github.com/GfellerLab/EPIC-ATAC).

Data availability

The   newly   generated   ATAC-Seq   data   have   been   deposited   on   Zenodo   (doi:

10.5281/zenodo.8358719). The other data related to this work are available in the supplementary

tables and on the Zenodo deposit (doi: 10.5281/zenodo.8358719).

Competing interests:

The authors declare that they have no competing interests.

Acknowledgement:

We thank the Lausanne Genomic Technologies Facility, University of Lausanne, Switzerland

(https://www.unil.ch/gtf/en/home.html) for the sequencing of the PBMC samples as well as Yan Liu,

Dana Moreno and Matei Teleman for testing the EPICATAC R package. Some of the illustrations were

created with BioRender.com.

Authors contributions:

Conceptualization: AAGG, JR, DG; Data curation: AAGG; Software: AAGG, JR, DG; Experiments: MF,

CJ; Visualization: AAGG, JR, DG; Methodology: AAGG, JR, DG; Writing—original draft: AAGG, DG;

Writing—review and editing: all authors.

Figure legends:

Figure 1: Graphical description of the identification of cell-type specific marker peaks and reference

ATAC-Seq profiles included in the EPIC-ATAC framework. 1) 564 pure ATAC-Seq data of sorted cells

were collected to build reference profiles for cancer-relevant cell populations. 2) Cell-type specific

marker peaks were identified using differential accessibility analysis. 3) Markers with previously

observed chromatin accessibility in human healthy tissues were then excluded. 4) For tumor bulk

deconvolution, the set of remaining marker peaks was refined by selecting markers with correlated

behavior in tumor bulk samples. 5) The cell-type specific marker peaks and reference profiles were

finally integrated in the EPIC-ATAC framework to perform bulk ATAC-Seq deconvolution. Parts of this

figure were created with BioRender.com.

Figure 2: ATAC-Seq data from sorted cell populations reveal cell-type specific marker peaks and

reference profiles. A) Number of samples collected for each cell type. The colors correspond to the

different studies of origin. B) Representation of the collected samples in 2D using UMAP based on the

PBMC markers (left) and TME markers (right). Colors correspond to cell types. **C)** *Scaled averaged chromatin accessibility of the cell-type specific marker peaks (rows) in each cell type (columns) in the ATAC-Seq reference samples used to identify the marker peaks.* **D)** *Scaled averaged chromatin accessibility of the marker peaks in external ATAC-Seq data from samples of pure cell types excluded from the reference samples (see Material and Methods).* **E)** *Scaled averaged chromatin accessibility of the marker peaks in an external scATAC-Seq dataset (Human Atlas (2)).* **F)** *Distribution of the marker peak distances to the nearest transcription start site (TSS) (left panel) and the ChiPSeeker annotations (right panel).* **G)** *Significance (-log10(q.value)) of pathways (columns) enrichment test obtained using ChIP-Enrich on each set of cell-type specific marker peaks (rows). A subset of relevant enriched pathways is represented. Colors of the names of the pathways correspond to cell types where the pathways were found to be enriched. When pathways were significantly enriched in more than one set of peaks, pathways names are written in bold.*

Figure 3: EPIC-ATAC accurately estimates immune cell fractions in PBMC ATAC-Seq samples. A) *Schematic description of the experiment designed to validate the ATAC-Seq deconvolution on PBMC samples.* **B)** *Comparison between cell-type proportions predicted by EPIC-ATAC and the true proportions in the PBMC bulk dataset. Symbols correspond to donors.* **C)** *Comparison between the proportions of cell-types predicted by EPIC-ATAC and the true proportions in the PBMC pseudobulk dataset. Symbols correspond to pseudobulks.* **D)** *Pearson correlation (left) and RMSE (right) values obtained by each deconvolution tool on the PBMC bulk dataset. The EPIC-ATAC results are highlighted in red.* **E)** *Pearson correlation (left) and RMSE (right) values obtained by each deconvolution tool on the PBMC pseudobulk dataset. Parts of this figure (panel 1) were created with BioRender.com.*

Figure 4: EPIC-ATAC accurately predicts fractions of cancer and non-malignant cells in tumor samples. A) *Comparison between cell-type proportions estimated by EPIC-ATAC and true proportions for the basal cell carcinoma (top) and gynecological (bottom) pseudobulk datasets. Symbols*

*correspond to pseudobulks. **B)** Pearson's correlation and RMSE values obtained for the deconvolution*

*tools included in the benchmark. EPIC-ATAC is highlighted in red. **C)** Same analyses as in panels B,*

with the uncharacterized cell population excluded for the evaluation of the predictions accuracy. The

predicted and true proportions of the immune, stromal and vascular cell types were rescaled to sum

to 1.

Figure 5: T cell subtypes quantification reveals the ATAC-Seq deconvolution limits for closely

***related cell types. A)** Comparison of the proportions estimated by EPIC-ATAC and the true proportions*

for PBMC samples (PBMC experiment and PBMC pseudobulk samples combined) (top) and the basal

cell carcinoma pseudobulks (bottom). Predictions of the proportions of CD4+ and CD8+ T-cells were

obtained using the reference profiles based on the major cell types and subtype predictions using the

*reference profiles including the T-cell subtypes. **B)** Pearson's correlation values obtained by EPIC-ATAC*

in each cell type.

Figure 6: EPIC-ATAC accurately infers the immune contexture in a bulk ATAC-Seq breast cancer

***cohort. A)** Proportions of different cell types predicted by EPIC-ATAC in the samples stratified based*

*on two breast cancer subtypes. **B)** Proportions of different cell types predicted by EPIC-ATAC in the*

samples stratified based on three ER+/HER2- subgroups. Wilcoxon test p-values are represented at

the top of the boxplots.

Figure 7: EPIC-ATAC performs similarly to EPIC RNA-seq based deconvolution and better than gene

***activity based deconvolution.** Pearson's correlation (left) and RMSE (right) values comparing the*

proportions predicted by the ATAC-Seq deconvolution, the RNA-Seq deconvolution and the GA-based

RNA deconvolution and true cell-type proportions in the 100 pseudobulks simulated form the 10x

multiome PBMC dataset (62). Dots correspond to outlier pseudobulks.

Supplementary Figures: Additional file named Supplementary_figures.pdf

Supplementary Tables:

Sup. Table 1: Metadata of the ATAC-Seq samples used in the study

Sup. Table 2: Averaged chromatin accessibility of the PBMC marker peaks in each cell-type.

Sup. Table 3: Averaged chromatin accessibility of the TME marker peaks in each cell-type.

Sup. Table 4: Annotations of the cell-type specific PBMC marker peaks

Sup. Table 5: Annotations of the cell-type specific TME marker peaks

Sup. Table 6: GO pathways enriched in each set of cell-type specific PBMC marker peaks

Sup. Table 7: GO pathways enriched in each set of cell-type specific TME marker peaks

Sup. Table 8: Averaged chromatin accessibility of the PBMC marker peaks in each cell-type (T cells

subtypes included).

Sup. Table 9: Averaged chromatin accessibility of the TME marker peaks in each cell-type (T cells

subtypes included).

Sup. Table 10: Annotations of the cell-type specific PBMC marker peaks (T cells subtypes included).

Sup. Table 11: Annotations of the cell-type specific TME marker peaks (T cells subtypes included).

Sup. Table 12: GO pathways enriched in each set of cell-type specific PBMC marker peaks (T cell

subtypes).

Sup. Table 13: GO pathways enriched in each set of cell-type specific TME marker peaks (T cell

subtypes).

References:

1.   Klemm SL, Shipony Z, Greenleaf WJ. Chromatin accessibility and the regulatory epigenome.

Nat Rev Genet. 2019 Jan 23;20(4):207–20.

2.   Zhang K, Hocker JD, Miller M, Hou X, Chiou J, Poirion OB, et al. A single-cell atlas of chromatin

accessibility in the human genome. Cell. 2021;184(24):5985-6001.e19.

3.   Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ. Transposition of native chromatin

834   for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and

835   nucleosome position. Nat Methods. 2013 Oct 6;10(12):1213–8.

836   4.   Luo L, Gribskov M, Wang S. Bibliometric review of ATAC-Seq and its application in gene

837   expression. Brief Bioinform. 2022 Mar 7;

838   5.   Corces MR, Granja JM, Shams S, Louie BH, Seoane JA, Zhou W, et al. The chromatin

839   accessibility landscape of primary human cancers. Science. 2018 Oct 26;362(6413).

840   6.   Corces MR, Trevino AE, Hamilton EG, Greenside PG, Sinnott-Armstrong NA, Vesuna S, et al. An

841   improved ATAC-seq protocol reduces background and enables interrogation of frozen tissues.

842   Nat Methods. 2017 Oct 1;14(10):959–62.

843   7.   Grandi FC, Modi H, Kampman L, Corces MR. Chromatin accessibility profiling by ATAC-seq. Nat

844   Protoc. 2022 Apr 27;1518–52.

845   8.   Fridman WH, Pagès F, Sautès-Fridman C, Galon J. The immune contexture in human tumours:

846   impact on clinical outcome. Nat Rev Cancer. 2012 Mar 15;12(4):298–306.

847   9.   Fridman WH, Zitvogel L, Sautès-Fridman C, Kroemer G. The immune contexture in cancer

848   prognosis and treatment. Vol. 14, Nature Reviews Clinical Oncology. Nature Publishing Group;

849   2017. p. 717–34.

850   10.   de Visser KE, Joyce JA. The evolving tumor microenvironment: From cancer initiation to

851   metastatic outgrowth. Cancer Cell. 2023 Mar 13;41(3):374–403.

852   11.   Cusanovich DA, Daza R, Adey A, Pliner HA, Christiansen L, Gunderson KL, et al. Multiplex

853   single-cell profiling of chromatin accessibility by combinatorial cellular indexing. Science. 2015

854   May 22;348(6237):910–4.

855   12.   Lareau CA, Duarte FM, Chew JG, Kartha VK, Burkett ZD, Kohlway AS, et al. Droplet-based

856   combinatorial indexing for massive-scale single-cell chromatin accessibility. Nat Biotechnol.

857   2019 Jun 24;37(8):916–24.

13.   Satpathy AT, Granja JM, Yost KE, Qi Y, Meschi F, McDermott GP, et al. Massively parallel

single-cell chromatin landscapes of human immune cell development and intratumoral T cell

exhaustion. Nat Biotechnol. 2019 Aug 1;37(8):925–36.

14.   Avila Cobos F, Vandesompele J, Mestdagh P, De Preter K. Computational deconvolution of

transcriptomics data from mixed cell populations. Bioinformatics. 2018 Jun;34(11):1969–79.

15.   Avila Cobos F, Alquicira-Hernandez J, Powell JE, Mestdagh P, De Preter K. Benchmarking of cell

type deconvolution pipelines for transcriptomics data. Nat Commun. 2020;11(1):1–14.

16.   Sturm G, Finotello F, Petitprez F, Zhang JD, Baumbach J, Fridman WH, et al. Comprehensive

evaluation of transcriptome-based cell-type quantification methods for immuno-oncology.

Bioinformatics. 2019 Jul 5;35(14):i436–45.

17.   Racle J, de Jonge K, Baumgaertner P, Speiser DE, Gfeller D. Simultaneous enumeration of

cancer and immune cell types from bulk tumor gene expression data. Elife. 2017 Nov 13;6.

18.   Monaco G, Lee B, Xu W, Mustafah S, Hwang YY, Carré C, et al. RNA-Seq Signatures Normalized

by mRNA Abundance Allow Absolute Deconvolution of Human Immune Cell Types. Cell Rep.

2019 Feb 5;26(6):1627-1640.e7.

19.   Newman AM, Steen CB, Liu CL, Gentles AJ, Chaudhuri AA, Scherer F, et al. Determining cell

type abundance and expression from bulk tissues with digital cytometry. Nat Biotechnol.

2019;37(7):773–82.

20.   Li H, Sharma A, Luo K, Qin ZS, Sun X, Liu H. DeconPeaker, a Deconvolution Model to Identify

Cell Types Based on Chromatin Accessibility in ATAC-Seq Data of Mixture Samples. Front

Genet. 2020 Jun 8;11.

21.   Finotello F, Mayer C, Plattner C, Laschober G, Rieder Di, Hackl H, et al. Molecular and

pharmacological modulators of the tumor immune contexture revealed by deconvolution of

RNA-seq data. Genome Med. 2019 May 24;11:34.

22.   Becht E, Giraldo NA, Lacroix L, Buttard B, Elarouci N, Petitprez F, et al. Estimating the

population abundance of tissue-infiltrating immune and stromal cell populations using gene

expression. Genome Biol. 2016 Oct 20;17:218.

23.   Clarke J, Seol P, Clarke B. Statistical expression deconvolution from mixed tissue samples.

Bioinformatics. 2010;26(8):1043–9.

24.   Gosink MM, Petrie HT, Tsinoremas NF. Electronically subtracting expression patterns from a

mixed cell population. Bioinformatics. 2007 Dec 15;23(24):3328–34.

25.   Racle J, Gfeller D. EPIC: A tool to estimate the proportions of different cell types from bulk

gene expression data. In: Methods in Molecular Biology. Humana Press Inc.; 2020. p. 233–48.

26.   Newman AM, Liu CL, Green MR, Gentles AJ, Feng W, Xu Y, et al. Robust enumeration of cell

subsets from tissue expression profiles. Nat Methods. 2015 Mar 20;12(5):453–7.

27.   Li T, Fu J, Zeng Z, Cohen D, Li J, Chen Q, et al. TIMER2.0 for analysis of tumor-infiltrating

immune cells. Nucleic Acids Res. 2020 May 22;48:W509–14.

28.   Jimenez-Sanchez A, Cast O, Miller ML. Comprehensive benchmarking and integration of tumor

microenvironment cell estimation methods. Cancer Res. 2019 Dec 15;79(24):6238–46.

29.   Gong T, Szustakowski JD. DeconRNASeq: a statistical framework for deconvolution of

heterogeneous tissue samples based on mRNA-Seq data. Bioinformatics. 2013 Feb

21;29(8):1083–5.

30.   Chakravarthy A, Furness A, Joshi K, Ghorani E, Ford K, Ward MJ, et al. Pan-cancer

deconvolution of tumour composition using DNA methylation. Nat Commun. 2018 Aug 13;9.

31.   Teschendorff AE, Zhu T, Breeze CE, Beck S. EPISCORE: Cell type deconvolution of bulk tissue

DNA methylomes from single-cell RNA-Seq data. Genome Biol. 2020 Sep 4;21.

32.   Arneson D, Yang X, Wang K. MethylResolver—a method for deconvoluting bulk DNA

methylation profiles into known and unknown cell contents. Commun Biol. 2020;3.

33.   Zhang H, Cai R, Dai J, Sun W. EMeth: An EM algorithm for cell type decomposition based on

DNA methylation data. Sci Rep. 2021 Mar 11;11:5717.

34.   Feng S, Calinawan A, Pugliese P, Wang P, Ceccarelli M, Petralia F, et al. Decomprolute : A

benchmarking platform designed for multiomics-based tumor deconvolution. bioRxiv. 2023;

35.   Corces MR, Buenrostro JD, Wu B, Greenside PG, Chan SM, Koenig JL, et al. Lineage-specific

and single-cell chromatin accessibility charts human hematopoiesis and leukemia evolution.

Nat Genet. 2016 Aug 15;48:1193–203.

36.   Zeng W, Chen X, Duren Z, Wang Y, Jiang R, Wong WH. DC3 is a method for deconvolution and

coupled clustering from bulk and single-cell genomics data. Nat Commun. 2019 Oct

10;10:4613.

37.   Burdziak C, Azizi E, Prabhakaran S, Pe'er D. A Nonparametric Multi-view Model for Estimating

Cell Type-Specific Gene Regulatory Networks. ArXiv. 2019;

38.   Peng XL, Moffitt RA, Torphy RJ, Volmar KE, Yeh JJ. De novo compartment deconvolution and

weight estimation of tumor samples using DECODER. Nat Commun. 2019 Oct 18;10(1):4729.

39.   Calderon D, Nguyen MLT, Mezger A, Kathiria A, Müller F, Nguyen V, et al. Landscape of

stimulation-responsive chromatin across diverse human immune cells. Nat Genet. 2019 Sep

30;51(10):1494–505.

40.   Zhang P, Amarasinghe HE, Whalley JP, Tay C, Fang H, Migliorini G, et al. Epigenomic analysis

reveals a dynamic and context-specific macrophage enhancer landscape associated with

innate immune activation and tolerance. Genome Biol. 2022 Jun 24;23(1):136.

41.   Liu Q, Zaba L, Satpathy AT, Longmire M, Zhang W, Li K, et al. Chromatin accessibility

landscapes of skin cells in systemic sclerosis nominate dendritic cells in disease pathogenesis.

Nat Commun. 2020 Nov 17;11(1):5843.

42.   Mumbach MR, Satpathy AT, Boyle EA, Dai C, Gowen BG, Cho SW, et al. Enhancer connectome

in primary human cells identifies target genes of disease-associated DNA elements. Nat

Genet. 2017 Sep 15;49(11):1602–12.

43.   Giles JR, Manne S, Freilich E, Oldridge DA, Baxter AE, George S, et al. Human epigenetic and

transcriptional T cell differentiation atlas for identifying functional T cell-specific enhancers.

Immunity. 2022 Mar;55(3):557-574.e7.

44.   Leylek R, Alcántara-Hernández M, Granja JM, Chavez M, Perez K, Diaz OR, et al. Chromatin

Landscape Underpinning Human Dendritic Cell Heterogeneity. Cell Rep. 2020 Sep

22;32(12):108180.

45.   Trizzino M, Zucco A, Deliard S, Wang F, Barbieri E, Veglia F, et al. EGR1 is a gatekeeper of

inflammatory enhancers in human macrophages. Sci Adv. 2021 Jan 13;7(3).

46.   Ram-Mohan N, Thair SA, Litzenburger UM, Cogill S, Andini N, Yang X, et al. Profiling chromatin

accessibility responses in human neutrophils with sensitive pathogen detection. Life Sci

Alliance. 2021 Jun 18;4(8).

47.   Perez C, Botta C, Zabaleta A, Puig N, Cedena M-T, Goicoechea I, et al. Immunogenomic

identification and characterization of granulocytic myeloid-derived suppressor cells in

multiple myeloma. Blood. 2020 Jul 9;136(2):199–209.

48.   Ge X, Frank-Bertoncelj M, Klein K, McGovern A, Kuret T, Houtman M, et al. Functional

genomics atlas of synovial fibroblasts defining rheumatoid arthritis heritability. Genome Biol.

2021 Aug 25;22(1):247.

49.   Xin J, Zhang H, He Y, Duren Z, Bai C, Chen L, et al. Chromatin accessibility landscape and

regulatory network of high-altitude hypoxia adaptation. Nat Commun. 2020 Oct 1;11(1):4928.

50.   The ENCODE Project Consortium, Moore JE, Purcaro MJ, Pratt HE, Epstein CB, Shoresh N, et al.

Expanded encyclopaedias of DNA elements in the human and mouse genomes. Nature. 2020

Jul 29;583:699–710.

51.   Rozowsky J, Gao J, Borsari B, Guigó R, Gingeras TR, Gerstein M, et al. The EN-TEx resource of

multi-tissue personal epigenomes & variant-impact models. Cell. 2023;186(7):1493-1511.e40.

52.   Qiu Y, Wang J, Lei J, Roeder K. Identification of cell-type-specific marker genes from co-

expression patterns in tissue samples. Bioinformatics. 2021 Oct 11;37(19):3228–34.

53.   Yu G, Wang LG, He QY. ChIPseeker: an R/Bioconductor package for ChIP peak annotation,

comparison and visualization. Bioinformatics. 2015 Jul 15;31(14):2382–3.

54.   Welch RP, Lee C, Imbriano PM, Patil S, Weymouth TE, Smith RA, et al. ChIP-Enrich: gene set

enrichment testing for ChIP-seq data. Nucleic Acids Res. 2014 Sep 1;42(13):e105.

55.   Franzén O, Gan LM, Björkegren JLM. PanglaoDB: a web server for exploration of mouse and

human single-cell RNA sequencing data. Database. 2019;2019:baz046.

56.   Hu C, Li T, Xu Y, Zhang X, Li F, Bai J, et al. CellMarker 2.0: an updated database of manually

curated cell markers in human/mouse and web tools based on scRNA-seq data. Nucleic Acids

Res. 2023 Jan 6;51(D1):D870–6.

57.   Castro-Mondragon JA, Riudavets-Puig R, Rauluseviciute I, Berhanu Lemma R, Turchi L, Blanc-

Mathieu R, et al. JASPAR 2022: the 9th release of the open-access database of transcription

factor binding profiles. Nucleic Acids Res. 2022 Jan 7;50(D1):D165–73.

58.   Stuart T, Srivastava A, Madad S, Lareau CA, Satija R. Single-cell chromatin state analysis with

Signac. Nat Methods. 2021 Nov 1;18:1333–41.

59.   Machlab D, Burger L, Soneson C, Rijli FM, Schübeler D, Stadler MB. monaLisa: an

R/Bioconductor package for identifying regulatory motifs. Bioinformatics. 2022 Feb

22;38(9):2624–5.

60.   Hammal F, De Langen P, Bergon A, Lopez F, Ballester B. ReMap 2022: a database of Human,

Mouse, Drosophila and Arabidopsis regulatory regions from an integrative analysis of DNA-

binding sequencing experiments. Nucleic Acids Res. 2022 Jan 7;50(D1):D316–25.

61.   Granja JM, Klemm S, McGinnis LM, Kathiria AS, Mezger A, Corces MR, et al. Single-cell

multiomic analysis identifies regulatory programs in mixed-phenotype acute leukemia. Nat

Biotechnol. 2019 Dec 1;37(12):1458–65.

62.   10x Genomics. PBMC from a Healthy Donor - Granulocytes Removed Through Cell Sorting

(10k) [Internet]. 2021 [cited 2023 Sep 6]. Available from:

https://www.10xgenomics.com/resources/datasets/pbmc-from-a-healthy-donor-

granulocytes-removed-through-cell-sorting-10-k-1-standard-2-0-0

63.   Regner MJ, Wisniewska K, Garcia-Recio S, Thennavan A, Mendez-Giraldez R, Malladi VS, et al.

A multi-omic single-cell landscape of human gynecologic malignancies. Mol Cell.

2021;81(23):4924-4941.e10.

64.   Kumegawa K, Saeki S, Takahashi Y, Yang L, Osako T, Nakadai T, et al. Chromatin profile-based

identification of a novel ER-positive breast cancer subgroup with reduced ER-responsive

element accessibility. Br J Cancer. 2023 Feb 1;128(7):1208–22.

65. Jin H, Liu Z. A benchmark for RNA-seq deconvolution analysis under dynamic testing

environments. Genome Biol. 2021 Apr 12;22:102.

66. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set

enrichment analysis: a knowledge-based approach for interpreting genome-wide expression

profiles. PNAS. 2005 Sep 30;102(43):15545–50.

67. Jiang Y, Hu Z, Jiang J, Zhu A, Zhang Y, Lynch AW, et al. scATAnno: Automated Cell Type

Annotation for single-cell ATAC Sequencing Data. bioRxiv. 2023 Jun 15;

68. Watt S, Vasquez L, Walter K, Mann AL, Kundu K, Chen L, et al. Genetic perturbation of PU.1

binding and chromatin looping at neutrophil enhancers associates with autoimmune disease.

Nat Commun. 2021 Apr 16;12:2298.

69. Geng D, Zheng L, Srivastava R, Asprodites N, Velasco-Gonzalez C, Davila E. When Toll-like

receptor and T-cell receptor signals collide: a mechanism for enhanced CD8 T-cell effector

function. Blood. 2010 Nov 4;116(18):3494–504.

70. Javaid N, Choi S. Toll-like Receptors from the Perspective of Cancer Treatment. Cancers

(Basel). 2020 Jan 27;12(2):297.

71. Kalafati L, Kourtzelis I, Schulte-Schrepping J, Li X, Hatzioannou A, Grinenko T, et al. Innate

Immune Training of Granulopoiesis Promotes Anti-tumor Activity. Cell. 2020 Oct

29;183(3):771-785.e12.

72. Sanseviero E. NK Cell-Fc Receptors Advance Tumor Immunotherapy. J Clin Med. 2019 Oct

12;8(10):1667.

73. Bonnema JD, Karnitz LM, Schoon RA, Abraham RT, Leibson PJ. Fc receptor stimulation of

phosphatidylinositol 3-kinase in natural killer cells is associated with protein kinase C-

independent granule release and cell-mediated cytotoxicity. J Exp Med. 1994 Oct

1;180(4):1427–35.

74.   Smith JP, Corces MR, Xu J, Reuter VP, Chang HY, Sheffield NC. PEPATAC: an optimized pipeline

for ATAC-seq data analysis with serial alignments. NAR Genom Bioinform. 2021 Dec;3(4).

75.   Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, et al. Model-based analysis

of ChIP-Seq (MACS). Genome Biol. 2008 Sep 17;9(9):R137.

76.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning

sequence reads to genomic features. Bioinformatics. 2014 Nov 13;30(7):923–30.

77.   Mayer M. R package ' splitTools ': Tools for Data Splitting. R package version 1.0.1.

https://cran.r-project.org/web/packages/splitTools/index.html. 2023.

78.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential expression

analyses for RNA-sequencing and microarray studies. Nucleic Acids Res. 2015 Apr

20;43(7):e47.

79.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential

expression analysis of digital gene expression data. Bioinformatics. 2010 Jan 1;26(1):139–40.

80.   Law CW, Chen Y, Shi W, Smyth GK. Voom: Precision weights unlock linear model analysis tools

for RNA-seq read counts. Genome Biol. 2014 Feb 3;15:R29.

81.   Lau JW, Lehnert E, Sethi A, Malhotra R, Kaushik G, Onder Z, et al. The cancer genomics cloud:

Collaborative, reproducible, and democratized - A new paradigm in large-scale computational

research. Cancer Res. 2017 Nov 1;77(21):e3–6.

82.   Kuhn M. Building Predictive Models in R Using the caret Package. J Stat Softw. 2008 Nov

10;28(5):1–26.

83.   Risso D, Schwartz K, Sherlock G, Dudoit S. GC-Content Normalization for RNA-Seq Data. BMC

Bioinformatics. 2011 Dec 17;12:480.

84.   Ucar D, Márquez EJ, Chung CH, Marches R, Rossi RJ, Uyar A, et al. The chromatin accessibility

signature of human immune aging stems from CD8+ T cells. J Exp Med. 2017 Sep

13;214(10):3123–44.

85.   Carvalho K, Rebboah E, Jansen C, Williams K, Dowey A, McGill C, et al. Uncovering the Gene

Regulatory Networks Underlying Macrophage Polarization Through Comparative Analysis of

Bulk and Single-Cell Data. bioRxiv. 2021 Jan 21;

86.   Granja JM, Corces MR, Pierce SE, Bagdatli ST, Choudhry H, Chang HY, et al. ArchR is a scalable

software package for integrative single-cell chromatin accessibility analysis. Nat Genet. 2021

Feb 25;53:403–11.

87.   Kumegawa K. ATAC-seq data of 42 BC samples as SummarizedExperiment object with count

matrix, normalized count matrix, peak info, and clinical info [Internet]. 2023. Available from:

https://doi.org/10.6084/m9.figshare.21992609.v1

1049

List of abbreviations:

ATAC: Assay for Transposase-Accessible chromatin

CBP(s): chromatin binding protein(s)

ChIP-seq: chromatin immunoprecipitation followed by sequencing

DC: dendritic cells

NK: natural killer cells

PCA: principal component analysis

RMSE: root mean squared error

TCGA : The Cancer Genome Atlas

TF(s): transcription factor(s)

TSS: transcription start site

UMAP: Uniform Manifold Approximation

1062

Main Figures.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Robust estimation of cancer and immune cell-type proportions from bulk tumor ATAC-Seq data.

Aurélie AG Gabriel[1,2,3,4], Julien Racle[1,2,3,4], Maryline Falquet[3,5,6,7], Camilla Jandus[3,5,6,7], David Gfeller[1,2,3,4,*]

- Department of Oncology, Ludwig Institute for Cancer Research, University of Lausanne, Lausanne, Switzerland
- Agora Cancer Research Centre, Lausanne, Switzerland
- Swiss Cancer Center Leman (SCCL), Switzerland
- Swiss Institute of Bioinformatics (SIB), Lausanne, Switzerland.
- Department of Pathology and Immunology Faculty of Medicine, University of Geneva, Geneva, Switzerland
- Ludwig Institute for Cancer Research, Lausanne Branch, Lausanne, Switzerland
- Geneva Center for Inflammation Research, Geneva, Switzerland

* Corresponding author: david.gfeller@unil.ch

Supplementary Figure 1: Chromatin accessibility signal of each marker peak (rows) in each reference sample and in the ENCODE samples from diverse tissues (columns).

Supplementary Figure 2: A) Comparison of the cell-type proportions estimated by the different deconvolution methods included in the benchmark (y axis) and the true proportions (x axis) in the PBMC experiment samples (top panel) and in the PBMC pseudobulk samples (bottom panel). **B)** Comparison of the cell-type proportions estimated by CIBERSORTx and DeconPeaker using reference profiles built by each tools using our collection of pure ATAC-Seq samples and the true proportions in the PBMC experiment samples (top panel) and in the PBMC pseudobulk samples (bottom panel). **C)** Pearson's correlation coefficient and RMSE values associated to the comparison of the cell-type proportions estimated by each deconvolution tool in each cell type of the PBMC datasets, colors correspond to cell types and shapes to the PBMC dataset. Note that for quanTIseq, gray dots correspond to uncharacterized cells since the prediction of an uncharacterized cell type can not be turned off using the quanTIseq function from the quantiseqr R package. *P*-values resulting from a paired wilcoxon test are represented for each pair of comparison between tools. Significance levels are represented as follow: *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$.

Supplementary Figure 3: Pearson's correlation (left panel) and RMSE values (right panel) associated to the comparison of the cell-type proportions estimated by each deconvolution tool (rows) included in the benchmark and the true cell-type proportions. The comparisons were made in each cell-type (columns) separately. Significance level of the correlations are represented as follow: *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$.

Supplementary Figure 4: A) Comparison of the cell-type proportions estimated by the different deconvolution methods using our reference profiles and the true proportion in the basal cell carcinoma samples (top panel) and in the gynecological cancer samples (bottom panel). In each panel, the top row includes the uncharacterized cells while the bottom row excludes them. **B)** Comparison of the cell-type proportions estimated by CIBERSORTx and DeconPeaker using reference profiles built by each tools using our collection of pure ATAC-Seq samples and the true proportion in the basal cell carcinoma samples (left panel) and in the gynecological cancer samples (right panel). In each panel, the top row includes the uncharacterized cells while the bottom row excludes them.

Supplementary Figure 5: Top panel: RMSE values associated to the comparison of the true proportions and the cell-type proportions estimated by each deconvolution tool in each cell type. For each cell type, all samples from each cancer pseudobulk datasets were considered, *i.e.*, colors correspond to cell types and shapes to datasets. *P*-values resulting from a paired wilcoxon test are represented for each pair of comparison between tools. Significance levels are represented as follow: *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$. Bottom panel: Considering that not all deconvolution tools are able to predict the proportion of uncharacterized cells, the same analyis as in the top panel was performed without considering the uncharacterized cells predictions.

A

Basal cell carcinoma

B

Gynecological cancers

Supplementary Figure 6: Pearson's correlation (left panel) and RMSE values (right panel) associated to the comparison of the cell-type proportions estimated by each deconvolution tool (rows) included in the benchmark and the true cell-type proportions. The comparisons were made in each cell-type (columns) separately. Panel **A** and **B** are associated to the benchmark performed on the basal cell carcinoma dataset and the gynecological cancer dataset respectively. Significance level of the correlations are represented as follow: *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$.

Supplementary Figure 7: **A** Scaled averaged chromatin accessibility of each cell-type specific marker peak in each cell-type obtained based on the ATAC-Seq reference samples used to identify the marker peaks. **B** Scaled averaged chromatin accessibility of each cell-type specific marker peak in each cell-type from the sorted samples excluded from the reference samples (See Material and Methods).

Supplementary Figure 8: Left panels: Scatter plots representing proportions estimated by the different deconvolution tools included in the benchmark (y-axis) as a function of the true proportions (x-axis) for PBMC samples (top row) and the basal cell carcinoma pseudobulks (bottom row). CD4+ and CD8+ T-cells predictions were obtained using the reference profiles based on the major cell-types and the CD4+ and CD8+ T-cell subtypes predictions using the reference profiles including the T-cells subtypes. Right panels: Barplots representing Pearson's correlation values (x-axis) obtained based on the comparison of the predictions and true proportions in each cell-type (y-axis).