In vitro reconstitution reveals membrane clustering and double-stranded RNA recruitment by the enteroviral AAA+ ATPase 2C

Kasturika Shankar[1,2,3,4], Marie N. Sorin[1,2,3,4], Oskar Skoglund[1,2,3,4], Selma Dahmane[1,2,3,4], Himanshu Sharma[1,2,3,4], Josy ter Beek[1,2,4], Louise Nenzén[1,2,3,4], Lars-Anders Carlson[1,2,3,4],*

1. Department of Medical Biochemistry and Biophysics, Umeå University, Umeå, Sweden

2. Wallenberg Centre for Molecular Medicine, Umeå University, Umeå, Sweden

3. Laboratory for Molecular Infection Medicine Sweden (MIMS), Umeå University, Umeå, Sweden

4. Umeå Centre for Microbial Research (UCMR), Umeå University, Umeå, Sweden

*corresponding author: lars-anders.carlson@umu.se

Keywords: enterovirus, picornavirus, poliovirus, AAA+ ATPase, cryo-electron tomography, membrane, in vitro reconstitution, 2C

Abstract

Enteroviruses are a vast genus of positive-sense RNA viruses that cause diseases ranging from
common cold to poliomyelitis and viral myocarditis. They encode a membrane-bound AAA+ ATPase,
2C, that has been suggested to serve several roles in virus replication, e.g. as an RNA helicase and
capsid assembly factor. As an oligomeric peripheral membrane protein, the low solubility of 2C has
hitherto impaired biochemical studies of the full-length protein. Here, we report a biochemical
reconstitution poliovirus 2C association with membranes. We show that a conserved glycine divides
the N-terminal membrane-binding domain of 2C into two amphipathic helix regions, AH1 and AH2.
AH2 is the main mediator of 2C oligomerization, and is sufficient for its membrane binding. AH1 is
the main mediator of a novel function of 2C: clustering of membranes. Cryo-electron tomography
reveal that several 2C copies mediate this function by localizing to vesicle-vesicle interfaces. 2C-
mediated clustering is partially outcompeted by nucleic acids, suggesting a way by which 2C can
switch from an early role in coalescing replication organelles, and recruiting lipid droplets to them, to
a later role where 2C assists RNA replication and particle assembly. 2C is sufficient to recruit RNA to
membranes, with a preference for double-stranded RNA, thus being a candidate for the factor that
localizes viral RNA replication to membranes. Finally, the *in vitro* reconstitution revealed that full-
length 2C localized to a membrane has weak RNA chaperoning activity, dependent on having a free N
terminus, but does not possess ATP-dependent helicase activity. Together, this study suggests two
novel roles for 2C in membrane clustering and double-stranded RNA recruitment to membranes, and
calls into question a role of 2C as an RNA helicase. The reconstitution of functional, 2C-decorated
vesicles provides a platform for further experimental studies into this protein and its roles in
enterovirus replication.

Introduction

The *Enterovirus* genus (family: *Picornaviridae*) are positive-sense single-stranded RNA (+ssRNA) viruses, whose ~7.5kb genome is encapsidated in a 30 nm non-enveloped, icosahedral capsid [1, 2]. There are hundreds of known human enteroviruses, responsible for diseases like poliomyelitis (poliovirus, PV), viral myocarditis (Coxsackievirus B3, CVB3), common cold (Rhinoviruses, RVs) and hand, foot and mouth disease (e.g., Coxsackievirus A16 and Enterovirus 71, EV71) [3-5]. In addition, their high mutation rate and frequent recombinations mean that novel enteroviruses emerge regularly [6-8]. PV, which causes the paralytic disease poliomyelitis, has been a major focus of enterovirus research and worldwide vaccination efforts. Since the launch of the Global Polio Eradication Initiative in 1988, the number of poliomyelitis cases have reduced significantly, but it continues to haunt a few countries as an endemic disease [9, 10]. There is limited progress in the production of vaccines against the many non-polio enteroviruses that cause frequent outbreaks in different parts of the world [11]. Development of pan-enteroviral drugs that inhibit conserved steps of the replication cycle is thus of importance not only for the last phase of polio eradication, but also for minimizing the risk associated with future enterovirus outbreaks, as well as for treating chronic enterovirus infections [12-15].

Enterovirus genome replication is a highly conserved process requiring the formation of cytoplasmic replication organelles (ROs) [16-18]. The ROs are formed from extensively modified cytoplasmic membranes which are turned into an assembly line for viral RNA and virus particles [19, 20]. RO membranes are obtained from several cellular sources. Initially, they come from the disassembly of the Golgi apparatus as well as from utilization of cellular cholesterol [21, 22]. In addition, ROs are enriched by fatty acids from lipid droplets, through RO-LD contacts mediated by the viral protein 2C [23]. At later time points, induction of macro-autophagy (henceforth: autophagy) provides another source of RO membranes in the form of double-membrane vesicles [24-26]. The direct dependence of enterovirus RNA replication on membranes is supported e.g. by electron microscopy autoradiography, showing that newly synthesized RNA co-localizes with RO membranes [27]. Further, subcellular fractions active in viral RNA synthesis contain membranes, and their activity is inhibited if the membranes are disintegrated by detergents [28, 29]. How viral RNA, viral proteins, and host factors arrange spatially on the cytoplasmic face of RO membranes is not known.

The enterovirus genome has one major open reading frame which is translated as a single major polyprotein, with some enteroviruses also having a smaller accessory open reading frame [30]. The major polyprotein is subsequently cleaved into eleven individual proteins by the viral proteases 2A and 3C (Fig. 1A). The non-structural proteins, contained in polyprotein regions P2 and P3, encode functions necessary for RO formation and genome replication (Fig. 1A). Amongst these is the protein 2C, a conserved, multifunctional ATPase of the AAA+ superfamily [31-33]. 2C is essential for both RNA replication and virion assembly, but its precise roles in these processes are unknown [34-38]. 2C has a central ATPase domain followed by a Zn finger and a C-terminal α-helix which engages a hydrophobic pocket of another 2C monomer to aid its oligomerization [39-42] (Fig. 1A). ATP hydrolysis by 2C is stimulated by its oligomerization [41, 43], and by binding to single-stranded RNA [44]. Interestingly, even though homologous proteins from other viruses are helicases, most publications report that 2C only has ATP-independent RNA chaperoning activity, and little-to-no ATP-dependent RNA helicase activity[44, 45]. However, it must be noted that biochemical studies on 2C have to date been carried out in simplified systems, using e.g. truncated 2C constructs, fusion proteins, and/or omitting the membrane platform on which 2C is located during infection.

The conserved, N-terminal membrane-binding domain (MBD) of 2C is necessary for virus replication [46]. It has been suggested to contain an amphipathic helix, and has been shown to be necessary and sufficient for membrane association when overexpressed in cells [46-48]. The MBD has further been described to mediate the hexamerization of poliovirus 2C [43], while the C-terminal α-helix appears sufficient to mediate hexamerization of 2C from some other enteroviruses [40]. Due to the

hydrophobic nature of the MBD, full-length 2C tends to aggregate during purification, thus prohibiting
its use in biochemical experiments. This problem has typically been circumvented by working with a
truncated 2C construct lacking the MBD [39-41], or by using high concentrations of detergents [44].
Alternatively, fusing the N-terminus of full-length 2C to the large, hydrophilic maltose-binding protein
(MBP) has also been reported to result in a soluble protein [43]. While all of these studies have
uncovered important aspects of 2C's function, they were not able to address the mechanisms of its
membrane binding, nor its function as a full-length protein on a membrane.

Here, we biochemically reconstituted the binding of purified, full-length poliovirus 2C to synthetic
vesicles. We suggest that the MBD is in fact composed of two amphipathic helix segments, AH1 and
AH2, interrupted by an absolutely conserved glycine residue. Both AH1 and AH2 are necessary for
complete hexamerization of 2C, whereas AH2 is sufficient for its membrane localization. A novel
function for AH1 is identified in tethering membranes together, and we show that 2C's membrane
tethering can be partially outcompeted by nucleic acid. We show that 2C is sufficient to recruit and
retain RNA to membranes, and that it has a strong preference for retaining double-stranded RNA at a
membrane. Together, these findings provide a biochemical basis for several roles of 2C in enterovirus
replication, and establishes a more realistic biochemical system as a basis for further investigations.

Results

The oligomerization state of 2C is controlled by its N-terminal, bipartite amphipathic helix

The membrane-binding domain (MBD) of 2C (Fig. 1A) consists of residues ~1-40 and is suggested to
be an amphipathic helix [46-48], but details of its structure are not known. We took a computational
approach to studying the MBD across enteroviruses. A multiple sequence alignment revealed that
several hydrophobic residues (W4, L5, F8, G17, W20, I25, F28, I29, W31, L32 and I36), polar
residues (S3, K6, K7, K24, K27, and K35) and P38 are broadly conserved between enterovirus species
(Fig. 1B). We next studied the amphipathic nature of the 1-40 region by calculating its hydrophobic
moment (μH) in a sliding window of 18 residues. A high value for hydrophobic moment means that a
helix is amphipathic (i.e. that one side is more hydrophobic than the other). The calculation suggested
that the MBD may, in fact, consist of two separate amphipathic helices, with the space between the
amphipathic regions coinciding with the conserved residue G17 (Fig. 1C). The amphipathic nature of
these two regions was corroborated by a helical wheel plot (Fig.S1A). Since glycine is a helix breaker
residue, we hypothesized that G17 in fact divides the MBD into two amphipathic helices, AH1 and
AH2. AlphaFold2 [49] structure predictions of 2C(1-40) of several enteroviruses indeed suggested that
a kink at G17 is a conserved structural feature (Fig. 1D). In addition, the structure predictions indicate
that AH1 has a hydrophobic surface which is offset from AH2 [50] (Fig. S1B). Together, these data
suggest that the MBD of 2C consists of two amphipathic helices, AH1 and AH2, and we thus set out
to study their individual roles in the protein.

Recombinant full-length 2C has been reported to be insoluble in the absence of detergents unless its
N-terminus is fused to a solubility-enhancing tag [39, 44]. We thus purified the N-terminally
His$_6$MBP-tagged, full-length poliovirus 2C protein (residues 1-329) and its N-terminally truncated
variants 2C(ΔAH1) (residues 12-329), 2C(ΔMBD) (residues 41-329) and 2C(Δ115) (residues 116-
329) (Fig. 2A, B, Fig. S2). The two first constructs correspond to the truncation of AH1, and AH1-2,
respectively, whereas the third construct corresponds to a soluble fragment which has previously been
crystallized [39]. Full-length 2C is a hexamer [39, 41, 43, 51]. We wanted to study the individual
roles of the AH1 and AH2 regions in hexamer assembly. Notably, every N-terminal truncation led to a
gradual shift towards higher elution volumes in size exclusion chromatography (Fig. 2B), indicating
that every part of the MBD plays a role in 2C oligomerization. To further dissect the role of AH1 and
AH2, we determined the molecular mass distribution of full-length and truncated 2C constructs. Mass

photometry of MBP-2C and MBP-2C(ΔAH1) at 100 nM revealed presence of monomeric protein, but
also high-mass peaks (Fig. 2C,D). For the full-length protein, the high-mass peak closely corresponds
to a hexamer, whereas the 2C(ΔAH1) high-mass peak corresponds to a pentamer, or e.g. a mix of
hexamer and a smaller species. The shorter constructs 2C(41-329) and 2C(116-329) were stable in
solution even after removal of the MBP tag, and their masses were analyzed at 38 μM and 134 μM,
respectively, by size exclusion chromatography coupled to multi-angle light scattering (SEC-MALS).
The mass estimates throughout the predominant peak for two consecutive runs were 43 kDa and 52
kDa for 2C(41-329), i.e. a mass intermediate between monomer (at 32.8 kDa) and dimer (Fig. 2E),
possibly due to an equilibrium between the two states. The predominant peak for two SEC-MALS
runs for 2C(116-329) gave an estimated mass of 27 kDa and 22 kDa, consistent with the expected 23.8
kDa of a monomer (Fig. 2F). The MBP-fused versions of these constructs both had SEC-MALS mass
estimates corresponding to monomers (Fig. S3A-B).

Taken together, the conserved glycine 17 divides the MBD into two amphipathic helices, AH1 and
AH2, with offset hydrophobic faces. Both AH1 and AH2 are important for hexamerization of 2C in
solution, and the deletion of the entire MBD only leaves 2C with a weak propensity for dimerization.

The AH2 region mediates charge-independent binding of 2C to membranes

We next wanted to study the individual roles of AH1 and AH2 in 2C's membrane binding. To this
end, we reconstituted the process *in vitro* using synthetic liposomes, and the purified 2C constructs
described above. We devised an assay in which MBP-fused 2C is first mixed with small or large
unilamellar vesicles (SUVs and LUVs, respectively), after which TEV protease is added to cleave off
the MBP tag and allow 2C to bind the liposomes (Fig. 3A). The 2C-decorated liposomes are then
subjected to a flotation assay, in which the liposome-containing upper fraction and liposome-free
lower fraction are separated. 2C and liposomes are labelled with different fluorophores (ATTO488
maleimide conjugated to cysteines, and ATTO647N conjugated to DOPE, respectively), allowing
quantification of 2C and lipids in the fractions (Fig. 3A). The lipid fluorescence readings in the upper
fraction were consistent across all experiments, allowing for quantitative comparisons of the amount
of protein bound to liposomes (Fig. S4). We first used SUVs composed of net uncharged lipids: 75
mol% 1-palmitoyl-2-oleoyl-glycero-3-phosphocholine (POPC), 25 mol% 1-palmitoyl-2-oleoyl-sn-
glycero-3-phosphoethanolamine (POPE). The SUVs had an average size of ~30 nm as determined by
dynamic light scattering (DLS) (Fig. S5). The addition of MBP-2C and TEV to the SUVs led to the
formation of two distinct SUV bands in the upper fraction (Fig. 3B, Fig. S6A). 88±11% of full-length
2C were found in the upper fraction, with most protein in the uppermost of the two bands (Fig. S6B).
MBP-fused 2C was able to associate with SUVs to a similar degree as cleaved 2C, indicating that the
MBD is exposed, also in the MBP fusion protein. However, later experiments would show that
removal of the MBP tag was necessary for functional association of 2C with membranes. Upon
deletion of AH1, 81±16 % of 2C(ΔAH1) was still found in the upper fraction, whereas deletion of the
entire MBD (AH1-AH2) led to only 10±7.4 % of 2C(41-329) being found in the upper fraction (Fig.
3C). Thus, the AH2 region is the main mediator of 2C membrane association.

We then wished to study if 2C has a preference for a given membrane charge or curvature. In side-by-
side flotation experiments, we quantitated 2C binding to the POPC/POPE (net uncharged) SUVs
described above, and SUVs consisting of 50 mol% phosphatidyl serine (POPS, net charge -1). Full-
length 2C and 2C(ΔAH1) showed no preference for the negatively charged POPS, whereas the binding
of 2C(41-329) increased from 10±7.4 % to 38±7% when POPS was included in the SUVs (Fig. 3D).
To determine if 2C has a membrane curvature preference, we compared 2C binding to the sonicated
POPC/POPE SUVs, with an average size of ~50 nm, and larger (and hence lower-curvature) LUVs
with the same lipid composition and an average diameter of ~2500 nm. Full-length 2C and 2C(ΔAH1)
showed a slight decrease in the binding to the lower-curvature LUVs (Fig. 3E), but this decrease was

no longer significant after normalizing for the amount of lipid present in the top fraction (Fig. S4F).In
summary, these data show that the AH2 region mediates strong 2C membrane binding, which is
neither dependent on high membrane curvature nor on negatively charged lipids.

2C mediates lipid clustering, which is enhanced by the AH1 region

In virus-infected cells, 2C is present at contact sites of ROs and lipid droplets[23], and we suspected
that the high-density band of 2C-coated SUVs in our flotation experiments may be an *in vitro*
recapitulation of 2C-mediated lipid tethering (Fig. 3B, Fig. S6B). We used DLS to investigate lipid
clustering by 2C. As described above, the MBP tag was cleaved from 2C in the presence of SUVs.
After a 30 min incubation, the size distribution of the sample was measured using DLS. The sonicated
SUVs were around 40 nm in size, whereas the addition of TEV protease and 0.8 μM MBP-2C to these
SUVs led to the formation of vesicle clusters with an average diameter of ~1 μm (Fig. 4A). 2C(ΔAH1)
also clustered SUVs (Fig. 4A). Defining a liposome cluster as a particle with a diameter of >100 nm(
2.5 times the average SUV size), we compared the percentage of clusters in reactions containing
different concentrations of 2C and 2C(ΔAH1). Already at 0.8 μM protein – the concentration used for
the flotation assays – full-length 2C generated significantly more clusters than 2C(ΔAH1) (11.5±4.0%
and 4.6±1.7%, respectively, Fig. 4B). At 3 μM 2C, the difference was even more pronounced, with
47.0±13% of liposomes being present in clusters with 2C and 12±13% for 2C(ΔAH1) (Fig. 4B). To
characterize the 2C-mediated vesicle clusters at nanometer resolution, we then performed cryo-
electron tomography (cryo-ET) on the 3 μM reaction with full-length 2C. This allowed direct
visualization of how 2C tethers liposomes together to form larger clusters (Fig. 4C). Typically,
multiple 2C densities were present on the imaged SUVs, and they were mostly found at the interface
between SUVs (Fig. 4C). Taken together, 2C clusters liposomes *in vitro*, and the AH1 region is
necessary for effective clustering.

**Nucleic acid binding competes with 2C membrane clustering, and 2C preferentially recruits
double-stranded RNA to membranes**

The enterovirus genome is replicated on the cytoplasmic face of the RO membrane [19, 52]. Since 2C
is a AAA+ protein and a putative RNA helicase [33, 36, 42, 44, 45, 53-56], we wondered whether 2C
is sufficient to recruit and retain RNA at the membrane, and whether nucleic acid binding may affect
2C's lipid clustering activity. To address the second question, we repeated the lipid clustering
experiment using full-length 2C, in the presence of increasing concentrations of nucleic acid. It was
previously reported that both DNA and RNA bind at the same site on 2C, and that initial binding is
dependent on the phosphodiester backbone [44]. Hence, we used single-stranded DNA (ssDNA)
oligonucleotide and measured the amount of clustering using DLS (Fig. 5A). Upon increasing the
concentration of ssDNA there was a consistent decrease in the percentage of clustered lipids, albeit
only statistically significant at 100 μM ssDNA concentration (Fig. 5B). This indicates that nucleic acid
binding partially outcompetes the membrane tethering activity of 2C. To directly test the hypothesis
that 2C recruits RNA to the membrane, and to investigate if it possesses any selectivity, we devised an
alternative SUV flotation assay. In this assay, 2C was no longer fluorophore-labelled. Instead, a
fluorophore-labeled RNA (single-stranded or double-stranded) was added in the first step of the
flotation assay, allowing quantitation of the amount of RNA co-migrating with the SUVs (Fig. 5C,
Fig. S7). 2C was able to recruit both ssRNA and dsRNA to SUVs and retain it at the membrane
throughout the ultracentrifugation of the flotation experiment (Fig. 5D). At identical starting
concentrations, dsRNA was recruited significantly more than ssRNA (Fig. 5D). The recruitment was
strictly dependent on 2C having a free MDB since omission of TEV protease completely abrogated the
RNA recruitment (Fig. 5D). Taken together, 2C recruits and retains nucleic acid to membranes, with a
preference for dsRNA over ssRNA, and nucleic acid binding reduces 2C-mediated membrane
clustering.

Membrane-bound, full-length 2C is an ATPase and an ATP-independent RNA chaperone

Several studies have sought to determine whether 2C is an active RNA helicase, but they were generally conducted with protein that was truncated and/or not membrane-associated. Having reconstituted the binding of full-length 2C to membranes, and shown that it recruits RNA to membranes, we set out to use this more realistic system to revisit the question of 2C's helicase activity. We conducted an RNA helicase assay using 2C and 2C(41-329) in the presence of SUVs. The MBP-fused proteins were incubated with a dsRNA with 5' overhang, and either with or without TEV protease and in the presence or absence of ATP. The unwinding of dsRNA to ssRNA was monitored on a native polyacrylamide gel (Fig. 6A). This revealed that the release of 2C's N terminus from MBP y the TEV protease was absolutely required for any RNA unwinding activity (Fig. 6A-B). With the N terminus released, both 2C and 2C(41-329) showed partial RNA unwinding activity in the absence of ATP, indicative of ATP-independent RNA chaperoning activity (Fig. 6A-B). However, the addition of Mg^{2+}-ATP did not lead to increased unwinding of the dsRNA as expected from an RNA helicase. In fact, the presence of Mg^{2+}-ATP instead led to less conversion of dsRNA to ssRNA (Fig. 6A-B). This was consistent between full-length and truncated 2C, and the same was also seen when a dsRNA with 3' overhang was used (Fig. 6C-D). In stark contrast, when the ATPase activity of 2C was measured under similar conditions, only the full-length protein was found to have appreciable ATPase activity, and this was independent of whether it was fused to MBP, or if there was a membrane present (Fig. 6E). Notably, this correlates with the conditions for hexamerization of 2C (Fig. 2) but not with the conditions under which 2C displayed RNA chaperoning activity (Fig. 6E). In summary, membrane-bound full length 2C is an active ATPase, but its RNA chaperoning activity is not ATP-dependent. The ATPase activity appears to correlate with the hexamerization of 2C, whereas the ATP-dependent RNA chaperoning activity correlates with the liberation of its N-terminus.

AH1 contributes to viral RNA replication to a similar degree as the entire MBD

The biochemical reconstitution allowed us to study the separate contributions of AH1 and AH2 to processes such as membrane binding and membrane clustering. To investigate their relative contribution to viral RNA replication in cells, we used poliovirus replicons where the capsid proteins were replaced by Renilla luciferase. As opposed to the biochemical experiments, the 2C truncations in replicons (Δ(2-11) and Δ(2-40)) retained the staring glycine of 2C since it is a core part of the viral protease 3C cleavage site (Fig. 7A). 12 hours after transfecting replicon RNAs into cells, they were harvested, and the luciferase signal was quantified. The luminescence from both the Δ(2-11) and Δ(2-40) replicons was significantly reduced, by approximately two orders of magnitude for the replicon containing unaltered, full-length 2C. Thus, although AH1 appeared to have more subtle roles in the biochemical experiments (e.g. in vesicle clustering), it is indispensable for effective viral RNA replication in cells.

Discussion

With fewer than 7,000 coding nucleotides, the *Enterovirus* genome can sustain a full viral replication cycle by encoding multifunctional proteins, and by hijacking host-cell functions. A universally present and highly conserved enteroviral protein is the AAA+ ATPase 2C, whose suggested roles range from membrane remodeling to RNA helicase and capsid morphogenesis. The nature of 2C as an oligomerizing, peripheral membrane protein has impeded biochemical studies, that have typically been conducted on truncated protein not engaging a membrane. Here we described the *in vitro* reconstitution of purified, full-length 2C's binding to membranes. We used this more realistic, yet

completely defined, system to investigate several functions of 2C. In summary, our findings are
consistent with the model shown in Fig. 8.

We showed that the N-terminal MBD is divided into two amphipathic helix segments, AH1 and AH2,
separated by a highly conserved glycine residue (Fig. 1C, Fig. S1). AH2 appears to be the main
mediator of 2C's hexamerization in solution (Fig. 2), and is sufficient for membrane binding (Fig. 3C).
Of note, removal of the N-terminal maltose-binding protein (MBP) tag was not necessary for 2C to
bind membranes in our assay, but it was strictly required for 2C to recruit RNA to membranes (Fig.
5D), and for its RNA chaperoning activity (Fig. 6A). Whereas the N-terminal MBP tag is an artificial
addition to aid purification of recombinant 2C, the fusion of 2C's N-terminus to another protein is, in
fact, relevant to infection. 2C is initially excised from the viral polyprotein in the form of the stable
precursor 2BC, in which the N terminus of 2C is still fused to the viral protein 2B [57]. Hence, our
finding that 2C only binds RNA after its N terminus is liberated may be indicative of what happens in
an infected cell when 2BC is processed to 2B and 2C.

Our study revealed a novel role for 2C in clustering vesicles, a role mediated by AH1 (Fig. 4). This
finding suggest a biochemical mechanism for the 2C-mediated recruitment of lipid droplets to
replication organelles (ROs)[23]. 2C-mediated membrane clustering may also be involved in
generating the tightly stacked membrane tubes that make up the ROs at early time points [20, 58].
Notably, the deletion of AH1 led to a stark reduction of poliovirus genome replication, similar in
effect to the deletion of the entire MBD (Fig. 7). This may indicate that the lipid-clustering activity of
2C is central to enterovirus replication, although it cannot be ruled out that AH1 deletion also affects
other aspects of the replication such as polyprotein processing. The decrease in 2C-mediated
membrane clustering seen with increasing nucleic acid concentration (Fig. 5A-B) suggests a balance
between 2C's role in membrane clustering at early time points, and a preferential role in RNA
replication and particle assembly at later time points (Fig. 8). In this context, the ability of 2C to bind
membranes independent of curvature and lipid species (Fig. 3D-E) fits its multitude of roles in binding
to different types of RO membranes and lipid droplets.

The localization of viral RNA replication to RO membranes is well established, but the molecular
mechanism behind the localization is poorly understood. An electrostatic interaction between the viral
polymerase 3D and anionic lipids may be part of the mechanism[59]. Here, we show that 2C is able to
recruit and retain RNA at membranes, with a selectivity for dsRNA (Fig. 5D). Long double-stranded
segments are not prevalent amongst cellular RNAs, but it is the form of the viral template RNA while
replicated. Thus, 2C could selectively localize viral template RNA to the ROs. On the other hand, full-
length, membrane-bound 2C had no ATP-dependent RNA helicase activity, and only weak ATP-
independent RNA chaperone activity (Fig. 6A-D). This is in line with most previous studies on
truncated and membrane-free 2C, and with structural models of the 2C hexamer indicating that its
central pore would be too narrow and have an incompatible electrostatic potential to support RNA
translocation [40, 41, 44]. The conditions for 2C ATPase activity were different from the conditions
that permitted RNA chaperoning: effective ATPase activity required the MBD, but did not require
removal of the N-terminal MBP tag (Fig. 6E). One possible interpretation of these data is that 2C
ATPase activity requires hexamerization, whereas its RNA binding, and its weak RNA chaperoning
activity, requires a free N-terminus. A recent study found that MBD-deleted 2C has RNA-stimulated
ATPase activity, which would be consistent with the above interpretation of our data if assuming that
RNA engagement restores hexamerization of 2C(41-329). The convergence of our work and previous
studies in suggesting that 2C is not a helicase can be interpreted in the light of the reported dsRNA-
unwinding activity of the polymerase 3D [60].

Taken together, our reconstitution of 2C membrane binding allowed us to shed light on several
functions of 2C: it is able to bind vesicles with no apparent preference for curvature and lipid
composition, it can cluster vesicles, and selectively recruit dsRNA to their surface without possessing
helicase activity. In spite of this, major questions remain unanswered about the role of 2C in

enterovirus RNA replication and particle assembly. Nothing is known about the supramolecular
arrangement of the RNA replication machinery on membranes, for instance how 2C, the polymerase
3D and host RNA-binding proteins such as PCBP1/2 are arranged [61]. Also, 2C is known to play a
role in capsid assembly and is suggested to be part of a "tether complex" enabling the RNA loading of
new capsids on RO membranes [19, 58], but biochemical mechanisms underlying this process are
largely unknown. The reconstitution of 2C-decorated vesicles presented here is a stepping stone
towards a more complete *in vitro* reconstitution of such complex processes.

Materials and methods

Expression and purification of poliovirus 2C

A codon-optimized synthetic gene corresponding to full-length 2C from poliovirus type 1 Mahoney
(GenBank: V01149.1) was inserted into 1M expression plasmid from Macrolab (University of
California, Berkeley, USA) with a TEV protease-cleavable, N-terminal His$_6$-MBP tag. The plasmid
was transformed to *E. coli* BL21(DE3) competent cells and grown overnight on LB-agar plates
containing 50ug/ml Kanamycin (KAN). A single colony was used to inoculate LB/KAN media, this
primary culture was grown for 5 hours. Then, a one-liter culture of LB/KAN was inoculated with 1ml
of the primary culture. The cells were grown at 37°C, induced at $OD_{600}=2$ by the addition of 0.5 mM
isopropyl-beta-D-thiogalacopyranoside, then incubated overnight at 25°C. Cells were collected by
centrifugation, and resuspended in lysis buffer (50 mM Tris, pH 8.5), 500 mM NaCl, 20 mM
Imidazole, 10% (v/v) glycerol, 36 μM NP40, 5 mM MgCl$_2$, 0.1 mM THP) containing protease
inhibitor cocktail (40 mM Phenylmethylsulfonyl fluoride, 1 mM Leupeptin and 200 mM Benzamidine
HCl) and DNase. The cells were lysed using a continuous flow cell disruptor (Constant Systems). The
lysate was cleared by centrifugation at 63,000*g at 4°C for 1 hour and the supernatant was loaded onto
Ni-NTA (nitrilotriacetic acid) resin and eluted with elution buffer (50 mM Tris (pH 8.5), 500 mM
NaCl, 250 mM Imidazole, 10% (v/v) glycerol, 36 μM NP40, 5mM MgCl$_2$, 0.1 mM THP). The protein
was then loaded onto a HiTrap Q HP Column (GE Healthcare) and eluted using a gradient from buffer
QA (50 mM Tris, pH 8.5), 50 mM NaCl, 10% (v/v) glycerol, 36 μM NP40, 5 mM MgCl$_2$, 0.1 mM
THP) and buffer QB (50 mM Tris (pH 8.5), 1 M NaCl, 10% (v/v) glycerol, 36 μM NP40, 5 mM
MgCl$_2$, 0.1 mM THP). The fractions containing the protein was loaded onto a 24 ml Superose 6
Increase 10/300 GL (Cytiva) that was equilibrated with SEC buffer (50 mM Tris-HCl (pH 7.4), 150
mM potassium acetate, and 0.1 mM THP). Purified 2C was then aliquoted, flash frozen with liquid N$_2$,
and stored at -80°C. All the other constructs of the protein were purified in a similar way.

To prepare 2C(41-329) and 2C(116-329) without His$_6$-MBP tag, the tagged proteins were diluted to
0.1mg/ml after the HiTrap Q HP column using buffer QA described above and the proteins were
cleaved at 4°C overnight by adding TEV protease in a molar ration TEV:2C of 1: 5 and 1:15 for
2C(41-329) and 2C(116-329), respectively. Uncleaved protein and His$_6$-MBP were removed by
incubating the mixture with NiNTA resin followed by passing of the protein over a 1ml MBPTrap
(Cytiva) column and then a HiTrap Heparin HP (Cytiva) using buffer QA and QB. The fractions
containing the protein were loaded onto a Superdex 75 Increase 10/300 GL (Cytiva) equilibrated with
SEC buffer. Purified cleaved 2C was aliquoted, flash frozen with liquid N$_2$, and stored at -80°C.

Fluorophore labeling of proteins

To label the proteins, after the HiTrap Q HP Column, the proteins were incubated with an equimolar
amount of Atto488 maleimide (Sigma-Aldrich) at 4°C overnight. Next day, excess dye was quenched
using 25 mM Dithiothreitol (DTT) and the proteins were loaded onto a Superose 6 Increase 10/300 GL
(Cytiva) column, pooled and snap frozen as described above for unlabeled protein. The labeling
efficiency was estimated by measuring the absorption of the labeled protein at 280 nm and at the
fluorophore absorption maximum. It was typically 0.5-1.1 fluorophores/2C monomer.

SEC-MALS

All the samples were run on an ÄKTA Pure (Cytiva) with a Superdex 200 Increase 10/300 GL column, equilibrated in SEC buffer, coupled to a light scattering (Wyatt Treas II) and refractive index (Wyatt Optilab T-Rex) detector. Bovine serum albumin (BSA) was used to calibrate the system. 150 to 500 µl (0.6 to 1 mg/ml) of protein was injected at a flow rate of 400 µl/min. Data were analyzed using Astra software (version 7.2.2; Wyatt Technology).

Mass photometry

Proteins were diluted to 100 nM in 50 mM Tris pH 7.4 and applied to a glass coverslip and recorded on Mass Photometer[62] (OneMP, Refeyn Ltd) for 60 seconds. The data were analyzed using DiscoverMP (Refeyn Ltd) and the molecular weight was obtained by contrast comparison with known mass standards measured on the same day and were repeated three times, and visualized as mean +/- standard deviation.

Multiple sequence alignment

Multiple sequence alignment of amino acids 1-40 was performed on EV71 (GenBank: AAB39969.1), CVB3 (GenBank: AFD33642.1), PV type 1 (MAHONEY STRAIN) (GenBank: V01148.1), EV D68 (GenBank: ALE66301.1), RV A2 STRAIN USA/2018/CA-RGDS-1062 (GenBank: MN228695.1) and RV C15 (GenBank: ACZ67658.1) using Clustal Omega[63] with default parameters.

Amphipathic helix prediction

The physicochemical properties 2C amino acids 1-40 from EV71 (GenBank: AAB39969.1), CVB3 (GenBank: AFD33642.1), PV type 1 (Mahoney strain) (GenBank: V01148.1), EV D68 (GenBank: ALE66301.1), RV A2 STRAIN USA/2018/CA-RGDS-1062 (GenBank: MN228695.1) and RV C15 (GenBank: ACZ67658.1) were determined using heliquest (https://heliquest.ipmc.cnrs.fr/) [64] using alpha-helix as helix type and window size of one turn (18 residues). The mean amphipathic moment or transverse hydrophobic moment (μH) was plotted against the amino acid at the center of the analysis window.

Structure predictions

All structures were predicted using Alphafold 2 implemented in COLABFOLD ((https://colab.research.google.com/github/sokrypton/ColabFold/blob/main/AlphaFold2.ipynb) [49] with default settings. The cartoon representations were generated using CHIMERAX [50].

Preparation of small unilamellar vesicles (SUVs)

1-palmitoyl-2-oleoyl-sn-glycero-3-phospho-L-serine (POPS), 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphoethanolamine (POPE) and 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (POPC) were purchased from Avanti Polar lipids. Total 1 mg/ml of lipids containing POPC (25 mol %), POPE (25 mol %), POPS (25 mol %) or POPC (75 mol %), POPE (25 mol %), and 1, 2-Dioleoyl-sn-glycero-3-phosphoethanolamine labeled with Atto 647N (0.1 mol %) was dried onto a borosilicate glass tube wall with a nitrogen stream and kept overnight in vacuum desiccator for complete removal of solvents. 500 µl of SEC buffer was added into the mixture. The tube was then vortexed to resuspend the lipids and then, kept in a water bath at 42°C for 15 min. Then, the lipids were transferred to an Eppendorf tube for sonicating with a tip sonicator (10 seconds pulse, 20 seconds rest, and 20% amplitude) until a clear solution was observed. Then, the tube was centrifuged at 21,000*g at 4°C for 20 min. Then, 300 µl of supernatant was carefully removed, and the quality of the preparation was checked using dynamic light scattering as described below. SUVs were kept at 4°C until use, at most for 3 days.

Preparation of large unilamellar vesicles (LUVs)

Total 1 mg/ml of lipids containing POPC (75 mol %), POPE (25 mol %) and1, 2-Dioleoyl-sn-glycero-
3-phosphoethanolamine labeled with Atto 647N (0.1 mol %) was dried onto a borosilicate glass tube
wall with a nitrogen stream and kept overnight in vacuum desiccator. Then, 500 μl of SEC buffer was
added into the mixture. The tube was then vortexed to resuspend the lipids and to break the multi-
lamellar vesicles, the lipids were subjected to 12 freeze-thaw cycles by moving it between a 42°C
water bath and liquid nitrogen. Then, this solution was passed 21 times through an extrusion
membrane (Whatman Nuclepore) with pore size of 2 μm on a mini-extruder (Avanti Polar Lipids) to
get a clear solution comprising of LUVs. The quality of the preparation was checked using Dynamic
light scattering as described below. LUVs were kept at 4°C until use, at most for 3 days.

Vesicle flotation assay

A reaction mixture of 100 μl was prepared containing 0.8 μM Atto488 maleimide His$_6$MBP-2C, 1
mg/ml SUVs or LUVs and X μM TEV protease was incubated for 2 hours at room temperature. Then,
this reaction mixture was mixed with 500 μl of 48% (w/v) OptiPrep density gradient medium (Merck
(Sigma-Aldrich)) and this mixture was carefully placed at the bottom of an Open-Top Thin wall Ultra-
Clear Tube (Beckman Coulter) and was overlaid with 800 μl of 30% (w/v) OptiPrep followed by
another layer of 800 μl of 20% (w/v) OptiPrep and finally 2 ml of SEC buffer. The tubes were then
ultra-centrifuged at 364,000*g for 3 hours at 4°C. After centrifugation different fractions were
carefully taken out from the tubes and fluorescence was measured using Agilent BioTek Synergy
fluorimeter.

Flotation assay with fluorescently labeled RNA

All oligonucleotides used in this study are listed in Table 1. 100 μl of reaction mixture was prepared as
described above, except that unlabeled 2C was used, and a final concentration of 500 nM of single-
stranded 5' TET-labeled RNA(5'CCUCUAACCACAGUCUGAUC 3') (Eurofins Genomics) or
dsRNA (described below) was added. The flotation assay was performed as described above and the
TET fluorescence was measured. To prepare dsRNA, 10uM of 5' TET labelled RNA
(5'CCUCUAACCACAGUCUGAUC 3') was mixed with its 10uM unlabeled complementary strand
in a buffer containing 50mM HEPES pH7.2, 100mM KCl and RNase secure. The mixture was heated
to 95°C and then the temperature was decreased by 0.5°C /minute until 22°C. The annealed dsRNA
was then placed overnight at 4°C and then at -20°C until use. The extent of annealing was checked
using a 16% acrylamide non-denaturing RNA gel. A standard curve was prepared using fluorescence
reading for ssRNA and dsRNA and the extent of binding of RNA to 2C was quantified using this
standard curve.

Dynamic light scattering

100 μl of reaction mixture was prepared containing 0.8 μM or 3 μM 2C, 1 mg/ml SUVs, and TEV
protease was incubated for 30 min at room temperature. Then, this mixture was transferred into a
cuvette and the particle size was determined by dynamic light scattering using a Zetasizer Nano-ZS
(Malvern Instruments, Worcestershire, UK) at room temperature. To check the effect of ssDNA on
clustering, increasing concentration of ssDNA oligo (5' TGAATGCGGCTAATCCCAACCT 3')
(Eurofins Genomics) was added in to the mix containing TEV protease, 1 mg/ml SUVs followed by
addition of protein and the particle size was determined as described above.

Sample preparation for cryo-electron tomography

456 A reaction mixture of 100 μl was prepared containing 3 μM His$_6$MBP-2C, 1 mg/ml SUVs and 4 μM
457 TEV protease was incubated for 30 min at room temperature. 3 μl of this mixture was blotted on a
458 glow-discharged Quantifoil R 2/1 Copper 300 mesh grid (Electron Microscopy Sciences) and plunge-
459 frozen in liquid ethane using a Vitrobot plunge freezer (ThermoFisher Scientific) at 22°C, 80%
460 humidity, with blot force -20, blot time 2 seconds and wait time 40 seconds.

Cryo-electron tomography data collection and processing

Tomographic tilt series were collected using a Titan Krios (ThermoFisher Scientific) operated at 300 kV, with a 100 μm objective aperture, Selectris energy filter and Falcon 4i detector (ThermoFisher Scientific). Tilt series were acquired in dose-symmetric mode, using SerialEM version 4. The following parameters were used for acquisition: 26,000 x nominal magnification corresponding to a specimen pixel size of 4.7 Å; defocus range of -3 μm to -8 μm, $\pm$ 63° tilt-range with tilt increment of 3°; total electron dose ~116 e$^-$/Å^2. At each tilt angle, the exposure was saved as an EER movie. The EER files were converted to TIFF files using IMOD [65], gain-reference corrected, and motion corrected using MotionCor2 [66]. The tilt series were then processed and aligned using IMOD[65]. CTF was estimated using CTFFIND4 [67] and corrected using phase flipping as implemented in IMOD's CTFPHASEFLIP.

The isotropic resolution and signal-to noise ratio of the tomograms were improved using the de-noising deep-learning software IsoNet [68]. The filtered tomograms were used for segmentation using Amira (Thermo Fisher Scientific) for the representation of SUVs and 2C.

Colorimetric ATPase assay

Malachite Green Phosphate Assay Kit (MAK307, Sigma-Aldrich) was used for the colorimetric ATPase assay. For the enzymatic reaction 1.2 μM of His$_6$MBP-2C (1-329) or (41-329), 2.9 μM TEV protease, 0.5 mg/ml SUV (POPC (25 mol %), POPE (25 mol %) and 0.4 mM ATP was mixed in buffer containing 40 mM HEPES pH7.5, 2 mM DTT and 12 mM NaCl and was incubated for 2 hours at 37 °C. The samples were diluted 20 times and plated in a 96 well plate and mixed with 20 μl working reagent supplied with the kit. The mixture was incubated at 37°C for 30 min after which the absorbance was measured at 620 nm according to manufacturer's instructions. A standard curved was prepared using the provided standard phosphate in same buffer as the reaction. This standard curve was used to determine the phosphate liberated in the ATPase assay.

Double-stranded RNA unwinding assay

The single stranded RNA (ssRNA) oligos (Eurofins Genomics) were annealed in molar ratio 1:1.5 (28 nucleotide labeled oligomer (5' AAAAAAAAAAAACCAGGCGACAUCAGCG-FAM3'): 16 oligomer un-labeled (5'CGCUGAUGUCGCCUGG3')) in a buffer containing 10 mM HEPES-NaOH (pH 7.2) and 10 mM KCl. This mix was heated to 95°C for 1 minute and then cooled down to 22°C at the rate of 1°C per minute. The sample was then kept at 22°C for one hour. The thus prepared dsRNA with 5' overhang was kept at 4°C overnight before freezing at -20°C. The dsRNA with 3' overhang was made similarly using 28 nucleotide labeled oligomer (FAM-5'CCAGGCGACAUCAGCGAAAAAAAAAAAA3') and mixing with 16 oligomer un-labeled (5'CGCUGAUGUCGCCUGG3'). Chaperone activity assay was performed in reaction buffer containing 40 mM HEPES pH7.5, 2 mM DTT and 15mM NaCl, 0.5 μl RNAse inhibitor, 0.8 mg/ml of SUVs (POPC (25 mol %), POPE (25 mol %) (the SUVs were prepared using the reaction buffer), 1.2 μM MBP-2C , 3.8 μM of TEV protease, 40 nM dsRNA template, 800 nM ssRNA trap strand (28 nucleotide un-labeled oligomer). This mixture was incubated at 37°C for 20 min, followed by addition of 3.5 mM magnesium-ATP to start the reaction. The reaction was incubated for 2 hours before adding 5 μl of quenching buffer consisting of 100 mM Tris pH 7.5, 0.1% bromophenol blue, 1% SDS, 50 mM EDTA, and 50% glycerol. The samples were run on 16% native polyacrylamide gels for 2 hours at 100 V and scanned using an Amersham Typhoon fluorescence gel scanner. The gel scans were quantified using ImageJ, measuring the intensity of the bands at the ssRNA position as defined by the ssRNA control and subtracting the intensity of the dsRNA control as background.

Luciferase assay

The P2-P3 region from poliovirus type 1 Mahoney (GenBank: V01149.1) was inserted into a vector containing *Renilla* luciferase reporter on the N-terminus. Truncated 2C versions (Δ2-11 and Δ2-40) were also constructed. The first glycine residue of 2C was not truncated in order to preserve the 3C protease cleavage site. The plasmids were linearized using ApaI restriction enzyme followed by *in vitro* transcription using MEGAscript™ T7 Transcription Kit (Invitrogen) according to manufacturer's protocol. The quality of the RNA was checked using 0.5% denaturing agarose gel. For transfection, 1 μg RNA was used and transfected in HeLa cells using TransIT®-mRNA Transfection Kit (mirusbio) according to manufacturer's instruction. The cells were harvested for luciferase assay after incubation at 37°C for 12 hrs. The same procedure was repeated for GFP-CVB3 replicon RNA which was used as transfection control (negative control for luciferase signal). The cells were lysed using lysis buffer supplied with Renilla Luciferase Assay Kit 2.0 (Biotium). 20 μl of cell lysate was mixed with 100 μl of freshly prepared working solution and the luminescence was recorded using a TECAN spectrophotometer.

Graphs and statistical procedure

Statistical significance tests were performed using student's T-test as implemented in GraphPad Prism v10.0.1. Error bars represent standard deviation. P value style: ; ns: $p > 0.05$; ; *: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$; ****: $p < 0.0001$

Data availability

The cryo-electron tomogram shown in Fig. 4C has been deposited to the Electron Microscopy Data Bank with accession code EMD-XXXXX.

Acknowledgements

We would like to thank Madeleine Ramstedt (Department of Chemistry, Umeå University) for the introduction to the Zetasizer Malvern Nanoz used for DLS measurements and Mikael Lindberg (Protein expertise platform, Umeå University, part of Protein Production Sweden) for cloning the poliovirus replicons. Cryo-ET was performed at the Umeå Center for Electron Microscopy (UCEM) a SciLifeLab National Cryo-EM facility part of the National Microscopy Infrastructure, NMI (VR-RFI 2016–00968). This project was funded by a Human Frontier Science Program Career Development Award (CDA00047/2017 C) to LAC, a Kempe foundations postdoctoral fellowship to MNS, an MSCA fellowship to HS ("MsInfection", grant agreement ID: 101033469), the Knut and Alice Wallenberg Foundation (through the Wallenberg Center for Molecular Medicine Umeå), and the Swedish research council (grants 2018–05851 and 2021–01145).

Figures and Tables

544 **Figure 1: The membrane-binding region of 2C consists of two amphipathic helix regions, separated by a**

545 **conserved glycine. (A)** Schematic representation of the *Enterovirus* genome, the translated polyprotein with

546 individual proteins marked, and the 2C protein with known domains marked. **(B)** Multiple sequence alignment of

547 the membrane-binding domain (MBD, residues 1-40) of 2C from representative genotypes of different

548 *Enterovirus* species. Conserved polar and hydrophobic residues are highlighted in grey (s: small residue, t: turn

549 like residue, h: hydrophobic residue, p: polar residue, l: aliphatic residue, +: positively charged residue, -:

550 negatively charged residue). The conserved glycine (number 17 in poliovirus 2C) is highlighted in pink. **(C)**

551 Hydrophobic moment (μH), a measure of the local amphiphilicity of a helix in a 18-residue sliding window, is

552 plotted for representative genotypes of different enterovirus species. The higher the μH value, the more an

553 amphipathic the helix at that position. The position of the conserved glycine 17 is marked. **(D)** Alphafold2

554 structure prediction of the MBD of 2C from representative genotypes of different enterovirus species. The

555 conserved glycine 17 is highlighted in pink separating AH1 and AH2.

556

Figure S1: Amphipathic nature of the 2C membrane-binding domain. (A) Helical wheel of the AH1 and

AH2 regions of poliovirus type 1 Mahoney 2C (GenBank: V01148.1) showing the distribution of hydrophobic

and hydrophilic residues. Conserved residues are numbered. **(B)** Isosurface representations of Alphafold2-

predicted structures of the MBD of different enterovirus genotypes showing the distribution of hydrophilic (blue)

and hydrophobic (yellow) residues. For each structure prediction, two views related by the indicated rotation are

shown.

Figure 2: The membrane-binding domain regulates 2C oligomerization. (A) Schematic representation of all the His$_6$-MBP-(TEV) constructs used in the paper: full-length 2C, and N-terminally truncated versions: 2C(ΔAH1) corresponding to deletion of AH1, 2C(41-329) corresponding to deletion of AH1-2 (entire MBD), and 2C(116-329) corresponding to deletion of MBD and part of the ATPase domain (comparable to published crystal structure)[39]. **(B)** Elution profile of the constructs described in panel A on a Superose 6 Increase size-exclusion column. The peaks were centered at the following elution volumes: 12.5 ml for MBP-2C, 13.5 ml for MBP-2C(ΔAH1), 16.5 ml for MBP-2C(41-329), and 17.0 ml for MBP-2C(116-329). The column void volume is indicated with V$_0$. **(C-D)** Mass photometry data showing the molecular mass distribution at 100 nM concentration of C: MBP-2C(ΔAH1) and D: MBP-2C(ΔAH1). Error bars represent a standard deviation of three repeats of the experiment. **(E-F)** SEC-MALS data showing estimated molecular mass of MBP-free 2C(41-329) and 2C(116-329) at concentrations of 38 and 134 μM, respectively. Throughout the main elution peak, 2C(41-329) had an estimated mass of 43 and 52 kDa for two consecutive runs, whereas 2C(116-329) had an estimated mass of 22 and 27 kDa for two consecutive runs.

Figure S2: SDS PAGE gel showing different 2C constructs. lane 1: molecular mass ladder with masses
indicated to its left. lane 2: His$_6$-MBP-(TEV)-full-length 2C, and N-terminally truncated versions: lane 3: His$_6$-
MBP-(TEV)-2C(ΔAH1), lane 4: His$_6$-MBP-(TEV)-2C(41-329), and lane 5: His$_6$-MBP-(TEV)-2C(116-329).

Figure S3: SEC-MALS mass analysis of MBP-fused truncated 2C constructs. SEC-MALS data showing estimated molecular mass throughout the main elution peak of (**A**) MBP-2C(41-329) and (**B**) MBP-2C(116-329) at concentrations of 38 and 134μM, respectively. The calculated masses are shown as a thick black line. MBP-2C(41-329) had an estimated mass of 78 kDa, whereas MBP-2C(116-329) had an estimated mass of 66 kDa.

Figure 3: The AH2 region is critical for 2C's membrane binding. (A) Schematic representation of the
flotation assay used to quantify the membrane binding property of 2C. MBP-2C is fluorophore-labeled with
ATTO488 on exposed cysteines, which only exist in the 2C part of the fusion protein. Small unilamellar vesicles
(SUVs) or large unilamellar vesicles (LUVs) – collectively: liposomes – are labelled with DOPE-coupled
ATTO647N, thus having a distinct fluorescence signal from the protein. The MBP tag is cleaved from MBP-2C
using TEV protease in the presence of liposomes, allowing the liberated MBD to directly associate with the
liposomes. 2C-decorated liposomes are layered at the bottom of a density gradient and subjected to ultra-
centrifugation. The amount of liposomes present in the top fraction, and their bound 2C, is quantified in separate
channels of a fluorimeter. 2C hexamers are dark red, MBP blue. **(B)** Photograph of a representative centrifuge
tube after the flotation assay with 2C, showing two distinct lipid bands in the upper fraction. **(C)** Flotation assays
were conducted as shown in panel A, with SUVs composed of 75 mol% POPC and 25 mol% POPE, having an
average diameter of typically ~50 nm as measured by DLS. 2C fluorescence was measured separately in upper
and lower fractions, and the percentage present in the upper fraction is plotted for the different 2C constructs.
Error bars represent a standard deviation of three repeats of the experiment. Statistical significance by unpaired
two-tailed Student's t test; ns > 0.05; ***p < 0.001. **(D)** Flotation assay as in C, but with varying SUV lipid
composition. PE/PC: 25 mol% POPC, 75 mol% POPE. PE/PC/PS: 25 mol% POPC, 25 mol% POPE, 50 mol%
POPS. Error bars represent a standard deviation of three repeats of the experiment. Statistical significance by
unpaired two-tailed Student's t test; ns >0.05; **p < 0.01. **(E)** Flotation assay as in C, but with varying liposome
size. SUV: sonicated SUVs, ~50 nm diameter. LUV: extruded LUVs, ~2500 nm diameter. Error bars represent a
standard deviation of three repeats of the experiment. Statistical significance by unpaired two-tailed Student's t-
test; ns > 0.05.

Figure S4: Quantitation of lipid fluorescence in flotation experiments. (A-C) Lipid fluorescence signal in the upper fraction of the assays shown in Fig. 3C-E. Error bars represent a standard deviation of three repeats of the experiment. Statistical significance by unpaired two-tailed Student's t test; ns > 0.05. **(D-E)** Fraction 2C recruited to the upper fraction, normalized by amount of lipid in the upper fraction, for the assays shown in Fig. 3C-E. Error bars represent a standard deviation of three repeats of the experiment. Statistical significance by unpaired two-tailed Student's t-test; ns > 0.05; **p < 0.01.

Figure S5: Size distribution of SUVs. Dynamic light scattering graph showing the size distribution of PE/PC

SUV (size distribution by volume). Average size of the SUVs was calculated to be 30 nm.

Figure S6: Two distinct lipid bands in the upper fraction in 2C flotation assays. (A) Photograph of

representative centrifuge tubes after the flotation assay with SUVs only (single lipid band), and SUVs + TEV

protease + MBP-2C (two lipid bands). **(B)** Photograph of centrifuge tube, indicating the upper fractions A, B,

and the lower fraction. The percentage of the total 2C fluorescence in fractions A and B are shown. Error bars

represent the standard deviation of three repeats of the experiment. Statistical significance by unpaired two-tailed

Student's t-test; **p < 0.01; ****p < 0.0001.

Figure 4: 2C has lipid clustering activity that is enhanced by the AH1 region. (A) Representative dynamic

light scattering (DLS) graphs (size distribution by volume), showing the size distribution of the PE/PC SUV

preparation, and appearance of additional peaks corresponding to vesicle clusters, 30 min after adding TEV

protease and 3 μM MBP-2C (red solid line) or MBP-2C(ΔAH1) (orange dotted line) to SUVs (blue solid line).

(B) Percentage of the total vesicle volume present in clusters (defined as particles with diameter > 100 nm), 30

min after adding TEV protease and 0.8 or 3 μM MBP-2C or MBP-2C(ΔAH1). Dots represent the average of

triplicate measurements on the same SUV preparation, bars the average and error bars one standard deviation of

the three independent replicates. Statistical significance by unpaired two-tailed Student's t-test; ns >0.05;

p < 0.01 **(C) Computational slice through a representative cryo-electron tomogram of PE/PC SUVs, 30 min

after adding TEV protease and 3 μM MBP-2C. Densities corresponding to 2C are pointed by red triangles. The

right panel shows the corresponding three-dimensional segmentation showing 2C (red) and SUVs (yellow).

Scale bar, 50 nm.

Figure 5: 2C preferentially recruits double-stranded RNA to membranes. (A) Representative DLS graphs

(size distribution by volume), performed as for Fig. 4A with 3 μM 2C, but with the indicated concentrations of

ssDNA oligonucleotide added. **(B)** Percentage of vesicle volume present as clusters (evaluated as for Fig. 4B),

30 min after addition of TEV protease, MBP-2C, and the indicated concentration of ssDNA oligonucleotide.

Dots represent the average of triplicate measurements on the same SUV preparation, bars the average and error

bars one standard deviation of the three independent replicates. Statistical significance by unpaired two-tailed

Student's t test; ns >0.05; *p < 0.05. **(C)** Schematic representation of the flotation assay used to quantify RNA

recruitment to membranes by 2C. The experiment is essentially performed as in Fig. 3A, except that MBP-2C is

unlabeled, and the RNA (single-stranded, or annealed with a complementary strand for double-stranded) is

labeled with TET fluorophore at its 5′ end (Fig. S8). Membrane-bound RNA is quantitated by measuring TET

fluorescence in the upper fraction **(D)** Fluorescence from ssRNA and dsRNA in top fraction in the experiment

outlined in (C). Dots represent the average of triplicate measurements on the same SUV preparation, bars the

average and error bars one standard deviation of the three independent replicates. Statistical significance by

unpaired two-tailed Student's t-test; ***p < 0.001.

Figure S7: Characterization of fluorescent ssRNA and dsRNA. Fluorescence scan of non-denaturing
polyacrylamide gel showing 5'TET-labelled ssRNA and 5'TET-labelled dsRNA.

Figure 6: Membrane-bound, full-length 2C is an ATP-independent RNA chaperone. (A) Native
polyacrylamide gel resolving dsRNA and ssRNA after a helicase assay. MBP-2C and MBP-2C(41-329) were
incubated with dsRNA with a 5'overhang in the presence of SUVs, and presence or absence of TEV protease
and Mg^{2+}-ATP. Lanes 1 and 2 from the left were loaded with ssRNA control and dsRNA substrate, respectively.
The remaining lanes were loaded with RNA helicase reactions taken at the end of the two hour incubation, with
proteins and reactants as indicated in the figure. A representative gel is shown, the experiment was performed in
three independent replicates. **(B)** The percentage of total RNA signal (dsRNA and ssRNA) present as ssRNA is
plotted for selected lanes of the gel in (A), and two independent replicates of the same experiment. Dots
represent individual experiments, bars the average and error bars one standard deviation. The dsRNA lane was
subtracted as background. Statistical significance by unpaired two-tailed Student's t-test; $*p < 0.05$; $**p < 0.01$;
$****p < 0.0001$. **(C)** Native polyacrylamide gel on a similar experiment to (A), but with a 3' overhang on the
dsRNA. Lanes are labeled analogously to (A). **(D)** The percentage of total RNA signal (dsRNA and ssRNA)
present as ssRNA is plotted for selected lanes of the gel in (C), and two independent replicates of the same
experiment. Dots represent individual experiments, bars the average and error bars one standard deviation of the
three independent replicates. The dsRNA lane was subtracted as background. Statistical significance by unpaired
two-tailed Student's t test; $ns > 0.05$. **(E)** Colorimetric ATPase assay. MBP-2C and MBP-2C(41-329) were
incubated with 0.4 mM ATP in the presence and absence of SUV+TEV protease. The amount of NTP
hydrolyzed was determined and plotted. Dots represent individual experiments, bars the average and error bars
one standard deviation. Statistical significance by unpaired two-tailed Student's t-test; $ns > 0.05$; $****p < 0.0001$.

Figure 7: Truncation of AH1 or the MBD drastically reduces poliovirus replication. (A) Schematic
representation of poliovirus replicons and their use. The replicons contained Renilla luciferase in place of the
capsid region of the polyprotein, and the indicated deletions in the MBD of 2C. The first residue of 2C, glycine,
was kept in both deletions since it is a core part of the protease cleavage site. In vitro-transcribed RNAs from
these replicons were transfected into HeLa cells and luminescence was recorded 12 hours post-transfection. **(B)**
Luminescence recorded for cells transfected with different poliovirus replicons. Dots represent individual
experiments, bars the average, and error bars one standard deviation of the three independent replicates.
Statistical significance by unpaired two-tailed Student's t-test; **$p < 0.01$.

Figure 8: A model for the multiple roles of 2C in enterovirus replication. The model places the biochemical
findings of this study into a suggested cellular context. At an early stage of infection, 2C serves to cluster
replication organelle membranes and lipid droplets, while tethering the first viral dsRNA to the membrane. As
the replication progresses, 2C is increasingly loaded with viral dsRNA and thus is a key component in localizing
viral RNA replication to the replication organelle membranes. Our data suggest that 2C is an active ATPase but
not and RNA helicase – a role which may be played by the polymerase 3D. Although not addressed in this study,
the literature strongly suggests an additional role for 2C in capsid assembly and RNA loading, consistent with it
being part of a "tethering complex" which tethers newly synthesized capsids to replication organelle membranes.

709 **Table 1: List of all the oligonucleotides used**

Oligo	Sequence 5'-3'	Type	Label
1	CCUCUAACCACAGUCUGAUC (20)	RNA	5' Tetrachlorofluorescein (TET)
2	GAUCAGACUGUGG UUAGAGG (20)	RNA	-
3	TGAATGCGGCTAATCCCAACCT (22)	DNA	-
4	AAAAAAAAAAAACCAGGCGACAUCAGCG (28)	RNA	3' Carboxyfluorescein (FAM)
5	CCAGGCGACAUCAGCGAAAAAAAAAAAA(28)	RNA	5' Carboxyfluorescein (FAM)
6	CGCUGAUGUCGCCUGG (16)	RNA	-

Supplementary movie 1: Slices through cryo-electron tomogram showing vesicle clustering by 2C. The

movie loops consecutive xy-planes of the cryo-electron tomogram shown in Fig. 4C.

Supplementary movie 2. Surface model of cryo-electron tomogram showing vesicle clustering by 2C. The

movie shows different views of the surface model of the tomogram shown in Fig. 4C.

References

1.	Hyypia T, Hovi T, Knowles NJ, Stanway G. Classification of enteroviruses based on
molecular and biological properties. J Gen Virol. 1997;78 (Pt 1):1-11. doi: 10.1099/0022-1317-78-1-1.
PubMed PMID: 9010279.
2.	Jiang P, Liu Y, Ma HC, Paul AV, Wimmer E. Picornavirus morphogenesis. Microbiol
Mol Biol Rev. 2014;78(3):418-37. Epub 2014/09/04. doi: 10.1128/MMBR.00012-14. PubMed PMID:
25184560; PubMed Central PMCID: PMCPMC4187686.
3.	Adams MJ, Lefkowitz EJ, King AM, Bamford DH, Breitbart M, Davison AJ, et al.
Ratification vote on taxonomic proposals to the International Committee on Taxonomy of Viruses
(2015). Arch Virol. 2015;160(7):1837-50. doi: 10.1007/s00705-015-2425-z. PubMed PMID: 25913692.
4.	Murphy OC, Messacar K, Benson L, Bove R, Carpenter JL, Crawford T, et al. Acute
flaccid myelitis: cause, diagnosis, and management. Lancet. 2021;397(10271):334-46. Epub 20201223.
doi: 10.1016/S0140-6736(20)32723-9. PubMed PMID: 33357469; PubMed Central PMCID:
PMCPMC7909727.
5.	Holm-Hansen CC, Midgley SE, Fischer TK. Global emergence of enterovirus D68: a
systematic review. Lancet Infect Dis. 2016;16(5):e64-e75. Epub 20160224. doi: 10.1016/S1473-
3099(15)00543-5. PubMed PMID: 26929196.
6.	Lukashev AN, Lashkevich VA, Ivanova OE, Koroleva GA, Hinkkanen AE, Ilonen J.
Recombination in circulating enteroviruses. J Virol. 2003;77(19):10423-31. doi:
10.1128/jvi.77.19.10423-10431.2003. PubMed PMID: 12970427; PubMed Central PMCID:
PMCPMC228507.
7.	Chevaliez S, Szendroi A, Caro V, Balanant J, Guillot S, Berencsi G, et al. Molecular
comparison of echovirus 11 strains circulating in Europe during an epidemic of multisystem
hemorrhagic disease of infants indicates that evolution generally occurs by recombination. Virology.
2004;325(1):56-70. doi: 10.1016/j.virol.2004.04.026. PubMed PMID: 15231386.
8.	Lukashev AN. Role of recombination in evolution of enteroviruses. Rev Med Virol.
2005;15(3):157-67. doi: 10.1002/rmv.457. PubMed PMID: 15578739.
9.	Martini M, Orsini D. The fight against poliomyelitis through the history: past, present
and hopes for the future. Albert Sabin's missing Nobel and his "gift to all the world's children". Vaccine.
2022;40(47):6802-5. Epub 20221014. doi: 10.1016/j.vaccine.2022.09.088. PubMed PMID: 36244880.
10.	Thompson KM, Lauring AS, Pollard AJ, Andino R, Bandyopadhyay AS, Berkley S, et
al. Polio eradication: Addressing the hurdles on the last mile. Cell. 2023;186(1):1-4. doi:
10.1016/j.cell.2022.12.021. PubMed PMID: 36608647.
11.	Pons-Salort M, Parker EP, Grassly NC. The epidemiology of non-polio enteroviruses:
recent advances and outstanding questions. Curr Opin Infect Dis. 2015;28(5):479-87. doi:
10.1097/QCO.0000000000000187. PubMed PMID: 26203854; PubMed Central PMCID:
PMCPMC6624138.
12.	Chapman NM, Kim KS. Persistent coxsackievirus infection: enterovirus persistence in
chronic myocarditis and dilated cardiomyopathy. Curr Top Microbiol Immunol. 2008;323:275-92. doi:
10.1007/978-3-540-75546-3_13. PubMed PMID: 18357775.
13.	Anasir MI, Zarif F, Poh CL. Antivirals blocking entry of enteroviruses and therapeutic
potential. J Biomed Sci. 2021;28(1):10. Epub 20210115. doi: 10.1186/s12929-021-00708-8. PubMed
PMID: 33451326; PubMed Central PMCID: PMCPMC7811253.
14.	Sun J, Yogarajah T, Lee RCH, Kaur P, Inoue M, Tan YW, et al. Drug repurposing of
pyrimidine analogs as potent antiviral compounds against human enterovirus A71 infection with
potential clinical applications. Sci Rep. 2020;10(1):8159. Epub 20200518. doi: 10.1038/s41598-020-
65152-4. PubMed PMID: 32424333; PubMed Central PMCID: PMCPMC7235037.
15.	Kejriwal R, Evans T, Calabrese J, Swistak L, Alexandrescu L, Cohen M, et al.
Development of Enterovirus Antiviral Agents That Target the Viral 2C Protein. ChemMedChem.
2023;18(10):e202200541. Epub 20230310. doi: 10.1002/cmdc.202200541. PubMed PMID: 36792530.
16.	Caliguiri LA, Tamm I. Membranous structures associated with translation and
transcription of poliovirus RNA. Science. 1969;166(3907):885-6. doi: 10.1126/science.166.3907.885.
PubMed PMID: 4310189.

17.   Bienz K, Egger D, Rasser Y, Bossart W. Kinetics and location of poliovirus
macromolecular synthesis in correlation to virus-induced cytopathology. Virology. 1980;100(2):390-9.
doi: 10.1016/0042-6822(80)90530-9. PubMed PMID: 6243433.
18.   Bienz K, Egger D, Rasser Y, Bossart W. Intracellular distribution of poliovirus proteins
and the induction of virus-specific cytoplasmic structures. Virology. 1983;131(1):39-48. doi:
10.1016/0042-6822(83)90531-7. PubMed PMID: 6316654.
19.   Dahmane S, Kerviel A, Morado DR, Shankar K, Ahlman B, Lazarou M, et al. Membrane-
assisted assembly and selective secretory autophagy of enteroviruses. Nature communications.
2022;13(1):5986. Epub 2022/10/11. doi: 10.1038/s41467-022-33483-7. PubMed PMID: 36216808;
PubMed Central PMCID: PMCPMC9550805.
20.   Limpens RW, van der Schaar HM, Kumar D, Koster AJ, Snijder EJ, van Kuppeveld FJ,
et al. The transformation of enterovirus replication structures: a three-dimensional study of single- and
double-membrane compartments. mBio. 2011;2(5). Epub 2011/10/06. doi: 10.1128/mBio.00166-11.
PubMed PMID: 21972238; PubMed Central PMCID: PMCPMC3187575.
21.   Strating JR, van der Linden L, Albulescu L, Bigay J, Arita M, Delang L, et al.
Itraconazole inhibits enterovirus replication by targeting the oxysterol-binding protein. Cell Rep.
2015;10(4):600-15. Epub 20150129. doi: 10.1016/j.celrep.2014.12.054. PubMed PMID: 25640182;
PubMed Central PMCID: PMCPMC4383725.
22.   Ilnytska O, Santiana M, Hsu NY, Du WL, Chen YH, Viktorova EG, et al. Enteroviruses
harness the cellular endocytic machinery to remodel the host cell cholesterol landscape for effective
viral replication. Cell Host Microbe. 2013;14(3):281-93. doi: 10.1016/j.chom.2013.08.002. PubMed
PMID: 24034614; PubMed Central PMCID: PMCPMC3802520.
23.   Laufman O, Perrino J, Andino R. Viral Generated Inter-Organelle Contacts Redirect
Lipid Flux for Genome Replication. Cell. 2019;178(2):275-89 e16. Epub 2019/06/18. doi:
10.1016/j.cell.2019.05.030. PubMed PMID: 31204099; PubMed Central PMCID: PMCPMC7077330.
24.   Schlegel A, Giddings TH, Jr., Ladinsky MS, Kirkegaard K. Cellular origin and
ultrastructure of membranes induced during poliovirus infection. J Virol. 1996;70(10):6576-88. doi:
10.1128/JVI.70.10.6576-6588.1996. PubMed PMID: 8794292; PubMed Central PMCID:
PMCPMC190698.
25.   Hsu NY, Ilnytska O, Belov G, Santiana M, Chen YH, Takvorian PM, et al. Viral
reorganization of the secretory pathway generates distinct organelles for RNA replication. Cell.
2010;141(5):799-811. Epub 2010/06/01. doi: S0092-8674(10)00369-7 [pii]

10.1016/j.cell.2010.03.050. PubMed PMID: 20510927; PubMed Central PMCID: PMC2982146.
26.   Miller S, Krijnse-Locker J. Modification of intracellular membrane structures for virus
replication. Nature reviews. 2008;6(5):363-74. Epub 2008/04/17. doi: nrmicro1890 [pii]

10.1038/nrmicro1890. PubMed PMID: 18414501.
27.   Melia CE, van der Schaar HM, de Jong AWM, Lyoo HR, Snijder EJ, Koster AJ, et al.
The Origin, Dynamic Morphology, and PI4P-Independent Formation of Encephalomyocarditis Virus
Replication Organelles. mBio. 2018;9(2). Epub 20180417. doi: 10.1128/mBio.00420-18. PubMed
PMID: 29666283; PubMed Central PMCID: PMCPMC5904412.
28.   Bienz K, Egger D, Pfister T, Troxler M. Structural and functional characterization of the
poliovirus replication complex. J Virol. 1992;66(5):2740-7. doi: 10.1128/JVI.66.5.2740-2747.1992.
PubMed PMID: 1313898; PubMed Central PMCID: PMCPMC241029.
29.   Takeda N, Kuhn RJ, Yang CF, Takegami T, Wimmer E. Initiation of poliovirus plus-
strand RNA synthesis in a membrane complex of infected HeLa cells. J Virol. 1986;60(1):43-53. doi:
10.1128/JVI.60.1.43-53.1986. PubMed PMID: 3018300; PubMed Central PMCID: PMCPMC253900.
30.   Guo H, Li Y, Liu G, Jiang Y, Shen S, Bi R, et al. A second open reading frame in human
enterovirus determines viral replication in intestinal epithelial cells. Nat Commun. 2019;10(1):4066.
Epub 20190906. doi: 10.1038/s41467-019-12040-9. PubMed PMID: 31492846; PubMed Central
PMCID: PMCPMC6731315.
31.   Gorbalenya AE, Koonin EV, Wolf YI. A new superfamily of putative NTP-binding
domains encoded by genomes of small DNA and RNA viruses. FEBS Lett. 1990;262(1):145-8. doi:
10.1016/0014-5793(90)80175-i. PubMed PMID: 2156730.

32.   Pfister T, Wimmer E. Characterization of the nucleoside triphosphatase activity of
poliovirus protein 2C reveals a mechanism by which guanidine inhibits poliovirus replication. J Biol
Chem. 1999;274(11):6992-7001. doi: 10.1074/jbc.274.11.6992. PubMed PMID: 10066753.
33.   Teterina NL, Kean KM, Gorbalenya AE, Agol VI, Girard M. Analysis of the functional
significance of amino acid residues in the putative NTP-binding pattern of the poliovirus 2C protein. J
Gen Virol. 1992;73 (Pt 8):1977-86. doi: 10.1099/0022-1317-73-8-1977. PubMed PMID: 1322957.
34.   Liu Y, Wang C, Mueller S, Paul AV, Wimmer E, Jiang P. Direct interaction between two
viral proteins, the nonstructural protein 2C and the capsid protein VP3, is required for enterovirus
morphogenesis.   PLoS   Pathog.   2010;6(8):e1001066.   Epub   20100826.   doi:
10.1371/journal.ppat.1001066.   PubMed   PMID: 20865167;   PubMed   Central   PMCID:
PMCPMC2928791.
35.   Vance LM, Moscufo N, Chow M, Heinz BA. Poliovirus 2C region functions during
encapsidation of viral RNA. J Virol. 1997;71(11):8759-65. Epub 1997/10/29. PubMed PMID: 9343235;
PubMed Central PMCID: PMCPMC192341.
36.   Li JP, Baltimore D. Isolation of poliovirus 2C mutants defective in viral RNA synthesis.
J Virol. 1988;62(11):4016-21. doi: 10.1128/JVI.62.11.4016-4021.1988. PubMed PMID: 2845120;
PubMed Central PMCID: PMCPMC253830.
37.   Pincus SE, Diamond DC, Emini EA, Wimmer E. Guanidine-selected mutants of
poliovirus: mapping of point mutations to polypeptide 2C. J Virol. 1986;57(2):638-46. doi:
10.1128/JVI.57.2.638-646.1986.   PubMed   PMID: 3003395;   PubMed   Central   PMCID:
PMCPMC252779.
38.   Tolskaya EA, Romanova LI, Kolesnikova MS, Gmyl AP, Gorbalenya AE, Agol VI.
Genetic studies on the poliovirus 2C protein, an NTPase. A plausible mechanism of guanidine effect on
the 2C function and evidence for the importance of 2C oligomerization. J Mol Biol. 1994;236(5):1310-
23. doi: 10.1016/0022-2836(94)90060-4. PubMed PMID: 8126722.
39.   Guan H, Tian J, Zhang C, Qin B, Cui S. Crystal structure of a soluble fragment of
poliovirus   2CATPase.   PLoS   Pathog.   2018;14(9):e1007304.   Epub   20180919.   doi:
10.1371/journal.ppat.1007304.   PubMed   PMID: 30231078;   PubMed   Central   PMCID:
PMCPMC6166989.
40.   Guan H, Tian J, Qin B, Wojdyla JA, Wang B, Zhao Z, et al. Crystal structure of 2C
helicase   from   enterovirus   71.   Sci   Adv.   2017;3(4):e1602573.   Epub   2017/05/17.   doi:
10.1126/sciadv.1602573. PubMed PMID: 28508043; PubMed Central PMCID: PMCPMC5409451.
41.   Hurdiss DL, El Kazzi P, Bauer L, Papageorgiou N, Ferron FP, Donselaar T, et al.
Fluoxetine targets an allosteric site in the enterovirus 2C AAA+ ATPase and stabilizes a ring-shaped
hexameric   complex.   Sci   Adv.   2022;8(1):eabj7615.   Epub   20220105.   doi: 10.1126/sciadv.abj7615.
PubMed PMID: 34985963; PubMed Central PMCID: PMCPMC8730599.
42.   Pfister T, Jones KW, Wimmer E. A cysteine-rich motif in poliovirus protein 2C(ATPase)
is   involved   in   RNA   replication   and   binds   zinc   in   vitro.   J   Virol.   2000;74(1):334-43.   doi:
10.1128/jvi.74.1.334-343.2000.   PubMed   PMID: 10590122;   PubMed   Central   PMCID:
PMCPMC111544.
43.   Adams P, Kandiah E, Effantin G, Steven AC, Ehrenfeld E. Poliovirus 2C protein forms
homo-oligomeric structures required for ATPase activity. J Biol Chem. 2009;284(33):22012-21. Epub
20090611. doi: 10.1074/jbc.M109.031807. PubMed PMID: 19520852; PubMed Central PMCID:
PMCPMC2755925.
44.   Yeager C, Carter G, Gohara DW, Yennawar NH, Enemark EJ, Arnold JJ, et al.
Enteroviral 2C protein is an RNA-stimulated ATPase and uses a two-step mechanism for binding to
RNA   and   ATP.   Nucleic   Acids   Res.   2022;50(20):11775-98.   Epub   2022/11/19.   doi:
10.1093/nar/gkac1054. PubMed PMID: 36399514; PubMed Central PMCID: PMCPMC9723501.
45.   Xia H, Wang P, Wang GC, Yang J, Sun X, Wu W, et al. Human Enterovirus
Nonstructural Protein 2CATPase Functions as Both an RNA Helicase and ATP-Independent RNA
Chaperone. PLoS Pathog. 2015;11(7):e1005067. Epub 20150728. doi: 10.1371/journal.ppat.1005067.
PubMed PMID: 26218680; PubMed Central PMCID: PMCPMC4517893.
46.   Paul AV, Molla A, Wimmer E. Studies of a putative amphipathic helix in the N-terminus
of poliovirus protein 2C. Virology. 1994;199(1):188-99. doi: 10.1006/viro.1994.1111. PubMed PMID:
8116242.

47.   Echeverri AC, Dasgupta A. Amino terminal regions of poliovirus 2C protein mediate
membrane binding. Virology. 1995;208(2):540-53. doi: 10.1006/viro.1995.1185. PubMed PMID:
7747426.
48.   Echeverri A, Banerjee R, Dasgupta A. Amino-terminal region of poliovirus 2C protein is
sufficient for membrane binding. Virus Res. 1998;54(2):217-23. doi: 10.1016/s0168-1702(98)00016-1.
PubMed PMID: 9696129.
49.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly
accurate protein structure prediction with AlphaFold. Nature. 2021;596(7873):583-9. Epub 20210715.
doi: 10.1038/s41586-021-03819-2. PubMed PMID: 34265844; PubMed Central PMCID:
PMCPMC8371605.
50.   Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, et al. UCSF
ChimeraX: Meeting modern challenges in visualization and analysis. Protein Sci. 2018;27(1):14-25.
Epub 2017/07/16. doi: 10.1002/pro.3235. PubMed PMID: 28710774; PubMed Central PMCID:
PMCPMC5734306.
51.   Papageorgiou N, Coutard B, Lantez V, Gautron E, Chauvet O, Baronti C, et al. The 2C
putative helicase of echovirus 30 adopts a hexameric ring-shaped structure. Acta Crystallogr D Biol
Crystallogr. 2010;66(Pt 10):1116-20. Epub 20100918. doi: 10.1107/S090744491002809X. PubMed
PMID: 20944244.
52.   Wolff G, Melia CE, Snijder EJ, Barcena M. Double-Membrane Vesicles as Platforms for
Viral Replication. Trends Microbiol. 2020;28(12):1022-33. Epub 2020/06/17. doi:
10.1016/j.tim.2020.05.009. PubMed PMID: 32536523; PubMed Central PMCID: PMCPMC7289118.
53.   Wang C, Jiang P, Sand C, Paul AV, Wimmer E. Alanine scanning of poliovirus
2CATPase reveals new genetic evidence that capsid protein/2CATPase interactions are essential for
morphogenesis. J Virol. 2012;86(18):9964-75. Epub 20120703. doi: 10.1128/JVI.00914-12. PubMed
PMID: 22761387; PubMed Central PMCID: PMCPMC3446611.
54.   Rodriguez PL, Carrasco L. Poliovirus protein 2C contains two regions involved in RNA
binding activity. J Biol Chem. 1995;270(17):10105-12. doi: 10.1074/jbc.270.17.10105. PubMed PMID:
7730315.
55.   Rodriguez PL, Carrasco L. Poliovirus protein 2C has ATPase and GTPase activities. J
Biol Chem. 1993;268(11):8105-10. PubMed PMID: 8385138.
56.   Mirzayan C, Wimmer E. Genetic analysis of an NTP-binding motif in poliovirus
polypeptide 2C. Virology. 1992;189(2):547-55. doi: 10.1016/0042-6822(92)90578-d. PubMed PMID:
1322588.
57.   Cohen L, Kean KM, Girard M, Van der Werf S. Effects of P2 cleavage site mutations on
poliovirus polyprotein processing. Virology. 1996;224(1):34-42. Epub 1996/10/01. doi:
10.1006/viro.1996.0504. PubMed PMID: 8862397.
58.   Dahmane S, Shankar K, Carlson LA. A 3D view of how enteroviruses hijack autophagy.
Autophagy. 2022:1-3. Epub 2022/12/07. doi: 10.1080/15548627.2022.2153572. PubMed PMID:
36471479.
59.   Dubankova A, Humpolickova J, Klima M, Boura E. Negative charge and membrane-
tethered viral 3B cooperate to recruit viral RNA dependent RNA polymerase 3D (pol). Scientific reports.
2017;7(1):17309. Epub 2017/12/13. doi: 10.1038/s41598-017-17621-6. PubMed PMID: 29230036;
PubMed Central PMCID: PMCPMC5725453.
60.   Cho MW, Richards OC, Dmitrieva TM, Agol V, Ehrenfeld E. RNA duplex unwinding
activity of poliovirus RNA-dependent RNA polymerase 3Dpol. J Virol. 1993;67(6):3010-8. Epub
1993/06/01. doi: 10.1128/JVI.67.6.3010-3018.1993. PubMed PMID: 8388485; PubMed Central
PMCID: PMCPMC237637.
61.   Baggen J, Thibaut HJ, Strating J, van Kuppeveld FJM. The life cycle of non-polio
enteroviruses and how to target it. Nature reviews. 2018;16(6):368-81. Epub 2018/04/08. doi:
10.1038/s41579-018-0005-4. PubMed PMID: 29626210.
62.   Young G, Hundt N, Cole D, Fineberg A, Andrecka J, Tyler A, et al. Quantitative mass
imaging of single biological macromolecules. Science. 2018;360(6387):423-7. doi:
10.1126/science.aar5839. PubMed PMID: 29700264; PubMed Central PMCID: PMCPMC6103225.
63.   Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation
of high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol. 2011;7:539.

Epub 20111011. doi: 10.1038/msb.2011.75. PubMed PMID: 21988835; PubMed Central PMCID: PMCPMC3261699.

64.   Gautier R, Douguet D, Antonny B, Drin G. HELIQUEST: a web server to screen sequences with specific alpha-helical properties. Bioinformatics. 2008;24(18):2101-2. Epub 20080728. doi: 10.1093/bioinformatics/btn392. PubMed PMID: 18662927.

65.   Mastronarde DN, Held SR. Automated tilt series alignment and tomographic reconstruction in IMOD. J Struct Biol. 2017;197(2):102-13. Epub 20160719. doi: 10.1016/j.jsb.2016.07.011. PubMed PMID: 27444392; PubMed Central PMCID: PMCPMC5247408.

66.   Zheng SQ, Palovcak E, Armache JP, Verba KA, Cheng Y, Agard DA. MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron microscopy. Nat Methods. 2017;14(4):331-2. Epub 2017/03/03. doi: 10.1038/nmeth.4193. PubMed PMID: 28250466; PubMed Central PMCID: PMCPMC5494038.

67.   Rohou A, Grigorieff N. CTFFIND4: Fast and accurate defocus estimation from electron micrographs. J Struct Biol. 2015;192(2):216-21. Epub 2015/08/19. doi: 10.1016/j.jsb.2015.08.008. PubMed PMID: 26278980; PubMed Central PMCID: PMCPMC6760662.

68.   Liu YT, Zhang H, Wang H, Tao CL, Bi GQ, Zhou ZH. Isotropic reconstruction for electron tomography with deep learning. Nat Commun. 2022;13(1):6482. Epub 20221029. doi: 10.1038/s41467-022-33957-8. PubMed PMID: 36309499; PubMed Central PMCID: PMCPMC9617606.