Human effector CD8[+] T cells with an exhausted-like phenotype control tumor growth *in vivo* in a humanized tumor model

Juliane Mietz[1], Meike Kaulfuss[1], Lukas Egli[1], Lennart Opitz[2],Christian Münz[3], Obinna Chijioke[1,4]

[1]Cellular Immunotherapy, Institute of Experimental Immunology, University of Zürich, Zürich, Switzerland
[2]Functional Genomics Center Zürich, University of Zürich/ETH Zürich, Zürich, Switzerland
[3]Viral Immunobiology, Institute of Experimental Immunology, University of Zürich, Zürich, Switzerland
[4]Institute of Medical Genetics and Pathology, University Hospital Basel, Basel, Switzerland

Corresponding author: Obinna Chijioke, Cellular Immunotherapy, Institute of Experimental Immunology, University of Zürich, Winterthurerstrasse 190, 8057 Zürich, Switzerland; Tel: +41 44 635 37 19, Email: chijioke@immunology.uzh.ch

Abstract

Background Humanized tumor models could be particularly valuable for cancer immunotherapy research, as they may better reflect human-specific aspects of the interfaces between tumor and immune system of human cancer. However, endogenous antitumor immunity in humanized models is still largely undefined.

Methods We established a novel autologous humanized mouse tumor model by using NSG mice reconstituted with human immune cells from hematopoietic progenitors and tumors generated from transformed autologous human B cells. We demonstrate growth of solid lymphoid tumors after subcutaneous implantation, infiltration by endogenous human immune cells and immunocompetence of the model.

Findings We found human T cell subsets described in human cancer, including progenitor exhausted (T_{pex}), terminally exhausted ($T_{ex-term}$) and tissue-resident (T_{RM}) cells in tumor-bearing humanized mice with accumulation of $T_{ex-term}$ and T_{RM} in the tumor. In addition, we identified tumor-reactive CD8$^+$ T cells through expression of CD137. This subpopulation of de novo arising human CD137$^+$ CD8$^+$ T cells displayed a highly proliferative, fully activated effector and exhausted-like phenotype with enhanced expression of activation and exhaustion markers like PD-1, CD39, CD160, TIM-3, TIGIT and TOX, the senescence marker CD57 (*B3GAT1*) and cytolytic effector molecules such as *PRF1*, *GZMH* and *NKG7*. Moreover, these CD137$^+$ CD8$^+$ T cells exhibited tumor-specific clonal expansion and presented signature overlap with tumor-reactive CD8$^+$ T cells described in human cancer. We demonstrate superior anticancer activity of this exhausted-like human CD8$^+$ T cell subset by adoptive transfer experiments using recipients bearing autologous human tumors. Mice adoptively transferred with CD137$^+$ CD8$^+$ T cells showed reduced tumor growth and higher CD8$^+$ T cell tumor infiltration, correlating with control of human tumors.

Interpretation We established an immunocompetent humanized tumor model, providing a tool for immunotherapy research and defined effective anticancer activity of human effector CD8$^+$ T cells with an exhausted-like phenotype, supporting clinical exploration of such cells in adoptive T cell therapies.

30   **Funding** Swiss Cancer Research foundation.

32   **Keywords:** Human cancer immunology; Humanized cancer models; Phenotypic markers; T

33   cells; Exhaustion; Adoptive cell therapy

Research in context

Evidence before this study

Antitumor immune responses and outcome of immunotherapeutic interventions are not always consistent between mouse models of cancer and data available in humans. This may be due to species-specific differences, therefore models with a potential for better translatability are needed, such as humanized mouse models. However, there is limited data on human antitumor T cell immunity in humanized mice.

Added value of this study

In this study, we established an immunocompetent humanized tumor model that recapitulates hallmarks of human antitumor T cell responses, offering the possibility for further translational investigation of the interface between human tumors and endogenous anticancer immunity. Furthermore, using functional *in vitro* assays and adoptive transfer, our study demonstrates the key importance of human effector CD8[+] T cells with an activated and exhausted-like phenotype in the antitumor immune response.

Implications of all the available evidence

The autologous humanized tumor model provided in this study can serve as a tool to elucidate human-specific immune features. By bridging a gap between syngeneic mouse tumor models and human-specific antitumor immune responses, the model may help open up avenues for greater translatability of preclinical data. Our findings suggest that exhausted-like effector CD8[+] T cells can be harnessed for clinical development of adoptive T cell therapies.

Introduction

The characterization of phenotypic traits that can faithfully identify tumor-specific CD8[+] T cells with functional anticancer activity is an area of intense research. Recent studies have shown tumor-reactive CD8[+] T cells to be rare in patients with various cancers and thus challenging to explore (1,2). Preclinical studies in mice have shown that CD8[+] T cells displaying an exhausted-like phenotype have decreased effector function (3). Reinvigoration of exhausted antitumor T cells is thought to be a key component of response to immune checkpoint blockade. Nevertheless, studies in individuals with cancer have correlated exhausted-like phenotypic states of CD8[+] T cells with tumor reactivity and clinical benefit (4,5).

Undoubtedly, tumor models using inbred and transgenic mice have contributed substantially to our understanding of tumor immunobiology and have been fundamental to the development of cancer immunotherapies that are now used on a routine basis in the clinic (6,7). However, human clinical studies (8) have shown that the outcome of cancer immunotherapies can significantly diverge from those of mouse models of cancer (9). This might be due to a relative lack of genetic diversity of mouse models compared to humans (10) but also to differences in aspects of gene expression networks (11), genomic organization (12) and tissue structure of the immune system of mice and humans (13). Xenograft mouse models using adoptive transfer of human immune cells rely on immunodeficient hosts, precluding analysis of the impact of endogenous adaptive antitumor immunity. Furthermore, in most preclinical studies investigating novel immunotherapy targets, monoclonal T cells with a single antigen specificity are explored (14,15) while polyclonal T cell responses might be a biomarker for favorable patient outcome to immunotherapy (16) and thus be more informative.

Complementing existing mouse tumor models, the use of mice with reconstituted human immune system components (HIS mice) holds hitherto underexplored translational potential to advance human cancer immunology (17). Yet, endogenous antitumor immunity in these models has remained largely undefined, mainly due to allogeneic reactivity of the reconstituted human T cells against allogeneic tumors (18), making a distinction of alloreactivity from tumor-specific immune responses challenging.

In this study, we demonstrate that reconstitution of human immune system components

across independent donors can mount endogenous antitumor T cell responses to autologous

human tumor challenge in HIS mice. A polyclonal population of human effector CD8$^+$ T cells

with an activated and exhausted-like phenotype that arises de novo in tumor-bearing HIS mice,

resembling exhausted tumor-reactive lymphocytes found in patients with cancer, is able to

control autologous human tumor growth *in vivo*. Thus, we provide a tool to investigate human

antitumor T cell immunity and direct evidence of clinically relevant anticancer activity of

expanded human tumor-reactive effector CD8$^+$ T cells with an activated and exhausted-like,

non-progenitor-like phenotype.

Methods

HIS & NSG mice

NSG (NOD.Cg-Prkdc<scid>Il2rg<tm1Wjl>/SzJ (#005557)) mice were purchased from The Jackson Laboratory and bred and housed under specific pathogen-free conditions at the Laboratory Animal Services Center (LASC) Zurich, University of Zurich. To generate HIS mice, newborn pups (1-5 days old) were irradiated (1 Gy) and intrahepatically injected with 0.2×10^6 human $CD34^+$ hematopoietic progenitor cells derived from human fetal liver (Advanced Bioscience Resources, USA). $CD34^+$ cells were isolated by MACS technology (Miltenyi, Cat. 130-046-703) following the manufacturer's instructions. At 3 months of age, reconstitution of human immune cells was assessed in PBMC from tail vein blood by flow cytometry [19]. Reconstitution analysis included staining for human CD45, CD3, CD4, CD8, CD19, NKp46 and HLA-DR. Only mice with a sufficiently high frequency of human CD45 (>25% of leukocytes) were included in the experiments. Both female and male mice were used in the experiments and no sex-based analysis was performed. All animal experiments were performed according to approved licenses by the veterinary office of the canton of Zurich, Switzerland (ZH049/20 and ZH067/2023).

LCL generation and tumor model

LCL were generated from $CD19^+$ B cells derived from the same autologous human fetal liver tissue from which the $CD34^+$ cells for reconstitution of the human immune system in the same experiment were derived. $CD19^+$ cells were isolated from the HFL tissue's $CD34^-$ fraction by MACS technology (Miltenyi, Cat. 130-050-301) according to the manufacturer's instructions. 0.25×10^6 $CD19^+$ human B cells were transformed by infection with Epstein-Barr virus (EBV B95-8; produced as previously described [19]) at an MOI of 0.1 - 0.15 and cultured in RPMI/ 10% FCS/ 1% Penicillin/ Streptomycin.

LCL tumors were injected subcutaneously into the left flank under isoflurane narcosis. For generation and phenotyping of tumor-reactive T cells, 5×10^6 autologous LCL tumor cells were resuspended in PBS and right before injection mixed in a 1:1 V/V ratio with Corning® Matrigel® Growth Factor Reduced (GFR) Basement Membrane Matrix (Milian, Cat. 354230).

For adoptive cell transfer experiments, 2 x 10^6 LCL were injected s.c. in a 1:1 V/V mix with

Matrigel and three days after tumor injection, 2 x 10^6 T cells (HIS recipient mice) or 10 x 10^6

(NSG recipient mice) were adoptively transferred by tail vein injection. Tumor size was

monitored by calipering (3x/week or daily). Tumor volume was calculated as volume [mm^3] =

length [mm] x width^2 [mm^2] x 0.52. General health was monitored by weighing and health

parameter scoring (3x/week or daily, according to animal license). PBMC composition and

expansion of adoptively transferred T cells were monitored by weekly tail vein bleeding and

flow cytometric analysis. White blood cell (WBC) counts were measured from blood with an

automatic cell counting machine (DxH 500, Beckman Coulter). Animals were euthanized when

they met pre-defined termination criteria defined in the animal license.

T cell isolation and expansion

Spleens from LCL tumor-bearing and naïve HIS mice were harvested 16-18 days after s.c.

tumor injection. Splenocytes were labelled with fluorescently labelled antibodies and sorted by

fluorescent activated cell sorting (FACS). CD137$^+$, CD137$^-$ and CD137$^-$PD-1$^-$ CD8$^+$ T cells were

sorted from tumor-bearing HIS mice and bulk CD8$^+$ T cells were isolated from naïve HIS mice.

Sorted T cells were expanded by a rapid expansion protocol (REP), as described elsewhere

(20). REP medium consisted of a 1:1 mix of RPMI (Gibco, Cat. 7001612) and X-Vivo (Lonza,

Cat. BE02-060F) containing 10% FCS (Biochrome, Cat. S0615-500ML) and 1% Penicillin/

Streptomycin (Thermo Fisher, Cat. 7001592). After sorting, T cells were cultured in REP

medium with 30 ng/µl OKT3 (Miltenyi, Cat. 130-093-387), 3000 IU/ml IL-2 (Peprotech, Cat.

200-02) and a 200-fold excess of irradiated (50 Gy) allogeneic PBMC derived from 3 healthy

donors at a total cell density of 5 x 10^6 cells/ml. Starting at day 7, cells were adjusted with REP

medium to 0.5-1 x 10^6 cells/ml and fresh IL-2 (3000 IU/ml) was added every second day. Cells

were expanded for a total of 14-16 days until use for further experiments.

Cell isolation from organs

For reconstitution analysis and weekly bleeding, 100-150 µl of tail vein blood was collected

into EDTA tubes (BD Microtainer, BD, Cat. 365975) and erythrocytes were lysed by incubation

with lysis buffer. Leukocytes were stained for flow cytometric analysis. For PBMC isolation on

the day of sacrifice, blood was harvested by heart puncture and erythrocytes were lysed with

erythrocyte lysis buffer. For splenocyte isolation, spleens were removed and meshed through

a 70 µm cell strainer in PBS and lymphocytes were isolated by Ficoll-Paque (GE Healthcare,

17-5442-03). Single cell suspensions were washed and stained for flow cytometric analysis.

For tumor-infiltrating lymphocyte (TIL) isolation, tumors were removed from the flank and skin

was removed from the tumors. Tumors were cut into small pieces with scissors and then

incubated with digestion mix (DMEM (Life technologies, Cat. 7001566) containing 2% FCS

(Biochrome, Cat. S0615-500ML), 1 mg/ml Collagenase IV (Roche, Cat. 7002219), 10ug/ml

DNase I (Roche, Cat. 7002221) and 1.2 mM CaCl2; approx. 1.5 ml/tumor, incubation for 45

min, 37°C). The tumor digest was diluted with 20 ml RPMI or PBS and meshed over a 70 µm

cell strainer. Single cell suspensions were washed and then stained for flow cytometric

analysis.

Flow cytometry

For surface staining, cells were washed in PBS and stained with an antibody master mix for

20 min at 4°C in the dark. For intracellular staining, cells were first stained for surface antigens

and fixable live/dead marker, then fixed and permeabilized using the BD Pharmingen™

Transcription Factor Buffer Set (BD, Cat. 562574) according to the manufacturer's instructions,

then stained for intracellular or intranuclear antigens for 50-60 minutes. Cells were washed in

PBS and acquired on a BD LSRFortessa™ or Cytek Aurora.

Antibodies for reconstitution analysis of HIS mice (all antibodies are anti-human): CD45-Pacific

Blue (Biolegend, HI30, Cat. 304029), CD3-PE (Biolegend, UCHT1, Cat. 300408), CD4-APC-

Cy7 (Biolegend, RPA-T4, Cat. 300518), CD8-PerCP (Biolegend, SK1, Cat. 344708), HLA-DR-

FITC (Biolegend, L243, Cat. 307604), CD19-PE-Cy7 (Biolegend, HIB19, Cat. 302216),

NKp46-APC (BD, 9-E2, Cat. 558051), Zombie Aqua™ Fixable Viability Kit (Biolegend, Cat.

423101).

Antibodies for flow cytometry-based sorting of T cells: CD3-FITC (Biolegend, UCHT1, Cat.

300406), CD4-APC-Cy7 (Biolegend, RPA-T4, Cat. 300518), CD8-BV650 (Biolegend, SK1,

Cat. 344730), CD45-Pacific Blue (Biolegend, HI30, Cat. 304029), CD137-APC (Miltenyi,

REA765, Cat. 130-110-901), PD-1-PE-Dazzle (Biolegend, EH12.2H7, Cat. 329940), Zombie

Aqua™ Fixable Viability Kit (Biolegend, Cat. 423101).

Antibodies for phenotyping in Panel 1: CD45-Pacific Blue (Biolegend, HI30, Cat. 304029),

CD3-PerCP-Cy5.5 (Biolegend, UCHT1, Cat. 300429), CD4-APC-Cy7 (Biolegend, RPA-T4,

Cat. 300518), CD8-BV650 (Biolegend, SK1, Cat. 344730), CD45RA-BV785 (Biolegend,

HI100, 304139), CD62L-FITC (BD, DREG-56, Cat. 555543), CD19-PE-Cy7 (Biolegend,

HIB19, Cat. 302216), HLA-DR-BV605 (BD, g46-6, Cat. 562845), CD39-BV711 (Biolegend, A1,

Cat. 328228), CD69-Alexa Fluor 700 (Biolegend, FN50, Cat. 310921), PD-1-PE-Dazzle

(Biolegend, EH12.2H7, Cat. 329940), TCF1-PE (Biolegend, 7F11A10, Cat. 655207), CD137-

APC (Miltenyi, REA765, Cat. 130-110-901), Zombie Aqua™ Fixable Viability Kit (Biolegend,

Cat. 423101); in Panel 2: CD3-FITC (Biolegend, UCHT1, Cat. 300406), CD4-PE-Cy7

(Biolegend, RPA-T4, Cat. 300512), CD8-BV650 (Biolegend, SK1, Cat. 344730), CD45-BV711

(Biolegend, HI30, Cat. 304050), CD137-APC-Fire750 (Biolegend, 4B4-1, Cat. 309834),

Granzyme B-APC-Alexa700 (Biolegend, QA16A02, Cat. 372222), Ki-67-BV605 (Biolegend,

Ki-67, Cat. 350522), PD-1-PE-Dazzle (Biolegend, EH12.2H7, Cat. 329940), Perforin – BV421

(Biolegend, dG9, Cat. 308122), TCF1-PE (Biolegend, 7F11A10, Cat. 655207), TOX-APC

(Miltenyi, REA473, Cat. 130-118-335), Zombie Aqua™ Fixable Viability Kit (Biolegend, Cat.

423101); in Panel 3: CD45-BUV395 (BD, HI30, Cat. 563792), CD3-Pacific Blue (Invitrogen,

S4.1, Cat. MHCD0328), CD4-BUV495 (BD, SK3, Cat. 612936), CD8-BUV563 (BD, RPA-T8,

Cat. 612914), CD19-PE-Cy5 (Biolegend, HIB19, Cat. 302210), CD45RA-BV785 (Biolegend,

HI100, 304139), CD62L-BV605 (Biolegend, DREG-56, Cat. 304834), CD39-PE-Fire810

(Biolegend, A1, Cat. 328245), CD69-BUV805 (BD, FN50, Cat. 748763), CD137-APC (Miltenyi,

REA765, Cat. 130-110-901), PD-1-PE-Dazzle (Biolegend, EH12.2H7, Cat. 329940), HLA-DR-

FITC (Biolegend, LN3, Cat. 327006), TCF1-PE (Biolegend, 7F11A10, Cat. 655207), CD160-

PerCP-Cy5.5 (Biolegend, BY55, Cat. 341210), LAG3-BV650 (Biolegend, 11C3C65, Cat.

369316), TIM3 – Alexa Fluor 700 (R&D Systems, Clone 344823, Cat. FAB2365N) , Galectin-9

-PE-Cy7 (Biolegend, 9M1-3, Cat. 348916), TIGIT-APC-Cy7 (Biolegend, A15153G, Cat.

372734), Ki-67-BV510 (Biolegend, Ki-67, Cat. 350518), CD103-BV711 (Biolegend, Ber-ACT8,

Cat. 350222), LIVE/DEAD^{TM} Fixable Blue Dead Cell Stain Kit (Invitrogen, Cat. L23105).

Co-expression analysis

For co-expression analysis of flow cytometric data, cells were manually gated on live, singlet

human CD8$^+$ T cells. Cells were analyzed using CRUSTY webtool (21), where clustering was

performed using Phenograph algorithm and default settings for UMAP generation.

IFN-γ ELISpot and TNFα ELISA

For IFN-γ ELISpot, ELISpot plates (Merck, Cat.MAIPN4550) were activated with 35% ethanol,

washed and coated overnight with anti-IFN-γ coating antibody (mAb-1-D1K, Mabtech, Cat.

3420-3-1000). Plates were washed and T cells were co-cultured with autologous LCL in RPMI/

10% FCS/ 1% Pen/Strep at an effector: target ratio of 5:1 for 16-24 hours. After incubation,

supernatant was harvested, stored at -20°C and used for TNFα ELISA. Wells were washed

and incubated for 2 hours with biotinylated anti-IFN-γ detection antibody (mAb-7-B6-1-Biotin,

Mabtech, Cat. 3420-6-250). Afterwards, wells were washed and incubated with Streptavidin-

ALP (Mabtech, Cat. 3310-10) for 1 hour. Wells were then washed again and spots were

developed with filtered substrate solution BCIP/ NBT-plus (Mabtech, Cat. 3650-10). When

distinct spots were visible, the plate was washed and dried overnight. Plates were acquired

with an ELISpot reader (AID classic, AID GmbH) and spots were automatically counted using

AID ELISpot 8.0 software (AID GmbH).

TNFα ELISA was performed with the supernatant from T cell and LCL co-culture as described

above using the human TNFα ELISA^{BASIC} kit (Mabtech, Cat. 3512-1H-6), following

manufacturer's instructions. Plates were acquired using an ELISA reader (infinite M200Pro,

Tecan) and analyzed with the i-control 2.0 software (Tecan).

Immunohistochemistry

Tumors were fixed in 4% paraformaldehyde and embedded in paraffin. Sample embedding,

preparation and staining was performed by the Pathology department of the University Hospital

Basel. Slides were acquired on an automated slide scanning brightfield microscope (Vectra 3,

Akoya Biosciences) and quantified using InForm software (Akoya Biosciences).

Bulk RNAseq and TCR profiling

RNA was extracted from 5 x 10^6 T cells using the RNeasy® Mini Kit (Qiagen, Cat. 74106). RNA

extraction was performed following the instructions from the "Purification of Total RNA from

Animal Cells Using Spin Technology" protocol given in the RNeasy® Mini Handbook, including

homogenization with QIAshredders (Qiagen, Cat. 79654) and on-column DNA digestion with

the RNase-free DNase Set (Qiagen, Cat. 79254). All further steps were performed by the

Functional Genomics Center Zurich (FGCZ). The quality of the isolated RNA was determined

with a Qubit® (1.0) Fluorometer (Life Technologies, California, USA) and a Bioanalyzer 2100

(Agilent, Waldbronn, Germany). Only those samples with a 260 nm/280 nm ratio between 1.8–

2.1 and a 28S/18S ratio within 1.5–2 were further processed. The TruSeq RNA Sample Prep

Kit v2 (Illumina, Inc, California, USA) was used in the succeeding steps. Briefly, total RNA

samples (600 ng) were poly A enriched and then reverse-transcribed into double-stranded

cDNA. The cDNA samples were fragmented, end-repaired and polyadenylated before ligation

of TruSeq adapters containing the index for multiplexing fragments containing TruSeq adapters

on both ends were selectively enriched with PCR. The quality and quantity of the enriched

libraries were validated using Qubit® (1.0) Fluorometer and the Caliper GX LabChip® GX

(Caliper Life Sciences, Inc., USA). The product is a smear with an average fragment size of

approximately 260 bp. The libraries were normalized to 10nM in Tris-Cl 10 mM, pH8.5 with

0.1% Tween 20. After library quantification, libraries were prepared for loading accordingly to

the NovaSeq workflow with the NovaSeq6000 Reagent Kit (Illumina, Catalog

No. 20012865). Cluster generation and sequencing were performed on a NovaSeq6000

System with a run configuration of single end 100bp. RNA-seq data analysis consisted of the

following steps: The raw reads were first cleaned by removing adapter sequences, trimming

low quality ends, and filtering reads with low quality (phred quality <20) using Fastp (Version

0.20) (22). Sequence pseudo alignment of the resulting high-quality reads to the human

reference genome (build GRCh38.p13 and GENCODE gene models release 37) and

quantification of gene level expression was carried out using Kallisto (Version 0.46.1) (23).

Differential expression was computed using the generalized linear model implemented in the

Bioconductor package DESeq2 (R version: 4.2.2, DESeq2 version: 1.38.1) (24). Genes

showing altered expression with adjusted (Benjamini and Hochberg method) p-value < 0.05

were considered differentially expressed. RNA-seq expression data underwent preprocessing

to remove duplicated genes. Subsequently, we utilized the CreateSeuratObject() function

within the Bioconductor package Seurat (R version: 4.2.2, Seurat version: 5.0.1) (25) to

construct a Seurat object, facilitating downstream analyses. Normalization of the expression

data was performed using the NormalizeData() function, with experimental group assignments

explicitly indicated within the Seurat object. To delineate the gene signatures characterizing

distinct T cell phenotypes, we curated pertinent signatures from previous studies. Genes

absent in our expression data were systematically filtered out. Next, employing the

AddModuleScore() function, we computed module scores for each gene signature across

experimental groups and values were scaled for heatmap visualization. For the TCR analysis

the read alignment to the Human genome was performed with STAR (v2.7.10b) (26). TCR

sequences (CDR3 of TRA, TRB, TRG and TRD) were reconstructed from bulk RNAseq data

using TRUST4 (v1.0.10) (27). Out-of-frame sequences were excluded from the analysis. TCR

repertoire analysis was performed using the R package Immunarch (28).

Statistics

Statistical analysis was performed using Prism 10 Software (GraphPad Software, Inc). Figure

legends indicate which statistical test was performed.

Study approval

Animal experiments were conducted according to licenses approved by the veterinary office of

the canton of Zurich, Switzerland (ZH049/20 and ZH067/2023).

BioRender

Schematics were generated with a licensed version of BioRender.

Author contributions

Conceptualization, J.M. and O.C.; Methodology, J.M., M.K. and O.C.; Investigation, J.M., M.K.,

L.E., L.O. and O.C.; Writing – Original Draft: , J.M. and O.C.; Writing – Review & Editing, , J.M.

C.M. and O.C.; Funding Acquisition, O.C.; Resources, L.O. and C.M.; Supervision, O.C.

Declaration of Interests

The authors declare that they have no competing interests.

Acknowledgments

We would like to thank Fernando Canale for help with gene signature analysis and Daniel E.

Speiser for critically reading the manuscript. Funding: This research was supported by Swiss

Cancer Research foundation (KFS-5292-02-2021 to O.C.).

Data Sharing

Supplemental data are provided with this paper. All data are also available upon request.

Results

Establishment of an autologous HIS mouse tumor model

We established and characterized a novel tumor model in which immunocompetent, humanized mice are subcutaneously implanted with autologous human tumor cells. For this aim, we isolated CD34$^+$ hematopoietic progenitor cells (HPC) and CD19$^+$ B cells from the same human HPC donor tissue. CD34$^+$ HPCs were engrafted into NSG mice to reconstitute components of the human immune system (from here on referred to as human immune system (HIS) mice), while CD19$^+$ B cells were transformed into lymphoblastoid cell lines (LCL) by infection with the oncogenic Epstein-Barr virus (EBV) (**Figure 1A**).

To characterize LCL tumorigenicity and tumor growth in the presence or absence of an autologous human immune system, HIS mice reconstituted with HPCs autologous to the tumor or NSG mice were injected subcutaneously with LCLs. In both recipient strains, palpable tumors formed at the site of implantation. In HIS mice, in the presence of an endogenous immune system, tumors showed progressive growth until approximately day 18 and subsequently, a majority of tumors showed regression (**Figure 1B**). In contrast, in NSG mice without an endogenous immune system, tumors increased in size until termination criteria were reached (**Figure 1C**). These data suggested a functional antitumor response mediated by components of the reconstituted human immune system, i.e. immunocompetence, in our autologous HIS mouse tumor model.

Endogenous antitumor immune response in HIS mice

To investigate the tumor regression that we observed in the presence of an autologous human immune system, we next characterized the immune response in tumor-bearing HIS mice. Around the same time of tumor regression, we found weight loss in HIS mice (**Figure 1D**), but not in tumor-bearing NSG mice (**Figure 1E**). This indicated immunopathology related to the endogenous human antitumor immune response, reminiscent of cancer cachexia (29) rather than an effect of tumor growth per se. Additionally, we found increases in spleen-to-body weight ratio, total splenocyte count and white blood cell count in tumor-bearing HIS mice (**Figure 1, F-H**), suggesting immune cell proliferation occurring in response to tumor growth. Furthermore,

tumors in HIS mice showed infiltration by endogenous human immune cells, including CD8$^+$ T

cells (Supplemental Figure 1).

The frequency of human CD3$^+$ T cells in tumor-bearing HIS mice in spleen (**Figure 2A**)

and circulation (Supplemental Figure 2A) increased significantly compared to naïve HIS mice.

Moreover, tumor-bearing HIS mice showed an increased frequency of CD8$^+$ T cells, suggesting

preferential expansion (**Figure 2A** and Supplemental Figure 2A). These CD8$^+$ T cells exhibited

differentiation towards an effector memory (T$_{EM}$) phenotype and, to a lower degree, central

memory (T$_{CM}$) phenotype, whereas non-tumor bearing HIS mice had higher frequencies of

naïve T cells (T$_{naïve}$) (**Figure 2B** and Supplemental Figure 2B). Consistent with an exhausted

effector-like phenotype, CD8$^+$ T cells from tumor-bearing HIS mice also showed enhanced

expression of the exhaustion/activation markers PD-1, CD39 and HLA-DR (**Figure 2C** and

Supplemental Figure 2C) as well as trending towards increased TOX expression (**Figure 2D**

and Supplemental Figure 2D), while exhibiting significantly reduced levels of the

progenitor/stemness-related marker TCF1 (30) (**Figure 2D** and Supplemental Figure 2D).

Similar to the increased CD8$^+$ T cell frequencies in secondary lymphoid tissue (spleen)

and circulation, tumors in HIS mice exhibited predominant infiltration by CD8$^+$ T cells (**Figure**

2E and Supplemental Figure 1C). The majority of autologous tumor-infiltrating human CD8$^+$ T

cells had a T$_{EM}$-like phenotype, with a lower proportion of T$_{CM}$ and greatly reduced proportions

of T$_{EMRA}$ and naïve T cells (**Figure 2F**). Tumor-infiltrating CD8$^+$ T cells displayed high

expression of exhaustion and activation markers PD-1 (91.0%±12.1), CD39 (56.4%±22.8) and

HLA-DR (53.9%±22.4) (**Figure 2G**), low levels of TCF1 (5.6%±5.0) (**Figure 2H**), while a

fraction of tumor-infiltrating CD8$^+$ T cells expressed TOX (21.7%±14.81) (**Figure 2H**).

Dimensionality reduction by UMAP revealed obvious differences of CD8$^+$ T cell landscapes

between tumor-infiltrating lymphocytes (TIL) and spleen (**Figure 2I**). Together, the phenotype

of human CD8$^+$ T cells in tumor-bearing HIS mice was reminiscent of CD8$^+$ T cell phenotypes

described in human cancer (31,32).

Progenitor exhausted T cells (T$_{pex}$; TCF1$^+$PD-1$^+$) play an important role for immunotherapy

of human cancer. T$_{pex}$ have been described as precursors of exhausted cytotoxic T cells (T$_{ex}$;

TCF1$^-$PD-1$^{+\cdot}$, further subdivided into TCF1$^-$PD-1$^+$CD69$^-$ intermediate exhausted (T$_{ex-int}$) and

TCF1$^-$PD-1$^+$CD69$^+$ terminally exhausted (T$_{ex\text{-}term}$) T cells (33)), which exhibit effector function

but reduced self-renewal and long-term survival, making T$_{pex}$ cells an important reservoir of

antitumor T cells (33–35). Interestingly, we detected an increased fraction of CD8$^+$ T$_{pex}$ cells in

secondary lymphoid tissue (spleen) of tumor-bearing HIS mice compared with tumor (**Figure**

2J). This distribution recapitulates human cancers, where a higher frequency of T$_{pex}$ is found

in lymph nodes compared to the tumor site (34). In addition, we found a significantly higher

proportion of terminally exhausted CD8$^+$ T cells (T$_{ex\text{-}term}$; TCF1$^-$PD-1$^+$CD69$^+$) within the tumor

compared to the spleen, a secondary lymphoid organ (**Figure 2J**), again consistent with what

has been described in human cancer (34,35).

Tissue-resident-memory (T$_{RM}$) T cells (CD69$^+$CD103$^+$) within tumors are another T cell

subset of interest for immunotherapy research. The presence of T$_{RM}$ cells has been associated

with a favorable outcome in certain cancer types and based on their expression of immune

checkpoint molecules, tumor-reactive T$_{RM}$ cells also constitute a target for immune checkpoint

blockade (36). We detected T$_{RM}$-like (CD69$^+$CD103$^+$) CD8$^+$ T cells in tumor-bearing HIS mice

and found a significantly higher abundance in tumor compared with spleen (**Figure 2K**).

The presence and identification of these human CD8$^+$ T cell subsets, which are considered

relevant in human cancer and are thought to play critical roles in response to immunotherapy,

underscores the translational value of our humanized mouse model for immunotherapy

research.

Activated and exhausted-like, cytotoxic effector-like phenotype of human CD137$^+$ CD8$^+$

T cells

In human cancers, CD137 has been described as a marker for tumor-reactive T cells (37–39).

We similarly identified a population of human CD137$^+$ T cells in tumor-bearing mice (**Figure**

2L) and characterized the CD8$^+$ T cell population further. Likewise, PD-1 has been reported to

be a marker of tumor-reactive T cells in human cancers, both in circulation and within tumor-

infiltrating lymphocytes (20,40,41). We observed that of splenic CD8$^+$ T cells from tumor-

bearing HIS mice, only a small proportion (10.9%±11.2) were CD137 positive, whereas the

majority of CD8$^+$ T cells (83.5%±12.1) expressed PD-1 on their surface (**Figure 2M**). Though

there was a small but still significant difference between the overall high level of PD-1

expression on CD137$^+$CD8$^+$ and CD137$^-$CD8$^+$ T cells from tumor-bearing HIS mice, both these

populations expressed PD-1 significantly more than bulk CD8$^+$ T cells from naïve HIS mice

(**Figure 2N**). These data suggested that CD137 alone might not be sufficient to delineate

tumor-reactive T cell subsets. We therefore included a CD8$^+$ T cell subset negative for both

CD137 and PD-1 in our analysis to further explore and distinguish bona fide tumor-reactive

from bystander CD8$^+$ T cells in tumor-bearing HIS mice.

Indeed, compared to double negative CD137$^-$PD-1$^-$ CD8$^+$ T cells, CD137$^+$ CD8$^+$ T cells and

CD137$^-$ CD8$^+$ T cells displayed a T_{EM}-like phenotype, while the double negative CD137$^-$PD-1$^-$

population was enriched in naïve-like T cells (Supplemental Figure 2E). Within the splenic

CD137$^+$ CD8$^+$ T cell population, we found the highest expression of the exhaustion marker

CD39, which has been described as a marker of CD8$^+$ T cells with enhanced tumor reactivity

across different human cancers (1,42–46), the activation marker HLA-DR as well as the

activation and residency-related marker CD69 (47), while this cell subset also had the

strongest expression levels of Ki-67, indicating high proliferation (**Figure 2O** and Supplemental

Figure 2F). Additionally, within the CD137$^+$ CD8$^+$ T cell population, we observed significantly

increased expression of TOX, a transcription factor associated with exhaustion (48) and

drastically reduced levels of the progenitor/stemness-related transcriptional regulator TCF1

(30) (**Figure 2O** and Supplemental Figure 2G). Several makers related to T cell exhaustion

and activation (49,50) (TIGIT, TIM3, CD160, Galectin-9) were similarly upregulated on CD137$^+$

CD8$^+$ T cells (**Figure 2O** and Supplemental Figure 2H). Finally, CD137$^+$CD8$^+$ T cells displayed

the highest levels of the cytolytic molecules granzyme B and perforin compared with CD137$^-$

CD8$^+$ T cells or CD137$^-$PD-1$^-$ CD8$^+$ T cells from tumor-bearing HIS mice (**Figure 2O** and

Supplemental Figure 2I).

The CD137$^+$ CD8$^+$ T cell population contained a significantly higher proportion of $T_{ex-term}$

cells in both spleen and tumor and a significantly lower frequency of T_{pex} in the tumor relative

to the CD137$^-$ CD8$^+$ T cell population (Supplemental Figure 2J and 2K). As CD69 is expressed

on subsets of both exhausted as well as T_{RM}-like T cells (33), we quantified CD8$^+$ T_{RM}-like

(CD69$^+$CD103$^+$) T cells and found no difference between CD137$^+$ and CD137$^-$ T cells in the

spleen (Supplemental Figure 2L). In the tumor, where an overall higher proportion of T_{RM}-like

T cells was present compared to spleen (**Figure 2K**), the CD137$^+$ CD8$^+$ T cell population

contained significantly fewer T_{RM}-like T cells compared to CD137$^-$ CD8$^+$ T cells (Supplemental

Figure 2M). Thus, the majority of CD137$^+$ CD8$^+$ T cells did not exhibit a T_{RM}-like phenotype, but

consisted of more $T_{ex-term}$ cells relative to CD137$^-$ CD8$^+$ T cells with only a minor proportion of

T_{pex} cells.

Analysis of CD8$^+$ T cell clusters (Supplemental Figure 3A) from spleen and tumor

demonstrated that CD137 expressing clusters 2, 7 and 10 (Supplemental Figure 3, B and C)

co-expressed a large number of markers related to activation/exhaustion (TIGIT, TIM3, PD-1,

CD39, CD160). CD8$^+$ T cells in cluster 7, the majority of which was localized in the tumor

(Supplemental Figure 3B), additionally expressed LAG-3 (Supplemental Figure 3C). Moreover,

co-expression analysis confirmed the presence of a T_{pex}-like cluster (TCF1$^+$PD1$^+$CD69$^-$), which

was preferentially found in the spleen (cluster 3). Also, a T_{RM}-like cluster (CD69$^+$CD103$^+$) was

detected, which was preferentially found in tumor (cluster 5) (Supplemental Figure 3C).

Altogether, de novo arising highly proliferative CD137$^+$ CD8$^+$ T cells exhibited an activated

and exhausted-like, cytotoxic effector and least stem-like phenotype relative to the other

human CD8$^+$ T cell populations we identified in tumor-bearing HIS mice.

Maintained expansion capacity and enrichment of *in vitro* tumor reactivity within an

activated and exhausted-like human CD8$^+$ T cell subset

To investigate tumor reactivity of CD137$^+$ CD8$^+$ T cells relative to the other T cell subsets,

splenic T cells were expanded *ex vivo* (**Figure 3A**). Using a rapid expansion protocol (20), we

observed an average expansion of >1000-fold after 14 days for all subsets. Interestingly,

although terminally exhausted CD8$^+$ T cells have generally been thought to contain reduced

proliferative potential (51), there were no significant differences in the expansion capacity of

CD137$^+$ CD8$^+$ T cells displaying the most exhausted-like, $T_{ex-term}$ enriched phenotype (Figure

2O and Supplemental Figure 2, J and K) compared to the other CD8$^+$ T cell populations from

tumor-bearing HIS mice or bulk CD8$^+$ T cells from naïve HIS mice (**Figure 3B**). All of the *ex*

vivo expanded T cell subsets from tumor-bearing HIS mice exhibited differentiation biased to

an effector memory-like phenotype (**Figure 3C**). While prior to *ex-vivo* expansion both PD-1

and CD39 were strongly increased in CD137$^+$ CD8$^+$ T cells (Figure 2, O and N), both markers

were no longer significantly upregulated after expansion. In contrast, HLA-DR remained

upregulated on CD137$^+$ CD8$^+$ T cells relative to CD137$^-$PD1$^-$ CD8$^+$ and bulk CD8$^+$ T cells from

naïve mice (**Figure 3D**).

After co-culture with autologous tumor cells, expanded CD137$^+$ CD8$^+$ T cells showed the

highest IFN-γ production by ELISpot, which gradually decreased from CD137$^-$ CD8$^+$ T cells to

CD137$^-$PD-1$^-$ CD8$^+$ T cells (**Figure 3E**). Similarly, TNFα secretion was highest in supernatants

containing CD137$^+$ CD8$^+$ T cells after co-culture with autologous tumor cells and lowest in

CD137$^-$PD-1$^-$ CD8$^+$ T cells (**Figure 3F**).

Together, these data indicated enrichment of tumor recognition within CD137$^+$ CD8$^+$ T cells

that maintained *ex vivo* proliferative potential, while double negative CD137$^-$PD-1$^-$ CD8$^+$ T cells

behaved as bystander cells.

Transcriptome analysis reveals tumor-reactive signature in CD137$^+$ CD8$^+$ T cells

To gain insight into differentially expressed genes (DEG) in tumor-reactive CD137$^+$ CD8$^+$ T

cells versus CD137$^-$ CD8$^+$ T cells and bystander CD137$^-$PD-1$^-$ CD8$^+$ T cells as well as bulk

CD8$^+$ T cells from naïve HIS mice, whole transcriptome RNA sequencing of *ex vivo* expanded

T cell subsets was performed. We found a higher number of DEG for the tumor-reactive

(CD137$^+$) gene set versus bystander CD8$^+$ T cells from tumor-bearing HIS mice (200 DEG)

and versus bulk CD8$^+$ T cells from naïve HIS mice (300 DEG) in comparison to the number of

DEG we found for the comparison against CD137$^-$ CD8$^+$ T cells (53 DEG) (**Figure 4A**), the

subset with the second highest tumor reactivity *in vitro* (Figure 3, D and E). Principal

component analysis (PCA) was consistent with functional data of tumor-specific cytokine

production (Figure 3, D and E), showing that the 4 subsets clustered in a gradual way from

tumor-reactive (CD137$^+$) to less tumor-reactive (CD137$^-$) and bystander (CD137$^-$PD-1$^-$) CD8$^+$

T cells and finally to bulk CD8$^+$ T cells from naïve HIS mice, with the biggest overlap between

CD137$^-$PD-1$^-$ bystander CD8$^+$ T cells and bulk human CD8$^+$ T cells from naïve HIS mice

(**Figure 4B**).

The transcriptomic profile that we found in tumor-reactive CD137[+] CD8[+] T cells suggested

an exhausted/activated-like phenotype with increased expression of *TNFRSF9, LAG3,*

PDCD1, TIGIT and *HLA-DPA1/HLA-DQA1* (**Figure 4, C, E and F**). We also observed

enhanced expression of effector molecules associated with cytotoxicity (*GZMH, NKG7* and

PRF1) as well as *IFNG* (**Figure 4, C, E and F**). Expression of the naïve or stem-like genes

CCR7, *LEF1* and *BACH2* were diminished (**Figure 4, D-F**) while the senescence marker CD57

(52) (*B3GAT1*) and the transcription factor *EOMES* showed the highest transcript level in the

CD137[+] subset (**Figure 4, C and F**). Although EOMES has been shown to be involved in T cell

exhaustion, it has also been reported to be commonly co-expressed with effector molecules in

effector T cells and to be essential for anticancer T cell function (53,54).

Among upregulated Gene Ontology (GO) terms comparing tumor-reactive CD137[+] CD8[+]

T cells against the other three populations, we found the terms interferon-gamma mediated

signaling (GO:0060333), antigen processing and presentation (GO:0019882), antigen

processing and presentation of peptide or polysaccharide antigen via MHC class II

(GO:0002504), immune response (GO:0006955) and adaptive immune response

(GO:0002250) (**Figure 4G**). Upregulation of these pathways in the CD137[+] population

indicated stronger T cell activation and response in comparison to the other populations.

To determine T cell states after *ex vivo* expansion, we compared our transcriptome data

to published datasets of different T cell states from human cancers (31,46,55–59). A high

overlap of signatures originally found in tumor-specific CD8[+] T cells from human cancers

describing terminally exhausted, exhausted, activated, and progenitor exhausted T cell states

was found for CD137[+] CD8[+] T cells (**Figure 4H**). Overlap with these different signatures might

hint to a degree of heterogeneity within the CD137[+] CD8[+] T cell population after *ex vivo*

expansion but confirmed the overall activated/exhausted-like state of these cells that we

already observed before *ex vivo* expansion. Signatures for tissue-resident memory and CD8[+]

stem-like memory were downregulated in the CD137[+] CD8[+] population. Strikingly, comparing

tumor-reactive CD137[+] CD8[+] T cells to bystander CD137[-]PD-1[-] CD8[+] T cells or bulk CD8[+] T

cells from naïve HIS mice, gene sets that have been described to define tumor-reactive T cells

in patients with cancer were significantly upregulated ('MANA TIL' (58), 'tumor specific' (56))

(**Figure 4I**). These results suggested that the antitumor T cell response in autologous tumor-

bearing HIS mice mirrors relevant aspects of the T cell response in cancer patients, thus further

supporting the potential translational value of our model.

Correlation of clonal expansion with effector function

As another surrogate for tumor recognition, we analyzed clonality of the CD8$^+$ T cell receptor

(TCR) repertoire, anticipating tumor-reactive CD8$^+$ T cells would have higher clonal expansion

and thus lower diversity than bystander CD8$^+$ T cells. Indeed, we found the total number of

clonotypes tending to be the smallest in the tumor-reactive CD137$^+$ subset, and to gradually

increase from the less tumor-reactive CD137$^-$ subset to CD137$^-$PD-1$^-$ bystander CD8$^+$ T cells

to bulk CD8$^+$ T cells from naïve HIS mice (**Figure 5A**), following the gradient of tumor

recognition we observed by tumor-specific cytokine production *in vitro* (Figure 3, D and E).

Likewise, the estimated sample diversity for the tumor-reactive CD137$^+$ subset was lower than

for the other populations (**Figure 5B** and Supplemental Figure 4A). In line with this, the CD137$^+$

population had a reduced abundance of rare or very small clonotypes (**Figure 5C** and

Supplemental Figure 4, B and C). Conversely, tumor-reactive CD137$^+$ CD8$^+$ T cells from tumor-

bearing HIS mice demonstrated a significantly higher abundance of hyperexpanded clones

(**Figure 5D**) with the 20 biggest clones occupying a significantly larger repertoire space than

in the other populations, suggesting polyclonal expansion (Supplemental Figure 4B).

To track the overlap of clonotypes between individual experiments (i.e. donor dependency)

and populations, we next performed TCR repertoire overlap analysis (**Figure 5E**). We found

that the tumor-reactive CD137$^+$ population showed the biggest overlap with the CD137$^-$ subset

and only to a minor extend an overlap with CD137$^-$PD-1$^-$ bystander CD8$^+$ T cells or bulk CD8$^+$

T cells from naïve HIS mice (**Figure 5, E and G**). Also, analyzing the repertoire of the bulk

naïve CD8$^+$ T cell population for each experiment by cross-comparing to each population from

each experiment, the overlap with the CD137$^+$ population was significantly less than the

overlap with every other population (**Fig. 5F**). These findings suggested that clonotypes arising

due to the conditions of expansion were less present in the CD137$^+$ population and that other,

presumably tumor-reactive, clonotypes expanded.

In addition, we observed that the expansion of individual bona fide tumor-reactive

clonotypes was highly donor specific. When comparing the 10 most abundant clonotypes of

the CD137[+] population per experiment (i.e. per donor), almost no overlap with the CD137[+]

TCR repertoire of any other experiment was detected (**Figure 5G** and Supplemental Figure

4D). Finally, to explore an association of clonal expansion with functionality, we correlated the

abundance of the top 10 bona fide tumor-reactive clonotypes of the CD137[+] population with *in*

vitro tumor reactivity and found a highly significant correlation of clone size abundance and

tumor-specific IFN-γ production by CD8[+] T cells (**Figure 5H**). Overall, these data indicated that

CD137[+] CD8[+] T cells derived from tumor-bearing HIS mice expanded polyclonally in a human

donor- and tumor-specific manner.

Human CD8[+] T cells with an activated and exhausted-like phenotype reduce tumor

growth *in vivo*

We next performed adoptive cell transfer (ACT) experiments of *ex vivo* expanded CD8[+] T cells

in NSG mice previously engrafted with autologous tumors (**Figure 6A**). We initially selected

NSG mice as recipients to unambiguously track the transferred T cells and to be able to

distinguish their anticancer activity from the endogenous antitumor T cell response in HIS mice.

In NSG mice, only ACT with expanded CD137[+] CD8[+] T cells resulted in reduced tumor growth

in a substantial number of recipients, while transfer of bystander CD8[+] T cells from tumor-

bearing HIS mice or bulk CD8[+] T cells from naïve HIS mice did not (**Figure 6, B-D**). Specifically,

adoptive transfer of human CD137[+] CD8[+] T cells from tumor-bearing HIS donors resulted in a

partial response ($\geq$30% tumor reduction compared to 'tumor only' group) in almost half of

recipients (6 out of 13 tumor-bearing NSG recipients, 46%), while this was observed in only

13% of NSG mice treated with CD137[-]PD-1[-] CD8[+] T cells from tumor-bearing HIS mice (1/8)

and no such response (0%) was seen in NSG recipients of bulk CD8[+] T cells from naïve HIS

mice (0/6) (**Figure 6D**).

Furthermore, a higher frequency of T cells was found in tumor (**Figure 6, E and F**), blood

(Supplemental Figure 5A) and spleen (Supplemental Figure 5B) of tumor-bearing NSG mice

receiving CD137[+] CD8[+] T cells compared to recipients transferred with any of the other cell

populations. Flow cytometric analysis of tumor-infiltrating lymphocytes (TIL) revealed high

CD3$^+$ T cell frequencies (53.65%±21.94) after ACT of expanded CD137$^+$CD8$^+$T cells, but only

11.64%±11.0 after transfer of bystander CD8$^+$ T cells and 7.25%±5.38 after transfer of bulk

CD8$^+$ T cells from naïve HIS mice (**Figure 6E**). Likewise, immunohistochemistry of tumors of

NSG mice confirmed T cell infiltration selectively after ACT with CD137$^+$CD8$^+$ T cells (**Figure**

6F).

Interestingly, while infusion products mainly contained a mix of T$_{EM}$-like and T$_{EMRA}$CD8$^+$ T

cells (Figure 3C), almost exclusively T$_{EM}$-like cells were recovered from blood (Supplemental

Figure 5C), spleen (Supplemental Figure 5D) and TIL (Supplemental Figure 5E) of recipient

NSG mice. Of note, whereas PD-1 was highly expressed on splenic CD137$^+$CD8$^+$ T cells in

tumor-bearing HIS mice (96.49%±3.1) (Figure 2N), PD-1 expression on this population

decreased during antigen-independent *ex vivo* expansion (32.45%±29.17) (Supplemental

Figure 5F). However, after adoptive transfer of expanded CD137$^+$CD8$^+$ T cells into tumor-

bearing NSG mice, PD-1 expression among CD8$^+$ T cells increased again and was significantly

higher in TIL (96.7%±3.65) (**Figure 6G**), blood (Supplemental Figure 5G) and spleen

(Supplemental Figure 5H) compared to NSG mice transferred with bystander CD8$^+$ T cells or

bulk CD8$^+$ T cells. Apart from the superior tumor control, these data further indicated

enrichment of tumor reactivity within CD137$^+$CD8$^+$ T cells, with very high PD-1 expression

serving as a potential marker for productive tumor recognition, but not loss of effector function.

As another index of the anticancer activity of CD137$^+$CD8$^+$ T cells, the degree of tumor

infiltration with CD8$^+$ T cells after ACT was inversely correlated with tumor size (**Figure 6H**).

Thus, the higher frequency of CD8$^+$ T cells in tumors was likely driven by increased tumor

recognition of transferred CD137$^+$CD8$^+$ T cells, and these cells mediated tumor cell killing.

Lastly, ACT of tumor-reactive T cells into autologous tumor-bearing HIS recipient mice

(**Figure 6I**) presented an opportunity to model polyclonal TIL transfer therapy in cancer patients

(60) in an immunocompetent humanized tumor setting. As observed with NSG recipients,

transfer of polyclonal tumor-reactive CD137$^+$CD8$^+$ T cells into autologous tumor-bearing HIS

recipient mice reduced tumor growth (**Figure 6, J-L**). A partial response (≥30% tumor reduction

compared to 'tumor only' group) was observed in 77% of HIS mice (10/13) receiving adoptive

transfer of autologous human CD137$^+$ CD8$^+$ T cell derived from tumor-bearing HIS donors,

while no such response (0%) was observed in any of the HIS mice (0/9) adoptively transferred

with autologous CD137$^-$ or CD137$^-$PD-1$^-$ CD8$^+$ T cells derived from tumor-bearing HIS donors

(**Figure 6L**). Importantly, this was achieved without prior conditioning of HIS recipient mice or

cytokine support, and the number of transferred tumor-reactive CD137$^+$ CD8$^+$ T cells was

fivefold lower compared with transfer in NSG recipients. These data established superior

anticancer activity of CD137$^+$ CD8$^+$ T cells with an activated and exhausted-like phenotype in

the most complete humanized setting.

Altogether, these adoptive transfer experiments confirmed anticancer activity of human

CD137$^+$ CD8$^+$ T cells that developed in autologous tumor-bearing HIS mice, suggesting that

our model captured relevant aspects of the immunobiology of human antitumor T cell

responses.

Discussion

Using in-depth phenotypic analyses, we established the presence of major, widely recognized human CD8$^+$ T cell subsets known to be important in tumor immunology, like progenitor exhausted (T$_{pex}$; TCF1$^+$PD-1$^+$), terminally exhausted (T$_{ex-term}$; TCF1$^-$PD-1$^+$CD69$^+$) or tissue resident-like (T$_{RM}$; CD69$^+$CD103$^+$) T cells in our autologous humanized tumor model. Interestingly, we found CD8$^+$ T$_{pex}$ cells to be more abundant in secondary lymphoid tissue (spleen) compared to tumor and vice versa CD8$^+$ T$_{ex-term}$ cells to be more abundant in tumor compared to spleen. This seems consistent with the mechanisms of formation of antitumor immunity as currently understood (61). Likewise, tissue-resident-like (T$_{RM}$) T cells were enriched in the tumor relative to spleen, as would be expected. Furthermore, we identified a proliferative human CD8$^+$ T cell subset in tumor-bearing HIS mice that expressed activation and exhaustion markers including very high levels of PD-1 and CD39 while having low expression of the progenitor marker TCF1 but high expression of TOX. This CD137$^+$ CD8$^+$ T cell subset, while being PD-1^{hi}CD39$^+$, displayed increased abundance of a T$_{ex-term}$ phenotype with co-expression of inhibitory receptors like TIM-3, TIGIT, CD160 and galectin-9. At the same time, CD137$^+$ CD8$^+$ T cell contained fewer T$_{RM}$-like cells, which was reflected by a relative lack of a tissue-resident signature even after expansion. However, we note that the expansion, which was necessary to achieve the cell numbers required for adoptive transfer, may have led to some alteration of populations, as evidenced, for example, in the downregulation of PD-1 and CD39 on the protein level after expansion. Nevertheless, most activation and exhaustion markers were still increased on the transcript level in expanded CD137$^+$ CD8$^+$ T cells relative to the other populations. Furthermore, expanded CD137$^+$ CD8$^+$ T cells from tumor-bearing HIS mice showed enrichment in gene signatures of exhaustion. Functionally, CD137$^+$ CD8$^+$ T cells exhibited increased tumor reactivity as demonstrated by enhanced cytokine production after coculture with autologous tumor cells. Importantly, by means of adoptive transfer, we uniquely provide direct *in vivo* evidence of superior anticancer activity of these human tumor-reactive effector CD8$^+$ T cells displaying an activated and exhausted-like phenotypic state.

We have leveraged this human CD8$^+$ T cell subset arising de novo in tumor-bearing HIS mice for transcriptomic analysis and TCR profiling to demonstrate across donors overlap with

tumor-reactive CD8[+] T cell signatures from human cancers (56,58) as well as tumor-induced

polyclonal expansion. Polyclonality of neoantigen-specific CD8[+] T cells in blood and tumor has

recently been shown to be associated with response to PD-1 immunotherapy in patients with

melanoma (16). Congruence of our model data across donors with these clinical benchmarks

suggests robustness and translational validity of our findings.

Of note, CD39[+]CD8[+] T cells expressing high levels of PD-1 have been described to be

enriched for tumor reactivity and clonal proliferation, predicting efficacy of immunotherapy in

human lung cancer (46). Likewise, the abundance of intratumoral CD39[+]CD8[+] T cells

displaying upregulated exhaustion markers has been shown to be prognostic for improved

progression-free and overall survival in individuals with treatment-naïve early stage lung

cancer (62). Furthermore, PD-1[hi]CD39[+] CD8[+] T cells correlated with improved survival in

patients with endometrial cancer (45) and the signature of similar CD8[+] T cells with increased

TOX expression predicted survival in patients with breast cancer (44). Recently, pre-existing

exhausted-like tumor-reactive CD8[+] T cells have been associated with favorable outcome to

immunotherapy in head and neck cancer (63). On the other hand, higher abundance of TCF1[+]

stem-like CD8[+] T cells with a CD39[-]CD69[-] phenotype in TIL products was associated with

better response after TIL transfer in patients with melanoma (64). However, by single-cell

analysis of lung tumors, this phenotype could not be identified in neoantigen-reactive T cells

(65). While we have not addressed and thus cannot exclude a contribution of the minor TCF1[+]

fraction within the human tumor-reactive CD8[+] T cells in our model, our data emphasize the

key importance of exhausted-like effector CD8[+] T cells in the antitumor immune response.

The striking phenotypic similarity of tumor-reactive CD8[+] T cells that have been shown to

correlate with clinical benefit and the polyclonally expanded human tumor-reactive CD8[+] T cell

cells from tumor-bearing HIS mice that we demonstrated to possess superior *in vivo* anticancer

activity highlights the translational potential of our model. Hence, the novel immunocompetent

human tumor model that we developed using HIS mice with human immune cells reconstituted

from hematopoietic progenitor cells that are autologous to tumor might bridge a gap between

syngeneic mouse tumor models and human-specific antitumor immune responses (13,66). Our

model demonstrates hallmarks of human antitumor T cell responses and thus may provide a

tool that should allow further investigation of drivers of anticancer activity in human CD8[+] T cells in cancer immunotherapy research. Overall, the endogenous human immune system in HIS mice markedly influenced tumor and host, allowing for experimental interrogation of the interfaces of human tumor and autologous human antitumor immunity in a small animal model.

Together, fully activated effector and exhausted-like human CD8[+] T cells directly mediate anticancer activity. Our data support clinical exploration of exhausted-like effector CD8[+] T cell subsets alongside the recently more favored stem-cell like phenotype for use in adoptive transfer and engineered T cell therapies.

References

1. Simoni Y, Becht E, Fehlings M, Loh CY, Koo SL, Teng KWW, et al. Bystander CD8+ T cells are abundant and phenotypically distinct in human tumour infiltrates. Nature. 2018 May;557(7706):575–9.
2. Scheper W, Kelderman S, Fanchi LF, Linnemann C, Bendle G, de Rooij MAJ, et al. Low and variable tumor reactivity of the intratumoral TCR repertoire in human cancers. Nat Med. 2019 Jan;25(1):89–94.
3. Lan X, Zebley CC, Youngblood B. Cellular and molecular waypoints along the path of T cell exhaustion. Sci Immunol. 2023 Sep;8(87):eadg3868.
4. Lowery FJ, Krishna S, Yossef R, Parikh NB, Chatani PD, Zacharakis N, et al. Molecular signatures of antitumor neoantigen-reactive T cells from metastatic human cancers. Science. 2022 Feb 25;375(6583):877–84.
5. Oliveira G, Wu CJ. Dynamics and specificities of T cells in cancer immunotherapy. Nat Rev Cancer. 2023 May;23(5):295–316.
6. Korman AJ, Garrett-Thomson SC, Lonberg N. The foundations of immune checkpoint blockade and the ipilimumab approval decennial. Nat Rev Drug Discov. 2022 Jul;21(7):509–28.
7. Ribas A, Wolchok JD. Cancer immunotherapy using checkpoint blockade. Science. 2018 Mar 23;359(6382):1350–5.
8. Powles T, Yuen KC, Gillessen S, Kadel EE, Rathkopf D, Matsubara N, et al. Atezolizumab with enzalutamide versus enzalutamide alone in metastatic castration-resistant prostate cancer: a randomized phase 3 trial. Nat Med. 2022 Jan;28(1):144–53.
9. Guan X, Polesso F, Wang C, Sehrawat A, Hawkins RM, Murray SE, et al. Androgen receptor activity in T cells limits checkpoint blockade efficacy. Nature. 2022 Jun;606(7915):791–6.
10. Palmer AC, Izar B, Hwangbo H, Sorger PK. Predictable Clinical Benefits without Evidence of Synergy in Trials of Combination Therapies with Immune-Checkpoint Inhibitors. Clin Cancer Res. 2022 Jan 15;28(2):368–77.
11. Zong W, Rahman T, Zhu L, Zeng X, Zhang Y, Zou J, et al. Transcriptomic congruence analysis for evaluating model organisms. Proc Natl Acad Sci U S A. 2023 Feb 7;120(6):e2202584120.
12. Gilbertson SE, Walter HC, Gardner K, Wren SN, Vahedi G, Weinmann AS. Topologically associating domains are disrupted by evolutionary genome rearrangements forming species-specific enhancer connections in mice and humans. Cell Rep. 2022 May 3;39(5):110769.
13. Medetgul-Ernar K, Davis MM. Standing on the shoulders of mice. Immunity. 2022 Aug 9;55(8):1343–53.
14. Dong MB, Wang G, Chow RD, Ye L, Zhu L, Dai X, et al. Systematic Immunotherapy Target Discovery Using Genome-Scale In Vivo CRISPR Screens in CD8 T Cells. Cell. 2019 Aug 22;178(5):1189-1204.e23.
15. Chen Z, Arai E, Khan O, Zhang Z, Ngiow SF, He Y, et al. In vivo CD8+ T cell CRISPR screening reveals control by Fli1 in infection and cancer. Cell. 2021 Mar 4;184(5):1262-1280.e22.
16. Puig-Saus C, Sennino B, Peng S, Wang CL, Pan Z, Yuen B, et al. Neoantigen-targeted CD8+ T cell responses with PD-1 blockade therapy. Nature. 2023 Mar;615(7953):697–704.
17. De La Rochere P, Guil-Luna S, Decaudin D, Azar G, Sidhu SS, Piaggio E. Humanized Mice for the Study of Immuno-Oncology. Trends Immunol. 2018 Sep;39(9):748–63.
18. Hiwarkar P, Qasim W, Ricciardelli I, Gilmour K, Quezada S, Saudemont A, et al. Cord

blood T cells mediate enhanced antitumor effects compared with adult peripheral blood T cells. Blood. 2015 Dec 24;126(26):2882–91.

19. Chijioke O, Müller A, Feederle R, Barros MHM, Krieg C, Emmel V, et al. Human natural killer cells prevent infectious mononucleosis features by targeting lytic Epstein-Barr virus infection. Cell Rep. 2013 Dec 26;5(6):1489–98.

20. Gros A, Tran E, Parkhurst MR, Ilyas S, Pasetto A, Groh EM, et al. Recognition of human gastrointestinal cancer neoantigens by circulating PD-1+ lymphocytes. J Clin Invest. 2019 Nov 1;129(11):4992–5004.

21. Puccio S, Grillo G, Alvisi G, Scirgolea C, Galletti G, Mazza EMC, et al. CRUSTY: a versatile web platform for the rapid analysis and visualization of high-dimensional flow cytometry data. Nat Commun. 2023 Sep 4;14(1):5102.

22. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics. 2018 Sep 1;34(17):i884–90.

23. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq quantification. Nat Biotechnol. 2016 May;34(5):525–7.

24. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 2014 Dec;15(12):550.

25. Hao Y, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, et al. Dictionary learning for integrative, multimodal and scalable single-cell analysis. Nat Biotechnol. 2023 May 25;

26. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 2013 Jan 1;29(1):15–21.

27. Song L, Cohen D, Ouyang Z, Cao Y, Hu X, Liu XS. TRUST4: immune repertoire reconstruction from bulk and single-cell RNA-seq data. Nat Methods. 2021 Jun;18(6):627–30.

28. Team I. immunarch: an R package for painless bioinformatics analysis of T-cell and B-cell immune repertoires. Zenodo. 2019;10:5281.

29. Baazim H, Antonio-Herrera L, Bergthaler A. The interplay of immunology and cachexia in infection and cancer. Nat Rev Immunol. 2022 May;22(5):309–21.

30. Utzschneider DT, Charmoy M, Chennupati V, Pousse L, Ferreira DP, Calderon-Copete S, et al. T Cell Factor 1-Expressing Memory-like CD8(+) T Cells Sustain the Immune Response to Chronic Viral Infections. Immunity. 2016 Aug 16;45(2):415–27.

31. van der Leun AM, Thommen DS, Schumacher TN. CD8+ T cell states in human cancer: insights from single-cell analysis. Nat Rev Cancer. 2020 Apr;20(4):218–32.

32. Naulaerts S, Datsi A, Borras DM, Antoranz Martinez A, Messiaen J, Vanmeerbeek I, et al. Multiomics and spatial mapping characterizes human CD8+ T cell states in cancer. Sci Transl Med. 2023 Apr 12;15(691):eadd1016.

33. Giles JR, Globig AM, Kaech SM, Wherry EJ. CD8+ T cells in the cancer-immunity cycle. Immunity. 2023 Oct 10;56(10):2231–53.

34. Rahim MK, Okholm TLH, Jones KB, McCarthy EE, Liu CC, Yee JL, et al. Dynamic CD8+ T cell responses to cancer immunotherapy in human regional lymph nodes are disrupted in metastatic lymph nodes. Cell. 2023 Mar;186(6):1127-1143.e18.

35. Prokhnevska N, Cardenas MA, Valanparambil RM, Sobierajska E, Barwick BG, Jansen C, et al. CD8+ T cell activation in cancer comprises an initial activation phase in lymph nodes followed by effector differentiation within the tumor. Immunity. 2023 Jan 10;56(1):107-124.e5.

36. Virassamy B, Caramia F, Savas P, Sant S, Wang J, Christo SN, et al. Intratumoral CD8+ T cells with a tissue-resident memory phenotype mediate local immunity and immune checkpoint responses in breast cancer. Cancer Cell. 2023 Mar;41(3):585-601.e8.

37. Wolfl M, Kuball J, Ho WY, Nguyen H, Manley TJ, Bleakley M, et al. Activation-induced expression of CD137 permits detection, isolation, and expansion of the full repertoire of

CD8+ T cells responding to antigen without requiring knowledge of epitope specificities. Blood. 2007 Jul 1;110(1):201–10.

38. Ye Q, Song DG, Poussin M, Yamamoto T, Best A, Li C, et al. CD137 accurately identifies and enriches for naturally occurring tumor-reactive T cells in tumor. Clin Cancer Res. 2014 Jan 1;20(1):44–55.

39. Parkhurst M, Gros A, Pasetto A, Prickett T, Crystal JS, Robbins P, et al. Isolation of T-Cell Receptors Specifically Reactive with Mutated Tumor-Associated Antigens from Tumor-Infiltrating Lymphocytes Based on CD137 Expression. Clin Cancer Res. 2017 May 15;23(10):2491–505.

40. Gros A, Robbins PF, Yao X, Li YF, Turcotte S, Tran E, et al. PD-1 identifies the patient-specific CD8$^+$ tumor-reactive repertoire infiltrating human tumors. J Clin Invest. 2014 May;124(5):2246–59.

41. Thommen DS, Koelzer VH, Herzig P, Roller A, Trefny M, Dimeloe S, et al. A transcriptionally and functionally distinct PD-1+ CD8+ T cell pool with predictive potential in non-small-cell lung cancer treated with PD-1 blockade. Nat Med. 2018 Jul;24(7):994–1004.

42. Duhen T, Duhen R, Montler R, Moses J, Moudgil T, de Miranda NF, et al. Co-expression of CD39 and CD103 identifies tumor-reactive CD8 T cells in human solid tumors. Nat Commun. 2018 Jul 13;9(1):2724.

43. Yeong J, Suteja L, Simoni Y, Lau KW, Tan AC, Li HH, et al. Intratumoral CD39+CD8+ T Cells Predict Response to Programmed Cell Death Protein-1 or Programmed Death Ligand-1 Blockade in Patients With NSCLC. J Thorac Oncol. 2021 Aug;16(8):1349–58.

44. Lee YJ, Kim JY, Jeon SH, Nam H, Jung JH, Jeon M, et al. CD39+ tissue-resident memory CD8+ T cells with a clonal overlap across compartments mediate antitumor immunity in breast cancer. Sci Immunol. 2022 Aug 26;7(74):eabn8390.

45. Palomero J, Panisello C, Lozano-Rabella M, Tirtakasuma R, Díaz-Gómez J, Grases D, et al. Biomarkers of tumor-reactive CD4+ and CD8+ TILs associate with improved prognosis in endometrial cancer. J Immunother Cancer. 2022 Dec;10(12):e005443.

46. Chow A, Uddin FZ, Liu M, Dobrin A, Nabet BY, Mangarin L, et al. The ectonucleotidase CD39 identifies tumor-reactive CD8+ T cells predictive of immune checkpoint blockade efficacy in human lung cancer. Immunity. 2023 Jan 10;56(1):93-106.e6.

47. Buggert M, Price DA, Mackay LK, Betts MR. Human circulating and tissue-resident memory CD8+ T cells. Nat Immunol. 2023 Jul;24(7):1076–86.

48. Sekine T, Perez-Potti A, Nguyen S, Gorin JB, Wu VH, Gostick E, et al. TOX is expressed by exhausted and polyfunctional human effector memory CD8+ T cells. Sci Immunol. 2020 Jul 3;5(49):eaba7918.

49. Bozorgmehr N, Okoye I, Oyegbami O, Xu L, Fontaine A, Cox-Kennett N, et al. Expanded antigen-experienced CD160+CD8+effector T cells exhibit impaired effector functions in chronic lymphocytic leukemia. J Immunother Cancer. 2021 Apr;9(4):e002189.

50. Okoye I, Xu L, Motamedi M, Parashar P, Walker JW, Elahi S. Galectin-9 expression defines exhausted T cells and impaired cytotoxic NK cells in patients with virus-associated solid tumors. J Immunother Cancer. 2020 Dec;8(2):e001849.

51. Baessler A, Vignali DAA. T Cell Exhaustion. Annu Rev Immunol. 2024 Jan 2;

52. Brenchley JM, Karandikar NJ, Betts MR, Ambrozak DR, Hill BJ, Crotty LE, et al. Expression of CD57 defines replicative senescence and antigen-induced apoptotic death of CD8+ T cells. Blood. 2003 Apr 1;101(7):2711–20.

53. Llaó-Cid L, Roessner PM, Chapaprieta V, Öztürk S, Roider T, Bordas M, et al. EOMES is essential for antitumor activity of CD8+ T cells in chronic lymphocytic leukemia. Leukemia. 2021 Nov;35(11):3152–62.

54. Amaria RN, Postow M, Burton EM, Tetzlaff MT, Ross MI, Torres-Cabala C, et al. Neoadjuvant relatlimab and nivolumab in resectable melanoma. Nature. 2022

Nov;611(7934):155–60.

55. Zhang Y, Chen H, Mo H, Hu X, Gao R, Zhao Y, et al. Single-cell analyses reveal key immune cell subsets associated with response to PD-L1 blockade in triple-negative breast cancer. Cancer Cell. 2021 Dec 13;39(12):1578-1593.e8.

56. Oliveira G, Stromhaug K, Klaeger S, Kula T, Frederick DT, Le PM, et al. Phenotype, specificity and avidity of antitumour CD8+ T cells in melanoma. Nature. 2021 Aug;596(7870):119–25.

57. Guo X, Zhang Y, Zheng L, Zheng C, Song J, Zhang Q, et al. Global characterization of T cells in non-small-cell lung cancer by single-cell sequencing. Nat Med. 2018 Jul;24(7):978–85.

58. Caushi JX, Zhang J, Ji Z, Vaghasia A, Zhang B, Hsiue EHC, et al. Transcriptional programs of neoantigen-specific TIL in anti-PD-1-treated lung cancers. Nature. 2021 Aug;596(7870):126–32.

59. Clarke J, Panwar B, Madrigal A, Singh D, Gujar R, Wood O, et al. Single-cell transcriptomic analysis of tissue-resident memory T cells in human lung cancer. J Exp Med. 2019 Sep 2;216(9):2128–49.

60. Rohaan MW, Borch TH, van den Berg JH, Met Ö, Kessels R, Geukes Foppen MH, et al. Tumor-Infiltrating Lymphocyte Therapy or Ipilimumab in Advanced Melanoma. N Engl J Med. 2022 Dec 8;387(23):2113–25.

61. Mellman I, Chen DS, Powles T, Turley SJ. The cancer-immunity cycle: Indication, genotype, and immunotype. Immunity. 2023 Oct 10;56(10):2188–205.

62. Koppensteiner L, Mathieson L, Pattle S, Dorward DA, O'Connor R, Akram AR. Location of CD39+ T cell subpopulations within tumors predict differential outcomes in non-small cell lung cancer. J Immunother Cancer. 2023 Aug;11(8):e006770.

63. Oliveira G, Egloff AM, Afeyan AB, Wolff JO, Zeng Z, Chernock RD, et al. Preexisting tumor-resident T cells with cytotoxic potential associate with response to neoadjuvant anti-PD-1 in head and neck cancer. Sci Immunol. 2023 Sep 8;8(87):eadf4968.

64. Krishna S, Lowery FJ, Copeland AR, Bahadiroglu E, Mukherjee R, Jia L, et al. Stem-like CD8 T cells mediate response of adoptive cell immunotherapy against human cancer. Science. 2020 Dec 11;370(6522):1328–34.

65. Hanada KI, Zhao C, Gil-Hoyos R, Gartner JJ, Chow-Parmer C, Lowery FJ, et al. A phenotypic signature that identifies neoantigen-reactive T cells in fresh human lung cancers. Cancer Cell. 2022 May 9;40(5):479-493.e6.

66. Mestas J, Hughes CCW. Of mice and not men: differences between mouse and human immunology. J Immunol. 2004 Mar 1;172(5):2731–8.

Figure 1: LCL tumor growth kinetics and weight loss in autologous HIS mice and NSG mice. LCL tumors were injected subcutaneously into the flank of mice and tumor size was assessed by calipering. **A**, schematic of generation of HIS mice and autologous tumor cells and subsequent injection in HIS mice. **B**, tumor volume in HIS mice after implantation of the indicated number of LCLs. **C,** Tumor volume in NSG mice after implantation of the indicated number of LCLs. (D, E) Relative weight loss calculated based on the maximum weight during the experiment after tumor cell implantation of the indicated number of LCLs in HIS (**D**) or NSG (**E**) mice. **F**, ratio of spleen weight (mg) to body weight (g), **G**, total splenocyte number and **H**, white blood cell count in HIS mice injected with 5×10^6 autologous LCL tumor cells or non-tumor bearing HIS mice (naïve) sacrificed 15-20 days after tumor implantation. B-E, n=3 per tumor cell number, F-H, n(tumor-bearing)=16, n(naïve)=9 from 4 individual experiments,

marked by individual symbols. Unpaired t-test, data shown as box and whiskers with the box from the 25-75 percentile and the median is shown as a line within the box; whiskers are shown from minimum to maximum data point.

Figure 2: T cell response in tumor-bearing HIS mice. Splenocytes (A-D) and tumor-infiltrating lymphocytes (TIL) (E-H) were isolated from HIS mice 16-18 days after tumor implantation and analyzed by flow cytometry. **A,** frequency of splenic T cells in tumor-bearing

or naïve HIS mice; parent population refers to frequency (%) of $CD3^+$ T cells within human $CD45^+$ cells and $CD4^+$ and $CD8^+$ T cells within $CD3^+$ T cells. **B**, $CD8^+$ T cell differentiation defined as $T_{naïve}$ ($CD45RA^+CD62L^+$), T_{CM} ($CD45RA^-CD62L^+$), T_{EM} ($CD45RA^-CD62L^-$), T_{EMRA} ($CD45RA^+CD62L^-$) in tumor-bearing or naïve HIS mice. **C-D**, expression of indicated markers on $CD8^+$ T cells from spleen of tumor-bearing or naïve HIS mice. **E,** frequency of $CD4^+$ and $CD8^+$ T cells within TILs, gated on total $CD3^+$ T cells. **F**, $CD8^+$ T cell differentiation within TILs. **G-H,** expression of indicated markers on $CD8^+$ T cells within TILs. **I**, UMAP of $CD8^+$ T cells from spleen and TIL of tumor-bearing HIS mice, colored for tissue of origin. **J**, expression of PD-1 and TCF1 in spleen and TIL of tumor-bearing HIS mice and quantification of $CD8^+$ T_{pex} ($TCF1^+PD1^+$), T_{ex-int} ($TCF1^-PD1^+CD69^-$), $T_{ex-term}$ ($TCF1^-PD1^+CD69^+$) in both tissues. **K**, expression of CD69 and CD103 in spleen and tumor of tumor-bearing HIS mice and quantification of T_{RM}-like ($CD69^+CD103^+$) $CD8^+$ T cells in both tissues. **L,** expression of CD137 on splenic $CD8^+$ and $CD4^+$ T cells of tumor-bearing or naïve HIS mice. **M,** expression and quantification of PD-1 and CD137 on splenic $CD8^+$ T cells of tumor-bearing HIS mice. **N**, expression of PD-1 on splenic $CD137^+$ or $CD137^-$ $CD8^+$ T cells in tumor-bearing HIS mice or bulk $CD8^+$ T cells from naïve HIS mice. **O**, expression of individual markers on splenic $CD8^+$ T cell populations from tumor-bearing HIS mice based on expression of CD137 and PD-1. Data are pooled from at least 3 independent experiments, n=10-23 mice per group. For each experiment, a different HPC donor was used for HIS mouse reconstitution and generation of autologous tumor. Significance by paired t-test, one-way ANOVA or 2way ANOVA, as appropriate. Data from individual experiments are indicated by different symbols, with individual mice from the same experiment indicated by the same symbol.

Figure 3: Exhausted-like human CD8⁺ T cells derived from tumor-bearing HIS mice can be expanded *ex vivo* and exhibit superior tumor-specific cytokine production. A, schematic of generation, expansion and characterization of T cells from HIS mice bearing autologous LCL tumors. **B**, fold expansion of FACS sorted splenic CD8⁺ T cells from tumor-bearing HIS mice (CD137⁺, CD137⁻ and CD137⁻PD1⁻) or naïve HIS mice (bulk). **C**, CD8⁺ T cell differentiation after *ex vivo* expansion defined as Tnaïve (CD45RA⁺CD62L⁺), TCM (CD45RA⁻CD62L⁺), TEM (CD45RA⁻CD62L⁻), TEMRA (CD45RA⁺CD62L⁻). **D**, expression of indicated markers after *ex vivo* expansion. **E**, IFN-γ ELISpot of expanded T cells in 5:1 (E:T) co-culture with autologous LCL tumor cells for 24 hours. Spot count is normalized to the spots produced by expanded CD8⁺ bulk T cells from naïve HIS mice. **F**, TNFα ELISA of supernatant of expanded T cells in co-culture (5:1, E:T) with autologous LCL tumor cells for 24 hours. TNFα concentration is normalized to the TNFα secretion from expanded CD8⁺ bulk T cells derived from naïve mice. Data are pooled from at least 3 independent experiments, n=6-19 mice per

group. For each experiment, a different HPC donor was used for HIS mouse reconstitution and generation of autologous tumor. Significance by mixed-effects analysis (paired), RM one-way ANOVA or 2way ANOVA, as appropriate. Data from individual experiments are indicated by different symbols, with individual mice from the same experiment indicated by the same symbol.

Figure 4: Transcriptomic profiling of tumor-reactive CD137⁺ CD8⁺ T cells

Transcriptome analysis and pathway analysis of expanded CD8⁺ T cells from tumor-bearing

HIS mice (CD137⁺, CD137⁻ and CD137⁻PD1⁻) or naïve HIS mice (bulk). **A,** Upset plot

(intersect) showing **number** of differentially expressed genes between groups in bulk RNAseq. p(FDR) < 0.05, log2 FC > 1.5. **B**, PCA plot of RNAseq showing PC1 and PC2. **C,** Top 50 upregulated and **D,** downregulated genes of T cell subsets based on the DEG between CD137$^+$ vs. CD137$^-$PD1$^-$ CD8$^+$ T cells. p(FDR) < 0.05, log2 FC > 1.5. **E,** Volcano plot showing DEG between CD137$^+$ and CD137$^-$PD1$^-$ CD8$^+$ T cells with genes of interest highlighted in yellow. **F**, differential expression of genes of interest between groups. **G**, Overrepresentation analysis (ORA) of upregulated pathways in CD137$^+$ CD8$^+$ T cells. **H**, Gene module score analysis of published signatures described on the right. **I**, Gene set enrichment analysis (GSEA) of signatures described on the y-axis. Gene ratio (# genes related to GO term / total number of sig genes) is displayed on the x-axis. Signatures with an adjusted p-value <0.05 are highlighted with a red box. Shown are data from 4-5 individual experiments, each experiment with different human HPC donor for reconstitution of HIS mice and autologous tumor.

Figure 5: TCR profiling of tumor-reactive CD137⁺ CD8⁺ T cells. A, total number of individual

clonotypes found per population. Left: data from individual experiments, right: pooled data for

group analysis. Wilcoxon test. **B,** TCR (CDR3 of *TRA*, *TRB*, *TRG* and *TRD*) sequence sample

diversity estimation using Hill numbers method, with Q=1 describing the Shannon diversity. **C,**

rare clonal proportion showing the occupied repertoire space by clonotypes with defined

counts (1, 2-3, 4-10, etc.). Left: data from individual experiments, right: pooled data for group

analysis. Wilcoxon test. **D,** relative abundance of clonotypes with defined frequencies (size). Left: data from individual experiments, right: pooled data for group analysis. Wilcoxon test. **E,** repertoire overlap analysis cross-comparing every population from every experiment (each with different donor). **F**, repertoire overlap comparing the repertoire of bulk CD8$^+$ T cells from naïve HIS mice to the populations from tumor-bearing HIS mice from individual experiments (each with different donor; data from individual experiments are indicated by different symbols). Mixed effects analysis with Tukey's multiple comparisons test. **G,** tracking of clonotypes over populations. The top 10 most abundant clonotypes of the TCR repertoire of CD137$^+$ CD8$^+$ T cells from one representative experiment are shown. **H,** proportion of the top 10 most abundant clonotypes (from repertoires of CD137$^+$ CD8$^+$ T cells) in the repertoire of all populations, correlated with the spot count of IFN-γ ELISpot. Shown are data from 4 individual experiments, each experiment with different human HPC donor for reconstitution of HIS mice and autologous tumor.

Figure 6: CD137⁺ CD8⁺ T cells from tumor-bearing HIS mice show superior anticancer activity in NSG and HIS recipients bearing autologous tumors. A, schematic of generation of tumor-reactive T cells and subsequent ACT. NSG mice were injected with 2 x 10⁶ LCL s.c. in the flank and after three days, 10 x 10⁶ *ex vivo* expanded T cells were adoptively transferred intravenously. Transferred T cells and LCL tumors were autologous to each other. **B**, tumor volume on the day of sacrifice in NSG recipient mice after ACT of the indicated cell populations. **C,** waterfall plot of tumor size in NSG recipient mice of ACT on the day of sacrifice relative to the tumor volume of control mice (no ACT). Bars depict individual mice. **D,** proportion of NSG recipient mice showing partial response (≥ 30% tumor reduction compared to control NSG mice) after ACT of the indicated cell population. **E**, frequency of CD3⁺ T cells (% of human CD45⁺ cells) in TIL of NSG mice after ACT of the indicated cell populations, measured by flow cytometry. **F**, frequency of CD8⁺ T cells (% of total cells) in tumors of NSG mice after ACT of the indicated cell populations, measured by immunohistochemistry. **G**, frequency of PD-1 expression on CD8⁺ T cells in TIL of NSG mice after ACT of the indicated cell populations. **H,** correlation between tumor volume and infiltration of CD8⁺ T cells

(measured by IHC) in tumors of NSG mice after adoptive transfer of CD137$^+$ CD8$^+$ T cells. **I**, schematic of ACT. CD137$^+$ CD8$^+$ T cells, CD137$^-$ CD8$^+$ T cells and CD137$^-$ PD-1$^-$ CD8$^+$ T cells were isolated from spleen of tumor-bearing HIS mice or bulk CD8$^+$ T cells from spleen of naïve HIS mice and expanded *ex vivo*. Recipient HIS mice were injected with 2 x 10^6 LCL s.c. in the flank and after three days, *ex vivo* expanded T cells were adoptively transferred intravenously. Tumor-bearing HIS recipient mice received 2 x 10^6 T cells without prior conditioning/lymphodepletion. Donor and recipient HIS mice as well as LCL were autologous to each other. **J,** tumor volume on the day of sacrifice in HIS recipient mice after ACT of the indicated cell populations. **K,** waterfall plot of tumor size on the day of sacrifice of HIS mice receiving ACT relative to the tumor volume of control HIS mice (no ACT). Bars depict individual mice. **L**, proportion of HIS recipient mice showing partial response ($\geq$ 30% tumor reduction compared to control HIS mice) after ACT of the indicated cell population. Data are pooled from 1-3 independent experiments, n=3-13 per group. For each experiment, a different HPC donor was used for HIS mouse reconstitution and generation of autologous tumor. Significance by one-way ANOVA. Data from individual experiments are indicated by different symbols, with individual mice from the same experiment indicated by the same symbol.