Human effector CD8⁺ T cells with an exhausted-like phenotype control tumor growth *in vivo*

Juliane Mietz[1], Meike Kaulfuss[1], Lukas Egli[1], Lennart Opitz[2], Christian Münz[3], Obinna Chijioke[1,4]

[1]Cellular Immunotherapy, Institute of Experimental Immunology, University of Zürich, Zürich, Switzerland
[2]Functional Genomics Center Zürich, University of Zürich/ETH Zürich, Zürich, Switzerland
[3]Viral Immunobiology, Institute of Experimental Immunology, University of Zürich, Zürich, Switzerland
[4]Institute of Medical Genetics and Pathology, University Hospital Basel, Basel, Switzerland

Corresponding author: Obinna Chijioke, Cellular Immunotherapy, Institute of Experimental Immunology, University of Zürich, Winterthurerstrasse 190, 8057 Zürich, Switzerland; Tel: +41 44 635 37 19, Email: chijioke@immunology.uzh.ch

Abstract

The functional relevance of CD8$^+$ T cell phenotypes in human cancer is not well established. In preclinical studies, murine CD8$^+$ T cells exhibiting an exhausted-like phenotype have been described to have drastically decreased anticancer activity while non-exhausted TCF-1$^+$ progenitor CD8$^+$ T cells have been thought to be mainly responsible for driving protective antitumor responses. However, human data has mostly been correlative by nature.

By establishing a novel autologous humanized mouse tumor model, we found a CD8$^+$ dominated T cell expansion upon tumor challenge and identified tumor-reactive CD8$^+$ T cells through expression of CD137. This subpopulation of de novo arising human CD8$^+$ T cells displayed a highly proliferative, fully activated effector and exhausted-like phenotype with enhanced expression of exhaustion markers like PD-1, CD39 and TOX, the senescence marker CD57 (*B3GAT1*) and cytolytic effector molecules such as *PRF1*, *GZMH* and *NKG7*. Moreover, these CD137$^+$CD8$^+$ T cells exhibited tumor-specific clonal expansion and presented signature overlap with tumor-reactive CD8$^+$ T cells described in human cancer. We demonstrate superior anticancer activity of this exhausted-like human CD8$^+$ T cell subset by adoptive transfer experiments using recipients bearing autologous human tumors. Mice adoptively transferred with CD137$^+$CD8$^+$ T cells showed reduced tumor growth, higher CD8$^+$ T cell persistence as well as tumor infiltration, correlating with control of human tumors.

In summary, we defined effective anticancer activity of human effector CD8$^+$ T cells with an exhausted-like phenotype, supporting clinical exploration of such cells in adoptive T cell therapies.

Keywords: Human cancer immunology; Humanized cancer models; Phenotypic markers; T cells; Exhaustion; Adoptive cell therapy

Introduction

The characterization of phenotypic traits that can faithfully identify tumor-specific CD8[+] T cells with functional anticancer activity is an area of intense research. Recent studies have shown tumor-reactive CD8[+] T cells to be rare in patients with various cancers and thus challenging to explore[1,2]. While preclinical studies in mice have shown that CD8[+] T cells displaying an exhausted-like phenotype have decreased effector function[3], studies in individuals with cancer have correlated exhausted-like phenotypic states of CD8[+] T cells with tumor reactivity and clinical benefit[4,5].

Undoubtedly, tumor models using inbred and transgenic mice have contributed substantially to our understanding of tumor immunobiology and have been fundamental to the development of cancer immunotherapies that are now used on a routine basis in the clinic[6,7]. However, human clinical studies[8] have shown that the outcome of cancer immunotherapies can significantly diverge from those of mouse models of cancer[9]. This might be due to a relative lack of genetic diversity of mouse models compared to humans[10] but also to differences in aspects of gene expression networks[11], genomic organization[12] and tissue structure of the immune system of mice and humans[13]. Xenograft mouse models using adoptive transfer of human immune cells rely on immunodeficient hosts, precluding analysis of the impact of endogenous adaptive antitumor immunity. Furthermore, in most preclinical studies investigating novel immunotherapy targets, monoclonal T cells with a single antigen specificity are explored[14,15] while polyclonal T cell responses might be a biomarker for favorable patient outcome to immunotherapy[16] and thus be more informative.

Complementing existing mouse tumor models, the use of mice with reconstituted human immune system components (HIS mice) holds hitherto underexplored translational potential to advance human cancer immunology[17]. Yet, endogenous antitumor immunity in these models has remained largely undefined, mainly due to allogeneic reactivity of the reconstituted human T cells against allogeneic tumors[18], making a distinction of alloreactivity from tumor-specific immune responses challenging.

In this study, we demonstrate that reconstitution of human immune system components across independent donors can mount endogenous antitumor T cell responses to autologous

human tumor challenge in HIS mice. A polyclonal population of human effector CD8[+] T cells

with an exhausted-like phenotype that arises de novo in tumor-bearing HIS mice, resembling

exhausted tumor-reactive lymphocytes found in patients with cancer, is able to control

autologous human tumor growth *in vivo*. Thus, we provide direct evidence of clinically relevant

anticancer activity of expanded human tumor-reactive effector CD8[+] T cells with a senescent

and exhausted-like, non-progenitor-like phenotype.

Results

Establishment of an autologous HIS mouse tumor model

We established and characterized a novel tumor model in which immunocompetent, humanized mice are subcutaneously implanted with autologous human tumor cells. For this aim, we isolated CD34[+] hematopoietic progenitor cells (HPC) and CD19[+] B cells from the same human HPC donor tissue. CD34[+] HPCs were engrafted into NSG mice to reconstitute components of the human immune system (from here on referred to as human immune system (HIS) mice), while CD19[+] B cells were transformed into lymphoblastoid cell lines (LCL) by infection with the oncogenic Epstein-Barr virus (EBV) (**Figure 1A**).

To characterize LCL tumorigenicity and tumor growth in the presence or absence of an autologous human immune system, HIS mice reconstituted with HPCs autologous to the tumor or NSG mice were injected subcutaneously with LCLs. In both recipient strains, palpable tumors formed at the site of implantation. In HIS mice, in the presence of an endogenous immune system, tumors showed progressive growth until approximately day 18 and subsequently, a majority of tumors showed regression (**Figure 1B**). In contrast, in NSG mice without an endogenous immune system, tumors increased in size until termination criteria were reached (**Figure 1C**). These data suggested a functional antitumor response mediated by components of the reconstituted human immune system, i.e. immunocompetence, in our autologous HIS mouse tumor model.

Endogenous antitumor immune response in HIS mice

To investigate the tumor regression that we observed in the presence of an autologous human immune system, we next characterized the immune response in tumor-bearing HIS mice. Around the same time of tumor regression, we found weight loss in HIS mice (**Figure 1D**), but not in tumor-bearing NSG mice (**Figure 1E**). This indicated immunopathology related to the endogenous human antitumor immune response, reminiscent of cancer cachexia[19] rather than an effect of tumor growth per se. Additionally, we found increases in spleen-to-body weight ratio, total splenocyte count and white blood cell count in tumor-bearing HIS mice (**Figure 1, F-H**), suggesting immune cell proliferation occurring in response to tumor growth. Furthermore,

tumors in HIS mice showed infiltration by endogenous human immune cells, including CD8[+] T

cells (Supplemental Figure 1).

The frequency of human CD3[+] T cells in tumor-bearing HIS mice in spleen (**Figure 2A**)

and circulation (Supplemental Figure 2A) increased significantly compared to naïve HIS mice.

Moreover, tumor-bearing HIS mice showed an increased frequency of CD8[+] T cells, suggesting

preferential expansion (**Figure 2A** and Supplemental Figure 2A). These CD8[+] T cells exhibited

differentiation towards an effector memory (T_{EM}) phenotype and, to a lower degree, central

memory (T_{CM}) phenotype, whereas non-tumor bearing HIS mice had higher frequencies of

naïve T cells ($T_{naïve}$) (**Figure 2B** and Supplemental Figure 2B). Consistent with an exhausted

effector-like phenotype, CD8[+] T cells from tumor-bearing HIS mice also showed enhanced

expression of the exhaustion/activation markers PD-1, CD39 and HLA-DR (**Figure 2C** and

Supplemental Figure 2C) as well as trending towards increased TOX expression (**Figure 2D**

and Supplemental Figure 2D), while exhibiting significantly reduced levels of the

progenitor/stemness-related marker TCF-1[20] (**Figure 2D** and Supplemental Figure 2D).

Similar to the increased CD8[+] T cell frequencies in secondary lymphoid tissue (spleen)

and circulation, tumors in HIS mice exhibited predominant infiltration by CD8[+] T cells (**Figure**

2E and Supplemental Figure 1C). The majority of autologous tumor-infiltrating human CD8[+] T

cells had a T_{EM}-like phenotype, with a lower proportion of T_{CM} and greatly reduced proportions

of T_{EMRA} and naïve T cells (**Figure 2F**). Tumor-infiltrating CD8[+] T cells displayed high

expression of exhaustion and activation markers PD-1 (91.0%±12.1), CD39 (56.4%±22.8) and

HLA-DR (53.9%±22.4) (**Figure 2G**), low levels of TCF-1 (5.6%±5.0) (**Figure 2H**), while a

fraction of tumor-infiltrating CD8[+] T cells expressed TOX (21.7%±14.81) (**Figure 2H**). Together,

the phenotype of human CD8[+] T cells in tumor-bearing HIS mice was reminiscent of CD8[+] T

cell phenotypes described in human cancer[21,22].

Exhausted, cytotoxic effector-like phenotype of human CD137[+]CD8[+] T cells

In human cancers, CD137 has been described as a marker for tumor-reactive T cells[23–25]. We

similarly identified a population of human CD137[+] T cells in tumor-bearing mice (**Figure 2I**)

and characterized this CD8[+] T cell population further. Likewise, PD-1 has been reported to be

a marker of tumor-reactive T cells in human cancers, both in circulation and within tumor-infiltrating lymphocytes[26–28]. We observed that of splenic CD8$^+$ T cells from tumor-bearing HIS mice, only a small proportion (10.9%±11.2) were CD137 positive, whereas the majority of CD8$^+$ T cells (83.5%±12.1) expressed PD-1 on their surface (**Figure 2J**). Though there was a small but still significant difference between the overall high level of PD-1 expression on CD137$^+$CD8$^+$ and CD137$^-$CD8$^+$ T cells from tumor-bearing HIS mice, both these populations expressed PD-1 significantly more than bulk CD8$^+$ T cells from naïve HIS mice (**Figure 2K**). These data suggested that CD137 alone might not be sufficient to delineate tumor-reactive T cell subsets. We therefore included a CD8$^+$ T cell subset negative for both CD137 and PD-1 in our analysis to further explore and distinguish bona fide tumor-reactive from bystander CD8$^+$ T cells in tumor-bearing HIS mice.

Indeed, compared to double negative CD137$^-$PD-1$^-$CD8$^+$ T cells, CD137$^+$CD8$^+$ T cells and CD137$^-$CD8$^+$ T cells displayed a T$_{EM}$-like phenotype, while the double negative CD137$^-$PD-1$^-$ population was enriched in naïve-like T cells (**Figure 2L**). Within the splenic CD137$^+$CD8$^+$ T cell population, we found the highest expression of the exhaustion marker CD39, which has been described as a marker of CD8$^+$ T cells with enhanced tumor reactivity across different human cancers[1,29–33], the activation marker HLA-DR as well as the activation and residency-related marker CD69[34], while this cell subset also had the strongest expression levels of Ki-67, indicating high proliferation (**Figure 2M**). Additionally, we observed significantly increased expression of TOX, a transcription factor associated with exhaustion[35] and drastically reduced levels of the progenitor/stemness-related transcriptional regulator TCF-1[20] (**Figure 2N**). Finally, CD137$^+$CD8$^+$ T cells displayed the highest levels of the cytolytic molecules granzyme B and perforin compared with CD137$^-$CD8$^+$ T cells or CD137$^-$PD-1$^-$CD8$^+$ T cells from tumor-bearing HIS mice (**Figure 2O**). Altogether, de novo arising highly proliferative CD137$^+$CD8$^+$ T cells exhibited the most exhausted-like, cytotoxic effector and least stem-like phenotype relative to the other human CD8$^+$ T cell populations we identified in tumor-bearing HIS mice.

Maintained expansion capacity and enrichment of *in vitro* tumor reactivity within an exhausted-like human CD8$^+$ T cell subset

To investigate tumor reactivity of CD137$^+$CD8$^+$ T cells relative to the other T cell subsets,

splenic T cells were expanded *ex vivo* (**Figure 3A**). Using a rapid expansion protocol[28], we

observed an average expansion of >1000-fold after 14 days for all subsets. Interestingly,

although exhausted CD8$^+$ T cells have generally been thought to contain reduced proliferative

potential, there were no significant differences in the expansion capacity of CD137$^+$CD8$^+$ T

cells displaying the most exhausted-like phenotype (Figure 2, M-N) compared to the other

CD8$^+$ T cell populations from tumor-bearing HIS mice or bulk CD8$^+$ T cells from naïve HIS mice

(**Figure 3B**). All of the *ex vivo* expanded T cell subsets from tumor-bearing HIS mice exhibited

differentiation biased to an effector memory-like phenotype (**Figure 3C**).

After co-culture with autologous tumor cells, expanded CD137$^+$CD8$^+$ T cells showed the

highest IFN-γ production by ELISpot, which gradually decreased from CD137$^-$CD8$^+$ T cells to

CD137$^-$PD-1$^-$CD8$^+$ T cells (**Figure 3D**). Similarly, TNFα secretion was highest in supernatants

containing CD137$^+$CD8$^+$ T cells after co-culture with autologous tumor cells and lowest in

CD137$^-$PD-1$^-$CD8$^+$ T cells (**Figure 3E**). These data indicated enrichment of tumor recognition

within CD137$^+$CD8$^+$ T cells that maintained *ex vivo* proliferative potential, while double negative

CD137$^-$PD-1$^-$CD8$^+$ T cells behaved as bystander cells.

Transcriptome analysis reveals tumor-reactive signature in CD137$^+$CD8$^+$ T cells

To gain insight into differentially expressed genes (DEG) in tumor-reactive CD137$^+$CD8$^+$ T cells

versus CD137$^-$CD8$^+$ T cells and bystander CD137$^-$PD-1$^-$CD8$^+$ T cells as well as bulk CD8$^+$ T

cells from naïve HIS mice, whole transcriptome RNA sequencing of *ex vivo* expanded T cell

subsets was performed. We found a higher number of DEG for the tumor-reactive (CD137$^+$)

gene set versus bystander CD8$^+$ T cells from tumor-bearing HIS mice (200 DEG) and versus

bulk CD8$^+$ T cells from naïve HIS mice (300 DEG) in comparison to the number of DEG we

found for the comparison against CD137$^-$CD8$^+$ T cells (53 DEG) (**Figure 4A**), the subset with

the second highest tumor reactivity *in vitro* (Figure 3, D and E). Principal component analysis

(PCA) was consistent with functional data of tumor-specific cytokine production (Figure 3, D

and E), showing that the 4 subsets clustered in a gradual way from tumor-reactive (CD137$^+$)

to less tumor-reactive (CD137$^-$) and bystander (CD137$^-$PD-1$^-$) CD8$^+$ T cells and finally to bulk

CD8[+] T cells from naïve HIS mice, with the biggest overlap between CD137[-]PD-1[-] bystander

CD8[+] T cells and bulk human CD8[+] T cells from naïve HIS mice (**Figure 4B**).

The transcriptomic profile that we found in tumor-reactive CD137[+]CD8[+] T cells suggested

an exhausted/activated-like phenotype with increased expression of *TNFRSF9, LAG3,*

PDCD1, TIGIT and *HLA-DPA1/HLA-DQA1* (**Figure 4, C, E and F**). We also observed

enhanced expression of effector molecules associated with cytotoxicity (*GZMH, NKG7* and

PRF1) as well as *IFNG* (**Figure 4, C, E and F**). Expression of the naïve or stem-like genes

CCR7, *LEF1* and *BACH2* were diminished (**Figure 4, D-F**) while the senescence marker

CD57[36] (*B3GAT1*) and the transcription factor *EOMES* showed the highest transcript level in

the CD137[+] subset (**Figure 4, C and F**). Although EOMES has been shown to be involved in

T cell exhaustion, it has also been reported to be commonly co-expressed with effector

molecules in effector T cells and to be essential for anticancer T cell function[37,38].

Among upregulated Gene Ontology (GO) terms comparing tumor-reactive CD137[+]CD8[+] T

cells against the other three populations, we found the terms interferon-gamma mediated

signaling (GO:0060333), antigen processing and presentation (GO:0019882), antigen

processing and presentation of peptide or polysaccharide antigen via MHC class II

(GO:0002504), immune response (GO:0006955) and adaptive immune response

(GO:0002250) (**Figure 4G**). Upregulation of these pathways in the CD137[+] population

indicated stronger T cell activation and response in comparison to the other populations.

Next, we compared our transcriptome data to published datasets of different T cell states

from individuals with cancers[21,33,39–41]. Strikingly, comparing tumor-reactive CD137[+]CD8[+] T

cells to bystander CD137[-]PD-1[-]CD8[+] T cells or bulk CD8[+] T cells from naïve HIS mice, gene

sets that have been described to define tumor-reactive T cells in patients with cancer were

significantly upregulated ('MANA TIL'[40], 'tumor specific'[41]) (**Figure 4H**). These results

suggested that the antitumor T cell response in autologous tumor-bearing HIS mice mirrors

relevant aspects of the T cell response in cancer patients, thus further supporting the potential

translational value of our model.

Correlation of clonal expansion with effector function

As another surrogate for tumor recognition, we analyzed clonality of the CD8$^+$ T cell receptor (TCR) repertoire, anticipating tumor-reactive CD8$^+$ T cells would have higher clonal expansion and thus lower diversity than bystander CD8$^+$ T cells. Indeed, we found the total number of clonotypes tending to be the smallest in the tumor-reactive CD137$^+$ subset, and to gradually increase from the less tumor-reactive CD137$^-$ subset to CD137$^-$PD-1$^-$ bystander CD8$^+$ T cells to bulk CD8$^+$ T cells from naïve HIS mice (**Figure 5A**), following the gradient of tumor recognition we observed by tumor-specific cytokine production *in vitro* (Figure 3, D and E). Likewise, the estimated sample diversity for the tumor-reactive CD137$^+$ subset was lower than for the other populations (**Figure 5B** and Supplemental Figure 3A). In line with this, the CD137$^+$ population had a reduced abundance of rare or very small clonotypes (**Figure 5C** and Supplemental Figure 3, B and C). Conversely, tumor-reactive CD137$^+$ CD8$^+$ T cells from tumor-bearing HIS mice demonstrated a significantly higher abundance of hyperexpanded clones (**Figure 5D**) with the 20 biggest clones occupying a significantly larger repertoire space than in the other populations, suggesting polyclonal expansion (Supplemental Figure 3B).

To track the overlap of clonotypes between individual experiments (i.e. donor dependency) and populations, we next performed TCR repertoire overlap analysis (**Figure 5E**). We found that the tumor-reactive CD137$^+$ population showed the biggest overlap with the CD137$^-$ subset and only to a minor extend an overlap with CD137$^-$PD-1$^-$ bystander CD8$^+$ T cells or bulk CD8$^+$ T cells from naïve HIS mice (**Figure 5, E and G** and Supplemental Figure 3E). Also, analyzing the repertoire of the bulk naïve CD8$^+$ T cell population for each experiment by cross-comparing to each population from each experiment, the overlap with the CD137$^+$ population was significantly less than the overlap with every other population (**Fig. 5F**). These findings suggested that clonotypes arising due to the conditions of expansion were less present in the CD137$^+$ population and that other, presumably tumor-reactive, clonotypes expanded.

In addition, we observed that the expansion of individual bona fide tumor-reactive clonotypes was highly donor specific. When comparing the 10 most abundant clonotypes of the CD137$^+$ population per experiment (i.e. per donor), almost no overlap with the CD137$^+$ TCR repertoire of any other experiment was detected (**Figure 5G** and Supplemental Figure 3D). Finally, to explore an association of clonal expansion with functionality, we correlated the

abundance of the top 10 bona fide tumor-reactive clonotypes of the CD137$^+$ population with *in*

vitro tumor reactivity and found a highly significant correlation of clone size abundance and

tumor-specific IFN-γ production by CD8$^+$ T cells (**Figure 5H**). Overall, these data indicated that

CD137$^+$CD8$^+$ T cells derived from tumor-bearing HIS mice expanded polyclonally in a human

donor- and tumor-specific manner.

Human CD8$^+$ T cells with an exhausted-like phenotype reduce tumor growth *in vivo*

We next performed adoptive cell transfer (ACT) experiments of *ex vivo* expanded CD8$^+$ T cells

in NSG mice previously engrafted with autologous tumors (**Figure 6A**). We initially selected

NSG mice as recipients to unambiguously track the transferred T cells and to be able to

distinguish their anticancer activity from the endogenous antitumor T cell response in HIS mice.

In NSG mice, only ACT with expanded CD137$^+$CD8$^+$ T cells resulted in reduced tumor growth,

while transfer of bystander CD8$^+$ T cells from tumor-bearing HIS mice or bulk CD8$^+$ T cells from

naïve HIS mice did not (**Figure 6, B and C**).

Furthermore, a higher frequency of T cells was found in tumor (**Figure 6, D and E**), spleen

(Supplemental Figure 4A) and blood (Supplemental Figure 4B) of tumor-bearing NSG mice

receiving CD137$^+$CD8$^+$ T cells compared to recipients transferred with any of the other cell

populations. Flow cytometric analysis of tumor-infiltrating lymphocytes (TIL) revealed high

CD3$^+$ T cell frequencies (53.65%±21.94) after ACT of expanded CD137$^+$CD8$^+$ T cells, but only

11.64%±11.0 after transfer of bystander CD8$^+$ T cells and 7.25%±5.38 after transfer of bulk

CD8$^+$ T cells from naïve HIS mice (**Figure 6D**). Likewise, immunohistochemistry of tumors of

NSG mice confirmed T cell infiltration selectively after ACT with CD137$^+$CD8$^+$ T cells (**Figure**

6E). Thus, human exhausted-like CD137$^+$CD8$^+$ T cells exhibited the potential for persistence

and robust tumor infiltration.

Interestingly, while infusion products mainly contained a mix of T$_{EM}$-like and T$_{EMRA}$ CD8$^+$ T

cells (Figure 3C), almost exclusively T$_{EM}$-like cells were recovered from TIL (**Figure 6F**), spleen

(Supplemental Figure 4C) and blood (Supplemental Figure 4D) of recipient NSG mice. Of note,

whereas PD-1 was highly expressed on splenic CD137$^+$CD8$^+$ T cells in tumor-bearing HIS

mice (96.49%±3.1) (Figure 2K), PD-1 expression on this population decreased during antigen-

independent *ex vivo* expansion (31.81%±31.81) (Supplemental Figure 5A). However, after

adoptive transfer of expanded CD137[+]CD8[+] T cells into tumor-bearing NSG mice, PD-1

expression among CD8[+] T cells in TIL increased again (96.7%±3.65) and was significantly

higher compared with CD8[+] T cells recovered from TIL of NSG mice transferred with bystander

CD8[+] T cells or bulk CD8[+] T cells (Supplemental Figure 5B). Apart from the superior tumor

control, these data further indicated enrichment of tumor reactivity within CD137[+]CD8[+] T cells,

with very high PD-1 expression serving as a potential marker for productive tumor recognition,

but not loss of effector function. As another index of the anticancer activity of CD137[+]CD8[+] T

cells, the degree of tumor infiltration with CD8[+] T cells after ACT was inversely correlated with

tumor size (**Figure 6G**). Thus, the higher frequency of CD8[+] T cells in tumors was likely driven

by increased tumor recognition of transferred CD137[+]CD8[+] T cells, and these cells mediated

tumor cell killing.

Lastly, ACT of tumor-reactive T cells into autologous tumor-bearing HIS recipient mice

(**Figure 6H**) presented an opportunity to model polyclonal TIL transfer therapy in cancer

patients[42] in an immunocompetent humanized tumor setting. As observed with NSG recipients,

transfer of polyclonal tumor-reactive CD137[+]CD8[+] T cells into autologous tumor-bearing HIS

recipient mice reduced tumor growth (**Figure 6, I and J**). Importantly, this was achieved without

prior conditioning of HIS recipient mice or cytokine support, and the number of transferred

tumor-reactive CD137[+]CD8[+] T cells was fivefold lower compared with transfer in NSG

recipients. These data established superior anticancer activity of CD137[+]CD8[+] T cells with an

exhausted-like phenotype the most complete humanized setting.

Altogether, these adoptive transfer experiments confirmed anticancer activity of human

CD137[+]CD8[+] T cells that developed in autologous tumor-bearing HIS mice, suggesting that

our model captured relevant aspects of the immunobiology of human antitumor T cell

responses.

Discussion

Using in-depth phenotypic analyses, *in vitro* functional assays and adoptive transfer experiments, we identified a proliferative human CD8[+] T cell subset in tumor-bearing HIS mice that expressed senescence and exhaustion markers including very high levels of PD-1 and CD39 while having low expression of the progenitor marker TCF-1 but exhibited increased tumor reactivity. By means of adoptive transfer, we uniquely provide direct *in vivo* evidence of superior anticancer activity of these human tumor-reactive effector CD8[+] T cells displaying a senescent and exhausted-like phenotypic state.

We have leveraged this human CD8[+] T cell subset arising de novo in tumor-bearing HIS mice for transcriptomic analysis and TCR profiling to demonstrate across donors overlap with tumor-reactive CD8[+] T cell signatures from human cancers[40,41] as well as tumor-induced polyclonal expansion. Polyclonality of neoantigen-specific CD8[+] T cells in blood and tumor has recently been shown to be associated with response to PD-1 immunotherapy in patients with melanoma[16]. Congruence of our model data across donors with these clinical benchmarks suggests robustness and translational validity of our findings.

Of note, CD39[+]CD8[+] T cells expressing high levels of PD-1 have been described to be enriched for tumor reactivity and clonal proliferation, predicting efficacy of immunotherapy in human lung cancer[33]. Likewise, the abundance of intratumoral CD39[+]CD8[+] T cells displaying upregulated exhaustion markers has been shown to be prognostic for improved progression-free and overall survival in individuals with treatment-naïve early stage lung cancer[43]. Furthermore, PD-1[hi]CD39[+]CD8[+] T cells correlated with improved survival in patients with endometrial cancer[32] and the signature of similar CD8[+] T cells with increased TOX expression predicted survival in patients with breast cancer[31]. Recently, pre-existing exhausted-like tumor-reactive CD8[+] T cells have been associated with favorable outcome to immunotherapy in head and neck cancer[44]. On the other hand, higher abundance of TCF-1[+] stem-like CD8[+] T cells with a CD39[-]CD69[-] phenotype in TIL products was associated with better response after TIL transfer in patients with melanoma[45]. However, by single-cell analysis of lung tumors, this phenotype could not be identified in neoantigen-reactive T cells[46]. While we have not addressed and thus cannot exclude a contribution of the minor TCF-1[+] fraction within the

human tumor-reactive CD8[+] T cells in our model, our data emphasize the key importance of

exhausted-like effector CD8[+] T cells in the antitumor immune response.

The striking phenotypic similarity of tumor-reactive CD8[+] T cells that have been shown to

correlate with clinical benefit and the polyclonally expanded human tumor-reactive CD8[+] T cell

cells from tumor-bearing HIS mice that we demonstrated to possess superior *in vivo* anticancer

activity highlights the translational potential of our model. Hence, the novel immunocompetent

human tumor model that we developed using HIS mice with human immune cells reconstituted

from hematopoietic progenitor cells that are autologous to tumor might bridge a gap between

syngeneic mouse tumor models and human-specific antitumor immune responses[13,47]. Our

model demonstrates hallmarks of human antitumor T cell responses and thus may provide a

tool that should allow further investigation of drivers of anticancer activity in human CD8[+] T

cells. Overall, the endogenous human immune system in HIS mice markedly influenced tumor

and host, allowing for experimental interrogation of the interfaces of human tumor and

autologous human antitumor immunity in a small animal model.

Together, fully activated effector and exhausted-like human CD8[+] T cells directly mediate

anticancer activity. Our data support clinical exploration of exhausted-like effector CD8[+] T cell

subsets alongside the recently more favoured stem-like phenotype for use in adoptive transfer

and engineered T cell therapies.

Methods

HIS & NSG mice

NSG (NOD.Cg-Prkdc<scid>Il2rg<tm1Wjl>/SzJ (#005557)) mice were purchased from The Jackson Laboratory and bred and housed under specific pathogen-free conditions at the Laboratory Animal Services Center (LASC) Zurich, University of Zurich. To generate HIS mice, newborn pups (1-5 days old) were irradiated (1 Gy) and intrahepatically injected with 0.2×10^6 human CD34$^+$ hematopoietic progenitor cells derived from human fetal liver (Advanced Bioscience Resources, USA). CD34$^+$ cells were isolated by MACS technology (Miltenyi, Cat. 130-046-703) following the manufacturer's instructions. At 3 months of age, reconstitution of human immune cells was assessed in PBMC from tail vein blood by flow cytometry[48]. Reconstitution analysis included staining for human CD45, CD3, CD4, CD8, CD19, NKp46 and HLA-DR. Only mice with a sufficiently high frequency of human CD45 (>25% of leukocytes) were included in the experiments. All animal experiments were performed according to approved licenses by the veterinary office of the canton of Zurich, Switzerland (ZH049/20 and ZH067/2023).

LCL generation and tumor model

LCL were generated from CD19$^+$ B cells derived from the same autologous human fetal liver tissue from which the CD34$^+$ cells for reconstitution of the human immune system in the same experiment were derived. CD19$^+$ cells were isolated from the HFL tissue's CD34$^-$ fraction by MACS technology (Miltenyi, Cat. 130-050-301) according to the manufacturer's instructions. 0.25×10^6 CD19$^+$ human B cells were transformed by infection with Epstein-Barr virus (EBV B95-8; produced as previously described[48]) at an MOI of 0.1 - 0.15 and cultured in RPMI/ 10% FCS/ 1% Penicillin/ Streptomycin.

LCL tumors were injected subcutaneously into the left flank under isoflurane narcosis. For generation and phenotyping of tumor-reactive T cells, 5×10^6 autologous LCL tumor cells were resuspended in PBS and right before injection mixed in a 1:1 V/V ratio with Corning® Matrigel® Growth Factor Reduced (GFR) Basement Membrane Matrix (Milian, Cat. 354230).

For adoptive cell transfer experiments, 2 x 10^6 LCL were injected s.c. in a 1:1 V/V mix with

Matrigel and three days after tumor injection, 2 x 10^6 T cells (HIS recipient mice) or 10 x 10^6

(NSG recipient mice) were adoptively transferred by tail vein injection. Tumor size was

monitored by calipering (3x/week or daily). Tumor volume was calculated as volume [mm^3] =

length [mm] x width^2 [mm^2] x 0.52. General health was monitored by weighing and health

parameter scoring (3x/week or daily, according to animal license). PBMC composition and

expansion of adoptively transferred T cells were monitored by weekly tail vein bleeding and

flow cytometric analysis. White blood cell (WBC) counts were measured from blood with an

automatic cell counting machine (DxH 500, Beckman Coulter). Animals were euthanized when

they met pre-defined termination criteria defined in the animal license.

T cell isolation and expansion

Spleens from LCL tumor-bearing and naïve HIS mice were harvested 16-18 days after s.c.

tumor injection. Splenocytes were labelled with fluorescently labelled antibodies and sorted by

fluorescent activated cell sorting (FACS). CD137$^+$, CD137$^-$ and CD137$^-$PD-1$^-$ CD8$^+$ T cells were

sorted from tumor-bearing HIS mice and bulk CD8$^+$ T cells were isolated from naïve HIS mice.

Sorted T cells were expanded by a rapid expansion protocol (REP), as described elsewhere[28].

REP medium consisted of a 1:1 mix of RPMI (Gibco, Cat. 7001612) and X-Vivo (Lonza, Cat.

BE02-060F) containing 10% FCS (Biochrome, Cat. S0615-500ML) and 1% Penicillin/

Streptomycin (Thermo Fisher, Cat. 7001592). After sorting, T cells were cultured in REP

medium with 30 ng/µl OKT3 (Miltenyi, Cat. 130-093-387), 3000 IU/ml IL-2 (Peprotech, Cat.

200-02) and a 200-fold excess of irradiated (50 Gy) allogeneic PBMC derived from 3 healthy

donors at a total cell density of 5 x 10^6 cells/ml. Starting at day 7, cells were adjusted with REP

medium to 0.5-1 x 10^6 cells/ml and fresh IL-2 (3000 IU/ml) was added every second day. Cells

were expanded for a total of 14-16 days until use for further experiments.

Cell isolation from organs

For reconstitution analysis and weekly bleeding, 100-150 µl of tail vein blood was collected

into EDTA tubes (BD Microtainer, BD, Cat. 365975) and erythrocytes were lysed by incubation

with lysis buffer. Leukocytes were stained for flow cytometric analysis. For PBMC isolation on

the day of sacrifice, blood was harvested by heart puncture and erythrocytes were lysed with

erythrocyte lysis buffer. For splenocyte isolation, spleens were removed and meshed through

a 70 μm cell strainer in PBS and lymphocytes were isolated by Ficoll-Paque (GE Healthcare,

17-5442-03). Single cell suspensions were washed and stained for flow cytometric analysis.

For tumor-infiltrating lymphocyte (TIL) isolation, tumors were removed from the flank and skin

was removed from the tumors. Tumors were cut into small pieces with scissors and then

incubated with digestion mix (DMEM (Life technologies, Cat. 7001566) containing 2% FCS

(Biochrome, Cat. S0615-500ML), 1 mg/ml Collagenase IV (Roche, Cat. 7002219), 10ug/ml

DNase I (Roche, Cat. 7002221) and 1.2 mM CaCl2; approx. 1.5 ml/tumor, incubation for 45

min, 37°C). The tumor digest was diluted with 20 ml RPMI or PBS and meshed over a 70 μm

cell strainer. Single cell suspensions were washed and then stained for flow cytometric

analysis.

Flow cytometry

For surface staining, cells were washed in PBS and stained with an antibody master mix for

20 min at 4°C in the dark. For intracellular staining, cells were first stained for surface antigens

and fixable live/dead marker, then fixed and permeabilized using the BD Pharmingen™

Transcription Factor Buffer Set (BD, Cat. 562574) according to the manufacturer's instructions,

then stained for intracellular or intranuclear antigens for 50-60 minutes. Cells were washed in

PBS and acquired on a BD LSRFortessa™.

Antibodies for reconstitution analysis of HIS mice (all antibodies are anti-human): CD45-Pacific

Blue (Biolegend, HI30, Cat. 304029), CD3-PE (Biolegend, UCHT1, Cat. 300408), CD4-APC-

Cy7 (Biolegend, RPA-T4, Cat. 300518), CD8-PerCP (Biolegend, SK1, Cat. 344708), HLA-DR-

FITC (Biolegend, L243, Cat. 307604), CD19-PE-Cy7 (Biolegend, HIB19, Cat. 302216),

NKp46-APC (BD, 9-E2, Cat. 558051), Zombie Aqua™ Fixable Viability Kit (Biolegend, Cat.

423101).

Antibodies for flow cytometry-based sorting of T cells: CD3-FITC (Biolegend, UCHT1, Cat.

300406), CD4-APC-Cy7 (Biolegend, RPA-T4, Cat. 300518), CD8-BV650 (Biolegend, SK1,

Cat. 344730), CD45-Pacific Blue (Biolegend, HI30, Cat. 304029), CD137-APC (Miltenyi,

REA765, Cat. 130-110-901), PD-1-PE-Dazzle (Biolegend, EH12.2H7, Cat. 329940), Zombie

Aqua™ Fixable Viability Kit (Biolegend, Cat. 423101).

Antibodies for phenotyping in Panel 1: CD45-Pacific Blue (Biolegend, HI30, Cat. 304029),

CD3-PerCP-Cy5.5 (Biolegend, UCHT1, Cat. 300429), CD4-APC-Cy7 (Biolegend, RPA-T4,

Cat. 300518), CD8-BV650 (Biolegend, SK1, Cat. 344730), CD45RA-BV785 (Biolegend,

HI100, 304139), CD62L-FITC (BD, DREG-56, Cat. 555543), CD19-PE-Cy7 (Biolegend,

HIB19, Cat. 302216), HLA-DR-BV605 (BD, g46-6, Cat. 562845), CD39-BV711 (Biolegend, A1,

Cat. 328228), CD69-Alexa Fluor 700 (Biolegend, FN50, Cat. 310921), PD-1-PE-Dazzle

(Biolegend, EH12.2H7, Cat. 329940), TCF-1-PE (Biolegend, 7F11A10, Cat. 655207), CD137-

APC (Miltenyi, REA765, Cat. 130-110-901), Zombie Aqua™ Fixable Viability Kit (Biolegend,

Cat. 423101); in Panel 2: CD3-FITC (Biolegend, UCHT1, Cat. 300406), CD4-PE-Cy7

(Biolegend, RPA-T4, Cat. 300512), CD8-BV650 (Biolegend, SK1, Cat. 344730), CD45-BV711

(Biolegend, HI30, Cat. 304050), CD137-APC-Fire750 (Biolegend, 4B4-1, Cat. 309834),

Granzyme B-APC-Alexa700 (Biolegend, QA16A02, Cat. 372222), Ki-67-BV605 (Biolegend,

Ki-67, Cat. 350522), PD-1-PE-Dazzle (Biolegend, EH12.2H7, Cat. 329940), Perforin – BV421

(Biolegend, dG9, Cat. 308122), TCF-1-PE (Biolegend, 7F11A10, Cat. 655207), TOX-APC

(Miltenyi, REA473, Cat. 130-118-335), Zombie Aqua™ Fixable Viability Kit (Biolegend, Cat.

423101).

IFN-γ ELISpot and TNFα ELISA

For IFN-γ ELISpot, ELISpot plates (Merck, Cat.MAIPN4550) were activated with 35% ethanol,

washed and coated overnight with anti-IFN-γ coating antibody (mAb-1-D1K, Mabtech, Cat.

3420-3-1000). Plates were washed and T cells were co-cultured with autologous LCL in RPMI/

10% FCS/ 1% Pen/Strep at an effector: target ratio of 5:1 for 16-24 hours. After incubation,

supernatant was harvested, stored at -20°C and used for TNFα ELISA. Wells were washed

and incubated for 2 hours with biotinylated anti-IFN-γ detection antibody (mAb-7-B6-1-Biotin,

Mabtech, Cat. 3420-6-250). Afterwards, wells were washed and incubated with Streptavidin-

ALP (Mabtech, Cat. 3310-10) for 1 hour. Wells were then washed again and spots were

developed with filtered substrate solution BCIP/ NBT-plus (Mabtech, Cat. 3650-10). When

distinct spots were visible, the plate was washed and dried overnight. Plates were acquired

with an ELISpot reader (AID classic, AID GmbH) and spots were automatically counted using

AID ELISpot 8.0 software (AID GmbH).

TNFα ELISA was performed with the supernatant from T cell and LCL co-culture as described

above using the human TNFα ELISA[BASIC] kit (Mabtech, Cat. 3512-1H-6), following

manufacturer's instructions. Plates were acquired using an ELISA reader (infinite M200Pro,

Tecan) and analyzed with the i-control 2.0 software (Tecan).

Immunohistochemistry

Tumors were fixed in 4% paraformaldehyde and embedded in paraffin. Sample embedding,

preparation and staining was performed by the Pathology department of the University Hospital

Basel. Slides were acquired on an automated slide scanning brightfield microscope (Vectra 3,

Akoya Biosciences) and quantified using InForm software (Akoya Biosciences).

Bulk RNAseq and TCR profiling

RNA was extracted from 5 x 10^6 T cells using the RNeasy® Mini Kit (Qiagen, Cat. 74106). RNA

extraction was performed following the instructions from the "Purification of Total RNA from

Animal Cells Using Spin Technology" protocol given in the RNeasy® Mini Handbook, including

homogenization with QIAshredders (Qiagen, Cat. 79654) and on-column DNA digestion with

the RNase-free DNase Set (Qiagen, Cat. 79254). All further steps were performed by the

Functional Genomics Center Zurich (FGCZ). The quality of the isolated RNA was determined

with a Qubit® (1.0) Fluorometer (Life Technologies, California, USA) and a Bioanalyzer 2100

(Agilent, Waldbronn, Germany). Only those samples with a 260 nm/280 nm ratio between 1.8–

2.1 and a 28S/18S ratio within 1.5–2 were further processed. The TruSeq RNA Sample Prep

Kit v2 (Illumina, Inc, California, USA) was used in the succeeding steps. Briefly, total RNA

samples (600 ng) were poly A enriched and then reverse-transcribed into double-stranded

cDNA. The cDNA samples were fragmented, end-repaired and polyadenylated before ligation

of TruSeq adapters containing the index for multiplexing fragments containing TruSeq adapters

on both ends were selectively enriched with PCR. The quality and quantity of the enriched

libraries were validated using Qubit® (1.0) Fluorometer and the Caliper GX LabChip® GX

(Caliper Life Sciences, Inc., USA). The product is a smear with an average fragment size of

approximately 260 bp. The libraries were normalized to 10nM in Tris-Cl 10 mM, pH8.5 with

0.1% Tween 20. After library quantification, libraries were prepared for loading accordingly to

the NovaSeq workflow with the NovaSeq6000 Reagent Kit (Illumina, Catalog

No. 20012865). Cluster generation and sequencing were performed on a NovaSeq6000

System with a run configuration of single end 100bp. RNA-seq data analysis consisted of the

following steps: The raw reads were first cleaned by removing adapter sequences, trimming

low quality ends, and filtering reads with low quality (phred quality <20) using Fastp (Version

0.20)[49]. Sequence pseudo alignment of the resulting high-quality reads to the human reference

genome (build GRCh38.p13 and GENCODE gene models release 37) and quantification of

gene level expression was carried out using Kallisto (Version 0.46.1)[50]. Differential expression

was computed using the generalized linear model implemented in the Bioconductor package

DESeq2 (R version: 4.2.2, DESeq2 version: 1.38.1)[51]. Genes showing altered expression with

adjusted (Benjamini and Hochberg method) p-value < 0.05 were considered differentially

expressed. For the TCR analysis the read alignment to the Human genome was performed

with STAR (v2.7.10b)[52]. TCR sequences (CDR3 of TRA, TRB, TRG and TRD) were

reconstructed from bulk RNAseq data using TRUST4 (v1.0.10)[53]. Out-of-frame sequences

were excluded from the analysis. TCR repertoire analysis was performed using the R package

Immunarch[54].

Statistics

Statistical analysis was performed using Prism 10 Software (GraphPad Software, Inc). Figure

legends indicate which statistical test was performed.

Study approval

Animal experiments were conducted according to licenses approved by the veterinary office of

the canton of Zurich, Switzerland (ZH049/20 and ZH067/2023).

BioRender

Schematics were generated with a licensed version of BioRender.com.

Author contributions

J.M. and O.C. conceived, designed, and evaluated the experiments. J.M. performed the analysis. M.K. and L.E. assisted with the in vivo experiments. L.O. provided bioinformatic analysis of transcriptomic data. O.C. supervised and C.M. cosupervised this work. J.M. and O.C. wrote the manuscript. All authors approved the manuscript.

Declaration of Interests

The authors declare that they have no competing interests.

Acknowledgments

This research was supported by Swiss Cancer Research foundation (KFS-5292-02-2021 to O.C.).

Data Sharing

Supplemental data are provided with this paper. All data are also available from the corresponding author upon reasonable request.

References

1. Simoni, Y. *et al.* Bystander CD8+ T cells are abundant and phenotypically distinct in human tumour infiltrates. *Nature* **557**, 575–579 (2018).

2. Scheper, W. *et al.* Low and variable tumor reactivity of the intratumoral TCR repertoire in human cancers. *Nat Med* **25**, 89–94 (2019).

3. Lan, X., Zebley, C. C. & Youngblood, B. Cellular and molecular waypoints along the path of T cell exhaustion. *Sci Immunol* **8**, eadg3868 (2023).

4. Lowery, F. J. *et al.* Molecular signatures of antitumor neoantigen-reactive T cells from metastatic human cancers. *Science* **375**, 877–884 (2022).

5. Oliveira, G. & Wu, C. J. Dynamics and specificities of T cells in cancer immunotherapy. *Nat Rev Cancer* **23**, 295–316 (2023).

6. Korman, A. J., Garrett-Thomson, S. C. & Lonberg, N. The foundations of immune checkpoint blockade and the ipilimumab approval decennial. *Nat Rev Drug Discov* **21**, 509–528 (2022).

7. Ribas, A. & Wolchok, J. D. Cancer immunotherapy using checkpoint blockade. *Science* **359**, 1350–1355 (2018).

8. Powles, T. *et al.* Atezolizumab with enzalutamide versus enzalutamide alone in metastatic castration-resistant prostate cancer: a randomized phase 3 trial. *Nat Med* **28**, 144–153 (2022).

9. Guan, X. *et al.* Androgen receptor activity in T cells limits checkpoint blockade efficacy. *Nature* **606**, 791–796 (2022).

10. Palmer, A. C., Izar, B., Hwangbo, H. & Sorger, P. K. Predictable Clinical Benefits without Evidence of Synergy in Trials of Combination Therapies with Immune-Checkpoint Inhibitors. *Clin Cancer Res* **28**, 368–377 (2022).

11. Zong, W. *et al.* Transcriptomic congruence analysis for evaluating model organisms. *Proc Natl Acad Sci U S A* **120**, e2202584120 (2023).

12. Gilbertson, S. E. *et al.* Topologically associating domains are disrupted by evolutionary genome rearrangements forming species-specific enhancer connections in mice and humans. *Cell Rep* **39**, 110769 (2022).

13. Medetgul-Ernar, K. & Davis, M. M. Standing on the shoulders of mice. *Immunity* **55**, 1343–1353 (2022).

14. Dong, M. B. *et al.* Systematic Immunotherapy Target Discovery Using Genome-Scale In Vivo CRISPR Screens in CD8 T Cells. *Cell* **178**, 1189-1204.e23 (2019).

15. Chen, Z. *et al.* In vivo CD8+ T cell CRISPR screening reveals control by Fli1 in infection and cancer. *Cell* **184**, 1262-1280.e22 (2021).

16. Puig-Saus, C. *et al.* Neoantigen-targeted CD8+ T cell responses with PD-1 blockade therapy. *Nature* **615**, 697–704 (2023).

17. De La Rochere, P. *et al.* Humanized Mice for the Study of Immuno-Oncology. *Trends Immunol* **39**, 748–763 (2018).

18. Hiwarkar, P. *et al.* Cord blood T cells mediate enhanced antitumor effects compared with adult peripheral blood T cells. *Blood* **126**, 2882–2891 (2015).

19. Baazim, H., Antonio-Herrera, L. & Bergthaler, A. The interplay of immunology and cachexia in infection and cancer. *Nat Rev Immunol* **22**, 309–321 (2022).

20. Utzschneider, D. T. *et al.* T Cell Factor 1-Expressing Memory-like CD8(+) T Cells Sustain the Immune Response to Chronic Viral Infections. *Immunity* **45**, 415–427 (2016).

21. van der Leun, A. M., Thommen, D. S. & Schumacher, T. N. CD8+ T cell states in human cancer: insights from single-cell analysis. *Nat Rev Cancer* **20**, 218–232 (2020).

22. Naulaerts, S. *et al.* Multiomics and spatial mapping characterizes human CD8+ T cell states in cancer. *Sci Transl Med* **15**, eadd1016 (2023).

23. Wolfl, M. *et al.* Activation-induced expression of CD137 permits detection, isolation, and expansion of the full repertoire of CD8+ T cells responding to antigen without requiring knowledge of epitope specificities. *Blood* **110**, 201–210 (2007).

24. Ye, Q. *et al.* CD137 accurately identifies and enriches for naturally occurring tumor-reactive T cells in tumor. *Clin Cancer Res* **20**, 44–55 (2014).

25. Parkhurst, M. *et al.* Isolation of T-Cell Receptors Specifically Reactive with Mutated Tumor-Associated Antigens from Tumor-Infiltrating Lymphocytes Based on CD137 Expression. *Clin Cancer Res* **23**, 2491–2505 (2017).

26. Gros, A. *et al.* PD-1 identifies the patient-specific CD8[+] tumor-reactive repertoire infiltrating human tumors. *J Clin Invest* **124**, 2246–2259 (2014).

27. Thommen, D. S. *et al.* A transcriptionally and functionally distinct PD-1+ CD8+ T cell pool with predictive potential in non-small-cell lung cancer treated with PD-1 blockade. *Nat Med* **24**, 994–1004 (2018).

28. Gros, A. *et al.* Recognition of human gastrointestinal cancer neoantigens by circulating PD-1+ lymphocytes. *J Clin Invest* **129**, 4992–5004 (2019).

29. Duhen, T. *et al.* Co-expression of CD39 and CD103 identifies tumor-reactive CD8 T cells in human solid tumors. *Nat Commun* **9**, 2724 (2018).

30. Yeong, J. *et al.* Intratumoral CD39+CD8+ T Cells Predict Response to Programmed Cell Death Protein-1 or Programmed Death Ligand-1 Blockade in Patients With NSCLC. *J Thorac Oncol* **16**, 1349–1358 (2021).

31. Lee, Y. J. *et al.* CD39+ tissue-resident memory CD8+ T cells with a clonal overlap across compartments mediate antitumor immunity in breast cancer. *Sci Immunol* **7**, eabn8390 (2022).

32. Palomero, J. *et al.* Biomarkers of tumor-reactive CD4+ and CD8+ TILs associate with improved prognosis in endometrial cancer. *J Immunother Cancer* **10**, e005443 (2022).

33. Chow, A. *et al.* The ectonucleotidase CD39 identifies tumor-reactive CD8+ T cells predictive of immune checkpoint blockade efficacy in human lung cancer. *Immunity* **56**, 93-106.e6 (2023).

34. Buggert, M., Price, D. A., Mackay, L. K. & Betts, M. R. Human circulating and tissue-resident memory CD8+ T cells. *Nat Immunol* **24**, 1076–1086 (2023).

35. Sekine, T. *et al.* TOX is expressed by exhausted and polyfunctional human effector memory CD8+ T cells. *Sci Immunol* **5**, eaba7918 (2020).

36. Brenchley, J. M. *et al.* Expression of CD57 defines replicative senescence and antigen-induced apoptotic death of CD8+ T cells. *Blood* **101**, 2711–2720 (2003).

37. Llaó-Cid, L. *et al.* EOMES is essential for antitumor activity of CD8+ T cells in chronic lymphocytic leukemia. *Leukemia* **35**, 3152–3162 (2021).

38. Amaria, R. N. *et al.* Neoadjuvant relatlimab and nivolumab in resectable melanoma. *Nature* **611**, 155–160 (2022).

39. Guo, X. *et al.* Global characterization of T cells in non-small-cell lung cancer by single-cell sequencing. *Nat Med* **24**, 978–985 (2018).

40. Caushi, J. X. *et al.* Transcriptional programs of neoantigen-specific TIL in anti-PD-1-treated lung cancers. *Nature* **596**, 126–132 (2021).

41. Oliveira, G. *et al.* Phenotype, specificity and avidity of antitumour CD8+ T cells in melanoma. *Nature* **596**, 119–125 (2021).

42. Rohaan, M. W. *et al.* Tumor-Infiltrating Lymphocyte Therapy or Ipilimumab in Advanced Melanoma. *N Engl J Med* **387**, 2113–2125 (2022).

43. Koppensteiner, L. *et al.* Location of CD39+ T cell subpopulations within tumors predict differential outcomes in non-small cell lung cancer. *J Immunother Cancer* **11**, e006770 (2023).

44. Oliveira, G. *et al.* Preexisting tumor-resident T cells with cytotoxic potential associate with response to neoadjuvant anti-PD-1 in head and neck cancer. *Sci Immunol* **8**, eadf4968 (2023).

45. Krishna, S. *et al.* Stem-like CD8 T cells mediate response of adoptive cell immunotherapy against human cancer. *Science* **370**, 1328–1334 (2020).

46. Hanada, K.-I. *et al.* A phenotypic signature that identifies neoantigen-reactive T cells in fresh human lung cancers. *Cancer Cell* **40**, 479-493.e6 (2022).

47. Mestas, J. & Hughes, C. C. W. Of mice and not men: differences between mouse and human immunology. *J Immunol* **172**, 2731–2738 (2004).

48. Chijioke, O. *et al.* Human natural killer cells prevent infectious mononucleosis features by targeting lytic Epstein-Barr virus infection. *Cell Rep* **5**, 1489–1498 (2013).

49. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

50. Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol* **34**, 525–527 (2016).

51. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

52. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

53. Song, L. *et al.* TRUST4: immune repertoire reconstruction from bulk and single-cell RNA-seq data. *Nat Methods* **18**, 627–630 (2021).

54. Team, I. immunarch: an R package for painless bioinformatics analysis of T-cell and B-cell immune repertoires. *Zenodo* **10**, 5281 (2019).

Figure 1: LCL tumor growth kinetics and weight loss in autologous HIS mice and NSG

mice. LCL tumors were injected subcutaneously into the flank of mice and tumor size was

assessed by calipering. **A**, schematic of generation of HIS mice and autologous tumor cells

and subsequent injection in HIS mice. **B**, tumor volume in HIS mice after implantation of the

indicated number of LCLs. **C,** Tumor volume in NSG mice after implantation of the indicated

number of LCLs. (D, E) Relative weight loss calculated based on the maximum weight during

the experiment after tumor cell implantation of the indicated number of LCLs in HIS (**D**) or NSG

(**E**) mice. **F**, ratio of spleen weight (mg) to body weight (g), **G**, total splenocyte number and **H**,

white blood cell count in HIS mice injected with 5 x 10^6 autologous LCL tumor cells or non-

589   tumor bearing HIS mice (naïve) sacrificed 15-20 days after tumor implantation. B-E, n=3 per

590   tumor cell number, F-H, n(tumor-bearing)=16, n(naïve)=9 from 4 individual experiments,

591   marked by individual symbols. Unpaired t-test, data shown as box and whiskers with the box

592   from the 25-75 percentile and the median is shown as a line within the box; whiskers are shown

593   from minimum to maximum data point.

Figure 2: T cell response in tumor-bearing HIS mice. Splenocytes (A-D) and tumor-infiltrating lymphocytes (TIL) (E-H) were isolated from HIS mice 16-18 days after tumor implantation and analyzed by flow cytometry. **A,** frequency of splenic T cells in tumor-bearing or naïve HIS mice; parent population refers to frequency (%) of CD3+ T cells within human CD45+ cells and CD4+ and CD8+ T cells within CD3+ T cells. **B**, CD8+ T cell differentiation defined as $T_{naïve}$ (CD45RA+CD62L+), T_{CM} (CD45RA-CD62L+), T_{EM} (CD45RA-CD62L-), T_{EMRA} (CD45RA+CD62L-) in tumor-bearing or naïve HIS mice. **C-D**, expression of indicated markers on CD8+ T cells from spleen of tumor-bearing or naïve HIS mice. **E,** frequency of CD4+ and

CD8$^+$ T cells within TILs, gated on total CD3$^+$ T cells. **F,** CD8$^+$ T cell differentiation within TILs.

G-H, expression of indicated markers on CD8$^+$ T cells within TILs. **I,** expression of CD137 on

splenic CD8$^+$ and CD4$^+$ T cells of tumor-bearing or naïve HIS mice. **J,** expression of PD-1 and

CD137 on splenic CD8$^+$ T cells of tumor-bearing mice. **K**, expression of PD-1 on splenic CD8$^+$

T cells based on CD137 in tumor-bearing HIS mice or bulk CD8$^+$ T cells from naïve HIS mice.

L, CD8$^+$ T cell differentiation in spleen of tumor-bearing HIS mice. **M-O**, expression of indicated

markers on splenocyte-derived CD8$^+$ T cell populations based on expression of CD137 and

PD-1. Data are pooled from at least 3 independent experiments, n=10-23 mice per group. For

each experiment, a different HPC donor was used for HIS mouse reconstitution and generation

of autologous tumor. Significance by paired t-test, one-way ANOVA or 2way ANOVA, as

appropriate. Data from individual experiments are indicated by different symbols.

Figure 3: Exhausted-like human CD8+ T cells derived from tumor-bearing HIS mice can be expanded *ex vivo* and exhibit superior tumor-specific cytokine production. A, schematic of generation, expansion and characterization of T cells from HIS mice bearing autologous LCL tumors. **B**, fold expansion of FACS sorted splenic CD8+ T cells from tumor-bearing HIS mice (CD137+, CD137- and CD137-PD1-) or naïve HIS mice (bulk). **C**, CD8+ T cell differentiation after *ex vivo* expansion defined as $T_{naïve}$ (CD45RA+CD62L+), T_{CM} (CD45RA-CD62L+), T_{EM} (CD45RA-CD62L-), T_{EMRA} (CD45RA+CD62L-). **D**, IFN-γ ELISpot of expanded T cells in 5:1 (E:T) co-culture with autologous LCL tumor cells for 24 hours. Spot count is normalized to the spots produced by expanded CD8+ bulk T cells from naïve HIS mice. **E**, TNFα ELISA of supernatant of expanded T cells in co-culture (5:1, E:T) with autologous LCL tumor cells for 24 hours. TNFα concentration is normalized to the TNFα secretion from expanded CD8+ bulk T cells derived from naïve mice. Data are pooled from at least 3 independent experiments, n=6-19 mice per group. For each experiment, a different HPC donor was used for HIS mouse reconstitution and generation of autologous tumor. Significance by mixed-effects analysis (paired), RM one-way ANOVA or 2way ANOVA, as appropriate. Data from individual experiments are indicated by different symbols.

Figure 4: Transcriptomic profiling of tumor-reactive CD137$^+$CD8$^+$ T cells

Transcriptome analysis and pathway analysis of expanded CD8$^+$ T cells from tumor-bearing HIS mice (CD137$^+$, CD137$^-$ and CD137$^-$PD1$^-$) or naïve HIS mice (bulk). **A,** Upset plot (intersect) showing number of differentially expressed genes between groups in bulk RNAseq. p(FDR) < 0.05, log2 FC > 1.5. **B**, PCA plot of RNAseq showing PC1 and PC2. **C,** Top 50 upregulated and **D,** downregulated genes of T cell subsets based on the DEG between CD137$^+$ vs. CD137$^-$ PD1$^-$ CD8$^+$ T cells. p(FDR) < 0.05, log2 FC > 1.5. **E,** Volcano plot showing DEG between CD137$^+$ and CD137$^-$PD1$^-$ CD8$^+$ T cells with genes of interest highlighted in yellow. **F**, differential expression of genes of interest between groups. **G**, Overrepresentation analysis (ORA) of upregulated pathways in CD137$^+$CD8$^+$ T cells. **H**, Gene set enrichment analysis (GSEA) of signatures described on the y-axis. Gene ratio (# genes related to GO term / total number of sig genes) is displayed on the x-axis. Signatures with an adjusted p-value <0.05 are highlighted with a red box. Shown are data from 5 individual experiments, each experiment with different human HPC donor for reconstitution of HIS mice and autologous tumor.

Figure 5: TCR profiling of tumor-reactive CD137⁺CD8⁺ T cells. A, total number of individual

clonotypes found per population. Left: data from individual experiments, right: pooled data for

group analysis. Wilcoxon test. **B,** TCR (CDR3 of *TRA*, *TRB*, *TRG* and *TRD*) sequence sample

diversity estimation using Hill numbers method, with Q=1 describing the Shannon diversity. **C,**

rare clonal proportion showing the occupied repertoire space by clonotypes with defined

counts (1, 2-3, 4-10, etc.). Left: data from individual experiments, right: pooled data for group

analysis. Wilcoxon test. **D,** relative abundance of clonotypes with defined frequencies (size).

Left: data from individual experiments, right: pooled data for group analysis. Wilcoxon test. **E,**

repertoire overlap analysis cross-comparing every population from every experiment (each

with different donor). **F**, repertoire overlap comparing the repertoire of bulk CD8$^+$ T cells from

naïve HIS mice to the populations from tumor-bearing HIS mice from individual experiments

(each with different donor; data from individual experiments are indicated by different symbols).

Mixed effects analysis with Tukey's multiple comparisons test. **G**, tracking of clonotypes over

populations. The top 10 most abundant clonotypes of the TCR repertoire of CD137$^+$CD8$^+$ T

cells from one representative experiment are shown. **H**, proportion of the top 10 most abundant

clonotypes (from repertoires of CD137$^+$CD8$^+$ T cells) in the repertoire of all populations,

correlated with the spot count of IFN-γ ELISpot. Shown are data from 4 individual experiments,

each experiment with different human HPC donor for reconstitution of HIS mice and autologous

tumor.

Figure 6: CD137⁺CD8⁺ T cells from tumor-bearing HIS mice show superior anticancer activity in NSG and HIS recipients bearing autologous tumors. A, schematic of generation of tumor-reactive T cells and subsequent ACT. NSG mice were injected with 2 x 10⁶ LCL s.c. in the flank and after three days, 10 x 10⁶ *ex vivo* expanded T cells were adoptively transferred intravenously. Transferred T cells and LCL tumors were autologous to each other. **B,** tumor volume on the day of sacrifice in NSG recipient mice after ACT of the indicated cell populations. **C,** waterfall plot of tumor size in NSG recipient mice of ACT on the day of sacrifice relative to the tumor volume of control mice (no ACT). Bars depict individual mice. **D,** frequency of CD3⁺ T cells (% of human CD45⁺ cells) in TIL from NSG mice after ACT of the indicated cell populations, measured by flow cytometry. **E,** frequency of CD8⁺ T cells (% of total cells) in tumors of NSG mice after ACT of the indicated cell populations, measured by immunohistochemistry. **F**, differentiation of CD8⁺ T cells in TIL of NSG mice after ACT of the indicated cell populations defined as T$_{naïve}$ (CD45RA⁺CD62L⁺), T$_{CM}$ (CD45RA⁻CD62L⁺), T$_{EM}$ (CD45RA⁻CD62L⁻) and T$_{EMRA}$ (CD45RA⁺CD62L⁻). **G,** correlation between tumor volume and infiltration of CD8⁺ T cells (measured by IHC) in tumors of NSG mice after adoptive transfer of CD137⁺CD8⁺ T cells. **H**, schematic of ACT. CD137⁺CD8⁺ T cells, CD137⁻CD8⁺ T cells and

CD137⁻ PD-1⁻CD8⁺ T cells were isolated from spleen of tumor-bearing HIS mice or bulk CD8⁺

T cells from spleen of naïve HIS mice and expanded *ex vivo*. Recipient HIS mice were injected

with 2 x 10^6 LCL s.c. in the flank and after three days, *ex vivo* expanded T cells were adoptively

transferred intravenously. Tumor-bearing HIS recipient mice received 2 x 10^6 T cells without

prior conditioning/lymphodepletion. Donor and recipient HIS mice as well as LCL were

autologous to each other. **I,** tumor volume on the day of sacrifice in HIS recipient mice after

ACT of the indicated cell populations. **J,** waterfall plot of tumor size on the day of sacrifice of

HIS mice receiving ACT relative to the tumor volume of control HIS mice (no ACT). Bars depict

individual mice. B-G: Data are pooled from 2-3 independent experiments, n=6-13 per group. I,

data are pooled from 2-3 independent experiments (n=6-13 per group); J, data are from 1-3

independent experiments (n=3-13 per group). For each experiment, a with different HPC donor

was used for HIS mouse reconstitution and generation of autologous tumor. Significance by

one-way ANOVA. Data from individual experiments are indicated by different symbols.