Disrupted endosomal trafficking of the Vangl-Celsr polarity complex underlies congenital anomalies in trachea-esophageal morphogenesis

Nicole A. Edwards[1*], Adhish Kashyap[1], Alissa Warren[1], Zachary N. Agricola[1,2], Alan P. Kenny[2], Yufeng Shen[4,5], Wendy K. Chung[6,7], Aaron M. Zorn[1,3,8*]

[1]Center for Stem Cell and Organoid Medicine (CuSTOM), Perinatal Institute, Division of Developmental Biology, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, 45208 USA

[2]Perinatal Institute, Division of Neonatology, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, 45208 USA

[3]University of Cincinnati, College of Medicine, Cincinnati, OH, USA

[4]Department of Systems Biology, Columbia University Irving Medical Center, New York, NY, 10032 USA

[5]Department of Biomedical Informatics, Columbia University Irving Medical Center, New York, NY, 10032 USA

[6]Department of Pediatrics, Boston Children's Hospital, Harvard Medical School, Boston, MA, 02115 USA

[7]Departments of Pediatrics and Medicine, Columbia University Irving Medical Center, New York, NY

[8]Lead contact

*Correspondence: aaron.zorn@cchmc.org, nicole.edwards@cchmc.org

Summary

Disruptions in foregut morphogenesis can result in life-threatening conditions where the trachea

and esophagus fail to separate properly, such as esophageal atresia (EA) and

tracheoesophageal fistulas (TEF). The developmental basis of these congenital anomalies is

poorly understood, but recent genome sequencing reveals that *de novo* variants in intracellular

trafficking genes are enriched in EA/TEF patients. Here we show that mutation of orthologous

genes in *Xenopus* disrupts trachea-esophageal separation similar to EA/TEF patients. We show

that the Rab11a recycling endosome pathway is required to localize Vangl-Celsr polarity

complexes at the cell surface where opposite sides of the common foregut tube fuse. Partial

loss of endosome trafficking or the Vangl/Celsr complex disrupts epithelial polarity and cell

division orientation. Mutant cells accumulate at the fusion point, fail to downregulate cadherin,

and do not separate into distinct trachea and esophagus. These data provide new insights into

the mechanisms of congenital anomalies and general paradigms of tissue fusion during

organogenesis.

39
Introduction

Esophageal atresia (EA) and tracheoesophageal fistulas (TEF), which affect approximately 1 in

3000 newborns, are life-threatening congenital anomalies that arise when the embryonic foregut

does not separate correctly into distinct trachea and esophageal tubes[1-3]. This results in

esophageal discontinuity (EA) and/or a pathological connection between the trachea and

esophagus (TEF). Rarer are laryngeal-tracheal-esophageal-clefts where in some extreme cases

the foregut fails to separate along its entire length, essentially resulting in an extended complete

TEF[4]. Roughly half of EA/TEF patients also have variably penetrant co-occurring anomalies in

other organs including gastrointestinal, pulmonary, cardiovascular, renal and the central

nervous system, suggesting undefined syndromes [5,6]. The genetic etiology of

tracheoesophageal anomalies is poorly understood and is associated with spontaneous *de novo*

variants, chromosomal rearrangements, copy number variation, indels, and single nucleotide

changes in affected newborns[1,7]. Known variants only account for 10-15% of cases world-wide

with the remaining having no identified risk gene. Even when the causative variants are known,

why they result in tracheoesophageal anomalies is unclear since the cellular mechanisms

governing foregut morphogenesis are poorly understood.

Tracheoesophageal morphogenesis begins when the foregut endoderm is patterned along the

dorsal-ventral axis by an evolutionarily conserved Hedgehog-Wnt-Bmp signaling network

resulting in dorsal Sox2+ esophageal progenitors and ventral Nkx2-1+ respiratory progenitors[8-

11]. After patterning, the foregut constricts at the midline and a population of epithelial cells on

either side of the foregut tube that express both dorsal and ventral markers (Sox2/Nkx2-1/Islet1)

come together and adhere forming a transient epithelial bilayer[12,13]. Epithelial cells at the fusion

point then progressively lose adhesion to one another and integrate into either the ventral or

dorsal foregut as the surrounding Foxf1+ splanchnic mesoderm invades between the nascent

trachea and esophagus[13]. Thus, the single foregut epithelial tube fuses, reorganizes its cellular

junctions at the luminal/apical surface, then undergoes fission with coordinated cell

rearrangements to reconnect the epithelia into two separate tubes.

How cell signaling pathways and transcription factors regulate these cellular processes is poorly

understood. Similar epithelial fusion events occur in other tissues including the heart, oral

palate, eyes, and neural tube; organ systems that are frequently disrupted in complex EA/TEF

patients, suggesting common biological mechanisms[14]. Studies in these other organs indicate

that dynamic changes in tissue and cell polarity are necessary for proper morphogenesis [14,15].

For example, vertebrate neural tube closure and morphogenesis of the *Drosophila* retina both

require the spatial-temporal coordination of apical-basal and planar cell polarity (PCP) proteins,

but how polarity and junctional proteins are localized and coordinated to regulate epithelial

fusion and remodeling is unclear [16-19].

We recently found in *Xenopus* that downstream of Hedgehog signaling, Rab11+ recycling

endosomes somehow regulate tracheoesophageal morphogenesis [13]. Rab11 is a key small

GTPase in the recycling endosome pathway that cells use to transport newly synthesized

proteins from the Golgi to plasma membrane and to change the composition of specific plasma

membrane domains by internalizing receptors and other cell surface proteins [20]. How recycling

endosomes regulate foregut epithelial remodeling was unclear, but intriguingly, whole genome

sequencing analysis revealed that EA/TEF patients with complex multi-organ phenotypes are

enriched for potentially damaging *de novo* variants in genes related to endocytosis and

intracellular trafficking [21]. These two independent observations led us to test the hypothesis that

endosome trafficking is a crucial morphogenetic bottleneck in tracheoesophageal separation

and to define the underlying mechanisms.

Here we demonstrate that the Rab11a recycling endosome pathway is required upstream of the

Vangl-Celsr PCP complex for proper fission of the foregut tube. Mechanistically, Rab11a

localizes the transmembrane protein Vangl2 specifically to the midline luminal cell surface

where opposite sides of the foregut touch, upregulate cadherin, and adhere to one another to

form the transient epithelial bilayer. Rab11a and Vangl-Celsr complexes enriched at the contact

surface appear to confer an "apical memory" essential to maintain the transient bilayer and its

subsequent resolution. A partial loss of 1) core endosome trafficking components, 2) orthologs

of EA/TEF patient risk genes, or 3) the Vangl-Celsr proteins disrupts cell polarity and the

orientation of cell division in the transient bilayer. Mutant cells accumulate in a disorganized

septum, fail to downregulate cadherin, and do not properly separate into distinct trachea and

esophagus. This unexpected Vangl/Celsr activity is distinct from their well-known Wnt/PCP role

in orienting cells within the plane of the epithelium and is an unusual twist on their typical

function in defining apical-basal epithelial polarity. Together these results indicate that failure to

maintain localized endosome-mediated epithelial polarity during foregut morphogenesis may be

a common disease mechanism underlying EA/TEF in many patients and may have broad

implications for tissue fusion events in the development of other organs.

Results

Variants in endosome pathway genes are enriched in EA/TEF patients and cause TEF in

Xenopus

Genome sequencing recently identified a cohort of complex EA/TEF patients with potentially

gene damaging and missense *de novo* heterozygous variants in 55 genes involved in

intracellular trafficking and endocytosis [21]. A meta-analysis of experimentally validated protein-

protein interactions from the STRING database [22] revealed that many of the proteins encoded

by patient risk genes interact with core recycling endosome components (Figure 1A), supporting

recent evidence that this pathway has an important role in tracheoesophageal morphogenesis

[13]. To determine if these patient variants were likely to be causative risk alleles, we mutated

orthologous genes in *Xenopus tropicalis* using F0 CRISPR-Cas9 gene editing, which

recapitulates the *de novo* variants in newborn patients. We designed CRISPR guide RNAs

(gRNAs) to induce indel mutations early in the coding sequence to disrupt protein function, and

injected Cas9-gRNA complexes into one cell-stage embryos. After of three days of

development, embryos were genotyped, and tracheoesophageal morphogenesis was assayed

by immunostaining and confocal microscopy (Figure 1B). If the embryos had EA/TEF-like

phenotypes, then the patient variant is likely a risk variant.

Figure 1. Mutating EA/TEF patient endosome trafficking genes causes TEDs in *Xenopus*.
A) STRING Database interactome of endosome-related proteins with potentially pathogenic
variants identified in EA/TEF patients together with a curated list of core endosome pathway
proteins and putative protein cargo. B) Experimental design of F0 CRISPR-Cas9 *X. tropicalis*
mutagenesis screen to validate candidate risk genes. C) Confocal images of CRISPR-Cas9 *X.*
tropicalis F0 mutants at NF44 stained for Sox2 (green), Foxf1 (red), and Nkx2-1 (purple). Hashed
yellow lines indicate the trachea (t) and esophageal (e) lumens and arrows indicate trachea-
esophageal fistulas. Asterisks indicate presence of esophageal dysmorphology. Numbers
indicate the number of CRISPR mutant tadpoles with TEF compared to the total mutants
screened. Scale bars are 50 µm.

We found that mutations in 10/19 orthologs of endosome-related patient risk genes (*abra,*
ap1g2, arhgap21, atp6v1b1, celsr2, itgb4, itsn1, lamb4, rapgef3 and *rasal2*) when mutated in
Xenopus result in variably penetrant tracheoesophageal anomalies in *Xenopus* (Figure 1C and
Table S1). This was also true for F0 mutations in the known EA/TEF risk gene *sox2* and
mutations in genes encoding key GTPases in endosome trafficking: *dynamin-2* (*dnm2),* which
regulates vesicle formation at the cell surface, *rab5a*, which regulates early endosome
trafficking, and *rab11a* involved in recycling endosomes that transports vesicles to the cell
surface (Table S1). Genotyping of individual embryos revealed a 30-90% damaging indel rate,
suggesting a significant loss of protein but not complete null mutants (Table S1 and Figure S1).
In each case, the foregut was correctly patterned into a ventral Nkx2-1+ trachea domain and a
dorsal Sox2+ esophageal domain with surrounding Foxf1+ mesenchyme, and the foregut
underwent initial constriction at the midline with the walls of the foregut epithelia coming
together. However, in many embryos the foregut failed to separate into distinct tubes and
exhibited a persistent fistula (Figure 1C, arrows). Some patient genes (*rapgef3, arhgap21,*
rasal2, ap1g2, atp6v1b1) when mutated also exhibited a dysmorphic esophagus ranging from
stenosis to a thick multilayered epithelium with multiple or occluded esophageal lumens (Figure
1C, asterisks), consistent with EA in patients. These results demonstrate that endosome
trafficking is required for tracheoesophageal morphogenesis in *Xenopus* and support the
likelihood that variants in endosome-related genes may have caused EA/TEF in some patients.
157
Dynamic endosome localization and epithelial remodeling during foregut morphogenesis
To better understand the cellular mechanisms by which endosome trafficking regulates
Xenopus foregut morphogenesis, we performed a detailed immunostaining analysis over the 24-
hour period when the single foregut separates into distinct esophageal and tracheal tubes
(Figure 2). At the onset of tracheoesophageal morphogenesis (stage NF37), the endodermal
foregut tube is a typical apical-basal polarized pseudostratified columnar epithelium. Cdh1 (E-
cadherin) is enriched in adherens junctions and along the basal-lateral cell borders, and tight
junctions and apical polarity cell proteins like aPKC and Par3 are localized to the luminal/apical
cell surface, whilst laminin lines the basal surface contacting the surrounding mesoderm (Figure
2A,E and Figure S2A-C). As the foregut constricts in the midline (stage NF37) and the opposing
epithelia touch, apical proteins aPKC and Par3 are downregulated and Tjp1+ tight junctions are
lost (Figure 2F,J and Figure S2D-G). This is coincident with an upregulation of Cdh1 at the
170
171

Figure 2. Dynamic endosome localization and epithelial remodeling during tracheoesophageal morphogenesis. A-J) Immunostaining of aPKC, Cdh1, Dnm2, Rab5a, and Rab11a in the NF37 foregut during epithelial fusion. K-T) Immunostaining of the NF41 foregut before and during resolution of the epithelial bilayer. E, J, O, T) Diagrams depicting the temporal-spatial dynamics of Cdh1 and Rab11a subcellular localization during tracheoesophageal morphogenesis. U) Quantification of Cdh1 immunostaining intensity in the membrane before and after fusion in the foregut (mean ± min/max, 2W-ANOVA, ****p<0.0001, n=4-6 embryos analyzed). V) Quantification of Rab11a immunostaining intensity before and after foregut fusion (mean ± min/max, 2W-ANOVA, ****p<0.0001, n=4-6 embryos analyzed). W) Quantification of Cdh1 signal intensity in the bilayer before and during resolution (mean ± min/max, Student's t-test, **p<0.01, n=4-6 embryos analyzed). X) Quantification of Rab11a signal intensity in the bilayer before and during resolution (mean± min/max, Student's t-test, **p<0.01, n=4-6 embryos analyzed). Y) A model of how endosome trafficking may mediate Cdh1 re-localization during epithelial fusion.

contact point (Figure 2F,U), which presumably helps mediate adhesion of the two sides and the

formation of a well-organized, but transient, columnar epithelial bilayer by NF39 (Figure 2K, O,

Figure S2J). After 12 hours, at NF41-42, columnar cells in the middle of the epithelial bilayer

downregulate Cdh1, round up, lose attachment to their neighbors, and appear to integrate into

either the trachea or esophagus as mesenchyme cells invade to complete separation of the

foregut (Figure 2P,T,W, Figure S2K-L,O).

Co-staining for the endosome pathway GTPases Dnm2, Rab5a, and Rab11a revealed intriguing

patterns of subcellular localization during this epithelial remodeling (Figure 2). Throughout the

process of foregut fusion, all epithelial cells (dorsal, ventral, and midline) have Dnm2 enriched at

the apical-lateral membrane and subapical Rab5a+ early endosome vesicles (Figure 2B-C,G,-

H, Figure S2H-I). In contrast, the recycling endosome Rab11a only becomes robustly enriched

at the luminal contact point where the two sides of the foregut epithelia touched (Figure

2D,I,J,V). Many Rab11a+ puncta at the contact point co-stain for Cdh1 with robust co-

localization persisting at the adhesion surface between the two epithelial in the bilayer

consistent with previous reports[13]. At NF41-42, when the epithelial bilayer progressively

resolves, Rab11a and Cdh1 levels significantly drop as the columnar cells round up and lose

attachment to their neighbors (Figure 2N,S,X). At this time Dnm2 is also observed in the

mesenchyme surrounding septum, while Rab5a marks submembrane vesicular structures

(Figure 2L-M,Q-R and Figure S2M-N). Collectively these analyses reveal dynamic localized

endosomal trafficking during the epithelial remodeling that separates the trachea and

esophagus (Figure 2Y).

Endosome trafficking is required for maintenance and resolution of a transient epithelial

bilayer during tracheoesophageal morphogenesis

We next determined which of the epithelial remodeling events driving tracheoesophageal

morphogenesis require endosome trafficking. A time course immunostaining of *dnm2*, *rab5a*,

and *rab11a* mutant embryos showed that initial foregut constriction and epithelial fusion with

downregulation of aPKC and upregulation of Cdh1 at the contact surface occurred normally

(Figure 3A, Figure S3B-C). However, mutant embryos did not form or maintain an organized

columnar epithelia bilayer between the presumptive trachea and esophagus. Rather, the mutant

septa were disorganized and 4-6 cells wide with cells in the middle being round rather than

columnar (Figure 3B,D). Moreover, at NF41 the mutant septum did not downregulate Cdh1 and

Figure 3. Disrupting endosome trafficking causes TEF in *Xenopus*. A) At NF37 the septum in *dnm2*, *rab5a*, and *rab11a* CRISPR mutants undergoes epithelial fusion and down regulate apical polarity proteins similarly to controls. B) At NF41 the epithelial bilayer in endosome mutants is disorganized compared to controls. C) *dnm2*, *rab5a*, and *rab11a* CRISPR mutants exhibit TEF.

Scale bars are 50 μm. D) The mutant epithelia septa have significantly more cells across the width
of the tissue compared to controls (mean ± SEM, * p<0.05 1W-ANOVA, ** p < 0.01, N=6-11
embryos). E) There are no significant (n. s.) differences in proliferation (pHH3+/total) or F) total
cells in the septum of mutant embryos compared to controls (mean ± min/max, 1W-ANOVA, n=4-
6 embryos). G) 3D surface renderings and measurements of the epithelial septum showing that
mutants are significantly wider (x axis) and shorter in the rostral-caudal (z axis) axis compared to
controls (mean length ± SEM, Student's t-test compared to control lengths, *p<0.05, **p<0.01,
n=4-6 embryos per condition). Scale bars are 30 μm.
264
265 resolve into distinct trachea and esophageal tubes, but remained a disorganized epithelial
266 mass, with frequent blisters, resulting in a blind ended fistula (Figure 3C, Figure S3D, Table S1).
267
268 A proportion of endosome mutants also exhibited a dysmorphic esophagus and occluded
269 tracheal lumens (Figure 3B, Table S1). These results indicate that endosome trafficking is
270 required to maintain an organized epithelial bilayer, and suggest that the accumulation of too
271 many cells with persistent Cdh1 expression disrupts the resolution into distinct trachea and
272 esophageal tubes.
273
To determine if the accumulation of cells was due to elevated cell proliferation, we quantified
confocal Z-stacks of the entire foregut region and found that neither the total number of cells in
the trachea-esophageal septum nor the rate of proliferation (phosphorylated histone H3
staining) was significantly increased in endosomal mutants compared to controls (Figure 3E-F).
Three dimensional reconstructions demonstrated that the mutant septa were wider (x-axis) and
shorter in the rostral-caudal axis (z-axis) compared to controls but were similar in the dorsal-
ventral axis (y-axis) (Figure 3G). This indicates that mutant cells somehow accumulate across
the width of the epithelia septum at the expense of fewer cells along the rostral-caudal
extension of the separating trachea and esophagus.
283
Endosome trafficking controls the orientation of division in the foregut epithelial bilayer
Since polarized epithelial cells are known to divide into the plane of the tissue to maintain a
single layer organization, we considered the possibility that in the mutant embryos, cells might
aberrantly divide into the middle of the septum contributing to the cell build-up (Figure 4A). To
assess the plane of cell division, we measured the angle of the mitotic spindle relative to the
basal membrane in the septum of endosomal mutants and controls. While cells in the epithelial
bilayer of control embryos divided into the plane of the epithelia as expected, with an average
spindle orientation of 0-15°, *dnm2*, *rab5a,* and *rab11a* mutants exhibited a more random
292

Figure 4. Endosome trafficking controls polarity of epithelial cell division. A) A potential
model of altered cell division orientation in endosome mutants causes cells to accumulate in the
epithelial septum. B) Immunostaining of the mitotic spindle (Tuba4a and phospho-H3 histone)
was using to measure the angle of cell division. C) Distribution of mitotic spindle angles is
random in endosome mutant epithelial cells compared to controls, which consistently divide
between 0-15° (n=3-15 spindles per embryo, Fisher's exact test, *p<0.05, **p<0.01, N=4-6
embryos. D-G) Endosome mutants have ectopic Laminin deposits (asterisks) inside the
blistered disorganized epithelial. Scale bars are 50 µm.

distribution of mitotic spindle angles (Figure 4B,C). This defect was localized to the septum, as

the mitotic spindle angles in the trachea and esophagus were not significantly different between

mutants and controls (Figure S4). Thus, loss of endosomal trafficking causes aberrant cell

division into the center of the transient epithelial bilayer.

To test whether these changes in the orientation of cell division were linked to altered apical-

basal cell polarity in the septum, we performed immunostaining for laminin, which is normally

secreted from the basal side of epithelia into the extracellular matrix. We observed that mutant

embryos exhibited ectopic laminin deposition in the center of the disorganized septum

surrounding the round central cells and in the lumens of blisters within the tissue (Figure 4D-G).

These observations are consistent with disrupted endosome trafficking leading to loss of apical-

basal polarity in the transient epithelial bilayer, randomized cell division, and cells accumulating

in the mutant tracheoesophageal septum.

Endosome trafficking localizes Vangl2 to maintain apical memory in the epithelial bilayer

During tracheoesophageal morphogenesis, when the foregut epithelial cells from either side

touch one another, they lose apical markers (aPKC, Par3) and upregulate Cdh1 (typically basal

-lateral) as they adhere and form the transient bilayer. Despite this apparent loss of apical

identity, the cells must somehow retain some apical-basal information to form an organized

bilayer and undergo cell division in the plane of the epithelium – a process that we found

requires endosome trafficking. One candidate pathway that might regulate this process is the

Vangl-Celsr polarity complex. In addition to their well-known role in the Wnt/PCP pathway where

heterotypic cell-cell interactions between transmembrane Vangl, Celsr, and Frizzled proteins

govern cell orientation in the plane of the epithelia[23], Vangl and Celsr are also involved in

specifying apical-basal polarity in epithelia by recruiting apical Par3 and aPKC whilst restricting

Scrib to the basolateral domain [16,24,25]. Furthermore, apical Vangl-Celsr complexes are reported

to be internalized by endosomes during polarized cell divisions of the fetal mouse epidermis [26],

suggesting a potential mechanistic link in the foregut.

To assess whether Vangl2 and Celsr1 are involved in endosome-dependent epithelial

remodeling in the foregut, we performed immunostaining in combination with Rab11a. Prior to

foregut fusion, a low level of diffuse Vangl2 was observed throughout the foregut epithelia with

no obvious difference between apical and basolateral cell domains (Figure 5A,F, Figure S5A),

but as the midline epithelia touched, Vangl2 became progressively enriched and co-localized

with Rab11a puncta at the apical contact point (Figure 5B,E,G). Once the transient bilayer

formed, robust Rab11a-Vangl2 co-localization was observed at the adherent interface between

the two epithelial (Figure 5C,E,H) but not at the luminal/apical cells surface of the dorsal

esophagus or ventral trachea (Figure 5A-D, Figure S5B). This was somewhat unexpected since

Vangl2 normally recruits apical Par and aPKC proteins. However, in this case, Rab11-Vangl2

co-localization is associated with a loss of apical markers and the accumulation of basal-lateral

Cdh1. At NF41, Vangl2 and Rab11a levels at the cell membrane were downregulated as cells in

the bilayer lose contact with one another (Figure 5D-E). In contrast to this dynamic Vangl2

subcellular localization, Celsr1 remained around the entire juxta-membrane region of the

epithelial bilayer cells throughout tracheoesophageal morphogenesis, suggesting that it only

complexes with Vangl2 as the midline epithelial cells come in contact. (Figure S5C-H).

Figure 5. Rab11a recycling endosomes localize Vangl2 to maintain apical memory in the trachea-esophageal septum. A-D) Time course of Vangl2-Rab11a immunostaining during *Xenopus* tracheoesophageal separation. E) Dynamic Vangl2-Rab11a co-localization at the epithelial interface (Mean Pearson colocalization coefficient ± SEM, *** p<0.001, **** p<0.0001 1W-ANOVA, n=6 cells per embryo, N=5 embryos per stage). F-I) Diagrams depicting Vangl2-Rab11a localization during foregut separation. J-M) Vangl2 is mis-localized in septum epithelial cells of endosome mutants. N) Endosome mutant septa have significantly lower Vangl2-Rab11a colocalization relative to controls (mean ± SEM, **** p <0.0001 1W-ANOVA, n=5 cells per embryo, N=4-6 embryos per condition). O-R) CRISPR F0 mutation of patient variant orthologous genes in *Xenopus* (*ap3d1, itsn1, arhgap21*, and *atp6v1b1)* disrupts Vangl2-Rab11a co-localization in the TE septum. Scale bars are 50 µm (20µm in insets).

Considering that endosomes can internalize cell surface Vangl in the fetal mouse epidermis [26],

these data suggested that Vangl2 might be a cargo of the Rab11+ recycling endosomes,

transporting it to the cell surface where the epithelia fuse. To test this possibility, we examined

Vangl2 localization in the endosome mutants. Indeed Vangl2-Rab11a co-localization at the

epithelia contact was lost in the disorganized multilayer septum of *dnm2*, *rab5a* and *rab11a*

mutants (Figure 5J-N, Figure S5I). However, Celsr1 localization and apical Vangl2 in the

esophagus were unaffected (Figure 5K-M, asterisks, Figure S5J-N), indicating that recycling

endosomes are specifically required to localize Vangl2 in the fusing midline.

Mutation of EA/TEF patient-like genes in *Xenopus* disrupted Rab11-Vangl2 localization

We next assessed whether a similar mechanism could account for the TEF observed when we

mutated the *Xenopus* orthologs of endosome pathway risk genes from patients. These included

AP3D1, *ARHGAP21*, and *ITSN1* which encode adaptors and CDC42-GEFs respectively during

endocytic vesicle formation, as well as *ATP6V1B1*, which encodes a vacuolar ATPase subunit

required for acidification of endosomal vesicles [27-31]. As predicted, F0 CRISPR mutants of

ap3d1, *arhgap21*, *atp6v1b1* and *itsn1* all exhibited a loss of Vangl2-Rab11a co-localization and

a disorganized persistent septum (Figure 5O-R), consistent with these proteins functioning

upstream of Rab11 in the recycling endosome pathway.

Together these results indicate that during tracheoesophageal morphogenesis dynamic

recycling endosome activity is required to localize Vangl2 at the cell surface where the opposing

midline epithelia adhere to one another. Vangl2 appears to define an "apical memory" in the

epithelial bilayer cells which have lost aPKC and upregulate Cadherin, a membrane

configuration more typical of adherent basal-lateral cell-cell contacts. This unexpected behavior

is distinct from the typical roles for Vangl2 in apical-basal and planar cell polarity.

Loss of Vangl or Celsr disrupts tracheoesophageal morphogenesis resulting in TEF

The results above imply that Vangl-Celsr is required for tracheoesophageal morphogenesis.

Interestingly, one patient from our EA/TEF cohort had a heterozygous *de novo* damaging

missense variant in *CELSR2* predicted to alter an amino acid (asparagine to serine) in an N-

terminal cadherin domain [21,32]. Moreover, a search of clinical genetic databases and the

published literature revealed additional patients with sequence variants or copy-number

variation in *VANGL1*, *VANGL2*, *CELSR1*, and *CELSR2* who had EA/TEF or poorly defined

upper gastrointestinal and/or respiratory phenotypes all suggestive of structural foregut

anomalies (Table S2).

We therefore generated loss of function CRISPR-Cas9 F0 *Xenopus* mutants for *celsr2, celsr1,*

or *vangl2* with a 45-95% damaging indel rate, all of which exhibited a variably penetrant TEF

with a failure to separate the Sox2+ esophagus and Nkx2-1+ trachea at NF44 (Table S1).

Examination of the mutant foreguts at NF40, revealed a disorganized, thicker septum with round

cells in the middle and frequent blisters, phenocopying the endosome mutants (Figure 6A-H).

Moreover, the *celsr1 and celsr2* mutants exhibited a reduction in Vangl2 immunostaining,

consistent with previous reports that Celsr1 forms a heterotypic cell-cell complex with Vangl2[23].

Interestingly, in *vangl2* and *celsr1/2* mutants Rab11a was still localized to the contact surface in

middle the septum (Figure 6E-H), indicating that Vangl and Celsr are required for

tracheoesophageal morphogenesis downstream of recycling endosomes. Together our results

suggest that an unexpected but evolutionarily conserved endosome-cell polarity pathway

controls tracheoesophageal morphogenesis, and that disruptions to this process might

represent a previously unrecognized risk for EA/TEF in patients (Figure 6I).

Discussion

In this study we demonstrate that an endosome-cell polarity pathway is required for the

epithelial remodeling that drives *Xenopus* tracheoesophageal morphogenesis. We discovered

that Rab11+ recycling endosomes localize the transmembrane polarity protein Vangl2 to the

contact surface where foregut epithelia fuse. This appears to confer an "apical memory" which is

essential to maintain an organized transient epithelial bilayer and its subsequent fission into

distinct trachea and esophagus (Figure 6I). This unexpected Rab11-Vangl2 activity is distinct

from its typical roles in planar cell polarity and apical basal polarity and may represent a general

paradigm for epithelial fusion and fission during morphogenesis.

Previous studies in *Xenopus* gastrulation and neural tube morphogenesis have shown that

Vangl2 functions upstream of Rab11 in classical Wnt/PCP and apical-basal polarity roles to

regulate convergent extension and apical constriction. Vangl2 controls Rab11 localization for

apical constriction of gastrula bottle cells and in the neural plate hinge point. In addition, Vangl2-

mediated planar cell polarization is required for medial-apical localization of Rab11 in

convergent extension and the closing neural tube [33-35]. In the developing mouse pancreas,

apical restriction of Vangl2 maintains epithelial integrity and survival of progenitor cells through

430
431
432 **Fig. 6. Loss of Vangl or Celsr disrupts tracheoesophageal morphogenesis resulting in TEF.**
433 (A-D) Cdh1 immunostaining in shows that *celsr2*, *vangl2*, and *celsr1* mutants have TEF and (E-
434 H) disorganized epithelial septa that fail to co-localize Vangl2-Rab11a. Scale bars are 50 μm. I)
435 Model of how endosome trafficking of Vangl2 maintains apical-basal polarity information and the
436 proper tissue organization required for tracheoesophageal morphogenesis.
437
438

modulating ROCK signaling activity, independent of apical-basal cell polarity[36]. Loss of

Rab11a/b in the pancreas leads to highly disorganized epithelial defects due to the mis-

localization of apical-basal proteins laminin, Par3, and aPKC, but this study did not examine

whether this was due to Vangl mis-localization [37]. Interestingly, Vangl2 has a PCP-independent

role in the developing lung mesenchyme during branching morphogenesis [38,39]. Neither the

looptail mutation nor conditional knockout of Vangl2 using ShhCre impairs trachea-esophageal

separation, however, a different Cre driver with earlier recombination in the foregut might lead to

separation anomalies. One example where Rab11 functions upstream of Vangl2, like we have

observed, is during mitosis of the mouse fetal epidermis, where endosomes internalize apical-

basal polarized Vangl, Celsr, and Fzd proteins which are later recycled back to the appropriate

cell surface after cell division [26]. A similar mechanism might explain the maintenance of

polarized cell division in the transient epithelial bilayer of the foregut, but in the mouse epidermis

this is a classical polarized epithelium, and in the foregut Vangl2 confers "apical memory" to

cells that have otherwise lost apical character. Another distinction between what we have

described in the foregut and these other examples of Vangl-regulated epithelial polarity in

morphogenesis is that in the foregut the epithelia first fuses (like the neural tube and palate) but

then undergo fission. Other tissues that experience this type of dynamic fusion-fission

morphogenesis include separation of anorectal tubes in the hindgut and semicircular canals in

the ear [40,41,42]. We predict that similar Rab11a-Vangl2 mechanisms may act in those tissues.

Our data, together with previously published work, suggests that Vangl2 is a cargo of Rab11a+

endosomes that transport newly synthesized Vangl2 from the ER-Golgi to the fusion surface, or

recycle Vangl2 from basolateral membranes. We postulate that other endosomal cargo may

include Cadherins and Integrins that facilitate tissue adhesion, and Laminins that mediate

interaction with the adjacent mesoderm at the basal surface. Consistent with this possibility we

note that EA/TEF patients were identified with potentially damaging variants in *ITGB4* and

LAMB4 [21]. While initial Cdh1 (E-Cadherin) localization to the fusion surface was not affected by

disrupted endosome trafficking, the normal downregulation of Cdh1 at the fission step does not

occur in endosome or Vangl-Celsr mutants, which might explain why the foregut does not

separate properly. This suggests that endosomes may also be required to internalize cell

adhesion proteins during epithelial fission. One candidate pathway that might regulate

endosome-mediated loss of adhesion is the Eph/Ephrin pathway. Eph/Ephrin are known to

regulate tissue separation and cell sorting in many contexts[43] and *Efnb2* was recently shown to

be required for trachea-esophageal separation downstream of Nkx2-1[44,45]. We were not able to

assess Ephrin-Cadherin-endosome-Vangl interactions because effective Eph/Ephrin antibodies

are not currently available, however in the future it would be intriguing to generate transgenic

Xenopus or mice with tagged fusion proteins to test this possibility.

Endosome trafficking is a ubiquitous cell pathway required for homeostasis and many basic

cellular functions. Several pleiotropic diseases are associated with variants in endosome

trafficking genes [20], but interestingly many of these are rather specific to certain tissues. Not

surprisingly, total loss of endosome activity leads to early embryonic lethality in germline mutant

mice [46,47] and in *Xenopus* embryos results in gastrulation and neurulation defects [33]. Our F0

CRISPR mutagenesis on the other hand results in 30-90% damaging indels and therefore is

only a partial loss of function. Indeed, in our experiments endosome mutant embryos have a

significantly higher frequency of gastrulation and neurulation defects compared to controls

where the pigmentation gene *tyr* was mutated. In our experiments we selected those embryos

that survived to later stages. The fact that later embryos with partial loss of endosome activity

show specific defects suggests that some morphogenetic events like tracheoesophageal

separation represent developmental bottle necks that are particularly sensitive. This may

partially explain why EA/TEF patients frequently present with variants predicted to have partial

loss of function or hypomorphic variants in endosome and polarity pathway genes.

The EA/TEF risk genes that we validated in this study are predicted to function at various levels

of endosome trafficking pathways including formation of an endocytic vesicle (ITSN1, AP1G2),

cytoskeletal transport (ARHGAP21, ABRA, RASAL2, RAPGEF3), and endosomal acidification

(ATP6V1B1). We also discovered that a number of variants in CELSR1/2 and VANGL1/2,

including a damaging missense variant in CELSR2 in a patient from this cohort, are implicated

in EA/TEF and other foregut anomalies[21,48]. Furthermore, variants in VANGL1/2 underlie a

broad spectrum of developmental anomalies in humans[23] such as neural tube anomalies[49,50],

renal anomalies [51,52], congenital heart anomalies[53,54], and lung diseases[55], and may be trafficked

by endosomes to specific cellular domains during development of these organs. Complex

EA/TEF patents tend to have comorbidities in these same organs that experience tissue fusion

events during fetal development[21]. This suggests that disruption of a conserved endosome-

polarity pathway may account for pleiotropic phenotypes in many complex EA/TEF patients.

Together this suggests the provocative possibility of an endosomeopathy developmental

syndrome, akin to ciliopathies. Further comparative analysis of patient genetics and animal

models will improve our understanding of fundamental tissue fusion and fission events in

development and advance our knowledge of the developmental basis of congenital anomalies,

informing the care and treatment of endosomeopathy patients.

Acknowledgements

We thank the Bio-Imaging and Analysis Facility and Dr. Matthew Kofron for imaging and

analysis advice (RRID:SCR_022628), the Division of Veterinary Services, and the DNA

Sequencing Core at CCHMC (RRID:SCR_022630) for their services used in this study. We

thank S. Sokol for kindly providing the anti-Vangl2 antibody. We thank the members of the

CLEARconsortium.org for access to EA/TEF patient data and for helpful discussions. This work

is supported by NICHD P01HD093393 to YS, WKC, and AMZ, and a Canadian Institutes of

Health Research postdoctoral fellowship to NAE.

Author contributions

Conceptualization: NAE, AMZ. Formal analysis: NAE, AK, AMZ. Funding acquisition: YS, WKC,

AMZ. Investigation: NAE, AK, AW, ZA, APK. Methodology: NAE. Project administration: NAE,

AMZ. Supervision: NAE, AMZ. Validation: NAE. Visualization: NAE, AMZ. Writing – original

draft: NAE, AMZ. Writing – review & editing: all authors.

Declaration of interests

The authors declare no competing interests.

Methods

Experimental models

Xenopus experiments were performed using Cincinnati Children's Hospital Medical Center

(CCHMC) Institutional Animal Care and Use Committee (IACUC 2022-0026) approved

guidelines. Adult *Xenopus tropicalis* and *Xenopus laevis* frogs were purchased from NASCO

and Xenopus1 (USA) and housed in the CCHMC vivarium under standard housing conditions.

Xenopus embryos were obtained by *in vitro* fertilization according to standard procedures [56].

Sox2 germline mutant embryos were generated at the National *Xenopus* Resource according to

standard procedures (RRID:SCR_013731).

CRISPR-Cas9 gene editing and Xenopus embryo culture

Guide RNAs and genotyping primers were designed using CRISPRScan, InDelphi, and

CRISPOR predictive algorithms [57] and synthesized as altR-cRNA by IDT (sequences in

Resources Table and Table S3). After annealing with tracrRNA to create a single gRNA, gRNAs

were combined with recombinant Cas9 protein (PNA Biosciences) and injected into 1-2 cell

stage *Xenopus* embryos at a final concentration of 500-700pg gRNA and 1-1.5ng Cas9p.

Xenopus tropicalis embryos were raised at 25°C in 0.1X Modified Barth's solution (MBS) for

three days post fertilization before being anesthetized and fixed at NF44 in MEMFA (3.7%

formaldehyde, 0.1M MOPS, 2mM EGTA, 1mM $MgSO_4$) for 1hr at room temperature, then in

Dent's solution (80% methanol, 20% DMSO) for long term storage at -20°C. *Xenopus laevis*

embryos were raised between 15-21°C in 0.1X MBS for 3-5 days post fertilization (to stages

NF37-44), then anesthetized and fixed in MEMFA or 2% trichloroacetic acid (for endosome

proteins and Vangl2 immunostaining) for 30-40 mins at room temperature. Embryos were

immunostained and imaged by confocal microscopy as outlined in the next section prior to

genotyping. Genotyping for CRISPR-Cas9 editing efficiency was performed as described in

Zhong et al., 2022[21] using the primer sequences in Resources Table and Table S3. Percent

indel efficiency was determined by Sanger sequencing of PCR products and quantified using

Inference of CRISPR Edits (Synthego, USA). A minimum of 20 individual mutant embryos from

3 independent editing experiments were analyzed to identify a statistically significant frequency

of trachea-esophageal anomalies by Student's t-test from the 2% baseline detected in controls.

Whole mount immunostaining and confocal microscopy

Whole mount immunostaining on *Xenopus* tissue was performed as described in Nasr et al.,

2019[13] and Zhong et al., 2022[21] with the following modifications for imaging endosome and

polarity proteins: fixed embryos were rehydrated in PBS, embedded in 2% low melt agarose in

PBS, and sectioned with a Vibratome VT1000S (Leica) to obtain 200 µm tissue slices containing

the trachea and esophagus. Slices were permeabilized with 0.1% Triton X-100 in PBS (0.3%

Triton X-100 for anti-Vangl2), blocked in 5% normal donkey serum in PBS, then incubated in

primary antibody (1:100) in blocking solution overnight at 4°C. Slices were incubated in Alexa

fluor-conjugated secondary antibody (1:500) in blocking solution for 2 hours at room

temperature, then dehydrated in 100% methanol. For confocal microscopy, slices were cleared

in BABB and imaged on a Nikon A1 inverted laser scanning confocal microscope with a 10X,

20X water immersion, or 40X oil immersion objective (Nikon USA).

Image analyses

Signal intensity quantification: Signal intensity quantification was performed in ImageJ (NIH).

Regions of interest (ROI) were drawn around the apical and basal membranes, then mean gray

value was measured and plotted as mean intensity ± max/min. At least five cells were measured

per embryo in a minimum of five embryos per condition for quantification and statistical analysis

by 2W-ANOVA.

Prolate ellipticity and surface contact area: Prolate ellipticity and surface contact area

measurements were performed in Imaris (Bitplane). 3D cell volumes were rendered based on E-

Cadherin staining and surface models were generated. Prolate ellipticity was measured from

rendered surfaces of epithelial cells and plotted as mean ± max/min. Values approaching 1

have elongated shapes while values approaching 0 have spherical shapes. Surface-surface

contact area between the esophagus and trachea-esophageal septum, within the trachea-

esophageal septum, and between the trachea and trachea-esophageal septum was measured

from rendered surfaces of epithelial cells using the Surface-Surface Contact Area XTension and

plotted as the mean ± max/min. At least five cells were measured per embryo in a minimum of

five embryos per condition for quantification and statistical analysis by unpaired Student's t-test

(prolate ellipticity) and 1W-ANOVA (surface contact area).

Three-dimensional measurements of axes in the foregut: Three dimensional lengths of x

(medial), y (dorsal-ventral), and z (rostral-caudal) axes were performed in Imaris (Bitplane). 3D

cell volumes were rendered based on E-Cadherin staining and surface models were generated.

Lengths were measured from rendered surfaces of epithelial cells and plotted as the mean

length of each axis ± standard error of the mean. At least five embryos were measured for

quantification and statistical analysis by unpaired Student's t-test.

Mitotic spindle orientation: Mitotic spindle orientation angles were obtained in ImageJ (NIH) as

described in Franco and Carmena, 2019[58]. Only images with z-sections through the axial axis of

Xenopus embryos containing clearly identifiable mitotic spindles stained with anti-α-tubulin were

analyzed for spindle angles. One line of the angle was drawn through linking both centrosomes,

and the second line of the angle was drawn parallel to the basement membrane of the epithelial

cell. Measurements were collected and wind rose plots were generated in Excel. At least three

mitotic spindles were measured in a minimum of 5 embryos for quantification and statistical

analysis by Fisher's exact test.

Colocalization: Pearson colocalization coefficients between Vangl2/Rab11a, Cdh1/Rab11a, and

Celsr1/Rab11a immunostaining signals were calculated in Nikon Elements (Nikon USA). ROIs

were drawn around an entire cell, and the Pearson colocalization coefficient was determined. A

coefficient of 1 indicates two channels whose fluorescence intensities are perfectly linearly

related. At least six cells were measured in a minimum of 5 embryos for quantification and

statistical analyses by 1W-ANOVA.

Immunoblotting

Whole Xenopus embryos or microdissected anterior foregut tissue (from pharynx to stomach)

were pooled and lysed in Triton X-100 lysis buffer (0.5% Triton X-100, 0.1M HEPES, 150 mM

NaCl, 2mM EDTA pH 8.0, 2mM EGTA) containing protease inhibitors (ThermoFisher). Protein

lysates were combined with 2X Laemmli buffer and 0.1M DTT, boiled, and separated by 4-12%

Bis-Tris gel electrophoresis (ThermoFisher). For HRP detection, membranes were blocked in

5% BSA, then incubated in primary antibody (1:1000) overnight in blocking solution at 4°C.

Membranes were incubated in HRP-conjugated secondary antibody (1:5000) in blocking

solution for 1 hour at room temperature, then visualized by addition of ECL substrate

(Amersham, GE) and imaged using a ChemiDoc (BioRad). Western blot quantification was

performed in ImageLab (BioRad). For fluorescence-based detection, membranes were blocked

in Odyssey blocking buffer (LI-COR), then incubated in primary antibody (1:1000) overnight in

blocking buffer at 4°C. Membranes were incubated in IRDye-conjugated secondary antibodies

(1:5000) in blocking buffer for 1 hour at room temperature, then visualized using an Odyssey

Dlx Imager (LI-COR). Quantification was performed with LI-COR Acquisition software.

Clinical genetic database mining

Deidentified case data for variants in CELSR1/2 and VANGL1/2 were searched in NCBI

PubMed and the DECIPHER database (https://www.deciphergenomics.org). Results from the

DECIPHER database were filtered for EA/TEF related phenotypes including "Feeding difficulties

in infancy", "Abnormal esophagus morphology", "Respiratory distress", "Laryngeal stenosis",

and "Respiratory insufficiency".

<u>**Resources**</u>

Antibodies		
Rabbit anti-PKC zeta	Abcam	ab59364; RRID:AB_944858
Mouse anti-Cdh1	Developmental Studies Hybridoma Bank	5D3 (deposited to the DSHB by Gumbiner, B.M.); RRID:AB_528116

Rabbit anti-Dnm2	Abcam	Ab65556; RRID:AB_1603781
Rabbit anti-Rab5a	Cell Signaling Technology	3547; RRID:AB_2300649
Rabbit anti-Rab11a	Thermo Fisher Scientific	700184; RRID:AB_2532295
Mouse anti-Rab11a	BD Biosciences	610657; RRID:AB_397984
Rabbit anti-Tjp1 (ZO-1)	Thermo Fisher Scientific	61-7300; RRID:AB_138452
Rabbit anti-Par3	Millipore	07-330; RRID:AB_2101325
Mouse anti-Tuba4b	Sigma	T9026; RRID:AB_477593
Rabbit anti-Lama1	Sigma	L9393; RRID:AB_477163
Rabbit anti-pHH3 (Ser10)	Cell Signaling Technology	9701; RRID:AB_331535
Rabbit anti-Vangl2	S. Sokol lab	RRID:AB_2744499
Rabbit anti-Celsr1	BiCell Scientific	50801
Mouse anti-Sox2	Abcam	ab79351; RRID:AB_10710406
Rabbit anti-Nkx2-1	Santa Cruz	sc-13040; RRID:AB_793532
Goat anti-Foxf1	R&D Systems	AF4798; RRID:AB_2105588
Rabbit anti-Itsn1	Millipore	ABS984
Chemicals		
Cas9 protein	PNA Biosciences	CP01-50
Experimental models: organisms/strains		
Xenopus laevis females	NASCO	LM00531
Xenopus laevis males	NASCO	LM00715
Xenopus tropicalis females	Xenopus1	
Xenopus tropicalis males	Xenopus1	
Xenopus tropicalis sox2 mutant embryos	National Xenopus Resource	NXR_3051; Xtr.sox2[tmHorb]
Oligonucleotides *(See also Table S3)*		
Rab11a.S gRNA GAGATTACTCTTCCCCACAC	IDT	Nasr et al., 2019[13]
Rab5a gRNA CGACTGCAGATTCTCCTAAA	IDT	This study
Dnm2 gRNA CGGAGACCGAGTGCAGCCAT	IDT	This study
Slc45a2 gRNA GGTTACATAGGCTGCCTCCA	IDT	Delay et al., 2018[59]
Genotyping, Rab11a.S forward primer ACAGTGTGCATAGAATTTGGCG	IDT	This study
Genotyping, Rab11a.S reverse primer CGTGTGTTGCCTAGAGGGAC	IDT	This study
Genotyping, Dnm2.S forward primer GGAGCCAGCGAGAAACACTTA	IDT	This study
Genotyping, Dnm2.S reverse primer GCGACAATATCAACCCACCAA	IDT	This study
Genotyping, Dnm2.L forward primer	IDT	This study

AGGGAAACACTTACCGGCCTA		
Genotyping, Dnm2.L reverse primer GAAACACAACCAATCACAATCCCT	IDT	This study
Genotyping, Rab5a.S forward primer ACCCTGGCAAACTTTGTGGT	IDT	This study
Genotyping, Rab5a.L forward primer GAGCATCCAACCCACCCTTT		This study
Genotyping, Rab5a reverse primer ACTGTCCCTTTACGAAGCGA	IDT	This study
Software		
NIS Elements	Nikon	RRID:SCR_014329
ImageJ	NIH	RRID:SCR_003070
IMARIS	Oxford Instruments	RRID:SCR_007370
Surface-Surface Contact Area XTension	Oxford Instruments	https://imaris.oxinst.com/open/view/surface-surface-contact-area
Prism 9	GraphPad	RRID:SCR_002798
Inference of CRISPR edits (ICE)	Synthego	http://www.ice.synthego.com
ImageLab	BioRad	RRID:SCR_014210

References

1. Edwards, N.A., Shacham-Silverberg, V., Weitz, L., Kingma, P.S., Shen, Y., Wells, J.M., Chung, W.K., and Zorn, A.M. (2021). Developmental basis of trachea-esophageal birth defects. Dev Biol *477*, 85-97. 10.1016/j.ydbio.2021.05.015.

2. Raad, S., David, A., Que, J., and Faure, C. (2020). Genetic Mouse Models and Induced Pluripotent Stem Cells for Studying Tracheal-Esophageal Separation and Esophageal Development. Stem Cells Dev *29*, 953-966. 10.1089/scd.2020.0075.

3. van Lennep, M., Singendonk, M.M.J., Dall'Oglio, L., Gottrand, F., Krishnan, U., Terheggen-Lagro, S.W.J., Omari, T.I., Benninga, M.A., and van Wijk, M.P. (2019). Oesophageal atresia. Nat Rev Dis Primers *5*, 26. 10.1038/s41572-019-0077-0.

4. Chitkara, A.E., Tadros, M., Kim, H.J., and Harley, E.H. (2003). Complete laryngotracheoesophageal cleft: complicated management issues. Laryngoscope *113*, 1314-1320. 10.1097/00005537-200308000-00010.

5. Bednarczyk, D., Sasiadek, M.M., and Smigiel, R. (2013). Chromosome aberrations and gene mutations in patients with esophageal atresia. J Pediatr Gastroenterol Nutr *57*, 688-693. 10.1097/MPG.0b013e3182a373dc.

6.   Stoll, C., Alembik, Y., Dott, B., and Roth, M.P. (2017). Associated anomalies in cases with
esophageal atresia. Am J Med Genet A *173*, 2139-2157. 10.1002/ajmg.a.38303.

7.   Brosens, E., Ploeg, M., van Bever, Y., Koopmans, A.E., H, I.J., Rottier, R.J., Wijnen, R., Tibboel,
D., and de Klein, A. (2014). Clinical and etiological heterogeneity in patients with tracheo-
esophageal malformations and associated anomalies. Eur J Med Genet *57*, 440-452.
10.1016/j.ejmg.2014.05.009.

8.   Li, Y., Litingtung, Y., Ten Dijke, P., and Chiang, C. (2007). Aberrant Bmp signaling and notochord
delamination in the pathogenesis of esophageal atresia. Dev Dyn *236*, 746-754.
10.1002/dvdy.21075.

9.   Goss, A.M., Tian, Y., Tsukiyama, T., Cohen, E.D., Zhou, D., Lu, M.M., Yamaguchi, T.P., and
Morrisey, E.E. (2009). Wnt2/2b and beta-catenin signaling are necessary and sufficient to specify
lung progenitors in the foregut. Dev Cell *17*, 290-298. 10.1016/j.devcel.2009.06.005.

10.   Harris-Johnson, K.S., Domyan, E.T., Vezina, C.M., and Sun, X. (2009). beta-Catenin promotes
respiratory progenitor identity in mouse foregut. Proc Natl Acad Sci U S A *106*, 16287-16292.
10.1073/pnas.0902274106.

11.   Rankin, S.A., Han, L., McCracken, K.W., Kenny, A.P., Anglin, C.T., Grigg, E.A., Crawford, C.M.,
Wells, J.M., Shannon, J.M., and Zorn, A.M. (2016). A Retinoic Acid-Hedgehog Cascade
Coordinates Mesoderm-Inducing Signals and Endoderm Competence during Lung Specification.
Cell Rep *16*, 66-78. 10.1016/j.celrep.2016.05.060.

12.   Kim, E., Jiang, M., Huang, H., Zhang, Y., Tjota, N., Gao, X., Robert, J., Gilmore, N., Gan, L., and
Que, J. (2019). Isl1 Regulation of Nkx2.1 in the Early Foregut Epithelium Is Required for Trachea-
Esophageal Separation and Lung Lobation. Dev Cell *51*, 675-683 e674.
10.1016/j.devcel.2019.11.002.

13.   Nasr, T., Mancini, P., Rankin, S.A., Edwards, N.A., Agricola, Z.N., Kenny, A.P., Kinney, J.L.,
Daniels, K., Vardanyan, J., Han, L., et al. (2019). Endosome-Mediated Epithelial Remodeling
Downstream of Hedgehog-Gli Is Required for Tracheoesophageal Separation. Dev Cell *51*, 665-
674 e666. 10.1016/j.devcel.2019.11.003.

14.   Cote, L.E., and Feldman, J.L. (2022). Won't You be My Neighbor: How Epithelial Cells Connect
684   Together to Build Global Tissue Polarity. Front Cell Dev Biol *10*, 887107.
685   10.3389/fcell.2022.887107.

686   15.   Pickett, M.A., Naturale, V.F., and Feldman, J.L. (2019). A Polarizing Issue: Diversity in the
687   Mechanisms Underlying Apico-Basolateral Polarization In Vivo. Annu Rev Cell Dev Biol *35*, 285-
688   308. 10.1146/annurev-cellbio-100818-125134.

689   16.   Aigouy, B., and Le Bivic, A. (2016). The PCP pathway regulates Baz planar distribution in
690   epithelial cells. Sci Rep *6*, 33420. 10.1038/srep33420.

691   17.   Murdoch, J.N., Doudney, K., Paternotte, C., Copp, A.J., and Stanier, P. (2001). Severe neural
692   tube defects in the loop-tail mouse result from mutation of Lpp1, a novel gene involved in floor
693   plate specification. Hum Mol Genet *10*, 2593-2601. 10.1093/hmg/10.22.2593.

694   18.   Butler, M.T., and Wallingford, J.B. (2018). Spatial and temporal analysis of PCP protein dynamics
695   during neural tube closure. Elife *7*. 10.7554/eLife.36456.

696   19.   Torban, E., Wang, H.J., Patenaude, A.M., Riccomagno, M., Daniels, E., Epstein, D., and Gros, P.
697   (2007). Tissue, cellular and sub-cellular localization of the Vangl2 protein during embryonic
698   development: effect of the Lp mutation. Gene Expr Patterns *7*, 346-354.
699   10.1016/j.modgep.2006.07.007.

700   20.   Yarwood, R., Hellicar, J., Woodman, P.G., and Lowe, M. (2020). Membrane trafficking in health
701   and disease. Dis Model Mech *13*. 10.1242/dmm.043448.

702   21.   Zhong, G., Ahimaz, P., Edwards, N.A., Hagen, J.J., Faure, C., Lu, Q., Kingma, P., Middlesworth,
703   W., Khlevner, J., El Fiky, M., et al. (2022). Identification and validation of candidate risk genes in
704   endocytic vesicular trafficking associated with esophageal atresia and tracheoesophageal
705   fistulas. HGG Adv *3*, 100107. 10.1016/j.xhgg.2022.100107.

706   22.   Szklarczyk, D., Kirsch, R., Koutrouli, M., Nastou, K., Mehryary, F., Hachilif, R., Gable, A.L., Fang,
707   T., Doncheva, N.T., Pyysalo, S., et al. (2023). The STRING database in 2023: protein-protein
708   association networks and functional enrichment analyses for any sequenced genome of interest.
709   Nucleic Acids Res *51*, D638-D646. 10.1093/nar/gkac1000.

23.   Butler, M.T., and Wallingford, J.B. (2017). Planar cell polarity in development and disease. Nat
Rev Mol Cell Biol *18*, 375-388. 10.1038/nrm.2017.11.

24.   Panzica, D.A., Findlay, A.S., van Ladesteijn, R., and Collinson, J.M. (2019). The core planar cell
polarity gene, Vangl2, maintains apical-basal organisation of the corneal epithelium. J Anat *234*,
106-119. 10.1111/joa.12676.

25.   vandenBerg, A.L., and Sassoon, D.A. (2009). Non-canonical Wnt signaling regulates cell polarity
in female reproductive tract development via van gogh-like 2. Development *136*, 1559-1570.
10.1242/dev.034066.

26.   Devenport, D., Oristian, D., Heller, E., and Fuchs, E. (2011). Mitotic internalization of planar cell
polarity proteins preserves tissue polarity. Nat Cell Biol *13*, 893-902. 10.1038/ncb2284.

27.   Yang, Q., Li, G., Singh, S.K., Alexander, E.A., and Schwartz, J.H. (2006). Vacuolar H+ -ATPase
B1 subunit mutations that cause inherited distal renal tubular acidosis affect proton pump
assembly and trafficking in inner medullary collecting duct cells. J Am Soc Nephrol *17*, 1858-
1866. 10.1681/ASN.2005121277.

28.   Dong, X., Li, H., Derdowski, A., Ding, L., Burnett, A., Chen, X., Peters, T.R., Dermody, T.S.,
Woodruff, E., Wang, J.J., and Spearman, P. (2005). AP-3 directs the intracellular trafficking of
HIV-1 Gag and plays a key role in particle assembly. Cell *120*, 663-674.
10.1016/j.cell.2004.12.023.

29.   Kantheti, P., Qiao, X., Diaz, M.E., Peden, A.A., Meyer, G.E., Carskadon, S.L., Kapfhamer, D.,
Sufalko, D., Robinson, M.S., Noebels, J.L., and Burmeister, M. (1998). Mutation in AP-3 delta in
the mocha mouse links endosomal transport to storage deficiency in platelets, melanosomes, and
synaptic vesicles. Neuron *21*, 111-122. 10.1016/s0896-6273(00)80519-x.

30.   Dubois, T., Paléotti, O., Mironov, A.A., Fraisier, V., Stradal, T.E., De Matteis, M.A., Franco, M.,
and Chavrier, P. (2005). Golgi-localized GAP for Cdc42 functions downstream of ARF1 to control
Arp2/3 complex and F-actin dynamics. Nat Cell Biol *7*, 353-364. 10.1038/ncb1244.

31.   Gryaznova, T., Gubar, O., Burdyniuk, M., Kropyvko, S., and Rynditch, A. (2018). WIP/ITSN1
complex is involved in cellular vesicle trafficking and formation of filopodia-like protrusions. Gene
674, 49-56. 10.1016/j.gene.2018.06.078.

32.   Wang, J., Ahimaz, P.R., Hashemifar, S., Khlevner, J., Picoraro, J.A., Middlesworth, W., Elfiky,
739   M.M., Que, J., Shen, Y., and Chung, W.K. (2021). Novel candidate genes in esophageal
740   atresia/tracheoesophageal fistula identified by exome sequencing. Eur J Hum Genet *29*, 122-130.
741   10.1038/s41431-020-0680-2.

742   33.   Lee, J.Y., and Harland, R.M. (2010). Endocytosis is required for efficient apical constriction during
743   Xenopus gastrulation. Curr Biol *20*, 253-258. 10.1016/j.cub.2009.12.021.

744   34.   Ossipova, O., Kim, K., Lake, B.B., Itoh, K., Ioannou, A., and Sokol, S.Y. (2014). Role of Rab11 in
745   planar cell polarity and apical constriction during vertebrate neural tube closure. Nat Commun *5*,
746   3734. 10.1038/ncomms4734.

747   35.   Ossipova, O., Chuykin, I., Chu, C.W., and Sokol, S.Y. (2015). Vangl2 cooperates with Rab11 and
748   Myosin V to regulate apical constriction during vertebrate gastrulation. Development *142*, 99-107.
749   10.1242/dev.111161.

750   36.   Flasse, L., Yennek, S., Cortijo, C., Barandiaran, I.S., Kraus, M.R., and Grapin-Botton, A. (2020).
751   Apical Restriction of the Planar Cell Polarity Component VANGL in Pancreatic Ducts Is Required
752   to Maintain Epithelial Integrity. Cell Rep *31*, 107677. 10.1016/j.celrep.2020.107677.

753   37.   Barlow, H.R., Ahuja, N., Bierschenk, T., Htike, Y., Fassetta, L., Azizoglu, D.B., Flores, J., Gao, N.,
754   De la, O.S., Sneddon, J.B., et al. (2023). Rab11 is essential to pancreas morphogenesis, lumen
755   formation and endocrine mass. Dev Biol *499*, 59-74. 10.1016/j.ydbio.2023.05.002.

756   38.   Paramore, S.V., Trenado-Yuste, C., Sharan, R., Devenport, D., and Nelson, C.M. (2022). Vangl
757   facilitates mesenchymal thinning during lung sacculation independently of Celsr. bioRxiv.
758   10.1101/2022.12.28.522148.

759   39.   Paramore, S.V., Goodwin, K., Devenport, D., and Nelson, C.M. (2023). Mesenchymal
760   *Vangl* facilitates airway elongation and widening independently of the planar cell
761   polarity complex. bioRxiv, 2023.2007.2003.547543. 10.1101/2023.07.03.547543.

762   40.   Seifert, A.W., Harfe, B.D., and Cohn, M.J. (2008). Cell lineage analysis demonstrates an
763   endodermal origin of the distal urethra and perineum. Dev Biol *318*, 143-152.
764   10.1016/j.ydbio.2008.03.017.

41.   Martin, P., and Swanson, G.J. (1993). Descriptive and experimental analysis of the epithelial
remodellings that control semicircular canal formation in the developing mouse inner ear. Dev
Biol *159*, 549-558. 10.1006/dbio.1993.1263.

42.   Alsina, B., and Whitfield, T.T. (2017). Sculpting the labyrinth: Morphogenesis of the developing
inner ear. Semin Cell Dev Biol *65*, 47-59. 10.1016/j.semcdb.2016.09.015.

43.   Niethamer, T.K., and Bush, J.O. (2019). Getting direction(s): The Eph/ephrin signaling system in
cell positioning. Dev Biol *447*, 42-57. 10.1016/j.ydbio.2018.01.012.

44.   Lewis, A.E., Kuwahara, A., Franzosi, J., and Bush, J.O. (2022). Tracheal separation is driven by
NKX2-1-mediated repression of Efnb2 and regulation of endodermal cell sorting. Cell Rep *38*,
110510. 10.1016/j.celrep.2022.110510.

45.   Lewis, A.E., Hwa, J., Wang, R., Soriano, P., and Bush, J.O. (2015). Neural crest defects in
ephrin-B2 mutant mice are non-autonomous and originate from defects in the vasculature. Dev
Biol *406*, 186-195. 10.1016/j.ydbio.2015.08.021.

46.   Yu, S., Yehia, G., Wang, J., Stypulkowski, E., Sakamori, R., Jiang, P., Hernandez-Enriquez, B.,
Tran, T.S., Bonder, E.M., Guo, W., and Gao, N. (2014). Global ablation of the mouse Rab11a
gene impairs early embryogenesis and matrix metalloproteinase secretion. J Biol Chem *289*,
32030-32043. 10.1074/jbc.M113.538223.

47.   Ferguson, S.M., Raimondi, A., Paradise, S., Shen, H., Mesaki, K., Ferguson, A., Destaing, O.,
Ko, G., Takasaki, J., Cremona, O., et al. (2009). Coordinated actions of actin and BAR proteins
upstream of dynamin at endocytic clathrin-coated pits. Dev Cell *17*, 811-822.
10.1016/j.devcel.2009.11.005.

48.   Zhang, R., Gehlen, J., Kawalia, A., Melissari, M.T., Dakal, T.C., Menon, A.M., Hofele, J.,
Riedhammer, K., Waffenschmidt, L., Fabian, J., et al. (2020). Human exome and mouse
embryonic expression data implicate ZFHX3, TRPS1, and CHD7 in human esophageal atresia.
PLoS One *15*, e0234246. 10.1371/journal.pone.0234246.

49.   Iliescu, A., Gravel, M., Horth, C., and Gros, P. (2014). Independent mutations at Arg181 and
Arg274 of Vangl proteins that are associated with neural tube defects in humans decrease protein
stability and impair membrane targeting. Biochemistry *53*, 5356-5364. 10.1021/bi500400g.

50. Merello, E., Mascelli, S., Raso, A., Piatelli, G., Consales, A., Cama, A., Kibar, Z., Capra, V., and
Marco, P.D. (2015). Expanding the mutational spectrum associated to neural tube defects:
literature revision and description of novel VANGL1 mutations. Birth Defects Res A Clin Mol
Teratol *103*, 51-61. 10.1002/bdra.23305.

51. Papakrivopoulou, E., Vasilopoulou, E., Lindenmeyer, M.T., Pacheco, S., Brzoska, H.L., Price,
K.L., Kolatsi-Joannou, M., White, K.E., Henderson, D.J., Dean, C.H., et al. (2018). Vangl2, a
planar cell polarity molecule, is implicated in irreversible and reversible kidney glomerular injury. J
Pathol *246*, 485-496. 10.1002/path.5158.

52. Schnell, U., and Carroll, T.J. (2016). Planar cell polarity of the kidney. Exp Cell Res *343*, 258-266.
10.1016/j.yexcr.2014.11.003.

53. Yuan, Y., Gao, Y., Wang, H., Ma, X., Ma, D., and Huang, G. (2014). Promoter methylation and
expression of the VANGL2 gene in the myocardium of pediatric patients with tetralogy of fallot.
Birth Defects Res A Clin Mol Teratol *100*, 973-984. 10.1002/bdra.23291.

54. Li, D., and Wang, J. (2018). Planar Cell Polarity Signaling in Mammalian Cardiac Morphogenesis.
Pediatr Cardiol *39*, 1052-1062. 10.1007/s00246-018-1860-5.

55. Yates, L.L., and Dean, C.H. (2011). Planar polarity: A new player in both lung development and
disease. Organogenesis *7*, 209-216. 10.4161/org.7.3.18462.

56. Wlizla, M., McNamara, S., and Horb, M.E. (2018). Generation and Care of Xenopus laevis and
Xenopus tropicalis Embryos. Methods Mol Biol *1865*, 19-32. 10.1007/978-1-4939-8784-9_2.

57. Naert, T., Tulkens, D., Edwards, N.A., Carron, M., Shaidani, N.I., Wlizla, M., Boel, A., Demuynck,
S., Horb, M.E., Coucke, P., et al. (2020). Maximizing CRISPR/Cas9 phenotype penetrance
applying predictive modeling of editing outcomes in Xenopus and zebrafish embryos. Sci Rep *10*,
14662. 10.1038/s41598-020-71412-0.

58. Franco, M., and Carmena, A. (2019). Measurement of Mitotic Spindle Angle and Mitotic Cell
Distance in Fixed Tissue of Drosophila Larval Brains. Bio Protoc *9*, e3432.
10.21769/BioProtoc.3432.

59.   DeLay, B.D., Corkins, M.E., Hanania, H.L., Salanga, M., Deng, J.M., Sudou, N., Taira, M., Horb,

M.E., and Miller, R.K. (2018). Tissue-Specific Gene Inactivation in Xenopus laevis: Knockout of

lhx1 in the Kidney with CRISPR/Cas9. Genetics *208*, 673-686. 10.1534/genetics.117.300468.

824
825
826 **Supplemental Figure 1 (Related to Figure 1) F0 CRISPR-Cas9 editing reduces protein**
827 **abundance.** Western blots of Xenopus CRISPR mutants show that mutations produced by
828 gRNAs reduce but not completely ablate the protein abundance of Dnm2, Rab5a, Rab11a, and
829 Itsn1 in F0 *X. laevis* mutants. Relative protein abundance indicated by numbers above
830 immunoblots, with control protein abundance normalized to 1.
831

832
833
Supplemental Figure 2 (Related to Figure 2) Dynamic cell shape and polarity changes
during trachea-esophageal separation. A-C) Before fusion the foregut is a pseudostratified
epithelium with apical tight junctions (Tjp1) and Par3. D-F) After apical fusion, epithelial cells

remodel to upregulate apical E-Cadherin (Cdh1) and downregulate Tjp1 and Par3. G)
Quantification of aPKC downregulation after epithelial fusion (mean ± min/max, ****p<0.00001,
2W-ANOVA, n=6 cells per embryo, N=5 embryos). H) Quantification of Dnm2 intensity after
epithelial fusion (mean ± min/max, 2W-ANOVA, n=6 cells per embryo, N=5 embryos). I)
Quantification of Rab5a intensity after epithelial fusion (mean ± min/max, 2W-ANOVA, n=6 cells
per embryo, N=5 embryos). J) Before resolution of the septum, the foregut epithelium remodels
to a bilayered columnar epithelium with basally positioned nuclei. K) The septum resolves after
downregulation of E-Cadherin and integration of septum cells into either the trachea or
esophagus. L) Prolate ellipticity is significantly decreased in cells actively resolving in the septum
(mean ± min/max, *p<0.05, Student's t-test, n=6 cells per embryo, N=5 embryos). M)
Quantification of Dnm2 intensity before and after resolution (mean ± min/max, Student's t-test,
n=6 cells per embryo, N=5 embryos) N) Quantification of Rab5a intensity before and after
resolution (mean ± min/max, *p<0.05, Student's t-test, n=6 cells per embryo, N=5 embryos). O)
3D cell-surface rendering of the resolving septum with esophageal cells in green, septum cells in
yellow, tracheal cells in purple. Septum cells significantly decrease surface contact area with each
other compared to the contact between septum cells and esophageal cells (mean ± min/max, ***
p<0.001 1W-ANOVA, n=6 cells per embryo, N=9 embryos). Scale bars are 50 µm.
854

855
856
Supplemental Figure 3 (Related to Figure 3) Endosome mutants undergo epithelial fusion.
A) Unilateral CRISPR injection at the 2-cell embryo stage of gRNAs against *dnm2*, *rab5a*, and
rab11a show downregulation of fluorescence intensity compared to the control, non-injected side.
B) *dnm2*, *rab5a*, and *rab11a* mutant Xenopus embryos undergo epithelial fusion at stage NF37
comparably to controls, significantly downregulating aPKC and upregulating Cdh1 (C) after
epithelial fusion (mean ± min/max, Student's t-test, n=6 cells per embryo, N=5 embryos). D)
Mutant TE septum epithelial cells do not downregulate Cdh1 during at the point of tissue resolution
compared to controls (mean ± SEM, * p<0.05 1W-ANOVA, n=6 cells per embryo, N=5-10
embryos).
866

867
868
869
Supplemental Figure 4 (Related to Figure 4). Mitotic spindle angles are not significantly
changed compared to controls in the trachea and esophageal epithelia of endosome
mutants. A) Control embryo trachea and esophagus epithelia most frequently divide between 0-
15°C relative to the plane of the tissue. B-D) Endosome mutant trachea and esophagus epithelia
division angles are not significantly altered (Fisher's exact test, n.s. p>0.05, 3-5 cells measured
per embryo, 5 embryos measured per condition).
876

Supplemental Figure 5 (Related to Figure 5) Celsr1 is localized to the periphery of cells
during tracheoesophageal separation. A) Quantification of Vangl2 intensity before and after
epithelial fusion (mean ± min/max, 2W-ANOVA, ****$p < 0.00001$. n=3-5 cells, 5 embryos analyzed).
B) Immunostaining of Vangl2 in the trachea and esophageal epithelium. Vangl2 is localized
apically in the lumen-facing esophageal epithelia but not in the more basal cell layers and does
not colocalize with Rab11a staining (inset). C) Celsr1 does not significantly localize with Rab11a
after epithelial fusion and is localized around the periphery of epithelial and mesenchymal cells
during tracheoesophageal development. D-F) Time course of Celsr1 immunostaining during
Xenopus trachea-esophageal morphogenesis. H) Diagrams depicting Celsr1/Rab11a localization
during foregut separation. I) Vangl2 intensity at the apical membrane is decreased in dnm2,
rab5a, and rab11a CRISPR mutants, while Rab11a remains localized to the apical membranes
(mean ± min/max, 2W-ANOVA, ****$p < 0.0001$, ***$p < 0.001$, **$p < 0.01$, n=3-5 cells per embryo, 5
embryos analyzed). J-N) Celsr1:Rab11a colocalization is significantly decreased in *rab5a*
Xenopus mutants, but Celsr1 cellular localization is not significantly altered in *dnm2*, *rab5a*, *and*
rab11a Xenopus mutants (mean ± min/max, 1W-ANOVA, **$p < 0.01$. n=3-5 cells, 5 embryos
analyzed).

Xenopus ortholog of patient gene	Protein Function	Frequency of alleles with damaging indels	Mean frequency of trachea-esophageal defect (±SEM, n, N, p value)	Xenopus Phenotype
sox2^	Transcription factor	100% (germline)	100% (±0, 14, 2, p<0.0001)***	TEF – 14/14
sox2^	Transcription factor	91%	63% (±10, 17, 3, p<0.0001)***	TEF – 11/11
atp6v1b1	Intracellular transport	42-92%	44% (±19,29, 2, p=0.0024)**	TEF – 4/14 Dysmorphic esophagus – 9/14
abra^	Rho-actin regulation	20-98%	42% (±8, 30, 4, p=0.0009)**	TEF – 10/12 Dysmorphic esophagus – 4/12
itsn1^	Endocytosis, Cc42 GEF	51-92%	42% (±13, 26, 3, p=0.0038)**	TEF – 6/11 Dysmorphic esophagus – 5/11
arhgap21^	Rho family GTPase	31-76%	39% (±7, 31, 4, p=0.0006)**	TEF – 5/11 Dysmorphic esophagus – 8/11
itgb4^	Integrin adhesion, putative cargo	42-85%	34% (±8, 45, 5, p=0.0063)**	TEF – 9/13 Dysmorphic esophagus – 4/13
lamb4	Laminin subunit	26-76%	30% (±19, 32, 3, p=0.0121)*	TEF - 9/10 Dysmorphic esophagus – 1/10
celsr2^	Cell polarity, putative cargo	43-75%	28% (±5, 30, 2, p=0.0041)**	TEF - 6/8 Dysmorphic esophagus – 2/8
rasal2	Ras GTPase-like	40-81%	27% (±2, 22, 3, p=0.0005)***	TEF - 5/6 Dysmorphic esophagus – 5/6
ap1g2^	Endocytosis	35-87%	20% (±3, 40, 5, p=0.0038)**	TEF – 7/7
rapgef3^	Ras signaling	39-87%	17% (±3, 32, 3, p=0.0360)*	TEF – 4/4 Dysmorphic esophagus – 2/4
ap3d1	Intracellular transport	70-90%	15% (±7, 21, 2, p=0.0808)	-
cldn10	Tight junctions	85-86%	14% (±7, 34, 3, p=0.1106)	-
pcdh1^	Cell adhesion, putative cargo	60-76%	13% (±7, 39, 3, p=0.1361)	-
itpr1	Inositol triphosphate receptor	93%	8% (±14, 25, 3, p=0.4262)	-
rasa2	Ras GTPase	60-93%	9% (±8, 38, 4, p=0.1950)	-
add1	Cytoskeleton	89%	5% (±5, 37, 3, p=0.6608)	-
rab3gap2^	Endocytosis	81%	5% (±6, 31, 3, p=0.5948)	-
arhgap17^	Rho signaling	78%	4% (±4, 31, 5, p=0.7341)	-
akap13	Rho guanine exchange factor	85%	6% (±6, 29, 3 p=0.5165)	-
tyr (control)	Pigmentation	n/d	2% (±2, 36, 6)	-

Endosome genes	Protein Function	Frequency of alleles with damaging indels	Mean frequency of trachea-esophageal defect (±SEM, n, N, p value)	Xenopus Phenotype
X.t. dnm2	Endocytosis	62-98%	71% (±9, 18, 2, p<0.0001)***	TEF – 7/18 Dysmorphic esophagus – 5/18
X.t. rab5a	Early endosome GTPase	45-75%	50% (10, 1)	TEF – 4/10 Dysmorphic esophagus – 2/10
X.t. rab11a	Recycling endosome GTPase	42-92%	39% (±1, 13, 2, p=0.0003)***	TEF – 1/13 Dysmorphic esophagus – 5/18
X.l. dnm2.L + dnm2.S	Endocytosis	47-93% (L) 50-94% (S)	95% (±4, 22, 4, p<0.00001)****	TEF – 19/20 Dysmorphic esophagus – 11/20
X.l. rab5a.L + rab5a.S	Early endosome GTPase	45-74% (L) 46-93% (S)	88% (±7, 32, 5, p=0.0002)***	TEF – 24/28 Dysmorphic esophagus – 8/28
X.l. rab11a.S	Recycling endosome GTPase	65-95%	75% (±14, 12, 3, p=0.0065)**	TEF - 7/9 Dysmorphic esophagus – 3/9
X.l. vangl2.L + vangl2.S	Cell polarity, putative cargo	56-95% (L) 46-81% (S)	32% (±13, 22, 2, p=0.05)	TEF – 5/7 Dysmorphic esophagus – 1/7
X.l. celsr1.L + celsr1.S	Cell polarity, putative cargo	77-95% (L) 84-93% (S)	42% (±9, 17, 3, p=0.0096)**	TEF - 6/7 Dysmorphic esophagus – 1/7

Supplemental Table 1.

^reported in Zhong et al., HGG Advances 20222
TEF – trachea-esophageal fistula
SEM – standard error of the mean, n=number of embryos assayed, N=number of independent experiments
p value determined by unpaired Student's t-test compared to *tyr* controls

Gene	Patient genotype-phenotype	Reference
CELSR1	DNV – EA	Zhang et al., 2020[48]
	CNV – feeding difficulties in infancy, abnormal esophagus morphology, respiratory distress	DECIPHER 1937 (+10 patients), 254458, 397368 (+3 patients)
CELSR2	DNV – EA/TEF	Zhong et al., 2022[21]
VANGL1	CNV– feeding difficulties in infancy	DECIPHER 263708
VANGL2	CNV – laryngeal stenosis, respiratory insufficiency, feeding difficulties in infancy	DECIPHER 285906, 289199, 263708

Supplemental Table 2.

DNV – *de novo* variant; CNV – copy number variant; data taken from the DECIPHER database https://www.deciphergenomics.org