Endocannabinoids and their receptors modulate endometriosis pathogenesis

and immune response

Harshavardhan Lingegowda[1], Katherine B Zutautas[1], Yuhong Wei[2], Priyanka Yolmo[1,3], Danielle

J Sisnett[1], Alison McCallion[1], Madhuri Koti[1,3], and Chandrakant Tayade[1]

[1] Department of Biomedical and Molecular Sciences, Queen's University, Kingston, ON, Canada

[2] Rosalind and Morris Goodman Cancer Institute, McGill University, Montreal, Quebec, Canada

[3] Division of Cancer Biology and Genetics, Queen's University, Kingston, ON, Canada.

Funding: This research was supported with funds from Canadian Institutes of Health Research

(CIHR)

Corresponding Author:

Dr. Chandrakant Tayade

Department of Biomedical and Molecular Sciences,

Queen's University, Kingston, ON, K7L 3N6

Email: tayadec@queensu.ca

Phone: 613-533-6354

Abstract

Endometriosis (EM), characterized by the presence of endometrial-like tissue outside the uterus, is the leading cause of chronic pelvic pain and infertility in females of reproductive age. Despite its high prevalence, the molecular mechanisms underlying EM pathogenesis remain poorly understood. The endocannabinoid system (ECS) is known to influence several cardinal features of this complex disease including pain, vascularization, and overall lesion survival, but the exact mechanisms are not known. Utilizing CNR1 knockout (k/o), CNR2 k/o and wild type (WT) mouse models of EM, we reveal contributions of ECS and these receptors in disease initiation, progression, and immune modulation. Particularly, we identified EM-specific T cell dysfunction in the CNR2 k/o mouse model of EM. We also demonstrate the impact of decidualization-induced changes on ECS components, and the unique disease-associated transcriptional landscape of ECS components in EM. Imaging Mass Cytometry (IMC) analysis revealed distinct features of the microenvironment between CNR1, CNR2, and WT genotypes in the presence or absence of decidualization. This study, for the first time provides an in-depth analysis of the involvement of the ECS in EM pathogenesis and lays the foundation for the development of novel therapeutic interventions to alleviate the burden of this debilitating condition.

Introduction

Endometriosis (EM) is a chronic gynecological disorder characterized by the presence and growth of endometrial like tissue outside the uterus, referred to as ectopic lesions. Despite its global impact on approximately 200 million individuals and the profound reduction in their quality of life, the exact origins of EM remain elusive[1]. Accumulating evidence, including our previous studies, highlight that components of the endocannabinoid system (ECS) are dysregulated within EM lesion microenvironment as well as in the systemic circulation of EM patients[2–4]. The ECS is a complex signaling network comprised of canonical receptors (CNR1 and CNR2) and endocannabinoid (EC) ligands, along with non-canonical extended signaling network of ligands and enzymes (extensively reviewed elsewhere[5]). CNR1 and CNR2 are primarily expressed in nerve tissues, immune cells, and reproductive tissues, where they regulate various physiological processes, including pain perception, immune responses, and reproductive functions[6]. Consequently, EM pathogenesis has been postulated as a consequence of EC deficiency[7,8].

Even though the precise etiology of EM is not known, the widely accepted Sampson's theory of retrograde menstruation suggests that EM lesions originate from refluxed, endometrial fragments deposited during menstruation[9]. Both pregnancy and menstruation depend on spontaneous decidualization of endometrial stroma that is extensively remodelled under the influence of hormones, growth factors, and select cytokines that orchestrate immune cell recruitment and vascular adaptions[10]. There is clear evidence that EM patients have defects in eutopic endometrium, including differential expression of key endometrial receptivity markers such as leukemia inhibitory factor (*LIF*), protein arginine methyltransferase 5 (*PRMT5*), and homeobox protein hox-A10 (*HOXA10*), that have been associated with EM and subsequent infertility[11–13]. Evidence also suggests that components of the ECS, including CNR1 and CNR2, are important in

maintaining tissue integrity during decidualization and successful implantation of the embryo.

Indeed, several reports indicate that mice lacking cannabinoid receptors, CNR1 and CNR2,

displayed impaired implantation, increased pregnancy failure rates, heightened edema, and

inadequate primary decidual zone formation, highlighting the crucial role of ECS signaling in

successful decidualization, implantation, and pregnancy[14–16].

In EM, CNR1 and CNR2 activation aids in controlling lesion proliferation, pain, and

vascularization[17,18]. Keeping in view dysregulated ECS signalling and their central role in

decidualization and fertility, we hypothesize that altered CNR1 and CNR2 expression will disrupt

ECS signaling dynamics, leading to further lesion development. Furthermore, involvement of ECS

in modulating immune response and homeostasis, may disrupt the immune dynamics and foster

lesion establishment.

We conducted a comprehensive investigation into the role of the dysregulated ECS in EM

establishment and progression by utilizing CNR1 k/o and CNR2 k/o mouse models. To address

the underlying causes of ECS dysfunction, we induced artificial decidualization in WT, CNR1 k/o,

and CNR2 k/o mice and used the endometrial fragments from decidualized (DD) and

undecidualized (UnD) uterine horns to induce EM in recipient mice of their respective genotypes.

Furthermore, we explored the immunomodulatory potential of the ECS in EM, shedding light on

how alterations in EC signaling may influence immune cell behavior within the localized

peritoneal milieu in mice induced with EM. Our study contributes to the foundational knowledge

around ECS dysregulation in EM and paves way for potential therapeutic strategies targeting ECS

for disease management.

Methods

Animals

Experiments described in this work were approved by the Queen's University Institutional

Animal Care Committee as per the guidelines provided by the Canadian Council of Animal Care.

All animals were assigned randomly to the surgical procedures. All studies were performed using

adult female mice at ages between 7 to 10 weeks. CNR1 k/o (B6.129P2(C)-Cnr1tm1.1Ltz/J) and

CNR2 k/o (B6.129P2-Cnr2tm1Dgen/J) male and female breeder mice were obtained from

Jackson Laboratory (Bar Harbor, USA) and were housed in the Queen's University animal

facility. CNR1 k/o and CNR2 k/o experimental female mice were obtained by trio breeding with

their respective homozygous k/o male counterparts. C57BL/6j (WT) control female mice and

vasectomized male mice at 7 to 10 weeks were obtained from Jackson Laboratory. All breeder

mice were housed in standard breeding cages in a barrier facility and the experimental animals

were housed in a conventional holding area. Animals were housed at constant temperature

$(22 \pm 1 \, ^{\circ}C)$ and relative humidity (50%), with a 12:12 h light:dark cycle (light on 07.00–19.00 h).

Food and water were available *ad libitum*. All experimental animals were acclimatized at the

conventional housing facility for 1 week before starting the experiments.

In vivo **decidualization**

In this study, we have used a modified syngeneic mouse model of EM, where the donor

fragments were obtained from artificially decidualized uterine horns. The method of artificial

decidualization used in this study has been previously established and utilized by several

research studies[11,19,20]. To artificially induce decidualization, female mice were allowed to mate

with vasectomized male mice to induce pseudopregnancy. After day 4 of pseudopregnancy,

female mice were subjected to laparotomy to receive a 30μL injection of sesame seed oil (S3547,

Sigma, USA), intra-luminally into one uterine horn to induce DD. The contralateral, uninjected

horn served as an UnD control. After sesame seed oil injection, animals were rested for 4 days,

after which the DD was successfully induced in one uterine horn as shown in **Figure 1A**. These

uterine horns were utilized as donor fragments to induce EM in recipient mice of their respective

genotype. **Figure 1B** shows the representative images of EM lesions 7 days post-surgical

induction.

Mouse model of EM

EM was surgically induced as described previously[2,17]. Two independent groups (DD and UnD)

per genotype were used in this study (n = 8-16). Briefly, the DD and UnD uterine horns from the

donor mice were harvested, and uterine horns were longitudinally dissected to reveal the

endometrium. Uterine fragments were obtained using a 3.0 mm epidermal biopsy punch (33-32,

Integra™ Miltex®, USA). Recipient mice were anesthetized under 3.5% isoflurane vaporizer

anesthesia to make a midline incision in the abdomen (n = 8-16) and two 3.0 mm DD or UnD

uterine fragments were implanted on the right inner peritoneal wall using a veterinary grade

tissue bonding glue (1469SB, 3M, USA). WT, CNR1 k/o, and CNR2 k/o control groups (n = 4)

were sham operated with a midline incision in the abdomen without implantation of uterine

fragments. Mice were sacrificed 7 days after EM induction surgery since the focus of the study

was on the earlier time point of EM initiation after induction. Based on the previous studies, 7

days after EM induction appears to be the log phase of tissue repair and immune response[21].

Blood was harvested through cardiac puncture to assess EC ligands. Peritoneal fluid (PF) was

collected by injecting 3 ml of ice-cold phosphate buffered saline (PBS) into the peritoneal cavity.

Spleens were collected in ice-cold RPMI media (11875093, ThermoFisher, Canada) before

processing to obtain single cell suspension. EM lesions were either snap frozen in liquid nitrogen

and stored at -80°C or processed using 4% paraformaldehyde overnight (12–20 h), kept at 4°C in

70% ethanol, and then embedded in paraffin.

Lipid extraction and targeted mass spectrometry

Plasma was undiluted and ~ 10mg of tissue per sample were homogenized with RIPA buffer

(89900, ThermoFisher, Canada) with 1:100 of protease inhibitor cocktail (535140-1ML, Sigma,

Canada) to obtain tissue lysates. Both plasma and tissue lysates obtained were individually

subjected to solid phase extraction (SPE). Internal standards, both deuterated and non-deuterated

used in the study to assess the ECS ligands were purchased from Cayman Chemicals, USA

(**Supplementary Table 1**). Aliquots of 100µL of plasma and tissue lysates were added to a

protein precipitation plate (CE0-7565-R, Phenomenex, USA) along with 200µL of cold

acetonitrile containing the deuterated internal standards. The filtrate was diluted with 500µL of

water and submitted to SPE extraction on an Oasis HLB 96-Well Plate (WAT058951, Waters,

Canada). Samples were washed with 60% methanol prior to elution with acetonitrile. The eluate

was dried, reconstituted in 100µL of mobile phase A and analyzed by liquid chromatography–

mass spectrometry (LC-MS as described in **Supplementary Table 2**). The endogenous

concentration for the four compounds in human plasma were calculated by standard addition.

Flow cytometry

Single-cell suspensions were prepared from murine spleens by mechanical dissociation, RBC

lysis, and centrifugation. Splenic cells and cells from PF were resuspended in staining buffer

(PBS with 2% fetal bovine serum) at a concentration of 0.5×10^6 cells/mL. All antibodies used

for flow cytometry analyses were purchased from BioLegend, USA, unless otherwise mentioned.

The antibodies included CD45-FITC (103107), CD3-BV510 (100234), CD4-BV785 (100551),

CD8-BV605 (100744), CD11b-AF700 (101222), F4/80-PE/Cy7 (123114), NK1.1-APC/Cy7

(108724), and CD19-PE/Dazzle 594 (115554). Staining was performed following the

manufacturer's recommendations. For each sample, 50µL of the antibody cocktail was added to

50μL of cell suspension in 96 well plates. The mixture was incubated at 4°C for 20 mins in the

dark, along with anti-CD16/32 Fc block antibody (101319). After incubation, cells were washed

twice with staining buffer and centrifuged before fixing the cells with fixation buffer (00-

8222/49, ThermoFisher, Canada). Flow cytometry analysis was carried out using a Beckman

Coulter CytoFlex S flow cytometer. Compensation controls were established using single

antibody-stained cells. Isotype controls provided baseline levels of non-specific staining and cell

populations were defined using fluorescence minus one (FMO) control. Data analysis employed

FlowJo software (v 10.9; FlowJo, USA) as well as SPECTRE (v 1.0) computational toolkit in R

(v 4.2.3) to obtain t-distributed stochastic neighbor embedding (t-SNE) plots based on the

unsupervised flowSOM clusters generated by marker expression.

***In-vitro* T-cell functional assay**

Total CD3+ T cells were isolated from splenocytes of naive WT and CNR2 k/o mice using a

negative selection magnetic kit (19851A, StemCell, Canada) following the manufacturer's

instructions. All recombinant proteins and compounds were purchased from Biolegend, USA,

unless otherwise mentioned. Subsequently, 250,000 T cells per well were seeded into a 96-well

plate coated with anti-mouse CD3 [(2 μg/ml), (100340)]. RPMI-1640 media supplemented with

rmIL-2 [(10 ng/ml), (575404)], anti-mouse CD28 [(5 μg/ml), (102116)], 10% fetal bovine serum,

β-mercaptoethanol (50μM), and penicillin/streptomycin (100 U/ml) was used as the growth

medium. T cells were then activated non-specifically with or without the cell activation cocktail

consisting of Phorbol 12-myristate 13-acetate (PMA) and ionomycin [(50 ng/ml PMA and 1 μM

ionomycin), (423301)] in the presence and absence of TNFα [(100 ng/ml), (410-MT-010/C,

R&D Systems, USA)] to simulate a sterile inflammatory challenge. Following a 48-h incubation

period, brefeldin A [(10 μg/ml), (11861, Cayman Chemicals, USA)] was introduced to the cells

to measure intracellular interferon-gamma (IFNγ) levels at the 42-h time point. Flow cytometry

analysis was conducted using a panel of markers, purchased from Biolegend, USA unless

otherwise mentioned, including CD3e-FITC (100306), CD4-AF700 (100430), CD8-PE/fire700

(100792), Ki67-PB (151223), FoxP3-PE (126404), IFNγ -BV605 (505840), and Live/dead-

K0525 (L304966, ThermoFisher, Canada), to assess various T cell subsets and viability. Flow

cytometry staining and acquisition was carried out as described above with the addition of

permeabilization (00-8333-56, ThermoFisher, Canada) buffer to stain for intracellular markers

(Ki-67 and IFNγ), according to the manufacturer's instructions. Data analysis was performed

using FlowJo software and visualized using GraphPad Prism (v 9.5.1).

RNA isolation using RNeasy mini kit

Snap frozen UnD and DD EM lesions from WT, CNR1 k/o, and CNR2 k/o were homogenized,

and RNA was isolated using the RNeasy Mini Kit (74104, Qiagen, Canada), according to the

manufacturer's instructions. Briefly, ~20mg EM lesion tissues were placed individually in

PowerBead Ceramic Tubes (13113-50, Qiagen, Canada) along with lysis buffer. Lesions were

homogenized using a Bead Ruptor homogenizer (Omni International, USA) and lysates were

extracted after centrifugation at 10,000 RCF. Lysate was mixed with 70% ethanol, added to the

RNeasy spin column, and then centrifuged to bind the RNA to the column. Spin column was

washed twice, and RNA was isolated using elution buffer. Total RNA quality was measured

using the nanodrop spectrophotometer and stored at -80°C before shipping to BGI Global

(Boston, USA) for bulk RNA analysis. RNA integrity was determined using the Agilent 4150

TapeStation System Agilent, USA) for sample quality control and only samples with RNA

quality number ≥ 7 were considered for library preparation and further sequencing.

RNA library preparation, sequencing, and analysis

Library preparation began with mRNA enrichment using oligo dT beads, which selectively

capture mRNA molecules. Next, the enriched mRNA was fragmented, and first-strand cDNA

was synthesized using random N6 primers, followed by second-strand cDNA synthesis using

deoxyuridine triphosphate (dUTP). After cDNA synthesis, end repair was performed to generate

blunt ends, and 3' adenylation was carried out to facilitate adaptor ligation. Adaptors were ligated

to the 3' adenylated cDNA fragments. To enrich the cDNA library for sequencing, PCR

amplification was conducted. Prior to amplification, the dUTP-marked strand was specifically

degraded by Uracil-DNA-Glycosylase (UDG). The remaining first-strand cDNA was then

amplified using PCR primers. Following amplification, single-strand separation was achieved

through denaturation by heat. The single-stranded DNA was cyclized using a splint oligo and

DNA ligase. DNA nanoball synthesis was performed on the cyclized single-stranded DNA

templates. This process facilitated the generation of clonal DNA clusters, providing the material

necessary for subsequent sequencing. Sequencing was executed using the DNBSEQ Technology

platform. The prepared DNA libraries were loaded onto the DNBSEQ sequencer and sequenced

at an average depth of 30 million paired end reads (2 x 100) per library. The sequencing data was

filtered with SOAPnuke by removing reads containing sequencing adapter; removing reads

whose low-quality base ratio (base quality less than or equal to 15) is more than 20%, and

removing reads whose unknown base ('N' base) ratio is more than 5%. Next, clean reads were

obtained and stored in FASTQ format. Clean reads were mapped to the mouse reference genome

(NCBI: GRCm38.p6) using HISAT2 (v2.0.4)**.** The subsequent analysis and data mining were

performed on Dr. Tom Multi-omics Data mining system (https://biosys.bgi.com).

Imaging Mass Cytometry: labeling

A comprehensive panel of antibodies identifying innate and adaptive immune cell populations

and cell types that are integral to EM lesion microenvironment (**Supplementary Table 3)** was

designed and optimized as previously described[22,23]. The formalin-fixed paraffin-embedded

(FFPE) tissue sections (n = 2-3 per tissue type) underwent deparaffinization and heat-mediated

antigen retrieval on the Ventana Discovery Ultra auto-stainer platform (Roche Diagnostics,

Canada), following the below instructions. Initially, the slides were exposed to a temperature of

70 °C in a pre-formulated EZ Prep solution (Roche Diagnostics, Canada), followed by a

subsequent incubation at 95 °C in pre-formulated Cell Conditioning 1 solution (Roche

Diagnostics, Canada). Following this, the slides were washed in 1× PBS and then exposed to

Dako Serum-free Protein Block solution (Agilent, USA) for 45 min at room temperature. An

antibody cocktail, containing metal-conjugated antibodies, was prepared using Dako Antibody

Diluent (Agilent, USA) at specified dilutions. The primary antibodies within this cocktail were

applied to the slides and left to react overnight at 4 °C, after which the slides were washed with

0.2% Triton X-100 and 1× PBS. For the subsequent step, a secondary antibody cocktail

comprising metal-conjugated anti-biotin antibodies was created in Dako Antibody Diluent, at a

predetermined dilution. The slides were treated with this anti-biotin cocktail for 1-h at room

temperature and then washed with 0.2% Triton X-100 and 1× PBS. For counterstaining, the

slides were exposed to Cell-ID Intercalator-Ir (Fluidigm, Canada) diluted at a ratio of 1:400 in 1×

PBS for 30 min at room temperature. After a 5-min rinse with distilled water, the slides were air-

dried in preparation for imaging mass cytometry (IMC) acquisition. The Hyperion Imaging

System (Fluidigm, Canada) was employed for the IMC acquisition process.

Imaging Mass Cytometry: Data analysis

Lesions (n = 2-3 per tissue type) were grouped based on the origin of uterine fragments (i.e.,

UnD or DD) from three different genotypes (WT, CNR1 k/o, and CNR2 k/o). IMC data analysis

methods employed in this study follow established procedures as outlined in Steinbock toolkit

(Spatial Experiment v 1. 12. 0) for data preprocessing, image segmentation, and object

quantification[24]. Cell segmentation utilized a deep learning approach described by Greenwald et

al[25]. Briefly, dual-channel images were generated using nuclear and cytoplasmic markers,

representing respective signals. The DeepCell tool with Mesmer, a pre-trained deep learning

segmentation algorithm from TissueNet, was used to automate cell mask generation, requiring no

additional user input. Given the IMC data was acquired in batches, we performed batch effect

corrections using the harmony algorithm as described[26]. This involved iterative clustering and

correction of cell positions in the principal component analysis (PCA) space. Subsequently,

unsupervised PhenoGraph clustering in R (v 4. 3. 2) was used to categorize cell types. For this,

signals including αSMA, B220, CD19, β-catenin, CD3, CD4, CD8, CD11b, CD11c, CD31,

CD68, E-cadherin, MPO, pan-cytokeratin, and vimentin were utilized, employing a k-value of

60. To ascertain cell interactions, imcRtools (v 1. 8. 0) and cytomapper (v 1. 14. 0) in R were

employed for visualization. A permutation test evaluated interactions with neighboring cells.

Neighboring cells were defined as those within a 5-pixel radius (5 µm), and the

buildSpatialGraph function established the number of one cluster neighbors interacting with

another cluster. A default of 1000 permutations was set. Each iteration led to interaction score

and p-value computation, and the significant outcomes (at alpha 1% risk) were depicted in

heatmaps. To delineate spatial cellular neighborhoods, neighbor windows were computed,

representing the N nearest cells to each cell. This process followed previous protocols.

Employing imcRtools, cellular neighborhood grouping was conducted, leading to the

identification of 8 cellular neighborhoods in the lesions.

Statistics

Statistical analyses performed to compare the concentration of EC ligands through targeted LC-

MS and evaluation of immune cell population via flow cytometry were conducted using Prism

GraphPad. A one-way analysis of variance (ANOVA) was performed with Holm-Sidak post-hoc

test to determine the specific pairwise differences between the groups. For *in-vitro* T cell

functional assay, two-way ANOVA was performed using Tukey's post-hoc test to compare

within and between the groups. The significance level was set at $\alpha = 0.05$. Data are presented as

mean ± standard deviation (SD) unless otherwise stated.

Results

Ligands of the ECS are dysregulated in a mouse model of EM lacking CNR1 and CNR2

receptors

Based on our previous work demonstrating dysregulated ligands of the ECS in both patients and

our mouse model of EM[2], we first evaluated whether absence of CNR1 or CNR2 led to ECS

ligand alterations. To do this, we performed targeted mass spectrometry on plasma and EM

lesions obtained from CNR1 k/o, CNR2 k/o, and their WT controls. In these mice, EM was

induced using UnD and DD tissues obtained from their respective strains into matched recipients

(for example, UnD and DD from CNR1 k/o donor mice was implanted into CNR1 k/o recipient

mice). We detected some of the major EC ligands such as, 2-Arachidonoylglycerol (2-AG), N-

arachidonoylethanolamine (AEA), Palmitoylethanolamide (PEA) and Oleoylethanolamide

(OEA) in plasma and EM lesions from all genotypes. All identified ECS ligands are

predominantly anti-inflammatory and the range of 2-AG, AEA, PEA, and OEA in the plasma

and lesions were comparable to our previous study[2]. In the plasma, we found no significant

differences in ECS ligands across all groups (**Figure 1C-F**), which could be due to the rapid

homeostasis achieved in circulation[27]. However, in the lesion microenvironment, we captured

higher levels of several EC ligands (**Figure 1G-J**). In CNR1 k/o mice, significantly higher

concentrations of OEA were observed in the DD compared to UnD lesions (**Figure 1J**), and

overall, was on average two-fold higher compared to both lesions from WT and CNR2 k/o mice.

2-AG, which selectively binds to the CNR2 receptor was significantly higher in both the UnD

and DD EM lesions from CNR2 k/o mice (**Figure 1G**) compared to the CNR1 k/o counterparts.

This could indicate a compensatory response in the absence of CNR2. Together, these findings

provide insights into potential dysregulation of ECS ligands in the absence of CNR1 and CNR2

and their involvement in DD vs UnD scenario during EM lesions establishment.

Impact on gene expression and pathway alterations in EM lesions from mice in the absence

of CNR1 and CNR2

Next, we investigated the effects of CNR1 and CNR2 absence on the transcriptomic profile of

both UnD and DD EM lesions from their respective genotypes. Bulk RNA sequencing was

performed on both UnD and DD lesions from WT, CNR1 k/o, and CNR2 k/o mice as detailed

earlier to elucidate the molecular alterations associated with the disruption of these two-receptors

signaling. Differential expression analysis revealed changes in gene expression profiles among

the different genotypes and lesion types (**Figure 2A**). A total of 1100 and 639 differentially

expressed genes (DEGs) were found in both UnD and DD lesions of CNR1 k/o and CNR2 k/o

mice, respectively, compared to WT controls (UnD data is provided in **Supplementary Data 2**).

To gain insights into the biological implications of the observed gene expression changes, we

conducted Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis on

the DEGs identified in UnD and DD lesions of CNR1 k/o and CNR2 k/o mice compared to WT

mice. In the DD lesions from CNR1 k/o mice, KEGG pathway analysis revealed significant

alterations in several pathways (**Figure 2B**). Notably, the cell adhesion molecules pathway was

prominently affected, indicating a potential role for CNR1 in mediating cell-cell interactions and

tissue remodeling processes. Additionally, the cyclic adenosine monophosphate (cAMP)

signaling emerged as another negatively impacted pathway, implicating CNR1 in modulating

intracellular signaling cascades. In DD lesions from the CNR2 k/o mice, analysis highlighted

distinct pathways affected in the context of inflammation and EM (**Figure 2C**) including the

cytokine receptor interactions pathway, pointing to the involvement of CNR2 in immune

responses and inflammatory processes associated with EM. Furthermore, we captured alterations

in the steroid hormone biosynthesis pathway suggesting a role for CNR2 in hormone-related

mechanisms relevant to endometrial tissue development and homeostasis.

Next, we performed a subset analysis for genes directly involved in ECS signaling. A total of 59

key genes in ECS signaling were selected, as identified in a study by Tanaka et al[28]. Surprisingly,

despite the central role of CNR1 and CNR2 in ECS signaling, we found a limited number of

DEGs related to this system. Out of 59 genes directly associated with ECS, only 3 (*CNR1,*

PLCH1 and *PLCH2*) and 2 (*CNR2* and *PLAG2GE*) DEGs were identified in the DD lesions of

CNR1 k/o (**Figure 2D**) and CNR2 k/o (**Figure 2E**), respectively, compared to DD lesions from

WT controls. This result suggests that CNR1 and CNR2 modulate the EM microenvironment

through intricate interactions with other signaling pathways beyond the canonical ECS pathway.

A comprehensive gene ontology (GO) classification analysis on the 59 identified ECS genes

(**Figure 2F**) unveiled their multifaceted roles in reproductive functions, immune system

regulation, and cellular processes. The genes exhibited enrichment in molecular functions such

as receptor activity and lipid binding. In terms of cellular components, these genes were

associated with plasma membrane structures, and intracellular compartments, signifying their

diverse subcellular localization and potential involvement in dynamic cellular processes.

Furthermore, GO analysis highlighted their participation in biological processes such as immune

response modulation, lipid metabolism, cell communication, and intracellular signaling

pathways, indicating the broader impact of ECS beyond its canonical functions. The results of

the UnD comparisons between the genotypes are provided in the supplementary files

(**Supplementary Figure 1**). While the UnD lesions exhibited distinct gene expression patterns

compared to DD lesions, common trends in the effects of CNR1 k/o and CNR2 k/o on gene

expression were observed across both lesion types. Specifically, the cytokine receptor

interaction, complement cascade, and inflammatory mediator pathways were significantly altered

in the UnD lesions from CNR1 k/o and CNR2 k/o mice compared to UnD lesions from WT

mice. Overall, our findings highlight the significant impact of CNR1 and CNR2 k/o on gene

expression in EM lesions and the implications for EM pathogenesis.

Disruption of genes related to adaptive immune response in EM lesions without CNR1 and

CNR2

Building upon our previous investigation into the transcriptomic alterations, we conducted an in-

depth analysis of differentially expressed immune-related genes (as per InnateDB version 5.4) in

both UnD and DD lesions across all genotypes. Here, our analysis is focused on the immune-

related genes within DD lesions of CNR1 k/o and CNR2 k/o mice compared to WT controls.

Comparison of the UnD lesions of CNR1 k/o and CNR2 k/o with WT EM mice are included in

the supplementary files (**Supplementary Figure 2**) The differential expression bar plot (**Figure**

3A) provides representation of the upregulated and downregulated genes in each comparison.

The volcano plots for DD lesions from CNR1 k/o vs. WT (**Figure 3B**) and CNR2 k/o vs. WT

(**Figure 3C**) illustrate the various DEGs. Notably, CNR1 k/o DD lesions exhibited 39

downregulated (eg., *NLRP6* and *IL1a,* pivotal regulators of inflammatory response) and 14

upregulated genes (eg., *CXCL9* and *CXCL10,* chemokines involved in immune cell recruitment),

while CNR2 k/o DD lesions showed 40 downregulated (eg., *SIGLECG* and *IL6,* involved in

immune regulation and pro-inflammatory response) and 25 upregulated genes (eg., *C8a*, *C9,* and

MASP2, part of the complement system), highlighting substantial changes in gene expression

associated with CNR1 and CNR2 disruption compared to the DD lesions from WT mice. Of

particular interest, we observed significant downregulation of T cell-related genes (*CD3e*, *CD3g*,

GATA3, and *CTLA4*) in the CNR2 k/o DD lesions (**Supplementary Data 3**), aligning with the

CD3+ T cell dysfunction observed in the PF and spleen and further validations from *in-vitro*

functional assay (mentioned below). However, we did not find the same differences (T cell-

related genes) in the UnD lesions of CNR2 k/o mice. Moreover, UnD lesions of CNR2 k/o mice

showed significantly low number of DEGs (11 compared to 65 in the DD lesions from CNR2 k/o

mice) suggesting a decidualization dependent response (**Supplementary Data 3**). This

observation clearly emphasises a potential link between CNR2 dysfunction with decidualization

characterized by T cell signaling issues within the EM microenvironment. To understand

functional implications of the DEGs, we conducted KEGG pathway analysis on specific

differentially expressed immune genes in DD lesions from CNR1 k/o and CNR2 k/o mice.

Notably, in CNR1 k/o DD lesions compared to WT DD, the chemokine signaling pathway,

cytokine-cytokine receptor interaction, and toll-like receptor (TLR) signaling pathways were

negatively affected (**Figure 3D**). These findings provide insights into the impact of CNR1

disruption on immune cell function within the DD environment. Conversely, CNR2 k/o DD

lesions were associated with significant alterations in cytokine-cytokine receptor interaction,

Th17, Th1, and Th2 cell differentiation pathways (**Figure 3E**). These pathways are proven to be

involved in T cell development, differentiation, and effector functions, aligning with our

observed dysregulation of T cell-related genes. Together, these findings further elucidate the

roles of CNR1 and CNR2 in modulating immune responses within the context of EM.

Multispectral flow cytometry revealed altered immune cell profiles in a mouse model of

EM lacking CNR1 and CNR2

Immune dysregulation is recognized as a crucial factor in the pathogenesis of EM[29]. To elucidate

the impact of CNR1 and CNR2 absence on immune cell populations, we performed multispectral

immune profiling in splenocytes and cells from the PF from WT, CNR1 k/o, and CNR2 k/o mice

harboring UnD and DD lesions. Representative gating panels of PF CD3 (y axis) vs CD11b (x

axis) cells (**Figure 4A-E**) illustrate distinct profiles among WT, CNR1 k/o, and CNR2 k/o mice

with EM, as well as CNR2 k/o naïve, non-operated mice, and CNR2 k/o sham-operated controls.

Strikingly, CD3+ T cell populations were nearly absent in the PF of CNR2 k/o mice with EM,

regardless of lesion types (UnD and DD), when compared to other groups (**Figure 4C and F**).

This trend further extended to CD4+ helper T cells and CD8+ cytotoxic T cells (**Figure 4G-H**).

CNR1 k/o mice with DD lesions exhibited significantly reduced CD3+ (**Figure 4F**), CD4+

(**Figure 4G**), and CD8+ (**Figure 4H**) T cell frequencies compared to their UnD counterparts, as

well as lower CD19+ B cells and NK1.1+ NK cells (**Figure 4K**) populations, compared to WT

and CNR2 k/o mice. Concomitant with the reduction of T cell subsets, an increase in CD11b

monocytes was observed in the PF of CNR2 k/o mice with UnD and DD lesions compared to

WT and CNR1 k/o mice (**Figure 4I**). Similarly, CNR1 k/o mice with DD lesions displayed

higher monocyte/macrophage populations compared to their UnD counterparts (**Figure 4I**).

Furthermore, WT mice with DD lesions demonstrated significantly lower CD3+ T cell

frequencies compared to their UnD counterparts (**Figure 4F**), suggestive of a decidualization-

associated effect. Splenocytes exhibited analogous trends (**Figure 4L**), as depicted by tSNE plots

(bar plots and contour gating plots in **Supplementary Figure 3 and 4**). These alterations in

immune cell numbers reinforce the influence of CNR1 and CNR2 dysregulation and

decidualization on immune cell populations, confirmed both locally in PF and systemically in the

spleen.

T cells from CNR2 k/o mice exhibit impaired viability upon TCR activation

To validate the functional consequences of CNR2 deficiency on T cell behavior, we conducted a

series of *in-vitro* assays using T cells isolated from splenocytes of naïve WT and CNR2 k/o

mice. CD3+ T cells were activated non-specifically, with or without PMA/ionomycin cocktail, in

the presence or absence of tumor necrosis factor alpha (TNFα) to create a sterile inflammatory

challenge.

We observed a significant reduction in the viability of total CD3+ T cells from CNR2 k/o mice

upon activation with PMA/ionomycin (gating strategies in **Supplementary Figure 5**) compared

to media controls (**Figure 5A** and **B**). In contrast, WT CD3+ T cells activated with

PMA/ionomycin, with or without TNFα, exhibited no significant difference in viability when

compared to the media control (**Figure 5A** and **B**). This observation aligns with our *in-vivo*

findings, whereby CD3+ T cells from CNR2 k/o mice with EM exhibited a significant reduction

in both the splenic and PF population, but not in the SHAM-operated mice, emphasizing the EM-

specific nature of this effect. Additionally, the overall reduced viability of CD3+ T cells of

CNR2 k/o mice upon PMA/ionomycin activation led to decrease in proliferation of CD4+ T cells

(**Figure 5C**) but not of CD8+ T cells (**Figure 5D**). However, our results indicated that although

CNR2-deficient T cells displayed reduced viability upon activation, they exhibited higher levels

of IFNγ production compared to CD3+ T cells from WT mice, suggesting their functional

competence (**Supplementary Figure 5**). These findings shed light on the intricate role of CNR2

in modulating T cell responses, with potential implications for immune dysregulation.

Spatial cell analysis of EM lesions from WT, CNR1 k/o, and CNR2 k/o mice reveal altered

immune cell populations

To gain a comprehensive insight into the spatial distribution of immune cells and stromal cell

types within EM lesion architecture of UnD and DD lesions of WT, CNR1 k/o, and CNR2 k/o

mice, we employed IMC analysis. This approach aimed to elucidate the impact of CNR1 and

CNR2 absence on the cellular composition and organization of EM lesions in mice. The

schematic representation of the IMC procedure (**Figure 6A**) outlines the steps involved in this

analysis. Following the acquisition of two regions of interest (ROI) per section (based on the

H&E stain), single-cell segmentation (**Figure 6B**) and subsequent segmentation quality

assessment (**Figure 6C**) were performed. After batch effect correction of samples, non-linear

dimensionality reduction of the sample type revealed a distinct expression pattern of immune

cells and cell state markers between UnD and DD lesions, as well as differences between the

genotypes (**Figure 6D**). After unsupervised phenotyping and labeling of the different cell types,

uniform manifold approximation and projection (UMAP) dimensionality reduction further

highlighted the key differences in cell composition between UnD and DD lesions (**Figure 6E**).

Overall, combined expression of the cell types of DD lesions from all three genotypes exhibited

increased stromal compartments, decreased epithelial cells, and heightened macrophage

infiltration compared to the expression of cell types from UnD lesions. Representative images

illustrate the distribution of different cell types based on unsupervised clustering and labeling

(**Figure 7A**).

Further analysis of the cell type distribution (**Figure 7B**) through bar plots unveiled several

differences. Although not significant, due to relatively low number of biological replicates, T cell

expression was increased (CD4+ helper T cells and CD8+ cytotoxic T cells) in both UnD and

DD lesions of CNR1 k/o and CNR2 k/o mice compared to WT (**Figure 7C**). This observation

highlights that resident T cells are not impacted in the absence of CNR1 and CNR2 within the

endometriotic milieu. Intriguingly, in EM lesions from CNR1 k/o and CNR2 k/o, we saw

significantly elevated expression of monocytes/macrophages (**Figure 7D**), stromal cells (**Figure**

7E) and hallmarks of EM such as proliferation and vascularization (**Figure 7F**) demonstrating an

altered microenvironment in the absence of these receptors. To comprehend cell-cell interactions

and their implications, we conducted cellular neighborhood (CN) analysis. This approach

grouped cells based on information within their direct spatial vicinity and identified intricate

spatial relationships among diverse cell types within the lesion microenvironment. This analysis

revealed distinct clustering patterns across different cell types within the lesion architecture of

the DD lesions from WT, CNR1 k/o, and CNR2 k/o mice (**Supplementary Figure 6**). Immune

cells predominantly clustered together in CN 4, while other cell types (stroma, epithelial cells,

and vasculature) exhibited distinct clustering patterns across CN 6, 3, and 8 in DD lesions

(**Figure 7G**). Although most of the immune cell types clustered together in the UnD lesions, cell

types of the lesion architecture clustered distinctly compared to the DD lesions (**Supplementary**

Figure 6). This clustering emphasizes the interplay between immune cells and the broader

cellular components of the lesions. In summary, our comprehensive investigation has unveiled

intricate spatial relationships among immune cells and diverse cell types within EM lesions in

mice. The observed alterations in T cell expression, coupled with stromal dynamics, in CNR1

k/o and CNR2 k/o lesions underscore the pivotal roles of these receptors in shaping the

endometriotic microenvironment.

Discussion

Emerging evidence from our previously reported findings and others, implicated dysregulation of

ECS, comprising CNR1 and CNR2 canonical receptors along with their EC ligands, in EM

pathophysiology[2,30]. The ECS is involved in several physiological processes including (but not

limited to) pain perception, immune regulation, and reproductive functions[31]; CNR1 and CNR2

are expressed in immune cells, nerve tissues, and serve as critical regulators of reproductive

processes including decidualization and embryo implantation[32,33]. The etiology of EM has been

speculated to be routed in defective decidualization and retrograde menstruation of the

endometrial fragments combined with ECS dysregulation[8,9,34]. While it is plausible that decreased

ECS function influences EM lesion initiation, progression, and severe pain experience, it is not

clear whether ECS dysfunction actively contributes to EM pathogenesis, or whether it represents

a secondary consequence of alterations occurring within the refluxed endometrial tissue, leading

to establishment of EM lesions.

Keeping this central dogma in view and to provide insights into early events of EM pathogenesis,

we induced EM in both CNR1 k/o and CNR2 k/o mice utilizing syngeneic, DD and UnD uterine

endometrial fragments. Absence of CNR1 and CNR2 did not influence systemic levels of ECS

ligands but the lesion microenvironment displayed significant changes in the levels of OEA and

PEA, suggesting a tissue-specific response.

One intriguing aspect of ECS involvement in EM is its role in decidualization, a process pivotal

for uterine receptivity to embryo implantation and successful pregnancy, that may also contribute

to EM establishment[35]. Although both CNR1 and CNR2 are active in decidualization, CNR1

may have a more prominent role. Absence of CNR1 and CNR2 show compromised

decidualization in mice in a CNR1-dependent manner validated through *in-vitro* studies[36].

Similarly, in our study, EM lesions (both UnD and DD) from mice lacking CNR1 showed

significantly more DEGs (2088), compared to CNR2 (287) and WT (2). Genes essential for

decidualization such as *IGFBP2*, *BMP3*, *PTGDR*, *WNT7a,* and *ESR1* were downregulated in the

DD, EM lesions from CNR1 k/o mice compared to their UnD counterparts. This further

reinforces the role of CNR1 in the uterine and EM lesion microenvironment, including their role

in decidualization response. Moreover, the interplay between CNR1 and CNR2 are important

since CNR2 contribute to immunomodulation, which is a key process during decidualization[37,38].

Even though retrograde menstruation is considered the main mechanism by which endometrial

fragments reach the peritoneal cavity and implant to form EM lesions, the retrograded menstrual

debris itself does not undergo decidualization. However, some EM lesions in the peritoneal

cavity, under the influence of estrogen and progesterone, can undergo decidualization as the

lesions can exist in different evolutionary stages, from active red lesions to quiescent white

lesions[39].

Given the complexity of ECS signalling and compensatory mechanisms, we focused our

investigation on the immune dysregulation aspect of EM pathophysiology. Our findings suggest

that altered ECS dynamics during decidualization disrupt ECS signaling, leading to

dysregulation of immune responses and aberrant cellular behavior. Indeed, immune dysfunction

is a hallmark of EM[29,40], and ECS could play a crucial role in shaping immune responses

particularly through its impact on T cell function yet there is no clear evidence. Alterations in T

cell populations and functions have been associated with EM progression, suggesting their vital

role in EM pathogenesis and maintenance[41,42]. Our flow cytometry analysis revealed significant

alterations in immune cell populations in mice bearing EM lesions, with a notable absence of

CD3+ T cells, CD4+ helper T cells, and CD8+ cytotoxic T cells specifically in CNR2 k/o mice.

Mechanistic *in-vitro* studies further confirmed an aberrant T cell response in CNR2 k/o mice, as

with T cell receptor activation and stimulation there was decreased viability. Combined, our

findings show that CNR2 is critical in T cell survival upon TCR activation with pathogen-

associated molecular patterns (PAMPs)/danger-associated molecular patterns (DAMPs) or

antigen-mediated signals. These findings also shed light on a previously unrecognized role of

CNR2 in EM-associated adaptive immune dysfunction given the critical role of T cells in

immune surveillance and regulation. Furthermore, speculation of EM being a cause of ECS

dysfunction could be of importance since CNR2 was found to be reduced in the lesions of EM

patients, as shown by our previous study[2]. In addition, the bulk RNA sequencing strengthens the

finding of dysregulation of T cell-related genes in CNR2 k/o EM lesions. The observed

downregulation of T cell-related genes such as *CD3e*, *CD3g*, *GATA3*, and *CTLA4* is consistent

with the diminished CD3+ T cell populations and highlights the relevance of CNR2 in T cell-

mediated immune responses within the endometriotic microenvironment.

KEGG pathway analysis of differentially expressed immune-related genes in CNR2 k/o DD

lesions further revealed that Th1, Th2, and Th17 cell differentiation pathways were impacted,

and the previous literature confirms dysregulation of these pathways in EM pathophysiology[43,44].

Additionally, our *in-vitro* functional assay showing CD4+ helper T cells being affected

(proliferation) more than the CD8+ cytotoxic T cell subset further adds to the subset specific

behaviour of CNR2. Our results provide a missing link between the ECS and immune system

functioning during EM.

To gain insights into the early features of lesion initiation and establishment, we conducted IMC

analysis of EM lesions across genotypes and different lesion types (DD and UnD). DD lesions

from CNR1 and CNR2 k/o mice showed higher T cell residing in the lesions with increased

stromal compartments and monocytes/macrophages population compared to WT lesions. Stromal

cells contribute to the early development of EM lesions by promoting inflammation,

angiogenesis, fibrosis, and immune modulation[45–47]. Additionally, the interaction between

macrophages and stromal cells is important in EM, with the NLRP3 inflammasome playing a

role in lesion development[48]. Studies have implicated macrophages in EM lesion growth where

they support angiogenesis (formation of blood vessels) by producing pro-angiogenic factors such

as vascular endothelial growth factor (VEGF)[49–51]. Given the combined increase in proliferation,

endothelial marker, and monocytes/macrophages in EM lesions from mice potentially indicates

that they could modulate the early lesion microenvironment in the event of CNR1 and CNR2

dysregulation. Based on the proportion of these macrophages to certain phenotypes, such as M1

or M2, would dictate lesion development and subsequent progression. Further studies are

required to tease out molecular interactions of CNR1 and CNR2 with specific immune cell

subsets in a complex EM lesion microenvironment and determine how it contributes to

establishment of blood supply and lesion survival.

The dysregulation of the ECS, as evidenced by the altered expression of CNR1 and CNR2 in our

mouse model of EM, appears to have far-reaching consequences on the cellular and molecular

landscape of endometriotic lesions. Since CNR1 is widely expressed in the central nervous

system where it regulates pain perception, mood, and other neurological processes, the

dysregulation of CNR1 signaling may impact the sensory innervation and pain responses

associated with the EM[31]. Additionally, CNR1 has been shown to modulate immune cell

function, including the production of pro-inflammatory cytokines[52]. The loss of CNR1 signaling

in immune cells infiltrating the endometriotic lesions likely contribute to the altered immune

profiles observed in our study. In contrast, CNR2 is predominantly expressed in immune cells,

such as macrophages, lymphocytes, and natural killer cells[53]. This receptor plays a crucial role in

the regulation of immune responses, including the modulation of cytokine production, cell

migration, and proliferation. The significant reduction in CD3+ T cells in the peritoneal cavity

and spleen of CNR2 knockout mice suggests that the loss of CNR2 may have a profound impact

on T cell homeostasis and function in EM dependent manner, as this was not the case in sham

operated mice. This impairment in T cell function has direct consequences on the adaptive

immune response, as evidenced by the altered gene expression profiles and pathways related to

immune function in our transcriptomic analysis. But we haven't conducted specific mechanistic

experiments to tease out interplay between immune response and endometriosis lesion

development in our model systems.

Several limitations should be acknowledged in our study. Firstly, understanding the homeostatic

aspects of ECS, both with and without the presence of CNR1 and CNR2, remains a complex

challenge. While our global k/o mouse models provide valuable insights, further research

utilizing targeted k/o specific to the uterus could offer a more precise understanding of their

contributions to uterine physiology and further implications in EM establishment. The use of

mouse models to study EM has inherent limitations due to species differences and the inability to

fully recapitulate the human disease. The molecular and cellular complexities associated with

DD and UnD endometrial tissues, as well as the timing of EM induction in these mice, may not

perfectly mirror the human condition. No significant baseline alterations were observed in the

immune profile between the CNR1 and CNR2 k/o and the WT mice without EM induction.

However, the uterine environment has not been assessed to understand the baseline immune

profile between the k/o and WT mice. These limitations emphasize the need for future

investigations to enhance the translational relevance of our findings and further our

understanding of the complex interplay between ECS, decidualization, and EM pathogenesis.

In conclusion, our study offers evidence for the involvement of CNR1 and CNR2 dysregulation

in EM pathogenesis. Through an integrative analysis of transcriptomic profiles, immune cell

dynamics, and spatial relationships within EM lesions from mice, we unveil the intricate

interactions between ECS, immune responses, and cellular changes in EM. By identifying

potential mechanisms through which ECS disruption could impact EM, our research provides a

foundation for the development of targeted therapies addressing the ECS's influence on EM.

These findings will advance our understanding of EM and lead to innovative therapeutic

strategies to manage this complex disorder.

Data availability

Bulk mRNA sequencing data is provided in the supplementary files (**Supplementary Data 4**)

and IMC data generated in this study has been deposited in Mendeley Data

(https://data.mendeley.com/datasets/2ptns5yhzh/2).

Code availability

All the codes used in the current study are from previously published articles, as cited in the

article. Authors do not report original code.

Acknowledgements

We thank Dr Alexandra Furtos and Karine Gilbert from the Regional Mass Spectrometry Centre

at Université de Montréal for designing and performing mass-spectrometry evaluation; Dr

Yuhong Wei and the Single Cell and Imaging Mass Cytometry Platform (SCIMAP) at McGill

University Goodman Cancer Institute for processing, labeling, and acquiring samples for IMC

imaging; Brittney Armitage-Brown from the Animal Care Services at Queen's University for

breeding mice utilized in this study. This work was supported by funding from the Canadian

Institutes of Health Research (CIHR-394340) to C. T and M. K.

Author information

Authors and Affiliations

Contributions

H. L., and C. T. designed and conceived experiments; H. L., K. Z., Y. W., P. Y., D. S., and A.

M., conducted experiments. H. L., K. Z., and P. Y., analyzed results; H. L., and C. T., wrote the

manuscript, generated the figures, and wrote the figure legend. C. T., and M. K., supervised the

project and obtained research funding. All authors discussed the results and commented on the

manuscript.

Competing interests

The authors declare no competing interests.

References

1. Giudice, L. C. & Kao, L. C. Endometriosis. TL - 364. *Lancet* (2004) doi:10.1016/S0140-6736(04)17403-5.
2. Lingegowda, H. *et al.* Implications of dysregulated endogenous cannabinoid family members in the pathophysiology of endometriosis. *F S Sci* **2**, 419–430 (2021).
3. Bilgic, E. *et al.* Endocannabinoids modulate apoptosis in endometriosis and adenomyosis. *Acta Histochem* **119**, 523–532 (2017).
4. Shen, X. *et al.* Decreased expression of cannabinoid receptors in the eutopic and ectopic endometrium of patients with adenomyosis. *Biomed Res Int* **2019**, 1–8 (2019).
5. Di Marzo, V., Bifulco, M. & De Petrocellis, L. The endocannabinoid system and its therapeutic exploitation. *Nat Rev Drug Discov* **3**, 771–784 (2004).
6. Zou, S. & Kumar, U. Cannabinoid receptors and the endocannabinoid system: Signaling and function in the central nervous system. *Int J Mol Sci* **19**, (2018).
7. Russo, E. B. Clinical Endocannabinoid Deficiency Reconsidered: Current Research Supports the Theory in Migraine, Fibromyalgia, Irritable Bowel, and Other Treatment-Resistant Syndromes. *Cannabis Cannabinoid Res* **1**, 154–165 (2016).
8. Lingegowda, H. *et al.* Role of the endocannabinoid system in the pathophysiology of endometriosis and therapeutic implications. *J Cannabis Res* **4**, 1–13 (2022).
9. Sampson, J. A. Peritoneal endometriosis due to the menstrual dissemination of endometrial tissue into the peritoneal cavity. *Am J Obstet Gynecol* **14**, 422–469 (1927).
10. Gellersen, B. & Brosens, J. J. Cyclic Decidualization of the Human Endometrium in Reproductive Health and Failure. *Endocr Rev* **35**, 851–905 (2014).
11. Cai, X. *et al.* Endometrial stromal PRMT5 plays a crucial role in decidualization by regulating NF-κB signaling in endometriosis. *Cell Death Discovery 2022 8:1* **8**, 1–12 (2022).
12. Tomassetti, C. & D'Hooghe, T. Endometriosis and infertility: Insights into the causal link and management strategies. *Best Pract Res Clin Obstet Gynaecol* **51**, 25–33 (2018).
13. Zutautas, K. B. *et al.* The dysregulation of leukemia inhibitory factor and its implications for endometriosis pathophysiology. *Front Immunol* **14**, (2023).
14. Wang, H., Xie, H. & Dey, S. K. Loss of Cannabinoid Receptor CB1 Induces Preterm Birth. *PLoS One* **3**, e3320 (2008).
15. Li, Y., Bian, F., Sun, X. & Dey, S. K. Mice Missing Cnr1 and Cnr2 Show Implantation Defects. *Endocrinology* **160**, 938–946 (2019).
16. Sun, X. *et al.* Endocannabinoid signaling directs differentiation of trophoblast cell lineages and placentation. *Proc Natl Acad Sci U S A* **107**, 16887–16892 (2010).
17. Lingegowda, H. *et al.* Synthetic Cannabinoid Agonist WIN 55212-2 Targets Proliferation, Angiogenesis, and Apoptosis via MAPK/AKT Signaling in Human Endometriotic Cell Lines and a Murine Model of Endometriosis. *Frontiers in Reproductive Health* **0**, 63 (2021).

662 663	18.	Leconte, M. *et al.* Antiproliferative effects of cannabinoid agonists on deep infiltrating endometriosis. *American Journal of Pathology* **177**, 2963–2970 (2010).
664 665 666	19.	Benson, G. V. *et al.* Mechanisms of reduced fertility in Hoxa-10 mutant mice: Uterine homeosis and loss of maternal Hoxa-10 expression. *Development* **122**, (1996).
667 668 669	20.	Liu, M. *et al.* Menin directs regionalized decidual transformation through epigenetically setting PTX3 to balance FGF and BMP signaling. *Nat Commun* **13**, (2022).
670 671	21.	Symons, L. K. *et al.* Neutrophil recruitment and function in endometriosis patients and a syngeneic murine model. *FASEB J* (2020) doi:10.1096/fj.201902272R.
672 673 674	22.	McDowell, S. A. C. *et al.* Neutrophil oxidative stress mediates obesity-associated vascular dysfunction and metastatic transmigration. *Nature Cancer 2021 2:5* **2**, 545–562 (2021).
675 676 677	23.	Sorin, M. *et al.* Original research: Single-cell spatial landscape of immunotherapy response reveals mechanisms of CXCL13 enhanced antitumor immunity. *J Immunother Cancer* **11**, 5545 (2023).
678 679	24.	Windhager, J., Bodenmiller, B. & Eling, N. An end-to-end workflow for multiplexed image processing and analysis. *bioRxiv* (2021).
680 681 682	25.	Greenwald, N. F. *et al.* Whole-cell segmentation of tissue images with human-level performance using large-scale data annotation and deep learning. *Nat Biotechnol* **40**, (2022).
683 684	26.	Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with Harmony. *Nat Methods* **16**, (2019).
685 686	27.	Lu, H. C. & MacKie, K. An introduction to the endogenous cannabinoid system. *Biol Psychiatry* **79**, 516–525 (2016).
687 688	28.	Tanaka, K. *et al.* Gene expression of the endocannabinoid system in endometrium through menstrual cycle. *Sci Rep* **12**, (2022).
689 690	29.	Symons, L. K. *et al.* The Immunopathophysiology of Endometriosis. *Trends Mol Med* **24**, 748–762 (2018).
691 692 693	30.	Sanchez, A. M. *et al.* Elevated Systemic Levels of Endocannabinoids and Related Mediators Across the Menstrual Cycle in Women with Endometriosis. *Reproductive Sciences* **23**, 1071–1079 (2016).
694 695	31.	de Fonseca, F. R. *et al.* The endocannabinoid system: Physiology and pharmacology. *Alcohol and Alcoholism* **40**, 2–14 (2005).
696 697	32.	Di Blasio, A. M., Vignali, M. & Gentilini, D. The endocannabinoid pathway and the female reproductive organs. *J Mol Endocrinol* **50**, R1–R9 (2012).
698 699 700	33.	Walker, O. L. S., Holloway, A. C. & Raha, S. The role of the endocannabinoid system in female reproductive tissues. *Journal of Ovarian Research* Preprint at https://doi.org/10.1186/s13048-018-0478-9 (2019).
701 702 703 704	34.	Maia, J., Fonseca, B. M., Teixeira, N. & Correia-Da-Silva, G. The fundamental role of the endocannabinoid system in endometrium and placenta: Implications in pathophysiological aspects of uterine and pregnancy disorders. *Hum Reprod Update* **26**, (2020).

35.   Correa, F., Wolfson, M. L., Valchi, P., Aisemberg, J. & Franchi, A. M.
Endocannabinoid system and pregnancy. *Reproduction* **152**, R191–R200 (2016).

36.   Li, Y., Dewar, A., Kim, Y. S., Dey, S. K. & Sun, X. Pregnancy success in mice requires
appropriate cannabinoid receptor signaling for primary decidua formation. *Elife* **9**,
1–29 (2020).

37.   Turcotte, C., Blanchet, M. R., Laviolette, M. & Flamand, N. The CB2 receptor and its
role as a regulator of inflammation. *Cellular and Molecular Life Sciences* **73**, 4449–
4470 (2016).

38.   Taylor, A. H. *et al.* Endocannabinoids and pregnancy. *Clinica Chimica Acta* **411**,
921–930 (2010).

39.   Maggiore, U. L. R. *et al.* A systematic review on endometriosis during pregnancy:
Diagnosis, misdiagnosis, complications and outcomes. *Hum Reprod Update* **22**,
(2016).

40.   Ahn, S. H. *et al. Pathophysiology and Immune Dysfunction in Endometriosis*.
BioMed Research International vol. 2015 (Hindawi Publishing Corporation, 2015).

41.   Tanaka, Y. *et al.* Exacerbation of endometriosis due to regulatory t-cell dysfunction.
Journal of Clinical Endocrinology and Metabolism (2017) doi:10.1210/jc.2017-
00052.

42.   Kyama, C. M., Debrock, S., Mwenda, J. M. & D'Hooghe, T. M. Potential involvement
of the immune system in the development of endometriosis. *Reproductive Biology*
and Endocrinology Preprint at https://doi.org/10.1186/1477-7827-1-123 (2003).

43.   Chang, L. Y., Shan, J., Hou, X. X., Li, D. J. & Wang, X. Q. Synergy between Th1 and
Th2 responses during endometriosis: A review of current understanding. *Journal of*
Reproductive Immunology vol. 158 Preprint at
https://doi.org/10.1016/j.jri.2023.103975 (2023).

44.   Andreoli, C. G. *et al.* T helper (Th)1, Th2, and Th17 interleukin pathways in infertile
patients with minimal/mild endometriosis. *Fertil Steril* **95**, 2477–2480 (2011).

45.   McKinnon, B. D. *et al.* Altered differentiation of endometrial mesenchymal stromal
fibroblasts is associated with endometriosis susceptibility. *Communications Biology*
2022 5:1 **5**, 1–14 (2022).

46.   Queckbörner, S., Syk Lundberg, E., Gemzell-Danielsson, K. & Davies, L. C.
Endometrial stromal cells exhibit a distinct phenotypic and immunomodulatory
profile. *Stem Cell Res Ther* **11**, 1–15 (2020).

47.   Zondervan, K. T. *et al.* Endometriosis. *Nature Reviews Disease Primers 2018 4:1* **4**,
1–25 (2018).

48.   Guo, X. *et al.* NLRP3 Inflammasome Activation of Mast Cells by Estrogen via the
Nuclear-Initiated Signaling Pathway Contributes to the Development of
Endometriosis. *Front Immunol* **12**, 749979 (2021).

49.   Ding, Y., Song, N. & Luo, Y. Role of bone marrow-derived cells in angiogenesis:
Focus on macrophages and pericytes. *Cancer Microenvironment* **5**, 225–236
(2012).

50.   Sun, H. *et al.* Macrophages alternatively activated by endometriosis-exosomes
contribute to the development of lesions in mice. *MHR: Basic science of*
reproductive medicine **25**, 5–16 (2019).

51. McLaren, J. *et al.* Vascular endothelial growth factor is produced by peritoneal fluid
macrophages in endometriosis and is regulated by ovarian steroids. *Journal of*
Clinical Investigation **98**, 482 (1996).
52. Klein, T. W. Cannabinoid-based drugs as anti-inflammatory therapeutics. *Nature*
Reviews Immunology Preprint at https://doi.org/10.1038/nri1602 (2005).
53. Turcotte, C., Blanchet, M. R., Laviolette, M. & Flamand, N. The CB2 receptor and its
role as a regulator of inflammation. *Cellular and Molecular Life Sciences* **73**, 4449
(2016).

Figure 1: Characterization of endocannabinoid ligands in a modified syngeneic mouse model of endometriosis

A Overview of the modified syngeneic mouse model of EM where pseudo-pregnant WT, CNR1 k/o, and CNR2 k/o mice were induced with DD by injecting sesame oil into the lumen of one uterine horn and the contralateral horn served as UnD control. Two, 3mm UnD and DD harvested fragments were implanted into their respective recipient mouse strain to induce EM. **B** Representative images of the EM lesions from WT, CNR1 k/o, and

 CNR2 k/o mice retrieved from the peritoneal cavity at end point (7 days post EM
induction surgery). **C-J** Bar plots (mean ± SD) showing the concentration of EC ligands
2AG, AEA, PEA, and OEA identified in the plasma and EM lesions from mice using
targeted LC-MS approach. **C-F** 2AG, AEA, PEA, and OEA were detected in plasma
samples without any significant differences between groups. **G, J** Significantly higher
concentration of 2AG was observed between the DD lesions of CNR2 k/o and CNR1 k/o
mice, and significantly higher levels of OEA in the DD lesions from CNR1 k/o mice
compared to DD lesions from WT mice. **H, I** AEA and PEA levels in the tissue samples
did not differ significantly between the comparison groups. *n* = 4-5 individual biological
samples per genotype. Statistical analyses were performed using the ordinary one-way
ANOVA with Holm-Sidak post hoc test. * *p* < 0.05 and ** *p* < 0.01.

779
Figure 2: Transcriptomic profiling of endometriosis-like lesions from CNR1 and
CNR2 knockout mice reveals extensive differential gene expression and altered
pathways
A Summary of the differentially expressed genes (DEGs) from bulk RNA sequencing
analysis conducted on both UnD and DD lesions from WT, CNR1 k/o, and CNR2 k/o
mice, revealing extensive changes in gene expression profiles among the different
genotypes and lesion types. A total of 1100 and 639 DEGs were identified in both UnD
and DD lesions of CNR1 k/o and CNR2 k/o mice, respectively, compared to WT controls.
B KEGG pathway analysis revealed significantly altered cell adhesion molecules and
cAMP signaling pathways in DD lesions of CNR1 k/o. **C** KEGG pathway analysis in DD

lesions of CNR2 k/o mice showed changes associated with cytokine receptor interactions
and steroid hormone biosynthesis pathways. **D, E** Venn diagrams showing the DEGs
among the 59 genes directly associated with the ECS, where we found limited DEGs in
DD lesions of CNR1 k/o (3) and CNR2 k/o mice (2), respectively. **F** A comprehensive
gene ontology analysis highlighting the roles of 59 ECS genes across diverse biological
processes (blue), cellular (orange), and molecular functions (light blue), accentuating their
broader impact beyond canonical ECS functions. Gene Number indicates the number of
DEGs enriched in pathway. Rich Ratio indicates the ratio of enriched DEGs to
background genes and Q-value indicates significance, with a value closer to zero being
more significant and is corrected by Benjamini-Hochberg method.

**Figure 3: Bulk RNA sequencing revealed alterations in immune-related gene
expression and pathway in EM lesions from CNR1 k/o and CNR2 k/o mice**
A Bar plot overview of the differentially expressed (DE) immune-related genes among
different genotypes and lesion types. **B, C** The volcano plots for DD lesions of CNR1 k/o
vs. WT and CNR2 k/o vs. WT, respectively, revealed 39 downregulated and 14
upregulated genes in CNR1 k/o DD lesions, while CNR2 k/o DD lesions exhibited 40
downregulated and 25 upregulated genes. Log2 fold change as the x-axis and log10 Q-
value (FDR adjusted) as the y-axis. Vertical dotted lines on the x-axis indicate ± 1-fold
change and vertical dotted line on the y-axis indicate Q-value of 0.05. **D, E** KEGG
pathway analysis of DEGs in CNR1 k/o DD lesions show significant alteration in the

813 chemokine signaling pathway, cytokine-cytokine receptor interaction, and toll-like
814 receptor signaling pathways, while in CNR2 k/o DD lesions, alterations were observed in
815 pathways related to cytokine-cytokine receptor interaction, Th17, Th1 and Th2 cell
816 differentiation.
817

818
Figure 4: Flow cytometry profiling of PF and splenic cells show altered immune cell
phenotypes in CNR1 k/o and CNR2 k/o mice with EM
A-E Gating panels of PF cells showing CD3 KO525-A on the y-axis vs CD11b APC-
A700-A on the x-axis among WT, CNR1 k/o, and CNR2 k/o mice with DD EM lesions, as
well as CNR2 k/o naive and CNR2 k/o sham-operated controls. **F** CD3+ total T cells in
the PF of CNR2 k/o mice with EM lesions, regardless of lesion types, were significantly
reduced compared to WT and CNR1 k/o mice with EM, as well as CNR2 k/o naïve and
sham operated mice. **G, H** This extended to the subsets of CD3+ T cells, CD4+, and
CD8+ T cells, respectively. CNR1 k/o mice with DD lesions also exhibited significantly
decreased CD3+ total T cells, CD4+ helper T cells, and CD8+ cytotoxic T cell frequencies
compared to their UnD counterparts. **I** CD11b+ monocyte/macrophage populations were
increased in the PF of CNR2 k/o mice with UnD and DD lesions compared to WT and
CNR1 k/o mice. CNR1 k/o mice with DD lesions displayed higher monocyte/macrophage
populations compared to their UnD counterparts. **K, J** CNR1 k/o mice with DD lesions
exhibited lower CD19+ B cells and NK1.1+ NK cell populations compared to WT and
CNR2 k/o mice. **L** Immune cell populations in splenocytes were analogous to findings

from PF cells, depicted by tSNE plots. *n* = 5-7 individual biological samples per
genotype. Statistical analyses were performed using the ordinary one-way ANOVA with
Holm-Sidak post hoc test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ and **** $p < 0.0001$.
Data presented as mean ± SD.

840
Figure 5: *In-vitro* validation of CNR2 deficiency on CD3 T cell viability and
functionality in conditions representative of EM lesion microenvironment
A Representative gating of percentage live, CD3+ total T cells from WT and CNR2 k/o
mice activated with or without PMA/Ionomycin cocktail in the presence or absence of
TNFα and media control. Graph with live gating shows count on the y axis and live/dead-
KO525 marker on the x axis. **B** Bar graphs of percentage live population of CD3+ total T
cells from CNR2 k/o mice show a significant decrease in the viability of cells activated
with PMA/ionomycin with or without TNFα. Whereas, no significant changes were
observed in the CD3+ total T cells from WT mice as well as from CNR2 k/o mice in

media. **C** Activation of CD3+ T cells of CNR2 k/o mice with PMA/ionomycin affected proliferation of CD4+ helper T cells specifically, with or without the presence of TNFα when compared to both their media control and WT controls. **D** No significant changes were observed in the proliferative CD8+ cytotoxic T cells from CNR2 k/o mice compared to their WT controls across different activation and non-activation groups. Ordinary two-way ANOVA with Tukey's post hoc test was performed to assess statistical significance. **p<0.05, **** $p < 0.0001$. Data presented as mean ± SD.

Figure 6: Imaging mass cytometry spatial profiling of immune cell distribution and cellular patterns in EM lesions in CNR1 k/o, CNR2 k/o, and WT mice
A The IMC data collection and analysis workflow outlines the steps involved in gaining comprehensive insights into the spatial distribution of immune cells and relevant cell types within UnD and DD EM-like lesions of WT, CNR1 k/o, and CNR2 k/o mice. **B, C** Representative images showing the single-cell segmentation performed following the

acquisition of 2 regions of interest (ROI) per section (3 biological samples per genotype)
and segmentation quality of the data after segmentation analysis was conducted,
respectively. **D** Non-linear dimensionality reduction after batch effect correction showed
distinct expression patterns of immune cells and cell state markers between UnD and DD
lesions. DD lesions from the CNR1 k/o and CNR2 k/o mice showed expression pattern
that was significantly different from the DD lesions of WT mice, as well as compared to
UnD lesions among different genotypes. **E** UMAP dimensionality reduction highlighted
key cell types and differences in composition between UnD and DD lesions. DD lesions
exhibited increased stroma and fibroblasts, decreased epithelial cells, and heightened
macrophage infiltration compared to UnD lesions.
875

Figure 7: Imaging mass cytometry revealed altered cellular composition and neighborhoods in EM lesions from CNR1 k/o, CNR2 k/o, and WT mice
A Representative image showing the distribution of different cell types within EM lesions (n = 2-3 per tissue type) based on unsupervised clustering and labeling. **B** Stacked bar plots reveal the distribution of various cell types within UnD and DD lesions of CNR1 k/o and CNR2 k/o mice compared to WT mice. **C** Increased CD3+ T cells and CD4+ helper T cells expression was found in the DD, EM lesions from CNR1 and CNR2 k/o mice compared to WT mice highlighting that T cells residing in the lesions were not affected. **D** CD11b+ monocyte and F4/80+ macrophage expression was significantly increased in the DD, EM lesions from mice lacking CNR1 and CNR2 compared to the WT controls. **E** Vimentin expressing stromal compartments that predominantly make up the EM lesions were also significantly increased in the DD lesions from CNR1 k/o and CNR2 k/o mice compared to WT mice. **F** Hallmark features of EM lesions, such as proliferation (Ki67+) and vascularization (CD31+) were significantly increased in the DD, EM lesions from mice lacking CNR1 and CNR2 compared to the WT controls. Combined, it highlights the effect on early lesion development and further progression through sustained proliferation due to dysregulated CNR1 and CNR2. **G** Heatmap representation of the CN analysis show distinct clustering patterns observed in the DD lesions among the different genotypes, where immune cells mainly clustered together in CN 4, while other cell types such as stroma, epithelial cells, and vasculature exhibited distinct clustering patterns across CN 6, 3, and 8, respectively.

Standards	Catalog number
2-Arachidonoyl Glycerol-d5	362162
2-Arachidonoyl Glycerol-d8	362160
Oleoyl Ethanolamide	90265
Palmitoyl Ethanolamide-d4	10007824
Arachidonoyl Ethanolamide-d8	390050
Oleoyl Ethanolamide-d4	9000552
Arachidonoyl Ethanolamide	90050
Palmitoleoyl Ethanolamide	10965
Arachidonoyl Ethanolamide-d4	10011178
2-Arachidonoyl Glycerol	62160

900 **Supplementary Table 1: Deuterated and non-deuterated standards used for mass**
901 **spectrometry analysis of the ECS ligands in plasma and tissues.**

Instrumentation	LC-MS : QQQ Agilent 6495 coupled with an Infinity 1260 HPLC system
LC Method	Column: CSH C18 2.1x100mm 3.5µm de Waters Mobile Phase A: 50/50 H2O/ACN +0.4 %FA Mobile Phase B: 90/10 IPA/ACN +0.4% FA Gradient: 30% to 45% B in 10min. Total run time: 20 min Volume of injection: 10µL
MS Method	ESI positive ionization MRM Transitions: AEA à 348.3 -> 62.1 AEA-d4 à 352.0 -> 66.0 OEA à 326.3 -> 62.0 OEA-d4 à 330.3 -> 66.0 PEA à 300.3 -> 62.0 PEA-d4 à 304.3 -> 62.0 2-AG à 379.3 -> 91.0 2-AG d5 à 384.3 -> 91.0

904 **Supplementary Table 2: Experimental conditions for ECS ligand identification**
905 **through LC-MS.**

	Mass Channel	Antibody Target	Clone	Provider	Dilution
1	141Pr	aSMA	polyclone	Abcam	1:100
2	142Nd	CD45R (B220)	RA3-6B2	Abcam	1:100
3	143Nd	Vimentin	D21H3	Cell Signaling Technology	1:200
4	144Nd				
5	145Nd	NK1.1	E6Y9G	Cell Signaling Technology	1:50
6	146Nd	CD31	SZ31	Cedarlane labs	1:100
7	147Sm	CD45	D3F8Q	Cell Signaling Technology	1:50
8	148Nd	Pan-CK	AE1/AE3	Standard BioTools	1:100
9	149Sm	CD11b	EPR1344	Standard BioTools	1:100
10	150Nd	CD19-Biotin	6OMP31	ThermoFisher	1:100
11	151Eu	CD44	E7K2Y	Cell Signaling Technology	1:100
12	152Sm				
13	153Eu	CD8a	D4W2Z	Cell Signaling Technology	1:100
14	154Sm	CD11c	D1V9Y	Cell Signaling Technology	1:50
15	155Gd				
16	156Gd	PD-L1	D5V3B	Cell Signaling Technology	1:50
17	158Gd	FoxP3	D6O8R	Cell Signaling Technology	1:50
18	159Tb	F4/80	D2S9R	Cell Signaling Technology	1:50
19	160Gd				
20	161Dy				
21	162Dy	CD3e	D4V8L	Cell Signaling Technology	1:100
22	163Dy				
23	164Dy	CD4	4SM95	ThermoFisher	1:50
24	165Ho	CD68	EPR23917-164	Abcam	1:100
25	166Er	Ly6G	1A8	Standard BioTools	1:50
26	167Er	PD-1	polyclonal	Bio-Techne	1:50
27	168Er	Ki67	B56	Standard BioTools	1:100
28	169Tm	CD206	polyclonal	Cell Signaling Technology	1:100
29	170Er	CD3	CD3-12	Abcam	1:50

30	171Yb	GranzymB	D6E9W	Cell Signaling Technology	1:100
31	172Yb	cleaved Casp3	Asp175	Cell Signaling Technology	1:100
32	173Yb				
33	174Yb	MHC-II	M5/114.15.2	Bio-Techne	1:100
34	175Lu	pS6	N7-548	Standard BioTools	1:100
35	176Yb	HH3	D1H2	Standard BioTools	1:300

908 **Supplementary Table 3: IMC antibodies used for the current study with their**
909 **respective dilutions**

911
Supplementary Figure 1: DEGs of bulk RNA sequencing between the UnD lesions
from CNR1 k/o and CNR2 k/o compared to WT controls.
A, **B** The volcano plots for DD lesions of CNR1 k/o and CNR2 k/o compared to WT,
respectively. Log2 fold change is represented on the x-axis and log10 Q-value (FDR
adjusted) as the y-axis. Vertical dotted lines on the x-axis indicate ± 1-fold change and
vertical dotted line on the y-axis indicate Q-value of 0.05. **C, D** KEGG pathway analysis
of DEGs of CNR1 k/o UnD lesions and CNR2 k/o UnD lesions compared to WT controls,
respectively. Gene Number represents the number of DEGs enriched in the pathway. Rich
Ratio shows the ratio of enriched DEGs to background genes and Q-value indicates
significance, with a value closer to zero being more significant and is corrected by
Benjamini-Hochberg method.
923
924

925
926 **Supplementary Figure 2**: **KEGG pathway analysis of immune specific DEGs.**
927 **A, B** Immune specific genes that were differentially expressed were subjected to KEGG
928 pathway analysis between CNR1 k/o UnD lesions and CNR2 k/o UnD lesions compared
929 to WT UnD controls, respectively.
930

931
932 **Supplementary Figure 3**: **Gating panel representing T cells from splenocytes.**
933 **A-G**, Splenocytes stained for CD3 on the y-axis and CD11b on the x-axis among WT,
934 CNR1 k/o, and CNR2 k/o mice with EM like condition, as well as CNR1 k/o and CNR2
935 k/o naive and sham-operated controls.

936
937 **Supplementary Figure 4**: **Flow cytometry analysis of immune cell phenotypes in the**
938 **splenocytes of WT, CNR1 k/o and CNR2 k/o mice with EMS.**
939 **A-F** Bar plot (mean ± SD) representation of splenocytes (SP) stained for CD3+ total T
940 cells, CD4+ helper T cells, CD8+ cytotoxic T cells, CD19+ B cells, CD11b+ monocytes
941 and F4/80 monocytes/macrophages of all the genotypes with EM lesions, respectively. $n =$
942 5-7 individual biological samples per genotype. Statistical analyses were performed using
943 the ordinary one-way ANOVA with Holm-Sidak post hoc test. * $p < 0.05$, ** $p < 0.01$, ***
944 $p < 0.001$ and **** $p < 0.0001$
945
946

A

B

C   **D**

947
Supplementary Figure 5: **In vitro functional assay and flow cytometry evaluation of**
activated CD3 T cells from naïve WT and CNR2 k/o mice.
A, **B** Flow cytometry gating strategies for CD3+ T cells to identifiy the different
phenotypes and functional state, respectively. **C**, Bar plots representing the percentage
positive live cells that are double positive for proliferation (Ki67) and IFNγ markers. **D**,
Bar plots showing the percentage positive live cells that are negative for Ki67 and positive
for IFNγ. Statistical analyses were performed using the ordinary one-way ANOVA with
Holm-Sidak post hoc test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ and **** $p < 0.0001$.
Data presented as mean ± SD.
957

958
Supplementary Figure 6: **IMC and cellular neighbourhoods (CN) in UnD, EMS**
lesions from CNR1 k/o, CNR2 k/o and WT mice.
A, B Representative image of 8 distinct cell types from CN analysis of DD and UnD
lesions from WT, CNR1 k/o, and CNR2 k/o mice, respectively. **C** Heatmap representation
of CN analysis shows distinct clustering patterns observed in the UnD lesions among the
different genotypes. The clustering reveals distinct spatial patterns of immune cell
populations within the UnD lesions, which appear to differ from the observations in
Figure 7G. This suggests potential spatial heterogeneity in the immune landscape of EM
like lesions under conditions of decidualization.
968