Title

1
2 Autophagosome development and chloroplast segmentation occur synchronously for piecemeal
3 degradation of chloroplasts

Authors

4
5 Masanori Izumi[a,b,1], Sakuya Nakamura[a], Kohei Otomo[c,d,e,f,g], Hiroyuki Ishida[h], Jun Hidema[i],
6 Tomomi Nemoto[c,d,e,f], Shinya Hagihara[a],

Affiliations

7
[a]Center for Sustainable Resource Science (CSRS), RIKEN, 351-0198 Wako, Japan
[b]Frontier Research Institute for Interdisciplinary Sciences (FRIS), Tohoku University, 980-8578
Sendai, Japan
[c]Exploratory Research Center on Life and Living Systems (ExCELLs), National Institutes of
Natural Sciences, 444-8787 Okazaki, Japan
[d]National Institute for Physiological Sciences, National Institutes of Natural Sciences, 444-8787
Okazaki, Japan
[e]The Graduate University for Advanced Studies (SOKENDAI), 444-8787 Okazaki, Japan
[f]Research Institute for Electronic Science, Hokkaido University, 001-0020 Sapporo, Japan
[g]Graduate School of Medicine, Juntendo University, 113-8421 Tokyo, Japan
[h]Graduate School of Agricultural Science, Tohoku University, 980-8572 Sendai, Japan
[i]Graduate School of Life Sciences, Tohoku University, 980-8577 Sendai, Japan
[1]Address correspondence to: masanori.izumi@riken.jp

Author contributions

21
22 M.I. conceived the study. M.I. performed biochemical assays. M.I., K.O, and T.N. performed
23 two-photon excitation microscopy observations. M.I., S.N., H.I., J.H., and S.H. performed
24 confocal microscopy studies. M.I. wrote the manuscript with the support of all authors.

Funding

25
26 This work was supported, in part, by KAKENHI (grant numbers JP16H06280, JP18H04852,
27 JP20H04916 to M.I., JP20H05352, JP22H04660 to S.N., JP22H04627 to H.I.), the Cooperative
28 Research Program of "NJRC Mater. & Dev." (to M.I.), and the Joint Research by Exploratory
29 Research Center on Life and Living Systems (ExCELLS program number 20-314 to M.I.).

Abstract

30
Plants distribute many nutrients to chloroplasts during leaf development and maturation. When
leaves senesce or experience sugar starvation, the autophagy machinery degrades chloroplast
proteins to facilitate efficient nutrient reuse. Here, we report on the intracellular dynamics of an
autophagy pathway responsible for piecemeal degradation of chloroplast components. Through
live-cell monitoring of chloroplast morphology, we observed the formation of chloroplast
budding structures in sugar-starved leaves. These buds were then released and incorporated into
the vacuolar lumen as an autophagic cargo termed a Rubisco-containing body. The budding
structures did not accumulate in mutants of core autophagy machinery, suggesting that
autophagosome creation is required for forming chloroplast buds. Simultaneous tracking of
chloroplast morphology and autophagosome development revealed that the isolation membranes
of autophagosomes interact closely with part of the chloroplast surface before forming
chloroplast buds. Chloroplasts then protrude at the site associated with the isolation membranes,
which divide synchronously with autophagosome maturation. This autophagy-related division
does not require DYNAMIN-RELATED PROTEIN 5B, which constitutes the division ring for
chloroplast proliferation in growing leaves. An unidentified division machinery may thus
fragment chloroplasts for degradation in coordination with the development of the chloroplast-
associated isolation membrane.

Introduction

Organelle morphology changes dynamically in response to fluctuations in cell functions, developmental stages, and environmental cues. Plastids are plant-specific organelles that differentiate into chloroplasts in green tissues to perform photosynthesis. Plastids further serve as hubs for metabolic pathways such as the biosynthesis of amino acids and plant hormones. Dynamic changes in plastid morphology also act as switches initiating biological programs (Osteryoung and Pyke, 2014). For instance, chloroplasts in leaf pavement cells form stroma-filled tubular extensions termed stromules to activate programmed cell death as an immune response upon perception of an invading pathogen, in a process regulated by the plant hormone salicylic acid (SA) (Caplan et al., 2015). Stromules are thin, tubular structures of plastid stroma less than 1 μm in diameter (Hanson and Sattarzadeh, 2011).

Leaf aging is closely associated with morphological and functional changes of plastids. At the early stage of leaf development, the undifferentiated form of plastids, termed proplastids, are converted to green chloroplasts as their population expands through active division. In Arabidopsis (*Arabidopsis thaliana*), the coordination of cytoplasmic ring structures comprising DYNAMIN-RELATED PROTEIN 5B (DRP5B), also called ACCUMULATION AND REPLICATION OF CHLOROPLASTS 5 (ARC5), and intrachloroplastic ring structures consisting of FILAMENTING TEMPERATURE-SENSITIVE Z (FTSZ) proteins mediates chloroplast division (Chen et al., 2018). In mature leaves, spherical chloroplasts occupy almost all of the cytoplasm and manifest high photosynthetic activity. When older leaves later senesce, the size and number of chloroplasts gradually decrease (Martinoia et al., 1983; Mae et al., 1984). Younger leaves closer to the shoot apex sometimes shade older leaves beneath them, thus perturbing photosynthesis by decreasing the amount of light reaching those leaves and causing local sugar starvation. In such sugar-starved leaves, senescence is accelerated to facilitate the relocation of nutrients into newly developing tissues (Weaver and Amasino, 2001; Ono et al., 2013; Law et al., 2018). Autophagy is an intracellular degradation machinery that contributes to the decline in chloroplast volume and numbers during leaf senescence (Wada et al., 2009).

Autophagy is a ubiquitous mechanism that transports cytoplasmic components to the vacuolar/lysosomal compartment for degradation in eukaryotic cells. The major autophagy pathway known as macroautophagy, or simply autophagy, begins with the assembly of a membrane termed the isolation membrane (also known as a phagophore) that forms a double-membrane structure termed the autophagosome and sequesters a portion of the cytoplasm. Core AUTOPHAGY (ATG) proteins (ATG1–10, 12–14, 16, and 18) are highly conserved among plants, yeasts, and mammals (Yoshimoto and Ohsumi, 2018; Nakamura et al., 2021a) and are required for the initiation and elongation of the isolation membrane (Nakatogawa, 2020). The outer membrane of autophagosomes fuses with the vacuolar/lysosomal membrane, resulting in the formation of autophagic bodies whose cargos are digested by vacuolar/lysosomal hydrolases. Our previous studies identified two types of autophagy for chloroplast degradation in mature leaves (Ishida et al., 2014; Nakamura and Izumi, 2018). Senescence and sugar starvation preferentially activate piecemeal degradation of chloroplast components involving a specific type of autophagosomal cargo termed a Rubisco-containing body (RCB), leading to smaller chloroplasts (Ishida et al., 2008; Wada et al., 2009; Izumi et al., 2015). The second type of autophagy, termed chlorophagy, removes entire unnecessary chloroplasts (Izumi et al., 2017; Nakamura et al., 2018), thereby modulating the number of chloroplasts in a cell (Wada et al., 2009; Izumi et al., 2017). Chlorophagy is likely another type of autophagy, termed microautophagy, during which the vacuolar membrane directly participates in sequestering the degradation target in the absence of encapsulation by autophagosomes (Nakamura et al., 2017; Lee et al., 2023).

Mitochondria are another type of energy-producing organelle derived from endosymbiosis. Dynamin-related proteins (DRPs) also mediate mitochondrial division (Giacomello et al., 2020); the involvement of DRP-mediated organelle division in

mitochondrion-targeted autophagy—termed mitophagy—in mammals and budding yeast
(*Saccharomyces cerevisiae*) has been under debate (Chen et al., 2022). Drp1 and Dnm1
(Dynamin-related 1) participate in mitochondrial fission in mammals and budding yeast,
respectively. Previous studies have suggested that segmented mitochondria resulting from
Drp1/Dnm1-mediated fission become a target for degradation by autophagosomes (Twig et al.,
2008; Rambold et al., 2011; Abeliovich et al., 2013). Notably, Drp1/Dnm1-independent
mitophagy has also been observed (Yamashita et al., 2016), whereby mitochondrial division
occurs concomitantly with the development of the mitochondrion-associated isolation
membrane, forming an autophagosome that specifically contains mitochondrial components
termed the mitophagosome. A recent study identified the mitochondrial intermembrane-space
protein mitofissin (also called Atg44), required for mitochondrial fission during yeast
mitophagy (Fukuda et al., 2023).
Plant autophagic bodies and RCBs are typically around 1 μm in diameter (Chiba et al.,
2003; Yoshimoto et al., 2004), which is smaller than chloroplasts. Thus, piecemeal-type
chloroplast autophagy via RCBs must start with division of the chloroplast segments that are to
be degraded. However, how a fragment of a chloroplast can be transported to the vacuole
remains poorly understood. Here, to characterize the underlying intracellular dynamics, we
performed high-resolution imaging analyses in living cells of Arabidopsis leaves. We used time-
lapse microscopy to observe the segmentation of chloroplast budding structures and their
transport to the vacuolar lumen. Use of multiple organelle markers revealed that the
development of the chloroplast-associated isolation membrane and the formation of RCBs occur
simultaneously. These morphological changes did not correspond to the emergence of stromules
or to DRP5b-mediated chloroplast division. Therefore, a previously undescribed chloroplast
division machinery may be in play during chloroplast autophagy.
123

Results

Chloroplast budding structures containing stroma and envelope are released in response
to sugar starvation
To observe the intracellular dynamics of the piecemeal-type of chloroplast autophagy, we
subjected the mature leaves of Arabidopsis plants to incubation in darkness, as this treatment
induces sugar starvation and accelerates leaf senescence, thus activating the RCB-mediated
chloroplast autophagy (Izumi et al., 2010). Exogenous application of concanamycin A (concA),
an inhibitor of vacuolar H^+-ATPase, allows the vacuolar accumulation of autophagic cargos,
including RCBs, facilitating their visualization (Yoshimoto et al., 2004; Ishida et al., 2008).
Following the excision and incubation in darkness of leaves harboring chloroplast stroma–
targeted fluorescent protein markers such as RUBISCO SMALL SUBUNIT (RBCS) fused to
monomeric red fluorescent protein (mRFP) in the presence of concA, we observed the
accumulation of many small puncta containing the RBCS-mRFP signal in the vacuole, i.e.,
RCBs (Figure 1–figure supplement 1A and 1C). Although co-incubation of dark-treated leaves
with concA and mineral nutrient–rich Murashige and Skoog (MS) salts did not block the RCB
accumulation, the addition of sucrose did, by rescuing the sugar-starved leaf (Figure 1–figure
supplement 1A and 1C). We detected no RCBs when we subjected the leaves of a mutant for the
core *ATG* gene *ATG7* to dark incubation in sucrose-free solution (Figure 1–figure supplement
1B and 1C). These observations indicate that RCBs are a type of autophagic cargo involved in
the degradation of chloroplast fragments in response to sugar starvation. Therefore, in this study,
we monitored chloroplast morphology in sugar-starved leaves resulting from their incubation in
darkness in a sugar-free solution.
We wished to monitor the changes in the three-dimensional (3D) morphology of
chloroplasts during autophagic degradation of chloroplast fragments. We thus used a two-
photon excitation microscope equipped with a confocal spinning-disk unit (Otomo et al., 2015),

Fig. 1

Figure 1. Chloroplast buds are released in sugar-starved leaves.
Time-lapse observations of 3D-reconstructed chloroplast morphology in Arabidopsis mesophyll cells accumulating chloroplast stroma–targeted fluorescent markers. A leaf from a plant accumulating chloroplast stroma–targeted GFP (CT-GFP) (A), RBCS-mRFP (B), or RBCS-EYFP (C) was incubated in sugar-free solution in darkness for 5 h from dawn (A), 21 h (B), or 24 h from the period in light (C) and then observed through a two-photon excitation microscope equipped with a confocal spinning-disk unit. Second rosette leaves from 20- to 22-d-old plants were used. Images in (A–C) are still frames from Videos 1, 2, and 3, respectively. Time scales above the images indicate the elapsed time from the start of the respective videos. Arrowheads indicate chloroplast budding structures. Scale bars, 5 µm. In (C), green, RBCS-EYFP; magenta, chlorophyll fluorescence. In the merged images, the overlapping regions of RBCS-EYFP and chlorophyll signals appear white.

Figure 1-figure supplement 1

Figure 1–figure supplement 1. Accumulation of chloroplast stroma components in the vacuole via autophagy.

Confocal images of mesophyll cells from wild-type (A) or *atg7* (B) plants accumulating the chloroplast stroma marker RBCS-mRFP. Second rosette leaves of 21-d-old plants were excised and incubated in 10 mM MES-NaOH containing 1 µM concanamycin A (concA) in darkness for 1 d. Sucrose (Suc) or Murashige and Skoog (MS) salts were added as an energy or nutrient source, respectively. Second rosette leaves from nontreated plants are shown as control. Green, RBCS-mRFP; magenta, chlorophyll fluorescence. Scale bars, 10 µm. The small puncta containing RBCS-mRFP without chlorophyll signal appear green and are Rubisco-containing bodies (RCBs) in the vacuole. (C) Number of accumulated RCBs from the observations described in (A) and (B). Different lowercase letters denote significant differences based on Tukey's test ($P < 0.05$). Values are means $\pm$ SE ($n = 4$). Dots represent individual data points.

Figure 1-figure supplement 2

Figure 1–figure supplement 2. The release of chloroplast buds by autophagy contributes to the decline in chloroplast stroma volume.
(A) Orthogonal projections produced from *z*-stack images (30 μm in depth) of mesophyll cells from WT or *atg7* leaves accumulating the chloroplast stroma marker RBCS-mRFP. Second rosette leaves of 21-d-old plants were excised at dawn and incubated in 10 mM MES-NaOH in the dark for 24 h. Mesophyll cells were observed before (0 h) and after (Dark 24 h) the treatment. Green, RBCS-mRFP. Scale bars, 10 μm. (B) Violin plot showing the volume of chloroplast stroma labeled by RBCS-mRFP, measured from the observations described in (A). Different lowercase letters denote significant differences based on Tukey's test ($P < 0.05$). Yellow bars indicate mean values. Five individual plants were observed, and the volumes of 564 (WT, 0 h), 554 (WT, dark 24 h), 597 (*atg7*, 0 h), and 725 (*atg7*, dark 24 h) chloroplasts were scored.

which facilitates the monitoring of changes in the 3D cellular structures of living plant cells (Sasaki et al., 2019; Nakamura et al., 2021b). Using dark-incubated leaves from transgenic plants harboring chloroplast-targeted *GFP* (*CT-GFP*) or *RBCS-mRFP* transgenes, we determined that mesophyll chloroplasts form a type of budding structure containing stromal components (Figure 1A and 1B, arrowheads). We observed these structures budding off from their associated chloroplasts within a few minutes (Figure 1A and 1B, Videos 1 and 2). Dual detection of the chloroplast stromal marker RBCS fused to enhanced yellow fluorescent protein (RBCS-EYFP) and of chlorophyll autofluorescence confirmed that the budding structure specifically contained stroma material (EYFP positive) without any chlorophyll signal, a marker of thylakoid membrane (Figure 1C, Video 3). These time-lapse observations reveal that chloroplasts form budding structures that are released in sugar-starved leaves. The chloroplast buds are cytoplasmic spherical bodies (approximately 0.5 to 1.5 μm in diameter) that are labeled by a chloroplast stroma marker and are adjacent to a main chloroplast body.

To estimate the impact of the release of chloroplast buds on the decline in chloroplast volume, we measured the volume of RBCS-mRFP signals in wild-type (WT) and *atg7* leaves before and after incubation in the dark for 24 h (Figure 1–figure supplement 2). Chloroplasts accumulate starch during the day and consume most of this starch during the night (Smith and Stitt, 2007). To minimize the effects of starch metabolism on the changes in chloroplast volume, we began sugar starvation treatment at the end of the night when a major portion of the starch has been consumed. After the treatment, stroma volume decreased in both genotypes (Figure 1–figure supplement 2B). However, the decline in volume was smaller in *atg7* leaves (14%) than in WT leaves (30%). The vacuolar transport of chloroplast budding structures as RCBs did not occur in *atg7* leaves (Figure 1–figure supplement 1). These results suggest that the release of chloroplast buds contributes to the decrease in chloroplast stroma volume in sugar-starved leaves.

We then performed a simultaneous detection of three organelle markers using conventional confocal microscopes to determine if the chloroplast buds contained membrane-bound proteins. Accordingly, we incubated leaves from plants accumulating the chloroplast outer envelope marker TRANSLOCON AT THE OUTER MEMBRANE OF CHLOROPLASTS 64 (TOC64) fused to mRFP and the stromal marker RBCS-GFP in darkness as described above. Under these conditions, we observed TOC64-mRFP signal surrounding a chloroplast bud (Figure 2A, arrowheads, Figure 2–figure supplement 1A). We then monitored chloroplasts in leaves accumulating the chloroplast inner envelope membrane protein K^+ EFFLUX ANTIPORTER 1 (KEA1) fused to mRFP along with RBCS-GFP after dark incubation. The chloroplast buds contained KEA1-mRFP signal but no chlorophyll signal (Figure 2–figure supplement 1B). We also captured the transport of an RCB with TOC64-mRFP or KEA1-mRFP signal (Figure 2B, Video 4, Figure 2–figure supplement 2, Video 5). These results indicate that the budding structures contained chloroplast envelope components. In another set of experiments, we generated transgenic plants accumulating the thylakoid membrane marker ATP synthase gamma subunit (ATPC1) fused to tagRFP and stromal RBCS-GFP. The ATPC1-tagRFP fluorescent signal fully overlapped with that of chlorophyll autofluorescence, with both originating from the thylakoid membrane (Figure 2C and 2D). In the chloroplast budding structures, however, we detected no ATPC1-tagRFP signal, in contrast to the strong stromal RBCS-GFP signal (Figure 2C, arrowheads, Figure 2–figure supplement 1C). In agreement with this observation, we captured a released RCB that did not contain any ATPC1-tagRFP or chlorophyll signals, but accumulated plentiful RBCS-GFP (Figure 2D, Video 6). These observations support the notion that the chloroplast buds containing stromal and envelope proteins are released as RCBs without thylakoid membranes. These live-cell imaging results are consistent with an immuno-electron microscopy study of RCBs in wheat (*Triticum aestivum*) leaves (Chiba et al., 2003).

Fig. 2

Figure 2. Chloroplast buds containing stroma and envelope components are released from the chloroplasts.

Time-lapse observations of Arabidopsis mesophyll cells accumulating the chloroplast stroma marker along with an envelope marker or a thylakoid membrane marker. Leaves accumulating stromal RBCS-GFP along with outer envelope–bound TOC64-mRFP (A and B) or with thylakoid membrane–bound ATPC1-tagRFP (C and D) were incubated in sugar-free solution in darkness for 5 to 9 h from dawn and then observed. Second rosette leaves from 21- to 22-d-old plants were used. Images in (B) or (D) are still frames from Videos 4 and 6, respectively. Time scales above the images indicate the elapsed time from the start of the respective videos. Arrowheads indicate chloroplast budding structures. Scale bars, 5 μm. Green, TOC64-mRFP or ATPC1-tagRFP; magenta, RBCS-GFP; orange, chlorophyll (Chl) fluorescence. The graphs in (A and C) show fluorescence intensities along the blue lines (a to b) in the magnified images of the area indicated by dashed blue boxes. The intensities are shown relative to the maximum intensity for each fluorescence channel, set to 1.

Figure 2-figure supplement 1

Figure 2–figure supplement 1. Chloroplast buds contain stroma and envelope proteins.
Second rosette leaves accumulating stromal RBCS-GFP along with outer envelope membrane–bound TOC64-mRFP (A), inner envelope membrane–bound KEA1-mRFP (B), or thylakoid membrane–bound ATPC1-tagRFP (C) were incubated in sugar-free solution in the dark for 5 to 9 h from dawn, and chloroplast budding structures were observed. Second rosette leaves from 21- to 22-d-old plants were used. Arrowheads indicate chloroplast budding structures. Four representative images of chloroplast buds are shown for each plant line. Scale bars, 5 µm. Green, TOC64-mRFP, KEA1-mRFP, or ATPC1-tagRFP; magenta, RBCS-GFP; orange, chlorophyll (Chl) fluorescence. The graphs show the relative ratio of GFP, RFP, or chlorophyll fluorescence intensity per unit area within chloroplast budding structures vs. their neighboring chloroplasts (bud/chloroplast). The intensities from 33 (A), 32 (B), or 30 (C) chloroplast buds from 8 individual plants were measured. Values are means ± SE (n = 30–33). Dots represent individual data points in each graph.

Figure 2-figure supplement 2

Figure 2–figure supplement 2. A released chloroplast bud contains an inner envelope marker protein.
Time-lapse observations of Arabidopsis mesophyll cells accumulating the chloroplast stroma marker RBCS-GFP along with inner envelope membrane–bound KEA1-mRFP. The second rosette leaf from a 22-d-old plant was incubated in sugar-free solution in the dark for 6 h from dawn and observed. The images are still frames from Video 5. Time scales above the images indicate the elapsed time from the start of the video. Arrowheads indicate chloroplast budding structures. Scale bars, 5 μm. Green, KEA1-mRFP; magenta, RBCS-GFP; orange, chlorophyll (Chl) fluorescence.

The vacuolar membrane dynamically interacts with chloroplast stromal components to incorporate them into the vacuolar lumen

We tracked the trafficking of RCBs in mesophyll cells further and obtained a two-dimensional (2D) sequential series of images of the release and transport of an RCB in leaves accumulating stromal CT-GFP (Figure 3A, Video 7). This time series included two phases during which the punctum moved quickly (Figure 3B, 24.4–34.8 s and 73.8–90.5 s). A tracking analysis for the RCB indicated that during the early phase (24.4–34.8 s), the punctum moved rather smoothly in one general direction (Figure 3C, Video 7); however, during the later phase (73.8–90.5 s), the punctum appeared to move more randomly, akin to Brownian movement (Figure 3C, Video 7). Other tracking series of dark-incubated leaves accumulating RBCS-tagRFP or RBCS-EYFP also showed RCBs starting to exhibit random movement: after 185 s in Figure 3–figure supplement 1A (Video 8) or after 25 s in Figure 3–figure supplement 1B (Video 9). Previous studies showed that autophagic bodies accumulating into the vacuolar lumen exhibit such random movements (Ishida et al., 2008). We thus reasoned that the incorporation of RCBs into the vacuolar lumen from the cytoplasm takes place when these structures start to exhibit random movement.

We therefore turned our attention to the dynamics of the vacuolar membrane when it is incorporating RCBs into the vacuolar lumen in transgenic plants accumulating the vacuolar membrane marker VACUOLAR H^+-PYROPHOSPHATASE 1 (VHP1) fused to mGFP (Segami et al., 2014) along with the stromal marker RBCS-mRFP (Figure 4). From one of our time-lapse imaging series, we identified still frames of an RCB present in the cytoplasm being engulfed by the vacuolar membrane (Figure 4A, open arrowheads from 25.4–30.5 s, Video 10). Later in the time series, we observed the opening of the part of the vacuolar membrane that had engulfed the RCB, resulting in the RCB being released into the vacuolar lumen (Figure 4A, filled arrowheads from 47.0–52.0 s, Video 10). We captured another instance of the dynamics of the vacuolar membrane in another time series, with the vacuolar membrane first surrounding the RCB (52.6–57.4 s) before releasing it (65.3–72.9 s) into the vacuolar lumen (Figure 4B, Video 11). Figure 4–figure supplement 1 and Videos 12–14 show similar membrane dynamics that were observed in three individual plants. An RCB marked by RBCS-mRFP was engulfed by the vacuolar membrane at 48.0–54.0 s (Figure 4–figure supplement 1A, Video 12), 6.0–15.0 s (Figure 4–figure supplement 1B, Video 13), and 111.0–123.0 s (Figure 4–figure supplement 1C, Video 14). The RCBs were then released into the vacuolar lumen. Our time-lapse imaging approach thus allowed us to successfully visualize the cytoplasm-to-vacuole transport progression of chloroplast stroma material via autophagy.

Formation and segmentation of chloroplast buds occur simultaneously with autophagosome maturation

Based on the observation of spherical structures budding off from chloroplasts (Figures 1–3), we hypothesized that these structures are formed in response to sugar starvation and are subsequently recognized and sequestered by the autophagosomal membranes. We thus expected the leaves of autophagy-deficient mutants to accumulate multiple extended structures of chloroplast stroma as chloroplast protrusions upon exposure to sugar starvation. This hypothesis seemed consistent with the observations of stromule-rich cells in hypocotyls or dark-incubated leaves of autophagy-deficient *atg5* or *atg7* mutants, compared with non-mutants (Ishida et al., 2008; Spitzer et al., 2015), assuming that the accumulation of chloroplast protrusions precedes stromule formation in *atg* mutants. To test this hypothesis, we subjected the leaves of *atg5* and *atg7* mutants, each defective in the function of a core ATG protein, to darkness-induced sugar starvation and treatment with concA for 1 d (Figure 5). We then evaluated the appearance of RCBs and chloroplast protrusions comprising extended structures of chloroplast stroma labeled by RBCS-mRFP, approximately 0.5 to 1.5 μm long. We detected no RCB in the leaves of either mutant (Figure 5A and 5B); however, in both the WT and the mutants, we noticed fewer chloroplast protrusions in dark-incubated leaves treated with concA compared with the

Fig. 3

Figure 3. Tracking the transport of a Rubisco-containing body.
A leaf accumulating chloroplast stroma–targeted GFP (CT-GFP) was incubated in sugar-free solution in darkness for 6 h from dawn, and the transport of a Rubisco-containing body (RCB) was tracked. The second rosette leaf from a 21-d-old plant was used. (A) Confocal images during the periods when the RCB moved quickly (24.4–34.8 and 73.8–90.5 s). Arrowheads indicate an RCB. The images are still frames from Video 7. Time scales above the images indicate the elapsed time from the start of the video. Green, CT-GFP; magenta, chlorophyll (Chl) fluorescence. Scale bar, 5 µm. (B) Calculated speed of the tracked RCB in (A). (C) The track of the RCB. The color of the track line changes over time, as indicated by the color bar. Scale bar, 2 µm.

Figure 3-figure supplement 1

Figure 3–figure supplement 1. Rubisco-containing bodies appear to begin random movement during their tracking.
A leaf accumulating RBCS-tagRFP (A) or RBCS-EYFP (B) was incubated in sugar-free solution in the dark for 7 to 8 h from dawn, and the transport of a Rubisco-containing body was tracked. Second rosette leaves from 22-d-old plants were used. Arrowheads indicate a tracked RCB. The images are still frames from Videos 8 (A) or 9 (B), respectively. The images show the time when RCBs appeared to begin random movement (185–197 s in A or 25–33 s in B). Time scales above the images indicate the elapsed time from the start of the respective videos. Green, RBCS-tagRFP or RBCS-EYFP; magenta, chlorophyll fluorescence. Scale bars, 5 µm. The color of the track line changes over time, as indicated by the color bars below the images.

Fig. 4

Figure 4. Dynamics of the vacuolar membrane during the incorporation of Rubisco-containing bodies.
Leaves accumulating the chloroplast stroma marker RBCS-mRFP along with the vacuolar membrane marker VHP1-mGFP were incubated in sugar-free solution in darkness for 6 to 8 h from dawn, and the behavior of cytosolic RCBs was monitored. Second rosette leaves from 21- or 23-d-old plants were used. The images when the vacuolar membrane engulfs an RCB (25.4–30.5 s in A and 52.6–57.4 s in B) and when an RCB is incorporated into the vacuolar lumen (47.0–52.0 s in A and 65.3–72.9 s in B) are shown. The images in (A) and (B) are still frames from Videos 10 and 11, respectively. Time scales above the images indicate the elapsed time from the start of the respective videos. Open arrowheads indicate an RCB engulfed by the vacuolar membrane. Closed arrowheads indicate the open site of the vacuolar membrane for the incorporation of an RCB. V indicates the region of the vacuolar lumen. Green, VHP1-mGFP; magenta, RBCS-mRFP. Scale bars, 5 µm.

Figure 4-figure supplement 1

Figure 4–figure supplement 1. Additional observations of the incorporation of Rubisco-containing bodies into the vacuolar lumen.
Additional time-lapse data of the experiments described in Figure 4. The second rosette leaves from 21- (A) or 24-d-old (B, C) plants were used. The images in (A), (B), and (C) are still frames from Videos 12, 13, and 14, respectively. Time scales above the images indicate the elapsed time from the start of the respective videos. Open arrowheads indicate an RCB engulfed by the vacuolar membrane. Closed arrowheads indicate the vacuolar membrane structure that engulfs an RCB. V indicates the region of the vacuolar lumen. Green, VHP1-mGFP; magenta, RBCS-mRFP. Scale bars, 5 μm.

Fig. 5

Figure 5. Autophagy deficiency does not increase the number of chloroplast protrusions during a 1-d dark treatment.
Leaves from wild-type (WT), *atg5*, or *atg7* plants accumulating the chloroplast stroma marker RBCS-mRFP were incubated in sugar-free solution containing 1 µM concanamycin A (concA) for 1 d in darkness. Second rosette leaves from 21-d-old plants were used. 2D images of mesophyll cells were acquired (A), and the number of accumulated RCBs in the vacuoles was scored (B). Small puncta containing RBCS-mRFP without the chlorophyll signal appear green and are RCBs in the vacuole. Leaves from untreated plants are shown as control. The appearance of chloroplast protrusions was observed from orthogonal projections created from *z*-stack images (15 µm in depth; C), and the proportion of chloroplasts forming protrusion structures out of 50 chloroplasts was scored (D). Scale bars, 10 µm. Green, RBCS-mRFP; magenta, chlorophyll fluorescence. Only the merged channels are shown. The overlapping regions of RBCS-mRFP and chlorophyll signals appear white. Arrowheads indicate the structures that were counted as a chloroplast protrusion in (D). Different lowercase letters denote significant differences based on Tukey's test ($P < 0.05$). Values are means ± SE ($n = 4$). Dots represent individual data points in each graph.

Figure 5-figure supplement 1

Figure 5–figure supplement 1. Chloroplast protrusions do not increase in *atg2* or *atg10* mutant leaves during a 1-d dark treatment.
The experiments described in Figure 5 were performed on leaves from wild-type (WT), *atg2*, or *atg10* plants accumulating the chloroplast stroma marker RBCS-mRFP. Scale bars, 10 μm. Green, RBCS-mRFP; magenta, chlorophyll fluorescence. Only the merged channels are shown. The overlapping regions of RBCS-mRFP and chlorophyll signals appear white. Small puncta containing RBCS-mRFP without chlorophyll signal appear green and are RCBs in the vacuole. Arrowheads indicate the structures that were counted as a chloroplast protrusion in (D). Different lowercase letters denote significant differences based on Tukey's test ($P < 0.05$). Values are means $\pm$ SE ($n = 4$). Dots represent individual data points in each graph.

respective untreated control leaves, contrary to our hypothesis (Figure 5C and 5D). We checked single mutants of *ATG2* and *ATG10* (*atg2* and *atg10*) to investigate other core ATGs and obtained similar results (Figure 5–figure supplement 1): The number of chloroplast protrusions did not increase in *atg2* or *atg10* leaves during a 1-d incubation in darkness.

We therefore posited that autophagosome formation might be required for the production of chloroplast buds. To test this idea, we observed the behavior of autophagosomes in transgenic plants accumulating the isolation membrane marker AUTOPHAGY8a (ATG8a) fused to GFP (as GFP-ATG8a) along with stromal RBCS-mRFP or chloroplast stroma–targeted (CT)-DsRed (Figure 6). Following incubation of leaves in darkness and confocal microscopy observation, we observed isolation membranes (marked by GFP-ATG8a) that were tightly associated with the chloroplast surface before any budding structure was visible (Figure 6A and 6B, 0 s). Over time, however, fluorescence monitoring of the same area revealed that the isolation membrane–associated site from the chloroplast side gradually protruded and eventually formed a spherical body containing RBCS-mRFP or CT-DsRed and surrounded by GFP-ATG8a signal, i.e., an autophagosome containing an RCB (Figure 6A and 6B, arrowheads; Videos 15 and 16).

Using the time-series data shown in Figure 6B (Video 16), we evaluated the time-dependent changes of the size of the chloroplast bud and the ratio between the major axis and the minor axis of the GFP-ATG8a-labeled autophagosome (Figure 6C). This ratio dropped to 1 as the isolation membrane became a spherical autophagosome. During this observation period, the ratio between the major and minor axes started to decrease at around 40 s and approached 1 at around 100 s. The chloroplast budding structure appeared at around 50 s and gradually increased in size until around 120 s. These data support the notion that the chloroplast bud appears as the isolation membrane develops and becomes a segmented RCB when the spherical autophagosome forms. We identified another example in an independent image series (Figure 6–figure supplement 1, Video 17) where the ratio between the major and minor axes of the isolation membrane started to decrease at approximately 40 s and the chloroplast bud appeared at around 50 s. In the third set of images, we observed the formation of chloroplast buds and later of RCBs near GFP-ATG8a-associated sites on multiple chloroplasts and sequentially (Figure 6–figure supplement 2, Video 18). Therefore, we conclude that the development of the chloroplast-associated isolation membrane precedes the production of chloroplast buds and of RCBs for piecemeal-type chloroplast autophagy.

We then evaluated the proportion of chloroplast buds associated with GFP-ATG8a signal when leaves accumulating RBCS-mRFP and GFP-ATG8a were incubated in the dark for 5 to 9 h from the end of night (Figure 6–figure supplement 3). Of the chloroplast buds examined, 64% were GFP-ATG8a positive (Figure 6–figure supplement 3A–3C). The formation of GFP-ATG8a-labeled chloroplast buds increased in response to the dark treatment (Figure 6–figure supplement 3D), but the buds did not appear in leaves from *atg7* plants exposed to dark treatment (Figure 6–figure supplement 3E). Therefore, a large portion of chloroplast buds may form synchronously with the development of the isolation membrane under sugar-starvation treatment. We then turned our attention to the isolation membrane and evaluated the proportion of GFP-ATG8a-labeled structures associated with chloroplasts (Figure 6–figure supplement 4). Of the isolation membranes examined, 23% localized on the chloroplast surface as flattened structures (Figure 6–figure supplement 4A and 4D), and 20% localized with chloroplast buds as spherical or curved structures (Figure 6–figure supplement 4B and 4D). Therefore, 43% of the GFP-ATG8a-labeled structures participated in chloroplast degradation during the dark treatment in this study. These results support the notion that chloroplast budding accompanied by autophagosome development substantially contributes to the degradation of chloroplast components in sugar-starved leaves.

Fig. 6

A

GFP-ATG8a	0 s	68 s	76 s	84 s	92 s	100 s	108 s	116 s

RBCS-mRFP

Merged

B

0 s   45 s   60 s   75 s   90 s   105 s   120 s   135 s

GFP-ATG8a

CT-DsRed

Merged

C

Major axis/minor axis (GFP-ATG8a)

Major axis — Minor axis
GFP-ATG8a

Area of chloroplast bud (μm², CT-DsRed)

Area of chloroplast bud
Chloroplast

Time (s)

Figure 6. The formation of a chloroplast bud and the maturation of the chloroplast-associated isolation membrane occur concomitantly.

Leaves accumulating the chloroplast stroma marker, RBCS-mRFP (A) or CT-DsRed (B), and the isolation membrane marker GFP-ATG8a were incubated in sugar-free solution in darkness for 6 h from dawn (A) or 21 h from the period in light (B) and then observed. Second rosette leaves from 23-d-old plants were used. Arrowheads indicate the position of the chloroplast-associated isolation membrane. Images in (A) and (B) are still frames from Videos 15 and 16, respectively. Time scales above the images indicate the elapsed time from the start of the respective videos. Green, GFP-ATG8a; magenta, RBCS-mRFP or CT-DsRed. Scale bars, 5 μm. (C) Time-dependent changes in the ratio of the major axis to the minor axis in the GFP-ATG8a-labeled isolation membrane (top) or in the area of the chloroplast budding structure (bottom) as measured from the images in (B).

Figure 6-figure supplement 1

Figure 6–figure supplement 1. Another observation of the budding of the isolation membrane–associated site in a chloroplast.
A leaf accumulating the chloroplast stroma marker CT-DsRed and the isolation membrane marker GFP-ATG8a was incubated in sugar-free solution in darkness for 6 h from dawn and then observed. A second rosette leaf from a 22-d-old plant was used. Arrowheads indicate the position of the chloroplast-associated isolation membrane. The images are still frames from Video 17. Time scales above the images indicate the elapsed time from the start of the video. Green, GFP-ATG8a; magenta, CT-DsRed. Scale bars, 5 μm. (B) Time-dependent changes in the ratio of the major axis to the minor axis in the GFP-ATG8a-labeled isolation membrane (top) or in the area of the chloroplast budding structure (bottom), as measured from the images in (A).

Figure 6-figure supplement 2

Figure 6–figure supplement 2. Chloroplast buds surrounded by the isolation membrane appear in multiple chloroplasts.
A leaf accumulating the chloroplast stroma marker RBCS-tagRFP and the isolation membrane marker GFP-ATG8a was incubated in sugar-free solution in darkness for 5 h from dawn and then observed. A second rosette leaf from a 21-d-old plant was used. White, yellow, and blue arrowheads indicate the isolation membrane–associated site in different chloroplasts, respectively. The images are still frames from Video 18. Time scales above the images indicate the elapsed time from the start of the video. Green, GFP-ATG8a; magenta, RBCS-tagRFP; orange, chlorophyll (Chl) fluorescence. Scale bars, 5 μm.

Figure 6-figure supplement 3

Figure 6–figure supplement 3. Proportion of chloroplast buds engulfed by GFP-ATG8a-labeled isolation membranes.
Leaves accumulating the chloroplast stroma marker RBCS-mRFP and the isolation membrane marker GFP-ATG8a were incubated in sugar-free solution in the dark for 5 to 9 h from dawn, and chloroplast budding structures were observed. Second rosette leaves from 22- to 23-d-old plants were used. (A and B) Four representative images of chloroplasts buds that are surrounded by GFP-ATG8a signals (A) or two representative images of chloroplast buds that are not labeled by GFP-ATG8a signals (B) are shown. Arrowheads indicate chloroplast budding structures. Green, GFP-ATG8a; magenta, RBCS-mRFP. Scale bars, 5 μm. (C) The proportions of chloroplast buds that are labeled by GFP-ATG8a signals (+) or not (−) were calculated from observations of 58 chloroplast buds from 8 individual plants. (D) The number of GFP-ATG8a-positive chloroplast buds, as shown in (A), were counted in the fixed regions from leaves incubated in the dark for 5 to 9 h from dawn (Dark) or from untreated leaves of plants grown under normal conditions (Control). Five individual plants were observed. (E) Leaves from wild-type (WT) or *atg7* plants accumulating RBCS-mRFP and GFP-ATG8a were incubated in the dark for 5 to 9 h from dawn, and the number of GFP-ATG8a-positive chloroplast buds was counted. Six individual plants were observed. Values are means $\pm$ SE (n = 5–6). Asterisks denote significant differences based on t-test (***, $P < 0.001$). Dots represent individual data points in each graph.

Figure 6-figure supplement 4

Figure 6–figure supplement 4. Proportion of GFP-ATG8a-labeled structures that are associated with chloroplasts. Leaves accumulating the chloroplast stroma marker RBCS-mRFP and the isolation membrane marker GFP-ATG8a were incubated in sugar-free solution in the dark for 5 to 9 h from dawn, and GFP-ATG8a-labeled structures in mesophyll cells were observed. Second rosette leaves from 22-d-old plants were used. (A to C) GFP-ATG8a-positive structures were classified into three types: flattened structures associated with the chloroplast surface (A), spherical or curved structures associated with chloroplast budding structures (B), and other structures (C). Four representative images of each type are shown. Green, GFP-ATG8a; magenta, RBCS-mRFP. Scale bars, 5 μm. (D) The proportions of the three types of GFP-ATG8a-labeled structures were calculated from observations of 157 GFP-ATG8a-labeled structures from 13 individual plants.

SA influences stromule formation in autophagy-deficient mutants

Our initial hypothesis was that stromules are related to RCBs: Stromule-rich cells appeared because of impaired RCB production in *atg* mutants. However, our imaging assays above revealed that RCBs emerge from mesophyll chloroplasts in the absence of prior formation of stromules or protrusions (Figure 6), pointing to the existence of another factor responsible for the elevated stromule formation seen in *atg* mutant leaves. A previous study reported the hyperaccumulation of SA in senescing Arabidopsis *atg* mutants (Yoshimoto et al., 2009). Since SA signaling stimulates immune responses, including stromule formation (Caplan et al., 2015), we speculated that SA might contribute to stromule formation in autophagy-deficient mutants. To investigate this possibility, we generated transgenic plants in the *atg5* mutant background accumulating stromal GFP (CT-GFP) and a construct carrying NahG, encoding a bacterial SA hydroxylase that catabolizes SA (Delaney, 1994; Yoshimoto et al., 2009). In these plant lines, we evaluated the frequency of chloroplasts forming stromules, which are thin, tubular structures of chloroplast stroma less than 1 μm in diameter (Hanson and Sattarzadeh, 2011; Brunkard et al., 2015). We examined chloroplasts in guard cells, as this cell type actively forms many more stromules than the mature chloroplasts of mesophyll cells (Ishida and Yoshimoto, 2008). Chloroplasts in the guard cells of the *atg5* mutant formed more stromules than WT and NahG-expressing plants (Figure 7A and 7B). The introduction of the NahG construct into the *atg5* mutant background (*atg5* NahG) largely abolished stromule formation (Figure 7A and 7B). To further explore the role of SA in stromule formation, we generated double mutants of *SALICYLIC ACID INDUCTION DEFICIENT 2* (*SID2*), encoding the SA biosynthetic enzyme ISOCHORISMATE SYNTHASE 1, and *ATG5* or *ATG7* (*sid2 atg5* and *sid2 atg7*). The *sid2* mutation also resulted in fewer stromules in the double mutant relative to the respective *atg5* and *atg7* single mutants in guard cells (Figure 7C and 7D).

We next observed stromule formation in mesophyll chloroplasts accumulating CT-GFP or RBCS-mRFP and found that many chloroplasts formed stromules in senescent leaves of 36-d-old *atg5* and *atg7* plants (Figure 7E and 7F, Figure 7–figure supplement 1). In cells from the third rosette leaves of 36-d-old plants accumulating CT-GFP, we observed stromules in 22.5% or 15.3% of chloroplasts from the *atg5* or *atg7* single mutants, respectively; however, the introduction of the *sid2* mutation in *sid2 atg5* and *sid2 atg7* led to fewer stromules (9.8% or 2.5% in *sid2 atg5* or *sid2 atg7*, respectively; Figure 7E and 7F). In the leaves of 20-d-old plants, we did not detect stromules in any genotype (Figure 7–figure supplement 2). These results support the notion that SA accumulation due to autophagy deficiency activates stromule formation during leaf senescence.

The frequency of stromules was higher in the *sid2 atg5* and *sid2 atg7* double mutants than in WT plants and the *sid2* single mutant (Figure 7). We measured the content of hydrogen peroxide (H_2O_2) in all plants, since reactive oxygen species also trigger stromule formation (Caplan et al., 2015). In 36-d-old plants, *atg5* or *atg7* leaves contained 2.7 or 2.8 times more H_2O_2 than WT leaves (Figure 7G). Notably, the presence of the *sid2* mutation in the *sid2 atg5* and *sid2 atg7* double mutants did not alleviate H_2O_2 accumulation (Figure 7G). Such ROS accumulation may thus be one factor responsible for the greater number of stromules in *sid2 atg5* and *sid2 atg7* compared with WT and *sid2* plants. Moreover, NahG expression did not counteract the increase in H_2O_2 level caused by the *atg5* mutation (Figure 7–figure supplement 3). ROS accumulation in *atg* leaves may not depend on stimulation of SA signaling. These results are consistent with the results of histochemical ROS staining of Arabidopsis leaves in a previous study (Yoshimoto et al., 2009).

DRP5b is not required for autophagy of chloroplast fragments

Simultaneous progression of autophagosome formation and organelle segmentation has been previously observed during mitophagy in mammals and budding yeast (Yamashita et al., 2016; Fukuda et al., 2023). This autophagy-related mitochondrial division does not require the

Fig. 7

Figure 7. Diminished salicylic acid signal suppresses stromule formation in autophagy-deficient mutants.
(A) Orthogonal projections produced from *z*-stack images (10 μm in depth) of guard cells from wild-type (WT), *atg5*, NahG, and *atg5* NahG leaves accumulating chloroplast stroma–targeted GFP (CT-GFP). Second rosette leaves from 13-d-old plants were used. Scale bars, 10 μm. (B) Percentage of chloroplasts forming stromules in 10 stomata, from the observations described in (A). (C) Orthogonal projections produced from *z*-stack images (10 μm in depth) of guard cells from WT, *atg5*, *atg7*, *sid2*, *sid2 atg5*, and *sid2 atg7* leaves accumulating CT-GFP. Scale bars, 10 μm. (D) Percentage of chloroplasts forming stromules in 10 stomata, from the observations described in (C). (E) Orthogonal projections produced from *z*-stack images (15 μm in depth) of mesophyll cells from WT, *atg5*, *atg7*, *sid2*, *sid2 atg5*, and *sid2 atg7* leaves accumulating CT-GFP. Third rosette leaves from 36-d-old plants were observed. Green, CT-GFP; magenta, chlorophyll fluorescence. Only the merged channels are shown. Scale bars, 20 μm. (F) Percentage of chloroplasts forming stromules out of 50 chloroplasts in mesophyll cells, from the observations described in (E). (G) Hydrogen peroxide (H_2O_2) content in third rosette leaves from 36-d-old WT, *atg5*, *atg7*, *sid2*, *sid2 atg5*, and *sid2 atg7* plants. Different lowercase letters denote significant differences based on Tukey's test ($P < 0.05$). Values are means $\pm$ SE ($n = 3$ in B and D or 4 in F and G). Dots represent individual data points in each graph.

Figure 7-figure supplement 1

Figure 7–figure supplement 1. Autophagy deficiency activates stromule formation from mesophyll chloroplasts accumulating RBCS-mRFP in senescing leaves.
(A) Orthogonal projections produced from *z*-stack images (15 μm in depth) of mesophyll cells from leaves of WT, *atg5*, and *atg7* plants accumulating RBCS-mRFP. Third rosette leaves from 36-d-old plants were observed. Green, RBCS-mRFP; magenta, chlorophyll fluorescence. Only the merged channels are shown. Spherical bodies only exhibiting chlorophyll signal represent a portion of chloroplasts in pavement cells. Scale bars, 20 μm. (B) Percentage of chloroplasts forming stromules out of 50 chloroplasts in mesophyll cells, from the observations described in (A). Different lowercase letters denote significant differences based on Tukey's test ($P < 0.05$). Values are means $\pm$ SE ($n = 4$). Dots represent individual data points in the graph.

Figure 7-figure supplement 2

Figure 7–figure supplement 2. Autophagy deficiency does not activate stromule formation from mesophyll chloroplasts in young leaves.
Orthogonal projections produced from *z*-stack images (15 μm in depth) of mesophyll cells from WT, *atg5*, *atg7*, *sid2*, *sid2 atg5*, and *sid2 atg7* leaves accumulating CT-GFP. Third rosette leaves from 20-d-old plants were observed. Green, CT-GFP; magenta, chlorophyll fluorescence. Only the merged channels are shown. Scale bars, 20 μm.

Figure 7-figure supplement 3

Figure 7–figure supplement 3. NahG expression does not counteract the increase in hydrogen peroxide content produced by the *atg5* mutation.

Hydrogen peroxide (H_2O_2) contents in third rosette leaves from 36-d-old plants. WT plants without any transgenic construct and CT-GFP-expressing plants in the WT, *atg5*, NahG, and *atg5* NahG backgrounds were used. Different lowercase letters denote significant differences based on Tukey's test ($P < 0.05$). Values are means $\pm$ SE ($n = 4$). Dots represent individual data points.

mammalian Drp1- or yeast Dnm1-dependent division machinery, respectively. Here, we
examined whether the DRP5B-mediated chloroplast division participates in chloroplast
autophagy and RCB production.
To this end, we generated *drp5b* mutant lines accumulating the stromal marker RBCS-
mRFP or stromal GFP (CT-GFP). Although the mature leaves of *drp5b* had larger chloroplasts
than WT leaves, reflective of the impaired chloroplast division during cell expansion typical of
this mutant, many RCBs accumulated when the leaves were incubated in darkness in the
presence of concA (Figure 8A and 8B, Figure 8–figure supplement 1). We detected no RCB in
the leaves of the *drp5b atg5* double mutant (Figure 8–figure supplement 1). These results
indicate that RCB-mediated chloroplast autophagy is active in the *drp5b* mutant. To obtain an
independent confirmation of this result, we performed an immunoblot assay of autophagy flux,
based on the detection of free mRFP derived from the vacuolar cleavage of RBCS-mRFP (Ono
et al., 2013). When we incubated the leaves of plants accumulating RBCS-mRFP in darkness
without concA, free mRFP levels increased in leaves of WT but not *atg5* or *atg7* plants (Figure
8C and 8D), indicating the occurrence of autophagy-dependent degradation of stromal proteins
in response to sugar starvation in WT plants. The accumulation of free mRFP in the *drp5b*
mutant was at least as high as that seen in WT plants (Figure 8C). The ratio of free mRFP to
RBCS-mRFP increased in WT and *drp5b* plants after dark treatment (6.1 or 4.1 times higher
compared with control conditions in WT plants or *drp5b*, respectively), consistent with
autophagy-mediated degradation of chloroplast fragments (Figure 8D). Confocal microscopy
observations of mesophyll cells also showed the spread of mRFP signal in the vacuolar lumen in
dark-treated leaves of WT and *drp5b* plants (Figure 8–figure supplement 2). These results
indicate that the autophagic degradation of chloroplast stroma is active in both genotypes. The
accumulation of free mRFP was 3.2 times higher in untreated leaves of *drp5b* compared with
untreated WT leaves (Figure 8D); therefore, the activity of chloroplast autophagy might be
constitutively higher in leaves of *drp5b* than in those of WT plants.
We observed the budding off of chloroplast fragments in leaves of the *drp5b* mutant.
We incubated a leaf from the *drp5b* mutant accumulating RBCS-mRFP in darkness, which
allowed us to observe the formation of a chloroplast bud and its release (Figure 9A, arrowheads;
Video 19). In a leaf of the *drp5b* mutant accumulating RBCS-mRFP and GFP-ATG8a, we
detected the vesiculation of a chloroplast fragment starting at the site to which the isolation
membranes, labeled by GFP-ATG8a, were tightly associated (Figure 9B). The isolation
membranes were anchored at two sites of a large chloroplast in a leaf of the *drp5b* mutant
(Figure 9B, white and blue arrowheads); the chloroplast gradually protruded before releasing
RCBs surrounded by autophagosomes (Figure 9B, Video 20). We conclude that DRP5B is
dispensable for the autophagy-related division of chloroplast fragments.

Discussion
390
Chloroplast division machinery functioning in piecemeal-type chloroplast autophagy
We previously demonstrated that chloroplast components in mesophyll cells are transported to
the vacuole as a type of autophagic cargo termed RCBs to facilitate amino acid recycling (Ishida
et al., 2008; Hirota et al., 2018). However, how a portion of chloroplasts is mobilized into the
vacuolar lumen remained uncertain. In this study, high-resolution time-lapse imaging techniques
allowed us to visualize the trafficking progression of chloroplast fragments from their parental
chloroplasts to the vacuolar lumen. Importantly, we established that the development of the
chloroplast-associated isolation membrane and the division of the chloroplast fragment occur
simultaneously. This autophagy-related chloroplast division does not rely on DRP5B-mediated
chloroplast division. Likewise, the formation of chloroplast buds is unlikely to be linked to the
formation of stromules. Therefore, an unknown division machinery may be required for the
autophagy-related division of chloroplast fragments. A similar organelle division mechanism
takes place during mitophagy in mammals and budding yeast. The site of a mitochondrion that

Fig. 8

Figure 8. DRP5b is dispensable for chloroplast autophagy in sugar-starved leaves.
(A) Confocal images of mesophyll cells from wild-type (WT) and *drp5b* leaves accumulating the stroma marker RBCS-mRFP. Second rosette leaves from 21-d-old plants were incubated in sugar-free solution containing 1 μM concanamycin A (concA) in darkness for 1 d. Green, RBCS-mRFP; magenta, chlorophyll fluorescence. Scale bars, 10 μm. (B) Number of accumulated RCBs in WT, *drp5b*, *atg5*, and *atg7* leaves, counted from the observations of leaves incubated in the dark with concA described in (A). Different lowercase letters denote significant differences based on Tukey's test ($P <$ 0.05). Values are means ± SE ($n = 4$–5). (C) Biochemical detection of autophagy flux for chloroplast stroma based on a free RFP assay. RFP and cFBPase (loading control) were detected by immunoblotting of soluble protein extracts from leaves of WT, *drp5b*, *atg5*, and *atg7* plants accumulating RBCS-mRFP. Protein extracts from either untreated control leaves (cont) or leaves after 2 d of incubation in darkness (dark) were used. Total protein was detected by Coomassie Brilliant Blue (CBB) staining as a loading control. The filled arrowhead indicates RBCS-mRFP fusion, and the open arrowhead indicates free mRFP derived from the cleavage of RBCS-mRFP. (D) Quantification of the free mRFP/RBCS-mRFP ratio shown relative to that of untreated WT plants, which was set to 1. Asterisks denote significant differences based on *t*-test (***, $P < 0.001$; n.s., not significant). Values are means ± SE ($n = 4$). Dots represent individual data points in each graph.

Figure 8–figure supplement 1

Figure 8–figure supplement 1. Production of Rubisco-containing bodies in *drp5b* mutants is ATG5 dependent.
(A) Confocal images of mesophyll cells from wild-type (WT), *atg5*, *drp5b*, and *drp5b atg5* leaves accumulating the stroma marker CT-GFP. Second rosette leaves from 21-d-old plants were incubated in sugar-free solution containing 1 µM concanamycin A (concA) in darkness for 1 d. Second rosette leaves from untreated plants were used as control. Green, CT-GFP; magenta, chlorophyll fluorescence. Only the merged channels are shown. Small puncta containing CT-GFP without chlorophyll signal appear as green and are RCBs in the vacuole. Scale bars, 10 µm. (B) Number of accumulated RCBs, counted from the observations of leaves incubated in the dark with concA described in (A). Different lowercase letters denote significant differences based on Tukey's test ($P < 0.05$). Values are means $\pm$ SE ($n = 5$). Dots represent individual data points.

Figure 8–figure supplement 2

Figure 8–figure supplement 2. Vacuolar accumulation of stromal marker proteins in sugar-starved leaves.
(A) Confocal images of mesophyll cells from wild-type (WT), *drp5b*, *atg5*, and *atg7* leaves accumulating the stroma marker RBCS-mRFP. Second rosette leaves from 21-d-old plants were incubated in sugar-free solution in darkness for 2 d. Second rosette leaves from untreated plants are shown as control. Green, RBCS-mRFP; magenta, chlorophyll fluorescence. Scale bars, 20 μm. (B) RFP intensity in the vacuolar lumen, as measured from the observations described in (A) and shown relative to that of untreated WT plants, which was set to 1. Asterisks denote significant differences based on *t*-test (***, $P < 0.001$; n.s., not significant). Values are means $\pm$ SE ($n = 4$). Dots represent individual data points.

Fig. 9

Figure 9. Formation and segmentation of chloroplast buds in leaves of the *drp5b* mutant.
(A) A leaf from the *drp5b* mutant accumulating the stroma marker RBCS-mRFP was incubated in sugar-free solution in darkness for 5 h from dawn and then observed. Arrowheads indicate a chloroplast bud. Green, RBCS-mRFP; magenta, chlorophyll fluorescence (Chl). (B) A leaf from the *drp5b* mutant accumulating the stroma marker RBCS-mRFP and the isolation membrane marker GFP-ATG8a were incubated in sugar-free solution in darkness for 6 h from dawn and then observed. Second rosette leaves from 21- to 22-d-old plants were used. Arrowheads indicate the position of the chloroplast-associated isolation membrane. White and blue arrowheads indicate different isolation membrane–associated sites in a chloroplast. Images in (A) and (B) are still frames from Videos 19 and 20, respectively. Time scales above the images indicate the elapsed time from the start of the respective videos. Green, GFP-ATG8a; magenta, RBCS-mRFP; orange, chlorophyll (Chl) fluorescence. Scale bars, 5 μm.

is associated with the isolation membrane protrudes and divides as the isolation membrane
develops and becomes an enclosed autophagosome (Yamashita et al., 2016). This type of
mitochondrial division is independent of Drp1 and Dnm1. In yeast, the mitochondrial
intermembrane-space protein mitofissin/Atg44 divides the mitochondrial fragment in
coordination with autophagosome maturation (Fukuda et al., 2023). A functionally equivalent
protein has not been identified in mammals.
Numerous studies have explored the molecular mechanisms of chloroplast division
during the development of juvenile leaves (Chen et al., 2018). For instance, in Arabidopsis
plants, ARC6 and PARALOG OF ARC6 (PARC6) mediate the tethering of the FTSZ ring on the
chloroplast inner envelope (Vitha et al., 2003). PLASTID DIVISION 1 (PDV1) and PDV2
recruit DRP5B to the division site on the chloroplast outer envelope (Miyagishima et al., 2006).
The ARC6–PDV2 and PARC6–PDV1 complexes may control the coordination of the stromal
FTSZ ring and the cytosolic DRP5B ring (Wang et al., 2017; Chen et al., 2018). Therefore, the
controlled division of plastids requires the cooperation of multiple protein complexes across the
inner and outer envelopes. Nevertheless, whether chloroplasts divide during their piecemeal
degradation has not been evaluated. It seems likely that autophagy-related division of
chloroplast fragments needs to be carefully regulated, as the size of the segment has to fit into
the autophagosome. Since the volume occupied by chloroplasts per cell is tightly regulated
during cell expansion (Pyke and Leech, 1994), the decline of chloroplast volume during
starvation or senescence might also be highly controlled. Further studies are needed to elucidate
the molecular machinery underlying the autophagy-related division of chloroplast fragments for
degradation.
In our evaluation of chloroplast buds associated with GFP-ATG8a in excised leaves
exposed to darkness (Figure 6–figure supplement 3), 36% of chloroplast budding structures
were not labeled by GFP-ATG8a. Since endogenous ATG8 family members (ATG8a–8i) were
functional, such structures might have formed due to the activities of non-labeled ATG8 proteins.
Another possibility is that autophagy-independent chloroplast budding might occur. For
example, stromule-related structures might form. CHLOROPLAST VESICULATION (CV)–
containing vesicles are also observed as autophagy-independent puncta released from
chloroplasts for degradation (Wang and Blumwald, 2014; Pan et al., 2023). However,
biochemical assays for chloroplast autophagy flux suggested that ATG-dependent pathways
were the main routes for the vacuolar degradation of stromal components during the darkness-
induced sugar starvation treatment used in this study (Figure 8). Previous studies have revealed
that CV plays a role in plant responses to abiotic stress, such as drought or salt stress (Wang and
Blumwald, 2014; Pan et al., 2023). A recent study suggested that autophagy, but not CV-
dependent pathways, is the main contributor to the adaptative response of metabolism to sugar
starvation (Barros et al., 2023). Overall, these insights support the notion that, during the early
phase of sugar starvation (such as 1 to 2 d of extended darkness), the autophagy-associated
machinery is mainly responsible for chloroplast segmentation and degradation. It is unclear
what type of process contributed to the decrease in chloroplast stroma volume in leaves of *atg7*
plants exposed to dark treatment (Figure 1–figure supplement 2).
An assay of autophagic flux of RBCS-mRFP protein in leaves from *drp5b* mutants
suggested the possibility that chloroplast autophagy activity is constitutively higher in leaves of
drp5b than in WT plants (Figure 8). The impaired chloroplast division due to the *drp5b*
mutation might affect photosynthesis-mediated energy production and reduce energy availability,
thereby activating chloroplast autophagy without dark treatment. As described in the
Introduction, entire organelle–type autophagy (chlorophagy) mediates the degradation of
chloroplasts in their entirety. Mesophyll cells in leaves of *drp5b* mutants contain few
chloroplasts (approximately 4 per cell), whereas those in WT leaves contain approximately 60
chloroplasts (Miyagishima et al., 2006). Since the volume in a cell occupied by chloroplasts is
similar between genotypes, chloroplasts from *drp5b* are much larger than WT chloroplasts.

Chloroplasts that are too large might show inhibited chlorophagy, resulting in the alternative
activation of piecemeal-type chloroplast autophagy. Analyzing how chloroplasts are degraded in
mutants of chloroplast division ring components might provide insight into the interplay
between entire organelle–type autophagy and piecemeal-type autophagy during chloroplast
degradation.
460
Autophagy receptor and adapter proteins for the degradation of mature chloroplasts
Organelle-selective autophagy is typically controlled by receptor proteins that
recognize the target organelles and act as a bridge between the organelles and isolation
membrane–anchored ATG8 (Farre and Subramani, 2016). The observation of a chloroplast-
associated isolation membrane (Figure 6) suggests that the receptors for chloroplast autophagy
connect the isolation membrane to the chloroplast envelope. Such interaction might facilitate the
localization of the isolation membrane on the region of subsequent chloroplast budding and its
development along the chloroplast surface. In budding yeast, the accumulation of the mitophagy
receptor Atg32 is required for the formation of the mitophagosome. Atg32 binds to the
mitochondrial outer membrane and interacts with Atg8 and Atg11, the latter being a scaffold
protein recruiting other ATG members, for autophagosome formation. In fission yeast
(*Schizosaccharomyces pombe*), Atg43 located on the mitochondrion outer membrane binds to
Atg8 to stabilize the autophagosomal membrane for mitophagosome formation during
starvation (Fukuda et al., 2020). In mammalian mitophagy, five types of mitophagy receptors
have been identified that bind to LC3 (light chain 3) or GABARAP proteins, the mammalian
orthologs of ATG8 (Onishi et al., 2021). Whether similar receptors work for chloroplast
autophagy in mature leaves has not been established. Atg39 is the receptor for nucleophagy in
budding yeast (Mochida et al., 2015), during which the condensation of Atg39 enables the
protrusion of the nuclear membrane to be sequestered by the autophagosome (Mochida et al.,
2022). The accumulation of receptor proteins might contribute to chloroplast budding to form
RCBs.
Another important regulator of the selective autophagy of organelles is the
ubiquitination of the targets of this process. One well-characterized process is mammalian
mitophagy regulated by PTEN-induced kinase 1 (PINK1) and Parkin (Ohnishi et al., 2021).
PINK1 accumulates on the surfaces of damaged mitochondria, facilitating the ubiquitination of
the mitochondria by the E3 ubiquitin ligase Parkin. Subsequently, autophagy adapter proteins
connect the ubiquitin chain to LC3/GABARAP proteins for the selective elimination of the
ubiquitinated mitochondria via autophagosomes (Lazarou et al., 2015). Five autophagy adaptor
proteins have been identified in mammals: p62/Sequestosome-1 (SQSTM1), Next to BRCA1
gene 1 (NBR1), Tax1-binding protein 1 (TAX1BP1), nuclear dot protein52 (NDP52), and
optineurin (OPTN). Among these, a plant ortholog of NBR1 has been identified (Svenning et al.,
2011). A recent study found that Arabidopsis NBR1 accumulates on damaged chloroplasts to
induce their removal in their entirety (Lee et al., 2023). NBR1 may participate in the recognition
of damaged chloroplasts during the entire organelle–type degradation of chloroplasts by
chlorophagy. Another study demonstrated that NBR1 regulates the selective degradation of the
outer envelope–bound TOC protein complexes that control cytoplasm-to-plastid protein import
(Wan et al., 2023). TOC proteins are ubiquitinated under stress conditions such as ultraviolet-B
exposure or high temperature and are then degraded by autophagosomes through NBR1-
mediated recognition. These findings suggest that NBR1 plays an important role in the
degradation of damaged chloroplast proteins. Whether ubiquitination and NBR1 participate in
the degradation of portions of chloroplasts via RCBs under sugar-starvation conditions has not
been evaluated.
503
Roles of stromules
The molecular functions of stromules are not completely understood. We predicted a functional

link between stromules and RCBs such that stromule-associated structures are vesiculated and
become RCBs for autophagic degradation. However, our imaging assays revealed that
chloroplast stromule formation is not essential for RCB formation in mesophyll cells (Figure 6).
The use of the *sid2* mutant and NahG transgenic lines indicated the close association between
enhanced SA signaling and the elevated stromule formation observed in *atg* mutants (Figure 7).
This finding is consistent with the suggested role of stromules in inducing programmed cell
death as an SA-dependent immune response (Caplan et al., 2015). We attribute the enhanced
stromule formation in *atg* mutant leaves to the higher SA and ROS levels, alone or in
combination with the excess stromal fraction accumulating in these mutants due to impaired
RCB formation.
In root cells, an association of the isolation membrane with stromules of non-green
plastids was reported (Spitzer et al., 2015). Stromule formation is more common in the non-
green plastids of tissues without photosynthetic activity than in the mature chloroplasts in leaf
mesophyll cells (Ishida and Yoshimoto, 2008; Hanson and Sattarzadeh, 2008). Therefore, the
interaction between stromules and the isolation membrane might be important for the efficient
degradation of non-green plastids by autophagy. Our study focused on the degradation
mechanism of mature chloroplasts in leaves, since mature chloroplasts are rich in nutrients and
amino acids and their degradation is particularly important for plant nutrient recycling (Makino
and Osmond, 1991). There might also be differences in the degradation mechanism of plastids
among cell types.
526
Intracellular dynamics for the transport of RCBs
Soluble *N*-ethylmaleimide-sensitive factor attachment protein receptors (SNAREs) mediate
versatile membrane fusion events (Ito and Uemura, 2022). Multiple SNARE proteins participate
in the fusion of the lysosomal/vacuolar membrane to the autophagosomal membrane in
mammals and budding yeast. A recent study reported the involvement of the Arabidopsis
SNARE proteins, VESICLE-ASSOCIATED MEMBRANE PROTEIN 724 (VAMP724) and
VAMP726, in autophagosome formation (He et al., 2022). In this study, we observed an
interesting morphology of vacuolar membranes that incorporate RCBs into the vacuole (Figure
4). A similar phenomenon was previously observed during the transport of pexophagosomes
(autophagosomes containing peroxisome components) in Arabidopsis leaves (Oikawa et al.,
2022). Thus, the transient engulfment of an autophagosome by the vacuolar membrane might be
a common phenomenon for vacuolar incorporation in leaf mesophyll cells. However, the
underlying mechanisms are unknown. In mammals, two autophagosomal membrane–anchored
SNAREs, Syntaxin 17 (Stx17) and Ykt6, independently contribute to autophagosome–lysosome
fusion (Itakura et al., 2012; Matsui et al., 2018). Stx17 translocates to the closed autophagosome
transiently, enabling its fusion with the lysosome (Tsuboyama et al., 2016). Such a translocation
system might allow the recruitment of the vacuolar membrane for its fusion with the mature
autophagosomes containing RCBs in plant cells.
545
Concluding remarks
The current study revealed how a type of piecemeal autophagy transports chloroplast stroma
and envelope components into the vacuole for degradation in mature Arabidopsis leaves. The
key event is the development of the isolation membrane along the chloroplast surface, which
may lead to the budding and segmentation of the membrane-contact site of a chloroplast.
Proteins mediating chloroplast segmentation and the interaction between the chloroplast surface
and the isolation membrane remain to be uncovered. This study developed a live-cell tracking
method for piecemeal-type chloroplast autophagy, which will help future studies elucidate the
underlying mechanisms of the intracellular dynamics of this type of autophagy.
555

Materials and Methods
Plant materials

Arabidopsis (*Arabidopsis thaliana*) plants from the Columbia accession (Col) were used in this study. Plants were grown in soil in growth chambers at 23°C under a 12-h-light/12-h-dark photoperiod with illumination from fluorescent lamps or LEDs (90–130 μmol m^{-2} s^{-1}). The T-DNA-insertion mutants for *ATG5* (*atg5-1*; SAIL_129_B07), *ATG7* (*atg7-2*; GABI_655B06), *ATG2* (*atg2-1*; SALK_076727), *ATG10* (*atg10-1*; SALK_084434), and *DRP5B* (*arc5-2*; SAIL_71_D11) have been described previously (Doelling et al., 2002; Thompson et al., 2005; Miyagishima et al., 2006; Phillips et al., 2008; Yoshimoto et al., 2009). The *sid2-2* mutant and a transgenic line expressing NahG have been reported previously (Delaney et al., 1994; Yoshimoto et al., 2009). The transgenic plants expressing a construct encoding chloroplast stroma–targeted GFP (GFP fused to the transit peptide of RECA protein) under the control of the cauliflower mosaic virus (CaMV) 35S promoter (*Pro35S:CT-GFP*; Kohler et al., 1997), chloroplast stroma–targeted DsRed (DsRed fused to the transit peptide of RECA protein) from the CaMV 35S promoter (*Pro35S:CT-DsRed*; Ishida et al., 2021), RBCS2B-GFP from the Arabidopsis *RBCS2B* promoter (*ProRBCS:RBCS-GFP*; Ishida et al., 2008), VHP1-mGFP from the Arabidopsis *VHP1* promoter (*ProVHP1:VHP1-mGFP*; Segami et al., 2014), GFP-ATG8a from the Arabidopsis *UBQ10* promoter (*ProUBQ:GFP-ATG8a*; Nakamura et al., 2021b), and TOC64-mRFP from the Arabidopsis *TOC64* promoter (*ProTOC64:TOC64-mRFP*; Kusano et al., 2023) have been reported in previous studies. Transgenic plants expressing *RBCS2B-EYFP*, *RBCS2B-mRFP*, or *RBCS2B-tagRFP* from the *RBCS2B* promoter (*ProRBCS:RBCS-EYFP*, *ProRBCS:RBCS-mRFP*, and *ProRBCS:RBCS-tagRFP*) were generated as follows. A genomic fragment encompassing the promoter region and the full-length coding region of *RBCS2B* (At5g38420), cloned into the pENTR/D/TOPO vector (Ishida et al., 2008), was inserted into the Gateway vectors pGWB540, pGWB553, or pGWB559 (Nakagawa et al., 2007) via LR clonase (Invitrogen) reaction to generate constructs encoding RBCS2B-EYFP, RBCS2B-mRFP, or RBCS2B-tagRFP, respectively.

Transgenic plants expressing *ATPC1-tagRFP* (*ProATPC1:ATPC1-tagRFP*) were produced as follows. A genomic fragment containing the promoter region and full-length coding region of *ATPC1* (At4g04640) was amplified from Col-0 genomic DNA by PCR using PrimeSTAR DNA polymerase (TaKaRa) and the primers ATPC1_F (CACCCATGGAGAGGGCTCGTACCTTAC) and ATPC1_R (AACCTGTGCATTAGCTCCAG), cloned into pENTR/D-TOPO (Invitrogen), and then recombined into the vector pGWB559 via LR reaction. Transgenic plants expressing *KEA1-mRFP* (*ProKEA1:KEA1-mRFP*) were produced as follows. A genomic fragment containing the promoter region and full-length coding region of *KEA1* (At1g01790) was amplified from Col-0 genomic DNA by PCR using the primers KEA1_F (AGGAACCAATTCAGTCGACTCATGATCATAACAAGTCTC) and KEA1_R (AAAGCTGGGTCTAGATATCCGATTACGACTGTGCCTCCTTC), cloned into pENTR1A (Invitrogen) using NEBuilder HiFi DNA Assembly master mix (New England Biolabs), and recombined into the vector pGWB553 via LR reaction. The resulting construct was introduced into Arabidopsis plants by the floral dip method (Clough and Bent, 1998) using Agrobacterium (*Agrobacterium tumefaciens*) strain GV3101. Transgenic plants expressing two types of fluorescent markers and mutant plants expressing fluorescent markers were generated by crossing or additional transformations.

Live-cell imaging by confocal microscopy

The second rosette leaves of 20- to 24-d-old plants were used for time-lapse imaging of living cells. The leaves were excised, infiltrated with sugar-free solutions, and incubated in darkness for 5 to 24 h, followed by confocal microscopy observations. When the dark treatment was started at the end of the night, the incubation time was around 5–9 h. When the treatment was

started during the light period, the incubation time was around 20–24 h. The sugar-free solutions
contained 10 mM MES-NaOH, pH 5.5, alone or with full-strength Murashige and Skoog salts
(Shiotani M. S.) or the photosynthesis inhibitor 3-(3,4-dichlorophenyl)-1,1-dimethylurea
(DCMU).
Time-lapse imaging in Figure 1 was performed as previously described with a two-
photon excitation confocal microscope with a spinning-disk unit (Otomo et al. 2015). The GFP
signal was excited by 920-nm femtosecond light pulses generated by a mode-locked titanium-
sapphire laser light source (Mai Tai eHP DeepSee; Spectra Physics). The YFP, RFP, and
chlorophyll signals were excited by 1040-nm femtosecond light pulses generated by an
ytterbium laser light source (femtoTrain; Spectra Physics). The fluorescent signals were
observed under an inverted microscope (IX-71; Olympus) equipped with a spinning-disk
scanner with 100-µm-wide pinholes aligned with a Nipkow disk (CSU-MPφ100; Yokogawa
Electric) and a water-immersion lens (UPLSAPO60XW, numerical aperture [NA] = 1.20,
Olympus). The fluorescence images were captured by an EM-CCD camera (EM-C2; Qimaging
or iXon Ultra 897; Andor Technology) through a bandpass filter for GFP and YFP (BrightLine
528/38; Semrock), RFP (D630/60M; Chroma Technology), or chlorophyll autofluorescence
(BrightLine 685/40; Semrock). For the simultaneous detection of YFP and chlorophyll signals,
fluorescence was detected through image-splitting optics (W-View Gemini; Hamamatsu
Photonics) including a dichroic mirror (FF580-FDi01-25 36; Semrock) and the bandpass filters.
Z-Scans were performed with a piezo actuator (P-721; PI). The acquired images were processed
and analyzed using NIS-Elements C software (Nikon) or Imaris software (Bitplane).
Time-lapse imaging analysis by confocal laser-scanning microscopy was performed
with LSM800 (Carl Zeiss), LSM880 (Carl Zeiss), LSM900 (Carl Zeiss), or SP8 (Leica) systems.
A water-immersion objective lens (C-Apochromat 40x, NA = 1.2 or LD C-Apochromat 63x, NA
= 1.15; Carl Zeiss) or an oil-immersion objective lens (HC PL APO 63x, NA = 1.40, Leica) was
used. Fast Airyscan mode was used for the observation displayed in Figure 4B with the
LSM880 system.
634
Quantification of microscopy images
Confocal images used for the quantification of RCB numbers (Figure 1–figure supplement 1,
Figure 5, Figure 8), chloroplast protrusions (Figure 5), stromules (Figure 7), or vacuolar RFP
intensity (Figure 8–figure supplement 2) were acquired with a C2 system (Nikon) equipped with
a water-immersion objective lens (CFI Apochromat LWD Lambda S 40XC, NA = 1.15; Nikon).
RFP emission was detected at 580–630 nm (bandpass filter RPB580–630; Omega optical) after
excitation with a 559.8-nm diode laser; chlorophyll autofluorescence was detected at 660–720
nm following excitation by a 636.5-nm diode laser; the two signals were detected
simultaneously. For the simultaneous detection of GFP and chlorophyll, GFP emission was
detected at 500–550 nm (bandpass filter RPB500–550; Omega Optical) following excitation by
a 489.6-nm diode laser; chlorophyll autofluorescence was detected at 660–720 nm following
excitation by a 489.6-nm diode laser.
For the quantification of RCBs and chloroplast protrusions, the second rosette leaves
of 21-d-old plants were excised, infiltrated with 10 mM MES-NaOH, pH 5.5, containing 1 µM
concanamycin A (Santa Cruz), and incubated for 1 d in darkness at 23°C. The stock solution
was 100 µM concanamycin A in DMSO. Sucrose (1%, w/v) or full-strength MS salts was added
as energy source or nutrients, respectively. The number of accumulated RCBs in a fixed area
(215.04 × 215. 04 µm each) was counted. The mean number from four areas of one second
rosette leaf per plant was calculated from four or five independent plants. The z-stack images of
the 3D region (215.04 × 215.04 × 15 µm each) were observed, and the proportion of
chloroplasts forming protrusions out of 50 chloroplasts was scored in the region. The protruding
structures of chloroplast stroma (approximately 0.5 to 1.5 µm long) containing the chloroplast
stroma marker RBCS-mRFP without the chlorophyll signal were defined as chloroplast

protrusions in Figure 5. The mean from two different regions of one second rosette leaf per plant
was calculated from four independent plants.
Chloroplast stromule formation in guard cells was observed in the second rosette
leaves of 13-d-old seedlings. The proportion of chloroplasts forming stromules in each pair of
guard cells was scored from the z-stack images ($251.04 \times 251.04 \times 20$ μm each). The mean
from 10 stomata of one second rosette leaf per plant was calculated from three individual
seedlings. Chloroplast stromule formation in mesophyll cells was observed in the third rosette
leaves of 20- or 36-d-old plants. The proportion of chloroplasts forming stromules out of 50
chloroplasts was scored in the z-stack images ($215.04 \times 215.04 \times 15$ μm each). The mean from
two different regions of one third rosette leaf per plant was calculated from four individual
plants. Thin, extended tubular structures (less than 1 μm in diameter) containing a chloroplast
stroma marker (CT-GFP or RBCS-mRFP) without chlorophyll signals were defined as
stromules, as described previously (Hanson and Sattarzadeh, 2011; Brunkard et al., 2015). For
the quantification of vacuolar RFP intensity, the second rosette leaves of 21-d-old plants were
excised and incubated in 10 mM MES-NaOH, pH 5.5, for 2 d in darkness at 23°C. The RFP
intensity in the central area of a mesophyll cell (15.2×15.2 μm each) was measured. The mean
from 12 cells of one second rosette leaf per plant was calculated from four individual plants.
Confocal images used to evaluate chloroplast stroma volume (Figure 1-figure
supplement 2), chloroplast buds, and GFP-ATG8a-labeled structures (Figure 2–figure
supplement 1; Figure 6–figure supplement 3 and 4) were acquired with an LSM900 (Carl Zeiss)
or SP8 (Leica) system. To quantify chloroplast stroma volume, the second rosette leaves of 21-
d-old plants accumulating RBCS-mRFP were excised at dawn, infiltrated with 10 mM MES-
NaOH, pH 5.5, and incubated for 24 h in the dark at 23°C. The z-stack images of the 3D region
($159.73 \times 159.73 \times 30$ μm each) were acquired, and the volumes of the respective chloroplasts
whose entire bodies were captured in the image and could be separated from neighboring
chloroplasts as a single structure were measured using Imaris software (Bitplane). Two different
regions of one second rosette leaf were observed in five individual plants. Leaves were observed
before incubation in the dark as a control. To quantify chloroplast buds, the second rosette
leaves of 21- to 23-d-old plants were excised at dawn, infiltrated with 10 mM MES-NaOH, pH
5.5, and incubated for 5 to 9 h in the dark at 23°C. The regions with RBCS-GFP, TOC64-mRFP,
KEA1-mRFP, ATPC1-tagRFP, or chlorophyll signals within a chloroplast bud and its
neighboring chloroplast or chloroplast envelope were selected as the regions of interest, and the
fluorescence intensities per unit area were measured in eight individual plants (Figure 2–figure
supplement 1). The number of chloroplast buds surrounded by GFP-ATG8a signal in four z-
stack images ($159.73 \times 159.73 \times 20$ μm each) of one leaf per plant was scored from five to six
individual plants (Figure 6–figure supplement 3D and 3E).
694
695
Protein analysis
The second and third rosette leaves of 21-d-old plants were incubated in darkness in 10 mM
MES-NaOH, pH 5.5, for 2 d and then frozen in liquid nitrogen. The leaves from 23-d-old,
untreated plants were used as control. The frozen leaves were homogenized using a tissue laser
(Qiagen) and a zirconium bead, before being resuspended in homogenization buffer containing
50 mM HEPES-NaOH, pH 7.5, 16 mM DTT, 10% (v/v) glycerol, and protease inhibitor cocktail
(Nacalai). Following centrifugation at 20,630 g for 10 min at 4°C, the protein amounts in the
supernatants were measured using a 660-nm Protein Assay Reagent (Pierce). The supernatants
were then mixed with an equal volume of SDS sample buffer containing 200 mM Tris-HCl, pH
8.5, 20% (v/v) glycerol, 2% (w/v) SDS, and 0.1 M DTT, and incubated for 5 min at 95°C. An
equal amount of protein was subjected to SDS-PAGE using TGX FastCast acrylamide gels
(Bio-Rad) and transferred to nitrocellulose membrane (Trans-blot turbo transfer pack; Bio-Rad).
An anti-RFP 1G9 clone antibody (1:2000, M204-3; MBL) and an anti-cFBPase antibody

(1:5000, AS04043; Agrisera) were used as primary antibodies. Goat anti-mouse IgG (H+L)
secondary antibody DyLight 800 4X PEG (1:10,000, SA5-35521; Invitrogen) or anti-rabbit
HRP secondary antibody (1:10,000, NA934; Cytiba) was used for RFP or cFBPase detection,
respectively. The chemiluminescence signals developed with SuperSignal West Dura Extended
Duration Substrate (Pierce) and the DyLight 800 fluorescent signals were detected by a
ChemiDoc MP system (Bio-Rad). The image processing and the quantification of band intensity
were performed using Image Lab Software (Bio-Rad).
716
717 **H_2O_2 measurements**
718 The amount of H_2O_2 in leaf lysates was measured as previously described (Chakraborty et al.,
719 2016) with an Amplex Red Hydrogen Peroxide/Peroxidase Assay Kit (Invitrogen) and an
720 Infinite 200 PRO plate reader (Tecan).
721
Statistical analysis
Statistical analysis in this study was performed with JMP14.3.0 software (SAS Institute).
Student's *t*-test or Tukey's test was used to compare paired samples or multiple samples,
respectively.

Acknowledgements

We thank Emi Sone, Izumi Fukuhara, and Mio Tokuda for their technical support. We thank Dr.
Kohki Yoshimoto for the use of *atg*, *sid2*, and NahG plants, Dr. Maureen R. Hanson for the use
of *Pro35S:CT-GFP* plants, Dr. Shoji Segami for the use of the *ProVHP1:VHP1-mGFP* construct,
and Dr. Tsuyoshi Nakagawa for the use of pGWB vectors.

Figure legends
733
Figure 1. Chloroplast buds are released in sugar-starved leaves.
Time-lapse observations of 3D-reconstructed chloroplast morphology in Arabidopsis mesophyll
cells accumulating chloroplast stroma–targeted fluorescent markers. A leaf from a plant
accumulating chloroplast stroma–targeted GFP (CT-GFP) (A), RBCS-mRFP (B), or RBCS-
EYFP (C) was incubated in sugar-free solution in darkness for 5 h from dawn (A), 21 h (B), or
24 h from the period in light (C) and then observed through a two-photon excitation microscope
equipped with a confocal spinning-disk unit. Second rosette leaves from 20- to 22-d-old plants
were used. Images in (A–C) are still frames from Videos 1, 2, and 3, respectively. Time scales
above the images indicate the elapsed time from the start of the respective videos. Arrowheads
indicate chloroplast budding structures. Scale bars, 5 µm. In (C), green, RBCS-EYFP; magenta,
chlorophyll fluorescence. In the merged images, the overlapping regions of RBCS-EYFP and
chlorophyll signals appear white.
746
Figure 1–figure supplement 1. Accumulation of chloroplast stroma components in the
vacuole via autophagy.
Confocal images of mesophyll cells from wild-type (A) or *atg7* (B) plants accumulating the
chloroplast stroma marker RBCS-mRFP. Second rosette leaves of 21-d-old plants were excised
and incubated in 10 mM MES-NaOH containing 1 µM concanamycin A (concA) in darkness for
1 d. Sucrose (Suc) or Murashige and Skoog (MS) salts were added as an energy or nutrient
source, respectively. Second rosette leaves from nontreated plants are shown as control. Green,
RBCS-mRFP; magenta, chlorophyll fluorescence. Scale bars, 10 µm. The small puncta
containing RBCS-mRFP without chlorophyll signal appear green and are Rubisco-containing
bodies (RCBs) in the vacuole. (C) Number of accumulated RCBs from the observations
described in (A) and (B). Different lowercase letters denote significant differences based on
Tukey's test ($P < 0.05$). Values are means ± SE ($n = 4$). Dots represent individual data points.

Figure 1–figure supplement 2. The release of chloroplast buds by autophagy contributes to the decline in chloroplast stroma volume.

(A) Orthogonal projections produced from *z*-stack images (30 μm in depth) of mesophyll cells from WT or *atg7* leaves accumulating the chloroplast stroma marker RBCS-mRFP. Second rosette leaves of 21-d-old plants were excised at dawn and incubated in 10 mM MES-NaOH in the dark for 24 h. Mesophyll cells were observed before (0 h) and after (Dark 24 h) the treatment. Green, RBCS-mRFP. Scale bars, 10 μm. (B) Violin plot showing the volume of chloroplast stroma labeled by RBCS-mRFP, measured from the observations described in (A). Different lowercase letters denote significant differences based on Tukey's test ($P < 0.05$). Yellow bars indicate mean values. Five individual plants were observed, and the volumes of 564 (WT, 0 h), 554 (WT, dark 24 h), 597 (*atg7*, 0 h), and 725 (*atg7*, dark 24 h) chloroplasts were scored.

Video 1. Release of a chloroplast bud as visualized by chloroplast stroma–targeted GFP.

The second rosette leaf from a 21-d-old plant accumulating the chloroplast stroma–targeted GFP (CT-GFP) was incubated in sugar-free solution in darkness for time-lapse imaging with a two-photon excitation microscope equipped with a confocal spinning-disk unit. Arrowhead indicates a chloroplast budding structure. 3D-reconstructed images (8 μm in depth) acquired about every 3 s are displayed at 10 frames/s. Scale bar, 5 μm. This video was used to generate Figure 1A.

Video 2. Release of a chloroplast bud as visualized by RBCS-mRFP

The second rosette leaf from a 22-d-old plant accumulating the chloroplast stroma marker RBCS-mRFP was incubated in sugar-free solution in darkness for time-lapse imaging with a two-photon excitation microscope equipped with a confocal spinning-disk unit. Arrowhead indicates a chloroplast budding structure. 3D-reconstructed images (2 μm in depth) acquired about every 1.2 s are displayed at 10 frames/s. Scale bar, 5 μm. This video was used to generate Figure 1B.

Video 3. Tracking of the stroma marker and chlorophyll fluorescence during the release of a chloroplast bud.

The second rosette leaf from a 20-d-old plant accumulating the chloroplast stroma marker RBCS-EYFP was incubated in sugar-free solution in darkness for time-lapse imaging with a two-photon excitation microscope equipped with a confocal spinning-disk unit. Arrowhead indicates a chloroplast budding structure. 3D-reconstructed images (8 μm in depth) acquired every 4 s are displayed at 10 frames/s. Scale bar, 5 μm. Green, RBCS-EYFP; magenta, chlorophyll fluorescence. Only the video of the merged channels is shown. The overlapping regions of RBCS-EYFP and chlorophyll signals appear white. This video was used to generate Figure 1C.

Figure 2. Chloroplast buds containing stroma and envelope components are released from the chloroplasts.

Time-lapse observations of Arabidopsis mesophyll cells accumulating the chloroplast stroma marker along with an envelope marker or a thylakoid membrane marker. Leaves accumulating stromal RBCS-GFP along with outer envelope–bound TOC64-mRFP (A and B) or with thylakoid membrane–bound ATPC1-tagRFP (C and D) were incubated in sugar-free solution in darkness for 5 to 9 h from dawn and then observed. Second rosette leaves from 21- to 22-d-old plants were used. Images in (B) or (D) are still frames from Videos 4 and 6, respectively. Time scales above the images indicate the elapsed time from the start of the respective videos. Arrowheads indicate chloroplast budding structures. Scale bars, 5 μm. Green, TOC64-mRFP or ATPC1-tagRFP; magenta, RBCS-GFP; orange, chlorophyll (Chl) fluorescence. The graphs in

(A and C) show fluorescence intensities along the blue lines (a to b) in the magnified images of the area indicated by dashed blue boxes. The intensities are shown relative to the maximum intensity for each fluorescence channel, set to 1.

Figure 2–figure supplement 1. Chloroplast buds contain stroma and envelope proteins.
Second rosette leaves accumulating stromal RBCS-GFP along with outer envelope membrane–bound TOC64-mRFP (A), inner envelope membrane–bound KEA1-mRFP (B), or thylakoid membrane–bound ATPC1-tagRFP (C) were incubated in sugar-free solution in the dark for 5 to 9 h from dawn, and chloroplast budding structures were observed. Second rosette leaves from 21- to 22-d-old plants were used. Arrowheads indicate chloroplast budding structures. Four representative images of chloroplast buds are shown for each plant line. Scale bars, 5 μm. Green, TOC64-mRFP, KEA1-mRFP, or ATPC1-tagRFP; magenta, RBCS-GFP; orange, chlorophyll (Chl) fluorescence. The graphs show the relative ratio of GFP, RFP, or chlorophyll fluorescence intensity per unit area within chloroplast budding structures vs. their neighboring chloroplasts (bud/chloroplast). The intensities from 33 (A), 32 (B), or 30 (C) chloroplast buds from 8 individual plants were measured. Values are means ± SE (n = 30–33). Dots represent individual data points in each graph.

Figure 2–figure supplement 2. A released chloroplast bud contains an inner envelope marker protein.
Time-lapse observations of Arabidopsis mesophyll cells accumulating the chloroplast stroma marker RBCS-GFP along with inner envelope membrane–bound KEA1-mRFP. The second rosette leaf from a 22-d-old plant was incubated in sugar-free solution in the dark for 6 h from dawn and observed. The images are still frames from Video 5. Time scales above the images indicate the elapsed time from the start of the video. Arrowheads indicate chloroplast budding structures. Scale bars, 5 μm. Green, KEA1-mRFP; magenta, RBCS-GFP; orange, chlorophyll (Chl) fluorescence.

Video 4. A released chloroplast bud contains the outer envelope marker TOC64-mRFP.
The second rosette leaf from a 21-d-old plant accumulating the chloroplast stroma marker RBCS-GFP and the outer envelope marker TOC64-mRFP was incubated in sugar-free solution in darkness for time-lapse imaging. Arrowheads indicate chloroplast budding structures. Images acquired every 2 s are displayed at 10 frames/s. Scale bar, 5 μm. Green, TOC64-mRFP; magenta, RBCS-GFP. The video contains TOC64-mRFP and the merged channels. This video was used to generate Figure 2B.

Video 5. A released chloroplast bud contains the inner envelope marker KEA1-mRFP.
The second rosette leaf from a 22-d-old plant accumulating the chloroplast stroma marker RBCS-GFP and the inner envelope marker KEA1-mRFP was incubated in sugar-free solution in the dark for time-lapse imaging. Arrowheads indicate chloroplast budding structures. Images acquired every 2 s are displayed at 10 frames/s. Scale bar, 5 μm. Green, KEA1-mRFP; magenta, RBCS-GFP. The video contains KEA1-mRFP and the merged channels. This video was used to generate Figure 2–figure supplement 2.

Video 6. A released chloroplast bud does not contain the thylakoid membrane marker ATPC1-tagRFP.
The second rosette leaf from a 21-d-old plant accumulating the chloroplast stroma marker RBCS-GFP and the thylakoid membrane marker ATPC1-tagRFP was incubated in sugar-free solution in darkness for time-lapse imaging. Arrowheads indicate chloroplast budding structures. Images acquired every 2 s are displayed at 10 frames/s. Scale bar, 5 μm. Green, ATPC1-tagRFP; magenta, RBCS-GFP. The video contains ATPC1-tagRFP and the merged channels. This video

861   was used to generate Figure 2D.
862

Figure 3. Tracking the transport of a Rubisco-containing body.
A leaf accumulating chloroplast stroma–targeted GFP (CT-GFP) was incubated in sugar-free
solution in darkness for 6 h from dawn, and the transport of a Rubisco-containing body (RCB)
was tracked. The second rosette leaf from a 21-d-old plant was used. (A) Confocal images
during the periods when the RCB moved quickly (24.4–34.8 and 73.8–90.5 s). Arrowheads
indicate an RCB. The images are still frames from Video 7. Time scales above the images
indicate the elapsed time from the start of the video. Green, CT-GFP; magenta, chlorophyll
(Chl) fluorescence. Scale bar, 5 µm. (B) Calculated speed of the tracked RCB in (A). (C) The
track of the RCB. The color of the track line changes over time, as indicated by the color bar.
Scale bar, 2 µm.

Video 7. Tracking of a Rubisco-containing body marked by CT-GFP.
The second rosette leaf from a 21-d-old plant accumulating chloroplast stroma–targeted GFP
(CT-GFP) was incubated in sugar-free solution in darkness for time-lapse monitoring of an RCB
marked by CT-GFP. Arrowhead indicates an RCB. Images acquired every 0.52 s are displayed
at 20 frames/s. Scale bar, 5 µm. Green, CT-GFP; magenta, chlorophyll fluorescence. Only the
video of the merged channels is shown. This video was used to generate Figure 3A.

**Figure 3–figure supplement 1. Rubisco-containing bodies appear to begin random
movement during their tracking.**
A leaf accumulating RBCS-tagRFP (A) or RBCS-EYFP (B) was incubated in sugar-free
solution in the dark for 7 to 8 h from dawn, and the transport of a Rubisco-containing body was
tracked. Second rosette leaves from 22-d-old plants were used. Arrowheads indicate a tracked
RCB. The images are still frames from Videos 8 (A) or 9 (B), respectively. The images show the
time when RCBs appeared to begin random movement (185–197 s in A or 25–33 s in B). Time
scales above the images indicate the elapsed time from the start of the respective videos. Green,
RBCS-tagRFP or RBCS-EYFP; magenta, chlorophyll fluorescence. Scale bars, 5 µm. The color
of the track line changes over time, as indicated by the color bars below the images.

Video 8. Tracking of a Rubisco-containing body marked by RBCS-tagRFP.
The second rosette leaf from a 22-d-old plant accumulating stromal RBCS-tagRFP was
incubated in sugar-free solution in the dark for time-lapse monitoring of an RCB marked by
RBCS-tagRFP. Arrowhead indicates an RCB. Images acquired every 1 s are displayed at 20
frames/s. Scale bar, 5 µm. Green, RBCS-tagRFP; magenta, chlorophyll fluorescence. Only the
video of the merged channels is shown. This video was used to generate Figure 3–figure
supplement 1A.

Video 9. Tracking of a Rubisco-containing body marked by RBCS-EYFP.
The second rosette leaf from a 22-d-old plant accumulating stromal RBCS-EYFP was incubated
in sugar-free solution in the dark for time-lapse monitoring of an RCB marked by RBCS-EYFP.
Arrowhead indicates an RCB. Images acquired every 1 s are displayed at 10 frames/s. Scale bar,
5 µm. Green, RBCS-EYFP; magenta, chlorophyll fluorescence. Only the video of the merged
channels is shown. This video was used to generate Figure 3–figure supplement 1B.

**Figure 4. Dynamics of the vacuolar membrane during the incorporation of Rubisco-
containing bodies.**
Leaves accumulating the chloroplast stroma marker RBCS-mRFP along with the vacuolar
membrane marker VHP1-mGFP were incubated in sugar-free solution in darkness for 6 to 8 h
from dawn, and the behavior of cytosolic RCBs was monitored. Second rosette leaves from 21-

or 23-d-old plants were used. The images when the vacuolar membrane engulfs an RCB (25.4–
30.5 s in A and 52.6–57.4 s in B) and when an RCB is incorporated into the vacuolar lumen
(47.0–52.0 s in A and 65.3–72.9 s in B) are shown. The images in (A) and (B) are still frames
from Videos 10 and 11, respectively. Time scales above the images indicate the elapsed time
from the start of the respective videos. Open arrowheads indicate an RCB engulfed by the
vacuolar membrane. Closed arrowheads indicate the open site of the vacuolar membrane for the
incorporation of an RCB. V indicates the region of the vacuolar lumen. Green, VHP1-mGFP;
magenta, RBCS-mRFP. Scale bars, 5 μm.

Video 10. Incorporation of a Rubisco-containing body into the vacuolar lumen.
The second rosette leaf from a 21-d-old plant accumulating the chloroplast stroma marker
RBCS-mRFP along with the vacuolar membrane marker VHP1-mGFP was incubated in sugar-
free solution in darkness for time-lapse imaging. Arrowhead indicates an RCB incorporated into
the vacuolar lumen. Images acquired every 1.27 s are displayed at 10 frames/s. Scale bar, 5 μm.
Green, VHP1-mGFP; magenta, RBCS-mRFP. Only the video of the merged channels is shown.
This video was used to generate Figure 4A.

Video 11. Video 2 of the vacuolar incorporation of a Rubisco-containing body.
930   Additional time-lapse data for the experiment described in Figure 4 and Video 10. The second
931   rosette leaf from a 23-d-old plant was used. Arrowhead indicates an RCB incorporated into the
932   vacuolar lumen. Images acquired every 0.317 s are displayed at 25 frames/s. Scale bar, 5 μm.
933   Green, VHP1-mGFP; magenta, RBCS-mRFP. Only the video of the merged channels is shown.
934   This video was used to generate Figure 4B.

**Figure 4–figure supplement 1. Additional observations of the incorporation of Rubisco-
containing bodies into the vacuolar lumen.**
Additional time-lapse data of the experiments described in Figure 4. The second rosette leaves
from 21- (A) or 24-d-old (B, C) plants were used. The images in (A), (B), and (C) are still
frames from Videos 12, 13, and 14, respectively. Time scales above the images indicate the
elapsed time from the start of the respective videos. Open arrowheads indicate an RCB engulfed
by the vacuolar membrane. Closed arrowheads indicate the vacuolar membrane structure that
engulfs an RCB. V indicates the region of the vacuolar lumen. Green, VHP1-mGFP; magenta,
RBCS-mRFP. Scale bars, 5 μm.

Video 12. Video 3 of the vacuolar incorporation of a Rubisco-containing body.
947   Additional time-lapse data for the experiment described in Figure 4 and Video 10. Arrowhead
948   indicates an RCB incorporated into the vacuolar lumen. Images acquired every 1.5 s are
949   displayed at 10 frames/s. Scale bar, 5 μm. Green, VHP1-mGFP; magenta, RBCS-mRFP. Only
950   the video of the merged channels is shown. This video was used to generate Figure 4–figure
951   supplement 1A.

Video 13. Video 4 of the vacuolar incorporation of a Rubisco-containing body.
954   Additional time-lapse data for the experiment described in Figure 4 and Video 10. Arrowhead
955   indicates an RCB incorporated into the vacuolar lumen. Images acquired every 1.5 s are
956   displayed at 10 frames/s. Scale bar, 5 μm. Green, VHP1-mGFP; magenta, RBCS-mRFP. Only
957   the video of the merged channels is shown. This video was used to generate Figure 4–figure
958   supplement 1B.

Video 14. Video 5 of the vacuolar incorporation of a Rubisco-containing body.
961   Additional time-lapse data for the experiment described in Figure 4 and Video 10. Arrowhead
962   indicates an RCB incorporated into the vacuolar lumen. Images acquired every 1.5 s are

963   displayed at 10 frames/s. Scale bar, 5 µm. Green, VHP1-mGFP; magenta, RBCS-mRFP. Only
964   the video of the merged channels is shown. This video was used to generate Figure 4–figure
965   supplement 1C.
966
Figure 5. Autophagy deficiency does not increase the number of chloroplast protrusions
during a 1-d dark treatment.
Leaves from wild-type (WT), *atg5*, or *atg7* plants accumulating the chloroplast stroma marker
RBCS-mRFP were incubated in sugar-free solution containing 1 µM concanamycin A (concA)
for 1 d in darkness. Second rosette leaves from 21-d-old plants were used. 2D images of
mesophyll cells were acquired (A), and the number of accumulated RCBs in the vacuoles was
scored (B). Small puncta containing RBCS-mRFP without the chlorophyll signal appear green
and are RCBs in the vacuole. Leaves from untreated plants are shown as control. The
appearance of chloroplast protrusions was observed from orthogonal projections created from *z*-
stack images (15 µm in depth; C), and the proportion of chloroplasts forming protrusion
structures out of 50 chloroplasts was scored (D). Scale bars, 10 µm. Green, RBCS-mRFP;
magenta, chlorophyll fluorescence. Only the merged channels are shown. The overlapping
regions of RBCS-mRFP and chlorophyll signals appear white. Arrowheads indicate the
structures that were counted as a chloroplast protrusion in (D). Different lowercase letters
denote significant differences based on Tukey's test ($P < 0.05$). Values are means ± SE ($n = 4$).
Dots represent individual data points in each graph.
983
Figure 5–figure supplement 1. Chloroplast protrusions do not increase in *atg2* or *atg10*
mutant leaves during a 1-d dark treatment.
The experiments described in Figure 5 were performed on leaves from wild-type (WT), *atg2*, or
atg10 plants accumulating the chloroplast stroma marker RBCS-mRFP. Scale bars, 10 µm.
Green, RBCS-mRFP; magenta, chlorophyll fluorescence. Only the merged channels are shown.
The overlapping regions of RBCS-mRFP and chlorophyll signals appear white. Small puncta
containing RBCS-mRFP without chlorophyll signal appear green and are RCBs in the vacuole.
Arrowheads indicate the structures that were counted as a chloroplast protrusion in (D).
Different lowercase letters denote significant differences based on Tukey's test ($P < 0.05$).
Values are means ± SE ($n = 4$). Dots represent individual data points in each graph.
994
Figure 6. The formation of a chloroplast bud and the maturation of the chloroplast-
associated isolation membrane occur concomitantly.
Leaves accumulating the chloroplast stroma marker, RBCS-mRFP (A) or CT-DsRed (B), and
the isolation membrane marker GFP-ATG8a were incubated in sugar-free solution in darkness
for 6 h from dawn (A) or 21 h from the period in light (B) and then observed. Second rosette
leaves from 23-d-old plants were used. Arrowheads indicate the position of the chloroplast-
associated isolation membrane. Images in (A) and (B) are still frames from Videos 15 and 16,
respectively. Time scales above the images indicate the elapsed time from the start of the
respective videos. Green, GFP-ATG8a; magenta, RBCS-mRFP or CT-DsRed. Scale bars, 5 µm.
(C) Time-dependent changes in the ratio of the major axis to the minor axis in the GFP-ATG8a-
labeled isolation membrane (top) or in the area of the chloroplast budding structure (bottom) as
measured from the images in (B).
1007
Video 15. Budding of the isolation membrane–associated site within a chloroplast.
The second rosette leaf from a 23-d-old plant accumulating the chloroplast stroma marker
RBCS-mRFP and the isolation membrane marker GFP-ATG8a was incubated in sugar-free
solution in darkness for time-lapse imaging. Arrowheads indicate the portion of the chloroplast-
associated isolation membrane. Images acquired every 4 s are displayed at 10 frames/s. Scale
bar, 5 µm. Green, GFP-ATG8a; magenta, RBCS-mRFP. The video contains RBCS-mRFP and

the merged channels. This video was used to generate Figure 6A.

Video 16. Autophagosome development and chloroplast segmentation occur concomitantly.
The second rosette leaf from a 23-d-old plant accumulating the chloroplast stroma–targeted
DsRed (CT-DsRed) and the isolation membrane marker GFP-ATG8a was incubated in sugar-
free solution in darkness for time-lapse imaging. Arrowhead indicates the portion of the
chloroplast-associated isolation membrane. Images acquired every 1 s are displayed at 20
frames/s. Scale bar, 5 μm. Green, GFP-ATG8a; magenta, CT-DsRed. Only the video of the
merged channels is shown. This video was used to generate Figure 6B.

Figure 6–figure supplement 1. Another observation of the budding of the isolation
membrane–associated site in a chloroplast.
A leaf accumulating the chloroplast stroma marker CT-DsRed and the isolation membrane
marker GFP-ATG8a was incubated in sugar-free solution in darkness for 6 h from dawn and
then observed. A second rosette leaf from a 22-d-old plant was used. Arrowheads indicate the
position of the chloroplast-associated isolation membrane. The images are still frames from
Video 17. Time scales above the images indicate the elapsed time from the start of the video.
Green, GFP-ATG8a; magenta, CT-DsRed. Scale bars, 5 μm. (B) Time-dependent changes in the
ratio of the major axis to the minor axis in the GFP-ATG8a-labeled isolation membrane (top) or
in the area of the chloroplast budding structure (bottom), as measured from the images in (A).

Figure 6–figure supplement 2. Chloroplast buds surrounded by the isolation membrane
appear in multiple chloroplasts.
A leaf accumulating the chloroplast stroma marker RBCS-tagRFP and the isolation membrane
marker GFP-ATG8a was incubated in sugar-free solution in darkness for 5 h from dawn and
then observed. A second rosette leaf from a 21-d-old plant was used. White, yellow, and blue
arrowheads indicate the isolation membrane–associated site in different chloroplasts,
respectively. The images are still frames from Video 18. Time scales above the images indicate
the elapsed time from the start of the video. Green, GFP-ATG8a; magenta, RBCS-tagRFP;
orange, chlorophyll (Chl) fluorescence. Scale bars, 5 μm.

Video 17. Another time-lapse assay showing the concomitant progression of
autophagosome development and chloroplast segmentation.
The second rosette leaf from a 22-d-od plant accumulating the chloroplast stroma–targeted
DsRed (CT-DsRed) and the isolation membrane marker GFP-ATG8a was incubated in sugar-
free solution in darkness for time-lapse imaging. Arrowhead indicates the portion of the
chloroplast-associated isolation membrane. Images acquired every 2 s are displayed at 10
frames/s. Scale bar, 5 μm. Green, GFP-ATG8a; magenta, CT-DsRed. Only the video of the
merged channels is shown. This video was used to generate Figure 6–figure supplement 1.

Video 18. Autophagy-related chloroplast segmentation occurs in sequence.
The second rosette leaf from a 21-d-old plant accumulating the chloroplast stroma marker
RBCS-tagRFP and the isolation membrane marker GFP-ATG8a was incubated in sugar-free
solution in darkness for time-lapse imaging. White, yellow, and blue arrowheads indicate the
isolation membrane–associated sites in different chloroplasts. Images acquired every 2 s are
displayed at 10 frames/s. Scale bar, 5 μm. Green, GFP-ATG8a; magenta, RBCS-tagRFP. Only
the video of the merged channels is shown. This video was used to generate Figure 6–figure
supplement 2.

Figure 6–figure supplement 3. Proportion of chloroplast buds engulfed by GFP-ATG8a-
labeled isolation membranes.

Leaves accumulating the chloroplast stroma marker RBCS-mRFP and the isolation membrane
marker GFP-ATG8a were incubated in sugar-free solution in the dark for 5 to 9 h from dawn,
and chloroplast budding structures were observed. Second rosette leaves from 22- to 23-d-old
plants were used. (A and B) Four representative images of chloroplasts buds that are surrounded
by GFP-ATG8a signals (A) or two representative images of chloroplast buds that are not labeled
by GFP-ATG8a signals (B) are shown. Arrowheads indicate chloroplast budding structures.
Green, GFP-ATG8a; magenta, RBCS-mRFP. Scale bars, 5 μm. (C) The proportions of
chloroplast buds that are labeled by GFP-ATG8a signals (+) or not (−) were calculated from
observations of 58 chloroplast buds from 8 individual plants. (D) The number of GFP-ATG8a-
positive chloroplast buds, as shown in (A), were counted in the fixed regions from leaves
incubated in the dark for 5 to 9 h from dawn (Dark) or from untreated leaves of plants grown
under normal conditions (Control). Five individual plants were observed. (E) Leaves from wild-
type (WT) or *atg7* plants accumulating RBCS-mRFP and GFP-ATG8a were incubated in the
dark for 5 to 9 h from dawn, and the number of GFP-ATG8a-positive chloroplast buds was
counted. Six individual plants were observed. Values are means ± SE (n = 5–6). Asterisks
denote significant differences based on t-test (***, P < 0.001). Dots represent individual data
points in each graph.
1082
Figure 6–figure supplement 4. Proportion of GFP-ATG8a-labeled structures that are
associated with chloroplasts.
Leaves accumulating the chloroplast stroma marker RBCS-mRFP and the isolation membrane
marker GFP-ATG8a were incubated in sugar-free solution in the dark for 5 to 9 h from dawn,
and GFP-ATG8a-labeled structures in mesophyll cells were observed. Second rosette leaves
from 22-d-old plants were used. (A to C) GFP-ATG8a-positive structures were classified into
three types: flattened structures associated with the chloroplast surface (A), spherical or curved
structures associated with chloroplast budding structures (B), and other structures (C). Four
representative images of each type are shown. Green, GFP-ATG8a; magenta, RBCS-mRFP.
Scale bars, 5 μm. (D) The proportions of the three types of GFP-ATG8a-labeled structures were
calculated from observations of 157 GFP-ATG8a-labeled structures from 13 individual plants.
1094
Figure 7. Diminished salicylic acid signal suppresses stromule formation in autophagy-
deficient mutants.
(A) Orthogonal projections produced from z-stack images (10 μm in depth) of guard cells from
wild-type (WT), *atg5*, NahG, and *atg5* NahG leaves accumulating chloroplast stroma–targeted
GFP (CT-GFP). Second rosette leaves from 13-d-old plants were used. Scale bars, 10 μm. (B)
Percentage of chloroplasts forming stromules in 10 stomata, from the observations described in
(A). (C) Orthogonal projections produced from z-stack images (10 μm in depth) of guard cells
from WT, *atg5*, *atg7*, *sid2*, *sid2 atg5*, and *sid2 atg7* leaves accumulating CT-GFP. Scale bars, 10
μm. (D) Percentage of chloroplasts forming stromules in 10 stomata, from the observations
described in (C). (E) Orthogonal projections produced from z-stack images (15 μm in depth) of
mesophyll cells from WT, *atg5*, *atg7*, *sid2*, *sid2 atg5*, and *sid2 atg7* leaves accumulating CT-
GFP. Third rosette leaves from 36-d-old plants were observed. Green, CT-GFP; magenta,
chlorophyll fluorescence. Only the merged channels are shown. Scale bars, 20 μm. (F)
Percentage of chloroplasts forming stromules out of 50 chloroplasts in mesophyll cells, from the
observations described in (E). (G) Hydrogen peroxide (H_2O_2) content in third rosette leaves
from 36-d-old WT, *atg5*, *atg7*, *sid2*, *sid2 atg5*, and *sid2 atg7* plants. Different lowercase letters
denote significant differences based on Tukey's test (P < 0.05). Values are means ± SE (n = 3 in
B and D or 4 in F and G). Dots represent individual data points in each graph.
1113
1114 **Figure 7–figure supplement 1. Autophagy deficiency activates stromule formation from**
1115 **mesophyll chloroplasts accumulating RBCS-mRFP in senescing leaves.**

(A) Orthogonal projections produced from *z*-stack images (15 µm in depth) of mesophyll cells from leaves of WT, *atg5*, and *atg7* plants accumulating RBCS-mRFP. Third rosette leaves from 36-d-old plants were observed. Green, RBCS-mRFP; magenta, chlorophyll fluorescence. Only the merged channels are shown. Spherical bodies only exhibiting chlorophyll signal represent a portion of chloroplasts in pavement cells. Scale bars, 20 µm. (B) Percentage of chloroplasts forming stromules out of 50 chloroplasts in mesophyll cells, from the observations described in (A). Different lowercase letters denote significant differences based on Tukey's test ($P < 0.05$). Values are means ± SE ($n = 4$). Dots represent individual data points in the graph.

Figure 7–figure supplement 2. Autophagy deficiency does not activate stromule formation from mesophyll chloroplasts in young leaves.
Orthogonal projections produced from *z*-stack images (15 µm in depth) of mesophyll cells from WT, *atg5*, *atg7*, *sid2*, *sid2 atg5*, and *sid2 atg7* leaves accumulating CT-GFP. Third rosette leaves from 20-d-old plants were observed. Green, CT-GFP; magenta, chlorophyll fluorescence. Only the merged channels are shown. Scale bars, 20 µm.

Figure 7–figure supplement 3. NahG expression does not counteract the increase in hydrogen peroxide content produced by the *atg5* mutation.
Hydrogen peroxide (H_2O_2) contents in third rosette leaves from 36-d-old plants. WT plants without any transgenic construct and CT-GFP-expressing plants in the WT, *atg5*, NahG, and *atg5* NahG backgrounds were used. Different lowercase letters denote significant differences based on Tukey's test ($P < 0.05$). Values are means ± SE ($n = 4$). Dots represent individual data points.

Figure 8. DRP5b is dispensable for chloroplast autophagy in sugar-starved leaves.
(A) Confocal images of mesophyll cells from wild-type (WT) and *drp5b* leaves accumulating the stroma marker RBCS-mRFP. Second rosette leaves from 21-d-old plants were incubated in sugar-free solution containing 1 µM concanamycin A (concA) in darkness for 1 d. Green, RBCS-mRFP; magenta, chlorophyll fluorescence. Scale bars, 10 µm. (B) Number of accumulated RCBs in WT, *drp5b*, *atg5*, and *atg7* leaves, counted from the observations of leaves incubated in the dark with concA described in (A). Different lowercase letters denote significant differences based on Tukey's test ($P < 0.05$). Values are means ± SE ($n = 4$–5). (C) Biochemical detection of autophagy flux for chloroplast stroma based on a free RFP assay. RFP and cFBPase (loading control) were detected by immunoblotting of soluble protein extracts from leaves of WT, *drp5b*, *atg5*, and *atg7* plants accumulating RBCS-mRFP. Protein extracts from either untreated control leaves (cont) or leaves after 2 d of incubation in darkness (dark) were used. Total protein was detected by Coomassie Brilliant Blue (CBB) staining as a loading control. The filled arrowhead indicates RBCS-mRFP fusion, and the open arrowhead indicates free mRFP derived from the cleavage of RBCS-mRFP. (D) Quantification of the free mRFP/RBCS-mRFP ratio shown relative to that of untreated WT plants, which was set to 1. Asterisks denote significant differences based on *t*-test (***, $P < 0.001$; n.s., not significant). Values are means ± SE ($n = 4$). Dots represent individual data points in each graph.

Figure 8–figure supplement 1. Production of Rubisco-containing bodies in *drp5b* mutants is ATG5 dependent.
(A) Confocal images of mesophyll cells from wild-type (WT), *atg5*, *drp5b*, and *drp5b atg5* leaves accumulating the stroma marker CT-GFP. Second rosette leaves from 21-d-old plants were incubated in sugar-free solution containing 1 µM concanamycin A (concA) in darkness for 1 d. Second rosette leaves from untreated plants were used as control. Green, CT-GFP; magenta, chlorophyll fluorescence. Only the merged channels are shown. Small puncta containing CT-GFP without chlorophyll signal appear as green and are RCBs in the vacuole. Scale bars, 10 µm.

(B) Number of accumulated RCBs, counted from the observations of leaves incubated in the dark with concA described in (A). Different lowercase letters denote significant differences based on Tukey's test ($P < 0.05$). Values are means ± SE ($n = 5$). Dots represent individual data points.

Figure 8–figure supplement 2. Vacuolar accumulation of stromal marker proteins in sugar-starved leaves.
(A) Confocal images of mesophyll cells from wild-type (WT), *drp5b*, *atg5*, and *atg7* leaves accumulating the stroma marker RBCS-mRFP. Second rosette leaves from 21-d-old plants were incubated in sugar-free solution in darkness for 2 d. Second rosette leaves from untreated plants are shown as control. Green, RBCS-mRFP; magenta, chlorophyll fluorescence. Scale bars, 20 μm. (B) RFP intensity in the vacuolar lumen, as measured from the observations described in (A) and shown relative to that of untreated WT plants, which was set to 1. Asterisks denote significant differences based on *t*-test (***, $P < 0.001$; n.s., not significant). Values are means ± SE ($n = 4$). Dots represent individual data points.

Figure 9. Formation and segmentation of chloroplast buds in leaves of the *drp5b* mutant.
(A) A leaf from the *drp5b* mutant accumulating the stroma marker RBCS-mRFP was incubated in sugar-free solution in darkness for 5 h from dawn and then observed. Arrowheads indicate a chloroplast bud. Green, RBCS-mRFP; magenta, chlorophyll fluorescence (Chl). (B) A leaf from the *drp5b* mutant accumulating the stroma marker RBCS-mRFP and the isolation membrane marker GFP-ATG8a were incubated in sugar-free solution in darkness for 6 h from dawn and then observed. Second rosette leaves from 21- to 22-d-old plants were used. Arrowheads indicate the position of the chloroplast-associated isolation membrane. White and blue arrowheads indicate different isolation membrane–associated sites in a chloroplast. Images in (A) and (B) are still frames from Videos 19 and 20, respectively. Time scales above the images indicate the elapsed time from the start of the respective videos. Green, GFP-ATG8a; magenta, RBCS-mRFP; orange, chlorophyll (Chl) fluorescence. Scale bars, 5 μm.

Video 19. Release of a chloroplast bud in a sugar-starved leaf of the *drp5b* mutant.
The second rosette leaf from a 21-d-old *drp5b* plant accumulating the chloroplast stroma marker RBCS-mRFP was incubated in sugar-free solution in darkness for time-lapse imaging. Arrowhead indicates a chloroplast bud. Images acquired every 1 s are displayed at 20 frames/s. Scale bar, 5 μm. Green, RBCS-mRFP; magenta, chlorophyll fluorescence. Only the video of the merged channels is shown. This video was used to generate Figure 9A.

Video 20. An enlarged chloroplast caused by the *drp5b* mutation forms RCBs along the isolation membrane–associated sites.
The second rosette leaf from a 22-d-old *drp5b* plant accumulating the chloroplast stroma marker RBCS-mRFP and the isolation membrane marker GFP-ATG8a was incubated in sugar-free solution in darkness for time-lapse imaging. White or blue arrowheads indicate different isolation membrane–associated sites in a chloroplast. Images acquired every 2 s are displayed at 20 frames/s. Scale bar, 5 μm. Green, GFP-ATG8a; magenta, RBCS-mRFP. Only the video of the merged channels is shown. This video was used to generate Figure 9B.

References
Abeliovich H, Zarei M, Rigbolt KTG, Youle RJ, Dengjel J (2013) Involvement of mitochondrial dynamics in the segregation of mitochondrial matrix proteins during stationary phase mitophagy. Nat Commun **4:** 2789
Barros JAS, Cavalcanti JHF, Pimentel KG, Magen S, Soroka Y, Weiss S, Medeiros DB,

Nunes-Nesi A, Fernie AR, Avin-Wittenberg T, Araújo WL (2023) The interplay
between autophagy and chloroplast vesiculation pathways under dark-induced
senescence. Plant Cell Environ **46:** 3721-3736.
Brunkard JO, Runkel AM, Zambryski PC (2015) Chloroplasts extend stromules
independently and in response to internal redox signals. Proc Natl Acad Sci U S A **112:**
10044-10049.
Caplan JL, Kumar AS, Park E, Padmanabhan MS, Hoban K, Modla S, Czymmek K,
Dinesh-Kumar SP (2015) Chloroplast stromules function during innate immunity. Dev
Cell **34:** 45-57
Chakraborty S, Hill AL, Shirsekar G, Afzal AJ, Wang GL, Mackey D, Bonello P (2016)
Quantification of hydrogen peroxide in plant tissues using Amplex Red. Methods **109:**
105-113
Chen C, MacCready JS, Ducat DC, Osteryoung KW (2018) The molecular machinery of
chloroplast division. Plant Physiol **176:** 138-151
Chen YF, Culetto E, Legouis R (2022) The strange case of Drp1 in autophagy: Jekyll and
Hyde? Bioessays **44:** e2100271
Chiba A, Ishida H, Nishizawa NK, Makino A, Mae T (2003) Exclusion of ribulose-1,5-
bisphosphate carboxylase/oxygenase from chloroplasts by specific bodies in naturally
senescing leaves of wheat. Plant Cell Physiol **44:** 914-921
Clough SJ, Bent AF (1998) Floral dip: a simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. Plant J **16:** 735-743
Delaney TP, Uknes S, Vernooij B, Friedrich L, Weymann K, Negrotto D, Gaffney T,
Gutrella M, Kessmann H, Ward E, Ryals J (1994) A central role of salicylic-acid in
plant-disease resistance. Science **266:** 1247-1250
Doelling JH, Walker JM, Friedman EM, Thompson AR, Vierstra RD (2002) The APG8/12-
activating enzyme APG7 is required for proper nutrient recycling and senescence in
Arabidopsis thaliana. J Biol Chem **277:** 33105-33114
Farre JC, Subramani S (2016) Mechanistic insights into selective autophagy pathways:
lessons from yeast. Nat Rev Mol Cell Bio **17:** 537-552
Fukuda T, Ebi Y, Saigusa T, Furukawa K, Yamashita SI, Inoue K, Kobayashi D, Yoshida Y,
Kanki T (2020) Atg43 tethers isolation membranes to mitochondria to promote
starvation-induced mitophagy in fission yeast. Elife **9:** e61245
Fukuda T, Furukawa K, Maruyama T, Yamashita SI, Noshiro D, Song C, Ogasawara Y,
Okuyama K, Alam JM, Hayatsu M, Saigusa T, Inoue K, Ikeda K, Takai A, Chen L,
Lahiri V, Okada Y, Shibata S, Murata K, Klionsky DJ, Noda NN, Kanki T (2023)
The mitochondrial intermembrane space protein mitofissin drives mitochondrial fission
required for mitophagy. Mol Cell **83:** 2045-2058 e2049
Giacomello M, Pyakurel A, Glytsou C, Scorrano L (2020) The cell biology of mitochondrial
membrane dynamics. Nat Rev Mol Cell Bio **21:** 204-224
Hanson MR, Sattarzadeh A (2008) Dynamic morphology of plastids and stromules in
angiosperm plants. Plant Cell Environ **31:** 646-657
Hanson MR, Sattarzadeh A (2011) Stromules: recent insights into a long neglected feature of
plastid morphology and function. Plant Physiol **155:** 1486-1492.
He YL, Gao JY, Luo MQ, Gao CJ, Lin YS, Wong HY, Cui Y, Zhuang XH, Jiang LW (2022)
VAMP724 and VAMP726 are involved in autophagosome formation in Arabidopsis
thaliana. Autophagy **19:**1406-1423
Hirota T, Izumi M, Wada S, Makino A, Ishida H (2018) Vacuolar protein degradation via
autophagy provides substrates to amino acid catabolic pathways as an adaptive response
to sugar starvation in *Arabidopsis thaliana*. Plant Cell Physiol **59:** 1363-1376
Ishida H, Izumi M, Wada S, Makino A (2014) Roles of autophagy in chloroplast recycling.
Biochim Biophys Acta **1837:** 512-521

Ishida H, Okashita Y, Ishida H, Hayashi M, Izumi M, Makino A, Bhuiyan NH, van Wijk KJ (2021) GFS9 Affects Piecemeal Autophagy of Plastids in Young Seedlings of Arabidopsis thaliana. Plant Cell Physiol **62**: 1372-1386

Ishida H, Yoshimoto K, Izumi M, Reisen D, Yano Y, Makino A, Ohsumi Y, Hanson MR, Mae T (2008) Mobilization of rubisco and stroma-localized fluorescent proteins of chloroplasts to the vacuole by an ATG gene-dependent autophagic process. Plant Physiol **148**: 142-155

Ishida H, Yoshimoto K (2008) Chloroplasts are partially mobilized to the vacuole by autophagy. Autophagy **4**: 961-962

Itakura E, Kishi-Itakura C, Mizushima N (2012) The hairpin-type tail-anchored SNARE Syntaxin 17 targets to autophagosomes for fusion with endosomes/lysosomes. Cell **151**: 1256-1269

Ito E, Uemura T (2022) RAB GTPases and SNAREs at the trans-Golgi network in plants. J Plant Res **135**: 389–403

Izumi M, Hidema J, Wada S, Kondo E, Kurusu T, Kuchitsu K, Makino A, Ishida H (2015) Establishment of monitoring methods for autophagy in rice reveals autophagic recycling of chloroplasts and root plastids during energy limitation. Plant Physiol **167**: 1307-1320

Izumi M, Ishida H, Nakamura S, Hidema J (2017) Entire photodamaged chloroplasts are transported to the central vacuole by autophagy. Plant Cell **29**: 377-394

Izumi M, Wada S, Makino A, Ishida H (2010) The autophagic degradation of chloroplasts via rubisco-containing bodies is specifically linked to leaf carbon status but not nitrogen status in Arabidopsis. Plant Physiol **154**: 1196-1209

Kohler RH, Cao J, Zipfel WR, Webb WW, Hanson MR (1997) Exchange of protein molecules through connections between higher plant plastids. Science **276**: 2039-2042

Kusano S, Nakamura S, Izumi M, Hagihara S (2023) A fluorescent molecular probe for live-cell imaging of starch granules. ChemRxib

Law SR, Chrobok D, Juvany M, Delhomme N, Linden P, Brouwer B, Ahad A, Moritz T, Jansson S, Gardestrom P, Keech O (2018) Darkened leaves use different metabolic strategies for senescence and survival. Plant Physiol **177**: 132-150

Lazarou M, Sliter DA, Kane LA, Sarraf SA, Wang CX, Burman JL, Sideris DP, Fogel AI, Youle RJ (2015) The ubiquitin kinase PINK1 recruits autophagy receptors to induce mitophagy. Nature **524**: 309-314.

Lee HN, Chacko JV, Solis AG, Chen KE, Barros JA, Signorelli S, Millar AH, Vierstra RD, Eliceiri KW, Otegui MS, Benitez-Alfonso Y (2023) The autophagy receptor NBR1 directs the clearance of photodamaged chloroplasts. Elife **12**: e86030.

Mae T, Kai N, Makino A, Ohira K (1984) Relation between ribulose bisphosphate carboxylase content and chloroplast number in naturally senescing primary leaves of wheat. Plant Cell Physiol **25**: 333-336

Makino A, Osmond B (1991) Effects of nitrogen nutrition on nitrogen partitioning between chloroplasts and mitochondria in pea and wheat. Plant Physiol **96**: 355-362

Martinoia E, Heck U, Dalling MJ, Matile P (1983) Changes in chloroplast number and chloroplast constituents in senescing barley leaves. Biochem Physiol Pfl **178**: 147-155

Matsui T, Jiang PD, Nakano S, Sakamaki Y, Yamamoto H, Mizushima N (2018) Autophagosomal YKT6 is required for fusion with lysosomes independently of syntaxin 17. J Cell Biol **217**: 2633-2645

Miyagishima SY, Froehlich JE, Osteryoung KW (2006) PDV1 and PDV2 mediate recruitment of the dynamin-related protein ARC5 to the plastid division site. Plant Cell **18**: 2517-2530

Mochida K, Oikawa Y, Kimura Y, Kirisako H, Hirano H, Ohsumi Y, Nakatogawa H (2015) Receptor-mediated selective autophagy degrades the endoplasmic reticulum and the nucleus. Nature **522**: 359-362

Mochida K, Otani T, Katsumata Y, Kirisako H, Kakuta C, Kotani T, Nakatogawa H (2022) Atg39 links and deforms the outer and inner nuclear membranes in selective autophagy of the nucleus. J Cell Biol **221:** e202103178

Nakagawa T, Suzuki T, Murata S, Nakamura S, Hino T, Maeo K, Tabata R, Kawai T, Tanaka K, Niwa Y, Watanabe Y, Nakamura K, Kimura T, Ishiguro S (2007) Improved Gateway binary vectors: high-performance vectors for creation of fusion constructs in transgenic analysis of plants. Bioscience, biotechnology, and biochemistry **71:** 2095-2100

Nakamura S, Hagihara S, Izumi M (2021a) Mitophagy in plants. BBA-Gen Subjects **1865:** 129916

Nakamura S, Hagihara S, Otomo K, Ishida H, Hidema J, Nemoto T, Izumi M (2021b) Autophagy contributes to the quality control of leaf mitochondria. Plant Cell Physiol **62:** 229–247

Nakamura S, Hidema J, Sakamoto W, Ishida H, Izumi M (2018) Selective elimination of membrane-damaged chloroplasts via microautophagy. Plant Physiol **177:** 1007-1026

Nakamura S, Izumi M (2018) Regulation of chlorophagy during photoinhibition and senescence: lessons from mitophagy. Plant Cell Physiol **59:** 1135-1143

Nakatogawa H (2020) Mechanisms governing autophagosome biogenesis. Nat Rev Mol Cell Bio **21:** 439-458

Oikawa K, Goto-Yamada S, Hayashi Y, Takahashi D, Kimori Y, Shibata M, Yoshimoto K, Takemiya A, Kondo M, Hikino K, Kato A, Shimoda K, Ueda H, Uemura M, Numata K, Ohsumi Y, Hara-Nishimura I, Mano S, Yamada K, Nishimura M (2022) Pexophagy suppresses ROS-induced damage in leaf cells under high-intensity light. Nat Commun **13:** 7493

Onishi M, Yamano K, Sato M, Matsuda N, Okamoto K (2021) Molecular mechanisms and physiological functions of mitophagy. EMBO J **40:** e104705.

Ono Y, Wada S, Izumi M, Makino A, Ishida H (2013) Evidence for contribution of autophagy to rubisco degradation during leaf senescence in *Arabidopsis thaliana*. Plant Cell Environ **36:** 1147-1159

Osteryoung KW, Pyke KA (2014) Division and dynamic morphology of plastids. Annu Rev Plant Biol **65:** 443-472

Otomo K, Hibi T, Murata T, Watanabe H, Kawakami R, Nakayama H, Hasebe M, Nemoto T (2015) Multi-point scanning two-photon excitation microscopy by utilizing a high-peak-power 1042-nm laser. Anal Sci **31:** 307-313

Pan T, Liu YX, Hu XF, Li PW, Lin CC, Tang YY, Tang W, Liu YS, Guo L, Kim C, Fang J, Lin HH, Wu ZH, Blumwald E, Wang SH (2023) Stress-induced endocytosis from chloroplast inner envelope membrane is mediated by CHLOROPLAST VESICULATION but inhibited by GAPC. Cell Rep **42:** 113208.

Phillips AR, Suttangkakul A, Vierstra RD (2008) The ATG12-conjugating enzyme ATG10 is essential for autophagic vesicle formation in *Arabidopsis thaliana*. Genetics **178:** 1339-1353

Pyke KA, Leech RM (1994) A Genetic-Analysis of Chloroplast Division and Expansion in *Arabidopsis thaliana*. Plant Physiol **104:** 201-207

Rambold AS, Kostelecky B, Elia N, Lippincott-Schwartz J (2011) Tubular network formation protects mitochondria from autophagosomal degradation during nutrient starvation. Proc Natl Acad Sci U S A **108:** 10190-10195

Sasaki T, Tsutsumi M, Otomo K, Murata T, Yagi N, Nakamura M, Nemoto T, Hasebe M, Oda Y (2019) A novel katanin-tethering machinery accelerates cytokinesis. Curr Biol **29:** 4060-+

Segami S, Makino S, Miyake A, Asaoka M, Maeshima M (2014) Dynamics of vacuoles and H+-pyrophosphatase visualized by monomeric green fluorescent protein in Arabidopsis:

artifactual bulbs and native intravacuolar spherical structures. Plant Cell **26**: 3416-3434

Smith AM, Stitt M (2007) Coordination of carbon supply and plant growth. Plant Cell Environ
30: 1126-1149.

Spitzer C, Li FQ, Buono R, Roschzttardtz H, Chung TJ, Zhang M, Osteryoung KW,
Vierstra RD, Otegui MS (2015) The endosomal protein CHARGED
MULTIVESICULAR BODY PROTEIN1 regulates the autophagic turnover of plastids
in Arabidopsis. Plant Cell **27**: 391-402

Svenning S, Lamark T, Krause K, Johansen T (2011) Plant NBR1 is a selective autophagy
substrate and a functional hybrid of the mammalian autophagic adapters NBR1 and
p62/SQSTM1. Autophagy **7**: 993-1010.

Thompson AR, Doelling JH, Suttangkakul A, Vierstra RD (2005) Autophagic nutrient
recycling in Arabidopsis directed by the ATG8 and ATG12 conjugation pathways. Plant
Physiol **138**: 2097-2110

Tsuboyama K, Koyama-Honda I, Sakamaki Y, Koike M, Morishita H, Mizushima N
(2016) The ATG conjugation systems are important for degradation of the inner
autophagosomal membrane. Science **354**: 1036-1041

Twig G, Elorza A, Molina AJA, Mohamed H, Wikstrom JD, Walzer G, Stiles L, Haigh SE,
Katz S, Las G, Alroy J, Wu M, Py BF, Yuan J, Deeney JT, Corkey BE, Shirihai OS
(2008) Fission and selective fusion govern mitochondrial segregation and elimination
by autophagy. EMBO J **27**: 433-446

Vitha S, Froehlich JE, Koksharova O, Pyke KA, van Erp H, Osteryoung KW (2003) ARC6
is a J-domain plastid division protein and an evolutionary descendant of the
cyanobacterial cell division protein Ftn2. Plant Cell **15**: 1918-1933

Wada S, Ishida H, Izumi M, Yoshimoto K, Ohsumi Y, Mae T, Makino A (2009) Autophagy
plays a role in chloroplast degradation during senescence in individually darkened
leaves. Plant Physiol **149**: 885-893

Wan C, Zhang H, Cheng HY, Sowden RG, Cai WJ, Jarvis RP, Ling QH (2023) Selective
autophagy regulates chloroplast protein import and promotes plant stress tolerance.
EMBO J **42**: e112534.

Wang SH, Blumwald E (2014) Stress-induced chloroplast degradation in Arabidopsis is
regulated via a process independent of autophagy and senescence-associated vacuoles.
Plant Cell **26**: 4875-4888.

Wang WH, Li JY, Sun QQ, Yu XY, Zhang WW, Jia N, An CJ, Li YQ, Dong YN, Han FJ,
Chang N, Liu XM, Zhu ZL, Yu Y, Fan SL, Yang MJ, Luo SZ, Gao HB, Feng Y
(2017) Structural insights into the coordination of plastid division by the ARC6-PDV2
complex. Nat Plants **3**: 17011

Weaver LM, Amasino RM (2001) Senescence is induced in individually darkened Arabidopsis
leaves but inhibited in whole darkened plants. Plant Physiol **127**: 876-886

Yamashita S, Jin XL, Furukawa K, Hamasaki M, Nezu A, Otera H, Saigusa T, Yoshimori
T, Sakai Y, Mihara K, Kanki T (2016) Mitochondrial division occurs concurrently
with autophagosome formation but independently of Drp1 during mitophagy. J Cell
Biol **215**: 649-665

Yoshimoto K, Hanaoka H, Sato S, Kato T, Tabata S, Noda T, Ohsumi Y (2004) Processing
of ATG8s, ubiquitin-like proteins, and their deconjugation by ATG4s are essential for
plant autophagy. Plant Cell **16**: 2967-2983

Yoshimoto K, Jikumaru Y, Kamiya Y, Kusano M, Consonni C, Panstruga R, Ohsumi Y,
Shirasu K (2009) Autophagy negatively regulates cell death by controlling NPR1-
dependent salicylic acid signaling during senescence and the innate immune response in
Arabidopsis. Plant Cell **21**: 2914-2927

Yoshimoto K, Ohsumi Y (2018) Unveiling the molecular mechanisms of plant autophagy-from
autophagosomes to vacuoles in plants. Plant Cell Physiol **59**: 1337-1344

1421