1 **Hidden evolutionary constraints dictate the retention of coronavirus accessory genes**
2

3 Stephen A. Goldstein[1], Teagan M. Feeley[1], Kristina M. Babler[1], Zoë A. Hilbert[1,2], Diane M.
4 Downhour[1,2], Niema Moshiri[3], Nels C. Elde[1,2]
5
6 [1]Department of Human Genetics, University of Utah School of Medicine, Salt Lake City, UT, USA
7
8 [2]Howard Hughes Medical Institute, 4000 Jones Bridge Road, Chevy Chase, MD 20815, USA
9
10 [3]Department of Computer Science and Engineering, University of California, San Diego, La Jolla,
11 CA 92093,
12

Summary

Coronaviruses exhibit many mechanisms of genetic innovation[1–5], including the acquisition of

accessory genes that originate by capture of cellular genes or through duplication of existing viral

genes[6,7]. Accessory genes influence viral host range and cellular tropism, but little is known about

how selection acts on these variable regions of virus genomes. We used experimental evolution

of mouse hepatitis virus (MHV) encoding a cellular AKAP7 phosphodiesterase and an inactive

native phosphodiesterase, NS2 (ref 8) to simulate the capture of a host gene and analyze its

evolution. After courses of serial infection, the gene encoding inactive NS2, ORF2, unexpectedly

remained intact, suggesting it is under cryptic constraint uncoupled from the function of NS2. In

contrast, AKAP7 was retained under strong selection but rapidly lost under relaxed selection.

Guided by the retention of ORF2 and similar patterns in related betacoronaviruses, we analyzed

ORF8 of SARS-CoV-2, which arose via gene duplication[6] and contains premature stop codons in

several globally successful lineages. As with MHV ORF2, the coding-defective SARS-CoV-2

ORF8 gene remains largely intact, mirroring patterns observed during MHV experimental

evolution, challenging assumptions on the dynamics of gene loss in virus genomes and extending

these findings to viruses currently adapting to humans.

Introduction

Coronaviruses have unusually large RNA genomes, ranging from ~27-32 kilobases[9,10]. The stability of these large genomes is facilitated by the incorporation of a proofreading exonuclease into the RNA polymerase complex, unique among RNA viruses, that enhances replication fidelity[11–13]. Consequently, coronaviruses can accommodate addition of genetic material in the form of accessory genes acquired via horizontal gene transfer[14–16] and gene duplication[6,17]. Evolutionary theory on the streamlining of RNA virus genomes[1,3,4] predicts that new genes, such as coronavirus accessory genes, should encode near-immediate fitness benefits to avoid deletion.

Experimental studies of tobacco etch virus (TEV) support the assumption that exogenous genetic material is rapidly purged under relaxed selection[2,3]. However, exogenous genes engineered into TEV, a small single-stranded positive sense RNA virus, substantially increase genome size, which might drive loss regardless of a potential fitness benefit offered by the gene product. To date, no comparable studies using animal RNA viruses to test the interplay between gain of beneficial coding sequence and selection on genome size have been reported. Therefore, our understanding of how viral genome architecture and content evolve is poor, relative to the intense focus on how viruses evolve by nucleotide substitution. Given the major evolutionary leaps afforded by gene rearrangements, gains, and losses this represents a significant gap in our fundamental understanding of viral evolution.

To explore how selection acts on newly acquired viral genes we used experimental evolution of mouse hepatitis virus (MHV), a prototypical betacoronavirus, the genus that includes SARS and MERS-related coronaviruses. ORF2 of MHV and related viruses encodes a 2'-5' phosphodiesterase (PDE), NS2, that antagonizes the OAS-RNase L antiviral pathway. We used recombinant MHV encoding a cellular PDE, the AKAP7 central domain (AKAP7), as a functional

replacement for an inactive NS2 (ref 8) with a substitution at position 126 (His126Arg), that results

in restricted viral replication[18]. Viral PDEs were derived via capture of cellular AKAP7-like

genes[19,20] so using this virus, MHV[AKAP7], simulates a horizontal transfer event giving rise to a novel

viral gene. Critically, the AKAP7 PDE is inserted in place of ORF4 (ref 8), allowing us to probe

the relationship between selection and gene retention without substantial expansion of the

genome.

We found that the retention of coronavirus accessory gene sequences in different selective

environments does not conform to expectations of genome streamlining. Under relaxed selection

in the absence of active OAS-RNase L immune pressure, AKAP7 was rapidly lost, consistent with

the impact of selective constraints on genome size. However, in striking contrast, MHV ORF2 was

retained under all conditions tested, despite encoding an inactive PDE. In parallel, we found that

ORF8 of SARS-CoV-2 is largely intact even in lineages where it has lost coding capacity[21]. These

results suggest fundamental and widespread constraints on streamlining of viral genomes due to

genetic information with hidden functions beyond protein coding capacity.

Results

Cell type-dependent selective pressure on viral phosphodiesterases

To test the impacts of distinct selective pressures on coronavirus genome composition, we used

a recombinant mouse hepatitis virus (MHV) that has an inactivating H126R amino acid change in

NS2, its native PDE, which is functionally complemented by insertion of the coding sequence for

the cellular AKAP7 central domain PDE (AKAP7)[8] (**Figure 1A-B**). The AKAP7 gene inserted into

MHV is 618 nucleotides and replaces the defective 321 nucleotide ORF4, so expansion of the

genome is minimal (~1%). MHV with inactive NS2 (MHV[NS2Mut]) is restricted in mouse primary bone

marrow-derived macrophages (BMDMs) by OAS-RNase L (ref 18), while AKAP7 inserted into

MHV (MHV[AKAP7]) compensates for the inactive NS2 and restores replication to wild-type virus

levels in BMDMs and mouse liver. MHV[NS2mut] is restricted in immortalized primary macrophages[22],

but replicates to wild-type levels in L2 mouse fibroblasts (**Figure 1C**), providing differential

selection for courses of experimental evolution[18]. Similarly, MHV[AKAP7] replicated to high titers

whereas MHV encoding an inactive AKAP7 PDE (MHV[AKAP7mut]) was significantly restricted

(**Figure 1D**), demonstrating that these cells have an active OAS-RNase L pathway, which we

hypothesized imposes selective pressure to retain an active PDE. In contrast, MHV[AKAP7] and

MHV[AKAP7mut] replicated to an equivalent level in L2 fibroblasts, establishing these cells as a model

for MHV[AKAP7] evolution under relaxed selection (**Figure 1E**).

Deletion of AKAP7 occurs rapidly under relaxed selection

Our experimental evolution workflow uses serial passage in macrophages and L2 cells to model

evolution of a horizontally acquired PDE under different conditions (**Figure 1E**). We plaque

purified passage 0 (p0) MHV[AKAP7] and passaged this virus in triplicate in each cell type at a

multiplicity of infection (MOI) of 0.01. At each passage we titrated virus replication by plaque assay

to calculate the MOI for the subsequent passage and collected total RNA for analysis of AKAP7

by PCR and Sanger sequencing. In macrophages, full-length AKAP7 was retained through ten

passages (**Figure 1F, Figure S1A-B**), whereas in L2 cells, smaller PCR amplicons appeared by

passage three or four (**Figure 1G, Figure S1C-D**). The smaller bands became increasingly

prominent through p10, while the full length AKAP7 band faded in intensity, indicating viruses with

the full-length gene were disappearing from the population (**Figure 1G, Figure S1C-D**). We

cloned and sequenced full-length bands from macrophages and L2 cells, and smaller bands from

L2 cells at each passage.

Figure 1. Experimental evolution of mouse hepatitis virus reveals hidden selective constraints. A) Schematic of the MHV genome with NS2, the protein encoded by ORF8 indicated. B) Schematic of the MHV^{AKAP7} genome, with AKAP7 inserted in place of ORF4. C) Replication of wild-type MHV and MHV^{NS2mut} in macrophages (p=0.002) and L2 fibroblasts (p=0.968) 24 hpi, infected at an MOI of 0.01. One representative experiment of three is shown. D) Replication of MHV^{AKP7} and MHV^{AKAP7mut} in macrophages (p=0.002) and L2 cells (p=0.681) 24 hpi, infected at an MOI of 0.01. One representative experiment of three is shown. E) Schematic of the experimental evolution protocol. F-G) PCR analysis of AKAP7 at passage 1 to 10 in macrophages and L2 fibroblasts, respectively. The black arrow indicates full-length intact AKAP7, while red arrows indicate AKAP7 amplicons with deletions. H) Replication in macrophages 24 hpi (MOI=0.01) of p0 MHVAKAP7 and p10 virus populations from L2 fibroblasts and macrophages. Population 1 p-value=0.0257, Population 2 p-value=0.0058, Population 3 p-value=0.0054. Statistical significance determined by unpaired t-test. I) Replication of plaque purified p10 isolates from L2 fibroblasts (gray bars) and macrophages (black bars), in macrophages 24 hpi (MOI=0.01). J-K) PCR analysis of ORF2 in macrophages and L2 fibroblasts, respectively.

AKAP7 from both cell types contained no mutations (**Supplementary File 1**). The smaller bands

replacing full-length AKAP7 during L2 passage featured substantial deletions (**Supplementary**

File 1). The AKAP7 deletion variants did not undergo mutation, consistent with high fidelity

coronavirus replication and revealed a lower barrier to genetic change by deletion than

substitution, which has been described in SARS-CoV-2 as well[23]. Virus populations recovered at

p10 from L2s exhibited significantly restricted replication relative to macrophage-passaged virus

populations (**Figure 1H, Figure S2A-B**), demonstrating that despite the heterogeneity in AKAP7

deletion patterns between replicates, all three MHV^{AKAP7} populations lost the ability to effectively

suppress OAS-RNase L.

We then plaque purified virus isolates from replicate three p10 virus populations from both cell

types and used these isolates to infect macrophages and L2 cells. Consistent with the loss of

AKAP7 during L2 passage, p10 plaque purified isolates from these cells exhibited reduced

replication in macrophages but replicated to wild-type titers in L2 fibroblasts (**Figure 1I, Figure**

S2C-E), with some experiments showing a growth advantage for AKAP7-deleted isolates, which

warrants further study (**Figure S2C-E**). The rapid loss of AKAP7 under relaxed selection is

consistent with an evolutionary model wherein newly encoded viral proteins viral genes must

provide a near-immediate advantage for their parent gene to escape deletion.

Inactive MHV ORF2 is retained under strong and relaxed selection

Like the loss of AKAP7 under relaxed selection, there is an *a priori* prediction that ORF2, which

encodes an inactive PDE in MHV^{AKAP7}, would be also rapidly lost from the MHV genome. ORF2,

encoding the H126R inactivating substitution in NS2 should be dispensable in both macrophages

and L2 fibroblasts. Surprisingly, PCR of ORF2 (**Figure 1J-K**) at each serial passage in

macrophages and L2s, and sequencing of ORF2 after 10 passages, revealed no deletions novel

mutations, or reversion of the H126R substitution (**Supplementary File 2**). Given the unexpected

retention of inactive ORF2, we conducted five additional passages in both cell types to provide

additional opportunity for deletion. We sequenced the macrophage-passaged viruses again at

p15 to confirm there was no reversion at amino acid position 126 that might suggest ORF2

retention in these cells, even out to passage 15, was due to OAS-RNase L-mediated selection

(**Figure 1I-J; Supplementary File 2**). This unpredicted retention of ORF2 raised the possibility

that protein-coding coronavirus genes might be regularly retained under constraint distinct from

protein function.

To investigate whether similar patterns of constraint exist in the wild, we analyzed ORF2 from

other betacoronaviruses in the same subgenus as MHV. Rabbit coronavirus HKU14 (ref 24) has

a premature stop codon mutation truncating NS2 to 43 amino acids and additional downstream

stop codons in the same frame. However, the entire nucleotide sequence encoding full-length

NS2 is present in 4/5 HKU14 sequences in NCBI, while the other has ~100 bp of missing

sequence (**Supplementary FIle 3**). In contrast, ORF2 in porcine hemagglutinating

encephalomyelitis virus (PHEV) appears much more evolutionarily malleable[25–28], with various

isolates containing ORF2 with premature stop codons or deletions (**Supplementary File 3**).

These findings are broadly consistent with constraints on deletion of ORF2, with the heterogeneity

in PHEV suggesting there are dispensable genomic regions.

Long read sequencing confirms retention of ORF2 after experimental evolution

To more extensively analyze ORF2 and AKAP7 genes after experimental evolution we performed

Oxford Nanopore direct cDNA sequencing of plaque purified MHV[AKAP7] p0, five isolates after 10

passages in L2 cells and p10 isolate from macrophage infections. PCR analysis of the plaque

purified isolates showed AKAP7 was uniformly intact in macrophage-derived isolates but deleted

in L2-derived plaque isolates (**FigureS3A-B**). Mean read length across direct cDNA Nanopore

sequencing runs ranged from 1,065 to 1,528 bases and average coverage was 2,603x to 27,296x

per nucleotide of the MHV^{AKAP7} genome with a mean of 8,943x. Across all sequencing runs ~70-

80% of reads aligned to

the MHV^{AKAP7} genome

(**Supplementary File 5**).

To quantify changes in

ORF2 and AKAP7 gene

content after ten

passages and control for

variable overall

sequencing depth across

runs, we normalized

average coverage of

ORF2 and AKAP7 to

average coverage of

ORF1ab for each isolate.

Consistent with PCR

Figure 2. Nanopore direct cDNA sequencing shows retention of ORF2 and near-complete loss of AKAP7 following serial passage. A-C) Relative coverage depth plots of ORF2 in purified plaque isolates from passage 0 MHV^{AKAP7}, and passage 10 macrophage and L2 fibroblast purified plaques. D-E) Relative coverage depth plots of AKAP7 in purified plaque isolates from passage 0 MHV^{AKAP7}, and passage 10 macrophage and L2 fibroblast purified plaques.

analysis, average relative coverage of ORF2 was the same in plaque purified passage 10 isolates

from macrophages and L2 fibroblasts (**Figure 2A-C; Figure S3C-F**), demonstrating that this

genomic region is under unanticipated constraint. The average relative AKAP7 coverage for p0

and macrophage p10 were 11.2 and 27.8, respectively (**Figure 2D-E**). In contrast, the average

relative AKAP7 coverage for L2 p10 isolates was 0.332, a 33-fold decrease from p0 consistent

with PCR analysis. (**Figure 2F; S3G-J**).

SARS-CoV-2 ORF8 is retained despite loss of coding capacity

Our analysis of ORF2 in MHV-like viruses suggested the constraints we identified experimentally

also exist in virus populations in the wild. However, the strength of this analysis was limited by

the relative paucity of genomic data available for animal viruses. In contrast, the SARS-CoV-2

dataset in the Global Initiative on Sharing All Influenza Data (GISAID) database contains more

than 16 million genomes, offering an unprecedented window into viral evolution. Analogous to the

inactivation of MHV ORF2 in our system, SARS-CoV-2 ORF8 has acquired premature stop

codons in numerous globally successful lineages, including B.1.1.7, XBB.1, XBB.1.5, XBB.1.9,

and XBB.1.16 (ref 21), and a mutation in the ORF8 transcriptional regulatory sequence (TRS) in

the BA.5 lineage that substantially reduces ORF8 subgenomic RNA and protein synthesis[29].

ORF8 has been of interest since early in the SARS-CoV-2 pandemic due to a small deletion that

emerged in its SARS-CoV ortholog in 2003 [30], an early cluster of infections in Singapore involving

an ORF8 deletion variant[31,32], as well as debate over its role during SARS-CoV-2 infection[6,32–39].

Despite the intense focus on ORF8 function and loss of full-length protein synthesis, very little

attention has been devoted to the evolution of the gene itself.

We downloaded all sequences assigned to the B.1.1.7, BA.5, XBB.1, XBB.1.5, XBB.1.9, and

XBB.1.16 lineages from GISAID and applied a cutoff date to exclude any samples without

metadata or with metadata suggesting they are incorrectly assigned (**Figure S4**). For each

lineage, we further defined a start-date for circulation and calculated the date at which 50% and

90% of sequences assigned to this lineage were collected, as of November 29, 2023. We further

excluded any sequences with any ambiguous nucleotides in the ORF8 gene to eliminate

confounding of our analysis by poor-quality sequences, which might obscure *bona fide* deletions.

We calculated a deletion length distribution of ORF8 in B.1.1.7 (1,097,307 sequences), BA.5

(20,833 sequences), XBB.1 (22,031 sequences), XBB.1.5 (166,068 sequences), XBB.1.9.1

(19.990 sequences), and XBB.1.16 (19,119 sequences) (**Figure 3C**). Across all lineages 96% of

deletions were smaller than ten nucleotides, ranging from a high of 98.15% of deletions smaller

than ten nucleotides for B.1.1.7 to a low of 86.6% of deletions smaller than ten nucleotides for

XBB.1.16. (**Figure 3C, Supplementary File 5**). We then analyzed all sequences in the dataset

to determine whether any regions of ORF8 were particularly deletion-tolerant or resistant. While

XBB.1.5 ORF8 deletions were enriched in the 5' half of the gene (**Figure 3D, S5B)**, deletions

were uniformly rare at all

positions across the other

lineages we analyzed

(**Figure 3D-F, S5A-D)**

except for a single site at

the 3' end that we speculate

may impact sgRNA

synthesis of downstream

ORFs N and 9b.

To determine whether

ORF8 in these lineages

degraded over time, we

calculated the percent of

ORF8 gene content in

these lineages, sorted by

collection date, and plotted

these values for all

Figure 3. ORF8 is retained in SARS-CoV-2 lineages despite mutations that disrupt protein coding. A) Schematic of SARS-CoV-2 genome. B) Schematic (with ORF8 expanded, not to scale) of ORF8, with mutations resulting in loss of subgenomic mRNA synthesis (BA.5) or an early stop codon (B.1.1.7 and XBB.1.x) indicated. C) Deletion length distribution - this displays the size of all deletions in ORF8 in each lineage from 0 to >100 nucleotides. D-F) Proportion of genomes from lineages with a deletion at each position in SARS-CoV-2 ORF8.

genomes of all six lineages (**Figure 4A-F**). To determine how the appearance of ORF8 deletions

in these lineages related to their epidemiological rise and fall, we calculated the dates at which

50% (dashed vertical line) and 90% (solid vertical line) of sequences assigned to each lineage

had been collected. For all lineages, genomes with deletions were rare outside this window,

generally occurring early after the appearance of a lineage and going extinct during lineage

growth or emerging very late when the lineage was already in steep decline.

Figure 4. SARS-CoV-2 ORF8 is recurrently retained following acquisition of an early stop codon.
A-F) Line plots of percent nucleotide content for the indicated SARS-CoV-2 lineages over time throughout the course of their circulation in humans. Dark lines are the mean percentage for all sequences collected on that date, and faded lines are the 95% confidence interval. Dashed vertical line is the date by which 50% of sequence assigned to the designated lineage were collected, and the solid line is the 90% cutoff.

Discussion

Findings from this study challenge a simple evolutionary model wherein the fate of protein coding

genes hinges only on protein function. The sustained, unexpected retention of ORF2 in both

macrophages and L2 fibroblasts through fifteen passages suggests a complex interplay of

selective pressure acting on viral coding sequence. In L2s the PDE activity of ORF2 is

dispensable, and the stability of the H126R substitution during experimental evolution in

macrophages is consistent with AKAP7 fully counteracting OAS-RNase L activity. While it is

formally possible that an unknown secondary role for the ORF2-encoded protein NS2 antagonizes

innate immunity, no such function has been identified and the protein contains no identifiable

domains other than the PDE.

The difference in genomic location between ORF2 and AKAP7 may be relevant to their respective

fates. AKAP7 is located upstream of ORF5a, which encodes an accessory gene that promotes

resistance to interferon signaling[41], whereas ORF2 immediately precedes the hemagglutinin-

esterase (HE) and spike structural genes. HE subgenomic mRNA is not expressed from the MHV

A59 genome[42,43], so it is unlikely that the selection against ORF2 deletion relates to transcription

of the immediate downstream gene.

Under a conventional model of selection favoring genome streamlining, viruses with ORF8

deleted would quickly dominate populations soon after the potential to encode full-length protein

was lost. The exact nature of the evolutionary constraint on MHV ORF2 and SARS-CoV-2 ORF8

is not yet clear. A leading possibility is that at least some coronavirus genes contain regulatory

functions in addition to their protein coding capacity that tune the transcription of other genes.

MHV ORF2 is upstream of spike, albeit with the defective HE gene intervening, and SARS-CoV-

2 ORF8 immediately precedes nucleocapsid, which has multiple functions key to promoting viral

fitness[44]. If accessory genes promote transcription of spike, nucleocapsid, and other subgenomic

RNAs, loss of such genes might reduce viral fitness and lineages containing such deletions would

go extinct. Increased transcription of subgenomic RNAs encoding innate immune antagonists has

been linked to enhanced suppression of innate immunity in major SARS-CoV-2 variants of

concern[45], which is consistent with our hypothesis and analysis of SARS-CoV-2 ORF8 retention.

The loss of ORF2 in HKU1 and the observation of some ORF8 deletion SARS-CoV-2 viruses

indicates that patterns of constraint may be subtle or absent in some genomic regions. In the case

of SARS-CoV-2, the apparent decreased fitness of ORF8-deletion viruses might be balanced by

adaptations that compensate for loss of ORF8, affording fitness benefits of genome compression

while shedding the deleterious impact of losing the gene. Continued surveillance of SARS-CoV-

2 might reveal such variants over and provide more insight into fundamental factors driving

coronavirus evolution.

Identifying additional patterns of evolutionary constraint may reveal other sequence features

impacting viral fitness. Patterns of selection can be prioritized for mechanistic studies using

reverse genetic strategies established for MHV and SARS-CoV-2, opening a bridge between

experimentation and computational analysis of viral evolution. Experimentally dissecting fitness

impacts will benefit from future competition studies. For example, the loss of ORF2 in HKU1

results in viable viruses and fitness defects of $MHV^{\Delta ORF2}$ might appear only in direct competition

with wild-type viruses and not in parallel courses of experimental evolution. Overall, this work

demonstrates the power of merging experimental approaches with computational analysis of

massive datasets to identify patterns of constraint influencing the evolution of emerging and

pandemic viruses.

Materials and Methods

Cells and Viruses

Immortalized primary macrophages were provided by Sunny Shin[22], 17-Clone 1 and L2 cells were provided by Susan Weiss. Macrophages were cultured in RPMI-1640 supplemented with 10% FBS, 1% L-glutamine, and 1% Penicillin/Streptomycin, while 17-Clone 1 and L2 cells were cultured in DMEM supplemented the same way.

All viruses used in this study were provided by Susan Weiss at the University of Pennsylvania. Wild-type MHV and MHV^{NS2mut} were previously described[18] as were MHV^{AKAP7} and MHV^{AKAP7mut8}. Viruses were received as seed aliquots and 100 μl of each virus was added to one T75 flask containing a confluent monolayer of 17-Clone 1 cells in 2 ml volume. Flasks were incubated for one hour at 37°C, after which 10 ml of fresh media was added. Once widespread cytopathic effect was present (18-24 hours post-infection) the flasks were put through three freeze-thaw cycles and the supernatant was clarified via centrifugation and aliquoted for later use.

Virus infections and plaque assays

One day after seeding in 12-well plates macrophages or L2 cells were infected at an MOI of 0.01 in 200 μl total volume and incubated for one hour at 37 °C with rocking every 15 minutes. After one hour cells were washed 3 times with PBS and 1 ml of fresh media (2% FBS) was added to each well. 24 hours post-infection 300 μl of supernatant was collected and stored at -80°C. We also harvested total RNA for additional analyses (described below). Initial infections, all serial passages, and endpoint replication comparisons were done at an MOI of 0.01. Viral titers were quantified by plaque assay on confluent L2 cell monolayers. Supernatants collected 24 hours post-infection were serially diluted 1:10 to 10^{-8}, inoculated onto L2 monolayers and incubated for 1 hour at 37 °C with rocking every 15 minutes. After one hour 3 mL of semi-solid agarose overlay was added to each well and the plates returned to 37 °C for 24 hours. 24 hours post-infection the

agarose overlay was removed and the monolayers were washed with PBS and stained with 0.5%

crystal violet plus 20% methanol. The stain was washed off with water and plaques counted and

recorded for analysis.

Statistical testing

Statistical testing of viral titers (**Figure 1C-D, H-I; Figure S2**) was conducted in GraphPad Prism

10, via unpaired t-test.

RNA extraction and cDNA synthesis

Total RNA was harvested 24 hours post-infection on each serial passage with the Zymo Research

Quick-RNA Miniprep Kit (Cat. #R1057) following the manufacturer's protocol. To increase yield

we eluted the RNA in a reduced volume of 25 µl nuclease-free water. RNA quantification and

quality assessment was conducted using a Synergy HT BioTek plate reader. cDNA was

synthesized using the Thermo Scientific Maxima First Strand cDNA Synthesis Kit with dsDNase

(Lot. 2664799) with 1 µg of input RNA.

PCR and Sanger sequencing

At each passage the AKAP7 central domain and ORF2 genes from MHV^{AKAP7} were analyzed by

PCR. We used Phusion Flash HiFi Master Mix (Cat # F548S) in 20 µl/reaction with 2 µl of cDNA

template. Thirty cycles of PCR were performed. Cycling conditions were: **1)** Denaturation, 98^{o}C

for 10 seconds **2) a)** Denaturation (98^{o}C for 2 seconds) **b)** Annealing (63^{o}C for 10 seconds) **c)**

Extension (72^{o}C for 30 seconds) **d)** Final extension (72^{o}C for 2 minutes) **3)** Hold (4^{o}C).

PCR products were visualized by agarose gel (1%) electrophoresis then extracted and purified

using Zymoclean Gel DNA Recovery Kit (Zymo Research Lot. #213587) following manufacturer

protocols. DNA was eluted in 10 µl. Purified PCR products were prepared for Sanger Sequencing

by TOPO cloning using the Invitrogen TA Cloning™ Kit, with pCR™2.1 Vector (Cat. #K202020)

following manufacturer protocols. Plasmid DNA was extracted using the Zippy Plasmid Miniprep

Kit (Zymo Research Cat. # 11-308) following manufacturer protocol with elution 75 µl nuclease-

free water. Plasmid DNA was quantified with a Synergy HT BioTek plate reader and sent for

sequencing to the University of Utah DNA sequencing core or Genewiz.

Sequencing primers

Primers	Forward	Reverse
AKAP7	5'-**ATTGCTACCTGGCCCCG**-3'	5'-**CTAGGGTCTTAGGCCCAAATG**-3'
MHV ORF2	5'-**ATGGCCTTTGCTGACAAGCCTAAT**-3'	5'-**TCAACACATACAACCCTTCATTCT**-3'
M13	5'-**GTAAAACGACGGCCAG**-3'	5'-**CAGGAAACAGCTATGAC**-3'

Sequence analysis

353   Sequence files were imported into Geneious and aligned to either the AKAP7 or ORF2 coding

354   sequences using the MAFFT Geneious plug-in [46]. HKU1, HKU14, and PHEV multiple sequence

355   alignments were likewise generated using the MAFFT Geneious plug-in with default parameters.

Oxford Nanopore direct cDNA sequencing

1 mL of media containing a plaque isolate was used to infect 2x 10 cm dishes of desired cell type

(macrophages or L2). 18-20 hours post-infection total RNA was collected using Zymo Research

Quick-RNA Miniprep Kit (Cat. #R1057). The same lysis buffer (1 ml) was used on both dishes/per

infection to maximize yield. Polyadenylated RNA was extracted using the Invitrogen Poly(A)Purist

MAG Kit (Cat # AM1922) following the manufacturer protocol. 100 ng PolyA+ RNA was prepared

for sequencing using the Direct cDNA Sequencing Kit SQK-DCS109 from Oxford Nanopore

Technologies, according to the manufacturer's instructions. The prepared cDNA libraries were

sequenced using the MinION Mk1C with FLO-MIN106D flow cells, version 9.4.1. Fastq files were

generated from fast5 files using the high-accuracy model of Guppy basecaller, version 6.5.7 with

the following parameters hac_guppy -c dna_r9.4.1_450bps_hac.cfg -x auto --compress_fastq.

Direct cDNA sequencing coverage analysis

Fastq files from Nanopore sequencing runs were individually aligned to the MHV^{AKAP7} (passage

0) reference genome. Alignments were performed using minimap2 (v.2.23) and subsequently

sorted and indexed with samtools (v.1.16). Coverage at every position across the genome for

each sample was calculated using the bedtools (v.2.26.0) genomecov command. Coverage was

then normalized for each sample to the average coverage across ORF1ab (nucleotide positions

211-21746 in the reference genome), which contains the non-amplified reading frame, allowing

for comparison across sequencing runs and samples. A rolling average of this normalized read

depth was calculated across the genome in 25 bp windows and plotted with ggplot2 in R. For

AKAP7 and ORF2 analysis, rolling averages were recalculated for the region containing the gene

sequence ± 300 bp of flanking sequence on either side and plotted identically to the genome wide

coverage plots. Fastq sequencing files are available in the Sequence Read Archive at BioProject

ID PRJNA1031673.

SARS-CoV-2 ORF8 Deletion Analysis

All   methods   and   code   used   to   conduct   this   analysis   are   available   at

https://github.com/niemasd/SC2-Deletion-Analysis.

Acknowledgements

This work was supported by NIH grants F32AI152341 to SAG and R35GM134936 to NCE. We

thank Susan Weiss and Sunny Shin at the University of Pennsylvania for reagents. We thank

Tom Peacock at the Pirbright Institute and David Bauer from the Francis Crick Institute for helpful

discussions.

Author Contributions

SAG: project conceptualization, experimental design, experimentation, data analysis, manuscript

writing, manuscript editing

TMF: experimentation, data analysis

KMB: experimentation, data analysis, schematic design, manuscript editing

ZAH: AKAP7 and ORF2 coverage quantification and analysis

DMD: sequencing library preparation, direct cDNA sequencing and analysis

NM: SARS-CoV-2 ORF8 deletion analysis

NCE: project conceptualization, data analysis, manuscript writing, manuscript editing

References

1. Willemsen, A. & Zwart, M. P. On the stability of sequences inserted into viral genomes.

Virus Evol **5**, vez045 (2019).

2. Zwart, M. P., Willemsen, A., Daròs, J.-A. & Elena, S. F. Experimental evolution of

pseudogenization and gene loss in a plant RNA virus. *Mol. Biol. Evol.* **31**, 121–134 (2014).

3. Willemsen, A., Zwart, M. P., Ambrós, S., Carrasco, J. L. & Elena, S. F. 2b or Not 2b:

Experimental Evolution of Functional Exogenous Sequences in a Plant RNA Virus.

Genome Biol. Evol. **9**, 297–310 (2017).

4. Belshaw, R., Pybus, O. G. & Rambaut, A. The evolution of genome compression and

genomic novelty in RNA viruses. *Genome Res.* **17**, 1496–1504 (2007).

5.   Simon-Loriere, E., Holmes, E. C. & Pagán, I. The effect of gene overlapping on the rate of

RNA virus evolution. *Mol. Biol. Evol.* **30**, 1916–1928 (2013).

6.   Tan, Y., Schneider, T., Leong, M., Aravind, L. & Zhang, D. Novel Immunoglobulin Domain

Proteins Provide Insights into Evolution and Pathogenesis of SARS-CoV-2-Related

Viruses. *MBio* **11**, (2020).

7.   Gorbalenya, A. E., Enjuanes, L., Ziebuhr, J. & Snijder, E. J. Nidovirales: evolving the

largest RNA virus genome. *Virus Res.* **117**, 17–37 (2006).

8.   Gusho, E. *et al.* Murine AKAP7 has a 2',5'-phosphodiesterase domain that can complement

an inactive murine coronavirus ns2 gene. *MBio* **5**, e01312–e1314 (2014).

9.   Debat, H. J. Expanding the size limit of RNA viruses: Evidence of a novel divergent

nidovirus in California sea hare, with a ~35.9 kb virus genome. *bioRxiv* 307678 (2018)

doi:10.1101/307678.

10. Saberi, A., Gulyaeva, A. A., Brubacher, J. L., Newmark, P. A. & Gorbalenya, A. E. A

planarian nidovirus expands the limits of RNA genome size. *PLoS Pathog.* **14**, e1007314

(2018).

11. Smith, E. C. & Denison, M. R. Coronaviruses as DNA wannabes: a new model for the

regulation of RNA virus replication fidelity. *PLoS Pathog.* **9**, e1003760 (2013).

12. Eckerle, L. D., Lu, X., Sperry, S. M., Choi, L. & Denison, M. R. High Fidelity of Murine

Hepatitis Virus Replication Is Decreased in nsp14 Exoribonuclease Mutants. *J. Virol.* **81**,

12135–12144 (2007).

13. Smith, E. C. & Denison, M. R. Implications of altered replication fidelity on the evolution and

pathogenesis of coronaviruses. *Curr. Opin. Virol.* **2**, 519–524 (2012).

14. Goldstein, S. A. & Elde, N. C. Recurrent Viral Capture of a Host-Derived Immune

Antagonist. *bioRxiv* 2023.05.12.540623 (2023) doi:10.1101/2023.05.12.540623.

15. Mazumder, R., Iyer, L. M., Vasudevan, S. & Aravind, L. Detection of novel members,

structure-function analysis and evolutionary classification of the 2H phosphoesterase

superfamily. *Nucleic Acids Res.* **30**, 5229–5243 (2002).

16. Asthana, A., Gaughan, C., Dong, B., Weiss, S. R. & Silverman, R. H. Specificity and

Mechanism of Coronavirus, Rotavirus, and Mammalian Two-Histidine Phosphoesterases

That Antagonize Antiviral Innate Immunity. *MBio* **12**, e0178121 (2021).

17. Neches, R. Y., Kyrpides, N. C. & Ouzounis, C. A. Atypical Divergence of SARS-CoV-2 Orf8

from Orf7a within the Coronavirus Lineage Suggests Potential Stealthy Viral Strategies in

Immune Evasion. *MBio* **12**, (2021).

18. Zhao, L. *et al.* Antagonism of the interferon-induced OAS-RNase L pathway by murine

coronavirus ns2 protein is required for virus replication and liver pathology. *Cell Host*

Microbe **11**, 607–616 (2012).

19. Goldstein, S. A. & Elde, N. C. Recurrent viral capture of cellular phosphodiesterases that

antagonize OAS-RNase L. *Proc. Natl. Acad. Sci. U. S. A.* **121**, e2312691121 (2024).

20. Nomburg, J., Price, N. & Doudna, J. A. Birth of new protein folds and functions in the

virome. *bioRxiv* 2024.01.22.576744 (2024) doi:10.1101/2024.01.22.576744.

21. Wagner, C. *et al.* Positive selection underlies repeated knockout of ORF8 in SARS-CoV-2

evolution. *bioRxiv* (2023) doi:10.1101/2023.09.21.23295927.

22. Blasi, E., Radzioch, D., Merletti, L. & Varesio, L. Generation of macrophage cell line from

fresh bone marrow cells with a myc/raf recombinant retrovirus. *Cancer Biochem. Biophys.*

10, 303–317 (1989).

23. McCarthy, K. R. *et al.* Recurrent deletions in the SARS-CoV-2 spike glycoprotein drive

antibody escape. *Science* **371**, 1139–1142 (2021).

24. Lau, S. K. P. *et al.* Isolation and characterization of a novel Betacoronavirus subgroup A

coronavirus, rabbit coronavirus HKU14, from domestic rabbits. *J. Virol.* **86**, 5481–5496

(2012).

25. He, W.-T. *et al.* Newly identified lineages of porcine hemagglutinating encephalomyelitis

virus exhibit respiratory phenotype. *Virus Evol* **9**, vead051 (2023).

26. Vijgen, L. *et al.* Evolutionary history of the closely related group 2 coronaviruses: porcine

hemagglutinating encephalomyelitis virus, bovine coronavirus, and human coronavirus

OC43. *J. Virol.* **80**, 7270–7274 (2006).

27. Shi, J. *et al.* Genomic characterization and pathogenicity of a porcine hemagglutinating

encephalomyelitis virus strain isolated in China. *Virus Genes* **54**, 672–683 (2018).

28. Rho, S. *et al.* Detection and genetic analysis of porcine hemagglutinating encephalomyelitis

virus in South Korea. *Virus Genes* **42**, 90–96 (2011).

29. Bouhaddou, M. *et al.* SARS-CoV-2 variants evolve convergent strategies to remodel the

host response. *Cell* **186**, 4597–4614.e26 (2023).

30. Muth, D. *et al.* Attenuation of replication by a 29 nucleotide deletion in SARS-coronavirus

acquired during the early stages of human-to-human transmission. *Sci. Rep.* **8**, (2018).

31. Young, B. E. *et al.* Effects of a major deletion in the SARS-CoV-2 genome on the severity

of infection and the inflammatory response: an observational cohort study. *Lancet* **396**,

603–611 (2020).

32. Su, Y. C. F. *et al.* Discovery and Genomic Characterization of a 382-Nucleotide Deletion in

ORF7b and ORF8 during the Early Evolution of SARS-CoV-2. *MBio* **11**, e01610–20 (2020).

33. Flower, T. G. *et al.* Structure of SARS-CoV-2 ORF8, a rapidly evolving immune evasion

protein. *Proc. Natl. Acad. Sci. U. S. A.* **118**, (2021).

34. Gamage, A. M. *et al.* Infection of human Nasal Epithelial Cells with SARS-CoV-2 and a

382-nt deletion isolate lacking ORF8 reveals similar viral kinetics and host transcriptional

profiles. *PLoS Pathog.* **16**, e1009130–e1009130 (2020).

35. Bello-Perez, M. *et al.* SARS-CoV-2 ORF8 accessory protein is a virulence factor. *MBio*

e0045123 (2023).

36. Wang, X. *et al.* SARS-CoV-2 ORF8 Protein Induces Endoplasmic Reticulum Stress-like

Responses and Facilitates Virus Replication by Triggering Calnexin: an Unbiased Study. *J.*

Virol. **97**, e0001123 (2023).

37. Tamura, T. *et al.* Virological characteristics of the SARS-CoV-2 XBB.1.5 variant. *bioRxiv*

2023.08.16.553332 (2023) doi:10.1101/2023.08.16.553332.

38. Rodriguez-Rodriguez, B. A. *et al.* A neonatal mouse model characterizes transmissibility of

SARS-CoV-2 variants and reveals a role for ORF8. *Nat. Commun.* **14**, 3026 (2023).

39. Ponde, N. O. *et al.* SARS-CoV-2 ORF8 Mediates Signals in Macrophages and Monocytes

through MyD88 Independently of the IL-17 Receptor. *J. Immunol.* **211**, 252–260 (2023).

40. Fehr, A. R. & Perlman, S. Coronaviruses: An Overview of Their Replication and

Pathogenesis. in 1–23 (Springer New York, 2015).

41. Koetzner, C. A. *et al.* Accessory protein 5a is a major antagonist of the antiviral action of

interferon against murine coronavirus. *J. Virol.* **84**, 8262–8274 (2010).

42. Shieh, C. K. *et al.* Identification of a new transcriptional initiation site and the corresponding

functional gene 2b in the murine coronavirus RNA genome. *J. Virol.* **63**, 3729–3736 (1989).

43. Kazi, L., Lissenberg, A., Watson, R., de Groot, R. J. & Weiss, S. R. Expression of

hemagglutinin esterase protein from recombinant mouse hepatitis virus enhances

neurovirulence. *J. Virol.* **79**, 15064–15073 (2005).

44. Johnson, B. A. *et al.* Nucleocapsid mutations in SARS-CoV-2 augment replication and

pathogenesis. *PLoS Pathog.* **18**, e1010627 (2022).

45. Thorne, L. G. *et al.* Evolution of enhanced innate immune evasion by SARS-CoV-2. *Nature*

602, 487–495 (2022).

46. Katoh, K., Misawa, K., Kuma, K.-I. & Miyata, T. MAFFT: a novel method for rapid multiple

sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* **30**, 3059–3066

(2002).

513