Rho GTPase signaling and mDia facilitate endocytosis via presynaptic actin

Kristine Oevel[1], Svea Hohensee[1], Atul Kumar[2], Irving Rosas-Brugada[1#], Francesca Bartolini[2],

Tolga Soykan[1*], and Volker Haucke[1,3,4*]

[1] Leibniz-Forschungsinstitut für Molekulare Pharmakologie (FMP), 13125 Berlin, Germany.

[2] Department of Pathology and Cell Biology, Columbia University Medical Center, New York,

NY 10032, USA.

[3] Faculty of Biology, Chemistry, Pharmacy, Freie Universität Berlin, 14195 Berlin, Germany.

[4] NeuroCure Cluster of Excellence, Charité Universitätsmedizin Berlin, 10117 Berlin, Germany.

*Correspondence: haucke@fmp-berlin.de (V.H.) or soykan@fmp-berlin.de (T.S.).

deceased

ABSTRACT

Neurotransmission at synapses is mediated by the fusion and subsequent endocytosis of synaptic

vesicle membranes. Actin has been suggested to be required for presynaptic endocytosis but the

mechanisms that control actin polymerization and its mode of action within presynaptic nerve

terminals remain poorly understood. We combine optical recordings of presynaptic membrane

dynamics and ultrastructural analysis with genetic and pharmacological manipulations to

demonstrate that presynaptic endocytosis is controlled by actin regulatory diaphanous-related

formins mDia1/3 and Rho family GTPase signaling. We show that impaired presynaptic actin

assembly in the near absence of mDia1/3 and reduced RhoA activity is partly compensated by

hyperactivation of Rac1. Inhibition of Rac1 signaling further aggravates impaired presynaptic

endocytosis elicited by loss of mDia1/3. Our data suggest that interdependent mDia1/3-Rho and

Rac1 signaling pathways cooperatively act to facilitate synaptic vesicle endocytosis by controlling

presynaptic F-actin.

INTRODUCTION

Synaptic transmission relies on the release of neurotransmitters by calcium-triggered exocytic fusion of synaptic vesicles (SVs) at specialized active zone release sites within presynaptic nerve terminals. Following fusion, compensatory endocytosis retrieves SV proteins and lipids from the presynaptic plasma membrane and SVs are reformed (Chanaday *et al*, 2019; Gan & Watanabe, 2018; Kononenko & Haucke, 2015; Saheki & De Camilli, 2012; Soykan *et al*, 2016), e.g. by clathrin-mediated vesicle budding (Kononenko *et al*, 2014; Watanabe *et al*, 2014) (but see (Wu *et al*, 2014) for evidence in favor of clathrin-independent SV reformation). Endocytosis relies on mechanical forces to enable membrane deformation and eventually fission (Engqvist-Goldstein & Drubin, 2003). In many biological systems, actin and actin-associated myosin motors appear to provide such mechanical force to facilitate vesicle formation (Anes *et al*, 2003; Marston *et al*, 2003). For example, the function of actin for endocytosis is well-established in yeast, whereas differential requirements for actin in distinct forms of endocytosis have been described in various types of mammalian cells including neurons (Boulant *et al*, 2011; Merrifield *et al*, 2005; Saffarian *et al*, 2009; Saheki & De Camilli, 2012). Pharmacological inhibition of actin assembly by latrunculin has been shown to cause the accumulation of endocytic intermediates at lamprey giant synapses (Shupliakov *et al*, 2002) and interfere with ultrafast endocytosis in response to single optical action potentials (APs) at hippocampal synapses (Watanabe *et al*, 2013; Watanabe *et al*., 2014; Wu & Chan, 2022) and with fast endocytosis at cerebellar mossy fiber boutons (Delvendahl *et al*, 2016). The same treatment does not hamper endocytosis of SV proteins at hippocampal synapses stimulated with trains of APs (Sankaranarayanan *et al*, 2003; Soykan *et al*, 2017). Genetic interference with actin function by conditional knockout (KO) of β- or γ-actin genes has suggested a crucial role for actin in all kinetically distinguishable forms of endocytosis (Wu *et al*, 2016) at hippocampal synapses and the calyx of Held, a fast giant synapse in the auditory

brain stem (Borst & Soria van Hoeve, 2012). However, in the same preparations actin loss also

affected SV exocytosis (Wu *et al.*, 2016), suggesting a more general requirement of actin for

presynaptic function. Consistently, actin has been shown to surround clusters of reserve pool SVs

in lamprey (Bloom *et al*, 2003), to facilitate the replenishment of fast releasing vesicles in various

models (Sakaba *et al*, 2013; Sakaba & Neher, 2003), to be involved in Tau pathology (Zhou *et al*,

2017), to induce spine growth and plasticity, (Cingolani & Goda, 2008) and to steer neuronal

migration (Shinohara *et al*, 2012) among various other roles.

We have previously demonstrated that pharmacological interference with the function of

formins, a group of Rho family-associated proteins that nucleate linear filamentous (F) actin,

kinetically delays SV endocytosis at hippocampal boutons and blocks compensatory endocytosis

at the calyx of Held in pre-hearing rats (Soykan *et al.*, 2017). Capacitance measurements at the

calyx of Held in post-hearing rodents suggest a less stringent requirement for formin-mediated

actin assembly for endocytosis (Hori *et al*, 2022). The latter observation may reflect the operation

of compensatory pathways for actin assembly, e.g. via Rac1 or Cdc42 Rho-family GTPases that

promote branched actin networks (Eisenmann *et al*, 2005; Goode & Eck, 2007; Hodge & Ridley,

2016).

In contrast to the vast body of literature regarding the role of actin and actin-associated

proteins at the postsynapse [(Cingolani & Goda, 2008; Colgan & Yasuda, 2014) and references

therein], most notably the actin-rich mesh of the postsynaptic density of glutamatergic synapses,

comparably little is known about the signaling pathways (e.g. via guanine nucleotide exchange

factors and GTPase activating proteins for Rho family small GTPases (Muller *et al*, 2020)) that

mediate actin assembly at the presynapse. Recent work has suggested that presynaptic Rac1

negatively regulates synaptic strength and release probability by altering SV priming and

replenishment at central synapses (O'Neil *et al*, 2021) including the calyx of Held (Keine *et al*,

2022). These data suggest a negative regulatory role for Rac1 in SV exocytosis. The abundance of

postsynaptic actin (Chen *et al*, 2020; Cingolani & Goda, 2008; Colgan & Yasuda, 2014) has

hampered the analysis of the nanoscale localization of actin at presynaptic nerve terminals and of

the actin-regulatory proteins that control its dynamics. KO of Dynamin, the main enzyme for

endocytic membrane fission (Ferguson *et al*, 2007; Imoto *et al*, 2022; Raimondi *et al*, 2011), has

been shown to cause the accumulation of F-actin at and around stalled endocytic intermediates in

non-neuronal cells (Ferguson *et al*, 2009) but this phenotype seems less overt at hippocampal

synapses (Raimondi *et al.*, 2011). Hence, actin in addition to a possible local role at endocytic

invaginations might serve a more general function in endocytosis at synapses, for example by

modulating plasma membrane tension or contractility, by providing a restrictive membrane

scaffold that promotes membrane fission by keeping endocytic invaginations under longitudinal

tension (Boulant *et al.*, 2011; Roux *et al*, 2006; Wu & Chan, 2022), or via formation of an F-actin

ring around the active zone that mechanically couples exocytic membrane compression to

endocytic pit formation (Ogunmowo *et al*, 2023).

Here we combine optical recordings of SV exo-/ endocytosis and ultrastructural analysis

with genetic and pharmacological manipulations to show that interdependent mDia1/3-Rho and

Rac1 signaling pathways cooperate to facilitate SV endocytosis by controlling presynaptic F-actin.

RESULTS

Actin dynamics and actin nucleating mDia1/3 proteins facilitate presynaptic endocytosis and SV

recycling

Previous studies using pharmacological inhibitors of actin polymerization and

depolymerization have yielded inconclusive, often conflicting data regarding the function of actin

in SV endocytosis and recycling in different models (Bleckert *et al*, 2012; Del Signore *et al*, 2021;

Piriya Ananda Babu *et al*, 2020; Sankaranarayanan *et al.*, 2003; Shupliakov *et al.*, 2002; Wu &

Chan, 2022). In contrast, genetic ablation of β- or γ-actin genes in mouse neurons suggests that

actin is required for all forms of endocytosis at several types of synapses (Wu *et al.*, 2016).

Prompted by these findings we revisited the role of actin in presynaptic endocytosis by analyzing

the effects of impaired F-actin dynamics in the combined presence of the G-actin sequestering

drug latrunculin A, the F-actin stabilizer jasplakinolide, and Y-27632, an inhibitor of ROCK kinase

signaling. This cocktail was shown to preserve the existing cytoskeleton architechture while

blocking actin assembly, disassembly and rearrangement (Peng *et al*, 2011). We optically recorded

the stimulation-induced exo-endocytosis of the SV protein Synaptophysin (Syph) fused to a pH-

sensitive super-ecliptic green fluorescent protein (pHluorin) (Miesenbock *et al*, 1998) that is de-

quenched during exocytosis and undergoes re-quenching as SVs are internalized and re-acidified

in hippocampal neurons at physiological temperature. Under these conditions (i.e. trains of APs,

37°C), SV endocytosis occurs on a timescale of > 10 seconds, e.g. a timescale that is slower than

re-quenching of pHluorin due to reacidification (Lopez-Hernandez *et al*, 2022; Soykan *et al.*,

2017). Hence, the decay of pHluorin signals can serve as a measure of the time course of SV

endocytosis. Perturbation of actin dynamics in the combined presence of latrunculin A,

jasplakinolide, and Y-27632 significantly slowed down the endocytic retrieval of exogenously

expressed Syph-pHluorin (Figure 1A,B). Application of Y-27632 alone or a combination of

latrunculin A and jasplakinolide had no effect on SV endocytosis kinetics, whereas combined use

of Y-27632 together with jasplakinolide displayed a mild inhibitory effect (Figure 1- Supplement

1C-F). Impaired actin dynamics in the presence of latrunculin A, jasplakinolide, and Y-27632 did

not impact the apparent level of SV exocytosis indicated by the maximal amplitude of Syph-

pHluorin fluorescence (Figure 1-Supplement 1A). As optical imaging of pHluorin reporters may

lead to artifacts, we analyzed the internalization kinetics of the endogenous vesicular γ-

aminobutyric acid transporter (vGAT) using antibodies directed against its luminal domain

coupled to the pH-sensitive fluorophore CypHer5E (Hua *et al*, 2011; Lopez-Hernandez *et al.*,

2022). The cyanine-based dye CypHer5E is quenched at neutral pH but exhibits bright

fluorescence when present in the acidic lumen of SVs and, thus can serve as a tracer for the exo-

endocytic cycling of endogenous SV proteins. Perturbation of actin dynamics significantly delayed

the endocytic retrieval of endogenous vGAT in response to train stimulation with 200 APs at

physiological temperature (Figure 1C,D), consistent with the results from exogenously expressed

Syph-pHluorin. vGAT exocytosis proceeded unperturbed (Figure 1-Supplement 1B). Hence, a

dynamic actin cytoskeleton facilitates presynaptic endocytosis in hippocampal neurons.

Previous studies have suggested that formins, in particular mDia1, promote SV

endocytosis (Soykan *et al.*, 2017), likely by inducing F-actin assembly (Ganguly *et al*, 2015). We

confirmed that shRNA-mediated depletion of mDia1 from hippocampal neurons led to

significantly delayed Syph-pHluorin endocytosis and re-acidification in response to stimulation

with 200 APs. Re-expression of shRNA-resistant wild-type mDia1 fully restored normal

endocytosis kinetics (Figure 1E,F). Impaired SV endocytosis in lentivirus-mediated mDia1-

depleted neurons (Figure 1-Supplement 1H) was also observed, if SV exo-/ endocytosis was

triggered by stimulation with 40 APs or 80 APs and probed with vesicular glutamate transporter 1

(vGLUT1)-pHluorin (Figure 1G,H and Figure 1-Supplement 1I,K). Exocytic fusion of vGLUT1-

pHluorin-containing SVs was unaffected (Figure 1-Supplement 1G). Co-depletion of the closely

related mDia3 isoform tended to further aggravate this phenotype (Figure 1G,H and Figure 1-

Supplement 1I,K) without impacting the apparent number of fused SVs (Figure 1- Supplement

1J,L), but not vGAT exocytosis (Figure 1-Supplement 1M). Furthermore, mDia1/3 depletion

significantly delayed the endocytic retrieval of endogenous vGAT (Figure 1I,J). Conversely,

application of IMM-01, a small molecule activator of mDia-related formins that acts by relieving

autoinhibition (Lash *et al*, 2013) (Figure 1-Supplement 1N), led to a moderate but significant

acceleration of SV endocytosis monitored by vGLUT1-pHluorin (Figure 1K,L) but no change in

apparent SV fusion (Figure 1- Supplement 1O). Ultrastructural analysis of hippocampal synapses

from mDia1/3-depleted neurons by electron microscopy revealed a reduction in SV density

compared to controls (Figure 2A,B). This phenotype was rescued upon prior silencing of

endogenous neuronal network activity in the presence of tetrodotoxin (TTX) (Figure 2C and Figure

2-Supplement 1A,B). A similar, slightly more pronounced depletion of SVs was observed in

hippocampal synapses from mDia1 knockout (KO) mice (Peng *et al*, 2007; Peng *et al*, 2003;

Shinohara *et al.*, 2012) (Figure 2D,E).

These data are consistent with a role for actin dynamics (see Fig. 1) and actin nucleating

mDia1/3 proteins in presynaptic endocytosis and SV recycling.

mDia facilitates SV endocytosis by regulating presynaptic F-actin

mDia1 in addition to its Rho binding domain and a C-terminal actin-assembling formin

homology 2 (FH2) domain comprises an unstructured N-terminal region involved in membrane

binding (Eisenmann *et al.*, 2005; Ramalingam *et al*, 2010) (Figure 3A). This architecture suggests

that mDia specifically promotes the nucleation of unbranched actin filaments at membranes, for

example to promote endocytosis at the presynaptic plasma membrane. To test this hypothesis, we

assayed the ability of mutant mDia1 lacking its N-terminal membrane-binding region (mDia1-ΔN)

(Ramalingam *et al.*, 2010) to rescue defective SV endocytosis in hippocampal neurons depleted of

endogenous mDia1. Mutant mDia1-ΔN displayed reduced association with membranes in

transfected HEK293T cells (Figure 3-Supplement 1A,B) and was unable to restore normal kinetics

of Syph-pHluorin endocytosis in hippocampal neurons transfected with mDia1-specific

microRNA (Figure 3B and Figure 3-Supplement 1C). Hence, mDia may act on SV endocytosis by

specifically promoting actin assembly at presynaptic membranes. We probed this model at several

levels. We analyzed the nanoscale localization and distribution of mDia1 via multicolor time-gated

stimulated emission depletion (gSTED) microscopy. Hippocampal neurons were stained for

mDia1, the presynaptic active zone marker Bassoon, and post-synaptic Homer1. The distributions

of all three proteins along line profiles perpendicular to the synaptic cleft (i.e. the space between

Bassoon and Homer1 clusters) were analyzed (Gerth *et al*, 2017) (Figure 3-Supplement 1H). This

analysis revealed that endogenous mDia1 is primarily localized to the presynaptic compartment,

where it is concentrated at or very close to the plasma membrane (Figure 3C,D), i.e. < 50 nm away

from the presynaptic membrane (see (Dani *et al*, 2010)). Prompted by these results we dissected

the interactome of mDia1 by unbiased quantitative proteomic analyses of immunoprecipiates from

detergent-extracted mouse synaptosomes (Figure 3-Supplement 1D). In addition to actin, we

identified a variety of additional actin-regulatory proteins including Myosin IIB (Chandrasekar *et*

al, 2013) and the small GTPase Rac1 (Figure 3E). We confirmed the association of mDia1 with

actin and Myosin IIB (Figure 3-Supplement 1E) and found Myosin IIB to be concentrated at or

near the presynaptic plasma membrane (Figure 3-Supplement 1F,G) akin to the localization of

mDia1. The association of mDia1 with Myosin IIB is consistent with the inhibitory effects of the

Myosin II inhibitor blebbistatin on SV endocytosis (Chandrasekar *et al.*, 2013; Soykan *et al.*,

2017). Interestingly, we also observed an enrichment of several endocytic proteins previously

implicated in SV recycling in mDia1 immunoprecipitates. These include the lipid phosphatase

Synaptojanin1, the bin-amphiphysin-rvs (BAR) domain proteins Amphiphysin, Endophilin A1,

and PACSIN1/2 and, the GTPase Dynamin1 (Chanaday *et al.*, 2019; Gan & Watanabe, 2018;

Saheki & De Camilli, 2012; Soykan *et al.*, 2016) (Figure 3E) as confirmed by immunoblotting

(Figure 3F). Based on the association of mDia1 with endocytic proteins including Dynamin1, we

hypothesized that mDia1 might get stalled at endocytic intermediates that accumulate under

conditions of perturbed Dynamin function (Ferguson *et al.*, 2009; Raimondi *et al.*, 2011).

Application of the partially non-selective Dynamin inhibitor Dynasore (Macia *et al*, 2006) (Figure

3C,H) or expression of dominant-negative mutant Dynamin1 K44A (Figure 3G,I) led to a

significant accumulation of mDia1 at presynaptic sites marked by Bassoon. mDia1 levels at the

postsynapse were also increased (Figure 3-Supplement 1I). Dynasore did not impact the levels

(Figure 3- Supplement 1J) or distribution of Bassoon or Homer1.

Based on these results we next investigated the relationship between mDia1/3 and

presynaptic actin. Multicolor gSTED imaging confirmed that F-actin labeled by phalloidin was

highly abundant at the postsynaptic compartment. In addition, F-actin was also visible, albeit at

reduced levels, in presynaptic boutons marked by Bassoon (Figure 4A,B). Inhibition of Dynamin

in the presence of Dynasore (Figure 4C and Figure 4-Supplement 1A) or expression of Dynamin1

K44A (Figure 4-Supplement 1B,C) significantly elevated presynaptic F-actin levels within the

Bassoon area. These data are consistent with a role for presynaptic actin in SV endocytosis. We

directly tested the role of mDia in presynaptic actin nucleation and actin-dependent SV endocytosis

in several ways. First, we generated a mutant version of mDia1 that lacks the ability to nucleate

F-actin due to site-specific inactivation of its FH2 domain (Daou *et al*, 2014; Higashi *et al*, 2008;

Xu *et al*, 2004). We found that actin polymerization-defective mutant mDia1 (K994A) failed to

restore delayed vGLUT1-pHluorin endocytosis in mDia1-depleted hippocampal neurons (Figure

4D and Figure 4-Supplement 1D). Second, we analyzed the effects of depleting hippocampal

neurons of mDia1/3 on presynaptic F-actin. We found that loss of mDia1/3 significantly reduced

the amount of detectable F-actin within the Bassoon area (Figure 4A,E). To unequivocally

distinguish between presynaptic actin at exo-/ endocytic sites and the abundant pool of

postsynaptic actin associated with the postsynaptic density, we used the ORANGE strategy for

CRISPR-based genome-engineering (Willems *et al*, 2020) to generate eGFP-β-actin knock-in (KI)

hippocampal neurons. Due to the low knockin efficiency, this strategy allowed us to visualize

endogenously tagged eGFP-β-actin by 2-color gSTED imaging exclusively in presynaptic boutons

marked by vGLUT1. This assay confirmed the significant reduction in endogenous β-actin at

presynapses from mDia1/3-depleted neurons (Figure 4F,G). Third, we reasoned that if loss of

presynaptic actin in mDia-depleted neurons was causal for the observed defect in SV endocytosis,

pharmacological stabilization of F-actin might rescue the endocytic phenotype. Treatment of

mDia1/3-depleted neurons with jasplakinolide indeed fully restored the kinetics of SV endocytosis

monitored by vGLUT1-pHluorin (Figure 4H and Figure 4-Supplement 1H) and rescued

presynaptic F-actin levels to that of controls (actin levels normalized to shCTR + DMSO set to

100: shCTR + DMSO = 100 ± 6.3; shmDia1+3 + DMSO = 47.7 ± 4.3; shCTR + Jasp = $150.6 \pm$

11.9; shmDia1+3 + Jasp = 94.3 ± 11.5) (Figure 4-Supplement 1G).

We conclude that mDia1/3 facilitates SV endocytosis by regulating presynaptic F-actin.

mDia1/3-Rho and Rac1 signaling pathways cooperatively act to facilitate presynaptic endocytosis

In non-neuronal cells mDia proteins have been shown to be disinhibited by active Rho

family GTPases, in particular RhoA (Otomo *et al*, 2005) (Figure 5A and Figure 1-Supplement

1N). We reasoned that Rho proteins might serve a similar function at presynaptic nerve terminals

in hippocampal neurons. Multicolor gSTED imaging revealed that endogenous RhoA was mostly

located within the presynaptic compartment (Figure 5B,C). Interference with Rho function by co-

expression of dominant-negative (DN) variants of RhoA and RhoB (Figure 5D and Figure 5-

Supplement 1A) or co-depletion of these Rho isoforms (Figure 5-Supplement 1C) delayed SV

endocytosis in response to stimulation with 200 APs without impacting the apparent levels of

exocytosis (Figure 5- Supplement 1B,D). These results are consistent with a model in which active

Rho promotes mDia function and, thereby F-actin nucleation.

F-actin dynamics are known to be controlled by interdependent signaling networks

characterized by feedback regulation between components within the same pathway and between

key regulatory switches (e.g. distinct Rho/Rac family members) that drive distinct forms of F-actin

(see below) (Lawson & Ridley, 2018). For example, active mDia is known to stimulate RhoA

activation (Kitzing *et al*, 2007) (Figure 5A). Consistently, we found that RhoA activity was

reduced in hippocampal neurons depleted of mDia1/3 (Figure 5E,F). Work in non-neuronal cells

has further revealed that RhoA activity antagonizes activation of Rac1 (Chauhan *et al*, 2011), a

key factor for promoting the formation of branched F-actin networks at the cell cortex (Hodge &

Ridley, 2016; Lawson & Ridley, 2018) (Figure 6A). We hypothesized that reciprocally

interdependent Rho/ Rac1 signaling might control presynaptic F-actin assembly and, thereby, SV

endocytosis. In agreement with the alleged antagonistic regulation of RhoA and Rac1 function

(Lawson & Ridley, 2018), we found Rac1 activity to be significantly elevated in hippocampal

neurons depleted of mDia1/3 (Figure 6B,C), e.g. under conditions of reduced Rho-GTP levels. A

similar increase in active Rac1 levels was observed upon pharmacological inhibition of RhoA/B

in the presence of Rhosin (Figure 6-Supplement 1A). Multicolor gSTED imaging showed that

Rac1 was equally distributed between pre- and postsynaptic compartments in unperturbed

hippocampal neurons (Figure 6D,E). Elevated active Rac1 levels might conceivably ameliorate

presynaptic endocytic phenotypes elicted by mDia1/3 loss. In support of this hypothesis, we found

that selective pharmacological inhibition of Rac1 reduced presynaptic actin levels (Figure 6-

Supplement 1B,C) and caused a delay in the endocytic retrieval of endogenous vGAT (Figure

6F,G). Moreover, Rac1 inhibition appeared to further aggravate impaired vGAT endocytosis in

mDia1/3-depleted neurons, although the effect remained below statistical significance. Expression

of constitutively active GTP-locked Rac1 (Rac1-CA) restored normal Syph-pHluorin endocytosis

kinetics in mDia1/3-depleted neurons, whereas Syph-pHluorin endocytosis was delayed upon

expression of dominant-negative Rac1 (Rac1-DN) and exacerbated the endocytic phenotype of

mDia1/3 loss (Figure 6H and Figure 6-Supplement 1E). No overt effect of Rac1 inhibition or

overexpression of either Rac1 form on SV exocytosis was observed (Figure 6-Supplement 1D,F).

Perturbation of the related Cdc42 protein, another actin regulatory GTPase (Lawson & Ridley,

2018) found at synapses (Figure 6-Supplement 1G-I), did not significantly affect the kinetics of

SV endocytosis in control or mDia1/3-depleted neurons (Figure 6- Supplement 1J,K). These data

suggest that interdependent mDia1/3 and Rac1-based pathways control SV endocytosis via

presynaptic actin.

We probed this model by analyzing the ultrastructure of synapses from hippocampal

neurons depleted of mDia1/3 or following inhibition of Rac1 activity. Synapses from mDia1/3-

depleted neurons displayed an accumulation of non-coated plasma membrane invaginations that

were found both in the vicinity of the active zone as well as distal from the synaptic contact area

(Figure 7A,B). Moreover, we observed elevated numbers of endosome-like vacuoles (ELVs)

(Figure 7A,C) that might serve as a donor membrane for SV reformation (Kononenko & Haucke,

2015; Kononenko *et al.*, 2014; Watanabe *et al.*, 2014) as suggested by the observed reduction in

SV numbers in mDia-depleted neurons (compare Figure 2). A prominent accumulation of plasma

membrane invaginations (Figure 7-Supplement 1A) and endosome-like vacuoles (Figure 7-

Supplement 1B) was similarly found in hippocampal neurons from mDia1 KO mice. Inhibition of

Rac1 function by EHT 1864 also led to the accumulation of non-coated plasma membrane

invaginations (Figure 7A,D), whereas the number of endosome-like vacuoles was not significantly

altered (Figure 7A, E). Pharmacological blockade of Rac1 activity in neurons depleted of mDia1/3

further increased the number of plasma membrane invaginations significantly, while the additional

effect on the number of endosome-like vacuoles remained insignificant (Figure 7F,G,H). mDia1/3

loss and Rac1 perturbation exhibited similar phenotypes with respect to the length and width of

tubular membrane invaginations (Figure 7-Supplement 1C,D).

Collectively, these findings demonstrate that mDia1/3-Rho and Rac1 signaling pathways

cooperatively act to facilitate presynaptic endocytosis and SV recycling.

DISCUSSION

We found that loss of mDia1 either alone or in conjunction with the closely related mDia3 isoform

slows the kinetics of SV endocytosis of exogenous pHluorin-tagged SV proteins and of

endogenous vGAT without perturbing exocytic SV fusion. These phenotypes are accompanied by

the accumulation of plasma membrane invaginations and vacuolar structures as well as an activity-

dependent reduction of the total SV pool (Figures 1, 2, and 7). Furthrmore, we observe that

interdependent mDia1/3-Rho and Rac1 signaling pathways cooperatively act to facilitate

presynaptic endocytosis (Figures 5, 6 and 7). Several lines of evidence indicate that these endocytic

phenotypes are a consequence of perturbed presynaptic actin levels or dynamics. (i) Endogenous

presynaptic F-actin is reduced in mDia1/3-depleted hippocampal synapses (Figure 4) and (ii) actin

polymerization-defective mDia1 fails to restore normal endocytosis kinetics in hippocampal

neurons depleted of mDia1 (Figure 4). Conversely, (iii) pharmacological stabilization of F-actin

by jasplakinolide (Figure 4) or (iv) expression of constitutively-active Rac1 to drive compensatory

actin polymerization (for example via other formins and/ or ARP2/3) (Figure 6) rescues

endocytosis in mDia1/3-KD hippocampal neurons. Moreover, (v) we find mDia1 (Figure 3) and

F-actin (Figure 4) to accumulate at presynapses under conditions of impaired dynamin-dependent

endocytosis. Together with the observation that inhibition of F-actin dynamics (Peng *et al.*, 2011)

in the combined presence of latrunculin A, jasplakinolide, and Y-27632 (Figure 1) slows the

kinetics of SV endocytosis, our data support the hypothesis that presynaptic actin facilitates SV

endocytosis downstream of mDia/ RhoA and Rac1-based signaling pathways.

Our data extend previous studies using conditional genetics that have identified an actin

requirement for SV endocytosis (Wu *et al.*, 2016). The endocytic phenotype described here for

loss of mDia1/3, however, is substantially milder than that elicited by complete loss of β– or γ-

actin, likely reflecting the fact that several partially redundant pathways cooperate to promote

F-actin assembly at synapses (see below). In contrast to β- or γ-actin KO hippocampal neurons

(Wu *et al.*, 2016) we did not observe strong exocytic depression in response to train stimulation in

mDia-depleted neurons (e.g. Figure 1-Supplement 1A-M) although we cannot rule out mild

differences in synaptic strength and/ or release probability as observed in Rac1 KO hippocampal

neurons and at the calyx of Held (Keine *et al.*, 2022; O'Neil *et al.*, 2021).

At the ultrastructural level, we find that genetic (KO) or lentiviral induced (KD) loss of

mDia1 or mDia1/3 partially phenocopies acute perturbation of formin-mediated actin assembly

(Soykan *et al.*, 2017) with respect to the accumulation of plasma membrane invaginations. Unlike

acute formin inhibition (Soykan *et al.*, 2017), we find elevated numbers of post-fission endosome-

like vacuoles in mDia-depleted hippocampal synapses. The latter correlate with a mild reduction

of SV numbers, suggesting that mDia may have a further role in SV reformation downstream of

the actual endocytosis reaction (Chanaday *et al.*, 2019; Gan & Watanabe, 2018; Kononenko *et al.*,

2014). The identified function of mDia1/3-mediated presynaptic actin assembly in presynaptic

endocytosis and SV recycling is consistent with the actin binding properties of many endocytic

proteins including Dynamin (Ferguson *et al.*, 2009; Gu *et al*, 2014), PACSINs (Kessels &

Qualmann, 2006), or Epsins, among others (McMahon & Boucrot, 2011; Merrifield *et al.*, 2005;

Saheki & De Camilli, 2012). How exactly actin functions to facilitate endocytosis of SV

membranes remains to be fully understood. The comparably small size of the membrane

invaginations that accumulate following pharmacological inhibition of formins (< 100 nm) or Rac1

or upon genetic loss of mDia1 (typically < 150 nm, although larger ones are observed) render an

actomyosin-based constriction mechanism for SV endocytosis unlikely (consistent with

(Sankaranarayanan *et al.*, 2003)). We rather favor a - possibly indirect - function of F-actin in

presynaptic endocytosis via formation of an F-actin ring that couples exocytic membrane

compression to endocytic pit formation (Ogunmowo *et al.*, 2023). Such a model is supported by

the abundance of large plasma membrane invaginations distal from the active zone where SV

fusion occurs at synapses following mDia1/3 and Rac1 loss of function (Figure 7). Consistent with

this, it has been shown that mDia1 controls presynaptic membrane contractility (Deguchi *et al*,

2016), e.g. via the recently identified presynaptic actin corrals visualized in genome-engineered

rat hippocampal neurons (Bingham *et al*, 2023).

A key result from our study is the finding that presynaptic endocytosis is regulated by

interdependent, yet partially redundant signaling for F-actin assembly downstream of Rho and

Rac1, e.g. pathways known to control linear and branched F-actin networks (Lawson & Ridley,

2018; Muller *et al.*, 2020) that were shown to co-exist within hippocampal presynaptic boutons

(Bingham *et al.*, 2023). We demonstrate that loss of mDia1/3 or inhibition of Rho cause the

hyperactivation of the actin-promoting GTPase Rac1 and that dual loss of mDia1/3 loss and Rac1

function results in synergistic inhibitory effects with respect to presynaptic endocytosis and

ultrastructural defects (Figures 5-7). These findings suggest that the signaling network that

regulates presynaptic actin assembly may have been evolutionary selected for plasticity and

resilience. A resilient network for the control of actin dynamics at the presynapse may also help

to explain the partially contradicting results from drug-based manipulations of actin and actin

regulatory factors in different models of presynaptic neurotransmission reported in the past

(Bleckert *et al.*, 2012; Del Signore *et al.*, 2021; Hori *et al.*, 2022; Piriya Ananda Babu *et al.*, 2020;

Richards *et al*, 2004; Sankaranarayanan *et al.*, 2003; Shupliakov *et al.*, 2002; Wu & Chan, 2022;

Wu *et al.*, 2016). Moreover, accumulating evidence indicates that different pools of actin might be

differentially amenable to drug treatments (Bleckert *et al.*, 2012), e.g. as a consequence of the

association of actin filaments with molecules such as tropomyosin (Gormal *et al*, 2017) that render

them resistant to drugs, and/ or variations in the fraction of F-actin *vs*. monomeric actin pools

(Higashida *et al*, 2008), providing a possible explanation for previously discrepant results.

A number of open questions remain. We predict that presynaptic actin assembly via mDia/

Rho and Rac1 facilitates all forms of presynaptic endocytosis that operate on timescales of

milliseconds to seconds (Chanaday & Kavalali, 2018; Clayton & Cousin, 2009; Watanabe &

Boucrot, 2017) and at many types of synapses (e.g. small central synapses, ribbon synapses, the

neuromuscular junction), but further studies will be needed to test this hypothesis. We note that

fast endophilin-mediated endocytosis (FEME), a process with resemblance to endocytosis at

synapses (Watanabe & Boucrot, 2017), has been shown to be differentially regulated by actin and

Rho GTPase family members (Boucrot *et al*, 2015; Renard *et al*, 2015). At present we do not

understand if and how RhoA/B and Rac1 GTPase activation is nested into the SV cycle and

whether these GTPases operate at the same or different nanoscale sites and on the same or different

membranes. Future studies will be required to tackle these questions.

MATERIALS AND METHODS

Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
genetic reagent (Mus musculus)	C57BL/6N wild type	Leibniz Research Institute for Molecular Pharmacology	RRID:IMSR_JAX:000664	
genetic reagent (Mus musculus)	mDia1 KO	Bartolini Lab, Columbia University		
cell line (Homo sapiens)	HEK293T	American Type Culture Collection	Cat# CRL-3216; RRID:CVCL_0063	
antibody	β-Actin (mouse monoclonal)	Sigma-Aldrich	Cat# A5441; RRID:AB_476744	IB (1:5000)
antibody	Bassoon (guinea pig monoclonal)	Synaptic Systems	Cat# 141 318; RRID:AB_2927388	IC (1:100)
antibody	CDC42 (rabbit polyclonal)	Abcam	Cat# ab64533; RRID:AB_1310067	IB (1:1000); IC (1:100)
antibody	Dynamin1 (rabbit polyclonal)	Pietro D. Camilli	Shupliakov et al., 1997; DOI: 10.1126/science.276.5310.259	IB (1:2000)
antibody	GFP (mouse monoclonal)	Thermo Fisher Scientific	Cat# A-11120; RRID:AB_221568	IC (1:2500)
antibody	Homer1 (mouse monoclonal)	Synaptic Systems	Cat# 160 011; RRID:AB_2120992	IC (1:200)

antibody	Homer1 (rabbit polyclonal)	Synaptic Systems	Cat# 160 003; RRID:AB_887730	IC (1:200)
antibody	LAMP1 (rabbit monoclonal)	Cell Signaling Technology	Cat# 9091; RRID:AB_2687579	IB (1:1000)
antibody	mDia1 (mouse monoclonal)	BD Biosciences	Cat# 610848; RRID:AB_398167	IB (1:500); IP (2μg)
antibody	mDia1 (rabbit monoclonal)	Abcam	Cat# ab129167; RRID:AB_11143749	IC (1:100)
antibody	mDia3 (rabbit polyclonal)	Sigma-Aldrich	Cat# HPA005647; RRID:AB_1078657	IB (1:1000)
antibody	Myosin IIb (mouse monoclonal)	Abcam	Cat# ab684; RRID:AB_305661	IC (1:100)
antibody	Myosin IIb (rabbit polyclonal)	Cell Signaling Technology	Cat# 3404; RRID:AB_126421	IB (1:2000)
antibody	Rac1 (mouse monoclonal)	BD Biosciences	Cat# 610650; RRID:AB_397977	IB (1:1000); IC (1:50)
antibody	RFP (rabbit polyclonal)	Takara Bio	Cat# 632496; RRID:AB_10013483	IC (1:500)
antibody	RhoA (rabbit monoclonal)	Cell Signaling Technology	Cat# 2117; RRID:AB_10693922	IB (1:1000); IC (1:100)
antibody	α-Tubulin (mouse monoclonal)	Sigma-Aldrich	Cat# T9026; RRID:AB_47759	IB (1:5000)

antibody	IgG control (mouse monoclonal)	Thermo Fisher Scientific	Cat# 31903; RRID:AB_1095 9891	IP (2 μg)
antibody	vGAT-CypHer5E (rabbit polyclonal)	Synaptic Systems	Cat# 131 103CpH; RRID:AB_2189 809	Live-imaging (1:500)
antibody	Anti-mouse IgG Atto542 (donkey polyclonal)	Martin Lehmann	Gonschior et al., 2022; https://doi.org/1 0.1038/s41467-022-32533-4	IC (1:400)
antibody	Anti-rabbit IgG Atto542 (donkey polyclonal)	Martin Lehmann	Gonschior et al., 2022; https://doi.org/1 0.1038/s41467-022-32533-4	IC (1:400)
antibody	Anti-mouse Alexa Fluor™ 594 (goat polyclonal)	Thermo Fisher Scientific	Cat# A-11032; RRID:AB_2534 091	IC (1:200)
antibody	Anti-rabbit Alexa Fluor™ 594 (goat polyclonal)	Thermo Fisher Scientific	Cat# A-11037; RRID:AB_2534 095	IC (1:200)
antibody	Anti-guinea pig Atto647N (camelid monoclonal)	Synaptic Systems	Cat# N0602-At647N-S; RRID:AB_2744 576	IC (1:200)
antibody	Anti-mouse Alexa Fluor™ 647 (goat polyclonal)	Thermo Fisher Scientific	Cat# A-21235; RRID:AB_2535 804	IC (1:200)

antibody	Anti-mouse IRDye® 680RD (goat polyclonal)	LI-COR Biosciences	Cat# 925-68070; RRID:AB_2651 128	IB (1:10000)
antibody	Anti-rabbit IRDye® 680RD (goat polyclonal)	LI-COR Biosciences	Cat# 926-68071; RRID:AB_1095 6166	IB (1:10000)
antibody	Anti-mouse IgG, IRDye® 800CW (goat polyclonal)	LI-COR Biosciences	Cat# 926-32210; RRID:AB_6218 42	IB (1:10000)
antibody	Anti-rabbit IgG, IRDye® 800CW (goat polyclonal)	LI-COR Biosciences	Cat# 926-32211; RRID:AB_6218 43)	IB (1:10000)
chemical compound, drug	Phalloidin-Alexa FluorTM 594	AAT Bioquest	Cat# ABD-23158	IC (1:1000)
recombinant DNA reagent	Synaptophysin-pHluorin	Leon Lagnado	Granseth et al., 2006; DOI: 10.1016/j.neuron.2006.08.029	Expresses rat Synaptophysin-2xpHluorin (inserted between Asn183 – Thr184) under a CMV promotor
recombinant DNA reagent	vGlut1-pHluorin	Volker Haucke	Bolz et al., 2023; https://doi.org/10.1016/j.neuron.2023.08.016	Expresses rat vGlut1-pHluorin (inserted between Gly99 – Gly100) under a hSyn1 promotor; lentiviral plasmid
recombinant DNA reagent	mCherry	Clontech	Cat# 632523	Expresses mCherry under a CMV promotor
recombinant DNA reagent	mDia1-WT-mCherry	This study	-	Expresses mouse mDia1-WT-mCherry under a CMV promotor
recombinant DNA reagent	mDia1-ΔN -mCherry	This study	-	Expresses truncation (first 60 AA) variant of mouse mDia1-mCherry under a CMV promotor

recombinant DNA reagent	mDia1-WT-SNAP	This study	-	Expresses mouse mDia1-WT-SNAP under a hSyn1 promotor; lentiviral plasmid
recombinant DNA reagent	mDia1-K994A-SNAP	This study	-	Expresses K994A variant of mDia1-SNAP under a hSyn1 promotor; lentiviral plasmid
recombinant DNA reagent	Dynamin1-WT-SNAP	This study	-	Expresses mouse Dynamin1-WT-SNAP under a hSyn1 promotor; lentiviral plasmid
recombinant DNA reagent	Dynamin1-K44A-SNAP	This study	-	Expresses K44A variant of mouse Dynamin1-SNAP under a hSyn1 promotor; lentiviral plasmid
recombinant DNA reagent	vGlut1-mCherry	Franck Polleux	Kwon et al., 2016; https://doi.org/10.1371/journal.pbio.1002516	Expresses rat vGlut1-mCherry under a CAG promotor
recombinant DNA reagent	RhoA-WT	Theofilos Papadopoulos	Reddy-Alla et al. 2010; https://doi.org/10.1111/j.1460-9568.2010.07149.x	Expresses human 3xHA-RhoA-WT under a CMV promotor
recombinant DNA reagent	RhoA-DN	Theofilos Papadopoulos	Reddy-Alla et al. 2010; https://doi.org/10.1111/j.1460-9568.2010.07149.x	Expresses T19N variant of human 3xHA-RhoA under a CMV promotor
recombinant DNA reagent	RhoB-WT	Harry Mellor	Mellor et al., 1998; https://doi.org/10.1074/jbc.273.9.4811	Expresses human myc-RhoB-WT under a CMV promotor

recombinant DNA reagent	RhoB-DN	Harry Mellor	Mellor et al., 1998; https://doi.org/10.1074/jbc.273.9.4811	Expresses T19N variant of human myc-RhoB under a CMV promotor
recombinant DNA reagent	shCTR (transfected)	Sigma-Aldrich	Cat# SHC016	No murine targets
recombinant DNA reagent	shmDia1 (transfected)	Sigma-Aldrich	Cat# TRCN0000108685	Targets 3'UTR of NM_007858
recombinant DNA reagent	shRhoA (transfected)	Sigma-Aldrich	Cat# TRCN0000302388	Targets CDS of NM_016802
recombinant DNA reagent	shRhoB (transfected)	Sigma-Aldrich	Cat# TRCN0000294874	Targets CDS of NM_007483
recombinant DNA reagent	shCTRmiR	This study	-	Expresses rat Synaptophysin-2xpHluorin under a CMV promotor and shRNA embedded into a microRNA (shRNAmiR) with no murine targets
recombinant DNA reagent	shmDia1miR	This study	-	Expresses rat Synaptophysin-2xpHluorin under a CMV promotor and shRNAmiR against the CDS of mouse mDia1
recombinant DNA reagent	shCTR (transduced)	Christian Rosenmund	Watanabe et al., 2014; https://doi.org/10.1038/nature13846	Expresses NLS-RFP or BFP under a hSyn1 promotor and shRNA against no murine target (msClathrin scrambled) under a U6 promotor
recombinant DNA reagent	shmDia1 (transduced)	This study	-	Expresses NLS-RFP or BFP under a hSyn1 promotor and shRNA against the 3'UTR of

				mouse mDia1 under a U6 promotor
recombinant DNA reagent	shmDia3 (transduced)	This study	-	Expresses NLS-RFP or BFP under a hSyn1 promotor and shRNA against the CDS of mouse mDia3 under a U6 promotor
recombinant DNA reagent	MD2.G	Addgene	Cat# 12259; RRID:Addgene _12259	Expresses lentiviral VSV-G envelope protein
recombinant DNA reagent	psPAX2	Addgene	Cat# 12260; RRID:Addgene _12260	Expresses lentiviral packaging protein
recombinant DNA reagent	pOrange GFP-β-Actin KI	Addgene	Cat#131479; RRID:Addgene _131479	gRNA and GFP donor for endogenous N-terminal tagging of β-Actin (amino acid position: D2)
chemical compound, drug	Dynasore	Sigma-Aldrich	Cat# D7693	80 μM 10 min
chemical compound, drug	EHT 1864	MedChemExpress	Cat# HY-16659	10 μM acute (CypHer); 2 h (IC/EM)
chemical compound, drug	IMM-01	Sigma-Aldrich	Cat# SML1064	10 μM acute
chemical compound, drug	Jasplakinolide	Sigma-Aldrich	Cat# J4580	8 μM acute (JLY); 1 μM 30 min
chemical compound, drug	Latrunculin A	Cayman Chemical	Cat# CAY10684	5 μM acute (JLY)
chemical compound, drug	ML141	MedChemExpress	Cat# HY-12755	10 μM acute
chemical compound, drug	Rhosin	MedChemExpress	Cat# HY-12646	10 μM 2 h
chemical compound, drug	Tetrodotoxin	Carl Roth	Cat# 6973.1	1 μM 36 h
chemical compound, drug	Y-27632	Tocris	Cat# 1254	1 μM acute (JLY)

commercial assay or kit	Rhotekin-Rho binding domain (RBD) beads	Cytoskeleton Inc.	Cat# RT02	60 μg
commercial assay or kit	PAK-p21 binding domain (PBD) beads	Cytoskeleton Inc.	Cat# PAK02	20 μg
commercial assay or kit	ProFection Mammalian Transfection System – Calcium Phosphate	Promega	Cat# E1200	
commercial assay or kit	Gibson Master Mix	New England Biolabs Inc.	Cat# E2611L	
commercial assay or kit	Q5 site-directed mutagenesis kit	New England Biolabs Inc.	Cat# E0552S	
software, algorithm	Fiji	NIH	RRID:SCR_002 285	
software, algorithm	Prism	GraphPad	RRID:SCR_002 798	Version 9.5.1.
software, algorithm	Image Lab Software	Bio-Rad	RRID:SCR_014 210	Version 6.0.1
software, algorithm	Image Studio Lite	LI-COR Biosciences	RRID:SCR_013 715	Version 5.2.5
software, algorithm	pHluorin ROI sector	Github	https://github.com/DennisVoll/pHluorin_ROI_selector/; RRID:SCR_002 630	
software, algorithm	SynActJ	Martin Lehmann	Schmied et al., 2021;	

| software, algorithm | Macro_plot_lineprofile_multicolor | Kees Straatman | https://doi.org/10.3389/fcomp.2021.777837 F. Gerth et al., 2019; https://doi.org/10.1016/j.str.2019.03.020 | |
| software, algorithm | MaxQuant | Jürgen Cox | https://www.maxquant.org/maxquant/ | Version 1.6.1.0 |

Materials

Animals

Primary neurons were obtained from either wild-type C57BL/6J (Charles River, RRID:

IMSR_JAX:000664) or mDia1 KO mice (Peng *et al.*, 2007; Peng *et al.*, 2003). All animal

experiments were reviewed and approved by the ethics committee of the *Landesamt für*

Gesundheit und Soziales (LAGeSo) Berlin or the Committee on the Ethics of Animal Experiments

of Columbia University and conducted according to the committees' guidelines (LAGeSo) or the

Guide for the Care and Use of Laboratory Animals of the National Institutes of Health (for mDia1

KO mice). At the facilities, animal care officers monitored compliance with all regulations. Mice

were group-housed under 12/12 h light/dark cycle with access to food and water *ad libitum*. Mice

from both genders were used and cultures were randomly allocated to experimental groups (e.g.,

different treatments). Multiple independent experiments using several biological replicates were

carried out as indicated in the Figure legends.

Antibodies

Antibodies and their working dilutions used in this study are denoted in the Key Resource Table

(IB: Immunoblot; IC: Immunocytochemistry; IP: Immunoprecipitation). Antibodies were stored

according to the manufacturer's recommendations. All secondary antibodies are species-specific

(highly cross-adsorbed).

Cell lines

Human embryonic kidney 293T (HEK293T) cells were obtained from the American Type Culture

Collection (Cat# CRL-3216; RRID:CVCL_0063). Cells were cultured in Dulbecco's modified

Eagle's medium supplemented with glucose (DMEM; 4.5 g/L; Thermo Fisher Scientific) and 10%

heat-inactivated fetal bovine serum (FBS; Gibco), penicillin (100 U/ml; Gibco), and streptomycin

(100 μg/ml; Gibco) at 37°C and 5% CO_2. Cells were routinely tested for mycoplasma

contamination.

Chemicals

Compounds were dissolved in dimethyl sulfoxide (DMSO), unless indicated otherwise, and

diluted 1: 1000 to their working concentrations (s. Key Resource Table). For acute

pharmacological treatment (in pHluorin/CypHer assays), drugs were added to the imaging buffer.

For longer incubations, the conditioned cell media was supplemented with the chemicals (s. Key

Resource Table for incubation time).

For silencing neuronal network activity, sodium channels were inhibited by addition of

tetrodotoxin (TTX; in 10 mM sodium acetate, pH 5.3) to the neuronal culture medium at day in

vitro (DIV) 12 for 36 h. As a control, cells from the same preparation were treated with equal

volumes of 10 mM sodium acetate (annotated as Vehicle).

Plasmids

All recombinant DNA reagents used for protein expression are listed in the Key Resource Table: Synaptophysin-pHluorin was a kind gift from Dr. L. Lagnado (Univ. of Sussex, UK), vGLUT1-pHluorin was generated in-house by Svenja Bolz as previously described, while vGLUT1-mCherry was kindly provided by Dr. Franck Polleux (New York, USA). HA-tagged RhoA-WT and RhoA-T19N were kind gifts from Dr. Theofilos Papadopoulos (Göttingen, Germany) and myc-tagged RhoB-WT and RhoB-T19N were kindly gifted by Dr. Harry Mellor (Bristol, UK). All other expression vectors were generated for this study. Plasmids based on pEGFP-C1, pmCherry-N1, pCAG and pcDNA3 utilising a CMV or CAG promotor (s. Key Resource table) were used for overexpression based on transfection of neuronal cells, while pFUGW vectors carrying a human Synapsin1 promotor (hSyn1) were used for the generation of lentiviral particles for the transduction of neurons (s. Key Resource Table annotated as "lentiviral plasmid"). In the course of this study several approaches to deplete mDia1 have been carried out and are annotated in the Figure legends: For knockdown of mDia1, RhoA and RhoB by transfection, commercially available lentiviral small hairpin RNA (shRNA) vectors based on the pLKO.1 backbone (s. Key Resource Table; annotated as "transfected") were purchased from Sigma-Aldrich. To reduce the amount of DNA needed for transfection to perform pHluorin assays, vectors expressing shRNA embedded into a microRNA (miRNA) context for mDia1 together with Synaptophysin-2x-pHluorin as a reporter were cloned based on pRRLsinPPT-emGFP-miR Control, a kind gift of Dr. Peter S. McPherson (Montreal, Canada). Finally, lentiviral vectors for knockdown of mDia1 and mDia3 (annotated in Key Resource Table as "transduced") were generated based on the backbone f(U6) sNLS-RFPw shCTR, a kind gift from Prof. Christian Rosenmund (Berlin, Germany). All vectors used for genetic depletion via RNA interference are listed in the Key Resource Table and

express gene-specific shRNA under a U6 promotor, that either targets the coding sequence (CDS)

or the 3'-untranslated region (3'-UTR).

Oligonucleotides

Plasmids generated for this study were cloned using oligonucleotides (BioTez GmbH, Berlin,

Germany) listed in Supporting Table 1 (lower case denotes nucleotides that do not anneal to the

backbone; underlined nucleotides denote sense and antisense sequences of shRNA).

Methods

Generation of expression plasmids

Expression plasmids generated for this study were cloned by PCR amplification (Phusion™ High-

Fidelity DNA polymerase) and restriction enzyme (Thermo Fisher Scientific; Fast digest) digest

according to the manufacturers' manual. pSNAP-N1 was generated in-house by Hannes Gonschior

by sub-cloning pSNAP$_f$ (New England BioLabs Inc., Cat#N9183S) via PCR and the restriction

enzymes *AgeI* and *NotI* into pmCherry-N1. pFUG_hSyn_MCS was cloned by Amirreza Ohadi to

enable simple insertion of SNAP-tagged proteins by inserting a multiple cloning site (MCS) on

annealed oligonucleotides into a pFUG_hSyn1 backbone (a gift from Christian Rosenmund,

Berlin, Germany) cut by *AgeI* and *EcoRI*. To generate mDia1-WT-mCherry, the coding sequence

of mDia1 was cut from mDia1-mEmerald-N1 (Addgene, Cat#54157) and pasted into pmCherry-

N1 by *AgeI* and *XhoI* digestion. mDia1-WT-SNAP was cloned by cutting out the coding sequence

of mDia1 from mDia1-mEmerald-N1 and pasting it into pSNAP-N1 by *AgeI* and *NheI* digest of

both vector and insert. To subclone mDia1-SNAP into a lentiviral vector the coding sequence of

mDia1-SNAP was pasted from mDia1-WT-SNAP-N1 into pFUG_hSyn1_MCS utilizing common

cut sites for *NheI* and *NotI* digest. For the generation of a lentiviral vector expressing Dynamin-

WT, the coding sequence of Dynamin1 was first extracted from Dynamin1-pmCherry-N1

(Addgene; Cat#27697) by *EcoRI* and *XmaI* digestion and cloned into pSNAP-N1 with the same

enzymes. Subsequently, Dynamin1-SNAP was isolated by *NheI* and *NotI* digest and transformed

into pFUG_hSyn1_MCS. The mDia1- ΔN truncation variant was generated by introducing a new

start codon on a primer shifted by 60 amino acids (s. Supporting Table 1), PCR amplifying the

truncated DNA and subcloning the template into pmCherry-N1 by *BglII* and *AgeI* digest. For

introducing point mutations in mDia1 (Lysine-994 to Alanine; K994A) and Dynamin1 (Lysine-44

to Alanine; K44A), the Q5 site-directed mutagenesis kit (New England Biolabs Inc.; E0552S) was

used according to the manufacturer's manual and oligonucleotides listed in Supporting Table 1.

All vectors were confirmed by Sanger sequencing (LGC Genomics, Berlin, Germany)

and amplified by self-made chemically competent TOP10 *E.coli* before purifying respective

endotoxin-free DNA by 2-propanole precipitation.

Isolation, culture and transfection of primary hippocampal neurons

Neuronal hippocampal cell cultures were prepared as described before (Lopez-Hernandez *et al.*,

2022; Soykan *et al.*, 2017). In short, hippocampi from postnatal mice (p0-p3) were surgically

isolated and dissociated into single cells by trypsin (5 g/L, Sigma-Aldrich) digestion. Neurons

(100000 cells/ well of a 6-well) were plated onto poly-L-lysine-coated coverslips and grown in

modified Eagle medium (MEM; Thermo Fisher) supplemented with 5% FCS and 2% B-27 (Gibco)

and maintained at 5% CO_2 and 37°C in humidified incubators. In addition, 2 μM cytosine β-D-

arabinofuranoside (AraC) was added to the cell culture media in the first 2 days in vitro (DIV) to

limit glial proliferation. For transient protein expression, neurons were transfected on DIV 7-9

utilizing a Calcium phosphate transfection kit (Promega; Cat# E1200): In brief, 1-6 μg plasmid

DNA (per well of a 6-well plate) were mixed with 250 mM calcium phosphate ($CaCl_2$) in ultrapure

nuclease-free water. The resulting solution was added to equal volumes of 2x 4-(2-hydroxyethyl)-

1-piperazineethanesulfonic acid buffered saline (2x HEPES; 100 µL) and incubated at room

temperature for 20 min. Resulting precipitates were added dropwise to cells that had been

transferred to osmolarity-adjusted Neurobasal-A (NBA; Gibco) media to induce starvation. After

incubation at 37°C and 5% CO_2 for 30 min, neurons were washed thrice with osmolarity-adjusted

Hank's balanced salt solutions (HBSS; Gibco) and transferred back to their original conditioned

media.

shRNA cloning and lentivirus production

Knockdown of proteins was achieved through RNA interference either by $CaCl_2$ transfection of

gene-specific shRNA encoding vectors (pLKO.1; Sigma-Aldrich or shRNAmiR) on DIV 7 or by

transduction of cells with lentiviral particles harboring the gene-specific shRNA on DIV 2. The

respective method is indicated in the Figure legends as *transfected* or *transduced*. To reduce the

amount of DNA needed for transfection and to improve neuronal health, shRNAmiR were

expressed from the 3'UTR of Synaptophysin-pHluorin based on (Ritter *et al*, 2017): The reporter

protein/synthetic cassette was generated by PCR amplifying Synaptophysin-2x-pHluorin with

oligos (s. Supporting Table 1) harboring *XbaI* and *SalI* sites. The eGFP in the RRLsinPPT-eGFP-

miRCTR plasmid (a kind gift from Dr. Peter S. McPherson, McGill University, Canada) was

replaced by the PCR product through similar restriction digest to yield shCTRmiR. miRmDia1

(start position 4892, targeting sequence matches open-reading frame) was designed with BLOCK-

iT™ (Thermo Fisher Scientific) and subcloned into shCTRmiR to yield shmDia1miR following

protocol in (Ritter *et al.*, 2017).

As both transfection strategies were limited by low efficiency, lentiviral knockdown was

carried out: Lentiviral particles were based on a shuttle vector (pFUGw) driving the expression of

a nuclear-targeted red fluorescent protein (NLS-RFP) under a human Synapsin1 (hSyn1) promotor

to monitor infection efficiency in neurons and a scrambled mouse shRNA against Clathrin without

any murine targets, which was used as the control virus (f(U6) sNLS-RFPw shCTR). To prevent

crosstalk between the NLS-RFP and other mCherry constructs, a similar backbone expressing the

blue fluorescent protein (BFP) as a reporter was generated in house by Klaas Ypermann: The

backbone was digested by *XbaI* and *PacI*, the hSyn1 promotor and mTagBFP (Addgene; #Cat

105772) were amplified with oligos (s. Supporting Table 1), gel extracted and assembled utilizing

a Gibson master mix (New England BioLabs Inc.; Cat#E2611L) to yield f(U6) BFP shCTR. To

knockdown mDia1, a shRNA sequence based on 5'-GCCTAAATGGTCAAGGAGATA-3' as the

sense nucleotide corresponding to the 3'UTR of mouse mDia1 (NM_007858.4; Sigma-Aldrich,

Cat# TRCN0000108685) was designed as an oligonucleotide with overhangs (s. Supporting Table

1) and annealed into the f(U6) sNLS-RFPw shCTR backbone cut with *BamHI* and *PacI* to yield

the vector f(U6) sNLS-RFPw/BFP shmDia1. For mDia3 shRNA, a sense sequence based on

5'-GCCCTAATCCAGAATCTTGTA-3' corresponding to nucleotides 2473-2493 of mouse

mDia3 (DIAPH2; NM_172493.2; Sigma-Aldrich, Cat# TRCN0000108782) was used to construct

f(U6) sNLS-RFPw/BFP shmDia3 as described above. Production of lentiviral particles was

conducted using the second-generation packaging system: In brief, HEK293T were co-transfected

with lentiviral shRNA constructs and the packaging plasmids psPAX2 (Addgene; Cat# 12260) and

MD2.G (Addgene; Cat# 12259) using $CaCl_2$. After 12 h, the cell media was replaced. Virus-

containing supernatants were collected at 48 and 72 h after transfection, filtered to remove cell

debris and particles were concentrated (30-fold) via low-speed centrifugation (Amicon Ultra-15,

Ultracel-100; Merck Millipore; Cat# UFC9100) before aliquoting and storage at -70°C. For all

experiments, an infection rate of over 95% was achieved at DIV 14-16.

Microscopy

Live-imaging of SV recycling

For live-imaging of Synaptophysin and vesicular glutamate transporter 1 (vGLUT1) recycling, SV

proteins fused to the green-fluorescent protein-based pH-sensitive fluorescent reporter pHluorin at

their luminal domain were overexpressed by plasmid transfection (Synaptophysin-pHluorin; DIV

7) or lentiviral transduction (vGLUT1-pHluorin; DIV 2).

For following endogenous vesicular gamma-aminobutyric acid transporter (vGAT) recycling,

spontaneously active synaptic boutons were labeled by incubating cells with CypHer5E-

conjugated antibodies against the luminal domain of vGAT (1:500 from a 1 mg/ml stock; Synaptic

Systems; Cat#131,103CpH) for 2 h in their respective conditioned culture media at 37 °C and 5 %

CO_2 prior to imaging.

To investigate kinetics of SV recycling, neurons at DIV 14-16 were placed into an RC-

47FSLP stimulation chamber (Warner Instruments) in osmolarity-adjusted imaging buffer [170

mM sodium chloride (NaCl), 20 mM N-Tris(hydroxyl-methyl)-methyl-2-aminoethane-sulphonic

acid (TES), 5 mM Glucose, 5 mM sodium hydrogencabonate ($NaHCO_3$), 3.5 mM potassium

chloride (KCl), 1.3 mM $CaCl_2$, 1.2 mM sodium sulfate (Na_2SO_4), 1.2 mM magnesium chloride

($MgCl_2$), 0.4 mM potassium dihydrogenphosphate (KH_2PO_4), 50 μM *(2R)*-amino-5-

phosphonovaleric acid (AP5) and 10 μM 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX); pH 7.4]

at 37°C (Tempcontrol 37-2 digital). Cells were subjected to electrical field-stimulation (MultiStim

System-D330; Digimeter Ltd.) with annotated stimulation trains to evoke action potentials (APs)

[40 Hz, 5s (200 APs), 40 Hz, 2s (80 APs), 20 Hz, 2s (40 APs); at 100 mA]. Following changes in

fluorescence were tracked by an inverted Zeiss Axiovert 200M microscope, equipped with a 40x

oil-immersion EC Plan Neofluar objective (NA 1.30), an EM-CCD camera (Evolve Delta 512)

and a pE-300[white] LED light source (CoolLED). The scanning format was set to 512 x 512 pixels

with 16-bit sampling. eGFP (Excitation: BP470-40; Emission: BP535-50; Zeiss filter set 38) or

Cy5 (Excitation: BP640-30; Emission: BP525-50; Zeiss filter set 50) filter sets were used for

pHluorin or CypHer assays, respectively. Images were acquired at 0.5 Hz frame rate for 100 s with

50 (Syph-pHluorin) or 100 (vGLUT1-pH, vGAT-CypHer) ms exposure with an electron

multiplying gain of 250 operated through Fiji-based MicroManager 4.11 software.

Analysis of responding boutons was performed through custom written macros to identify

regions of interest (ROIs) (https://github.com/DennisVoll/pHluorin_ROI_selector/) in an

automated manner using SynActJ (Schmied *et al*, 2021). Such analysis averaged fluorescence for

each time point in an image series (video) of at least > 20 responding boutons and corrected values

for background fluorescence yielding raw background-corrected fluorescence (F). The fold

increase of fluorescence after stimulation (F_{max}) can serve as a measure for exocytic fusion and is

calculated by normalising F by the mean intensity of F before stimulation (Baseline/basal

fluorescence (F_0) = mean intensity of first 5 frames for pHluorin; first 10 frames for CypHer) for

each time point (surface normalisation = F/F_0 for pHluorin; $F_0 - F_{min}$ for CypHer). To account for

boutons with varying pHluorin expression in one image series and to compare reacidification

kinetics, the fluorescence of each time point was subtracted by the basal fluorescence (F_0) to yield

ΔF ($F-F_0$), which was then normalised by its peak value (ΔF_{max}). Resulting peak normalised curves

($\Delta F/ \Delta F_{max}$) are annotated as norm. ΔF. Endocytic decay constants (τ) were calculated by averaging

and then fitting the norm. ΔF ($\Delta F/\Delta F_{max}$) traces of all videos in one condition (N) to a mono-

exponential decay curve [$y_0 + A*e^{(-t/\tau)}$] with the constraints of $y_0 = 1$ and offset = 0 in Prism 9

(Graphpad).

CypHer-traces had to be corrected for photobleaching: The decay constant of the bleaching curve

was determined by fitting the data points of the first 10 frames (20 s, prior to stimulation) to a

mono-exponential decay curve. The corresponding value of the photobleaching curve at a given

time *t* was added to the raw fluorescence intensity measured at each corresponding time point to

correct for the loss of intensity due to bleaching.

Immunocytochemistry

For immunostainings, neuronal cultures were chemically fixed on DIV 14-16 using 4 % p-

formaldehyde (PFA) and 4 % sucrose in phosphate buffered saline (PBS) for 15 min at room

temperature (RT). For experiments in Figure 3C, G-I, cultures were stimulated with a 40 Hz train

for 5 s in imaging buffer before immediate fixation. After fixation, cells were washed thrice with

PBS and incubated with permeabilization buffer (10% normal goat serum, 0.3 % Triton X-100 in

PBS) for 30 min at RT, followed by primary antibody incubation of proteins of interest in

permeabilization buffer at indicated dilutions (see Key Resource Table) at 4°C overnight.

Subsequently, unbound antibodies were removed by three PBS washes while bound antibodies

were decorated by corresponding fluorophore-coupled secondary antibodies in permeabilization

buffer for 1h at RT. For F-Actin staining, Phalloidin-Alexa Fluor^{TM} 594 (1:1000 stock; AAT

Bioquest; Cat# ABD-23158) was added in the secondary incubation step. Finally, neurons were

washed thrice with PBS, followed by two washes with ultra-pure water. Coverslips were dried for

2 h before mounting in ProLong^{TM} Gold Antifade (Thermo Fisher Scientific; #P36934) on glass

slides (Thermo Fisher Scientific; VWR; Cat#630-1985). The slides were cured for at least 72 h at

RT before imaging.

Multicolor time-gated STED imaging

STED images were acquired from fixed samples by a HC PL APO CS2 100 x oil objective (1.40

NA) on a Leica SP8 TCS STED 3x microscope (Leica Microsystems) equipped with a pulsed

white-light excitation laser (WLL; ~ 80 ps pulse width, 80 MHz repetition rate; NKT Photonics)

and a STED laser for depletion (775 nm). The scanning format was set to 1024 x 1024 pixels, with

8-bit sampling, 4x line averaging, 4x frame accumulation and 6 x optical zoom, yielding a final

pixel size of 18.9 nm. Three-color imaging was performed by sequentially exciting following

fluorophores (excitation filter = Exf; emission filter = Emf): Atto647N (Exf: 640 nm; Emf: 650-

700 nm); Alexa Fluor^{TM} 594 (Exf: 590 nm; Emf: 600-640 nm) and Atto542 (Exf: 540 nm; Emf:

550-580 nm) operated by the Leica Application Suite X (Leica Microsystems, 2020). For

stimulated depletion of the signal, the 775 nm STED laser was applied to all emissions. Detection

of the resulting signal was time-gated by 0.3-6 ns to allow enough time for stimulated depletion

and collected by two sensitive HyD detectors at appropriate spectral regions distinct from the

STED laser wavelength. Settings in independent experiments were similar between conditions to

allow quantification of signals. Raw data obtained from three-channel time-gated STED (gSTED)

imaging were analyzed with Fiji. For analysis, only synapses oriented in the xy-plane exhibiting a

clear separation of Bassoon (presynapse) and Homer1 (postsynapse) clusters were taken into

account. To determine synaptic localisation and presynaptic levels of proteins of interest,

multicolor line profiles were measured: A line (1.0 μm length, 0.4 μm width) perpendicular to the

synaptic cleft (space between Bassoon & Homer1 cluster) was drawn (Gerth *et al.*, 2017) and

fluorescence intensity of all three channels along this line was measured using a Fiji Macro

(Macro_plot_lineprofile_multicolor; Dr. Kees Straatman, University of Leicester, UK). Resulting

profiles were aligned to the maximum Bassoon intensity, which was set to 0 nm. For localization

analysis, all three profiles were normalized to their maxima, which were set to 1. For quantification

of presynaptic protein levels, only the fractions of the non-normalized line profiles of proteins of

interest overlapping with the normalized averaged Bassoon distribution (between 151.4 and -37.8

nm; Figure 3-Supplement 1H) were integrated. Intensities were normalized to controls (DMSO,

shCTR, WT) which were set to 100.

Imaging of endogenous β-Actin levels

To visualize endogenous β-Actin levels, eGFP knockin following the ORANGE method was performed (Willems *et al.*, 2020): Neurons were transfected with vGLUT1-mCherry (Franck Polleux) and pOrange GFP-β-Actin KI (Addgene; Cat#131479) by $CaCl_2$ on DIV 2. Directly following transfection, cells were transduced with shCTR or shmDia1+3 lentiviral particles carrying BFP as a reporter to prevent crosstalk between mCherry and NLS-RFP expression. Neuronal cultures were chemically fixed on DIV 15. After fixation, endogenous GFP and overexpressed vGLUT1-mCherry intensities were enhanced by additional antibody (anti-GFP and anti-RFP; see Key Resource Table) incubation according to the immunocytochemistry protocol described above. Images of neurons were taken on the STED microscope with and without stimulated depletion (confocal) with the same settings as indicated above. For analysis, confocal images were filtered by Gaussian blur (sigma = 2 pixel) and auto-thresholded using Otsu's method. Whenever confocal vGLUT1 signal co-localized with confocal Actin clusters, the area was chosen as a ROI. Actin intensity was then measured in the ROI applied to the STED image. The intensities of β-Actin were normalized to shCTR which was set to 100.

Transmission electron microscopy

To investigate synaptic effects of F-actin manipulations at the ultrastructural level, cells on coverslips were chemically fixed with 2 % glutaraldehyde in cacodylat buffer (CDB; 0.1 M sodium cacodylat) for 1 h at RT. Coverslips were washed thrice with CDB before osmification by 1 % (w/v) osmium tetroxide (OsO_4) and 1.5 % (w/v) potassium hexacyanoferrat ($K_3Fe(CN)_6$) in CBD for 1 h at 4°C. After postfixation, cells were stained with 1 % (w/v) uranyl acetate, dehydrated by methanol gradients and finally embedded by epoxy resin (Sigma Aldrich; Cat# 45359) infiltration. After polymerization (60°C, 30 h), coverslips were removed and ultra-thin (70 nm) sections were

cut and contrasted with 2 % (w/v) uranyl acetate and 80 mM lead citrate. 8-bit Images were

obtained on a Zeiss 900 transmission electron microscope equipped with Olympus MegaViewIII

or Olympus Morada G2 digital cameras at 30000 x magnification yielding a pixel size of 1.07 nm.

Subsequently, morphometry (density of synaptic vesicles, endosome-like vacuoles, clathrin-

coated vesicles, clathrin-coated pits and non-coated invaginations) was analyzed from synaptic

profiles with clearly distinguishable active zones and adjacent synaptic vesicles in a blinded

manner.

Biochemistry

Immunoblotting

To compare protein levels between experimental conditions, immunoblotting was performed:

Protein concentrations in lysates were measured by BCA assay (Thermo Fisher Scientific;

Cat#23227) and equal protein amounts were diluted in Laemmli sample buffer [final (1 x)

concentration: 31.5 mM 2-Amino-2-(hydroxymethyl)propane-1,3-diol (Tris), 1 % (w/v) sodium

dodecyl sulfate (SDS), 10 % (v/v) glycerol, 0.001% (w/v) 3,3-Bis(3,5-dibromo-4-hydroxyphenyl)-

2,1λ^6-benzoxathiole-1,1(3H)-dione (bromophenol blue), 5 % (v/v) 2-mercaptoethanol; pH 6.8]

and denatured at 55°C for 20 min (unless indicated otherwise). Samples were resolved by SDS-

polyacrylamide gel electrophoresis (SDS-PAGE) with self-made Bis(2-hydroxyethyl)amino-

tris(hydroxymethyl)methan (BisTris; 250 mM) based 4-20 % polyacrylamide (Rotiphorese Gel

30; Carl Roth) gradient gels and run (80-120 V; 90 min) in NuPAGE MOPS SDS running buffer

(Thermo Fisher Scientific; Cat#NP000102) using Mini-PROTEAN Tetra Vertical Electrophoresis

Cells (Bio-Rad, Cat#1658004). For membrane fractionation experiments, RFP-fluorescence of

mDia1 variants in respective fractions was imaged in-gel on a ChemiDoc XRS+ (Bio-Rad)

controlled by the Image Lab software (version 6.0.1) utilizing the Alexa546 preset (605/50 Filter3;

Light Green Epi illumination). Separated proteins were wet-blotted (110 V; 90 min; 4°C) on

fluorescence-optimized polyvinylidene difluoride (PVDF) membranes (Immobilon-FL; Merck;

IPFL00010) in transfer buffer (25 mM Tris (pH 7.6), 192 mM glycine, 20% (v/v) methanol, 0.03%

(w/v) SDS). Subsequently, membranes were blocked with blocking buffer [5 % bovine serum

albumin (BSA), in Tris-buffered saline (TBS) containing 0.01% Tween 20 (TBS-T)] for 1 h at RT

and incubated (4 °C; overnight) with primary antibodies under constant agitation at indicated

dilutions (s. Key Resource Table) in blocking buffer. Membranes were washed thrice with TBS-T

and incubated with corresponding pairs of IRDye 680RD- or 800CW-conjugated secondary

antibodies in TBS-T for 1 h at RT. After three washes with TBS-T, bound antibodies were

visualized by the Odyssey Fc Imaging System (LI-COR Biosciences) controlled and analyzed by

Image Studio Lite (Version 5.2.5). For colorimetric analysis of protein levels, intensity of bands

was measured by assigning shapes of equal size to all lanes at similar heights and subtracting the

individual background for each shape. Signals were normalized to controls on the same blot. The

PageRuler Prestained (Thermo Fisher Scientific; Cat#26616) was used as a ladder to control for

protein size.

702   Immunoprecipitation from synaptosomes

To immunoprecipitate synaptic mDia1, synaptosomes (P2') were prepared as follows: One mouse

brain (p28) was homogenized (900 rpm; 12 strokes with glass-teflon homogenizer) in 7 mL of ice-

cold homogenization buffer [(320 mM sucrose, 4 mM HEPES; pH 7.4) supplemented with

mammalian protease inhibitor cocktail (PIC)] at 4°C. Large cellular debris and nuclei were

sedimented by centrifugation at 900 g for 10 min at 4°C. The supernatant was further centrifuged

at 12500 g for 15 min at 4°C. The resulting pellet (P2) was resuspended in 15 mL of

homogenization buffer and pelleted at 12500 g for 15 min at 4°C yielding the crude synaptosomal

fraction P2'. Subsequently, the pellet was resuspended in 2 mL of immunoprecipitation buffer [20

mM HEPES, 130 mM NaCl, 2 mM $MgCl_2$, 1 % (w/v)

3-[Dimethyl[3-(3α,7α,12α-trihydroxy-5β-cholan-24-amido)propyl]azaniumyl]propane-1-

sulfonate (CHAPS), PIC, phosphatase inhibitor cocktail II and III (Sigma-Aldrich); pH 7.4] and

lysed for 30 min at 4°C under light agitation. The lysate was cleared by centrifugation at 15000 g

at 4°C and protein concentration was measured using the BCA assay.

For identification of the protein environment of mDia1, P2' lysate (2 mg; 2 g/L) was

incubated with either 2 µg of anti-mDia1 antibody (BD Biosciences, Cat# 610848) or equal

amounts of immunoglobulin G (IgG) isotype control mouse antibody (Thermo Fisher Scientific;

Cat# 31903; RRID:AB_10959891) for 1 h at 4°C before the addition of 25 µL of Pierce Protein

A/G Magnetic Beads (Thermo Fisher Scientific) for an additional 2 h under constant rotation.

Subsequently, unbound proteins were removed from the beads by three consecutive wash cycles

with immunoprecipitation buffer. Bound proteins were eluted and boiled (10 min; 95°C) in 2x

Laemmli buffer, resolved by SDS-PAGE and analyzed by immunoblotting.

LC-MS analysis of protein environment of mDia1

For the analysis of interaction partners of synaptic mDia1, mDia1 was immunoprecipitated from

synaptosomes as described above. Eluted proteins were reduced (5 mM dithiothreitol; 30 min at

55°C), alkylated (15 mM iodoacetamide; 20 min at RT in the dark), and submitted to LC-MS

analysis: Proteins were subjected to SDS-PAGE following excision of 3 bands per lane and in-gel

digestion of proteins by trypsin (1:100 (w/w); overnight at 37°C). Resulting tryptic peptides were

separated by reverse-phase high-performance liquid chromatography (RP-HPLC; Ultimate™

3000 RSLCnano system; Thermo Scientific) using a 50 cm analytical column (in-house packed

with Poroshell 120 EC-C18; 2.7 µm; Agilent Technologies) with a 120 min gradient. RP-HPLC

was coupled on-line to an Orbitrap Elite mass spectrometer (Thermo Fisher Scientific) that

performed precursor ion (MS1) scans at a mass resolution of 60000, while fragment ion (MS2)

scans were acquired with an automatic gain control (AGC) target of 5×10^3 and a maximum

injection time of 50 ms. Data analysis including label free quantification was performed with

MaxQuant (version 1.6.1.0) using the following parameters: The initial maximum mass deviation

of the precursor ions was set at 4.5 parts per million (ppm), and the maximum mass deviation of

the fragment ions was set at 0.5 Da. Cysteine carbamidometyl and propionamide, Methionine

oxidation and N-terminal acetylation were set as variable modifications. For identification of

proteins, data were searched against the SwissProt murine database (Mouse_2016oktuniprot-

proteome%3AUP000000589.fasta). False discovery rates were < 1 % at the protein level based on

matches to reversed sequences in the concatenated target-decoy database. The statistical analysis

was done with Perseus (version 1.6.7.0).

Effector pulldown assays

To analyze the activity of small Rho GTPases, pulldowns utilizing effector protein domains which

exclusively bind to their GTP-bound forms were performed: Neurons (200 000 cells) were washed

with ice-cold PBS and harvested in GTPase lysis buffer (50 mM HEPES, 500 mM NaCl, 10mM

$MgCl_2$, PIC, phosphatase Inhibitor cocktails II + III; pH 7.4). Cells were lysed for 5 min under

repetitive mixing before centrifugation (15000 g; 5 min; 4°C). Cleared lysates were incubated with

Rhotekin-Rho binding domain (RBD) beads (60 µg; Cytoskeleton Inc.) or with PAK-p21 binding

domain (PBD) beads (20 µg; Cytoskeleton Inc.) at 4°C under constant rotation to bind active RhoA

or Cdc42 and Rac1, respectively. After 2 h, the beads were pelleted by centrifugation (1000 g; 1

min; 4°C) and washed with washing buffer (50 mM HEPES, 150 mM NaCl, 10 mM $MgCl_2$; pH

7.4). After repeated centrifugation, unbound proteins were discarded with the supernatant, while

bound proteins were eluted from the beads by addition of 2 x Laemmli buffer and boiling (10 min;

95°C). Finally, activity of small Rho GTPases was resolved by SDS-PAGE and analyzed by

immunoblotting input and pulldown samples.

Membrane fractionation

To characterize membrane association of the truncation mDia1 mutant, HEK293T cells were

transfected with wild-type (mDia1-WT) or mDia1 truncation mutant (mDia1-ΔN) using $CaCl_2$.

48 h after transfection cells were washed with ice-cold PBS and harvested in ice-cold resuspension

buffer (20 mM HEPES, 130 mM NaCl, PIC; pH 7.4). Cells were lysed by forcing the suspension

through 18-gauge syringes to crack the plasma membrane in between three freeze-thaw cycles in

liquid nitrogen. Nuclei were removed by centrifugation at 1000 g for 5 min at 4°C. The supernatant

(total lysate) was sedimented at 100000 g for 30 min at 4°C to yield the membrane fraction as the

pellet and the cytosolic fraction as the high-speed supernatant. The pellet was resuspended in

resuspension buffer to the same volume of the cytosolic fraction. Equal volumes of all fractions

were analyzed by SDS-PAGE and immunoblotting to allow interpretation of protein enrichment

in cytosolic or membrane fractions with respect to the total lysate.

Statistical analysis

All data in this study are presented as the mean ± standard error of the mean (SEM) and was

obtained from N independent experiments with a total sample number of n (e.g., number of images,

videos, synapses, etc.) as annotated in the Figure legends. For analysis of protein levels in STED

microscopy and synaptic structures in EM, statistical differences between groups were calculated

considering n, while in all other experiments statistical differences were calculated between

independent experiments N (In pHluorin/CypHer assays, at least 20 responding boutons/video

were analyzed). For n > 100 or N > 5, data were tested for Gaussian distribution following

D'Agostino-Pearson tests to determine parametric versus non-parametric statistical testing. The

statistical significance between two groups was evaluated with either two-tailed unpaired student's

t-tests for normally distributed data or two-tailed unpaired Mann-Whitney tests, if data did not

follow Gaussian distribution. In experiments that necessitated normalization (to 100 or 1) before

analysis, one-sample t-tests or one-sample Wilcoxon rank tests were performed for normal and

non-normal distributed data, respectively. The statistical significance between more than two

experimental groups of normally distributed data was analyzed by one-way ANOVA, followed by

a Tukey's post hoc test, while Kruskal-Wallis tests with post hoc Dunn's multiple comparison test

were used when datasets did not follow Gaussian distribution.

Corresponding statistical tests are indicated in the Figure and significance levels are

annotated as asterisks (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and ****$p < 0.0001$). Differences that

are not significant are not stated or indicated as ns ($p > 0.05$). Statistical data evaluation was

performed using GraphPad Prism 9.5.1 (733) and all calculated p-values are annotated in the

source data table. All Figures were assembled using Affinity Designer (version 1.10.6.1665).

Data availability

All data generated and analyzed in this study are included in the manuscript and supporting files.

Numerical data as well as raw images of western blots are provided as source data files. Values

from the same experiment (N) are shaded in similar colors.

Contact for Reagent and Resource Sharing

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the corresponding contact V.H. (Haucke@fmp-berlin.de).

ACKNOWLEDGEMENTS

We are indebted to Sabine Hahn, Delia Löwe and Silke Zillmann for expert technical assistance

with the preparation of neuronal cultures. We further wish to thank Dr. Martin Lehmann (FMP,

Berlin), Hannah Gelhaus and Gresy Bregu (both FU Berlin) for aid with STED imaging, Dr.

Dmytro Puchkov (FMP Imaging Core Facility) for supervising electron microscopy analysis, and

Heike Stephanowitz and Prof. Fan Liu (FMP Core Facility Proteomics) for proteomic analyses.

Supported by grants from the Deutsche Forschungsgemeinschaft (SFB958/ TP A01) to V.H..

REFERENCES

Anes E, Kuhnel MP, Bos E, Moniz-Pereira J, Habermann A, Griffiths G (2003) Selected lipids

activate phagosome actin assembly and maturation resulting in killing of pathogenic mycobacteria.

Nat Cell Biol 5: 793-802

Bingham D, Jakobs CE, Wernert F, Boroni-Rueda F, Jullien N, Schentarra EM, Friedl K, Da Costa

Moura J, van Bommel DM, Caillol G *et al* (2023) Presynapses contain distinct actin

nanostructures. *J Cell Biol* 222

Bleckert A, Photowala H, Alford S (2012) Dual pools of actin at presynaptic terminals. *J*

Neurophysiol 107: 3479-3492

Bloom O, Evergren E, Tomilin N, Kjaerulff O, Low P, Brodin L, Pieribone VA, Greengard P,

Shupliakov O (2003) Colocalization of synapsin and actin during synaptic vesicle recycling. *J Cell*

Biol 161: 737-747

Borst JG, Soria van Hoeve J (2012) The calyx of Held synapse: from model synapse to auditory

relay. *Annu Rev Physiol* 74: 199-224

Boucrot E, Ferreira AP, Almeida-Souza L, Debard S, Vallis Y, Howard G, Bertot L, Sauvonnet

N, McMahon HT (2015) Endophilin marks and controls a clathrin-independent endocytic

pathway. *Nature* 517: 460-465

Boulant S, Kural C, Zeeh JC, Ubelmann F, Kirchhausen T (2011) Actin dynamics counteract

membrane tension during clathrin-mediated endocytosis. *Nat Cell Biol* 13: 1124-1131

Chanaday NL, Cousin MA, Milosevic I, Watanabe S, Morgan JR (2019) The Synaptic Vesicle

Cycle Revisited: New Insights into the Modes and Mechanisms. *J Neurosci* 39: 8209-8216

Chanaday NL, Kavalali ET (2018) Optical detection of three modes of endocytosis at hippocampal

synapses. *Elife* 7

Chandrasekar I, Huettner JE, Turney SG, Bridgman PC (2013) Myosin II regulates activity

dependent compensatory endocytosis at central synapses. *J Neurosci* 33: 16131-16145

Chauhan BK, Lou M, Zheng Y, Lang RA (2011) Balanced Rac1 and RhoA activities regulate cell

shape and drive invagination morphogenesis in epithelia. *Proc Natl Acad Sci U S A* 108: 18289-

18294

Chen X, Wu X, Wu H, Zhang M (2020) Phase separation at the synapse. *Nat Neurosci* 23: 301-

310

Cingolani LA, Goda Y (2008) Actin in action: the interplay between the actin cytoskeleton and

synaptic efficacy. *Nat Rev Neurosci* 9: 344-356

Clayton EL, Cousin MA (2009) The molecular physiology of activity-dependent bulk endocytosis

of synaptic vesicles. *J Neurochem* 111: 901-914

Colgan LA, Yasuda R (2014) Plasticity of dendritic spines: subcompartmentalization of signaling.

Annu Rev Physiol 76: 365-385

Dani A, Huang B, Bergan J, Dulac C, Zhuang X (2010) Superresolution imaging of chemical

synapses in the brain. *Neuron* 68: 843-856

Daou P, Hasan S, Breitsprecher D, Baudelet E, Camoin L, Audebert S, Goode BL, Badache A

(2014) Essential and nonredundant roles for Diaphanous formins in cortical microtubule capture

and directed cell migration. *Mol Biol Cell* 25: 658-668

Deguchi Y, Harada M, Shinohara R, Lazarus M, Cherasse Y, Urade Y, Yamada D, Sekiguchi M,

Watanabe D, Furuyashiki T *et al* (2016) mDia and ROCK Mediate Actin-Dependent Presynaptic

Remodeling Regulating Synaptic Efficacy and Anxiety. *Cell Rep* 17: 2405-2417

Del Signore SJ, Kelley CF, Messelaar EM, Lemos T, Marchan MF, Ermanoska B, Mund M, Fai

TG, Kaksonen M, Rodal AA (2021) An autoinhibitory clamp of actin assembly constrains and

directs synaptic endocytosis. *Elife* 10

Delvendahl I, Vyleta NP, von Gersdorff H, Hallermann S (2016) Fast, Temperature-Sensitive and

Clathrin-Independent Endocytosis at Central Synapses. *Neuron* 90: 492-498

Eisenmann KM, Peng J, Wallar BJ, Alberts AS (2005) Rho GTPase-formin pairs in cytoskeletal

remodelling. *Novartis Found Symp* 269: 206-218; discussion 219-230

Engqvist-Goldstein AE, Drubin DG (2003) Actin assembly and endocytosis: from yeast to

mammals. *Annu Rev Cell Dev Biol* 19: 287-332

Ferguson SM, Brasnjo G, Hayashi M, Wolfel M, Collesi C, Giovedi S, Raimondi A, Gong LW,

Ariel P, Paradise S *et al* (2007) A selective activity-dependent requirement for dynamin 1 in

synaptic vesicle endocytosis. *Science* 316: 570-574

Ferguson SM, Raimondi A, Paradise S, Shen H, Mesaki K, Ferguson A, Destaing O, Ko G,

Takasaki J, Cremona O *et al* (2009) Coordinated actions of actin and BAR proteins upstream of

dynamin at endocytic clathrin-coated pits. *Dev Cell* 17: 811-822

Gan Q, Watanabe S (2018) Synaptic Vesicle Endocytosis in Different Model Systems. *Front Cell*

Neurosci 12: 171

Ganguly A, Tang Y, Wang L, Ladt K, Loi J, Dargent B, Leterrier C, Roy S (2015) A dynamic

formin-dependent deep F-actin network in axons. *J Cell Biol* 210: 401-417

Gerth F, Japel M, Pechstein A, Kochlamazashvili G, Lehmann M, Puchkov D, Onofri F, Benfenati

F, Nikonenko AG, Fredrich K *et al* (2017) Intersectin associates with synapsin and regulates its

nanoscale localization and function. *Proc Natl Acad Sci U S A* 114: 12057-12062

Goode BL, Eck MJ (2007) Mechanism and function of formins in the control of actin assembly.

Annu Rev Biochem 76: 593-627

Gormal R, Valmas N, Fath T, Meunier F (2017) A role for tropomyosins in activity-dependent

bulk endocytosis? *Mol Cell Neurosci* 84: 112-118

Gu C, Chang J, Shchedrina VA, Pham VA, Hartwig JH, Suphamungmee W, Lehman W, Hyman

BT, Bacskai BJ, Sever S (2014) Regulation of dynamin oligomerization in cells: the role of

dynamin-actin interactions and its GTPase activity. *Traffic* 15: 819-838

Higashi T, Ikeda T, Shirakawa R, Kondo H, Kawato M, Horiguchi M, Okuda T, Okawa K, Fukai

S, Nureki O *et al* (2008) Biochemical characterization of the Rho GTPase-regulated actin assembly

by diaphanous-related formins, mDia1 and Daam1, in platelets. *J Biol Chem* 283: 8746-8755

Higashida C, Suetsugu S, Tsuji T, Monypenny J, Narumiya S, Watanabe N (2008) G-actin

regulates rapid induction of actin nucleation by mDia1 to restore cellular actin polymers. *J Cell*

Sci 121: 3403-3412

Hodge RG, Ridley AJ (2016) Regulating Rho GTPases and their regulators. *Nature reviews*

Molecular cell biology 17: 496-510

Hori T, Eguchi K, Wang HY, Miyasaka T, Guillaud L, Taoufiq Z, Mahapatra S, Yamada H, Takei

K, Takahashi T (2022) Microtubule assembly by tau impairs endocytosis and neurotransmission

via dynamin sequestration in Alzheimer's disease synapse model. *Elife* 11

Hua Y, Sinha R, Thiel CS, Schmidt R, Huve J, Martens H, Hell SW, Egner A, Klingauf J (2011)

A readily retrievable pool of synaptic vesicles. *Nat Neurosci* 14: 833-839

Imoto Y, Raychaudhuri S, Ma Y, Fenske P, Sandoval E, Itoh K, Blumrich EM, Matsubayashi HT,

Mamer L, Zarebidaki F *et al* (2022) Dynamin is primed at endocytic sites for ultrafast endocytosis.

Neuron 110: 2815-2835 e2813

Keine C, Al-Yaari M, Radulovic T, Thomas CI, Valino Ramos P, Guerrero-Given D, Ranjan M,

Taschenberger H, Kamasawa N, Young SM, Jr. (2022) Presynaptic Rac1 controls synaptic

strength through the regulation of synaptic vesicle priming. *Elife* 11

Kessels MM, Qualmann B (2006) Syndapin oligomers interconnect the machineries for endocytic

vesicle formation and actin polymerization. *J Biol Chem* 281: 13285-13299

Kitzing TM, Sahadevan AS, Brandt DT, Knieling H, Hannemann S, Fackler OT, Grosshans J,

Grosse R (2007) Positive feedback between Dia1, LARG, and RhoA regulates cell morphology

and invasion. *Genes Dev* 21: 1478-1483

Kononenko NL, Haucke V (2015) Molecular mechanisms of presynaptic membrane retrieval and

synaptic vesicle reformation. *Neuron* 85: 484-496

Kononenko NL, Puchkov D, Classen GA, Walter AM, Pechstein A, Sawade L, Kaempf N,

Trimbuch T, Lorenz D, Rosenmund C *et al* (2014) Clathrin/AP-2 mediate synaptic vesicle

reformation from endosome-like vacuoles but are not essential for membrane retrieval at central

synapses. *Neuron* 82: 981-988

Lash LL, Wallar BJ, Turner JD, Vroegop SM, Kilkuskie RE, Kitchen-Goosen SM, Xu HE, Alberts

AS (2013) Small-molecule intramimics of formin autoinhibition: a new strategy to target the

cytoskeletal remodeling machinery in cancer cells. *Cancer Res* 73: 6793-6803

Lawson CD, Ridley AJ (2018) Rho GTPase signaling complexes in cell migration and invasion. *J

Cell Biol* 217: 447-457

Lopez-Hernandez T, Takenaka KI, Mori Y, Kongpracha P, Nagamori S, Haucke V, Takamori S

(2022) Clathrin-independent endocytic retrieval of SV proteins mediated by the clathrin adaptor

AP-2 at mammalian central synapses. *Elife* 11

Macia E, Ehrlich M, Massol R, Boucrot E, Brunner C, Kirchhausen T (2006) Dynasore, a cell-

permeable inhibitor of dynamin. *Dev Cell* 10: 839-850

Marston DJ, Dickinson S, Nobes CD (2003) Rac-dependent trans-endocytosis of ephrinBs

regulates Eph-ephrin contact repulsion. *Nat Cell Biol* 5: 879-888

McMahon HT, Boucrot E (2011) Molecular mechanism and physiological functions of clathrin-

mediated endocytosis. *Nature reviews Molecular cell biology* 12: 517-533

Merrifield CJ, Perrais D, Zenisek D (2005) Coupling between clathrin-coated-pit invagination,

cortactin recruitment, and membrane scission observed in live cells. *Cell* 121: 593-606

Miesenbock G, De Angelis DA, Rothman JE (1998) Visualizing secretion and synaptic

transmission with pH-sensitive green fluorescent proteins. *Nature* 394: 192-195

Muller PM, Rademacher J, Bagshaw RD, Wortmann C, Barth C, van Unen J, Alp KM, Giudice

G, Eccles RL, Heinrich LE *et al* (2020) Systems analysis of RhoGEF and RhoGAP regulatory

proteins reveals spatially organized RAC1 signalling from integrin adhesions. *Nat Cell Biol* 22:

498-511

O'Neil SD, Racz B, Brown WE, Gao Y, Soderblom EJ, Yasuda R, Soderling SH (2021) Action

potential-coupled Rho GTPase signaling drives presynaptic plasticity. *Elife* 10

Ogunmowo TH, Jing H, Raychaudhuri S, Kusick GF, Imoto Y, Li S, Itoh K, Ma Y, Jafri H, Dalva

MB *et al* (2023) Membrane compression by synaptic vesicle exocytosis triggers ultrafast

endocytosis. *Nat Commun* 14: 2888

Otomo T, Otomo C, Tomchick DR, Machius M, Rosen MK (2005) Structural basis of Rho

GTPase-mediated activation of the formin mDia1. *Mol Cell* 18: 273-281

Peng GE, Wilson SR, Weiner OD (2011) A pharmacological cocktail for arresting actin dynamics

in living cells. *Mol Biol Cell* 22: 3986-3994

Peng J, Kitchen SM, West RA, Sigler R, Eisenmann KM, Alberts AS (2007) Myeloproliferative

defects following targeting of the Drf1 gene encoding the mammalian diaphanous related formin

mDia1. *Cancer Res* 67: 7565-7571

Peng J, Wallar BJ, Flanders A, Swiatek PJ, Alberts AS (2003) Disruption of the Diaphanous-

related formin Drf1 gene encoding mDia1 reveals a role for Drf3 as an effector for Cdc42. *Curr*

Biol 13: 534-545

Piriya Ananda Babu L, Wang HY, Eguchi K, Guillaud L, Takahashi T (2020) Microtubule and

Actin Differentially Regulate Synaptic Vesicle Cycling to Maintain High-Frequency

Neurotransmission. *J Neurosci* 40: 131-142

Raimondi A, Ferguson SM, Lou X, Armbruster M, Paradise S, Giovedi S, Messa M, Kono N,

Takasaki J, Cappello V *et al* (2011) Overlapping role of dynamin isoforms in synaptic vesicle

endocytosis. *Neuron* 70: 1100-1114

Ramalingam N, Zhao H, Breitsprecher D, Lappalainen P, Faix J, Schleicher M (2010)

Phospholipids regulate localization and activity of mDia1 formin. *Eur J Cell Biol* 89: 723-732

Renard HF, Simunovic M, Lemiere J, Boucrot E, Garcia-Castillo MD, Arumugam S, Chambon V,

Lamaze C, Wunder C, Kenworthy AK *et al* (2015) Endophilin-A2 functions in membrane scission

in clathrin-independent endocytosis. *Nature* 517: 493-496

Richards DA, Rizzoli SO, Betz WJ (2004) Effects of wortmannin and latrunculin A on slow

endocytosis at the frog neuromuscular junction. *J Physiol* 557: 77-91

Ritter B, Ferguson SM, De Camilli P, McPherson PS (2017) A lentiviral system for efficient

knockdown of proteins in neuronal cultures [version 1; referees: 2 approved]. *MNI Open Res* 1

Roux A, Uyhazi K, Frost A, De Camilli P (2006) GTP-dependent twisting of dynamin implicates

constriction and tension in membrane fission. *Nature* 441: 528-531

Saffarian S, Cocucci E, Kirchhausen T (2009) Distinct dynamics of endocytic clathrin-coated pits

and coated plaques. *PLoS Biol* 7: e1000191

Saheki Y, De Camilli P (2012) Synaptic vesicle endocytosis. *Cold Spring Harbor perspectives in*

biology 4: a005645

Sakaba T, Kononenko NL, Bacetic J, Pechstein A, Schmoranzer J, Yao L, Barth H, Shupliakov O,

Kobler O, Aktories K *et al* (2013) Fast neurotransmitter release regulated by the endocytic scaffold

intersectin. *Proc Natl Acad Sci U S A* 110: 8266-8271

Sakaba T, Neher E (2003) Involvement of actin polymerization in vesicle recruitment at the calyx

of Held synapse. *J Neurosci* 23: 837-846

Sankaranarayanan S, Atluri PP, Ryan TA (2003) Actin has a molecular scaffolding, not propulsive,

role in presynaptic function. *Nat Neurosci* 6: 127-135

Schmied C, Soykan T, Bolz S, Haucke V, Lehmann M (2021) SynActJ: Easy-to-Use Automated

Analysis of Synaptic Activity. *Front Comp Sci-Switz* 3

Shinohara R, Thumkeo D, Kamijo H, Kaneko N, Sawamoto K, Watanabe K, Takebayashi H,

Kiyonari H, Ishizaki T, Furuyashiki T *et al* (2012) A role for mDia, a Rho-regulated actin

nucleator, in tangential migration of interneuron precursors. *Nat Neurosci* 15: 373-380, S371-372

Shupliakov O, Bloom O, Gustafsson JS, Kjaerulff O, Low P, Tomilin N, Pieribone VA, Greengard

P, Brodin L (2002) Impaired recycling of synaptic vesicles after acute perturbation of the

presynaptic actin cytoskeleton. *Proc Natl Acad Sci U S A* 99: 14476-14481

Soykan T, Kaempf N, Sakaba T, Vollweiter D, Goerdeler F, Puchkov D, Kononenko NL, Haucke

V (2017) Synaptic Vesicle Endocytosis Occurs on Multiple Timescales and Is Mediated by

Formin-Dependent Actin Assembly. *Neuron* 93: 854-866 e854

Soykan T, Maritzen T, Haucke V (2016) Modes and mechanisms of synaptic vesicle recycling.

Curr Opin Neurobiol 39: 17-23

Watanabe S, Boucrot E (2017) Fast and ultrafast endocytosis. *Curr Opin Cell Biol* 47: 64-71

Watanabe S, Rost BR, Camacho-Perez M, Davis MW, Sohl-Kielczynski B, Rosenmund C,

Jorgensen EM (2013) Ultrafast endocytosis at mouse hippocampal synapses. *Nature* 504: 242-247

Watanabe S, Trimbuch T, Camacho-Perez M, Rost BR, Brokowski B, Sohl-Kielczynski B, Felies

A, Davis MW, Rosenmund C, Jorgensen EM (2014) Clathrin regenerates synaptic vesicles from

endosomes. *Nature* 515: 228-233

Willems J, de Jong APH, Scheefhals N, Mertens E, Catsburg LAE, Poorthuis RB, de Winter F,

Verhaagen J, Meye FJ, MacGillavry HD (2020) ORANGE: A CRISPR/Cas9-based genome

editing toolbox for epitope tagging of endogenous proteins in neurons. *PLoS Biol* 18: e3000665

Wu LG, Chan CY (2022) Multiple Roles of Actin in Exo- and Endocytosis. *Front Synaptic*

Neurosci 14: 841704

Wu XS, Lee SH, Sheng J, Zhang Z, Zhao WD, Wang D, Jin Y, Charnay P, Ervasti JM, Wu LG

(2016) Actin Is Crucial for All Kinetically Distinguishable Forms of Endocytosis at Synapses.

Neuron

Wu Y, O'Toole ET, Girard M, Ritter B, Messa M, Liu X, McPherson PS, Ferguson SM, De Camilli

P (2014) A dynamin 1-, dynamin 3- and clathrin-independent pathway of synaptic vesicle

recycling mediated by bulk endocytosis. *Elife* 3: e01621

Xu Y, Moseley JB, Sagot I, Poy F, Pellman D, Goode BL, Eck MJ (2004) Crystal structures of a

Formin Homology-2 domain reveal a tethered dimer architecture. *Cell* 116: 711-723

Zhou L, McInnes J, Wierda K, Holt M, Herrmann AG, Jackson RJ, Wang YC, Swerts J, Beyens

J, Miskiewicz K *et al* (2017) Tau association with synaptic vesicles causes presynaptic

dysfunction. *Nat Commun* 8: 15295

1018

1019

1020 **FIGURES**

Figure 1

1021

Figure 1 | Actin dynamics and actin-nucleating mDia1/3 proteins facilitate SV endocytosis.

(A) Averaged normalized Synaptophysin-pHluorin (Syph-pH) fluorescence traces from

transfected hippocampal neurons stimulated with 200 APs (40 Hz, 5s) at physiological temperature

(37.5°C). Neurons were treated with 0.1% DMSO or JLY cocktail (containing 8 µM

Jasplakinolide, 5 µM Latrunculin A and 10 µM Y-27632) as indicated. Data shown represent the

mean ± SEM. N = 4 independent experiments from n_{DMSO} = 23 videos; n_{JLY} = 36 videos.

(B) Endocytic decay constants (τ) of Synaptophysin-pHluorin traces in A: τ_{DMSO} = 29.1 ± 3.4 s;

τ_{JLY} = 55.8 ± 7.2 s; $p < 0.05$, two-tailed student's t-test. Data shown represent mean ± SEM.

(C) Averaged normalized bleach-corrected vGAT-CypHer fluorescence traces from hippocampal

neurons treated with DMSO or JLY cocktail in response to 200 AP (40 Hz, 5s) stimulation. Data

shown represent the mean ± SEM. N = 4 independent experiments from n_{DMSO} = 23 videos; n_{JLY}

= 29 videos.

(D) Endocytic decay constants of vGAT-CypHer traces in C: τ_{DMSO} = 10.9 ± 0.7 s, τ_{JLY} = 24.6 ±

2.0 s; $p < 0.001$, two-tailed unpaired student's t-test. Data shown represent the mean ± SEM.

(E) Averaged normalized Synaptophysin-pHluorin fluorescence traces from hippocampal neurons

transfected with shRNA-encoding plasmids against no mammalian target (shCTR) or mDia1

(shmDia1) in response to 200 AP (40 Hz, 5s) stimulation. Neurons were co-transfected with

mDia1-mCherry (mDia1-WT) or mCherry alone (shCTR & shmDia1) to exclude artefacts from

overexpression. Data shown represent the mean ± SEM. N = 3 independent experiments from

n_{shCTR} = 28 videos, $n_{shmDia1}$ = 21 videos, $n_{shmDia1 + mDia1-WT}$ = 21 videos.

(F) Endocytic decay constants of Synaptophysin-pHluorin traces in E: τ_{shCTR} = 29.7 ± 1.9 s;

$\tau_{shmDia1}$ = 64.7 ± 3.9 s; $\tau_{shmDia1 + mDia1-WT}$ = 30.6 ± 3.7 s; $p_{shCTR\ vs\ shmDia1} < 0.001$, $p_{shmDia1\ vs\ shmDia1 +}$

$_{mDia1-WT} < 0.001$, one-way ANOVA with Tukey's post-test. Data shown represent mean ± SEM.

(G) Endocytic decay constants of averaged normalized vGLUT1-pHluorin fluorescence traces

(Figure 1–Supplement 1I) from hippocampal neurons transduced with shCTR (τ_{shCTR} = 9.1 ± 0.8

s), shmDia1 ($\tau_{shmDia1}$ = 14.3 ± 1.5 s) or shmDia1+3 ($\tau_{shmDia1+3}$ = 16.4 ± 1.3 s) in response to 40 AP

(20 Hz, 2s) stimulation ($p_{shCTR\ vs\ shmDia1} < 0.05$, $p_{shCTR\ vs\ shmDia1+3} < 0.01$, one-way ANOVA with

Tukey's post-test). Data shown represent mean ± SEM. N = 4 independent experiments from n_{shCTR}

= 17 videos; $n_{shmDia1}$ = 19 videos; $n_{shmDia1+3}$ = 18 videos.

(H) Endocytic decay constants of averaged normalized vGLUT1-pHluorin fluorescence traces

(Figure 1–Supplement 1K) of neurons transduced with lentiviral vectors encoding shCTR (τ_{shCTR}

= 13.6 ± 1.6 s), shmDia1 ($\tau_{shmDia1}$ = 22.0 ± 3.2 s) or shmDia1+3 ($\tau_{shmDia1+3}$ = 26.9 ± 3.6 s) in

response to 80 AP (40 Hz, 2s) stimulation ($p_{shCTR \, vs \, shmDia1+3}$ < 0.05, one-way ANOVA with

Tukey's post-test). Data shown represent mean ± SEM. N = 4 independent experiments from n_{shCTR}

= 12 videos, $n_{shmDia1}$ = 15 videos, $n_{shmDia1+3}$ = 18 videos.

(I) Averaged normalized bleach-corrected vGAT-CypHer fluorescence traces from hippocampal

neurons transduced with shCTR or shmDia1+3 in response to 200 AP (40 Hz, 5s) stimulation.

Data shown represent the mean ± SEM. N = 8 independent experiments from n_{shCTR} = 37 videos,

$n_{shmDia1+3}$ = 35 videos.

(J) Endocytic decay constants of vGAT-CypHer traces in I: τ_{shCTR} = 14.1 ± 1.3 s; $\tau_{shmDia1+3}$ = 27.3

± 2.6 s; p < 0.001, two-tailed unpaired student's t-test. Data shown represent the mean ± SEM.

(K) Averaged normalized vGLUT1-pHluorin fluorescence traces from transduced neurons in

response to 80 AP (40 Hz, 2s) stimulation. Cells were treated with 0.1% DMSO or 10 μM mDia

activator (IMM) in the imaging buffer. Data shown represent mean ± SEM. N = 3 independent

experiments from n_{DMSO} = 18 videos; n_{IMM} = 16 videos.

(L) Endocytic decay constants of vGLUT1-pHluorin traces in K: τ_{DMSO} = 14.9 ± 0.8 s; τ_{IMM}= 9.8

± 0.5 s; p < 0.05, two-tailed unpaired student's t-test. Data shown represent mean ± SEM.

1069

Figure 2

Figure 2 | Loss of mDia1/3 causes an activity-dependent reduction of the SV pool.

(A) Representative synaptic electron micrographs from hippocampal neurons transduced with lentiviruses encoding shCTR or shmDia1+3. Postsynapse and postsynaptic cleft are colored in green and maroon, respectively. Scale bar, 250 nm.

(B) Average number of SVs per μm^2 in boutons from shCTR (92.2 ± 2.5) and shmDia1+3 (81.4 ± 2.9; $p < 0.0001$, Mann-Whitney test) treated neurons. Data shown represent mean $\pm$ SEM. N = 3 independent experiments from $n_{shCTR} = 326$ synapses, $n_{shmDia1+3} = 321$ synapses.

(C) Average number of SVs per μm^2 in synaptic boutons from hippocampal neurons transduced

with lentiviruses encoding shmDia1+3 and treated with 0.1 % Vehicle (10 μM NaOAc; 83.2 $\pm$

2.9) or 1 μM Tetrodotoxin (TTX; 93.8 $\pm$ 3.1; $p < 0.01$, Mann-Whitney test) for 36 h before

chemical fixation. Data shown represent mean $\pm$ SEM from two independent experiments and

$n_{Vehicle}$ = 225 synapses, n_{TTX} = 221 synapses. Representative synaptic electron micrographs are

shown in Figure 2–Supplement 1A. The dotted line represents the average SV numbers/ μm^2 in

shCTR boutons treated with DMSO from the same experiments as a reference (Figure 2-

Supplement 1A,B).

(D) Representative electron micrographs of synapses in hippocampal neurons from WT or mDia1

KO mice. Postsynapse and postsynaptic cleft are colored in green and maroon, respectively. Scale

bar, 250nm.

(E) Average number of SVs per μm^2 in WT (117.6 $\pm$ 5.3) and mDia1 KO (84.6 $\pm$ 5.2; $p < 0.0001$,

Mann-Whitney test) boutons. Data shown represent the mean $\pm$ SEM from n_{WT} = 103, n_{KO}= 96

synapses (N = 1).

1092

1093

Figure 3

1094

Figure 3 | mDia1 associates with endocytic proteins and localizes to presynaptic sites.

(A) Schematic representation of functional domains of mDia1. Rho-binding domain (RBD),

Diaphanous inhibitory domain (DID), Dimerization domain (DD), Coiled coil domain (CC),

Formin homology domain 1 (FH1), Formin homology domain 2 (FH2), Diaphanous autoinhibitory

domain (DAD). The unstructured N-terminus (first 60 amino acids) contains three basic stretches

and was truncated in the ΔN mutant.

(B) Endocytic decay constants of Synaptophysin-pHluorin traces (Figure 3-Supplement 1C) from

hippocampal neurons transfected with shRNAmiR against no mammalian target (shCTRmiR) or

mDia1 (shmDia1miR) in response to 200 AP (40 Hz, 5s) stimulation. For rescue experiments,

neurons were co-transfected with plasmids encoding mDia1-WT-mCherry ($\tau_{shmDia1miR + mDia1-WT}$ =

20.0 ± 0.8 s), mDia1-ΔN-mCherry ($\tau_{shmDia1miR + mDia1-\Delta N} = 34.5 \pm 2.9$ s) or mCherry alone ($\tau_{shCTRmiR}$

$= 21.8 \pm 1.1$ s, $\tau_{shmDia1miR} = 30.4 \pm 1.9$ s) to exclude artefacts from overexpression ($p_{shCTRmiR \; vs}$

$_{shmDia1miR} < 0.05$; $p_{shmDia1miR \; vs \; shmDia1miR + mDia1-WT} < 0.01$; $p_{shmDia1miR + mDia1-WT \; vs \; shmDia1miR + mDia1-\Delta N}$

< 0.01, one-way ANOVA with Tukey's post-test). Data shown represent mean $\pm$ SEM. N = 5

independent experiments from $n_{shCTRmiR} = 41$ videos, $n_{shmDia1miR} = 51$ videos, $n_{shmDia1miR + mDia1-WT}$

$= 35$ videos, $n_{shmDia1miR + mDia1-\Delta N} = 37$ videos.

(C) Representative three-channel time-gated STED images of synapses from hippocampal cultures

treated with 0.1% DMSO or 80 μM Dynasore for 10 min before fixation and immunostained for

Bassoon (presynaptic marker, magenta), mDia1 (cyan) and Homer1 (postsynaptic marker, green).

Scale bar, 250 nm.

(D) Averaged normalized line profiles for synaptic distribution of mDia1 and Homer1 relative to

Bassoon (Maximum set to 0 nm). Data represent mean $\pm$ SEM. N = 3 independent experiments

from n = 235 synapses.

(E) Volcano plot of proteins associating with synaptic mDia1 analyzed by label-free proteomics

of anti-mDia1 versus control (CTR) immunoprecipitates from detergent-extracted mouse

synaptosomes (P2' fraction). The logarithmic ratios of protein intensities are plotted against

negative logarithmic p-values derived from two-tailed student's t-test. N = 3 independent

experiments. Each dot represents one protein/gene. Selected cytoskeletal hits include: Actin,

Myosin IIB (MyoIIB) and Rac1. Selected endocytic hits include Amphiphysin ($p < 0.05$),

Dynamin1, Endophilin-A1, PACSIN1, PACSIN2 ($p < 0.05$) and Synaptojanin1.

(F) Endogenous immunoprecipitation of mDia1 from detergent-extracted mouse synaptosomes

(P2' fraction) using mDia1-specific antibodies. Immunoprecipitates were analyzed by

immunoblotting for mDia1, Dynamin1 and β-Actin.

(G) Representative three-channel time-gated STED images of synapses from hippocampal cultures

transduced with wildtype Dynamin1 (WT) or GTPase-deficient Dynamin1 (K44A) in response to

200 AP (40 Hz, 5s) stimulation. Cells were immunostained for Bassoon (magenta), mDia1 (cyan)

and Homer1 (green). Scale bar, 250 nm.

(H) Presynaptic mDia1 levels in synapses treated with 0.1% DMSO (100 ± 7.3) or 80 μM

Dynasore (145.8 ± 9.3; $p = 0.0001$; one sample Wilcoxon test) for 10 min in response to 200 AP

(40 Hz, 5s) stimulation. Absolute line profiles of mDia1 overlapping with Bassoon (presynapse)

distribution were integrated. Data shown are normalized to DMSO (set to 100) and expressed as

mean $\pm$ SEM. N = 3 independent experiments from n_{DMSO} = 92 synapses, $n_{Dynasore}$ = 135 synapses.

(I) Presynaptic mDia1 levels in synapses from hippocampal neurons transduced with wildtype

Dynamin1 (WT; 100 ± 6.2) or GTPase-deficient Dynamin1 (K44A; 142.9 ± 8.3, $p < 0.0001$, one

sample Wilcoxon test) in response to 200 AP (40 Hz, 5s) stimulation. Line profiles of mDia1

overlapping with Bassoon distribution were integrated. Data shown are normalized to Dynamin1-

WT (set to 100) and expressed as mean $\pm$ SEM. N = 2 independent experiments from n_{WT} = 43

synapses, n_{K44A} = 51 synapses.

1143

Figure 4

1144

Figure 4 | mDia facilitates SV endocytosis by regulating presynaptic F-actin.

(A) Representative three-channel time-gated STED images of synapses from hippocampal cultures transduced with shCTR or shmDia1+3, fixed and immunostained for Bassoon (presynaptic marker, magenta), F-Actin (cyan) and Homer1 (postsynaptic marker, green). Scale bar, 250 nm.

(B) Averaged normalized line profiles for synaptic distribution of F-Actin and Homer1 relative to Bassoon (Maximum set to 0 nm). Data are expressed as mean ± SEM. N = 4 independent experiments from n = 154 synapses.

(C) Presynaptic F-Actin levels in synapses treated with 0.1% DMSO (100 ± 4.8) or 80 μM Dynasore (134.7 ± 6.8; p = 0.001, one sample Wilcoxon test) for 10 min before fixation

(Representative images in Figure 4–Supplement 1A). Cells were immunostained for Bassoon

(magenta), F-Actin (cyan), and Homer1 (green). Absolute line profiles of F-Actin overlapping

with Bassoon (presynapse) distribution were integrated. Data shown are normalized to DMSO (set

to 100) and expressed as mean ± SEM. N = 3 independent experiments from n_{DMSO} = 207 synapses,

$n_{Dynasore}$ = 211 synapses.

(D) Endocytic decay constants of vGLUT1-pHluorin traces from hippocampal neurons transduced

with lentiviral particles encoding shCTR (τ_{shCTR} = 20.7 ± 0.9 s) or shmDia1 ($\tau_{shmDia1}$ = 26.4 ± 2.0

s) in response to 200 AP (40 Hz, 5s) stimulation. For rescue experiments, neurons were co-

transduced with lentiviruses encoding mDia1-WT-SNAP ($\tau_{shmDia1 + mDia1-WT}$ = 16.1 ± 1.9 s) or

mDia1-K994A-SNAP ($\tau_{shmDia1 + mDia1-K994A}$ = 29.0 ± 1.9 s) ($p_{shmDia1\ vs\ shmDia1 + mDia1-WT}$ < 0.01;

$p_{shmDia1 + mDia1-WT\ vs\ shmDia1 + mDia1-K994A}$ < 0.001, one-way ANOVA with Tukey's post-test). Data are

expressed as mean ± SEM. N = 6 independent experiments from n_{shCTR} = 21 videos; $n_{shmDia1}$ =

21 videos, $n_{shmDia1 + mDia1-WT}$ = 16 videos, $n_{shmDia1 + mDia1-K994A}$ = 19 videos.

(E) Presynaptic F-Actin levels in synapses from hippocampal cultures transduced with shCTR

(100 ± 6.4) or shmDia1+3 (58.1 ± 2.9; p < 0.001, one sample Wilcoxon test). Line profiles of F-

Actin overlapping with Bassoon (presynapse) distribution were integrated. Data shown are

normalized to shCTR (set to 100) and expressed as mean ± SEM. N = 4 independent experiments

from n_{shCTR} = 155 synapses, $n_{shmDia1+3}$ = 158 synapses.

(F) Representative confocal and two-channel time-gated STED images of endogenous β-Actin

(cyan) in vGLUT1 (magenta) positive boutons from hippocampal neurons transduced with

lentiviruses encoding shCTR or shmDia1+3. Scale bar, 250 nm.

(G) Analysis of presynaptic endogenous β-Actin levels in vGLUT1 positive boutons from shCTR

(100 ± 6.3) and shmDia1+3 (47.7 ± 4.3; p < 0.0001 one sample Wilcoxon test) transduced neurons.

β-Actin STED mean intensity was measured using confocal vGLUT1 signal as a mask. Data

shown are normalized to shCTR (set to 100) and expressed as mean ± SEM from two independent

experiments and n_{shCTR} = 67 synapses, $n_{shmDia1+3}$ = 53 synapses.

(H) Endocytic decay constants of vGLUT1-pHluorin traces (Figure 4–Supplement 1H) for

neurons transduced with shCTR or shmDia1+3 in response to 40 AP (20 Hz, 2s) stimulation.

Neurons were pre-incubated with 0.1 % DMSO or 1 μM Jasplakinolide (Jasp) for 30 min in the

media before imaging ($\tau_{shCTR + DMSO}$ = 13.4 ± 1.0 s, $\tau_{shCTR + Jasp}$ = 15.0 ± 2.2 s, $\tau_{shmDia1+3 + DMSO}$ =

25.0 ± 2.7 s, $\tau_{shmDia1+3 + Jasp}$ = 15.6 ± 2.4 s; $p_{shCTR\ vs\ shmDia1+3}$ < 0.01; $p_{shmDia1+3 + DMSO\ vs\ shmDia1+3 + Jasp}$

< 0.05, one-way ANOVA with Tukey's post-test). Data are expressed as mean ± SEM. N = 6

independent experiments from $n_{shCTR + DMSO}$ = 32 videos, $n_{shmDia1+3 + DMSO}$ = 35 videos, $n_{shCTR + Jasp}$

= 33 videos; $n_{shmDia1+3 + Jasp}$ = 34 videos).

1188

1189

Figure 5

1190

Figure 5 | RhoA/B facilitate presynaptic endocytosis and are regulated by mDia1/3.

(A) Schematic representation of activation of mDia1 by RhoA-GTP and positive feedback loop of mDia1 on RhoA-GTP levels through GEF stimulation.

(B) Representative three-channel time-gated STED image of a synapses from hippocampal cultures, fixed and immunostained for Bassoon (magenta), RhoA (cyan) and Homer1 (green). Scale bar, 250 nm.

(C) Averaged normalized line profiles for synaptic distribution of RhoA and Homer1 relative to Bassoon (Maximum set to 0 nm). Data are expressed as mean ± SEM. N = 5 independent experiments from n = 230 synapses.

(D) Endocytic decay constants of averaged normalized Synaptophysin-pHluorin fluorescence traces (Figure 5–Supplement 1A) in response to 200 AP (40 Hz, 5s) stimulation. Neurons were transfected with the annotated combinations of plasmids encoding wild-type (WT) or dominant-negative (DN, T19N mutation) RhoA and RhoB ($\tau_{\text{RhoA-WT + RhoB-WT}} = 18.4 \pm 0.7$ s, $\tau_{\text{RhoA-WT + RhoB-}}$

$_{DN} = 16.0 \pm 1.0$ s, $\tau_{RhoA\text{-}DN + RhoB\text{-}WT} = 19.8 \pm 2.4$ s, $\tau_{RhoA\text{-}DN + RhoB\text{-}DN} = 30.1 \pm 1.0$ s; $p_{RhoA\text{-}WT + RhoB\text{-}}$

$_{WT \text{ vs } RhoA\text{-}DN + RhoB\text{-}DN} < 0.01$, one-way ANOVA with Tukey's post-test). Data shown represent

mean $\pm$ SEM. N = 3 independent experiments from $n_{RhoA\text{-}WT + RhoB\text{-}WT} = 21$ videos, $n_{RhoA\text{-}DN + RhoB\text{-}}$

$_{WT} = 31$ videos, $n_{RhoA\text{-}WT + RhoB\text{-}DN} = 23$ videos, $n_{RhoA\text{-}DN + RhoB\text{-}DN} = 22$ videos.

(E) Analysis of RhoA activity by RhoA-GTP pulldown (PD) from whole-cell lysates (input) of

mouse hippocampal neurons expressing shCTR or shmDia1+3 using immobilized Rhotekin as a

bait. Samples were analyzed by immunoblotting for mDia1, mDia3, RhoA and Tubulin using

specific antibodies. Input, 10% of material used for the pulldown. Contrast of pulldown and input

blots was seperately adjusted for visualisation purposes.

(F) Densitometric quantification of RhoA-GTP normalized to total RhoA levels (input) in lysates

from neurons transduced with shCTR or shmDia1+3 (0.7 ± 0.0, $p < 0.001$, one sample t-test) from

immunoblots exemplified in E. Values for shCTR were set to 1. Data are expressed as mean $\pm$

SEM from N = 3 independent experiments.

1217

Figure 6

1218

Figure 6 | mDia1/3-Rho and Rac1 signaling facilitates presynaptic endocytosis.

(A) Schematic of the interplay between RhoA and Rac1 signaling via GTPase activating proteins

(GAPs) common for RhoA and Rac1.

(B) Analysis of Rac1 activity by Rac1-GTP pulldown (PD) from whole-cell lysates (input) of

mouse hippocampal neurons expressing shCTR or shmDia1+3 utilizing immobilized PAK as a

bait. Samples were analyzed by immunoblotting for mDia1, mDia3, Rac1 and Tubulin using

specific antibodies. Input, 10% of material used for the pulldown. Contrast of pulldown and input

blots was seperately adjusted for visualisation purposes.

(C) Densitometric quantification of Rac1-GTP normalized to total Rac1 levels (input) in lysates

from neurons transduced with shCTR or shmDia1+3 (2.2 ± 0.2; $p < 0.05$, one sample t-test) from

immunoblots exemplified in (B). Values for shCTR were set to 1. Data are expressed as mean ±

SEM from N = 3 independent experiments.

(D) Representative three-channel time-gated STED image of a synapses from hippocampal

cultures, fixed and immunostained for Bassoon (magenta), Rac1 (cyan) and Homer1 (green). Scale

bar, 250 nm.

(E) Averaged normalized line profiles for synaptic distribution of Rac1 and Homer1 relative to

Bassoon (Maximum set to 0 nm). Data represent mean ± SEM. N = 3 independent experiments

from n = 79 synapses.

(F) Averaged normalized vGAT-CypHer fluorescence traces for neurons transduced with shCTR

or shmDia1+3 in response to 200 AP (40 Hz, 5 s) stimulation. Cells were acutely treated with

0.1 % DMSO or 10 µM Rac1 Inhibitor (EHT 1864) in the imaging buffer. Data shown represent

the mean ± SEM. N = 8 independent experiments from $n_{shCTR + DMSO} = 46$ videos, $n_{shmDia1+3 + DMSO}$

$= 45$ videos, $n_{shCTR + EHT\ 1864} = 42$ videos, $n_{shmDia1+3 + EHT\ 1864} = 43$ videos.

(G) Endocytic decay constants of vGAT-CypHer traces in F: $\tau_{shCTR + DMSO} = 14.7 \pm 0.9$ s, $\tau_{shmDia1+3}$

$_{+ DMSO} = 27.5 \pm 2.3$ s, $\tau_{shCTR + EHT\ 1864} = 30.3 \pm 6.7$ s, $\tau_{shmDia1+3 + EHT\ 1864} = 41.0 \pm 4.3$ s; $p_{shCTR + DMSO}$

$_{vs\ shmDia1+3 + DMSO} < 0.05$, $p_{shCTR + DMSO\ vs\ shmDia1+3 + EHT\ 1864} < 0.0001$, Kruskal-Wallis test with Dunn's

post-test. Data represent mean ± SEM.

(H) Endocytic decay constants of Synaptophysin-pHluorin traces (Figure 6–Supplement 1E) of

neurons transduced with shCTR ($\tau_{shCTR} = 12.0 \pm 0.7$ s) or shmDia1+3 ($\tau_{shmDia1+3} = 22.7 \pm 2.0$ s) and

transfected with constitutively active Rac1 (Rac1-CA; Q61L variant; $\tau_{shCTR + Rac1-CA} = 13.6 \pm 1.2$

s, $\tau_{shmDia1+3 + Rac1-CA} = 13.3 \pm 1.4$ s) or dominant negative Rac1 (Rac1-DN; T17N variant; $\tau_{shCTR +}$

$_{Rac1-DN} = 27.8 \pm 1.3$ s, $\tau_{shmDia1+3 + Rac1-DN} = 33.4 \pm 1.6$ s) in response to 200 AP (40 Hz, 5s) stimulation

($p_{shCTR\ vs\ shmDia1+3} < 0.01$; $p_{shCTR\ vs\ shCTR + Rac1-DN} < 0.0001$, $p_{shCTR\ vs\ shmDia1+3 + Rac1-DN} < 0.01$, $p_{shmDia1+3}$

$_{vs\ shmDia1+3 + Rac1-DN} < 0.01$, one-way ANOVA with Tukey's post-test). Data are expressed as mean

$\pm$ SEM. N = 3 independent experiments from n_{shCTR} = 12 videos, $n_{shmDia1+3}$ = 23 videos; $n_{shCTR + Rac1-CA}$

= 10 videos, $n_{shmDia1+3 + Rac1-CA}$ = 14 videos, $n_{shCTR + Rac1-DN}$ = 9 videos; $n_{shmDia1+3 + Rac1-DN}$

= 13 videos.

1256

1257

Figure 7

1258

1259   **Figure 7 | Defects in presynaptic ultrastructure induced by interference with mDia1/3-Rho**

1260   **and Rac1 signaling.**

(A) Representative synaptic electron micrographs from hippocampal neurons transduced with lentiviruses encoding shCTR or shmDia1+3 and treated with 0.1 % DMSO or 10 µM Rac1 Inhibitor (EHT 1864) for 2 h before fixation. Invaginations and ELVs are colored in blue and yellow, while postsynapse and synaptic cleft are colored in green and maroon, respectively. Scale bar, 250 nm.

(B) Average number of invaginations per µm^2 in shCTR (0.5 ± 0.1) and shmDia1+3 (0.9 ± 0.1; $p < 0.0001$, Mann-Whitney test) boutons. Data represent mean ± SEM. N = 3 independent experiments from n_{shCTR} = 326 synapses, $n_{shmDia1+3}$ = 323 synapses.

(C) Average number of ELVs per µm^2 in shCTR (1.4 ± 0.1) and shmDia1+3 (2.7 ± 0.2; $p < 0.0001$, Mann-Whitney test) boutons. Data represent mean ± SEM. N = 3 independent experiments from n_{shCTR} = 326 synapses, $n_{shmDia1+3}$ = 323 synapses.

(D) Average number of invaginations per µm^2 in shCTR and shmDia1+3 boutons treated with 0.1 % DMSO (0.8 ± 0.1 for shCTR; 1.2 ± 0.1 for shmDia1+3, $p_{shCTR + DMSO \text{ vs } shmDia1+3 + DMSO} < 0.01$) or 10 µM EHT 1864 (1.8 ± 0.1 for shCTR, $p_{shCTR + DMSO \text{ vs } shCTR + EHT 1864} < 0.0001$; 1.9 ± 0.2 for shmDia1+3, $p_{shCTR + DMSO \text{ vs } shmDia1+3 + EHT 1864} < 0.0001$, $p_{shmDia1+3 + DMSO \text{ vs } shmDia1+3 + EHT 1864} < 0.05$, Kruskal-Wallis test with Dunn's post-test) for 2 h before fixation. Data represent mean ± SEM from $n_{shCTR + DMSO}$ = 144 synapses, $n_{shmDia1+3 + DMSO}$ = 143 synapses, $n_{shCTR + EHT 1864}$ = 136 synapses, $n_{shmDia1+3 + EHT 1864}$ = 153 synapses.

(E) Average number of ELVs per µm^2 in shCTR and shmDia1+3 boutons treated with 0.1 % DMSO (2.2 ± 0.2 for shCTR; 3.3 ± 0.3 for shmDia1+3, $p_{shCTR + DMSO \text{ vs } shmDia1+3 + DMSO} < 0.05$) or 10 µM EHT 1864 (2.6 ± 0.3 for shCTR; 4.3 ± 0.4 for shmDia1+3, $p_{shCTR + DMSO \text{ vs } shmDia1+3 + EHT 1864} < 0.001$, $p_{shCTR + EHT 1864 \text{ vs } shmDia1+3 + EHT 1864} < 0.01$, Kruskal-Wallis test with Dunn's post-test) for 2 h before fixation. Data represent mean ± SEM from $n_{shCTR + DMSO}$ = 144 synapses, $n_{shmDia1+3 + DMSO}$ = 143 synapses, $n_{shCTR + EHT 1864}$ = 136 synapses, $n_{shmDia1+3 + EHT 1864}$ = 153 synapses.

1285

1286

1287

LEGENDS for SUPPLEMENTS

List of supplemental figures

Figure 1–figure supplement 1 | Role of formins and mDia1/3 in SV endocytosis.

Figure 2–figure supplement 1 | SV depletion in mDia1/3-depleted neurons is activity

dependent.

Figure 3–figure supplement 1 | mDia1 binds membranes and localizes to presynaptic

endocytic sites.

Figure 4–figure supplement 1| mDia1 regulates presynaptic actin and SV endocytosis.

Figure 5–figure supplement 1 | RhoA/B regulate SV endocytosis.

Figure 6–figure supplement 1 | Cooperative action of mDia1/3 and Rac1 pathways in

presynaptic endocytosis.

Figure 7–figure supplement 1 | mDia1/3 and Rac1 cooperatively regulate the SV cycle and

presynaptic ultrastructure.

1301

1302 **Supporting Table 1 – Oligonucleotides used in this study**

1303 **Source Data Table**

1304 **Raw uncropped immunoblot images (merged PDF)**

1305

Figure 1 S1

1306

Figure 1-Supplement 1 | Role of formins and mDia1/3 in SV endocytosis.

(A) Maxima of background-corrected Syph-pHluorin fluorescence traces (surface normalized) for

neurons treated with 0.1 % DMSO (1.7 ± 0.1) or JLY cocktail (1.8 ± 0.2) in response to 200 AP

stimulation (40 Hz, 5 s). Data represent mean ± SEM. N = 4 independent experiments from n_{DMSO}

= 23 videos; n_{JLY} = 36 videos.

(B) Minima of background-corrected vGAT-CypHer fluorescence traces (surface normalization)

for neurons treated with 0.1 % DMSO or JLY cocktail (1.0 ± 0.1) in response to 200 AP stimulation

(40 Hz, 5 s). Data represent mean ± SEM. Values for DMSO were set to 1. N = 4 independent

experiments from n_{DMSO} = 23 videos; n_{JLY} = 29 videos.

(C) Averaged normalized Syph-pH fluorescence traces from transfected hippocampal neurons

treated with 0.1% DMSO, JY or JL combinations (containing 8 μM Jasplakinolide, 5 μM

Latrunculin A and 10 μM Y-27632) stimulated with 200 APs (40 Hz, 5s). Data shown represent

the mean ± SEM. N = 4 independent experiments from n_{DMSO} = 24 videos; n_{JY} = 19 videos; n_{JL} =

21 videos.

(D) Endocytic decay constants of Syph-pHluorin traces in C: τ_{DMSO} = 19.0 ± 1.9 s; τ_{JY} = 28.5 ± 1.6

s; τ_{JL} = 18.2 ± 1.6 s; $p_{DMSO\ vs\ JY}$ < 0.01, one-way ANOVA with Tukey's post-test. Data shown

represent mean ± SEM.

(E) Averaged normalized Syph-pH fluorescence traces from transfected hippocampal neurons

treated with 0.1% DMSO or 10 μM Y-27632 following 200 AP (40 Hz, 5s) stimulation. Data

shown represent the mean ± SEM. N = 4 independent experiments from n_{DMSO} = 25 videos; n_{Y} =

18 videos.

(F) Endocytic decay constants of Syph-pHluorin traces in C: τ_{DMSO} = 17.0 ± 2.2 s; τ_{Y} = 20.2 ± 4.3

s. Data shown represent mean ± SEM.

(G) Maxima of background-corrected Syph-pHluorin fluorescence traces (surface normalized) for

neurons transfected with shCTR (1.8 ± 0.1) or shmDia1 (1.8 ± 0.1) in response to 200 AP

stimulation (40 Hz, 5 s). Data represent mean ± SEM. N = 9 independent experiments from n_{shCTR} = 49 videos; $n_{shmDia1}$ = 42 videos.

(H) Analysis of knockdown efficiency of lentiviral particles carrying shRNA against no mammalian target (shCTR) or mDia1 and mDia3 (shmDia1+3) in mouse hippocampal cultures harvested 12 days after transduction. Protein abundance of mDia1, mDia3 and Tubulin were immunoblotted with specific antibodies.

(I) Averaged normalized vGLUT1-pHluorin fluorescence traces from stimulated (40 APs; 20 Hz, 2 s) hippocampal neurons transduced with lentiviruses encoding shCTR, shmDia1 or both shmDia1 and shmDia3 combined (shmDia1+3). Data represent mean ± SEM. N = 4 independent experiments from n_{shCTR} = 17 videos, $n_{shmDia1}$ = 19 videos, $n_{shmDia1+3}$ = 18 videos. The corresponding endocytic decay constants are shown in Figure 1G.

(J) Maxima of background-corrected vGLUT1-pHluorin fluorescence traces (surface normalized) for neurons transduced with shCTR (1.9 ± 0.1) or shmDia1+3 (1.9 ± 0.1) in response to 40 AP stimulation (20 Hz, 2 s). Data represent mean ± SEM. N = 22 independent experiments from n_{shCTR} = 105 videos and $n_{shmDia1+3}$ = 128 videos.**(K)** Averaged normalized vGLUT1-pHluorin fluorescence traces for neurons transduced with shCTR, shmDia1 or shmDia1+3 in response to 80 AP stimulation (40 Hz, 2 s). Data represent mean ± SEM. N = 4 independent experiments from n_{shCTR} = 12 videos; $n_{shmDia1}$ = 15 videos; $n_{shmDia1+3}$ = 18 videos. Corresponding endocytic decay constants are shown in Figure 1H.

(L) Maxima of background-corrected vGLUT1-pHluorin fluorescence traces (surface normalized) for neurons transduced with shCTR (2.8 ± 0.2), shmDia1 (2.6 ± 0.2), or shmDia1+3 (2.4 ± 0.1) in response to 80 AP stimulation (40 Hz, 2 s). Data represent mean ± SEM. N = 4 independent experiments from n_{shCTR} = 12 videos; $n_{shmDia1}$ = 15 videos and $n_{shmDia1+3}$ = 18 videos.

(M) Minima of background-corrected vGAT-CypHer fluorescence traces (surface normalized) for neurons transduced with shCTR or shmDia1+3 (1.0 ± 0.2) in response to 200 AP stimulation (40 Hz, 5 s). Data represent mean $\pm$ SEM. Values for shCTR were set to 1. N = 11 independent experiments from n_{shCTR} = 45 videos and $n_{shmDia1+3}$ = 42 videos.

(N) Schematic representation of the activation of mDia1. Binding of RhoA-GTP to the RBD domain (green) or application of mDia1 activator (IMM) competes with the intramolecular interaction of the N-terminal DID (yellow) with the C-terminal DAD (red) domain (see Figure 3A for domain structure). The release of autoinhibition leads to dimerization of mDia formins in solution.

(O) Maxima of background-corrected vGLUT1-pHluorin fluorescence traces (surface normalized) for neurons treated with 0.1 % DMSO (1.7 ± 0.1) or mDia activator (IMM; 1.6 ± 0.1) in response to 80 AP stimulation (40 Hz, 2 s). Data represent mean $\pm$ SEM. N = 3 independent experiments from n_{DMSO} = 18 videos; n_{IMM} = 16 videos.

Figure 2 S1

1373

Figure 2-Supplement 1 | SV depletion in mDia1/3-depleted neurons is activity dependent.

(A) Representative synaptic electron micrographs from neurons transduced with shCTR or shmDia1+3 and treated with 0.1% Vehicle or 1 μM TTX for 36 h before fixation. Postsynapse and postsynaptic cleft are colored in green and maroon, respectively. Scale bar, 250 nm.

(B) Average number of SV per μm^2 in synaptic boutons of neurons transduced with shCTR and treated with 0.1 % Vehicle (A; 90.7 ± 3.1) or 1 μM TTX (B; 98.4 ± 3.3) for 36 h before chemical fixation. Data shown represent the mean ± SEM from two independent experiments and $n_{Vehicle}$ = 180, n_{TTX} = 204 synapses.

1382

1383

Figure 3 S1

1384

Figure 3-Supplement 1 | mDia1 binds membranes and localizes to presynaptic endocytic sites.

(A) Membrane levels of mDia1-WT-mCherry versus mDia1-ΔN-mCherry proteins overexpressed

in HEK293T cells. Membrane and cytosolic cellular fractions were isolated by ultracentrifugation

and analyzed by immunoblotting with specific antibodies (LAMP1) and in-gel fluorescence of

mCherry tags.

(B) Densitometric quantification of mDia1-WT versus mDia1-ΔN (0.6 ± 0.1; $p < 0.05$, one sample

t-test) membrane-associated protein levels. Data shown are normalized to mDia1-WT (set to 1)

and expressed as mean ± SEM. Representative immunoblot is shown in A. N = 5 independent

experiments.

(C) Averaged normalized Synaptophysin-pHluorin fluorescence from stimulated (200 APs, 40 Hz,

5s) hippocampal neurons transfected with shCTRmiR or shmDia1miR. For rescue experiments,

neurons were co-transfected with plasmids encoding mDia1-WT-mCherry, mDia1-ΔN-mCherry

or mCherry alone (shCTRmiR & shmDia1miR). Endocytic decay constants are shown in Figure

3B.

(D) Full volcano plot of proteins from Figure 3E associating with synaptic mDia1 analyzed by

label-free proteomics of anti-mDia1 versus CTR immunoprecipitates from detergent-extracted

mouse synaptosomes (P2' fraction). The cyan dot shows the specific enrichment of mDia1 as the

bait protein of the immunoprecipitation ($p < 0.001$, two-tailed student's t-test). N = 3 independent

experiments.

(E) Endogenous co-immunoprecipitation of Myosin IIB by mDia1 from detergent-extracted

mouse synaptosomes (P2' fraction). Samples were analyzed by immunoblotting using specific

antibodies against mDia1, Myosin IIB (MyoIIB) and β-Actin.

(F) Representative three-channel time-gated STED image of a synapse from hippocampal cultures

fixed and immunostained for Bassoon (presynaptic marker, magenta), Myosin IIB (cyan) and

Homer1 (postsynaptic marker, green). Scale bar, 250 nm.

(G) Averaged normalized line profiles for synaptic distribution of Myosin IIB and Homer1 relative

to Bassoon (Maximum set to 0 nm). Data are expressed as mean ± SEM. N = 3 independent

experiments from n = 267 synapses.

(H) Rationale for quantification of presynaptic protein levels of interest. The presynapse was

defined by the normalized Bassoon distribution (purple fraction, cutoff at the crossection with the

Homer1 profile) and corresponding absolute individual synaptic line profiles were integrated.

(I) Postsynaptic F-Actin levels in synapses treated with 0.1% DMSO (42.6 ± 3.4) or 80 μM

Dynasore (56.5 ± 3.3; $p < 0.001$, Mann-Whitney test) for 10 min before fixation from Figure 3C.

Data shown are normalized to presynaptic DMSO values from Figure 3H (set to 100) and

expressed as mean ± SEM. N = 3 independent experiments from n_{DMSO} = 92 synapses, $n_{Dynasore}$ =

135 synapses.

(J) Quantification of Bassoon and Homer1 levels in synapses treated with 0.1% DMSO (100.0 ±

4.5 for Bassoon; 100.0 ± 4.3 for Homer1) or 80 μM Dynasore (103.7 ± 4.1 for Bassoon; 98.6 ±

5.3) for 10 min before fixation. Data shown are normalized to DMSO values (set to 100) and

expressed as mean ± SEM. N = 3 independent experiments from n_{DMSO} = 132 synapses, $n_{Dynasore}$ =

128 synapses.

1426

1427

1428

1429

Figure 4 S1

1430

1431 **Figure 4-Supplement 1 | mDia1 regulates presynaptic actin and SV endocytosis.**

1432 **(A)** Representative three-channel time-gated STED images of synapses from hippocampal cultures

1433 treated with 0.1% DMSO or 80 µM Dynasore for 10 min. Cells were fixed and stained for Bassoon

1434 (magenta), F-Actin (cyan) and Homer1 (green). Scale bar, 250 nm. Corresponding analysis of

1435 presynaptic F-Actin levels is shown in Figure 4C.

(B) Representative three-channel time-gated STED images of synapses from hippocampal cultures transduced with Dynamin1-WT or Dynamin1-K44A . Cells were fixed and stained for Bassoon (magenta), F-Actin (cyan) and Homer1 (green). Scale bar, 250 nm.

(C) Presynaptic F-Actin levels in synapses from neurons transduced with Dynamin1-WT (100 ± 5.9) or Dynamin1-K44A (119.8 ± 6.2, $p < 0.01$, one sample t-test) in B. Absolute line profiles of F-Actin overlapping with Bassoon (presynapse) distribution were integrated. Data shown are normalized to WT (set to 100) and expressed as mean $\pm$ SEM. $n_{WT} = 54$ synapses, $n_{K44A} = 49$ synapses.

(D) Averaged normalized vGLUT1-pHluorin fluorescence traces for neurons transduced with shCTR or shmDia1 in response to 200 AP (40 Hz, 5s) stimulation. For rescue purposes, cells were co-transduced with mDia1-WT-SNAP or mDia1-K994A-SNAP. Data are expressed as mean $\pm$ SEM. N = 6 independent experiments from $n_{shCTR} = 21$ videos; $n_{shmDia1} = 21$ videos; $n_{shmDia1 + mDia1-WT} = 16$ videos; $n_{shmDia1 + mDia1-K994A} = 19$ videos. Corresponding endocytic decay constants are shown in Figure 4D.

(E) Postsynaptic F-Actin levels in synapses transduced with shCTR (100.0 ± 6.4) or shmDia1+3 (89.3 ± 6.4) from Figure 4A,E. Data shown are normalized to shCTR values (set to 100) and expressed as mean $\pm$ SEM. N = 3 independent experiments from $n_{shCTR} = 206$ synapses, $n_{shmDia1+3} = 135$ synapses.

(F) Quantification of Bassoon and Homer1 levels in synapses transduced with shCTR (100.0 ± 4.7 for Bassoon; 100.0 ± 4.5 for Homer1) or shmDia1+3 (101.4 ± 4.8 for Bassoon; 92.4 ± 4.0). Data shown are normalized to DMSO values (set to 100) and expressed as mean $\pm$ SEM. N = 3 independent experiments from $n_{shCTR} = 158$ synapses and $n_{shmDia1+3} = 159$ synapses.

(G) Representative STED images of endogenous β-Actin in vGLUT1 positive synapses in hippocampal neurons transduced with shCTR or shmDia1+3 and treated with 0.1 % DMSO or

1 µM Jasplakinolide for 45 min. Neurons were co-transfected with pOrange-GFP-β-Actin knockin

and vGLUT1-mCherry plasmids before fixation and immunostaining. Scale bar, 2.5 µm.

(H) Averaged normalized vGLUT1-pHluorin fluorescence traces for neurons transduced with

shCTR or shmDia1+3 in response to 40 AP (20 Hz, 2s) stimulation. Neurons were pre-incubated

with 0.1 % DMSO or 1 µM Jasplakinolide (Jasp) for 30 min in the cell media before imaging.

Data are expressed as mean ± SEM. N = 6 independent experiments from $n_{shCTR + DMSO} = 32$ videos,

$n_{shmDia1+3 + DMSO} = 35$ videos, $n_{shCTR + Jasp} = 33$ videos; $n_{shmDia1+3 + Jasp} = 34$ videos. The corresponding

endocytic decay constants are shown in Figure 4I.

1468

1469

1470

Figure 5 S1

1471

1472

Figure 5-Supplement 1 | RhoA/B regulate SV endocytosis.

(A) Averaged normalized Synaptophysin-pHluorin fluorescence traces from stimulated (200 APs;

40 Hz, 5s) hippocampal neurons transfected with plasmids encoding the indicated combinations

of WT or DN RhoA and RhoB variants. Data represent mean ± SEM. N = 3 independent

experiments from $n_{RhoA-WT + RhoB-WT}$ = 21 videos, $n_{RhoA-DN + RhoB-WT}$ = 31 videos, $n_{RhoA-WT + RhoB-DN}$

= 23 videos, $n_{RhoA-DN + RhoB-DN}$ = 22 videos. Endocytic decay constants are shown in Figure 5D.

(B) Maxima of background-corrected Syph-pHluorin fluorescence traces (surface normalized) for

neurons transfected with indicated combinations of WT or DN RhoA and RhoB variants (1.9 ±

0.2 for RhoA-WT + RhoB-WT; 1.9 ± 0.1 for RhoA-WT + RhoB-DN; 1.8 ± 0.1 for RhoA-DN +

RhoB-WT; 1.7 ± 0.1 for RhoA-DN + RhoB-DN) in response to 200 AP stimulation (40 Hz, 5 s).

Data represent mean ± SEM.

(C) Averaged normalized Synaptophysin-pHluorin fluorescence traces from stimulated (200 APs;

40 Hz, 5s) hippocampal neurons transfected with shRNA against no mammalian target (shCTR)

or against RhoA and RhoB (shRhoA+B). Data represent mean ± SEM. N = 3 independent

experiments from n_{shCTR} = 27 videos, $n_{shRhoA+B}$ = 25 videos.

(D) Maxima of background-corrected Syph-pHluorin fluorescence traces (surface normalized) for

neurons transfected with shCTR (1.6 ± 0.2) or shRhoA+B (1.6 ± 0.1) in response to 200 AP

stimulation (40 Hz, 5 s). Data represent mean ± SEM.

1491

1492

Figure 6 S1

Figure 6-Supplement 1 | Cooperative action of mDia1/3 and Rac1 pathways in presynaptic endocytosis.

(A) Analysis of Rac1 activity by Rac1-GTP pulldown (PD) from whole-cell lysates (input) of mouse hippocampal cultures upon inhibition of Rho activity utilizing immobilized PAK as bait.

Cells were treated with 0.1 % DMSO or 10 µM Rho Inhibitor (Rhosin) for 2 h before harvest.

Samples were analyzed by immunoblotting for Rac1 and Tubulin using specific antibodies. Input,

10% of material used for the pulldown. Contrast of pulldown and input blots was seperatly adjusted

for visualisation purposes.

(B) Representative three-channel time-gated STED images of synapses from hippocampal cultures

treated with 0.1 % DMSO or 10 µM Rac1 Inhibitor (EHT 1864) for 2 h. Cells were fixed and

stained for Bassoon (magenta), F-Actin (cyan) and Homer1 (green). Scale bar, 250 nm.

(C) Presynaptic F-Actin levels in synapses of neurons treated with 0.1 % DMSO (100 ± 8.5) or

10 µM Rac1 Inhibitor (EHT 1864; 58.6 ± 6.5; $p < 0.0001$, one sample Wilcoxon test) for 2 h. Line

profiles of F-Actin overlapping with Bassoon (presynapse) distribution were integrated. Data

shown are normalized to DMSO (set to 100) and expressed as mean $\pm$ SEM. $n_{DMSO} = 30$, $n_{EHT\ 1864}$

$= 46$ from two independent experiments.

(D) Minima of background-corrected vGAT-CypHer fluorescence traces (surface normalized) for

neurons treated with 0.1 % DMSO (1.0 ± 0.2 for shmDia1+3) or 10 µM Rac1 Inhibitor (EHT 1864;

0.8 ± 0.1 for shCTR; 0.8 ± 0.1 for shmDia1+3) in response to 200 AP stimulation (40 Hz, 5 s).

Data represent mean $\pm$ SEM. Values were normalised to DMSO treated shCTR (set to 1). $N = 8$

independent experiments from $n_{shCTR + DMSO} = 46$ videos, $n_{shmDia1+3 + DMSO} = 45$ videos, $n_{shCTR + EHT}$

$_{1864} = 42$ videos, $n_{shmDia1+3 + EHT\ 1864} = 43$ videos.

(E) Averaged normalized Synaptophysin-pHluorin fluorescence traces from stimulated (200 APs;

40 Hz, 5s) hippocampal neurons transduced with lentiviruses encoding shCTR or shmDia1+3 and

transfected with plasmids for expression of constitutively-active Rac1 (Rac1-CA; Q61L variant)

or dominant-negative Rac1 (Rac1-DN; T17N variant). Data represent mean $\pm$ SEM. $N = 3$

independent experiments from $n_{shCTR} = 12$ videos, $n_{shmDia1+3} = 23$ videos, $n_{shCTR + Rac1-CA} = 10$

videos, $n_{shmDia1+3 + Rac1-CA} = 14$ videos, $n_{shCTR + Rac1-DN} = 9$ videos; $n_{shmDia1+3 + Rac1-DN} = 13$ videos.

The corresponding endocytic decay constants are shown in Figure 6H.

(F) Maxima of background-corrected Synaptophysin-pHluorin fluorescence traces (surface

normalized maximum values of traces shown in E) from stimulated (200 APs; 40 Hz, 5s)

hippocampal neurons transduced with lentiviruses encoding shCTR ($F_{max}/F_0 = 1.3 \pm 0.0$) or

shmDia1+3 ($F_{max}/F_0 = 1.5 \pm 0.0$) and transfected with plasmics encoding CA ($F_{max}/F_{0\ shCTR + Rac1-}$

$_{CA}$ = 1.4 ± 0.2; $F_{max}/F_{0\ shmDia1+3 + Rac1-CA} = 1.5 \pm 0.1$) or DN versions (($F_{max}/F_{0\ shCTR + Rac1-DN} = 1.2 \pm$

0.1; $F_{max}/F_{0\ shmDia1+3 + Rac1-DN} = 1.3 \pm 0.1$) of Rac1. Data represent mean $\pm$ SEM.

(G) Densitometric quantification of Cdc42-GTP normalized to total Cdc42 levels in lysates from

shmDia1+3 neurons (2.7 ± 0.6; $p < 0.05$, one sample t-test).Values for shCTR were set to 1. Data

are expressed as mean $\pm$ SEM from N = 3 independent experiments.

H) Representative three-channel time-gated STED image of a synapses from hippocampal mouse

cultures, fixed and immunostained for Bassoon (magenta), Cdc42 (cyan) and Homer1 (green).

Scale bar, 250 nm.

(I) Averaged normalized line profiles for synaptic distribution of Cdc42 and Homer1 relative to

Bassoon (Maximum set to 0 nm). Data are expressed as mean $\pm$ SEM (N = 3; n = 96 synapses).

(J) Averaged normalized vGAT-CypHer fluorescence traces for neurons transduced with shCTR

or shmDia1+3 in response to 200 AP (40 Hz, 5s) stimulation. Cells were acutely treated with 0.1

% DMSO or 10 µM Cdc42 Inhibitor (ML141) in the imaging buffer. Data shown represent the

mean $\pm$ SEM. N = 6 independent experiments from $n_{shCTR + DMSO} = 31$ videos, $n_{shmDia1+3 + DMSO} =$

33 videos, $n_{shmDia1+3 + ML141} = 32$ videos.

(K) Endocytic decay constants of vGAT-CypHer traces in J: $\tau_{shCTR + DMSO} = 15.6 \pm 1.0$ s, $\tau_{shmDia1+3}$

$_{+ DMSO} = 28.0 \pm 3.1$ s, $\tau_{shCTR + ML141} = 17.6 \pm 1.6$ s, $\tau_{shmDia1+3 + ML141} = 33.1 \pm 7.7$ s; $p_{shCTR + DMSO\ vs}$

$_{shmDia1+3 + DMSO} < 0.01$, Kruskal-Wallis test with Dunn's post-test. Data shown represent the mean

$\pm$ SEM. N = 6 independent experiments from $n_{shCTR + DMSO}$ = 31 videos, $n_{shmDia1+3 + DMSO}$ = 33

videos, $n_{shCTR + ML141}$ = 29 videos, $n_{shmDia1+3 + ML141}$ = 32 videos.

1547

1548

1549

Figure 7-Supplement 1 | mDia1/3 and Rac1 cooperatively regulate the SV cycle and

presynaptic ultrastructure.

(A) Average number of invaginations per μm^2 in WT (0.1 ± 0.1) and mDia1 KO (0.4 ± 0.1; p <

0.01, Mann-Whitney test) boutons. Data shown represent the mean $\pm$ SEM from $n_{WT} = 103$

synapses, $n_{KO} = 96$ synapses.

(B) Average number of ELVs per μm^2 in WT (1.3 ± 0.2) and mDia1KO (3.1 ± 0.5; p < 0.001,

Mann-Whitney test) boutons. Data shown represent the mean $\pm$ SEM from $n_{WT} = 103$ synapses,

$n_{KO} = 96$ synapses.

(C) Average invagination length in shCTR and shmDia1+3 boutons treated with 0.1 % DMSO

(97.6 ± 4.5 nm for shCTR; 136.8 ± 6.0 nm for shmDia1+3, $p_{shCTR + DMSO \, vs \, shmDia1+3 + DMSO} < 0.0001$)

or 10 μM EHT 1864 (130.9 ± 4.5 nm for shCTR, $p_{shCTR + DMSO \, vs \, shCTR + EHT \, 1864} < 0.001$; $143.1 \pm$

4.9 nm for shmDia1+3, $p_{shCTR + DMSO \, vs \, shmDia1+3 + EHT \, 1864} < 0.0001$, Kruskal-Wallis test with Dunn's

post-test) for 2 h before chemical fixation. Data represent mean $\pm$ SEM from $n_{shCTR + DMSO} = 77$

invaginations, $n_{shmDia1+3 + DMSO} = 141$ invaginations, $n_{shCTR + EHT \, 1864} = 176$ invaginations, $n_{shmDia1+3}$

$_{+ EHT \, 1864} = 189$ invaginations.

(D) Average invagination width in shCTR and shmDia1+3 boutons treated with 0.1 % DMSO

(124.5 ± 5.6 nm for shCTR; 184.1 ± 6.6 nm for shmDia1+3, $p_{shCTR + DMSO \text{ vs } shmDia1+3 + DMSO} <$

0.0001) or 10 μM EHT 1864 (179.0 ± 5.8 nm for shCTR, $p_{shCTR + DMSO \text{ vs } shCTR + EHT\ 1864} < 0.0001$;

191.0 ± 5.4 nm for shmDia1+3, $p_{shCTR + DMSO \text{ vs } shmDia1+3 + EHT\ 1864} < 0.0001$, Kruskal-Wallis test with

Dunn's post-test) for 2 h before chemical fixation. Data represent mean ± SEM from $n_{shCTR + DMSO}$

$= 77$ invaginations, $n_{shmDia1+3 + DMSO} = 141$ invaginations, $n_{shCTR + EHT\ 1864} = 176$ invaginations,

$n_{shmDia1+3 + EHT\ 1864} = 189$ invaginations.

1572
1573