TITLE

Downregulation of *Let-7* miRNA promotes Tc17 differentiation and emphysema via de-

repression of RORγt

AUTHORS

Phillip A. Erice[1,2], Xinyan Huang[2,6], Matthew J. Seasock[1,2], Matthew J. Robertson[3], Hui-Ying,

Tung[4], Melissa A. Perez-Negron[2], Shivani L. Lotlikar[2], David B Corry[2,4,6], Farrah Kheradmand[4,5,6],

Antony Rodriguez[2,6*]

[1]Immunology Graduate Program, Baylor College of Medicine, Houston, TX, 77030.

[2]Department of Medicine, Immunology & Allergy Rheumatology, Baylor College of Medicine
Houston TX, 77030.

[3]Dan Duncan Comprehensive Cancer Center, Baylor College of Medicine Houston, TX, 77030.

[4]Department of Pathology and Immunology, Baylor College of Medicine Houston, TX, 77030.

[5]Department of Medicine, Section of Pulmonary and Critical Care, Baylor College of Medicine.
Houston, TX, 77030.

[6]Center for Translational Research on Inflammatory Diseases, Michael E. Debakey, Baylor
College of Medicine, Houston, TX, 77030.

[7]Current address, Department of Pulmonary and Critical Care Medicine, The First Affiliated
Hospital of Sun Yat-sen University. Guangzhou, Guangdong Province, P.R. China

Author Correspondence: antony.rodriguez@bcm.edu
23

ABSTRACT

Environmental air irritants including nanosized carbon black (nCB) can drive systemic

inflammation, promoting chronic obstructive pulmonary disease (COPD) and emphysema

development. The *let-7* family of miRNAs is associated with IL-17-driven T cell inflammation, a

canonical signature of lung inflammation. Recent evidence suggests the *let-7* family is

downregulated in patients with COPD, however, how they cause emphysema remains unclear.

Here we show that overall expression of the *let-7* miRNA clusters, *let-7b/let-7c2* and *let-7a1/let-*

7f1/let-7d, are reduced in the lungs and T cells of smokers with emphysema as well as in mice

with cigarette smoke (CS)- or nCB-elicited emphysema. We demonstrate that loss of the *let-7b/let-*

7c2-cluster in T cells predisposed mice to exaggerated CS- or nCB-elicited emphysema.

Furthermore, ablation of the *let-7b/let-7c2-cluster* enhanced CD8$^+$IL17a$^+$ T cells (Tc17) formation

in emphysema development in mice. Additionally, transgenic mice overexpressing *let-7* in T cells

were resistant to Tc17 and CD4$^+$ T cells (Th17) development when exposed to nCB.

Mechanistically, our findings reveal the master regulator of Tc17/Th17 differentiation, RAR-related

orphan receptor gamma t (RORγt), as a direct target of *let-7* miRNA in T cells. Overall, our findings

support of the *let-7*/RORγt as a braking and driving regulatory circuit in the generation of Tc17

cells, suggesting novel therapeutic approaches for tempering the augmented IL-17-mediated

response in emphysema.

INTRODUCTION

Chronic obstructive pulmonary disease (COPD) ranks as the third leading cause of

mortality and is projected to account for over a billion deaths by the end of the twenty-first century

(GBD Chronic Respiratory Disease Collaborators, 2020; *Findings from the Global Burden of*

Disease Study 2017, 2019; Laniado-Laborín, 2009). Currently, there are no treatment options to

reverse emphysema, the most clinically significant variant of COPD, which often is progressive

despite smoking cessation (Bhavani et al., 2015; Anthonisen et al., 2002).

Inhalation of fine particulate matter smaller than 2.5 microns (PM$_{2.5}$) found in outdoor and

indoor air pollution as well as tobacco smoke are risk factors for COPD development (Adeloye et

al., 2022; Eisner et al., 2010; Hu et al., 2010). We have previously shown that nano-sized carbon

black (nCB), a noxious chemical constituent of $PM_{2.5}$ found in the lungs of smokers, activates

macrophages and dendritic cells orchestrating a pathogenic T cell-dependent inflammatory

response and emphysema in mice (Lu et al., 2015; You et al., 2015; Shan et al., 2009; C.-Y.

Chang et al., 2022).

Research over the last decade has pointed to the importance of dysfunctional

inflammatory T cells in human COPD lung tissue and animal models of emphysema (Grumelli et

al., 2004; Xu et al., 2012; Williams et al., 2021). Aberrant T cells are implicated in impaired host

defense, exaggerated inflammation, and loss of self-tolerance in COPD (Williams et al., 2021;

Chen et al., 2023; Hogg et al., 2004; Maeno et al., 2007; Xu et al., 2012). In this regard, we and

others have demonstrated the role and pathogenicity of activated IFN-γ and IL-17-secreting

subsets of CD4[+] and CD8[+] T lymphocytes including Th1, Th17, and Tc1 cells in clinical isolates

and in mice with COPD (Lu et al., 2015; You et al., 2015; Shan et al., 2009; S.-H. Lee et al., 2007;

Kheradmand et al., 2023). The IL-17-secreting Th17 cells are particularly important as they

promote the destruction of lung epithelium and recruitment of macrophages and neutrophils which

then release proteolytic enzymes such as matrix metalloproteinases (MMPs) involved in the

degradation of the lung structural matrix (Barnes, 2016; Hoenderdos & Condliffe, 2013). We

previously demonstrated that intranasal inhalation of nCB in mice is sufficient to induce

emphysema by stimulating T cell activation by dendritic cells and macrophages in mice. Moreover,

we found that genetic ablation of IL-17a can attenuate nCB- or cigarette smoke-induced alveolar

destruction and airway inflammation (Shan et al., 2012; You et al., 2015). More recently, IL-17A

and IL-17F secreting CD8[+] T cell (Tc17) subpopulation has been shown to play a critical role in

the pathogenesis of several autoimmune and inflammatory disorders (Globig et al., 2022; Huber

et al., 2013; Srenathan et al., 2016).

Both Th17 and Tc17, require the fate-deterministic transcription factor RAR-related orphan

receptor gamma t (RORγt, encoded by *Rorc*) for differentiation and production of IL-17a (Ivanov

et al., 2007). RORγt is the best-studied positive transcriptional regulator of IL-17a and IL-17f

(Ivanov et al., 2006). In accordance with the importance of IL-17a transcription, RORγt expression

has also been reported to be elevated in COPD patients and in mouse models of COPD (Chu et

al., 2011; Li et al., 2015). However, the upstream pathophysiologic mechanisms that contribute to

the induction of RORγt and differentiation of Tc17 cells in COPD have not been well elucidated.

We previously reported that *miR-22* inhibits HDAC4, promoting antigen-presenting cell

activation (APC) in the lungs and inducing Th17-mediated emphysema in response to CS or nCB

in mice (Lu et al., 2015). Additional miRNAs that control APC and/or T cell driven IL-17a$^+$

inflammation have been identified by others including the *let-7* miRNA family (Mai et al., 2012;

Angelou et al., 2020). MicroRNA expression-based studies have shown frequent downregulation

of members of the *let-7* miRNA family, including *let-7a, let-7b, let-7c*, *let-7d*, l*et-7e*, and *let-7f* in

human emphysematous lung tissue and in murine models of emphysema, but the mechanism(s)

of action remain ill-defined (Christenson et al., 2013; Pottelberge et al., 2011; Conickx et al., 2017;

Izzotti et al., 2009). *Let-7* microRNA genes are encoded across eight loci either as single genes

or as polycistronic clusters which have confounded their analysis *in vivo* (Rodriguez et al., 2004).

Previous studies used *Lin28b* transgenic overexpression in T cells to block the maturation and

processing of *the let-7* miRNA family. They showed an inhibitory role of *let-7* family in Th17-driven

response in murine model of experimental autoimmune encephalomyelitis (EAE) (Angelou et al.,

2020).

Here we found that *let-7* miRNA, notably the *let-7a3/let-7b* and *let-7a1/let-7f1/let-7d*

clusters, are suppressed in the T cells isolated from lungs of emphysema patients. Consistently,

the analogous murine *let-7b/let-7c2-* (*let-7bc2*) and *let-7a1/f1/d1-* (*let-7afd*) clusters were similarly

downregulated in pre-clinical emphysema models. We engineered mouse models with the

specific loss-of-function (LOF) mutations of the *let-7bc2* or *let-7afd* clusters (*let-7bc2^{LOF}*and *let-*

7afd^{LOF}, respectively) in T cells as well as an inducible *let-7g* gain-of-function (GOF) (*let-7^{GOF}*)

model to determine the T cell-intrinsic role of *let-7* miRNA in emphysema pathogenesis. Deletion

of *let-7* miRNA in T cells worsened alveolar damage elicited by inhalation of CS or nCB, and

increased infiltration of immune cells in the airways, including IL-17-producing CD8$^+$ T (Tc17) cells.

Mechanistically, we found that *let-7* controls type 17 differentiation by directly targeting the

lineage-determining transcription factor, RORγt. In support of this conclusion, *let-7^{GOF}* mice were

resistant to nCB-mediated induction of RORγt and Tc17 responses. Thus, we show a previously

unappreciated role for *let-7* miRNA as a repressor of RORγt and a molecular brake to the IL-17a-

mediated T cell inflammation in emphysema.

RESULTS

The *let-7bc2-* and *let-7afd-clusters* are downregulated in lungs and T cells in COPD.

To explore the involvement of *let-7* in emphysema, we scrutinized the genomic locations and

transcriptional annotation of *let-7* members frequently downregulated in lung T cells isolated from

smoker's lungs as well as mouse models of emphysema. This combined approach showed close

linkage and high conservation of two *Let-7* clusters encoded from long intergenic non-coding RNA

(linc)-like precursors in humans and mice (Figure 1A). To shed light on whether these *Let-7*

clusters are downregulated in patients with COPD, we analyzed a published (GSE57148) lung

RNA-seq dataset obtained from COPD (N=98 and control (N=91) subjects (Kim et al., 2015). Our

analysis identified significant downregulation of the *Mirlet7ahg* and *Mirlet7bhg* gene cluster

transcripts in COPD compared to control subjects (Figure 1B). We carried out quantitative PCR

(qPCR) detection of *Let-7a,* which is encoded by both clusters, in lung tissue samples of smokers

with emphysema and non-emphysema controls, detecting significant downregulation of *Let-7a* in

emphysema samples relative to controls (Figure 1C). Because *Let-7* has been shown to

participate in IL-17$^+$ T cell responses (Angelou et al., 2020; Guan et al., 2013; Newcomb et al.,

2015), we next sought to determine if the expression pattern of *Mirlet7ahg* and *Mirlet7bhg*-derived

Let-7 members are impaired in purified CD4$^+$ T cells from emphysematous lungs. In support of

our original hypothesis, the CD4$^+$ T cell expression of *Let-7a*, *Let-7b*, *Let-7d*, and *Let-7f* were all

inversely correlated with more severe emphysema distribution in the lungs as determined by CT

scan (Figure 1D).

Next, we elucidated *let-7a1/let-7f1/let-7d-* and *let-7b/let-7c2*-clusters expression (herein

referred to as *let-7afd* and *let-7bc2* respectively) in murine models of CS- or nCB-induced

emphysema respectively (Figure 1E). Paralleling our observations in human COPD and

emphysema, mice with CS- or nCB-induced emphysema exhibited reduced expression levels of

pri-let7afd and *pri-let7bc2* transcripts in the lung and from isolated lung CD4$^+$ and CD8$^+$ T cells

(Figure 1F,G,H). Collectively, our expression results indicate suppression of *let-7afd* and *let-7bc2*-

clusters in the lung and T cells in human and pre-clinical models of emphysema.

Conditional deletion of the *let7bc2*-cluster in T cells enhances nCB- or CS-induced

emphysema.

To investigate the *in vivo* requirement of the *let-7bc2-cluster* within T cells, we generated

conditional ready floxed mice (*let-7bc2$^{flox/flox}$*). We then crossed *let7-bc2$^{flox/flox}$* mice with *CD4-Cre*

mice to generate *let-7bc2$^{flox/flox}$*; *CD4-Cre* LOF mice (denoted as *let-7bc2^{LOF}* mice hereafter)

(Figure 2A). This approach allowed us to conditionally delete the *let-7bc2* cluster in all T cells

derived from the CD4$^+$CD8$^+$ double-positive (DP) stage (P. Lee et al., 2001; Shi & Petrie, 2012).

We confirmed that *let-7bc2^{LOF}* mice exhibit robust conditional deletion of the *let-7bc2* cluster in

DP thymocytes and peripheral CD8$^+$ T and CD4$^+$ T cells (Figure 2B and data not shown). Our *let-*

7bc2^{LOF} adult mice were born at the expected Mendelian frequency and did not show any overt

histopathologic or inflammatory changes in lungs histopathology up to 1 year of age in comparison

to *let-7bc2^{f/f}* control mice (Figure 1-figure supplement 1A,B,C,D and data not shown).

Furthermore, quantification of major immune populations and T cell subsets by flow cytometry in

let-7bc2^{LOF} were comparable to control mice under baseline conditions and with moderate aging

(Figure 1-figure supplement 1A,B,C,D).

We next exposed *let-7bc2^{LOF}* and *let-7bc2^{f/f}* control mice to nCB or CS and examined the

lungs under the context of experimental emphysema. Histomorphometry measurements of mean

linear intercept (MLI) from hematoxylin and eosin (H&E)-stained sections revealed that the

enlargement of alveolar spaces sustained from either nCB- or CS-exposure was exaggerated in

let-7bc2^{LOF} mice relative to controls (Figure 2C-E). Chronic inflammation in emphysema is

characterized by the recruitment of macrophages and neutrophils to the lung tissue and airways

(Peleman et al., 1999; Senior & Anthonisen, 1998). Internally consistent with MLI measurements,

let-7bc2^{LOF} mice treated with nCB showed significantly increased airway infiltration of

macrophages and neutrophils in BAL fluid as compared to wild-type control animals (Figure 2F).

Concomitant with these findings, expression levels of *Mmp9* and *Mmp12*, which are secreted by

macrophages and neutrophils to degrade elastin and mediate alveolar damage, were elevated in

airways of *let-7bc2^{LOF}* mice exposed to CS versus controls (Figure 2G). Collectively, our data

suggests that the *let-7bc2*-cluster within T cells protects by dampening airway destruction and

inflammation because the absence of this cluster worsens the severity of experimental

emphysema in mice.

The *let-7bc2* miRNA cluster negatively regulates T$_C$17 inflammation in emphysema.

We sought to identify the T cell-intrinsic mechanisms that underlie the exaggerated

inflammation observed in emphysematous *let7bc2^{LOF}* mice. We focused on the IL-17-mediated T

cell response because it promotes neutrophil and macrophage recruitment in the lungs (Beringer

et al., 2016; Veldhoen, 2017; Shan et al., 2012). Previously, we established the induction of

CD4$^+$IL17$^+$ (Th17) cells along with CD4$^+$IFNg$^+$ (Th1) cells in mice with chronic nCB exposure (You

et al., 2015), however whether nCB similarly induces CD8$^+$IL17a$^+$ T cells (Tc17) or cytotoxic T

cells (Tc1) had not been studied. The flow cytometric profiling of lung T cells revealed enriched

proportions and counts of Tc1 and Tc17 cells in mice with nCB-emphysema and we confirmed the

induction of Th17 and Th1 cells (Figure 3A-B control PBS and control nCB). These findings

suggests that nCB elicits both the type 17 and type 1 T cell responses, consistent with CS and

elastase pre-clinical models of emphysema (Zhang et al., 2019).

We next interrogated the regulatory role of the *let-7bc2*-cluster in the type 17 and type 1

responses generated from exposure to nCB. Interestingly, *let-7bc2^{LOF}* mice showed increased

CD8$^+$IL17a$^+$ Tc17 cells relative to nCB control animals. In contrast, CD8$^+$IFN$^+$ and GZMA$^+$ Tc1

populations remained unperturbed with absence of the *let-7bc2* cluster, suggestive of a more

refined regulatory role on Tc17 differentiation (Figure 3A-B). There were no significant differences

in either Th1 or Th17 cells when comparing nCB-treated *let-7bc2^{LOF}* to wild-type controls,

indicating the *let-7bc2* cluster was dispensable for their generation (Figure 3C). Regulatory T cells

form a dynamic axis with Tc17/Th17 cells and act as a counterbalance to lung inflammation in

emphysema (Duan et al., 2016; Jin et al., 2014). Therefore, we examined whether Tc17 cell

alterations were driven by the *let-7bc2* cluster acting on regulatory T cells (Tregs). The *let-7bc2^{LOF}*

mice showed no significant difference in Tregs subset relative to controls in our model (Figure

3D). Together, our data support the notion that deletion of the *let-7bc2*-cluster is insufficient to

provoke Tc17 cell generation under homeostatic conditions. However, under the context of chronic

inflammation in emphysema, the loss of *let-7bc2*-cluster is intrinsic for the potentiation of T cells

towards Tc17 differentiation.

The *let-7* family directly inhibits RORγt expression governing Tc17 differentiation in

emphysema.

We utilized the TargetScan predictive algorithm to identify putative *let-7* microRNA targets

that are known to control the IL-17-mediated T cell response (Agarwal et al., 2015). This analysis

revealed that the 3'UTR region of *Rorc*, encoding RORγt, contains an evolutionarily conserved

and complementary motif for the *let-7* miRNA family (Figure 4A). Thus, we examined if *let7bc2-*

cluster loss in T cells would stimulate and enhance RORγt. Initially, we carried out flow cytometric

quantification for RORγt in thymocyte, splenic, and lung T cells of naïve control and *let7bc2^{LOF}*

mice up to 6-months of age. Our interrogation of RORγt mean fluorescent intensity (MFI) by flow

cytometry showed induction of RORγt in single-positive CD8$^+$ and CD4$^+$ thymocytes, as well as

peripheral splenic CD8$^+$ and CD4$^+$ T cells (Figure 4B). However, RORγt levels appeared

unchanged in purified lung CD8$^+$ T cells and CD4$^+$ T cells of naive *let-7bc2^{LOF}* mice, alluding to a

compensatory effect in homeostatic lung T cells (Figure 4B). Since we and others have shown

that miRNAs are frequently associated with stress-dependent phenotypes, we posited that

emphysematous *let-7bc2^{LOF}* T cells are poised towards induction of RORγt and production of

IL17$^+$ subsets after challenge with nCB. Indeed, nCB-emphysematous *let-7bc^{LOF}* mice exhibited

enhanced RORγt protein levels in both CD8$^+$ and CD4$^+$ T cells relative to control mice with

emphysema (Figure 4C).

Because we had found that the *let-7afd*-cluster is downregulated in T cells isolated from

COPD lungs in human and mice, and that the *let-7* family operates with some functional

redundancy, we generated mice with conditional deletion of the *let7afd*-cluster in T cells (*let-7afd$^{f/f}$;*

CD4-Cre). The *let-7afd$^{f/f}$; CD4-Cre* (*let7afd^{LOF}*) mice aged up to 6-months did not exhibit lung

histopathology nor inflammatory changes (data not shown). Of particular interest, ablation of the

let-7afd-cluster enhanced levels of RORγt in thymic and peripheral T cells of mice (Figure 4D).

Overall, this indicates that independent *let-7* clusters restrain RORγt expression levels from

thymic development to peripheral T cells under homeostatic conditions. Next, we determined

whether loss of *let-7afd*-cluster in T cells likewise sensitizes mice towards induction of RORγt in

nCB-emphysema. Intranasal administration of nCB provoked increased RORγt expression in lung

T cells of *let-7afd^{LOF}* mice as compared to *let-7afd$^{f/f}$* control mice (Figure 4E), supporting

overlapping functionality between the *let-7bc2* and *let-7afd*-clusters in repression of RORγt within

T cells.

To confirm that the *let-7* family negatively regulates Tc17 cell differentiation, at least in part,

cell autonomously in CD8$^+$ T cells, we purified naïve CD8$^+$ T cells from *let-7bc2^{LOF}* and control

mice spleens and cultured these cells *in vitro* in the presence of Tc17 polarizing (TGFβ, IL-6, anti-

IFNγ, IL-23, and IL-1β) or Tc1 polarizing (IL-2) conditions (Flores-Santibáñez et al., 2018). Our

flow cytometric analysis confirmed the enhanced commitment of *let-7*-cluster deficient CD8$^+$ T

cells towards Tc17 cells and IL-17a$^+$ production relative to control CD8$^+$ T cells (Figure 5A-B).

Moreover, enhanced Tc17 cell differentiation mirrored the increased IL-17a detected in the

supernatant from *in vitro* polarized cells as quantified by ELISA (Figure 5C). Parallel assessment

of Tc1 differentiation did not detect a difference in CD8$^+$IFNγ$^+$ cells (Figure 5A and Figure 5D).

Altogether, these data recapitulated our *in vivo* findings that the *let-7bc2* cluster negatively

regulates Tc17 response but is dispensable in Tc1 cells. Finally, to determine whether Tc17

differentiation is likewise controlled by the *let-7afd* cluster, we cultured naive CD8$^+$ splenocytes

from *let-7afd^{LOF}* and controls under Tc17 conditions. As we had observed with *let-7bc2^{LOF}*,

absence of the *let-7afd* cluster in T cells further enhanced differentiation towards Tc17 cells as

quantified by flow cytometry and ELISA (Figure 5E-F).

Next, we focused on *Rorc* as a potential direct target of *let-7*, which could mechanistically

mediate enhanced Tc17 differentiation in *let7bc2^{LOF}* mice. Towards this objective, tested whether

let-7bc2^{LOF} naïve CD8$^+$ T cells show elevated RORγt expression under either Tc0 or Tc17

differentiation conditions. In agreement with enhanced Tc17 differentiation, RORγt expression

was differentially and significantly upregulated under both Tc0 and Tc17 differentiation conditions

in *let-7bc2^{LOF}* cells relative to controls (Figure 5G). To determine whether *let-7* directly represses

Rorc mRNA levels we cloned the 3'UTR of *Rorc* into luciferase constructs. These reporter assays

with *let-7b* expressing cells independently confirmed that *let-7b* represses *Rorc* (Figure 5H, left).

Furthermore, deletion of the putative *let-7* binding sequence (Figure 4A) abrogated repression

by *let-7b* (Figure 5H, right), thus confirming *Rorc* as a functional target of *let-7* miRNA. Overall,

these *in vitro* experiments readily recapitulated an upstream regulatory role for *let-7* in Tc17

differentiation, mediated in part, via direct suppression of RORγt.

Enforced expression of *let-7* tempers RORγt T cell expression levels in experimental

emphysema*.*

To explore a potential protective role of *let-7* miRNA in experimentally induced

emphysema, we generated mice which allowed for selective induction of *let-7* activity in T cells

using the published *rtTA-iLet7* mice (Zhu et al., 2011; Belteki et al., 2005). We bred the *rtTA-iLet7*

mice to *CD4-Cre* (here in referred to as *let-7^{GOF}*) to allow Cre-loxP/doxycycline dependent *let-7g*

overexpression in thymic DP-derived T cells (Figure 6A). Steady-state *let-7^{GOF}* and control (*rtTA-*

iLet7) mice were examined for compromised RORγt protein levels within thymocytes and

peripheral T cells. Providing further evidence of *let-7*-dependent regulation of *Rorc*, protein levels

of RORγt were suppressed in CD8$^+$ and CD4$^+$ T cells of *let-7^{GOF}* mice relative to controls (Figure

6B). To determine whether enforced expression of *let-7* offered protection from experimental

emphysema, *let-7^{GOF}* and control mice were treated with nCB and then examined for changes in

lung pathology and T cell type 17 responses. The *let-7^{GOF}* mice did not exhibit any signs of lung

inflammation or pathologic remodeling at baseline (Figure 6C-D and data not shown)

Histopathologic analysis revealed a comparable degree of lung alveolar distension via

morphometric measurements of MLI in nCB-treated *let-7^{GOF}* mice versus controls suggesting that

enforced *let-7* expression is insufficient to protect the lung from emphysema (Figure 6C-D). On

the other hand, evaluation of IL17$^+$ response and RORγt levels in emphysematous lung T cells

demonstrated that, in contrast to control nCB-treated mice, *let-7^{GOF}* mice exhibited dampened

lung Tc17 and Th17 cell populations and were resistant to the induction of RORγt after nCB-

exposure (Figure 6F-6E)*.* Taken together, our *let-7* LOF and GOF models demonstrate the

necessity and sufficiency of *let-7* miRNA to act as a molecular brake to the type 17 T cell response

through the direct regulation of RORγt, further our data suggests that nCB- or CS-mediated

suppression of this braking mechanism furthers inflammation and exacerbates emphysema

severity (Figure 6G).

DISCUSSION

MiRNA expression-based studies of COPD patients and mice exposed to CS have

reported downregulation of *let-7* miRNA expression in lung tissues (Conickx et al., 2017;

Christenson et al., 2013b; Schembri et al., 2009). We and others explored the consequence of

loss of *let-7* expression/activity with synthetic oligonucleotides, sponges, lentiviral antisense

knockdown, or via ectopic delivery of Lin28b (Polikepahad et al., 2010; Viswanathan et al., 2008;

Piskounova et al., 2011), but studies pinpointing the role of individual *let-7* clusters as potential

drivers of lung inflammation and COPD within T cells remained elusive. In the present study, we

established that the *let-7* miRNA family members encoded by the *let-7bc2-* and *let-7afd-*clusters

are downregulated in T cells from lungs of emphysema patients and emphysematous mice that

were exposed to CS or nCB. Correspondingly, we demonstrated that *in vivo* genetic ablation of

let-7bc2-cluster further sensitized mice to lung tissue destruction and emphysema upon treatment

with nCB or CS. Mechanistically, our studies suggests that *let-7* miRNA prevents the emergence

of CD8$^+$ T cell differentiation into Tc17 cells during emphysema in part, by directly silencing of

Rorc.

Tc17 cells are vital for defense against viral, fungal, and bacterial infections and they have

also been associated with inflammation in various human diseases such as multiple sclerosis,

inflammatory bowel disease, and cancer (Huber et al., 2013; Globig et al., 2022; Corgnac et al.,

2020). In accordance with the potential pathogenic role of Tc17 cells as drivers of COPD, several

studies detected increased cell numbers in airways and tissues of COPD patients as well as lungs

of smoke-exposed animal models (Chang et al., 2011; Zhou et al., 2020; Duan et al., 2013). Other

researchers also detected increased Tc17 subpopulations in tissues of COPD patients with

infectious microbial exacerbations. In our earlier work to define the adaptive T cell immune

responses in nCB induced COPD, we predominantly focused on the pathogenic role of Th17 cells,

but did not examine Tc17 cells (You et al., 2015). Here we expand upon our prior observations,

revealing that chronic exposure to nCB and elicitation of emphysema mice orchestrates the

emergence and accumulation of Tc17 cells which may act in parallel with Th17 cells to promote

tissue damage.

Although RORγt has been the subject of intense scrutiny and is subject to extensive

transcriptional and post-transcriptional regulatory control, to our knowledge this is the first

reported demonstration of miRNA-mediated gene silencing to RORγt. Our data also showed that

in vivo conditional genetic ablation of individual *let-7* clusters in T cells stimulates a rise in RORγt

protein expression in single-positive thymocytes and peripheral CD8$^+$ and CD4$^+$ T cells while

enforced *let-7* activity leads to partial repression of RORγt in T cells. Despite these alterations in

RORγt expression in our *let-7* T cell LOF and GOF mice, the mice did not exhibit spontaneous

gross phenotypes in thymus, spleens, or lungs at baseline. This may be due to the subtle and

modest expression thresholding of RORγt detected in mice and/or residual *let-7* expression in T

cells. On the other hand, and in agreement with our Tc17 and experimental emphysema data, we

observed enhanced RORγt expression and IL-17a$^+$CD8$^+$ T cells in lungs of *let-7* LOF mice after

treatment with nCB. We corroborated the importance of *let-7* activity in Tc17 differentiation of *ex*

vivo cultured CD8$^+$ T cells, as well as in the direct posttranscriptional control of RORγt, suggesting

that this defect, is in part, direct and cell autonomous. Interestingly, even though RORγt was

elevated in lung CD4$^+$ T cells to a similar extent as CD8$^+$ T cells, the enhanced IL17$^+$ responses

were limited to Tc17 subpopulation in the *let-7bc2*^{LOF} mice. Although, the *let-7afd*^{LOF} mice also

showed baseline induction of RORγt relative to controls in lung T cells the mice did not exhibit

changes in Tc17/Th17 subpopulations (data not shown). Nonetheless, it seems likely that under

different cellular contexts, the functions of *let-7*-clusters do not fully overlap in association with

differential thresholding of mRNA targets. Indeed, in B cells, the *let-7afd*-cluster is essential for an

antigen-specific antibody response, whereas the *let-7bc2*-cluster appears dispensable (Jiang et

al., 2018), iterating that differential physiological role of *let-7* clusters in the immune system and

validating an individual cluster approach in dissemination of mechanisms of *let-7* in T cells. RORγt

is a defining transcription factor of the IL-17-secreting subset of immune cells, which also includes

γδ17 T cells, NKT17 and type 3 innate lymphoid cells, populations that contribute to COPD

pathology (Yanagisawa et al., 2017). A limitation of our study is that we did not examine whether

these populations were impacted by nCB-emphysema and/or in the context of *let-7* LOF.

Published studies revealed a protective role of *let-7* family against Th17-driven pathogenic

response in EAE attributed in part to direct regulation of *IL-1 receptor 1* and *IL-23 receptor*

(Angelou et al., 2020). Lending some support to these observations, we detected subtle transcript

level induction of *Il23r* in *let-7bc2*^{LOF} *in vitro* polarized Tc17 cells (data not shown). However, prior

publications on the role of *let-7* in T cells made use of *in vivo Lin28b* transgenic overexpression

in immune cells to block maturation and activity of the entire *let-7* miRNA family. Furthermore,

Lin28b was recently reported to influence transcriptome-wide ribosome occupancy and global

miRNA biogenesis (Tan et al., 2019) which could account for differences in scope of *let-7* targets

in those studies and ours. Nonetheless, further studies will be required to ascertain whether other

targets of *let-7* beyond RORγt synergistically potentiate the *in vivo* Tc17-response and
emphysema phenotype in *let7-bc2^{LOF}* T cells.

Tc17 cells play a major role in microbial infections, providing a potent anti-viral response
(Hamada et al., 2009; Yeh et al., 2010), while viral infection has been an established factor in
COPD exacerbations (Hewitt et al., 2016; Wedzicha, 2004). It will be interesting to determine
whether loss of *let-7bc2* or *let-7afd*-cluster activity in the T cell compartment contributes to COPD
disease susceptibility in the context of viral exposure. Our experiments with *let-7* GOF were
partially successful in limiting the emergence of Tc17 and Th17 in nCB-elicited emphysema but it
did not protect the lung from alveolar remodeling. Additional, studies will be required to ascertain
whether interventions that enhance *let-7* activity are successful in acute models of CS exposure
and COPD or in other chronic inflammation diseases associated with dysregulation of RORγt and
IL17^{+} injury.

MATERIALS AND METHODS

Mice

Conditional knockout-ready floxed *let-7bc2* and *let-7afd* mice were generated using CRISPR gene
editing in an isogenic C57BL/6 genetic background and were sequence verified for rigor. Mice
were PCR genotyped from ear samples with primers flanking loxP sites (Supplementary Table).
The *let-7bc2^{flox/flox}*; *CD4-cre* and *let-7afd^{flox/flox}*; *CD4-cre* mice were PCR genotyped. The *R26-*
STOP-rtTA; *Col1a1-tet0-let-7* (*rtTA-iLet7)* mice were obtained from JAX Jax Stocks 023912 and
05670 and then bred to *CD4-Cre* were PCR genotyped with established JAX primers. Control
rtTA-iLet7 and the *let-7^{GOF}* mice were fed *ad libitum* with 200mg/kg of doxycycline-containing chow
(Bio-Serv S3888) at weaning age. Syngeneic littermates served as controls for all mouse
experiments. All mice were bred in the transgenic animal facility at Baylor College of Medicine. All
experimental protocols used in this study were approved by the Institutional Animal Care and Use

Committee of Baylor College of Medicine animal protocol (AN-7389) and followed the National

Research Council Guide for the Care and Use of Laboratory Animals.

Human emphysema tissue samples and T cell isolation

Lung tissues were obtained from a total of 28 non-atopic current or former smokers with significant

(>20 pack-years, one pack-year equals to smoking one pack of cigarettes per day each year)

history of smoking who were recruited into studies from the chest or surgical clinics at Michael E.

DeBakey Houston Veterans Affairs Medical Center hospitals (Shan et al., 2009). Human lung

single T cells were prepared from surgical resection and lungs in patients as previously described

by selection with biotin-labeled antibodies by autoMACs (Miltenyi Biotec) (Yuan et al., 2020;

Grumelli et al., 2004). Studies were approved by the Institutional Review Board at Baylor College

of Medicine and informed consent was obtained from all patients. Emphysema and non-

emphysema control patients were diagnosed from CT scans according to the criteria

recommended by the National Institutes of Health–World Health Organization workshop summary

(Pauwels et al., 2001).

Human lung transcriptome data

A publicly available RNA-seq dataset from a Korean cohort GSE57148 was selected for the

analysis (Kim et al., 2015). The raw FASTQ files of paired end reads representing the

transcriptome of control and cases were retrieved from the GEO database at the National Centre

for Biological Information (NCBI) through accession number GSE57148 and analyzed with R

package for differential expression.

Cigarette smoke exposure model of pulmonary emphysema

To promote emphysema, mice were exposed to cigarette smoke using our custom designed

whole-body inhalation system (Morales-Mantilla et al., 2020). In total, mice were exposed to four

cigarettes (Marlboro 100's; Philip Morris USA) per day, five days a week, for four months as

previously described (Morales-Mantilla et al., 2020, Shan et al., 2012).

nCB exposure model of pulmonary emphysema

Nano-sized particulate carbon black was prepared and administered as previously described (You

et al., 2015; Lu et al., 2015). Dried nCB nanoparticles were resuspended in sterile PBS to a

concentration of 10 mg/ml. Fifty μl of reconstituted nCB (0.5 mg) were intranasally delivered to

deeply anesthetized mice on a schedule of three times a week for four weeks (total delivered

dose of 6 mg). Lung histomorphometry and airway inflammation were assessed four weeks after

the final nCB challenge. For histomorphometric analysis, mice lungs were fixed with 10% neutral-

buffered formalin solution via a tracheal cannula at 25-cm H_2O pressure followed by paraffin

embedding and tissue sectioning and stained with hematoxylin and eosin. Mean linear intercept

(MLI) measurement of mouse lung morphometry were done as previously described (Shan et al.,

2014; Morales-Mantilla et al., 2020). Briefly, this was done in a blinded fashion to mice genotypes

from ten randomly selected fields of lung parenchyma sections. Paralleled lines were placed on

serial lung sections and MLI was calculated by multiplying the length and the number of lines per

field, divided by the number of intercepts (Morales-Mantilla et al., 2020).

BALF was collected by instilling and withdrawing 0.8 ml of sterile PBS twice through the

trachea. Total and differential cell counts in the BALF were determined with the standard

hemocytometer and HEMA3 staining (Biochemical Sciences Inc, Swedesboro, NJ) using 200 μL

of BALF for cytospin slide preparation (Morales-Mantilla et al., 2020; Lu et al., 2015).

Cell isolation from murine lung tissue

Mouse lung tissue were cut into 2-mm pieces and digested with collagenase type D (2 mg/ml;

Worthington) and deoxyribonuclease (DNase) I (0.04 mg/ml; Roche) for 1 hour in a 37°C

incubator. Single-cell suspensions from lung digest, spleen, and thymus were prepared by

mincing through 40-μm cell strainers then washing and resuspension in complete RPMI media.

Mouse lung and spleen single-cell suspensions were additionally overlaid on Lympholyte M cell

separation media (Cedarlane) as indicated in the manufacturer's protocol to purify lymphocytes.

For murine *let-7* expression studies, lung single-cell suspensions were labeled with anti-CD4[+] or

anti-CD8[+] magnetic beads and separated by autoMACS (Miltenyi Biotec), or CD4[+]CD8[+] double

positive cells purified from thymus single-cell suspensions by flow-cytometric sorting on FACS

Aria (BD Biosciences).

***In vitro* polarization of CD8[+] T cells**

CD8[+] naïve T cells were isolated from spleen using Mojosort Mouse CD8 Naïve T cell isolation

Kit (Biolegend) and adjusted to a concentration of $1.0x10^6$ cells/mL. Purified cells were activated

with plate-bound anti-CD3 (1.5μg/mL) and complete RPMI media containing anti-CD28

(1.5μg/mL) and β-mercaptoethanol (50nM) for Tc0 polarization, or further supplemented with Tc1

[IL-2 (10ng/mL)] or Tc17 [TGFβ (2ng/mL), IL-6 (20ng/mL), anti-IFNγ (10μg/mL), IL-23 (20ng/mL),

and IL-1β (5ng/mL)] polarization conditions for 72 hours (Flores-Santibáñez et al., 2018).

ELISA

Supernatant was collected from *in vitro* polarized murine CD8[+] T cells and centrifuged to remove

cellular debris. Cytokine levels of IL-17a and IFNγ were quantified from collected supernatant

using Mouse IL-17a Uncoated ELISA and Mouse IFN gamma Uncoated ELISA (Invitrogen) Kits,

respectively, per the manufacturer's instructions with colorimetric analysis by the Varioskan LUX

microplate reader (ThermoFisher).

Flow cytometric analysis

Cells used for *in vitro* or *in vivo* cytokine analysis were stimulated with PMA (20ng/mL; Sigma

Aldrich), Ionomycin (1μg/mL; Sigma Aldrich), and Brefeldin A (2μg/mL; Sigma Aldrich) for 4 hours

prior to flow staining (Lu et al., 2015). For intracellular staining, cells were fixed and permeabilized

using the Mouse FOXP3 Buffer Set (BD) per the manufacturer's protocol. The fluorophore-

conjugated antibodies used in this study were as follows: Live/Dead Fix Blue (Invitrogen), CD3

PerCPCy5.5 (Biolegend), TCRb PE/Cy7 (Biolegend), CD4 PB (Biolegend), CD4 AF700

(Biolegend), CD8 BV650 (Biolegend), CD25 BV421 (Biolegend), FOXP3 AF488 (Biolegend),

ROR gamma T PE (Invitrogen), TCF1 AF647 (Cell Signaling Technologies), TCF1 PE (Biolegend),

IFNγ AF647 (Biolegend), IL17a FITC (Biolegend), IL17a PE (ebioscience), PD1 BUV737 (BD),

TIM3 AF647 (R&D). Samples were analyzed using BD LSR II flow cytometer (BD Biosciences)

and FlowJo software (TreeStar),

RNA Isolation and Quantitative RT-PCR

RNA was isolated using miRNeasy (Qiagen) or RNeasy Mini Kit (Qiagen) in conjunction with the

RNase-Free DNase (Qiagen) according to the manufacturer's instructions. cDNA of miRNAs and

mRNAs were synthesized using TaqMan Advanced miRNA cDNA Synthesis Kit (ThermoFisher)

and High-Capacity cDNA Reverse Transcription Kit Real-Time PCR system (Applied Biosystems).

18S and snoRNA-202 were used to normalize mRNA and miRNA expression respectively.

Quantitative RT-PCR data were acquired on 7500 Real-Time PCR System or StepOne Real-Time

PCR System (Applied Biosystems) with the following TaqMan probes: *hsa-let-7a* [000377], *hsa-*

let-7b [000378], *hsa-let-7d* [002283], *hsa-let-7f* [000382], *pri-miR-let7a* [Mm03306744_pri], *pri-*

miR-let7d [Mm03306666_pri], *pri-miR-let7b* [Mm03306756_pri], *Mmp9* [Mm00442991], *Mmp12*

[Mm00500554].

Luciferase reporter assays

Genomic fragment containing the murine *Rorc* 3'UTR was cloned into psiCHECK2 luciferase

reporter plasmid (Promega). This construct was also used to generate the *let-7* 'seed' deletion

mutant derivative using the QuikChange Multi Site Mutagenesis Kit (catalog 200514-5,

Stratagene). 3T3 mouse embryonic fibroblasts (MEFs) were transfected using Oligofectamine

(Invitrogen) with 100 ng of psiCheck-2 plasmid containing wild-type or mutant 3'UTR, along with

the miRNA control or *let-7b* duplex (Dharmacon) at a final concentration of 6 nM (Gurha et al.,

2012). Reporter activity was detected with the Dual-Luciferase Reporter Assay System

(Promega).

Statistical analysis

Statistical analyses were performed using GraphPad Prism 10.0.1 software. Statistical

comparison between groups was performed using the unpaired Student's t-test, two-way analysis

of variance (ANOVA) with Tukey's or Sidak's correction, and Mann-Whitney Test when indicated.

A P-value less than 0.05 was considered statistically significant; ns indicates not significant.

Statistical significance values were set as $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, and $****p < 0.0001$.

Data are presented as means ± SEM. P-value and sample sizes (n) are indicated in the figure

legends.

COMPETING INTEREST STATEMENT

The authors declare no competing interests.

ACKNOWLEDGEMENTS

We thank Jason Heaney and Denise Lanza at BCM Genetically Engineered Rodents Core

(funded in part by NIH P30 CA125123); Patricia Castro at Tissue Acquisition and Pathology Core

(funded in part by P30 CA125123); and Joel M. Sederstrom at the BCM and Cell Sorting Core

with funding from the CPRIT Core Facility Support Award (CPRIT-RP180672) and NIH

(CA125123 and RR024574). This work was supported by grants from the NHLBI (R01HL140398

to AR), the Gilson Longenbaugh Foundation (to A.R.), and NIEHS (T32 ES027801 to PE).

AUTHOR CONTRIBUTIONS

P.E., X.H., D.B.C, F.K., and A.R. conceptualized experiments and interpreted results. P.E., X.H., M.J.S., H.T., M.A.P., and S.L.L. acquired the data, M.J.R. provided bioinformatics analyses. P.E. and A.R. wrote the manuscript.

REFERENCES

Agarwal, V., Bell, G. W., Nam, J.-W., & Bartel, D. P. (2015). Predicting effective microRNA target sites in mammalian mRNAs. *eLife*, *4*, e05005. https://doi.org/10.7554/eLife.05005

Angelou, C. C., Wells, A. C., Vijayaraghavan, J., Dougan, C. E., Lawlor, R., Iverson, E., Lazarevic, V., Kimura, M. Y., Peyton, S. R., Minter, L. M., Osborne, B. A., Pobezinskaya, E. L., & Pobezinsky, L. A. (2020). Differentiation of Pathogenic Th17 Cells Is Negatively Regulated by Let-7 MicroRNAs in a Mouse Model of Multiple Sclerosis. *Frontiers in Immunology*, *10*, 3125. https://doi.org/10.3389/fimmu.2019.03125

Barnes, P. J. (2016). Inflammatory mechanisms in patients with chronic obstructive pulmonary disease. *Journal of Allergy and Clinical Immunology*, *138*(1), 16–27. https://doi.org/10.1016/j.jaci.2016.05.011

Beringer, A., Noack, M., & Miossec, P. (2016). IL-17 in Chronic Inflammation: From Discovery to Targeting. *Trends in Molecular Medicine*, *22*(3), 230–241. https://doi.org/10.1016/j.molmed.2016.01.001

Chang, Y., Nadigel, J., Boulais, N., Bourbeau, J., Maltais, F., Eidelman, D. H., & Hamid, Q. (2011). CD8 positive T cells express IL-17 in patients with chronic obstructive pulmonary disease. *Respiratory Research*, *12*(1), 43. https://doi.org/10.1186/1465-9921-12-43

Christenson, S. A., Brandsma, C.-A., Campbell, J. D., Knight, D. A., Pechkovsky, D. V., Hogg,

        J. C., Timens, W., Postma, D. S., Lenburg, M., & Spira, A. (2013a). miR-638 regulates

        gene expression networks associated with emphysematous lung destruction. *Genome*

        Medicine, *5*(12), 114. https://doi.org/10.1186/gm519

Christenson, S. A., Brandsma, C.-A., Campbell, J. D., Knight, D. A., Pechkovsky, D. V., Hogg,

        J. C., Timens, W., Postma, D. S., Lenburg, M., & Spira, A. (2013b). miR-638 regulates

        gene expression networks associated with emphysematous lung destruction. *Genome*

        Medicine, *5*(12), 114. https://doi.org/10.1186/gm519

Chu, S., Zhong, X., Zhang, J., Lao, Q., He, Z., & Bai, J. (2011). The expression of Foxp3 and

        ROR gamma t in lung tissues from normal smokers and chronic obstructive pulmonary

        disease patients. *International Immunopharmacology*, *11*(11), 1780–1788.

        https://doi.org/10.1016/j.intimp.2011.06.010

Conickx, G., Avila Cobos, F., van den Berge, M., Faiz, A., Timens, W., Hiemstra, P. S., Joos, G.

        F., Brusselle, G. G., Mestdagh, P., & Bracke, K. R. (2017). microRNA profiling in lung

        tissue and bronchoalveolar lavage of cigarette smoke-exposed mice and in COPD

        patients: A translational approach. *Scientific Reports*, *7*(1), Article 1.

        https://doi.org/10.1038/s41598-017-13265-8

Corgnac, S., Malenica, I., Mezquita, L., Auclin, E., Voilin, E., Kacher, J., Halse, H., Grynszpan,

        L., Signolle, N., Dayris, T., Leclerc, M., Droin, N., de Montpréville, V., Mercier, O.,

        Validire, P., Scoazec, J.-Y., Massard, C., Chouaib, S., Planchard, D., … Mami-Chouaib,

        F. (2020). CD103+CD8+ TRM Cells Accumulate in Tumors of Anti-PD-1-Responder

        Lung Cancer Patients and Are Tumor-Reactive Lymphocytes Enriched with Tc17. *Cell*

        Reports Medicine, *1*(7), 100127. https://doi.org/10.1016/j.xcrm.2020.100127

Duan, M.-C., Tang, H.-J., Zhong, X.-N., & Huang, Y. (2013). Persistence of Th17/Tc17 Cell

Expression upon Smoking Cessation in Mice with Cigarette Smoke-Induced Emphysema.

Journal of Immunology Research, *2013*, e350727. https://doi.org/10.1155/2013/350727

Duan, M.-C., Zhang, J.-Q., Liang, Y., Liu, G.-N., Xiao, J., Tang, H.-J., & Liang, Y. (2016).

Infiltration of IL-17-Producing T Cells and Treg Cells in a Mouse Model of Smoke-

Induced Emphysema. *Inflammation*, *39*(4), 1334–1344. https://doi.org/10.1007/s10753-

016-0365-8

Findings from the Global Burden of Disease Study 2017. (2019, January 4). Institute for Health

Metrics and Evaluation. https://www.healthdata.org/policy-report/findings-global-

burden-disease-study-2017

Globig, A.-M., Hipp, A. V., Otto-Mora, P., Heeg, M., Mayer, L. S., Ehl, S., Schwacha, H.,

Bewtra, M., Tomov, V., Thimme, R., Hasselblatt, P., & Bengsch, B. (2022). High-

dimensional profiling reveals Tc17 cell enrichment in active Crohn's disease and

identifies a potentially targetable signature. *Nature Communications*, *13*(1), Article 1.

https://doi.org/10.1038/s41467-022-31229-z

Grumelli, S., Corry, D. B., Song, L.-Z., Song, L., Green, L., Huh, J., Hacken, J., Espada, R., Bag,

R., Lewis, D. E., & Kheradmand, F. (2004). An Immune Basis for Lung Parenchymal

Destruction in Chronic Obstructive Pulmonary Disease and Emphysema. *PLOS*

Medicine, *1*(1), e8. https://doi.org/10.1371/journal.pmed.0010008

Guan, H., Fan, D., Mrelashvili, D., Hao, H., Singh, N. P., Singh, U. P., Nagarkatti, P. S., &

Nagarkatti, M. (2013). MicroRNA let-7e is associated with the pathogenesis of

experimental autoimmune encephalomyelitis. *European Journal of Immunology*, *43*(1),

104–114. https://doi.org/10.1002/eji.201242702

Hamada, H., Garcia-Hernandez, M. de la L., Reome, J. B., Misra, S. K., Strutt, T. M.,

McKinstry, K. K., Cooper, A. M., Swain, S. L., & Dutton, R. W. (2009). Tc17, a Unique

Subset of CD8 T Cells That Can Protect against Lethal Influenza Challenge1. *The*

Journal of Immunology, *182*(6), 3469–3481. https://doi.org/10.4049/jimmunol.0801814

Hewitt, R., Farne, H., Ritchie, A., Luke, E., Johnston, S. L., & Mallia, P. (2016). The role of

viral infections in exacerbations of chronic obstructive pulmonary disease and asthma.

Therapeutic Advances in Respiratory Disease, *10*(2), 158–174.

https://doi.org/10.1177/1753465815618113

Hoenderdos, K., & Condliffe, A. (2013). The Neutrophil in Chronic Obstructive Pulmonary

Disease. Too Little, Too Late or Too Much, Too Soon? *American Journal of Respiratory*

Cell and Molecular Biology, *48*(5), 531–539. https://doi.org/10.1165/rcmb.2012-0492TR

Huber, M., Heink, S., Pagenstecher, A., Reinhard, K., Ritter, J., Visekruna, A., Guralnik, A.,

Bollig, N., Jeltsch, K., Heinemann, C., Wittmann, E., Buch, T., Costa, O. P. da, Brüstle,

A., Brenner, D., Mak, T. W., Mittrücker, H.-W., Tackenberg, B., Kamradt, T., & Lohoff,

M. (2013). IL-17A secretion by CD8$^+$ T cells supports Th17-mediated autoimmune

encephalomyelitis. *The Journal of Clinical Investigation*, *123*(1), 247–260.

https://doi.org/10.1172/JCI63681

Ivanov, I. I., McKenzie, B. S., Zhou, L., Tadokoro, C. E., Lepelley, A., Lafaille, J. J., Cua, D. J.,

& Littman, D. R. (2006). The orphan nuclear receptor RORgammat directs the

differentiation program of proinflammatory IL-17+ T helper cells. *Cell*, *126*(6), 1121–

1133. https://doi.org/10.1016/j.cell.2006.07.035

Ivanov, I. I., Zhou, L., & Littman, D. R. (2007). Transcriptional Regulation of Th17 Cell

Differentiation. *Seminars in Immunology*, *19*(6), 409–417.

https://doi.org/10.1016/j.smim.2007.10.011

Izzotti, A., Calin, G. A., Arrigo, P., Steele, V. E., Croce, C. M., & De Flora, S. (2009).

Downregulation of microRNA expression in the lungs of rats exposed to cigarette smoke.

The FASEB Journal, *23*(3), 806–812. https://doi.org/10.1096/fj.08-121384

Jiang, S., Yan, W., Wang, S. E., & Baltimore, D. (2018). Let-7 Suppresses B Cell Activation

through Restricting the Availability of Necessary Nutrients. *Cell Metabolism*, *27*(2), 393-

403.e4. https://doi.org/10.1016/j.cmet.2017.12.007

Jin, Y., Wan, Y., Chen, G., Chen, L., Zhang, M.-Q., Deng, L., Zhang, J.-C., Xiong, X.-Z., & Xin,

J.-B. (2014). Treg/IL-17 Ratio and Treg Differentiation in Patients with COPD. *PLOS*

ONE, *9*(10), e111044. https://doi.org/10.1371/journal.pone.0111044

Kheradmand, F., Shan, M., Xu, C., & Corry, D. B. (2012). Autoimmunity in chronic obstructive

pulmonary disease: Clinical and experimental evidence. *Expert Review of Clinical*

Immunology, *8*(3), 285–292. https://doi.org/10.1586/eci.12.7

Kim, W. J., Lim, J. H., Lee, J. S., Lee, S.-D., Kim, J. H., & Oh, Y.-M. (2015). Comprehensive

Analysis of Transcriptome Sequencing Data in the Lung Tissues of COPD Subjects.

International Journal of Genomics, *2015*, 206937. https://doi.org/10.1155/2015/206937

Laniado-Laborín, R. (2009). Smoking and Chronic Obstructive Pulmonary Disease (COPD).

Parallel Epidemics of the 21st Century. *International Journal of Environmental Research*

and Public Health, *6*(1), Article 1. https://doi.org/10.3390/ijerph6010209

Lee, P. P., Fitzpatrick, D. R., Beard, C., Jessup, H. K., Lehar, S., Makar, K. W., Pérez-Melgosa,

M., Sweetser, M. T., Schlissel, M. S., Nguyen, S., Cherry, S. R., Tsai, J. H., Tucker, S.

M., Weaver, W. M., Kelso, A., Jaenisch, R., & Wilson, C. B. (2001). A Critical Role for Dnmt1 and DNA Methylation in T Cell Development, Function, and Survival. *Immunity*, *15*(5), 763–774. https://doi.org/10.1016/S1074-7613(01)00227-8

Lee, S.-H., Goswami, S., Grudo, A., Song, L.-Z., Bandi, V., Goodnight-White, S., Green, L., Hacken-Bitar, J., Huh, J., Bakaeen, F., Coxson, H. O., Cogswell, S., Storness-Bliss, C., Corry, D. B., & Kheradmand, F. (2007). Antielastin autoimmunity in tobacco smoking-induced emphysema. *Nature Medicine*, *13*(5), 567–569. https://doi.org/10.1038/nm1583

Li, H., Liu, Q., Jiang, Y., Zhang, Y., Xiao, W., & Zhang, Y. (2015). Disruption of Th17/Treg Balance in the Sputum of Patients With Chronic Obstructive Pulmonary Disease. *The American Journal of the Medical Sciences*, *349*(5), 392–397. https://doi.org/10.1097/MAJ.0000000000000447

Lu, W., You, R., Yuan, X., Yang, T., Samuel, E. L. G., Marcano, D. C., Sikkema, W. K. A., Tour, J. M., Rodriguez, A., Kheradmand, F., & Corry, D. B. (2015). MicroRNA-22 Inhibits Histone Deacetylase 4 to Promote T Helper-17 Cell-Dependent Emphysema. *Nature Immunology*, *16*(11), 1185–1194. https://doi.org/10.1038/ni.3292

Mai, J., Virtue, A., Maley, E., Tran, T., Yin, Y., Meng, S., Pansuria, M., Jiang, X., Wang, H., & Yang, X.-F. (2012). MicroRNAs and other mechanisms regulate interleukin-17 cytokines and receptors. *Frontiers in Bioscience (Elite Edition)*, *4*, 1478–1495.

Morales-Mantilla, D. E., Huang, X., Erice, P., Porter, P., Zhang, Y., Figueroa, M., Chandra, J., King, K. Y., Kheradmand, F., & Rodríguez, A. (2020). Cigarette Smoke Exposure in Mice using a Whole-Body Inhalation System. *JoVE (Journal of Visualized Experiments)*, *164*, e61793. https://doi.org/10.3791/61793

Newcomb, D. C., Cephus, J. Y., Boswell, M. G., Fahrenholz, J. M., Langley, E. W., Feldman, A. S., Zhou, W., Dulek, D. E., Goleniewska, K., Woodward, K. B., Sevin, C. M., Hamilton, R. G., Kolls, J. K., & Peebles, R. S. (2015). Estrogen and progesterone decrease let-7f microRNA expression and increase IL-23/IL-23 receptor signaling and IL-17A production in patients with severe asthma. *Journal of Allergy and Clinical Immunology*, *136*(4), 1025-1034.e11. https://doi.org/10.1016/j.jaci.2015.05.046

Pauwels, R. A., Buist, A. S., Calverley, P. M. A., Jenkins, C. R., & Hurd, S. S. (2001). Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease. *American Journal of Respiratory and Critical Care Medicine*, *163*(5), 1256–1276. https://doi.org/10.1164/ajrccm.163.5.2101039

Peleman, R. A., Rytila, P. H., Kips, J. C., Joos, G. F., & Pauwels, R. A. (1999). The cellular composition of induced sputum in chronic obstructive pulmonary disease. *European Respiratory Journal*, *13*(4), 839–843. https://doi.org/10.1034/j.1399-3003.1999.13d24.x

Pottelberge, G. R. V., Mestdagh, P., Bracke, K. R., Thas, O., Durme, Y. M. T. A. van, Joos, G. F., Vandesompele, J., & Brusselle, G. G. (2011). MicroRNA Expression in Induced Sputum of Smokers and Patients with Chronic Obstructive Pulmonary Disease. *American Journal of Respiratory and Critical Care Medicine*, *183*(7), 898–906. https://doi.org/10.1164/rccm.201002-0304OC

Prevalence and attributable health burden of chronic respiratory diseases, 1990–2017: A systematic analysis for the Global Burden of Disease Study 2017. (2020). *The Lancet. Respiratory Medicine*, *8*(6), 585–596. https://doi.org/10.1016/S2213-2600(20)30105-3

Rodriguez, A., Griffiths-Jones, S., Ashurst, J. L., & Bradley, A. (2004). Identification of

Mammalian microRNA Host Genes and Transcription Units. *Genome Research*, *14*(10a),

1902–1910. https://doi.org/10.1101/gr.2722704

Schembri, F., Sridhar, S., Perdomo, C., Gustafson, A. M., Zhang, X., Ergun, A., Lu, J., Liu, G.,

Zhang, X., Bowers, J., Vaziri, C., Ott, K., Sensinger, K., Collins, J. J., Brody, J. S., Getts,

R., Lenburg, M. E., & Spira, A. (2009). MicroRNAs as modulators of smoking-induced

gene expression changes in human airway epithelium. *Proceedings of the National*

Academy of Sciences, *106*(7), 2319–2324. https://doi.org/10.1073/pnas.0806383106

Senior, R. M., & Anthonisen, N. R. (1998). Chronic Obstructive Pulmonary Disease (COPD).

American Journal of Respiratory and Critical Care Medicine, *157*(4), S139–S147.

https://doi.org/10.1164/ajrccm.157.4.nhlbi-12

Shan, M., Cheng, H.-F., Song, L., Roberts, L., Green, L., Hacken-Bitar, J., Huh, J., Bakaeen, F.,

Coxson, H. O., Storness-Bliss, C., Ramchandani, M., Lee, S.-H., Corry, D. B., &

Kheradmand, F. (2009). Lung Myeloid Dendritic Cells Coordinately Induce TH1 and

TH17 Responses in Human Emphysema. *Science Translational Medicine*, *1*(4), 4ra10-

4ra10. https://doi.org/10.1126/scitranlsmed.3000154

Shan, M., You, R., Yuan, X., Frazier, M. V., Porter, P., Seryshev, A., Hong, J.-S., Song, L.,

Zhang, Y., Hilsenbeck, S., Whitehead, L., Zarinkamar, N., Perusich, S., Corry, D. B., &

Kheradmand, F. (2014). Agonistic induction of PPARγ reverses cigarette smoke–induced

emphysema. *The Journal of Clinical Investigation*, *124*(3), 1371–1381.

https://doi.org/10.1172/JCI70587

Shan, M., Yuan, X., Song, L., Roberts, L., Zarinkamar, N., Seryshev, A., Zhang, Y., Hilsenbeck,

S., Chang, S.-H., Dong, C., Corry, D. B., & Kheradmand, F. (2012). Cigarette Smoke

Induction of Osteopontin (SPP1) Mediates TH17 Inflammation in Human and

Experimental Emphysema. *Science Translational Medicine*, *4*(117), 117ra9-117ra9.

https://doi.org/10.1126/scitranslmed.3003041

Shi, J., & Petrie, H. T. (2012). Activation Kinetics and Off-Target Effects of Thymus-Initiated

Cre Transgenes. *PLoS ONE*, *7*(10), e46590. https://doi.org/10.1371/journal.pone.0046590

Veldhoen, M. (2017). Interleukin 17 is a chief orchestrator of immunity. *Nature Immunology*,

18(6), Article 6. https://doi.org/10.1038/ni.3742

Wedzicha, J. A. (2004). Role of Viruses in Exacerbations of Chronic Obstructive Pulmonary

Disease. *Proceedings of the American Thoracic Society*, *1*(2), 115–120.

https://doi.org/10.1513/pats.2306030

Williams, M., Todd, I., & Fairclough, L. C. (2021). The role of CD8 + T lymphocytes in chronic

obstructive pulmonary disease: A systematic review. *Inflammation Research*, *70*(1), 11–

18. https://doi.org/10.1007/s00011-020-01408-z

Xu, C., Hesselbacher, S., Tsai, C.-L., Shan, M., Spitz, M., Scheurer, M., Roberts, L., Perusich,

S., Zarinkamar, N., Coxson, H., Krowchuk, N., Corry, D., & Kheradmand, F. (2012).

Autoreactive T Cells in Human Smokers is Predictive of Clinical Outcome. *Frontiers in*

Immunology, *3*. https://www.frontiersin.org/articles/10.3389/fimmu.2012.00267

Yeh, N., Glosson, N. L., Wang, N., Guindon, L., McKinley, C., Hamada, H., Li, Q., Dutton, R.

W., Shrikant, P., Zhou, B., Brutkiewicz, R. R., Blum, J. S., & Kaplan, M. H. (2010).

Tc17 Cells Are Capable of Mediating Immunity to Vaccinia Virus by Acquisition of a

Cytotoxic Phenotype. *The Journal of Immunology*, *185*(4), 2089–2098.

https://doi.org/10.4049/jimmunol.1000818

You, R., Lu, W., Shan, M., Berlin, J. M., Samuel, E. L., Marcano, D. C., Sun, Z., Sikkema, W.

K., Yuan, X., Song, L., Hendrix, A. Y., Tour, J. M., Corry, D. B., & Kheradmand, F.

(2015). Nanoparticulate carbon black in cigarette smoke induces DNA cleavage and

Th17-mediated emphysema. *eLife*, *4*, e09623. https://doi.org/10.7554/eLife.09623

Zhang, H., Zhou, X., Chen, X., Lin, Y., Qiu, S., Zhao, Y., Tang, Q., Liang, Y., & Zhong, X.

(2019). Rapamycin attenuates Tc1 and Tc17 cell responses in cigarette smoke-induced

emphysema in mice. *Inflammation Research*, *68*(11), 957–968.

https://doi.org/10.1007/s00011-019-01278-0

Zhou, J.-S., Li, Z.-Y., Xu, X.-C., Zhao, Y., Wang, Y., Chen, H.-P., Zhang, M., Wu, Y.-F., Lai,

T.-W., Di, C.-H., Dong, L.-L., Liu, J., Xuan, N.-X., Zhu, C., Wu, Y.-P., Huang, H.-Q.,

Yan, F.-G., Hua, W., Wang, Y., … Shen, H.-H. (2020). Cigarette smoke-initiated

autoimmunity facilitates sensitisation to elastin-induced COPD-like pathologies in mice.

European Respiratory Journal, *56*(3). https://doi.org/10.1183/13993003.00404-2020

Zhu, H., Shyh-Chang, N., Segrè, A. V., Shinoda, G., Shah, S. P., Einhorn, W. S., Takeuchi, A.,

Engreitz, J. M., Hagan, J. P., Kharas, M. G., Urbach, A., Thornton, J. E., Triboulet, R.,

Gregory, R. I., Altshuler, D., & Daley, G. Q. (2011). The Lin28/let-7 Axis Regulates

Glucose Metabolism. *Cell*, *147*(1), 81–94. https://doi.org/10.1016/j.cell.2011.08.033

Figure 1. Repression of *Let-7* miRNA gene clusters in lung T cells from COPD patients and murine models of emphysema. (A) Schematic representation of the polycistronic transcripts for the *Let-7a1/Let-7f1/Let-7d-* and *Let-7b/Let-7a3*-clusters in humans and *let-7a1/let-7f1/let-7/d-* and l*et-7b/let-7c2*-clusters in mice. (B) *in silico* analysis of *Mirlet7a1hg* and *Mirlet7bhg* from the publicly available lung transcriptome dataset from RNA-seq of COPD and control patients (GEO: GSE57148). (C) Quantitative RT-PCR (qPCR) of mature *Hsa-Let-7a* from resected lung tissue of COPD (n=15) and control subjects (n=11). (D) qPCR and regression analysis of *Hsa-Let-7a, Hsa-Let-7b, Hsa-Let-7d, and Hsa-Let-7f* expression to emphysema severity score based on CT: 0=no, 1=upper lobes only, 2=upper/middle lobes, 3=extensive pan lobular emphysema (n=19). (E) Schematic diagram of experimental emphysema in mice induced by either intranasal (i.n.) instillation of nCB or exposure to CS by whole-body inhalation (w.b.i.). (F-H) qPCR analysis for *pri-let-7a1/f1/d* and *pri-let-7b/c2* from lung tissue or lung-derived CD8+ and CD4+ T cells of mice with emphysema elicited by (F) nCB- or (G-H) CS (n=3-6 per group). Data are representative of three independent experiments displayed as mean±SEM. Mann-Whitney (B,C) or Student's t-test (F,G,H). *p < 0.05, **p < 0.01, ***p <0.001, ****p<0.0001.

Figure 2. Deletion of the *let7bc2* cluster in T cells enhances nCB- or CS-triggered emphysema. (A) Schematic representation of CD4-Cre (*let-7bc2^LOF^*) or *let-7bc2^f/f^* (Control) mice. (B) qPCR analysis of pri-miRNA transcript for *pri-let-7bc2* from flow-sorted live, TCRβ+, CD4+CD8+ double-positive (DP) thymocytes of control and *let-7bc2^LOF^* mice (n=3-5 per group). (C-G) Control and *let-7bc2^LOF^* mice were exposed to vehicle (PBS) or nCB for 4 weeks, or alternatively air or cigarette smoke by whole body inhalation of cigarette smoke (CS) for 16 weeks. (C) Representative H&E stained lung sections from PBS-, nCB-, or CS-exposed mice as indicated on each panel (x20 magnification; scale bars, 50μm). (D-E) Mean linear intercept (MLI) measurement of lung morphometry. (F) Total and differential cell counts from bronchoalveolar lavage (BAL) fluid from controls and nCB-emphysemic mice (n=4-7 per group). (G) *Mmp9* and *Mmp12* mRNA expression from BAL cells of air- and smoke-exposed control and *let-7bc2^LOF^* mice (n=4-6 per group). Data are representative of at least three independent experiments displayed as mean±SEM using Student's t-test (B) or two-way ANOVA with *post-hoc* Tukey correction (D,E,F,G). *p < 0.05, **p < 0.01, ***p <0.001, ****p<0.0001.

Figure 3. *In vivo* **T cell ablation of the** *let-7bc2***-cluster enhances Tc17 inflammatory response to nCB-emphysema.** Representative flow plots with percentage and counts of live TCRβ⁺ (A) CD8⁺IL-17a⁺ and CD8⁺IFNγ⁺, (B) CD8⁺IFNγ⁺GzmA⁺, (C) CD4⁺IL-17a⁺ and CD4⁺IFNγ⁺, and (D) CD4⁺ Foxp3⁺CD25⁺ cells from the lungs of control PBS vehicle- (n=5-6), control nCB- (n=6), and *let-7bc2*^LOF nCB-exposed mice. Data are representa-tive of three independent experiments displayed as mean±SEM using ANOVA with *post-hoc* Sidak correction. *p < 0.05, **p < 0.01, ***p <0.001, ****p<0.0001.

Figure 4. Deletion of either the *let7bc2-* or *let7afd*-cluster in T cells enhances RORγt expression *in vivo*. (A) Left: Schematic representation of the murine *Rorc* 3'UTR with *let-7* miRNA binding sites as identified by TargetScan. Right: Schematic of a conserved *let-7* miRNA target sequence in the 3'UTR of *Rorc*. (B-C) Flow analysis of RORγt expression by MFI quantification in live TCRβ⁺CD8⁺ or CD4⁺ T cells from indicated tissues of (B) naïve control and *let-7bc2^LOF* mice or (C) nCB-treated lungs by representative flow plot and MFI quantification (n=5 per group). (D) RORγt expression by MFI quantification in naïve mice *let-7afd^LOF* mice thymus, spleen, and lungs (n=3-4 per group), or (E) nCB-ex-posed lungs (n=5 per group). Data are representative of at least three independent experiments displayed as mean±SEM using student's t-test. *p < 0.05, **p < 0.01, ***p <0.001, ****p<0.0001.

Figure 5. *Let-7* restricts Tc17 *in vitro* differentiation in part via direct targeting of *Rorc* mRNA. (A) Representative flow plots of live TCRβ⁺ CD8⁺, IL-17a⁺ and IFNγ⁺ populations from Tc1 and Tc17 polarized naïve splenic CD8⁺ cells from control and *let-7bc2^LOF* mice and (B) quantification of CD8⁺IL-17a⁺ cells (n=5 per group). (C) ELISA of IL-17a from the supernatant of Tc1 and Tc17 polarized control and *let-7bc2^LOF* cells (n=5-6 per group). (D) Flow quantification of CD8⁺IFNγ⁺ populations in Tc1 and Tc17 polarized control and *let-7bc2^LOF* cells (n=5 per group). (E) Representative flow plots of CD8⁺IL-17a⁺ population frequency in Tc17 polarized cells indicated mice, and (E) quantification of CD8⁺IL-17a⁺ cells from control and *let-7afd^LOF* mice polarized under Tc1 or Tc17 conditions. (F) ELISA of IL-17a from control Tc1 (n=4), control Tc17 (n=4), and *let-7afd^LOF* Tc17 (n=3) polarized cells. (G) Representative flow plot and quantification of RORγt from Tc0 or Tc17 differentiated naïve splenic CD8⁺ T cells isolated from control and *let-7bc2^LOF* mice (n=5 per group). (H) Control (*Rorc* WT) or binding site mutant (*Rorc* Mut) 3' UTRs of *Rorc* were cloned downstream of the renilla luciferase reporter. Plasmids were cotransfected with either a control-miR (black bars) or *let-7b* mimic (blue bars) duplex into cultured cells. Reporter activity was measured 24 hours after transfection and normalized to firefly activity. Data are representative of three independent experiments (A-G) or carried out in triplicate (H) and displayed as mean±SEM using student's t-test. *p < 0.05, **p < 0.01, ***p <0.001, ****p<0.0001.

Figure 6. Enforced *let-7* expression in T cells restrains induction of RORγt and Tc17/Th17 inflammation in lungs of nCB-exposed mice. (A) Schematic outlining our T cell-inducible *let-7g* mouse model (*let-7^GOF*). (B) Flow analysis of RORγt expression in live, TCRβ+CD8+ or CD4+ T cells from (B) naïve control and *let-7^GOF* mice in thymus, spleen, and lungs (n=3-5 per group). (C) Control and *let-7^GOF* mice were treated with PBS vehicle or nCB then analyzed. Representative H&E-stained lung sections from PBS- and nCB-exposed mice as indicated on each panel (x20 magnification; scale bars, 50μm) (D) MLI measurements from indicated mice (n=5-6 per group). (E) Flow analysis of lungs gated on live TCRβ+ CD8+ or CD4+ cells for (E) IL-17a+ population frequency (n=3-4 per group) or (F) RORγt expression by representative flow plot and MFI quantification (n=4-5 per group). (G) Figure model for *let-7*/RORγt axis in emphysema pathogenesis. Data are representative of two or three independent experiments and displayed as mean±SEM using student's t-test (B) or two-way ANOVA with Tukey's multiple correction (C,D,E). *p < 0.05, **p < 0.01, ***p <0.001, ****p<0.0001.

Figure Supplement 1. T cell-specific deletion of the *let-7bc2*-cluster does not promote lung inflammation or pathology with moderate aging. (A) Representative H&E-stained lung sections from Control and *let-7bc2^LOF^* naive mice aged to one year (x20 magnification; scale bars, 50μm). (B) MLI measurements of indicated mice (n=5-6 per group). (C) Total and differential cell count from bronchoalveolar lavage (BAL) fluid from control and *let-7bc2^LOF^* naive mice (n=6 per group; Mac. = macrophages, Neut. = neutrophils). (D) Flow cytometric analysis of CD4+, CD8+, or DP T cells from the lungs, spleen, and thymus of control and *let-7bc2^LOF^* mice at steady-state (n=4-6 per group). (E) Immunophenotyping of Tc17, Tc1, Th17, Th1, and Tregs from lungs of naïve control and *let-7bc2^LOF^* mice (n=3-5 per group). Data are representative of three independent experiments and displayed as mean±SEM using student's t-test.

Table S1. Primers used in this study

Name	Sequence	Description
AFD-lox2-F	5'-GTTTTCTGAGGTGTGGGAG	let-7afd-cluster floxed allele Forward primer for genotyping
AFD-lox2-R	5'-AGTGGGATAGAAGGATCTC	let-7afd-cluster floxed allele Reverse primer for genotyping
BC-lox2-F	5'-GGACATGAGATCGCCAACC	let-7bc2-cluster floxed allele Forward primer for genotyping
BC-lox2-R	5'-TGGAAGCCAGTACTGTGCT	let-7bc2-cluster floxed allele Reverse primer for genotyping