Autism candidate gene *rbm-26* (*RBM26/27*) regulates MALS-1 to protect against mitochondrial dysfunction and axon degeneration during neurodevelopment.

Tamjid A Chowdhury[1], David A Luy[1], Garrett Scapellato[1] Dorian Farache[2,3], Amy SY Lee[2,3], Christopher C Quinn[1*]

[1] Department of Biological Sciences, University of Wisconsin-Milwaukee, Milwaukee, WI, USA.
[2] Department of Cell Biology, Harvard Medical School, Boston, MA, USA.
[3] Department of Cancer Immunology and Virology, Dana-Farber Cancer Institute, Boston, MA, USA.

[*]To whom correspondence should be addressed at: Department Biological Sciences
3209 N. Maryland Avenue
Milwaukee, WI 53211

EMAIL: quinnc@uwm.edu
Phone: (414)229-2602
Fax: (414)229-3926

Abstract

Mitochondrial dysfunction is thought to be a key component of neurodevelopmental
disorders such as autism, intellectual disability, and ADHD. However, little is known
about the molecular mechanisms that protect against mitochondrial dysfunction during
neurodevelopment. Here, we address this question through the investigation of *rbm-26*,
the *C. elegans* ortholog of the *RBM27* autism candidate gene, which encodes an RNA-
binding protein whose role in neurons is unknown. We report that RBM-26 (RBM26/27)
protects against axonal defects by negatively regulating expression of the MALS-1
(MALSU1) mitoribosomal assembly factor. Autism-associated missense variants in
RBM-26 cause a sharp decrease in RBM-26 protein expression along with defects in
in axon tiling and axon degeneration that occurs during larval development. Using a
biochemical screen, we identified the mRNA for the MALS-1 mitoribosomal assembly
factor as a binding partner for RBM-26. Loss of RBM-26 function causes a dramatic
overexpression of *mals-1* mRNA and MALS-1 protein. Moreover, genetic analysis
indicates that this overexpression of MALS-1 is responsible for the mitochondrial and
axon degeneration defects in *rbm-26* mutants. These observations reveal a mechanism
that regulates expression of a mitoribosomal assembly factor to protect against axon
degeneration during neurodevelopment.

Introduction

Emerging evidence suggests an association between mitochondria and
neurodevelopmental disorders including autism, intellectual disability, and attention
deficit hyperactivity disorder (ADHD). For example, certain mitochondrial DNA lineages
confer substantial risk for autism and ADHD [1, 2]. Likewise, heteroplasmic mtDNA
mutations are also a risk factor for autism and intellectual disability [3-5]. Consistent with
a causative role for mitochondria dysfunction in neurodevelopmental disorders, mice
with a hypomorphic missense mutation in the *ND6* mtDNA gene exhibit autism
endophenotypes and also produce excess ROS in the brain [6]. In this regard, it is
intriguing to note that elevated levels of reactive oxygen species (ROS) have also been
observed in autistic human brains [7]. Together, these observations suggest that
mitochondrial dysfunction plays a causative role in neurodevelopmental disorders.
However, the roles of mitochondrial dysfunction in neurodevelopmental disorders are
not well understood.

Some neurodevelopmental syndromes feature neurodegenerative phenotypes that
begin during the developmental time period. For example, loss of *EXOSC3* or *EXOSC9*
gene function is thought to cause a neurodevelopmental syndrome that includes
intellectual disability and axon degeneration during infancy [8-10]. Likewise, loss of the

GAN gene in humans causes the giant axonal neuropathy disease, featuring intellectual
disability and axon degeneration in children [11]. Moreover, loss of either the *SPTBN1*
or *ADD1* genes are thought to cause a neurodevelopmental syndrome that includes
autism and ADHD [12, 13]. In mouse models, variants in these genes cause defects in
axon development along with axon degeneration, consistent with a causative role for
these genes in the human syndrome [14, 15]. Together, these observations suggest
that neurodegeneration may contribute to some neurodevelopmental disorders.
However, little is known about the cellular and molecular mechanisms that protect
against neurodegeneration in infants and children.

The *RBM27* (*rbm-26/Swm/Rmn1*) gene encodes an RNA binding protein and is
considered an autism candidate gene by the Simmons Foundation Autism Research
Initiative (SFARI) [16]. In a cultured human cell line, RBM27 and its paralog RBM26 can
function with the PAXT connection, an adaptor complex for the nuclear RNA exosome
[17]. Likewise, in *S. pombe* the Rmn1 ortholog of RBM26/27 can also function as part of
an adaptor for the nuclear RNA exosome [18]. In *Drosophila,* the Swm ortholog of
RBM26/27 has also been implicated in RNA metabolism and the maintenance of adult
intestinal stem cells [19]. Despite these findings, the role of RBM27 and its orthologs in
neurons have not been investigated. Moreover, the identity of *RBM27* as an autism
candidate gene is based on de novo missense mutations that have not been tested for
gene-disrupting activity.

Here, we identify a mitochondria-protecting role for the RBM-26 ortholog of RBM26/27
that protects mitochondria and prevents axon degeneration during larval development.
We find that loss of RBM-26 function causes a decrease in mitochondrial density in the
axon, mitochondrial dysfunction in the cell body, axon degeneration, and transient
defects in axon tiling. These phenotypes are induced by an *rbm-26* null allele as well as
by alleles that are equivalent to autism-associated de novo missense variants in
RBM27, suggesting that these are likely gene-disrupting variants in humans.
Mechanistically, we identified the *mals-1* mRNA as a binding partner for RBM-26 and
found that RBM-26 negatively regulates the expression of the *mals-1* mRNA, which
encodes a mitoribosomal assembly factor. Moreover, gain-of-function and loss-of
function experiments indicate that MALS-1 is required for both the axon degeneration
phenotype and the reduction in axonal mitochondria that are caused by loss of RBM-26.
These observations indicate that RBM-26 negatively regulates MALS-1 expression to
protect against mitochondrial dysfunction and axon degeneration during development.
Based on our findings in *C. elegans*, along with prior work on RBM26/27 [17], we
speculate that RBM27 may function with the RNA exosome in humans to protect
against neurodevelopmental disorders.

Results

Expression of RBM-26 is disrupted by P80L and L13V missense mutations.

RBM27 has been identified as a candidate autism gene by the Simons Foundation
Autism Research Initiative (SFARI) [16]. The potential association of *RBM27* with
autism is based on the identification of 5 de novo variants in probands with autism or
other neurodevelopmental disorders, but none in unaffected siblings (Fig. 1A) [20-25].
However, the confidence in this association is limited by the fact that the identified
variants are all missense variants, rather than likely-gene-disrupting variants. Moreover,
a role for *RBM27* in neurodevelopment has not been investigated. To address these
issues, we used the *C. elegans rbm-26* ortholog of *RBM27.* We found that 2 of the 5 de
novo missense mutation sites in human RBM27 are conserved in *C. elegans rbm-26*.
These two conserved missense mutations cause L13V and P79L amino acid changes in
human RBM27, which are equivalent to L13V and P80L respectively in *C. elegans*
RBM-26 (Fig. 1B). Both of these mutations occur in the PWI-like domains and were
identified in genome sequencing studies of children with neurodevelopmental disorders
[20, 23].

To determine how the L13V and P80L variants affect RBM-26 protein expression, we
used CRISPR to introduce these mutations into the endogenous *rbm-26* gene and also
added a 3XFLAG tag to allow for detection by western blot. A sequence encoding
3XFLAG was inserted just upstream of the *rbm-26* stop codon, thereby creating the
rbm-26(cue25) allele, hereafter referred to as *rbm-26(3XFLAG).* Next, we used CRISPR
to introduce the P80L mutation into the *rbm-26(3XFLAG)* allele, thereby creating the
rbm-26(cue22cue25) allele, hereafter called *rbm-26(P80L::3XFLAG).* We also used
CRISPR to introduce the L13V mutation into the *rbm-26(3XFLAG)* allele, thereby
creating the *rbm-26(cue24cue25)* allele, hereafter called *rbm-26(L13V::3XFLAG).*

We next used western blotting to detect the RBM-26::3XFLAG, RBM-26
P80L::3XFLAG, and RBM-26 L13V::3XFLAG proteins **(Figure 1C, Figure S1)**. The
wild-type RBM-26::3XFLAG protein is predicted to produce proteins with molecular
weights around 75 and 82 KD. Consistent with this prediction, we found that the RBM-
26::3XFLAG protein was detected as two bands that migrate in between the 72KD and
95KD markers. In addition, we also observed a more intense bands just above the
56KD marker and just above the 95 KD marker as well as two fainter bands in between
the 34KD and 56KD markers. The identity of these additional bands are unknown,
though it they are likely the result of post-translational processing, or unpredicted splice
forms. We found that each of these RBM-26::3XFLAG bands were sharply reduced by
the P80L and L13V mutations **(Figure 1C,D)**. To determine if the P80L mutation can
reduce expression of RBM-26 in vivo, we used an endogenously tagged *rbm-26::scarlet*

gene **(Figure 1E,F)**. Consistent with our western blotting data, we found that the P80L
mutation substantially reduces expression of the RBM-26::Scarlet protein in vivo. These
observations indicate that the P80L and L13V mutations in *C. elegans rbm-26* sharply
reduce RBM-26 expression and imply that the P79L and L13V mutations in human
RBM27 are likely to be gene-disrupting.

Mutations in *rbm-26* cause developmental defects in the PLM axon.

To determine if RBM-26 can affect neuronal development, we used the PLM, a neuron
that has its cell body in the tail and extends its axon anteriorly along the body wall. The
PLM has three presynaptic regions and can also act as a sensor for light touch [26]. In
wild type, the PLM axon initially overlaps with the ALM axon during the L1 stage, but
this overlap is resolved in later larval stages [26].

To determine how loss of RBM-26 function affects the PLM, we used an *rbm-26(null)*
mutation, along with *rbm-26(P80L)* and *rbm-26(L13V)* mutations. The *rbm-26(gk910)*
null mutation consists of a 408 base pair deletion that removes part of exon 4 and exon
5 of *rbm-26* and is lethal. We maintained this *rbm-26(null)* allele over a balancer and
found that maternally rescued homozygote progeny are viable until the L3 stage. To
study the P80L missense mutation, we used CRISPR to create the *rbm-26(cue23)*
allele, hereafter referred to as *rbm-26(P80L)*. We also introduced the L13V missense
mutation into the *rbm-26* gene, creating the *rbm-26(cue34)* mutation, hereafter referred
to as *rbm-26(L13V)*. In contrast to the *rbm-26(null)* mutation, both the *rbm-26(P80L)* and
the *rbm-26(L13V)* mutations are viable.

We found that mutations in *rbm-26* cause overlap between the PLM and ALM axons,
hereafter referred to as the PLM/ALM overlap phenotype **(Figure 2A-D)**. In *rbm-26(null)*
mutants, a greater penetrance of PLM/ALM overlap was observed in the L1, L2 and L3
stages relative to wild type. Due to the lethal phenotype of the *rbm-26(null)* mutation, we
were unable to analyze PLM axons in in later stages. In the P80L and L13V mutants,
we also observed a greater penetrance of PLM/ALM overlap in the L2 and L3 stages
relative to wild type. However, we did not observe PLM/ALM overlap at later stages in
these missense mutations, suggesting that that these tiling defects resolve at later
developmental stages **(Figure 2D)**. Analysis of the PLM axon length to body length ratio
suggests that the PLM/ALM overlap phenotype in these three *rbm-26* mutants is caused
by overextension of the PLM axon **(Figure S2)**.

Mutations in *rbm-26* cause degeneration phenotypes in the PLM axon.

We found that the *rbm-26(null)*, *rbm-26(P80L)*, and the *rbm-26(L13V)* mutations all
cause degeneration phenotypes in the PLM axon **(Figures 2E-H and S3; Table S1)**. In
the *rbm-26(null)* mutants, we observed a very weak axon beading phenotype in the L2
stage that became much stronger in the L3 stage **(Figure 2F)**. The *rbm-26(P80L)*
mutation caused an axon beading phenotype that was very weak in the L2 stage,
increased in the L3 and L4 stages and became much stronger in the adult stage. We
also observed the beading phenotype in *rbm-26(L13V)* mutants, but at a substantially
lower penetrance relative to *rbm-26(P80L)*. In addition to beading we observed several
other degeneration phenotypes **(Table S1)**, including axon breaks **(Figure 2G,H)**. To
determine if loss of *rbm-26* function might cause defects in other neurons, we also
observed the PVD neuron and observed signs of degeneration in both its axon and
dendrites **(Figure S4)**. These degenerative phenotypes in PLM and PVD are similar to
those that have been reported in aging worms [27-29] and are reminiscent of
axonopathy that has been reported in human neurodegenerative disorders and
described as axonal swellings or spheroids [30-32]. Overall, these observations suggest
that RBM-26 protects against axon degeneration. Moreover, these results imply that the
P79L and L13V de novo missense mutations in human *RBM27* are likely to disrupt
RBM27 function, thereby supporting a role for this gene in neurodevelopmental
disorders.

**RBM-26 is expressed in multiple tissues and functions within neurons to protect
against axonal defects.**

To visualize the localization of RBM-26, we used CRISPR to insert the coding sequence
for a Scarlet::AID (auxin-inducible degron) tag into the 3′ end of the *rbm-26* gene,
thereby creating the *rbm-26(*syb2552) allele. We found that this allele expresses RBM-
26::Scarlet::AID protein in many tissues including neurons, hypodermis, muscle, and
intestine **(Figures 3A-C, S5, and S6)**. Within hypodermis and neurons, we noticed that
the RBM-26::Scarlet::AID protein was localized to the nucleus **(Figures 3A-C and S5)**.

We next asked where RBM-26 functions to protect against axon degeneration and
defects in axon development. For our initial approach to this question, we used the
auxin-inducible degron system to selectively degrade the RBM-26::Scarlet::AID protein
in in neurons, hypodermis, muscle, and intestine [33]. We found that degradation of
RBM-26::Scarlet::AID in neurons caused both PLM/ALM overlap defects **(Figure 3D)**
and the PLM beading phenotype **(Fig. 3E)**. However, degradation of RBM-
26::Scarlet::AID in hypodermis, muscle, or intestine failed to cause any of these

phenotypes **(Figures 3D-E)**. These phenotypes were dependent on auxin and the *rbm-*
26::scarlet::AID gene **(Figures S7 and S8)**. We also note that both of these phenotypes
occurred at a lower penetrance in these degradation experiments relative to
experiments with the *rbm-26* mutant alleles, suggesting incomplete degradation of the
RBM-26::Scarlet::AID protein.

To further test the locus of RBM-26 function, we conducted transgenic rescue
experiments. For the transgenic rescue experiments, we used a *Pmec-7::rbm-26*
transgene to selectively drive expression of RBM-26 in the touch receptor neurons
(Figures 3F-G). We found that this *Pmec-7::rbm-26* transgene could rescue the
PLM/ALM axon overlap phenotype in both the *rbm-26(P80L)* mutants and in the *rbm-*
26(L13V) mutants **(Figure 3F)**. Likewise, the *Pmec-7::rbm-26* transgene could also
rescue the beading phenotype in both the *rbm-26(P80L)* mutants and in the *rbm-*
26(L13V) mutants **(Figure 3G)**.These observations suggest that RBM-26 is broadly
expressed and that it functions cell-autonomously within neurons to protect against
axon degeneration and defects in axon development.

Loss of RBM-26 causes mitochondrial dysfunction.

Mitochondrial dysfunction is thought to be an early causative factor for neuronal
degeneration. Therefore, we asked if loss of RBM-26 function affects mitochondria in
the PLM. We used the third larval (L3) stage for these experiments, because the *rbm-*
26(null) mutants die after this stage. We found that mitochondria were localized
throughout wild-type PLM axons, with an average density of 9 per 100μm **(Fig. 4A,B,E)**.
The *rbm-26(null)* mutation caused a reduction of the mitochondrial density in the PLM
axon to about about 6 per 100μm **(Fig. 4C-E)**. Likewise, the *rbm-26(P80L)* and the *rbm-*
26(L13V) mutations also caused a reduction of the mitochondrial density in the PLM
axon **(Fig. 4E)**.To determine the timing of this phenotype, we analyzed *rbm-26(null)*
mutants at earlier larval stages and found that the phenotype begins in the L1 stage and
persists through the L3 stage, when these mutants die **(Figure 3F)**. We also found that
all three *rbm-26* alleles caused a reduction in the number of mitochondria in the
proximal axon (50 μM closest to the cell body; see Figure 4G). However, none of the
rbm-26 alleles affected the number of mitochondria in the distal axon (50 μM closest to
the axon tip; see Figure 4H). These observations suggest that RBM-26 is needed to
maintain a normal density of mitochondria in the proximal PLM axon. We speculate that
mitochondria in the proximal PLM axon might be more susceptible to damage because
of the higher level of axon transport that occurs in the proximal PLM axon [34].

Excess production of reactive oxidation species (ROS) is a key marker for mitochondrial
dysfunction [35, 36]. Therefore, we asked if the *rbm-26(null)* mutation causes an
increase in ROS production by mitochondria. For this experiment, we used mitoTimer,
an oxidation sensor that is tethered to mitochondria and changes irreversibly from green
to red as it becomes oxidized [37]. We analyzed the mitoTimer red:green ratio in the
PLM cell body because the mitoTImer signal in axons was not sufficient for quantitation.
We found that the *rbm-26(null)*, *rbm-26(P80L)*, and *rbm-26(L13V)* mutations all increase
the mitoTimer red:green ratio in the PLM cell body relative to wild type **(Figure 5A-E**
and Figure S9), suggesting that loss of RBM-26 function causes mitochondria to
produce excessive ROS and that RBM-26 is needed to maintain normal mitochondrial
function in neurons. Overall, these results suggest that loss of RBM-26 causes
dysfunction of mitochondria in the neuronal cell body, which could lead to a decrease in
mitochondria density in the axon.

RBM-26 binds to the *mals-1* mRNA and negatively regulates its expression.

To understand the molecular mechanisms of RBM-26 function, we conducted a screen
to identify RNAs that bind to RBM-26. For this screen, we used ultraviolet light to induce
RNA-protein crosslinking in strains expressing RBM-26::3XFLAG and in controls
expressing GFP::3XFLAG and Scarlet::3XFLAG (see methods for details). We then
used an anti-FLAG antibody to purify the RBM-26::3XFLAG::RNA and
Scarlet/GFP::3XFLAG::RNA complexes, followed by RNA-Seq analysis. We identified
63 RNAs that were significantly enriched in RBM-26::3XFLAG pulldowns relative to
Scarlet/GFP::3XFLAG controls **(Figure 6A and Table S2)**. Among these 63 RNAs we
found 26 protein-coding mRNAs, 14 non-coding RNAs, 8 piRNAs, 10 snoRNAs, 1 tRNA
and 1rRNA.

We focused on the *mals-1* mRNA because it had, by far, the lowest adjusted p-value
amongst the RBM-26-binding partners with human orthologs **(Table S2)**. The *mals-1*
mRNA encodes an ortholog of the mammalian MALSU1 mitochondrial ribosome
assembly factor [38]. Using qPCR, we found that loss of RBM-26 function causes an
increase in *mals-1* mRNA expression in *rbm-26(P80L)* and *rbm-26(L13V)* mutants
relative to wildtype **(Figure 6B)**. To analyze MALS-1 protein expression, we used
CRISPR to add a 3XFLAG tag to MALS-1, thereby creating the *mals-1(cue37)* allele,
which expresses the MALS-1::3XFLAG protein. Since the *mals-1* gene is tightly linked
to the *rbm-26* gene, we used CRISPR to introduce the P80L mutation into a strain
containing the *mals-1(cue37)* mutation, thus creating the *mals-1(cue37) rbm-26(cue42*
[P80L]) double mutant. By performing western blotting to analyze expression of MALS-
1::3XFLAG in these strains we found that the *rbm-26(P80L)* mutation substantially

enhances expression of the MALS-1::3XFLAG protein relative to wild type **(Figure 6C-**
D). We also obtained similar results with the *rbm-26(L13V)* mutation. These
observations indicate that RBM-26 negatively regulates the expression of *mals-1* mRNA
and MALS-1 protein.

Overexpression of MALS-1 phenocopies *rbm-26* loss of function.

Since we found that loss of RBM-26 function causes an increase in MALS-1 expression,
we next asked if overexpression of MALS-1 could cause axonal phenotypes. For this
experiment, we created the *cueSi36 IV* transgene, hereafter called *Pmec-7::mals-*
1::scarlet to overexpress MALS-1 in the PLM neuron. Consistent with work on
mammalian MALSU1, we found that this overexpressed MALS-1::Scarlet is localized to
mitochondria **(Figure 7A-C)**. We found that overexpression of MALS-1 causes an
increase in PLM/ALM overlap defects that are significantly greater than those observed
in controls **(Figure 7D)**. Overexpression of MALS-1 also caused the axon beading
phenotype **(Figure 7E)**. Finally, we found that overexpression of MALS-1 decreases the
average density of mitochondria in the PLM axon **(Figure 7F)**. These observations
suggest that overexpression of MALS-1 can cause defects that are similar to those
caused by loss of RBM-26 function.

MALS-1 is required for phenotypes caused by loss of RBM-26 function.

Since loss of *rbm-26* function causes axonal phenotypes and an increase in MALS-1
expression, we hypothesized that MALS-1 expression is causative for the axonal
phenotypes. To test this hypothesis, we asked if *mals-1* is required for the *rbm-26*
mutant phenotypes. For these experiments, we used the *mals-1(syb6330)* allele, a near
complete deletion of the *mals-1* coding sequence. We also used the *mals-1(tm12122)*
mutation, which is a partial deletion of the *mals-1* coding sequence that is predicted to
cause early termination of the MALS-1 protein.

To determine if *mals-1* is required for the *rbm-26* mutant phenotypes, we constructed
double mutants between alleles of *mals-1* and *rbm-26*. If the *rbm-26* loss of function
phenotypes are dependent on *mals-1*, we would expect that the *rbm-26* mutant
phenotypes would be suppressed in *rbm-26 mals-1* double mutants relative to *rbm-26*
single mutants.

Since the *mals-1* and *rbm-26* genes are tightly linked, we used CRISPR to introduce the
P80L mutation into a strain containing the *mals-1(syb6330)* mutation, thereby creating
the *rbm-26(cue40 [P80L]) mals-1(syb6330)* double mutant. We found that *rbm-*
26(cue40 [P80L]) mals-1(syb6330) double mutants have significantly lower penetrance
of axon beading relative to *rbm-26(P80L)* single mutants **(Figure 8A)**. Likewise, *rbm-*
26(cue40 [P80L]) mals-1(syb6330) double mutants also have significantly higher
mitochondrial density relative to *rbm-26(P80L)* single mutants **(Figure 8B)**. For the
PLM/ALM overlap phenotype, we would not expect to see a suppression of the *rbm-*
26(P80L) phenotype by *mals-1(syb6330)* because *mals-1(syb6330)* single mutants
already have this phenotype at a penetrance equal to *rbm-26(P80L)* mutants. However,
we did find that *rbm-26(cue40 [P80L]) mals-1(syb6330)* double mutants had a
penetrance of axon termination defects that was equal to *mals-1(syb6330)* single
mutants **(Figure 8C)**, consistent with the function of *mals-1* and *rbm-26* in a genetic
pathway. To further validate these findings, we also repeated these experiments with an
independent allele of *mals-1*, *mals-1(tm12122)* and found similar results **(Figure 8A-C)**.
We noticed that this *mals-1(tm12122)* allele had slightly higher penetrance of defects
relative to the *mals-1(syb6330)* mutant and speculate that this might be the result of an
incomplete deletion of the sequence coding for MALS-1 in the *syb12122* allele. Taken
together, these observations indicate that the axonal phenotypes of *rbm-26(P80L)* and
rbm-26(L13V) mutants are dependent on MALS-1 function. Moreover, these
observations are consistent with the hypothesis that RBM-26 prevents axon
degeneration by negatively regulating expression of MALS-1.

We also conducted western blotting as an alternative way to test the interaction
between RBM-26, MALS-1, and mitochondria **(Figure 8D-E)**. For this experiment, we
used the MRPL-58 mitiribosomal protein as a marker for mitochondria. We found that
the *rbm-26(P80L)* mutation causes a significant reduction in amount of MRPL-58
protein, consistent with a reduction in the number of mitochondria. We also found that
this reduction of MRPL-58 expression was suppressed in *rbm-26(P80L) mals-*
1(syb6330) double mutants. These observations are consistent with the idea that loss of
rbm-26 causes a reduction in mitochondria that is caused by excessive expression of
MALS-1.

Discussion

Here, we report that RBM-26 functions in neurons to regulate expression of the MALS-1
mitiribosome assembly factor. RBM-26 binds to the *mals-1* mRNA and sharply reduces
its expression. Disruptions in this process cause excessive MALS-1 expression, leading
to the disruption of mitochondrial function in the developing neuron. These observations

provide an example of how post-transcriptional regulation of an mRNA in the nucleus
can protect mitochondria in the axon to prevent axon degeneration during neuronal
development.

We also note that the degenerative and developmental defects caused by loss of *rbm-*
26 function in *C. elegans* larvae are reminiscent of defects in human infants who carry
mutations in various components of the RNA exosome [8-10, 39, 40]. Prior work in a
human cell line suggests that the RNA exosome functions with the RBM-26/27
orthologs of RBM-26 to degrade RNA [17]. Therefore, we speculate that human
RBM26/27 could function with the RNA exosome complex to protect against
neurodevelopmental defects and axon degeneration in infants.

Relationship between axon development defects and axon degeneration
phenotypes.

We have found that RBM-26 causes defects in both axon development and axon
degeneration. This is interesting, because the RBM26/27 orthologs of RBM-26 are
thought to function with the RNA exosome [17] and variants in the genes that encode
RNA exosome components have been implicated in neurodevelopmental syndromes
that also cause defects in both axon development and axon degeneration in children [8-
10, 39, 40] (see below for further discussion on the RNA exosome).

Axon development defects occur prior to axon degeneration defects in *rbm-26* mutants.
Loss of RBM-26 function causes a reduction in mitochondria density that begins during
early larval development, coincident with the start of the PLM/ALM overlap phenotype.
On the other hand, mitochondrial oxidation is minimal in early larval stages and peaks in
later larval stages, coincident with the axon degeneration phenotypes. Overall, these
observations suggest that the RBM-26 mutations begin to affect mitochondria in early
larval development, giving rise to defects in axon development. In later larval stages,
the mitochondrial damage becomes substantially worse, giving rise to axon
degeneration.

Axon tiling defects are transient in the P80L and L13V missense mutants. The
hypomorphic P80L and L13V missense mutations in RBM-26 cause axon tiling defects
that start in the early larval stages and are resolved by the 4th larval stage. By contrast,
the *rbm-26* null mutation produces the axon tiling defect at a higher penetrance and this
phenotype is not resolved. However, the null mutants die prior to the L4 stage, so we
are unable to determine if the phenotype would resolve in later stages. We note that in a
prior study, we found that neurodevelopmental disease-associated missense mutations

in *unc-16* also cause axon tiling defects that are resolved in later stages in *C. elegans*
despite causing severe intellectual disability in humans [41]. In this case, *unc-16* null
mutations are viable and produce an axon tiling phenotype that persists throughout the
life of the worm. On the other hand, a missense mutation in the *egl-19* ortholog of
CACNA1C produces phenotypes that persist throughout the worm's life [42, 43].
Overall, these observations suggest that some, but not all, neurodevelopmental patient-
derived mutations can have transient effects in *C. elegans*.

406 Emerging evidence for the role of the nuclear RNA exosome in the regulation of
407 neurodevelopment and neurodegeneration.

The negative regulation of *mals-1* mRNA by RBM-26 may occur through the process of
mRNA decay in the nucleus. This idea is supported by our finding that RBM-26 is
localized to the nucleus and negatively regulates *mals-1* mRNA. Moreover, prior work in
human cells has indicated that RBM26/27 functions with the PAXT complex, an adaptor
that delivers polyadenylated RNA to the nuclear RNA exosome for degradation [17]. A
similar complex exists in *S. pombe*, where the Rmn1 protein is an ortholog of RBM-26
and functions with the MTREC complex, which is orthologous to the PAXT complex
[18]. Thus, RBM-26 is part of a highly conserved complex that collaborates with the
nuclear RNA exosome to negatively regulate polyadenylated RNA.

Although most work on the nuclear RNA exosome complex has focused on non-coding
nuclear RNAs, emerging evidence suggests that this complex is also important for the
regulation of some mRNAs. For example, in *S. pombe*, the MTREC complex is
orthologous to the PAXT complex and is critical for the suppression of meiosis. In this
process, the MTREC complex binds to the mRNAs of meiotic genes and delivers them
to the nuclear RNA exosome for degradation [44]. In cultured human cells, the *PABPN1*
mRNA is degraded by the nuclear RNA exosome, thereby negatively regulating both the
PABPN1 mRNA and the PABPN protein [45]. In addition, knockdown of RBM26/27 or
other members of the PAXT complex in a human cell line can cause upregulation of
more than 1000 mRNAs, suggesting that the PAXT complex is able to regulate mRNAs
[17]. Consistent with this finding, a group of 360 full length protein-coding mRNAs were
identified that are regulated by the nuclear mRNA exosome in a human cell line [46].
Together, these observations indicate that the PAXT complex can function with the
nuclear RNA exosome to regulate some protein-coding mRNA transcripts.

The axon degeneration and axon development phenotypes caused by *rbm-26*
mutations are reminiscent of phenotypes caused by mutations in subunits of the human
RNA exosome complex. For example, mutations in the *EXOSC3*, *EXOSC8* and
EXOSC9 are thought to cause syndromes that include defects in brain development
such as hypoplasia of the cerebellum and the corpus callosum [8-10, 39, 40]. In

addition, these syndromes also involve progressive neurodegeneration of spinal
neurons that begins in infancy and resembles spinal motor atrophy (SMA). Consistent
with these observations, knockdown of EXOSC3, EXOSC8, and EXOSC9 in zebrafish
causes brain malformation and axon pathfinding defects [8, 39, 47]. Moreover,
mutations in EXOSC3 and EXOSC9 are also associated with intellectual disability in
humans [10, 40] and mutations in EXOSC8 causes behavioral defects in zebrafish [39].
Together, these observations suggest an important role in for the RNA exosome
complex in protecting against neurodegeneration and preventing neurodevelopmental
defects.

Potential role for MALS-1 overexpression in the disruption of mitoribosome biogenesis.

The biogenesis of mitoribosomes is a complex process that involves the assembly of 80
proteins and 2 rRNAs into a large subunit (mtLSU) and a small subunit (mtSSU). This
process is facilitated by a large number of mitoribosomal assembly factors including
RNA processing enzymes, GTPases, RNA helicases, and kinases. MALS-1 is an
ortholog of the human MALSU1 mitoribosomal assembly factor that functions as part of
the MALSU1:LOR8F8:mtACP anti-association module [48-50]. This module binds to the
mtLSU and blocks premature association of mtLSU with mtSSU. Once assembly of
mtLSU is complete, MALSU1:LOR8F8:mtACP is released, thereby allowing association
of mtLSU with mtSSU. Consistent with these observations, knockdown of MALSU1 in
cultured human cells causes mitochondrial dysfunction by disrupting the formation of
mitoribosomes [38].

Our results suggest that excessive expression of MALS-1 can also disrupt mitochondria.
We propose that this disruption of mitochondria occurs because excess MALS-1 can
disrupt mitoribosome assembly. This idea is also supported by our finding that loss of
rbm-26 function causes a very high level of MALS-1, the ortholog of MALSU1, a
negative regulator of mitoribosome assembly. Moreover, the axon and mitochondrial
defects caused by loss of RBM-26 function are dependent on MALS-1 expression. In
humans, under normal conditions, MALSU1 binds to mtLSU and prevents it from
prematurely assembling with mtSSU [48-50]. After completion of mtLSU assembly,
MALSU1 is released, thereby allowing for binding of mtLSU and mtSSU. Extrapolating
from the function of MALSU1 in humans, we hypothesize that the high level of MALS-1
expression in *rbm-26* mutants may prevent the dissociation of MALS-1 from the mtLSU,
thereby interfering with the association between mtLSU and mtSSU.

Although our work suggests that MALS-1 overexpression can cause PLM axon defects,
we also observe weaker defects with *mals-1* loss of function mutants. Consistent with
this finding, prior work has indicated that knockdown of MALSU1 disrupts mitoribosome
assembly and function in human cells [38]. This observation reflects the critical role of
MALSU1 in the assembly of the mitoribosomes. Thus, mitochondria can be disrupted by
either MALS-1 overexpression or MALS-1 loss of function. We note that the *rbm-26* loss
of function defects are substantially stronger than the *mals-1* loss of function defects.
Therefore, we propose that the disruption of post transcriptional regulation of
mitoribosomal assembly factors may represent a key vulnerability for mitoribosome-
associated disease.

Materials and Methods

482 **Materials and Methods**

483
Strains: The alleles used in this study are listed in **Table S3**. The *rbm-*
26(gk910) mutation is lethal and was maintained over the hT2 balancer marked with *bli-*
4 and *(qIs48)*. Worms used for phenotype analysis were maintained at 20°C on
nematode growth medium (NGM) agar plates seeded with OP50 using standard
procedures. Worms used for RNA Immunoprecipitation experiments and Western
Blotting were cultured on 8P plates seeded with *E. coli* NA22 strain as food source.

490
491 **Genome editing:** Genome editing for *mals-1(syb6330)* and *rbm-26(*syb2552 [*rbm-*
492 *26::scarlet::AID]*) were performed by SunyBiotech. All other genome editing was
493 performed in our laboratory using a previously described protocol [51, 52]. Guide RNAs
494 and homology directed repair (HDR) templates are listed in Table S4. CRISPR edits
495 were confirmed via PCR and sequencing.

496
Transgenes: The *cueSi36 IV* transgene encodes *Pmec7::mals-1::scarlet::tbb-2 3'UTR*,
cueSi52 IV transgene encodes *Pmec-7::rbm-26::scarlet::unc-54 3'UTR*, and the
cueSi35 IV transgene encodes *Pmec-7:mitoTimer::tbb-2 3'UTR*. A single copy of these
transgenes was inserted into the *jsTi1493* landing pad on chromosome IV using the Flp
recombinase mediated cassette exchange (RMCE) protocol [53]. The *muIs32* transgene
is an integrated multicopy array that encodes *Pmec7::GFP* and was used to visualize
the PLM axon. The *jsIs1073* transgene was obtained from Michael Nonet and encodes
Pmec-7::mts:tagRFP and expresses mitochondria-targeted TagRFP (mitoTagRFP). The
jsIs609 transgene was also obtained from Michael Nonet and encodes *Pmec-*
7::mts::GFP and expresses a mitochondria-targeted GFP (mitoGFP) in the PLM neuron.

507
Analysis of phenotypes: To quantify phenotypes, worms were mounted on a 5%
agarose pad, anesthetized with levamisole, and observed with a 40x objective on a

Zeiss Axio Imager M2 microscope as previously described [54-57]. For all experiments
we observed phenotypes in the L3 stage, since the *rbm-26(null)* mutants don't survive
beyond this stage. For many experiments, we also included observations at various
other stages from L1 to adult. PLM axons were visualized using the *muIs32 (Pmec-*
7::GFP) transgene. The PLM/ALM overlap phenotype was scored when the PLM axon
tip terminated anteriorly to the ALM cell body. We also captured DIC images of worms,
fluorescence images of PLM in the same worms and used ImageJ to quantify the length
of the worm and corresponding PLMs. Beading refers to focal enlargement or bubble-
like lesions which were at least twice the diameter of the axon in size as shown in
Figure S3.
520
Imaging: Larval stage images displayed in all figures were obtained by mounting on 5%
agarose pad, anesthetizing with levamisole, and imaging with a 60x water objective on
a Nikon ECLIPSE Ti microscope equipped with the X-Light V2 L-FOV spinning disk
system as previously described [58]. NIS-Elements software was used for image
acquisition. Image analysis was done with ImageJ or NIS-elements. Identical imaging
settings and look-up-tables were used for control and experimental groups. Imaging for
mitochondrial counts was done with a 40X oil objective on a Zeiss Axio Imager M2.
528
Analysis of mitoTimer: All images were captured with identical settings on a
spinning disk confocal system (see above). 8-bit images were opened in ImageJ in both
red and green channels, the PLM cell body was traced in one channel with the free
hand selection tool, and the tracing was restored in the other channel. Raw Integrated
Density (RawIntDen) and the area of the region of interest (ROI) was determined in
ImageJ. Similarly, RawIntDen and area of a defined background region was obtained.
RawIntDen was divided by the area of ROI to obtain total pixel/area. Corrected total
pixel/area was obtained by subtracting the background pixel/area from the ROI for both
red and green channels.
538
Auxin mediated protein degradation: Tissue-specific degradation of RBM-
26::Scarlet::AID was carried out using the auxin-induced degron (AID) system as
previously described [58]. Briefly, Naphthaleneacetic acid (K-NAA) (Phytotechnology
Laboratories) was dissolved in double distilled water to obtain 250 mM K-NAA solution
and filter-sterilized by passing through 25 mm sterile syringe filter (Pall Corporation).
Aliquots from sterile 250 mM K-NAA stock was added to autoclaved and slightly cooled
(60°C) NGM to obtain a concentration of 4 mM K-NAA and poured into plates. The next
day, 4mM K-NAA plates were seeded with 250 ul *E. coli* OP50 strain. These plates
were used within two weeks.
To obtain tissue specific degradation, homozygous lines were obtained by crossing
rbm-26(syb2552[rbm-26::Scarlet::AID]) with the following strains: *reSi1 [col-*
10p::TIR1::F2A::mTagBFP2::AID*::NLS::tbb-2 3'UTR] (for hypodermal degradation),
reSi7 [rgef-1p::TIR1::F2A::mTagBFP2::AID*::NLS::tbb-2 3'UTR] (for neuronal
degradation), *reSi3 [unc-54p::TIR1::F2A::mTagBFP2::AID*::NLS::tbb-2 3'UTR]* (for

degradation in muscles), and *reSi12 [ges-1p::TIR1::F2A::mTagBFP2::AID*::NLS::tbb-2*
3'UTR] (for degradation in intestine).
L4 stage *C. elegans* of different genotypes were placed on 4mM K-NAA plates, allowed
to grow and lay eggs, L3 stage progenies of the parents initially placed on 4mM K-NAA
plates were analyzed for PLM termination defects and PLM beading phenotypes.
Knockdown of the RBM-26::Scarlet::AID protein was verified visually. PLMs were
visualized with *Pmec-7::GFP* transgene. *rbm-26::Scarlet::AID* without any TIR1 cofactor
of AID made up the control group. Knockdown of the RBM-26::Scarlet::AID protein in
the PLM was quantified in the same manner as mitoTimer (Figure S10). To elaborate,
ImageJ was used to trace PLM in the green channel and the tracing was restored in the
red channel. The RawIntDen was recorded for both the channels and subtracted from
the background.
565
UV-crosslinking RNA Immunoprecipitation and RNA-Seq: Worms expressing
wbmIs67 transgene *(eft-3p::3X FLAG::Scarlet::unc-54 3'UTR)* and *wbmIs72* transgene
(pei-1p::3X FLAG::GFP::unc-54 3'UTR) were used as control group. Control and *rbm-*
26::3X FLAG worms were grown on NGM plates seeded with *E. coli* OP50 strain until
they became gravid, then they were washed in M9 (22mM KH2PO4; 22mM Na2HPO4;
85mM NaCl; 1mM MgSO4) and grown for 2 days on 8P plates seeded with *E. coli* strain
NA22. 25,000 mixed stages *C. elegans* were washed three times in M9 and placed in
an unseeded NGM plate (10 cm) for UV-crosslinking. Plate covers and liquid were
removed, and the NGM plates were placed in Spectrolinker X-1000 (Spectronics
Corporation) and irradiated with 254 nm UV at energy setting 3000. Worms were then
checked under the microscope to ensure that they were immobilized, following which
they were washed in M9, and 100,000 worms were collected in 1 ml of RIPA buffer (50
mM Tris HCl pH 7.4, 150 mM NaCl, 1% Nonidet NP-40, 0.1% SDS, 0.5% Sodium
deoxycholate, Roche cOmplete EDTA-free Protease Inhibitor Cocktail, 25U/ml RNase
inhibitor) and homogenized/lysed in BeadBug™3 Microtube homogenizer (40 second
homogenization with 1 min rest on ice) at 4°C with 3.2 mm diameter chrome steel beads
(8 beads per tube). Lysates were placed in fresh tubes, centrifuged at 16,000 x g for 15
minutes at 4°C. The supernatant was saved as lysate and quantified using Pierce BCA
Protein Assay kit. 100 ul of Pierce Anti-DYKDDDDK (anti-FLAG) magnetic agarose
resin was washed three times in RIPA buffer and added to 1 ml of protein lysate (2
mg/ml), incubated overnight at 4°C, and anti-flag agarose bound lysate was separated
with the help of a magnet while the supernatant was discarded. The agarose resin was
washed once with high salt buffer (50 mM Tris HCl pH 7.4, 1M NaCl, 1 mM EDTA, 1%
Nonidet NP-40, 0.1% SDS, 0.5% Sodium deoxycholate, Roche cOmplete EDTA-free
Protease Inhibitor Cocktail and RNase Inhibitor) and three times with low salt buffer (20
mM Tris HCl pH 7.4, 10mM MgCl2, 0.2% Tween-20, Roche Protease Inhibitor and
RNase Inhibitor). To elute RNA, the agarose resin was dissolved in Proteinase K buffer
(50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 1 mM MgCl2, 0.05% NP-40, 1% SDS) and
incubated with 1.2 mg/ml Proteinase K at 55°C for 30 mins with gentle agitation. RNA
was extracted with phenol:chloroform and precipitated with 0.25M Ammonium Acetate,

0.12M Lithium Chloride and 85% Ethanol. The pellets were washed with 75% ethanol,
dissolved in DEPC-treated water and sent to Azenta US, Inc. for library preparation and
RNA sequencing. Raw data obtained from Azenta US, Inc. was analyzed to identify
RNA interactors of RBM-26. For RNA-Sequencing pre-processing, adapters were
trimmed using Cutadapt [59], and contaminating rRNA reads were removed using
bowtie2 [60, 61]. Reads were mapped to the Caenorhabditis elegans transcriptome
(Caenorhabditis_elegans.WBcel235) using STAR [62] and differential gene analysis
was performed using the DESeq2 package [63] in R Studio. Mapped reads were sorted
and indexed using samtools [64].
605
Absolute quantification of *mals-1* mRNA: 300 N2 wild-type L4 worms and 300 *rbm-*
26 mutants [*rbm-26(cue23)* and *rbm-26(cue34)*] L4 worms were collected in Trizol.
Following 5 rounds of freeze thaw on dry ice with 1 min of vortexing during thawing,
RNA was extracted as per the manufacturer's protocol.
We purchased the following synthetic RNA from Azenta US, Inc:
GCACGAAGGAGCACAAAACGAAGCAAUGGAUGGUAUGUCAGUGAGGUGGAAAGA
GUUCAGGUGCACG The synthetic RNA was diluted in nuclease free water and the
following concentrations were confirmed with Qubit RNA Broad Range Assay Kit
(Thermo Fisher Scientific): 10 ng/ul, 1 ng/ul, 0.1 ng/ul, 0.01 ng/ul, 0.001 ng/ul, and
0.0001 ng/ul. The diluted RNA standards and 1 ug of total RNA extracted from wildtype
and *rbm-26* mutant L4 worms were reverse transcribed with 2µm reverse primer
(CGTGCACCTGAACTCTTTCC) and SuperscriptTM First-Strand Synthesis System
(Thermo Fisher Scientific) following the manufacturer's protocol. 1µl cDNA was used
with 10µl 2x SsoAdvanced Universal SYBR Green Supermix (Biorad) and 5 pmol of the
forward *mals-1* primer (TGCACGAAGGAGCACAAAAC) and the reverse *mals-1* primer
(CGTGCACCTGAACTCTTTCC). Fluorescence emitted during each cycle was
monitored using the CFX96 Touch Real-Timer PCR Detection System (Biorad).
Dissociation curve confirmed the presence of a single amplicon. Cycle-to-Threshold (Ct)
value of the RNA standards were used to plot a standard curve (R^2 = 0.995) (Figure
S11), which was then used to deduce mals-1 mRNA quantity. Data presented here
comes from three biological replicates.
627
Western Blotting: Worms were grown on NGM plates seeded with *E. coli OP50* until
gravid, washed in M9, and plated onto 8P plates seeded with *E. coli NA22*. 50,000
worms were collected in M9, washed three times and M9 was replaced with RIPA buffer
(50,000 worms/1 ml RIPA buffer). Worms were lysed in Bead Bug3TM Microtube
homogenizer at 4°C with 3.2 mm chrome steel beads (8 beads/ml RIPA buffer).
Following centrifugation at 16000 xg for 15 mins at 4°C, supernatant was saved
(snapfrozen) as total protein lysate and snap frozen. Total protein lysates were
quantified through BCA assay. 20 – 25 µg of total protein lysate was run on 12% SDS-
PAGE, transferred to nitrocellulose membrane (0.2 µm pore). For some experiments, 40
µg of total protein lysate was used. The nitrocellulose membrane was incubated with
No-stainTM protein labeling reagent for 10 minutes following the manufacturer's

instructions. The membrane was washed 2X in water and 3X in TBST (20 mM Tris, 150
mM NaCl, 0.1 % Tween-20). Then, the membrane was blocked in 5% non-fat milk in
TBST, probed with primary antibody, washed 3X in TBST, probed with secondary
antibody, washed in 3X in TBST, developed with Super Signal West PICO Plus
(Thermo Fisher Scientific) and imaged with iBright CL1500 (Thermo Fisher Scientific).
The iBright CL 1500 on-instrument analysis software and No-Stain Labeled Membrane
setting was used for normalization. iBright analysis software was used for band intensity
quantification, which was reevaluated with ImageJ. Antibodies used in this study are
listed in supplemental table S4.
648
649
650
651 **Acknowledgements**
652
We thank Michael Nonet, Shohei Mitani, and the Caenorhabditis Genetics Center
(funded by NIH P40 OD010440) for strains. We also thank Claire de la Cova and Mark
McBride for sharing equipment, Claire de la Cova for assistance with imaging
experiments and Scott Aoki for help with the UV-RIPseq experiments. This work was
funded by the National Institute of Mental Health grant R01MH119157 (to CCQ), the
National Institute of Neurological Disorders and Stroke grant R03NS101524 (to CCQ),
Discovery and Innovation Grant (101X432) from the University of Wisconsin -
Milwaukee (to TAC) and National Institutes of Health grant 1R35GM142527 (to ASYL).
This article does not represent the official views of the National Institutes of Health and
the authors bear sole responsibility for its content.
663
664
665
**Figure 1: Expression of RBM-26 is disrupted by P80L and L13V missense
mutations**. (A) Putative protein domains in human RBM27, and *C. elegans* RBM-26
proteins predicted by NCBI's CD-Search. CCCH-type Zn Finger domains (Green
Rectangles) and RNA Recognition Motifs (Gold Pentagons) were predicted with high
confidence by NCBI's CD-Search and SMART database. PWI superfamily (Blue
Hexagon) was predicted by NCBI's CD-Search with high confidence. Red arrows
indicate de novo missense variants identified in patients with autism or other
neurodevelopmental disorders. (B) Clustal Omega alignment of the N-terminus of the
human RBM27 and *C. elegans* RBM-26 proteins. Two de novo RBM27 variants, L13V
and P79L (red arrows), were found in regions conserved in *C. elegans* RBM-26. (C)
Representative western blot of three biological replicates showing expression of RBM-
26::3XFLAG, RBM-26 P80L::3XFLAG and RBM-26 L13V::3XFLAG proteins. 20 μg of
total protein lysate was loaded per well and specific proteins were detected with an anti-

FLAG antibody and enhanced chemiluminescence. Protein loading was quantified by
No-Stain™ Protein Labeling Reagent. (D) Densitometric quantification of protein bands.
Statistical significance was analyzed by one-way ANOVA with a Tukey post hoc test,
**p<0.01. (E) Example of endogenously tagged RBM-26 in wild type at the L3 stage. (F)
Example of endogenously tagged RBM-26 in *rbm-26(P80L)* mutant at the L3 stage.
RBM-26 was endogenously tagged with Scarlet using CRISPR to generate the *rbm-*
26(syb2552[rbm-26::Scarlet::AID]) allele. CRISPR was used to insert the P80L mutation
into *rbm-26 (syb2552)* to create the *rbm-26(cue48[rbm-26(P80L)] syb2552[rbm-*
26::Scarlet::AID]) allele.
688
689
690
Figure 2: Mutations in *rbm-26* cause axon degeneration and defects in axon tiling.
(A) Example of a normal PLM axon, where the PLM axon does not overlap with the ALM
axon (B) Diagram of the touch receptor neurons. (C) Example of an axon tiling defect
(PLM/ALM overlap phenotype), where the PLM axon overlaps with the ALM axon. Axon
beading can also be seen (asterisk). (D) Quantification of axon tiling defects in various
stages. (E) Example of axon beading phenotype (asterisks) in an *rbm-26(null)* mutant.
(F) Quantification of beading phenotype at various stages. (G) Example of the axonal
break phenotype in the PLM of a two-day old adult *rbm-26 (P80L)* mutant. (H)
Quantification of the axonal break phenotype at various stages. Asterisks indicate
statistically significant difference relative to wild type, Z-test for proportions (***
p<0.0001, ** p<0.01 and * p<0.05), and error bars represent the standard error of the
proportion. Scale bars are 10 µm. Axons are visualized with the *muIs32 (Pmec-7::gfp)*
transgene. Stages are labeled with L1(first larval stage), L2(second larval stage,
L3(third larval stage), L4(fourth larval stage), 1-Day Adult and 2-Day Adult. Alleles: *rbm-*
26(null) is *rbm-26(gk910)*; *rbm-26(P80L)* is *rbm-26(cue23)*; *rbm-26(L13V)* is *rbm-*
26(cue34).
707
Figure 3: RBM-26 is expressed in the PLM and functions within neurons to
protect against axon degeneration and prevent defects in axon tiling.
(A) PLM is identified by GFP expression driven by the *muIs32* transgene that encodes
Pmec-7::gfp. (B) Expression of endogenously tagged RBM-26. The endogenously
tagged allele of RBM-26 is *rbm-26 [syb2552(rbm-26::Scarlet::AID)].* A and B are merged
in C. (D & E) Neuron-specific degradation of RBM-26::Scarlet::AID causes the
PLM/ALM overlap phenotype and axon beading. However, degradation in hypodermis,
muscle and intestine do not cause these phenotypes. Control strain is *rbm-*

26::Scarlet::AID. (F & G) The PLM/ALM overlap and the beading phenotypes caused by
mutations in *rbm-26* are rescued by the expression of the *Pmec-7::rbm-26::Scarlet*
transgene, which expresses RBM-26 in PLM and other touch receptor neurons.
Asterisks indicate statistically significant difference, Z-test for proportions (* $p<0.05$,
***$p<0.001$), and error bars represent the standard error of the proportion. "n" is 150 for
in D and E and 200 in F and G. Touch receptor neurons were visualized in D – G with
the *muIs32* transgene.

723
724
725

Figure 4: Loss of RBM-26 function causes a reduction in the density of
mitochondria in the PLM axon.
(A-E) RBM-26 loss of function reduces the number of mitochondria in the PLM axon. (A-
B) Example of mitochondria (arrowheads) in an L3 wild-type PLM. (C-D) Example of
mitochondria in *rbm-26(null)* L3 PLM. PLM axons are visualized with the *muIs32*
transgene that encodes *Pmec-7::gfp* and mitochondria are visualized with *jsIs1073*
transgene that encodes *Pmec-7::mito:RFP*. B & D show mitochondria only. Images are
from the middle of the PLM axon. Scale bars are 10 µm. (E) Quantification of the
density of mitochondria (number of mitochondria per 100 µm of PLM axon) in axons at
the L3 stage (n ≥ 25). Statistical significance was analyzed by one-way ANOVA with a
Tukey post hoc test, **$p<0.01$. (F) Quantification of the density of mitochondria (number
of mitochondria per 100 µm) in L1, L2 and L3 PLM axons. 20 PLMs were analyzed for
L1 and L2, and 25 PLMs were analyzed for L3. Statistical significance was analyzed by
two-tailed Student's t-test. The reduction in the number of mitochondria in PLM axons in
the *rbm-26* mutants was limited to the proximal PLM axon (50µm closest to the cell
body) (G) and was not observed in the distal PLM axon (50 µm closest to the axon tip)
(H). 20 PLMs were observed for each genotype in G and H. Statistical significance was
analyzed by one-way ANOVA with a Tukey *post hoc* test, ** $p < 0.01$, ns = not
significant. Error bars in E – H are the standard error of mean. Alleles: *rbm-26(null)* is
rbm-26(gk910); *rbm-26(P80L)* is *rbm-26(cue23)*; *rbm-26(L13V)* is *rbm-26(cue34)*.

746
Figure 5: Loss of RBM-26 causes mitochondria dysfunction in PLM. RBM-26 loss of
function causes excess oxidative activity in mitochondria. To detect oxidative activity in
mitochondria, we used mitoTimer, which is a green fluorescent protein when newly
synthesized but shifts to red fluorescence when oxidized. (A-A'') Example of mitoTimer
expression in an L3 wild-type PLM cell body. (B-B'') Example of mitoTimer expression in an
L3 *rbm-26(null)* mutant PLM cell body. Scale bars are 4 µm. (C–E) Quantification of the ratio

of red to green mitoTimer expression in *wt* and *rbm-26* mutant PLM cell bodies in different
larval stages. Statistical significance was analyzed by one-way ANOVA with a Tukey post
hoc test (*p<0.05, **p<0.01 and ns = not significant). "n" is 30 for C and E, and 21 for D.
Error bars represent the standard error of mean. The mitoTimer transgene is *cueSi35 IV*
(Pmec-7::mitoTimer::tbb-2 3'UTR). Alleles: *rbm-26(null)* is *rbm-26(gk910)*; *rbm-*
26(P80L) is *rbm-26(cue23)*; *rbm-26(L13V)* is *rbm-26(cue34)*.
759
760   **Figure 6: RBM-26 binds to the *mals-1* mRNA and negatively regulates its**
761   **expression.**
762
(A) RNA interactors of RBM-26 as identified by UV-RIPseq (see methods for details).
Data generated from three biological replicates. (B) Loss of RBM-26 function increases
expression of the *mals-1* mRNA. Absolute levels of *mals-1* mRNA quantified using real-
time RT-PCR (see methods for details). Graph represents averages of three biological
replicates. Error bars are standard deviations. Asterisks indicate statistically significant
difference relative to wild type, two-tailed Student's t-test (***p<0.0001). (C) Loss of
RBM-26 function causes an increase in MALS-1 protein expression. Wildtype worms
are *mals-1(cue37)* that expresses MALS-1::3X FLAG. CRISPR was used to insert P80L
and L13V mutations into the *mals-1(cue37)* chromosome generating *rbm-26(cue40)*
mals-1(cue37) and *rbm-26(cue49) mals-1(cue37),* respectively. 20 µg of total protein
lysate from a mixed stage population was loaded per well in the top panel and specific
proteins were detected with an anti-FLAG antibody and enhanced chemiluminescence.
Loading was assessed with No-Stain™ protein labeling reagent. 40 µg of total protein
lysate was loaded for wild-type and *rbm-26(cue40) mals-1(cue37)* mutants in bottom
panels. (D) Densitometric quantification of MALS-1 protein levels in western blots. Data
are expressed as mean ± S.E.M. from three independent experiments. Asterisks
indicate statistically significant difference relative to wild type, two-tailed Student's t-test
(**p<0.01,*p<0.05)
781
782
783   **Figure 7: Overexpression of MALS-1 reduces mitochondria in the PLM axon and**
784   **causes axon degeneration and defects in axon tiling.**
785
(A-C) Overexpressed MALS-1::Scarlet protein colocalizes with mitochondria. (A)
Mitochondria visualized with the *jsIs609* transgene, which expresses a mitochondria-
targeted GFP (mitoGFP) in the PLM neuron. (B) MALS-1::Scarlet expressed by the
cueSi36 IV transgene. (C) Merged image showing colocalization of MALS-1::Scarlet

with mitoGFP. (D-F) Overexpression of MALS-1::Scarlet causes defects similar to those
caused by loss of RBM-26 function. (D) Axon tiling defects caused by MALS-1::Scarlet
overexpression. (E) PLM axon beading phenotype caused by MALS-1::Scarlet
overexpression. (F) Decrease in mitochondria density (number of mitochondria per 100
μm) in the PLM axon caused by MALS-1::Scarlet overexpression. Asterisks in panels D
and E indicate statistically significant difference relative to wild type, Z-test for
proportions (**p<0.01, ***p<0.0001). For panels D and E, error bars represent the
standard error of the proportion. Statistical significance in panel F was analyzed by
Student's t-test, ***p<0.001 and error bars represent the standard error of mean.
799
800
801   **Figure 8: Loss of MALS-1 function suppresses the mitochondrial and axon**
802   **degeneration phenotypes that are caused by loss of RBM-26 function.**
803
(A) Loss of *mals-1* function suppresses the PLM axon beading phenotype caused by
the *rbm-26* mutations. (B) Loss of *mals-1* function suppresses the reduction of
mitochondria caused by the *rbm-26* mutations. (C) Loss of *mals-1* function causes PLM
axon termination defect but fails to enhance PLM axon termination defects caused by
the *rbm-26* mutations. The *rbm-26(cue40 [P80L]) mals-1(syb6330)* double mutant was
created by introducing the P80L mutation into the *mals-1(syb6330)* mutant
chromosome. The *rbm-26(cue41 [P80L]) mals-1(tm12122)* double mutant was created
by introducing the P80L mutation into the *mals-1(tm12122)* mutant chromosome. The
rbm-26 (cue47 [L13V]) mals-1(syb6330) double mutant was created by introducing the
L13V mutation into the *mals-1(syb6330)* mutant chromosome. Asterisks in A and C
indicate statistically significant difference, Z-test for proportions (*p<0.05, **p<0.01, ***
p<0.0001). For panels A and C, error bars represent the standard error of the proportion
(n = 200). For panel B, error bars represent the standard error of the mean (n=25), and
statistical significance was analyzed by one-way ANOVA with a Tukey post hoc test
(*p<0.05 and **p<0.01). (D) Loss of RBM-26 protein expression causes a decrease in
mitoribosome expression, which was assayed with the MitoRibo-Tag system consisting
of the MRPL-58 mitoribosomal protein tagged at its C-terminus with 3XFLAG (MRPL-
58::3XFLAG). This MRPL-58::3XFLAG protein was expressed by the *mrpl-58(cue38)*
allele, which was created by CRISPR. *mrpl-58(cue38) rbm-26(cue39)* double mutant
was created by introducing the P80L mutation into *mrpl-58(cue38)* by CRISPR. CRISPR
was also used to insert a 3X FLAG tag at the C-terminus of MRPL-58 in *mals-*
1(syb6330) mutant to create *mrpl-58(cue50) mals-1(syb6330)*. Finally, *rbm-26 (P80L)*
mutation was inserted into *mrpl-58(cue50) mals-1(syb6330)* to create *rbm-26(cue51*

[P80L]) mrpl-58(cue50) mals-1(syb6330). 25 µg of total protein lysate from mixed stage
population were loaded per lane and specific proteins were detected with an anti-FLAG
antibody and enhanced chemiluminescence. Protein loading was analyzed by No-
stain^{TM} protein labeling reagent. (E) Densitometric quantification of MRPL-58 protein
levels in western blots. Data are expressed as mean ± S.E.M. from three biological
replicates. Asterisks indicate statistically significant difference relative to wild type,
Student's t-test (*$p<0.05$).
834
835

Literature Cited

1.　Chalkia, D., et al., *Association Between Mitochondrial DNA Haplogroup Variation and*
Autism Spectrum Disorders. JAMA Psychiatry, 2017. **74**(11): p. 1161-1168.
2.　Chang, X., et al., *Mitochondrial DNA haplogroups and risk of attention deficit and*
hyperactivity disorder in European Americans. Transl Psychiatry, 2020. **10**(1): p. 370.
3.　Valiente-Palleja, A., et al., *Genetic and clinical evidence of mitochondrial dysfunction in*
autism spectrum disorder and intellectual disability. Hum Mol Genet, 2018. **27**(5): p.
891-900.
4.　Wang, Y., et al., *Association of mitochondrial DNA content, heteroplasmies and inter-*
generational transmission with autism. Nat Commun, 2022. **13**(1): p. 3790.
5.　Wang, Y., M. Picard, and Z. Gu, *Genetic Evidence for Elevated Pathogenicity of*
Mitochondrial DNA Heteroplasmy in Autism Spectrum Disorder. PLoS Genet, 2016.
12(10): p. e1006391.
6.　Yardeni, T., et al., *An mtDNA mutant mouse demonstrates that mitochondrial deficiency*
can result in autism endophenotypes. Proc Natl Acad Sci U S A, 2021. **118**(6).
7.　Liu, X., et al., *Oxidative Stress in Autism Spectrum Disorder-Current Progress of*
Mechanisms and Biomarkers. Front Psychiatry, 2022. **13**: p. 813304.
8.　Burns, D.T., et al., *Variants in EXOSC9 Disrupt the RNA Exosome and Result in Cerebellar*
Atrophy with Spinal Motor Neuronopathy. Am J Hum Genet, 2018. **102**(5): p. 858-873.
9.　Wan, J., et al., *Mutations in the RNA exosome component gene EXOSC3 cause*
pontocerebellar hypoplasia and spinal motor neuron degeneration. Nat Genet, 2012.
44(6): p. 704-8.
10.　Zanni, G., et al., *Exome sequencing in a family with intellectual disability, early onset*
spasticity, and cerebellar atrophy detects a novel mutation in EXOSC3. Neurogenetics,
2013. **14**(3-4): p. 247-50.
11.　Shirakaki, S., R.R. Roshmi, and T. Yokota, *Genetic Approaches for the Treatment of Giant*
Axonal Neuropathy. J Pers Med, 2022. **13**(1).
12.　Cousin, M.A., et al., *Pathogenic SPTBN1 variants cause an autosomal dominant*
neurodevelopmental syndrome. Nat Genet, 2021. **53**(7): p. 1006-1021.

13. Qi, C., et al., *Variants in ADD1 cause intellectual disability, corpus callosum dysgenesis,*
and ventriculomegaly in humans. Genet Med, 2022. **24**(2): p. 319-331.
14. Leite, S.C., et al., *The Actin-Binding Protein alpha-Adducin Is Required for Maintaining*
Axon Diameter. Cell Rep, 2016. **15**(3): p. 490-498.
15. Lorenzo, D.N., et al., *betaII-spectrin promotes mouse brain connectivity through*
stabilizing axonal plasma membranes and enabling axonal organelle transport. Proc Natl
Acad Sci U S A, 2019. **116**(31): p. 15686-15695.
16. Abrahams, B.S., et al., *SFARI Gene 2.0: a community-driven knowledgebase for the*
autism spectrum disorders (ASDs). Mol Autism, 2013. **4**(1): p. 36.
17. Silla, T., et al., *The human ZC3H3 and RBM26/27 proteins are critical for PAXT-mediated*
nuclear RNA decay. Nucleic Acids Res, 2020. **48**(5): p. 2518-2530.
18. Zhou, Y., et al., *The fission yeast MTREC complex targets CUTs and unspliced pre-mRNAs*
to the nuclear exosome. Nat Commun, 2015. **6**: p. 7050.
19. Ariyapala, I.S., et al., *The RNA-binding protein Swm is critical for Drosophila*
melanogaster intestinal progenitor cell maintenance. Genetics, 2022. **222**(2).
20. Deciphering Developmental Disorders, S., *Prevalence and architecture of de novo*
mutations in developmental disorders. Nature, 2017. **542**(7642): p. 433-438.
21. Homsy, J., et al., *De novo mutations in congenital heart disease with*
neurodevelopmental and other congenital anomalies. Science, 2015. **350**(6265): p. 1262-
6.
22. Iossifov, I., et al., *The contribution of de novo coding mutations to autism spectrum*
disorder. Nature, 2014. **515**(7526): p. 216-21.
23. Krumm, N., et al., *Excess of rare, inherited truncating mutations in autism.* Nat Genet,
2015. **47**(6): p. 582-8.EXOSC
24. Turner, T.N., et al., *denovo-db: a compendium of human de novo variants.* Nucleic Acids
Res, 2017. **45**(D1): p. D804-D811.
25. Zhou, X., et al., *Integrating de novo and inherited variants in 42,607 autism cases*
identifies mutations in new moderate-risk genes. Nat Genet, 2022. **54**(9): p. 1305-1319.
26. Gallegos, M.E. and C.I. Bargmann, *Mechanosensory neurite termination and tiling*
depend on SAX-2 and the SAX-1 kinase. Neuron, 2004. **44**(2): p. 239-49.
27. Pan, C.L., et al., *Genetic analysis of age-dependent defects of the Caenorhabditis elegans*
touch receptor neurons. Proc Natl Acad Sci U S A, 2011. **108**(22): p. 9274-9.
28. Tank, E.M., K.E. Rodgers, and C. Kenyon, *Spontaneous age-related neurite branching in*
Caenorhabditis elegans. J Neurosci, 2011. **31**(25): p. 9279-88.
29. Toth, M.L., et al., *Neurite sprouting and synapse deterioration in the aging*
Caenorhabditis elegans nervous system. J Neurosci, 2012. **32**(26): p. 8778-90.
30. Burke, R.E. and K. O'Malley, *Axon degeneration in Parkinson's disease.* Exp Neurol, 2013.
246: p. 72-83.
31. Yong, Y., et al., *Axonal spheroids in neurodegeneration.* Mol Cell Neurosci, 2021. **117**: p.
103679.

32. Yuan, P., et al., *PLD3 affects axonal spheroids and network defects in Alzheimer's disease.*
Nature, 2022. **612**(7939): p. 328-337.
33. Martinez, M.A.Q. and D.Q. Matus, *Auxin-mediated Protein Degradation in*
Caenorhabditis elegans. Bio Protoc, 2020. **10**(8).
34. Drozd, C.J. and C.C. Quinn, *UNC-116 and UNC-16 function with the NEKL-3 kinase to*
promote axon targeting. Development, 2023. **150**(18).
35. Murphy, M.P., *How mitochondria produce reactive oxygen species.* Biochem J, 2009.
417(1): p. 1-13.
36. Schon, E.A. and S. Przedborski, *Mitochondria: the next (neurode)generation.* Neuron,
2011. **70**(6): p. 1033-53.
37. Laker, R.C., et al., *A novel MitoTimer reporter gene for mitochondrial content, structure,*
stress, and damage in vivo. J Biol Chem, 2014. **289**(17): p. 12005-12015.
38. Rorbach, J., P.A. Gammage, and M. Minczuk, *C7orf30 is necessary for biogenesis of the*
large subunit of the mitochondrial ribosome. Nucleic Acids Res, 2012. **40**(9): p. 4097-109.
39. Boczonadi, V., et al., *EXOSC8 mutations alter mRNA metabolism and cause*
hypomyelination with spinal muscular atrophy and cerebellar hypoplasia. Nat Commun,
2014. **5**: p. 4287.
40. Di Donato, N., et al., *Mutations in EXOSC2 are associated with a novel syndrome*
characterised by retinitis pigmentosa, progressive hearing loss, premature ageing, short
stature, mild intellectual disability and distinctive gestalt. J Med Genet, 2016. **53**(6): p.
419-25.
41. Drozd, C.J., T.A. Chowdhury, and C.C. Quinn, *UNC-16 interacts with LRK-1 and WDFY-3 to*
regulate the termination of axon growth. Genetics, 2024.
42. Buddell, T., et al., *An autism-causing calcium channel variant functions with selective*
autophagy to alter axon targeting and behavior. PLoS Genet, 2019. **15**(12): p. e1008488.
43. Buddell, T. and C.C. Quinn, *An autism-associated calcium channel variant causes defects*
in neuronal polarity in the ALM neuron of C. elegans. MicroPubl Biol, 2021. **2021**.
44. Harigaya, Y., et al., *Selective elimination of messenger RNA prevents an incidence of*
untimely meiosis. Nature, 2006. **442**(7098): p. 45-50.
45. Bergeron, D., et al., *Regulated Intron Retention and Nuclear Pre-mRNA Decay Contribute*
to PABPN1 Autoregulation. Mol Cell Biol, 2015. **35**(14): p. 2503-17.
46. Wu, M., et al., *The RNA exosome shapes the expression of key protein-coding genes.*
Nucleic Acids Res, 2020. **48**(15): p. 8509-8528.
47. Giunta, M., et al., *Altered RNA metabolism due to a homozygous RBM7 mutation in a*
patient with spinal motor neuropathy. Hum Mol Genet, 2016. **25**(14): p. 2985-2996.
48. Brown, A., et al., *Structures of the human mitochondrial ribosome in native states of*
assembly. Nat Struct Mol Biol, 2017. **24**(10): p. 866-869.
49. Cheng, J., O. Berninghausen, and R. Beckmann, *A distinct assembly pathway of the*
human 39S late pre-mitoribosome. Nat Commun, 2021. **12**(1): p. 4544.

50. Rebelo-Guiomar, P., et al., *A late-stage assembly checkpoint of the human mitochondrial*
ribosome large subunit. Nat Commun, 2022. **13**(1): p. 929.

51. Buddell, T. and C. Quinn, *A null allele in the wdfy-3 selective autophagy gene of C.*
elegans. MicroPubl Biol, 2022. **2022**.

52. Ghanta, K.S., T. Ishidate, and C.C. Mello, *Microinjection for precision genome editing in*
Caenorhabditis elegans. STAR Protoc, 2021. **2**(3): p. 100748.

53. Nonet, M.L., *Efficient Transgenesis in Caenorhabditis elegans Using Flp Recombinase-*
Mediated Cassette Exchange. Genetics, 2020. **215**(4): p. 903-921.

54. Xu, Y. and C.C. Quinn, *MIG-10 functions with ABI-1 to mediate the UNC-6 and SLT-1 axon*
guidance signaling pathways. PLoS Genet, 2012. **8**(11): p. e1003054.

55. Xu, Y. and C.C. Quinn, *Transition between synaptic branch formation and synaptogenesis*
is regulated by the lin-4 microRNA. Dev Biol, 2016. **420**(1): p. 60-66.

56. Xu, Y., et al., *Axon response to guidance cues is stimulated by acetylcholine in*
Caenorhabditis elegans. Genetics, 2011. **189**(3): p. 899-906.

57. Xu, Y., et al., *SYD-1C, UNC-40 (DCC) and SAX-3 (Robo) function interdependently to*
promote axon guidance by regulating the MIG-2 GTPase. PLoS Genet, 2015. **11**(4): p.
e1005185.

58. Fischer, N.C., et al., *The ANC-1 (Nesprin-1/2) organelle-anchoring protein functions*
through mitochondria to polarize axon growth in response to SLT-1. PLoS Genet, 2022.
18(11): p. e1010521.

59. Kechin, A., et al., *cutPrimers: A New Tool for Accurate Cutting of Primers from Reads of*
Targeted Next Generation Sequencing. J Comput Biol, 2017. **24**(11): p. 1138-1143.

60. Langmead, B. and S.L. Salzberg, *Fast gapped-read alignment with Bowtie 2.* Nat
Methods, 2012. **9**(4): p. 357-9.

61. Langmead, B., et al., *Scaling read aligners to hundreds of threads on general-purpose*
processors. Bioinformatics, 2019. **35**(3): p. 421-432.

62. Dobin, A., et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013. **29**(1): p.
15-21.

63. Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and dispersion*
for RNA-seq data with DESeq2. Genome Biol, 2014. **15**(12): p. 550.

64. Li, H., et al., *The Sequence Alignment/Map format and SAMtools.* Bioinformatics, 2009.
25(16): p. 2078-9.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure S1: Longer exposure of blot shown in figure 1C. Representative western blot of three biological replicates showing expression of RBM-26::3XFLAG, RBM-26 P80L::3XFLAG and RBM-26 L13V::3XFLAG proteins. 20 μg of total protein lysate was loaded per well and specific proteins were detected with an anti-FLAG antibody and enhanced chemiluminescence. Protein loading was quantified by No-Stain™ Protein Labeling Reagent.

Figure S2: Analysis of PLM axon length suggests that the *rbm-26* mutants cause overextension of the PLM axon. PLM length as a ratio of body length at L3 in wild type (wt), *rbm-26 (null), rbm-26 (P80L) and rbm-26 (L13V)* mutants. Error bars = SD. Statistical significance was analyzed by one-way ANOVA with a Tukey *post hoc* test, ** $p < 0.01$. n for wt = 24, n for *rbm-26 null* = 21, n for *rbm-26 (P80L)* = 20 and n for *rbm-26 (L13V)* = 22. Alleles: *rbm-26(null)* is *rbm-26(gk910)*; *rbm-26(P80L)* is *rbm-26(cue23)*; *rbm-26(L13V)* is *rbm-26(cue34)*

Figure S3: Examples of PLM degeneration phenotypes. (A) Example of beading phenotype. Only those focal enlargements along PLM axon that were at least twice the diameter of the axon were considered to be beads. (B) Example of blebbing phenotype. (C) Example of waviness in PLM axon. (D) Quantification of the bead density (beads per 100 um of PLM axon) observed in wild type (wt), *rbm-26(null)*, *rbm-26(P80L)* and *rbm-26(L13V)* at L3. Error bars are standard error of mean. Statistical significance was analyzed by one-way ANOVA with a Tukey *post hoc* test, * $p < 0.05$ (n = 28).

□ *wt*   ■ *rbm-26 (P80L)*

Figure S4. The *rbm-26(P80L)* mutation causes degeneration phenotypes in the axon and dendrites of the PVD neuron. (A) Wild type L4 PVD – primary (1°), secondary (2°), tertiary (3°), and quaternary (4°) dendrites can be seen. (B) Beading can be seen in tertiary dendrites and some quaternary dendrites are missing in *rbm-26(P80L)* L4 PVD neuron. (C) Quantification of phenotypes observed in *rbm-26(P80L)* mutants. 50 wildtype L4 PVD neurons were observed and 60 *rbm-26(P80L)* L4 PVD neurons were observed. PVD neurons were visualized by the *wdIs52 [F49H12.4::GFP + unc-119(+)]* transgene, which expresses GFP in PVD. Asterisks indicate statistically significant difference relative to wild type, Z-test for proportions (* p<0.05), and error bars represent the standard error of the proportion.

Figure S5. RBM-26 is localized to nuclei in hypodermal and neuronal cells.
RBM-26 is expressed in the nuclei of neurons and hypodermal cells. Neurons are identified by the pan-neuronal expression of GFP by the *evIs111* transgene (panels A and D). Hypodermal nuclei are identified by tissue-specific expression of blue fluorescence protein tagged with a nuclear localization signal and an auxin-induced degron *(col-10p::TIR1::F2A::mTagBFP2::AID::NLS::tbb-2 3'UTR)* (panels C and D). Panels B & D show endogenously tagged RBM-26*(rbm-26::Scarlet::AID)*. Panels A – C are merged in D. " N" in panels A, B, and D indicate neuron and "Hn" in panels B, C, and D indicate hypodermal nuclei.

Figure S6: RBM-26 is widely expressed. Representative Z-stack projection of endogenous RBM-26 protein tagged with Scarlet (RBM-26::Scarlet::AID) at larval stage L3.

Figure S7: The *rbm-26::Scarlet::AID* gene does not cause the PLM/ALM overlap phenotype (A) or the beading phenotype (B) in the absence of auxin. Worms expressing auxin inducible degron (AID) in frame with RBM-26 and tissue-specific TIR-1 as indicated on the X-axis were placed in Auxin free worm plates at L4 and their progeny were observed at L3. "n" is between 125 and 150. N.S. = not significant (Z-test for proportions). Error bars represent the standard error of the proportion.

Figure S8: Auxin treatment in the absence of RBM-26::Scarlet::AID does not cause the PLM/ALM axon overlap phenotype or the beading phenotype. L4 wild type or neuronal TIR-1 (without AID tagged RBM-26) were placed on 4 mM synthetic auxin coated plates, allowed to lay eggs and their L3 progenies were analyzed for (A) PLM termination defects and (B) beading phenotypes. "n" is 132 for wt and 138 for Neuronal TIR-1. N.S. = Not Significant (Z-test for proportions). Error bars represent the standard error of the proportion.

Figure S9. Example of mitoTimer expression in an L3 wild-type [A – C], *rbm-26(P80L)* [D-F] and *rbm-26(L13V)* [E-G] PLM cell bodies. mitoTimer is expressed by the transgene *cueSi35(Pmec-7::mitoTimer::tbb-2 3'UTR)*. Alleles: *rbm-26(P80L)* is *rbm-26(cue23)*; *rbm-26(L13V)* is *rbm-26(cue34).* Scale bars are 5 µm.

Figure S10: Degron mediated depletion of RBM-26 in PLM. L4 *rbm-26(syb2552); resi7* worms that express RBM-26::Scarlet::AID and F-box Transport Inhibitor Response 1 (TIR-1) in neurons were placed in normal NGM plates (A – C) or on plates coated with 4mM synthetic auxin (D–E), allowed to lay eggs and L3 progeny were analyzed through confocal microscopy. PLM neuron was visualized with the help of the *muIs32 (Pmec-7::gfp)* transgene. Scale bars are 10 µm. Auxin treatment causes depletion of RBM-26 protein, which was quantified in G as a ratio of RBM-26::Scarlet intensity relative to the GFP expression in PLM.

Figure S11: Example of RNA standards used to calculate the quantity of _mals-1_ mRNA present in 1 μg of total RNA in wild-type and _rbm-26_ mutants _(P80L and L13V)_ L4 _C. elegans_. Synthetic RNA corresponding to 68 nucleotides in the last exon and the 3'UTR of _mals-1_ mRNA was purchased from Azenta Inc. The RNA was diluted 1 ng/μl, 0.1 ng/μl, 0.01 ng/μl, 0.001 ng/μl, and 0.0001 ng/μl in nuclease free water. The dilutions were quantified in Qubit (Thermo Fisher Scientific) using RNA Qubit Broad Range kit (Thermo Fischer Scientific). 1, 0.1, 0.01, 0.001, 0.0001 ng of RNA standard was used in RT-qPCR reaction. Cycle to threshold (Ct) values obtained from RT-qPCR reaction was plotted against the known quantity of RNA used.

	Age	% Blebbing	% Beading	% Waviness	% Break	N
Wild type	L1	0.925926	1.851852	0	0	108
	L2	0	2.941176	0.980392	0	102
	L3	0	4.545455	1.515152	0	198
	L4	0	6.097561	1.829268	0	164
	1-DAY Adult	0	14.03509	4.385965	0	114
	2-DAY Adult	1.886792	17.92453	0	0.943396	106
rbm-26 (null)	L1	5	0	0	0	80
	L2	2.631579	11.40351	1.754386	0	114
	L3	0	32.36715	13.52657	1.449275	207
	L4	NA	NA	NA	NA	NA
	1-DAY Adult	NA	NA	NA	NA	NA
	2-DAY Adult	NA	NA	NA	NA	NA
rbm-26 (P80L)	L1	1.960784	8.823529	3.921569	0	102
	L2	1.6	15.2	1.6	0	125
	L3	0	24.37811	7.462687	1.492537	201
	L4	2.112676	26.40845	2.112676	2.112676	284
	1-DAY Adult	2.542373	33.89831	4.237288	1.694915	118
	2-DAY Adult	3.448276	62.6506	4.237288	6.024096	118
rbm-26 (L13V)	L1	0.980392	10.78431	2.941176	0	102
	L2	2	8.783784	0.666667	0	150
	L3	1.470588	13.23529	3.333333	0	204
	L4	0.78125	13.88889	2.5	0	128
	1-DAY Adult	0.847458	24.57627	3.389831	0.847458	118
	2-DAY Adult	3.508772	36.84211	3.508772	2.631579	132

Table S1: PLM phenotypes observed in larval stages L1 through L4, one- and two-day adults in wild-type, *rbm-26 (null)*, *rbm-26 (P80L)* and *rbm-26 (L13V)* worms. Maternally rescued *rbm-26 (null)* worms do not survive past L3.

Gene.name	Human Ortholog (HGNC Symbol)	log2FoldChange	padj
F59C6.15	NA	3.529339	1.70E-09
pud-2.1	NA	3.103615	5.63E-06
pud-2.2	NA	3.04405	7.45E-06
pud-1.1	NA	2.772255	2.98E-05
pud-1.2	NA	2.749208	3.21E-05
C23H5.8	NA	2.578636	0.00021
C37A2.9	NA	5.056751	0.000568
ilys-5	NA	2.848721	0.000745
K12H4.2 (mals-1)	MALSU1	1.983865	0.000826
rrn-2.1	NA	2.319809	0.002424
clec-85	NA	1.860425	0.00272
K07C5.11	NA	2.422354	0.002764
21ur-8178	NA	4.814433	0.003612
spp-4	NA	2.183194	0.005335
Y6B3B.13	NA	3.736574	0.00653
21ur-3462	NA	4.815693	0.009799
K02A11.4	NA	2.023582	0.011462
21ur-10303	NA	4.592398	0.01198
F43A11.8	NA	2.201064	0.01198
F33H12.7	NA	2.189781	0.014145
msp-81	NA	1.939555	0.01655
fip-7	NA	2.878975	0.017713
21ur-12103	NA	4.77819	0.019219
Y46G5A.40	NA	2.514422	0.021423
F55A8.4	NA	1.86247	0.022547
clc-1	NA	1.772934	0.023129
clec-209	NA	1.99952	0.026991
21ur-7931	NA	4.59147	0.027738
F59D8.3	NA	2.344256	0.027738
F22D6.16	NA	2.290677	0.027738
K07C5.18	NA	2.283691	0.027738
K06A4.7	NA	2.239518	0.027738
T19C4.17	NA	1.972172	0.027738
D1065.7	NA	2.118396	0.028351
C33H5.13	NA	1.929031	0.028351
dpy-4	MARCO	1.922891	0.028351
Y75B8A.28	NA	1.860966	0.028351
F28H7.3	NA	1.968765	0.029705
Y38H6C.23	NA	2.002049	0.029819
Y40B1A.5	NA	2.581652	0.030633
T27A10.8	NA	2.290521	0.031615
C08F11.116	NA	2.249682	0.032408
F16H6.10	NA	1.354888	0.032408
F56A4.2	NA	1.954246	0.035992
Y43F8C.22	NA	2.149066	0.03674
gst-22	GSTA1/GSTA2/GSTA3/GSTA4/GSTA5	1.583856	0.037515
W05H9.3	NA	1.974427	0.039296
F07H5.3	NA	1.942375	0.040116
F54B11.14	NA	1.465389	0.040116
C05E7.t2	NA	3.735348	0.041196
W02F12.11	NA	3.084148	0.041196
T19C9.8	NA	1.693645	0.041196
acdh-1	ACADS/ACADSB	2.136673	0.043721
lys-4	NA	2.048744	0.043721
21ur-7165	NA	4.475939	0.04407
F18C5.15	NA	2.463115	0.04407
H04D03.5	NA	2.223033	0.04407
JC8.15	NA	2.10709	0.04407
21ur-5454	NA	3.776464	0.045548
21ur-10438	NA	2.263752	0.045548
C18E9.14	NA	2.055965	0.046461
ZK546.18	NA	2.109542	0.049805
rme-2	LRP3/LRP12/LDLRAD3/LRP10	1.706674	0.049881

Table S2: RNA interactors of RBM-26 identified by RNA immunoprecipitation followed by RNASeq. Human orthologs were identified with Orhtolist2 (Woojin Kim, Ryan S Underwood, Iva Greenwald, Daniel D Shaye, OrthoList 2: A New Comparative Genomic Analysis of Human and *Caenorhabditis elegans* Genes, *Genetics*, Volume 210, Issue 2, 1 October 2018, Pages 445–461). NA = Human Ortholog not found.

Genotype	Source
rbm-26(cue23[rbm-26 (P80L)]) III	This Study
rbm-26(cue34[rbm-26 (L13V)]) III	This Study
rbm-26(cue25[rbm-26::3X FLAG]) III	This Study
rbm-26(cue22[rbm-26 (P80L)] cue25[rbm-26::3X FLAG]) III	This Study
rbm-26(cue24[rbm-26 (L13V)] cue25[rbm-26::3X FLAG]) III	This Study
mals-1(cue37[mals-1::3X FLAG]) III	This Study
mals-1(syb6330) III	SUNY Biotech
rbm-26(cue40[rbm-26(P80L)]) mals-1(syb6330) III	This Study
rbm-26(cue42[rbm-26(P80L)]) mals-1(cue37 [mals-1::3X FLAG]) III	This Study
mals-1(tm12122) III	Mitani Lab, International C. elegans Gene Knockout Consortium
rbm-26(cue 41[rbm-26(P80L)]) mals-1(tm12122) III	This Study
mrpl-58(cue38[mrpl-58::3XFLAG]) III	This Study
rbm-26(cue39[rbm-26 (P80L)]) mrpl-58(cue 38 [mrpl-58::3X FLAG]) III	This Study
rbm-26(cue 47[rbm-26 (L13V)]) mals-1(syb6330)] III	This Study
rbm-26(syb2552[rbm-26::Scarlet::AID]) III	SUNY Biotech
rbm-26(cue 48[rbm-26 (P80L)]) syb2552[rbm-26::Scarlet::AID]) III	This Study
cueSi35(Pmec-7::mitoTimer::tbb-2 3'UTR) IV	this study
cueSi36(Pmec-7::mals-1::scarlet::tbb-2 3'UTR) IV	this study
evIs111[F25B3.3::GFP + dpy-20(+)]; rbm-26(syb2552[rbm-26::Scarlet::AID]) III; reSi1[col-10p::TIR1::F2A::mTagBFP2::AID::NLS::tbb-2 3'UTR] I*	this study
muIs32[mec-7p::GFP + lin-15(+)]	CGC
muIs32[mec-7p::GFP + lin-15 (+)]; rbm-26(cue23[rbm-26 (P80L)]) III	this study
muIs32[mec-7p::GFP + lin-15 (+)]; rbm-26(gk910) III/hT2 [bli-4(e937) let-?(q782) qIs48] I;III	this study
muIs32[mec-7p::GFP + lin-15 (+)]; rbm-26(cue34 [rbm-26 (L13V)]) III	this study
muIs32[mec-7p::GFP + lin-15(+)]; rbm-26(syb2552 [rbm-26::Scarlet::AID]) III	this study
muIs32[mec-7p::GFP + lin-15(+)]; rbm-26(syb2552 [rbm-26::Scarlet::AID]) III; reSi1[col-10p::TIR1::F2A::mTagBFP2::AID::NLS::tbb-2 3'UTR] I*	this study
muIs32[mec-7p::GFP + lin-15(+)]; rbm-26(syb2552 [rbm-26::Scarlet::AID]) III; reSi7[rgef-1p::TIR1::F2A::mTagBFP2::AID::NLS::tbb-2 3'UTR] I*	this study
muIs32[mec-7p::GFP + lin-15(+)]; rbm-26(syb2552 [rbm-26::Scarlet::AID]) III; reSi3[unc-54p::TIR1::F2A::mTagBFP2::AID::NLS::tbb-2 3'UTR] I*	this study
muIs32[mec-7p::GFP + lin-15(+)]; rbm-26(syb2552 [rbm-26::Scarlet::AID]) III; reSi12[ges-1p::TIR1::F2A::mTagBFP2::AID::NLS::tbb-2 3'UTR] II*	this study
jsIs1073[mec-7p::TagRFP-mito::CBunc119]	Obtained from Michael Nonet
jsIs1073[mec-7p::TagRFP-mito ,CBunc119]; rbm-26(gk910) III/hT2[bli-4(e937) let-?(q782) qIs48] I;III	this study
jsIs609[mec7p::mtGFP + lin-15(+)] X	Obtained from Michael Nonet
jsIs609[mec7p::mtGFP + lin-15(+)] X; cueSi36(Pmec-7::mals-1::scarlet::tbb-2 3'UTR) IV	this study
cueSi35(Pmec-7::mitoTimer::tbb-2 3'UTR) IV; rbm-26(gk910) (III)/hT2 [bli-4(e937) let-?(q782) qIs48] (I;III)	this study
cueSi35(Pmec-7::mitoTimer::tbb-2 3'UTR) IV; rbm-26(cue23 [rbm-26(P80L)]) III	this study
*wbmIs67[eft-3p::3XFLAG::wrmScarlet::unc-54 3'UTR *wbmIs65] V*	CGC
*wbmIs72[pie-1p::3XFLAG::GFP::unc-54 3'UTR *wbmIs60] III*	CGC
cueSi52(Pmec-7::rbm-26::scarlet::unc-54 3'UTR) IV	this study
rbm-26(cue49[rbm-26(L13V)] mals-1(cue37[mals-1::3XFLAG]) III	this study
mrpl-58(cue50 [mrpl-58::3XFLAG]) mals-1(syb6330) III	this study
rbm-26(cue51[rbm-26(P80L)] mrpl-58(cue50 [mrpl-58::3XFLAG]) mals-1(syb6330) III	this study

Table S3: Strains used in this study

Table S4: Resource Tables

Primers used in figure 5B
TGCACGAAGGAGCACAAAAC (forward)
CGTGCACCTGAACTCTTTCC (reverse)

Antibodies used in this study

Antibody	Catalog Number	Manufacturer	Dilution
DYKDDDDK Tag Antibody	2368S	Cell Signaling Technology	1:1000
Goat Anti-Rabbit IgG H&L (HRP)	ab205718	Abcam	1: 5000

CRISPR reagents used in this study

CRISPR edit	guide RNA	Homology Directed Repair Template
rbm-26 (P80L)	CACAAGCAAAAGCTATATGC	GCAGTTGCTGAAGCAGAGGTAGGCGCTGTTGATGCAAGCATATAGCTTTTGCTAGTGAGTGCTTCGAAGACTTTGTCGACGAATGGAGCAGT
rbm-26 (L13V)	ATGAGGATGCTCTCTTCGAT	GGCTGTCTAATCAATGAAAAAACTTACATTGGCGAGAGCTCATCAGAGACCCAGTCGAAAAGAGCATCCTCATTGTCGATGTGCATGCTTTCTTTGATATT
rbm-26 (3X FLAG)	ACTAGGACTAATCATTTAGC	ATTTTGAGATCAAAGTTCTAATATAACTAGGACTATTTATCGTCATCGTCTTTGTAGTCGATATCGTGATCTTTGTAGTCGCCATCGTGATCTTTGTAGTCATCATTTAGCAGGTCATCTTCTTCGTCAGACTGAAATCTATCATTTAGCAGGTCATCTTCTTCGTCAGACTGAAATCT
mals-1 (3X FLAG)	AAAGGAAAACATTATCGTCT	CAATAATCTAAATGTTATAAATAACAAAAGGAAAACATTACTTGTCATCGTCGTCTTTGTAGTCGATGTCGTGATCCTTATAATCGCCGTCGTGGTCCTTGTAGTCTCGTCTCGGTGGCAGCAAAATCTTCTGTTTCTCCTCATCAATCTCG
mrpl-58 (3X FLAG)	GCTAATTAGCTAAAACTCAA	GGAAAAATAAAACACGAACAAAATTTCAGCTAATTAGCTACTTGTCATCGTCGTCTTTGTAGTCGATGTCGTGATCCTTATAATCGCCGTCGTGGTCCTTGTAGTCAAACTCAACGGCGGCTCGTCGCGACGCCTTTTTCTCACTCGTTCT