Title: The *R1441C-LRRK2* mutation induces myeloid immune cell exhaustion

in an age- and sex-dependent manner

Authors: Rebecca L., Wallings[1,2], Karen McFarland[2,3], Hannah A. Staley[1,2], Noelle

Neighbarger[1,2], Susen Schaake[4], Norbert Brüggemann[4,5], Simone Zittel[5], Tatiana Usnich[4,5],

Christine Klein[4], Esther M Sammler[6, 7] and Malú Gámez Tansey[1, 2, 3, 8]*

Affiliations:

[1] Department of Neuroscience, [2] Center for Translational Research in Neurodegenerative Disease,

University of Florida, College of Medicine, McKnight Brain Institute, Gainesville, Florida, USA

[3] Department of Neurology and Fixel Institute for Neurological Diseases, University of Florida

Health, Gainesville, Florida, USA

[4] Institute of Neurogenetics, University of Lübeck, Lübeck, Germany.

[5] Department of Neurology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

[6] MRC Protein Phosphorylation and Ubiquitylation Unit, School of Life Sciences, University of

Dundee, Dundee DD1 5EH, U.K. d.r.alessi@dundee.ac.uk e.m.sammler@dundee.ac.uk.

[7] Department of Neurology, School of Medicine, Ninewells Hospital, Ninewells Drive, Dundee

DD1 9SY, U.K.

[8] Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase,

MD, USA

*Corresponding author. Email: mgtansey@ufl.edu

Abstract: Considering age is the greatest risk factor for many neurodegenerative diseases, aging, in
particular aging of the immune system, is the most underappreciated and understudied contributing factor
in the neurodegeneration field. Genetic variation around the *LRRK2* gene affects risk of both familial and
sporadic Parkinson's disease (PD). The leucine-rich repeat kinase 2 (LRRK2) protein has been implicated
in peripheral immune signaling, however, the effects of an aging immune system on LRRK2 function have
been neglected to be considered. We demonstrate here that the *R1441C* mutation induces a hyper-
responsive phenotype in macrophages from young female mice, characterized by increased effector
functions, including stimulation-dependent antigen presentation, cytokine release, phagocytosis, and
lysosomal function. This is followed by age-acquired immune cell exhaustion in a Lrrk2-kinase-dependent
manner. Immune-exhausted macrophages exhibit suppressed antigen presentation and hypophagocytosis,
which is also demonstrated in myeloid cells from *R1441C* and *Y1699C*-PD patients. Our novel findings
that *LRRK2* mutations confer immunological advantage at a young age but may predispose the carrier to
age-acquired immune exhaustion have significant implications for LRRK2 biology and therapeutic
development. Indeed, LRRK2 has become an appealing target in PD, but our findings suggest that more
research is required to understand the cell-type specific consequences and optimal timing of LRRK2-
targeting therapeutics.

One Sentence Summary: The *R1441C-LRRK2* mutation causes an age-acquired immune cell
exhaustion in macrophages in sex-dependent manner

Word count: 9483

Main Text:

INTRODUCTION

Parkinson's disease (PD) is a common, progressive neurodegenerative disease, affecting around 1-2% of the population over the age of 65 (*1*). PD prevalence will likely increase two-fold by the year 2030 (*2*). In addition, the projected total economic burden will surpass $79 billion by 2037 (*3*), underscoring the need to reduce PD incidence and delay disease progression. The fact that there are currently no disease-modifying drugs for people with PD indicates that knowledge gaps still need to be closed to identify ways to cure or prevent this disease. Although the etiology of PD is largely unknown, it is thought to involve a complex interaction between various genetic and environmental factors (*4*). Although typically thought of as a disease limited to the central nervous system (CNS), evidence suggests a crucial and fundamental role of inflammation in both the CNS and the periphery in PD pathogenesis (*5*).

Pathogenic variants in the gene encoding leucine-rich repeat kinase 2 (LRRK2) account for the majority of familial PD patients (*6*). LRRK2 is a large, complex protein containing six functional domains, two of which have known enzymatic activities; a ROC domain (GTPase activity) and a MAPKK domain (kinase activity). There are numerous single-nucleotide substitutions in each of these domains that have been shown to be pathogenic (*7-14*). The second most prevalent mutation, *R1441C*, located in the GTPase ROC domain, is present in about 46% of familial Parkinson's disease that has an origin in the Basque region in Spain (*15*) and leads to decreased LRRK2 GTPase hydrolysis and increased LRRK2 protein kinase activity (*16, 17*).

LRRK2 is expressed in innate and adaptive immune cells and its expression is tightly regulated by immune stimulation (*18-22*). Furthermore, LRRK2 is a member of the receptor interacting protein (RIP) kinase family, which is a group of proteins that detect and respond to cellular stress by regulating cell death and activation of the immune system (*23*). LRRK2 function has also been implicated in pathogen control based on observations that LRRK2 variants confer differential susceptibility to a number of bacterial infections, including *Mycobacterium tuberculosis* (*24, 25*), *Salmonella Typhimurium* (*26*), and *Listeria monocytogenes* (*27*), as well as an increased risk of developing leprosy (*28*) and inflammatory bowel disease such as

Crohn's (*29*) and ulcerative colitis (*30*). Collectively such data strongly suggest a generalized role of

LRRK2 in immune system regulation.

Although research has focused primarily on neuroinflammation in PD over the last two decades, it is

becoming increasingly clear that peripheral inflammatory responses may play important roles in PD

pathogenesis (*31, 32*). For example, immune signaling plays a role in the regulation of neurodegeneration

in a *LRRK*2-PD pre-clinical model, with the *R1441C* mutation altering the type II interferon response in

peripheral immune cells (*22*). Furthermore, neutralization of IL-6 over-production in the periphery of these

mice prevents substantia nigra *pars compacta* (SNpc) dopaminergic (DA) neuron loss in the presence of

peripheral lipopolysaccharide (LPS) exposure (*33*). Importantly, immune dysfunction in the periphery takes

primacy over the CNS, as the presence of *Lrrk2* mutations in lymphocytes alone is sufficient to induce DA

neuron loss in the midbrain of LPS-treated mice (*33*). Such data suggest a pertinent role of the peripheral

immune system in *LRRK2*-PD, with immune dysfunction driving the development of the disease.

As well as a role in immune response, a vast amount of research has focused on the role of LRRK2 at the

lysosome. In immune cells, specifically, LRRK2 has been shown to mediate a number of lysosome-

dependent immune functions such as phagocytosis (*34*) and antigen presentation (*18, 21, 35*). Intriguingly,

LRRK2 has been shown to alter lysosomal function and autophagy in kidneys in an age-dependent, biphasic

manner, with enhanced autophagy levels in young *Lrrk2*-null mice, that are reduced and drop below control

levels once the mice reach 20 months of age (*36*). Despite this intriguing observation, few studies have

investigated the effects of *LRRK2* mutations on lysosomal processes in the context of aging. Considering

age is the greatest risk factor for many neurodegenerative diseases, aging, in particular aging of the immune

system (*5*), is the most underappreciated and understudied contributing factor to neurodegeneration in the

field.

To close this important gap, we assessed the effects of *LRRK2* mutations on immune responses and

lysosomal function in peripheral immune cells in young versus old mice. We used *R1441C-Lrrk2* mutant

knock-in (KI) mice based on previous findings demonstrating an effect of the *R1441C/G* mutation on

peripheral immune responses (*22, 33*) whilst evading limitations of an overexpression model and

investigating the effects of the *R1441C* mutation in physiologically relevant levels. Peritoneal macrophages

(pMacs) were isolated from these mice to examine the effects of the *R1441C* mutation on macrophage cell

function at different ages. pMacs are not a homogenous population of cells, but rather a mix of small and

large pMacs (SPMs and LPMs, respectively). LPMs are resident to the peritoneal cavity and are

traditionally thought of as anti-inflammatory, phagocytic and responsible for the presentation of exogenous

antigens (*37*). SPMs, on the other hand, are generated from bone-marrow-derived myeloid precursors which

migrate to the peritoneal cavity in response to infection, inflammatory stimuli, or thioglycolate, and present

a pro-inflammatory functional profile (*37*). We observed a bi-phasic, age-dependent effect of the *R1441C-*

Lrrk2 KI mice, with increased antigen presentation, production of anti-inflammatory cytokines, lysosomal

activity, and pathogen uptake in peripheral macrophages from young (2-3 months) *R1441C-Lrrk2* KI mice

examined *ex vivo*. Conversely, peripheral macrophages from aged (18-21 months) *R1441C-Lrrk2* KI mice

exhibited a diminished capacity to present antigens in the presence of inflammatory insults, decreased

lysosomal function and pathogen uptake, with concomitant increases in DNA fragmentation in the presence

of pathogens. Importantly, we demonstrate that this is a sex-dependent phenotype, with male *R1441C-Lrrk2*

KI failing to present this age-dependent phenotype. Furthermore, we demonstrate that the age-dependent

hypophagocytosis and decreased antigen presentation in the *R1441C-LRRK2* KI females is due to an

acquired immune cell exhaustion driven by increased Lrrk2 protein kinase activity and immune responses

whilst young. These phenotypes were also observed in human peripheral myeloid cells, with monocyte-

derived macrophages from *R1441C* and *Y1699C* patients exhibiting decreased pathogen uptake and

increased programmed cell death protein 1 (PD1) expression, consistent with immune cell exhaustion.

RESULTS

LRRK2 R1441C **mutant immune cells display age-dependent biphasic alteration in antigen**

presentation

LRRK2 protein expression has been shown to increase in immune cells from both murine pre-clinical

models and patient cells in response to inflammatory stimuli (*18, 21, 22, 38-40*). To determine if whether

this is age-dependent, Lrrk2 protein levels were assessed in pMacs from young and aged mice treated with

100U of IFNγ for 18h in the presence or absence of 100nM of the Lrrk2 kinase inhibitor, PF360 (PF-

06685360). In pMacs from young females, IFNγ treatment significantly increased total Lrrk2 protein levels

in both B6 and *R1441C* pMacs, with no effects of PF360 observed (Fig. 1A, B). Relative to pMacs from

young female mice, pMacs from aged females expressed increased total Lrrk2 protein levels under vehicle

stimulation, with aged *R1441C* pMacs expressing significantly less Lrrk2 protein levels relative to B6 aged

pMacs. Furthermore, pMacs from aged females did not exhibit a significant increase in Lrrk2 protein

expression with IFNγ treatment, with aged *R1441C* pMacs still expressing significantly less Lrrk2 protein

levels relative to B6 aged pMacs in IFNγ-treatment. In pMacs from young male mice, the same observations

were made as in the females (Fig. S1A, B). However, in aged male mice, although a lack of IFNγ-mediated

Lrrk2 increase was observed as seen in females, the effect of the *R1441C* mutation was not observed. These

observations indicate that Lrrk2 protein levels increase in macrophages as a function of age in both males

and females, however, the *R1441C* mutation limits this increase in a sex-dependent manner.

As pMacs are not a homogenous population of cells, as previously discussed, we used multi-color flow

cytometry-based methods to immunophenotype LPMs and SPMs, as the two populations can be

distinguished based on CD11b expression (Fig. S1C) (*37*). No differences were seen in LPM count between

genotypes in males or females at either young or aged (Fig. S1D-G). No significant differences in SPM

count were seen in pMacs from young females or males (Fig. S1H, I), nor aged males (Fig. S1J). However,

a significant increase in SPM count was observed in pMacs from *R1441C* aged females relative to B6

controls in IFNγ treatment conditions (Fig. S1K). As bone-marrow-derived myeloid precursors are known

to differentiate into SPMs upon inflammation, it may be, therefore, *R1441C* bone-marrow-derived myeloid

precursors present in these cultures have increased propensities to differentiate in such conditions.

To begin to investigate the effects of the *R1441C* mutation on pMac function at different ages, surface

MHC-II expression was assessed on both LPMs and SPMs via multi-color flow cytometry as a measure of

antigen presentation in both vehicle and IFNγ treated pMacs. In pMacs from young females, the *R1441C*

mutation increases surface MHC-II expression on LPMs and percentage of MHC-II$^+$ LPMs in both vehicle

and IFNγ treatments (Fig 1C, Fig. S1L). Interestingly, this effect of the *R1441C* mutation was not observed

in LPMs from young male mice (Fig. 1D, Sup. Fig 1M). Regarding surface MHC-II expression on SPMs,

a significant increase was seen in vehicle-stimulated SPMs from young *R1441C* females relative to B6

controls (Fig. S1N) which was not observed in SPMs from males (Fig. S1O). However, an increase in

MHC-II$^+$ SPM percentage was observed in both male and female *R1441C* pMacs (Fig. S1P, Q).

When assessing surface MHC-II on vehicle and IFNγ treated pMacs from aged females it was observed

that, although no effects of genotype were detectable in vehicle-treated LPMs, a significant reduction in

surface MHC-II expression on LPMs from *R1441C* females relative to B6 controls was evident in IFNγ-

treated conditions (Fig. 1E), indicative of decreased ability to engage in antigen presentation in these cells.

As observed in young mice, no effects of genotype were observed in LPMs from male mice (Fig. 1F)

suggesting a sex-dependent effect of the *R1441C* mutation on antigen presentation in macrophages. No

significant genotype effects were observed regarding percentage of MHC-II+ LPMs or MHC-II+ SPMs

(Fig. S1R-W).

With alterations in surface MHC-II expression, there is likely increased and decreased transport of MHC-

II-complexes to the cell surface in pMacs from young and aged *R1441C* females, respectively. To test this,

using pMacs treated with vehicle or IFNγ, we co-stained for extracellular vs intracellular MHC-II

(ExMHCII vs IcMHCII) and monitored expression via fluorescence microscopy. pMacs were co-treated

with 100nM PF360 to determine the effects of Lrrk2 kinase activity on surface MHC-II expression. PF360

treatment reduced levels of Lrrk2 phosphorylated at S935 (pS935 Lrrk2) in all groups (Fig. S2A, B). pMacs

from young *R1441C* females exhibited increased Ex:IcMHCII ratio relative to B6 controls upon IFNγ

treatment, which was ameliorated upon Lrrk2 kinase inhibition (Fig. S2C, D), with this treatment

decreasing Ex:IcMHCII ratio in both genotypes. Such data suggests that Lrrk2-kinase inhibition reduces the transport of IcMHCII to the cell surface to engage in antigen presentation. No effect of the *R1441C* mutation was observed in young males (Fig. S2E, F). Interestingly, in pMacs from aged *R1441C* females, a significant reduction in Ex:IcMHCII ratio was observed in IFNγ treatments relative to B6 controls, with no effect of the *R1441C* mutation observed in pMacs from aged males (Fig.1G-J). Lrrk2 kinase inhibition also had no significant effect on Ex:IcMHCII ratio in pMacs from *R1441C* mice, but significantly reduced Ex:IcMHCII ratio in pMacs from B6 controls. It seems therefore that the *R1441C* mutation exhibits a bi-phasic, age- and sex-dependent effect on antigen presentation in macrophages. The effect of Lrrk2 protein kinase inhibition on antigen presentation, however, does not appear to be bi-phasic nor age-dependent, with Lrrk2 kinase inhibition decreasing antigen presentation in pMacs from both young and aged mice.

LRRK2 R1441C **mutant immune cells display age-dependent biphasic alteration in cytokine release** *ex vivo* **and circulating cytokines in vivo**

To investigate the extent to which alterations in antigen presentation were accompanied by changes in cytokine release, media from vehicle- and IFNγ treated cells were collected and cytokine levels were quantified on a multiplexed immunoassay panel (Fig. 2A). In pMacs from young females, increases in IL6 and IL10 relative to B6 controls were observed, with no differences between genotypes in TNF release observed (Fig. 2B). IL6 and IL10 release were also increased in pMacs from young *R1441C* male mice relative to B6 controls with no differences between genotypes observed in TNF release (Fig. 2C). In pMacs from aged female mice, the genotype-dependent changes in IL6 seen in young mice were no longer observed (Fig. 2D) and a significant decrease in IL10 was observed, with a concomitant increase in TNF release, relative to B6 controls. With a decrease in the release of the anti-inflammatory cytokine, IL10, and an increase in the release of the pro-inflammatory cytokine, TNF, these findings indicate that pMacs from aged *R1441C* female mice promote a pro-inflammatory microenvironment. Interestingly, in pMacs from

aged male mice, the *R1441C*-associated increase in IL6 and IL10 seen in young mice was retained with age

but *R1441C* macrophages secreted more TNF than B6 controls after stimulation (Fig. 2E).

To determine if these alterations in cytokine release associated with the *R1441C* mutation *ex vivo* was also

detectable *in vivo*, plasma from whole blood was collected and cytokine levels were quantified (Fig. 2F).

A significant increase in IL10, IL6, and TNF was observed in plasma from young *R1441C* female and male

mice (Fig. 2G). Conversely, a decrease in circulating levels of IL10 was observed in the plasma of aged

R1441C female and male mice relative to B6 controls (Fig. 2H). Similarly, IL6 was also increased in aged

R1441C plasma relative to B6 controls, although this was only observed in females, and no significant

differences in TNF levels were observed.

Regarding other cytokines measured in media collected from pMacs, although significant effects of

genotype were observed, these differences were not as consistent between age and sex as was observed for

IL10, IL6 and TNF. For example, an increase in keratinocyte chemoattractant (KC)/growth-regulated

oncogene (GRO), a small cytokine that belongs to the CXC chemokine family, was observed in pMacs

from young *R1441C* females, which was not observed in pMacs from males (Fig. S3A). However, no

significant effects of genotype were observed in aged pMacs (Fig. S3B). Upon IFNγ treatment, pMacs from

young *R1441C* females exhibited decreased levels of the pro-inflammatory cytokine, IL1β (Fig. S3C)

which was not observed in pMacs from young male mice. Again, no significant differences between

genotypes were observed in aged pMacs (Fig. S3D). Although no significant differences between genotypes

were observed in the release of the anti-inflammatory cytokine, IL4, in young pMacs (Fig. S3E), a

significant decrease was observed in pMacs from aged *R1441C* females relative to B6 controls, with no

differences seen in aged males (Fig. S3F). Lastly, although no significant differences in IL5 were seen in

pMacs from young mice (Fig. S3G), a decrease in IL5 release was observed in aged pMacs from *R1441C*

females and males relative to B6 controls (Fig. S3H).

The R1441C Lrrk2 mutation is associated with alterations of monocyte populations and microglia activation in aged female mice

The observation of a biphasic immune response, with an initial hyperinflammation state proceeded by a subsequent downregulation of immune responses, prompted us to hypothesize the presence of the *R1441C* mutation elicited immune cell exhaustion. Immune exhaustion arises from chronic stimulation of immune cells, leading to a downregulation, or even absence, of an expected effector response (e.g. antigen presentation, cytokine release or phagocytosis) (*41*). It is known that monocytes pushed to exhaustion shift to express higher Ly6C levels and a reduction in Ly6C$^-$ monocytes under resting conditions (*42*). We therefore assessed monocyte sub-populations in the blood of aged mice to determine if a shift to an exhausted state could be observed. Indeed, we observed a detectable increase in Ly6C$^+$:Ly6C$^-$ in the peripheral blood mononuclear cells isolated from blood of aged *R1441C* female mice relative to B6 controls (Fig. S4A) with no significant change observed in aged male mice (Fig. S4B). Interestingly, no effects of genotype were observed in young mice (Fig. S4C, D), suggesting development of an age-acquired immune exhaustion phenotype in the *R1441C* females.

It has been suggested that circulating monocytes, disrupted in PD, may infiltrate the brain parenchyma and affect local glial and adaptive immune responses (*43*). To determine whether exhausted monocytes could be infiltrating the brain parenchyma of aged *R1441C* female mice, immune cells were isolated from whole brain and assessed via multi-color flow cytometry. Indeed, in aged *R1441C* females, an increase in overall Ly6C$^+$ monocytes in the brain parenchyma was observed relative to B6 controls, whilst MHC-II/Ly6C$^+$ were decreased (Fig. S4E, F), suggesting that whilst there is an increase in infiltrating Ly6C$^+$ monocytes in the brains of aged *R1441C* females, they exhibit the same decrease in antigen presenting capabilities to those observed in pMacs *ex vivo*. Importantly, alongside this increase in infiltrating monocytes into the brain parenchyma, a concomitant increase in MHC-II expression on microglia was observed in aged *R1441C* females relative to B6 controls (Fig. S4G) with no changes in overall microglia number (Sup. Fig 4H). No changes were observed in the brains of male mice (Fig. S4I-L).

LRRK2 R1441C mutant immune cells display age-dependent biphasic alterations in lysosomal activity

As previously discussed, loss of _Lrrk2_ leads to biphasic alterations in lysosomal function in kidneys (_36_). 250 Interestingly, lysosomal dysfunction has been implicated in immune cell exhaustion, with decreased 251 autophagy levels and lysosomal function being associated with increased T cell exhaustion (_44_). We 252 therefore hypothesized that the biphasic immune response observed in _R1441C_ females would be associated 253 with concomitant biphasic lysosomal function. 254

The Eα: YAe model is used to monitor the antigen presentation capabilities of cells by incubating an 255 endogenous peptide (Eα 52-68) which is subsequently phagocytosed, transported to the lysosome, loaded 256 onto an MHC-II complex at the lysosome and transported back to the plasma membrane for antigen 257 presentation (Fig. 3A). This Eα peptide-loaded MHC-II can subsequently be detected via flow cytometry 258 using the YAe antibody (_45_). We observed that YAe median fluorescence intensity (MFI) was significantly 259 increased in IFNγ-treated LPMs from young _R1441C_ females relative to B6 controls, which was not 260 observed in LPMs from young male mice (Fig. 3B, C). When co-treated with PF360, YAe MFI was 261 significantly reduced in LPMs of both sexes, ameliorating the increased YAe MFI observed in LPMs from 262 young _R1441C_ females. Antigen presentation and pathogen sensing require protease action and sufficient 263 lysosomal function in order to occur (_46_), which is why lysosomotropic agents have been shown to decrease 264 peptide-loaded-MHC-II surface expression in antigen presenting cells (_45_). Indeed, when co-treated with 265 the vacuolar H+ ATPase (V-ATPase) inhibitor, Bafilomycin A1, YAe MFI significantly decreased in both 266 _R1441C_ and B6 LPMs from both sexes (Fig. 3B, C). The same age-dependent biphasic phenotype in 267 _R1441C_ females was observed, with IFNγ-treated LPMs from aged females exhibiting decreased YAe MFI 268 relative to B6 controls which was not observed in aged male mice (Fig. 3D, E). PF360 significantly reduced 269 YAe in LPMs from aged B6 mice and aged _R1441C_ males but had no significant effect on LPMs from aged 270 _R1441C_ females. Furthermore, the same effect of Bafilomycin A1 was observed in all genotypes. Due to 271

this observation and the crucial role of the lysosome in antigen presentation, we next sought to probe the

effects of the *R1441C* mutation on lysosomal function in these pMacs.

pMacs were plated and incubated with the lysosomal degradation probe, DQ Red BSA, and the pan-

cathepsin probe, BMV109, and collected for flow cytometry to quantify lysosomal activity. It was observed

that LPMs from young *R1441C* females exhibited increased DQ-BSA and BMV109 MFI (Fig. 3F, G),

indicative of increased lysosomal protein degradation and pan-cathepsin activity, respectively. Conversely,

LPMs from aged *R1441C* females exhibited increased DQ-BSA and BMV109 MFI (Fig. 3H, I). No effects

of genotype were observed in LPMs from male mice (Fig. S5A-D). These observations made via flow

cytometry were confirmed with fluorescence microscopy (Fig. 3J-N; Fig. S5E-J). As was observed with

YAe measurements, Lrrk2 kinase inhibition via treatment with PF360 significantly reduced both DQ-BSA

and BMV109 MFI observed via microscopy in all groups except aged *R1441C* females, which exhibited

decreased measurement of lysosomal activity which were unaffected by Lrrk2 kinase inhibition.

_R1441C_ Lrrk2 mutation is association with age-acquired hypo-phagocytosis

Lysosomal function in macrophages is crucial for a number of other immune-related functions in myeloid

cells, including phagocytosis. Therefore, we posited that the observed biphasic effects of the *R1441C*

mutation on lysosomal function would have downstream consequences on phagocytosis in pMacs. To

explore this hypothesis, pMacs were plated and incubated with pHrodo™ Green *E. coli* BioParticles™,

after which uptake and fluorescence intensity were monitored over 5 hours. An increase in both GFP[+] cells

and GFP MFI was observed in pMac cultures from young *R1441C* females (Fig. 4A-C). In male mice,

although the percent of GFP[+] cells was not significantly different between the two genotypes, pMacs from

young *R1441C* males did exhibit increased GFP MFI relative to B6 controls (Fig. S6A-C). Lrrk2 kinase

inhibition with PF360 significantly reduced GFP MFI in both *R1441C* males and female pMacs, however

only pMacs from male *R1441C* mice exhibited significantly reduced percentage of GFP[+] cells in response

to Lrrk2 kinase inhibition. It appears, therefore, that in young mice, the *R1441C* mutation induces an

upregulation in phagocytic uptake in pMacs from both males and females, as indicated by increased GFP

MFI, which can be ameliorated upon Lrrk2 kinase inhibition.

In pMacs from aged females, a significant reduction in both percent GFP$^+$ cells and GFP MFI was observed

in pMacs from *R1441C* females relative to B6 controls (Fig. 4D-F). Although Lrrk2 kinase inhibition with

PF360 significantly increased percent GFP$^+$ cells in pMacs from *R1441C* females, no effect of Lrrk2 kinase

inhibition was observed in GFP MFI. pMacs from aged male *R1441C* mice exhibited the same decrease in

GFP$^+$ cells as females, relative to B6 controls, that was also partially ameliorated by Lrrk2 kinase inhibition

(Fig. S6D). Interestingly, pMacs from aged *R1441C* male mice exhibit increased GFP MFI relative to B6

controls (Fig. S6E, F) which was unaffected by Lrrk2 kinase inhibition. Collectively, it seems that pMacs

from *R1441C* females exhibit increased phagocytic uptake at a young age and develop an age-dependent

hypophagocytic phenotype. pMacs from *R1441C* males, however, exhibit this same increased phagocytic

uptake at a young age but retain this increase in old ages.

Hypophagocytic activity in *R1441C* macrophages is associated with aberrant pathogen-mediated

DNA damage

Our data thus far suggests a potential age-acquired immune cell exhaustion in macrophages from *R1441C*

female mice. In other disease states associated with cellular exhaustion, apoptotic depletion of immune cells

is also observed (*47*). To investigate the level of pathogen-induced apoptosis in pMacs from *R1441C* mice,

pMacs were plated in the presence of *E. coli* for 12 hours, harvested and stained using terminal

deoxynucleotidyl transferase dUTP nick end labeling (TUNEL), which detects DNA breaks formed during

the final phase of apoptosis, and TUNEL fluorescence was assessed via flow cytometry. LPMs from young

R1441C females exhibited decreased TUNEL MFI in response to *E. Coli* relative to B6 controls (Fig. 4G),

whilst an increase was observed in LPMs from aged *R1441C* females (Fig. 4H). Reduced apoptosis could

lead to prolonged presence of activated immune cells which should otherwise be eliminated, resulting in an

increased and prolonged inflammatory response and, if sustained for prolonged times, could lead to immune

cell exhaustion. Indeed, LPMs from young *R1441C* males exhibit increased TUNEL MFI indicative of

increased apoptosis in these cells (Fig. S6G), with no effect of genotype observed in aged males (Fig. S6H).

These novel findings support the conclusion that *R1441C* males evade this female-specific, age-acquired

immune exhaustion that is seen in females by increasing clearance of activated immune cells.

Immune exhaustion in *R1441C* Lrrk2 mutant immune cells is dependent on Lrrk2 kinase activity

Our data thus far suggest a potential age-acquired immune cell exhaustion in macrophages from *R1441C*

female mice. We hypothesized that this age-acquired immune exhaustion was driven by the initial

hyperinflammatory phase when these mice are young, characterized by increased cytokine release,

increased antigen presentation, and reduced apoptosis of activated immune cells. We therefore

hypothesized that pMacs from young *R1441C* female mice could be pushed to acquire this age-dependent

immune exhaustion phenotype if subjected to chronic overstimulation *ex vivo*. To test this, pMacs were

isolated and plated in the presence of a vehicle or 100ng/mL lipopolysaccharide (LPS) for a 5-day chronic

treatment (Fig. 5A). We first sought to understand how Lrrk2 expression is altered by LPS-induced

exhaustion and how age and the *R1441C* affects this. In pMacs from young females, an increase in Lrrk2

expression was observed in response to LPS-induced exhaustion (Fig. 5B, C). Importantly, there was no

significant effect of treatment in pMacs from aged female mice, with aged *R1441C* pMacs expressing

significantly less Lrrk2 relative to B6 controls. Interestingly, these same observations were made in pMacs

from aged male mice (Fig. S7A, B) suggesting that, with an additional chronic inflammatory stress, aged

R1441C males can exhibit some of the phenotypes observed in females.

A number of negative costimulatory markers that are associated with immune exhaustion have been

identified, including the immune checkpoint molecule, programmed cell death-1 (PD1) (*48*). PD1

expression was quantified via multi-color flow cytometry in pMacs from aged mice to determine if pMacs

from *R1441C* females were exhausted as hypothesized. Indeed, an increase in PD1$^+$ cells was observed in

both vehicle and LPS-exhausted *R1441C* LPMs relative to B6 controls (Fig. 5D). As this was seen in both

vehicle and LPS conditions, these exciting findings suggest that pMacs from *R1441C* females are already

exhibiting immune exhaustion markers, which is then exacerbated by chronic LPS-exhaustion.

Interestingly, LPMs from aged *R1441C* male mice, although no significant effect of genotype was observed

in vehicle conditions, when pushed to exhaustion with LPS, an increase in PD1$^+$ cells relative to B6 controls

was observed (Fig. S7C). The lack of differences in PD1$^+$ cells between genotypes in aged male mice

suggests there is no effect of the *R1441C* mutation on immune exhaustion in naïve animals. However, based

on the increased PD1$^+$ cell counts observed in *R1441C* males when treated with LPS for 5 days, we conclude

that a secondary, chronic inflammatory insult can push *R1441C* pMacs from male mice to acquire immune

exhaustion like that seen in female mice.

When macrophages from young *R1441C* female mice were assessed for PD1 expression after a 5-day

incubation with vehicle or LPS, a significant increase in PD1$^+$ cell count was observed in both treatment

conditions relative to B6 controls, with a significant increase detectable in LPS-treated conditions relative

to vehicle (Fig. 5E). Interestingly, when cells were pushed to exhaustion with Lrrk2 kinase inhibitor co-

treatment, a significant reduction in PD1$^+$ cell count was observed in *R1441C* cells. These data suggest that

R1441C macrophages from female mice may be susceptible to immune exhaustion in a Lrrk2-kinase

dependent manner. No significant differences between genotypes were observed in LPMs from young male

mice (Fig. S7D).

We next sought to determine if LPS-induced exhausted macrophages also exhibited the diminished antigen

presentation capabilities that we observe in the aged *R1441C* macrophages. After a 5-day treatment with

vehicle or LPS +/- PF30, macrophages were treated with IFNγ and MHC-II expression was quantified via

multi-color flow cytometry. In LPMs subjected to a 5-day vehicle pre-treatment, the same increased MHC-

II expression in *R1441C* macrophages relative to B6 controls was observed (Fig. 5F). As hypothesized, in

LPMs pushed to exhaustion with a 5-day LPS pre-treatment, a significantly muted response to IFNγ was

elicited in both genotypes; however, *R1441C* LPMs exhibited decreased MHC-II relative to B6 controls in

response to IFNγ, phenocopying what was observed in macrophages with aged mice. Importantly, this could

be ameliorated by co-treatment with PF360 during the 5-day pre-treatment, suggesting that the push to

immune exhaustion and vulnerability of *R1441C* macrophages is dependent on Lrrk2 protein kinase

activity. Furthermore, we also observed that pMacs from young *R1441C* females pushed to exhaustion with

a 5-day pre-treatment of LPS exhibited a significant reduction in phagocytosis as measured by uptake of

pHrodo™ Green *E. coli* relative to B6 controls detected by GFP[+] cell percentage and GFP MFI which was

ameliorated with PF360 co-treatment (Fig. 5G-I).

Based on the observation that the PD1[+] cell count was elevated in LPMs from aged *R1441C* males relative

to B6 controls, we hypothesized that a 5-day LPS treatment would induce immune exhaustion in pMacs

from aged *R1441C* males similar to that seen in female mice. After a 5-day treatment with vehicle or LPS

+/- PF30, macrophages from both young and aged male mice were treated with IFNγ and MHC-II

expression quantified via multi-color flow cytometry. No significant effects of genotype were observed in

LPMs from young male mice (Fig. S7D-E). However, in LPMs from aged males pushed to exhaustion with

5-day LPS pre-treatment, *R1441C* LPMs exhibited decreased MHC-II relative to B6 controls in response

to IFNγ, phenocopying what was observed in macrophages from aged female *R1441C* mice (Fig. S7F).

Notably, this could be rescued with PF360 co-treatment during the 5-day LPS pre-treatment. Additionally,

the same hypophagocytic phenotype observed in aged *R1441C* female pMacs was observed in males once

pushed to exhaustion with a 5-day LPS treatment (Fig. S7G-I). Collectively such data support the

hypothesis that, with a secondary hit in the form of a chronic, inflammatory stimulus, male *R1441C* mice

exhibit increased immune cell exhaustion relative to B6 controls, similar to that seen in naïve *R1441C*

female macrophages.

Transcriptomic profiling reveals age- and sex-specific alterations in critical effector function

pathways in *R1441C* macrophages

To explore the age and sexually dimorphic effects of the *R1441C* mutation on macrophage function, RNA

was harvested from pMacs post-vehicle or IFNγ treatment and processed for RNA sequencing (RNAseq).

A principal component analysis (PCA) was conducted from the resulting dataset which revealed that groups

in our dataset primarily cluster based on age and treatment with vehicle or IFNγ treatment (Fig. S8A, B).

We first sought to identify key cellular pathways altered under different treatment conditions in *R1441C*

pMacs relative to B6 controls. In pMacs from young females treated with IFNγ, Gene Ontology (GO)

enrichment analysis revealed that upregulated differentially expressed genes (DEGs) in *R1441C*

macrophages were enriched in pathways such as regulation of phagocytosis, T cell differentiation, cytokine

production and defense response to virus (Fig. S8C, D). In pMacs from aged females treated with IFNγ,

GO enrichment analysis showed downregulated DEGs in *R1441C* macrophages were enriched in pathways

such as detection of external biotic stimulus, acute inflammatory response, and regulation of MAP kinase

activity (Fig. S8E, F). Collectively, such observations support our previous data that shows an effect of

R1441C Lrrk2 on functional readouts such as increased phagocytosis and cytokine production in response

to IFNγ in young females, whilst in aged mice, effector functions responsible for defense responses are

significantly downregulated.

Overall, fewer genes were dysregulated in *R1441C* pMacs from young females treated with vehicle, relative

to pMacs treated with IFNγ (Sup. Fig.9A), although these DEGs were enriched in similar pathways (Fig.

S9B). Similar observations were made with pMacs treated with vehicle from aged females (Fig. S9C). As

Lrrk2 protein levels increase in response to IFNγ, it follows that DEGs driven by the *R1441C* mutation

would have a more significant biological impact in these conditions where Lrrk2 levels are increased.

Furthermore, given that the majority of phenotypes we observe in pMacs from *R1441C* females are seen in

response to an inflammatory insult or immunological challenge, such as IFNγ or *E. Coli,* it is logical that

more *R1441C*:B6 DEGs would be observed in these conditions relative to vehicle treatments. Few

R1441C:B6 DEGs were observed in pMacs from male mice at both ages and in both treatments (Fig. S9E-

I). This is unsurprising, given that many phenotypes were not observed in *R1441C* male mice. pMacs from

aged *R1441C* only begin to exhibit phenotypes similar to *R1441C* female mice when pushed to immune

cell exhaustion and re-challenged with IFNγ. It is therefore likely that to identify key cellular pathways that

are altered by the *R1441C* mutation in male mice, a second, chronic immune insult is needed to elicit those

changes.

R1441C Lrrk2 mutation alters transcriptional regulation of mitochondrial and oxidative stress genes

in macrophages in a sex- and age-dependent manner

Based on the observation that the *R1441C* mutation alters gene expression in pMacs from aged female

mice, we next sought to determine the key cellular pathways that this mutation altered in the aging process.

When comparing DEGs in pMacs from both B6 and *R1441C* aged mice treated with IFNγ relative to those

from young pMacs, there was a significant degree of overlap between the DEGs identified in the two

genotypes (Fig. 6A-C). However, 2012 genes were identified to be differentially expressed upon aging and

novel to *R1441C* pMacs. KEGG pathway enrichment analysis revealed that these 2012 genes unique to

aged R1441C pMacs were significantly enriched for pathways associated with neurodegenerative diseases,

such as Parkinson's disease, Alzheimer's disease and Huntington's disease (Fig. 8D). Similarly, GO

biological pathway enrichment analysis of these DEGs showed a significant enrichment in pathways related

to mitochondria, oxidative stress and metabolism, including mitochondrial respiratory chain complex I

assembly, glucose and pyruvate metabolic process and cellular respiration, which are all known key

lysosomal and mitochondrial functions. Many DEGs observed in the Parkinson's Disease KEGG pathway

were present in the GO biological pathways associated with mitochondrial function and oxidative stress,

including many *Nduf* and *Cox* genes, *Park7, Pink1* and *Prkn.*

When comparing DEGs in aged pMacs from both B6 and *R1441C* female mice treated with vehicle from,

unsurprisingly, different DEGs were identified relative to what was seen in IFNγ conditions (Fig. S10A-

D). As opposed to pathways related to mitochondria and oxidative induced by IFNγ conditions, DEGs novel

to *R1441C* in vehicle conditions were significantly enriched in pathways such as wound healing responses,

and Wnt, ERK and PI3K signaling. It seems, therefore, that at rest, or under vehicle-treatment conditions,

the *R1441C* mutation exerts its effects on aging through pathways crucial for metabolism, cell survival, and

cell growth. The Wnt signaling cascade is crucial in macrophages and serves as a regulator of macrophage

inflammation-resolving state (*49*). However, during an inflammatory insult, the *R1441C* mutation exerts its

effects on aging through pathways crucial for oxidative stress and cellular respiration.

Interestingly, when assessing DEGs in pMacs from aged male mice treated with IFNγ, 540 genes were

identified as novel and significantly upregulated in *R1441C* macrophages that were not significantly

affected by aging in B6 controls (Fig. S10E-G). Pathway enrichment analysis revealed that these 540

upregulated genes significantly enriched pathways related to type II interferon production, cytokine

production, and NFkB and JNK signaling (Fig. S10H). We also observed that, with the 5-day LPS

stimulation, pMacs from aged *R1441C* male mice could be pushed to exhibit phenotypes that paralleled

those seen in aged female *R1441C* macrophages. It has previously been reported that sustained and chronic

STAT1 signaling, known to be stimulated by cytokine signaling (*50*), is necessary to induce macrophage

exhaustion (*42*). Collectively, we interpret our findings that the upregulated DEGs, including driving

increased cytokine release in *R1441C* aged male macrophages, may predispose these macrophages to

become exhausted upon LPS-stimulation.

***R1441C*- and *Y1699C* LRRK2 monocyte-derived patient macrophages exhibit hypophagocytosis and**

increased markers for exhaustion

To assess the effects of PD-associated mutations on immune exhaustion in humans carrying *LRRK2*

mutations, pathogen phagocytosis was quantified in monocyte-derived macrophages from 2 PD-*Y1699C*, 1

PD-*R1441C*, 7 healthy controls, and 3 non-manifesting carriers (NMC) of the *R1441C* pathogenic variant

(Fig. 7A). Both *R1441C* and *Y1699C* variants are located in the GTPase domain of the LRRK2 protein, and

both have been shown to perturb LRRK2's GTPase activity and have been reported to elevate its kinase

activity (*8, 16, 17, 51, 52*). A hypophagocytic phenotype was observed in monocyte-derived macrophages

relative to healthy controls and *R1441C* NMCs, as indicated by a reduced number of GFP$^+$ cells (Fig. 7B,

C), with no differences in GFP MFI observed between groups (Fig. S11A). Next, increased markers of

immune exhaustion accompanied reduced rates of phagocytosis, PD1 expression was assessed in peripheral

blood mononuclear cells (PBMCs). Consistent with our mouse data, we observed that PBMCs from both

PD-*Y1699C* and PD-*R1441C* patients exhibited increased PD1$^+$ cell count and expression levels (Fig. 7D,

E). When quantifying PD1 expression in specific immune-cell subsets (Fig. S11B), it was determined that

this overall increase in PD1 expression was driven by expression in classical monocytes, with increased

PD1 expression observed in PD-*Y1699C* and *R1441C* classical monocytes relative to healthy controls and

R1441C NMCs (Fig. 7F), with no significant differences observed in other cell types (Fig. S11C-E).

Similarly, increased PD1 expression was evident in monocyte-derived macrophages from PD-*Y1699C* and

R1441C patients relative to healthy controls and *R1441C* NMCs (Fig. 7G). Collectively, the immune

exhaustion phenotype observed in aged *R1441C* female mice is also associated with an exhaustion

phenotype in cell from patients harboring the *LRRK2-Y1699C* or *R1441C* mutation.

DISCUSSION

Considering age is the greatest risk factor for many neurodegenerative diseases, aging, in particular aging

of the immune system, is the most underappreciated and understudied contributing factor in the

neurodegeneration field. Our novel findings reveal a bi-phasic age- and sex-dependent effect of the *R1441C*

mutation in peripheral immune cells, with increased cytokine release, antigen presentation, phagocytosis

and lysosomal activity in macrophages from young female *R1441C* mice, with many of these phenotypes

ameliorated by Lrrk2 protein kinase inhibition. Our findings further reveal an age-acquired immune cell

exhaustion in macrophages from *R1441C* females associated with increased Lrrk2 protein kinase activity,

hypophagocytosis, decreased antigen presentation, and reduced lysosomal pan-cathepsin activation, protein

degradation and increased pathogen-mediated DNA breakage, and increased PD1 expression indicative of

immune cell exhaustion. Importantly, this immune cell exhaustion phenotype was also observed in

monocytes and macrophages from human subjects with PD and harboring the *R1441C* and the *Y1699C*

mutation. Furthermore, it was demonstrated that, although macrophages from male mice did not exhibit

this bi-phasic age-dependent phenotype, they could be pushed to exhaustion with chronic LPS stimulation,

suggesting that, along with age, a secondary inflammatory insult is required to observe immune cell

exhaustion phenotypes in macrophages from *R1441C* males. Finally, it was demonstrated that there is an

age-acquired dysregulation in the expression of genes associated with PD risk, mitochondrial function, and

oxidative stress in macrophages from female *R1441C* mice.

Consistent with findings in *ex vivo* studies in macrophages, we consistently observed elevated levels of

both IL10 and IL6 in circulating plasma from young *R1441C* mice. It has recently been shown that

RAW264.7 cells expressing *T1348N-LRRK2*, an artificial P-Loop null mutation that disrupts GTP binding

(*53, 54*), produce significantly less IL10 relative to wild-type cells in response to LPS and Zymosan (*55*).

IL10 is produced by macrophages and is critical in limiting immune-mediated pathology (*56*), and it was

therefore suggested by Nazish *et al* that there may be a neuroprotective role of LRRK2 in immune signaling

through regulation of IL10 secretion. In addition, we also observed elevations in IL6, which is known to be

promptly and transiently produced in response to infections and tissue injuries during host defense through

the stimulation of immune reactions (*57*). These observations implicate a protective role of Lrrk2 in young

R1441C mice. The pro-inflammatory cytokine, TNF, was also increased in circulating plasma of young

R1441C mice, a compensatory increase that may also contribute to organisms ability to fight infection given

that mice lacking TNF exhibit increased susceptibility to infection with increased bacterial load (*58*). In

support of the protective role of LRRK2 in immune signaling, LRRK2 has been demonstrated to provide

increased pathogen control of *Salmonella typhimurium (26, 59)* and *Listeria monocytogenes* (*27*).

Consistent with these reports, we observed increased *E. coli* uptake and decreased *E. coli*-mediated DNA

damage in macrophages from young *R1441C* female mice, providing further support for a protective role

of LRRK2 in immune signaling and defense against pathogens, at least in a young organism. Given that

LRRK2 mutations are associated with not only PD, but also Crohn's Disease (CD) (*2, 60, 61*) and leprosy

(*62*), it is difficult to understand why these mutations would have persisted in the human population. Given

the evidence discussed and the data presented here, it is plausible to speculate that these mutations may

have persisted due to the immunological advantage they confer to young organisms, including protection

against certain infections.

For the first time, we have reported here an age-dependent, bi-phasic effect of LRRK2 on immune cell

function, with the *R1441C* mutation inducing myeloid cell immune exhaustion in a sex-dependent manner.

The concept of immune cell exhaustion was first described as the clonal deletion of virus-specific CD8[+] T-

cells that occurs during high-grade chronic infections, and in T-cells is characterized by the stepwise and

progressive loss of T-cell functions (*63*). Regarding exhausted myeloid cells, pathogenic inflammation and

immunosuppression, characterized by decreased antigen presentation, hypophagocytosis and altered

cytokine release, have been described more recently (*42*). The immune system is composed of numerous

cell types with varying states of activation, and cell-to-cell communication during immune responses is

crucial for host defense and resolution of inflammation. The observations made here in innate immune cells,

therefore, likely have downstream effects on myeloid cell communication with adaptive immune cell types

in the periphery. Indeed, alterations in the adaptive immune system have been reported in PD patients. For

example, a meta-analysis of 943 cases of PD identified a reduction in circulating CD4[+] T cells in PD-

patients (*64, 65*). More specifically, HLA-DR[+] T cells and CD45RO[+] memory T cells have been shown to

be increased in people with PD relative to healthy age- and sex-matched controls, while naive CD4[+] T cells

are reduced (*65-67*). CD4[+]FOXP3[+] Treg cells have increased suppressive activity in patients with PD (*68*).

Interestingly, one characteristic of immune exhaustion is the dysregulated cross-talk between CD4[+] and

CD8[+] T-cells due to the suppressive action of Tregs (*69*). Furthermore, Tregs directly inhibit the antigen-

presenting function of monocytes and macrophages (*70*). Further research is required to understand the role

of cell-to-cell communication in these myeloid cell phenotypes, and if immune cell exhaustion driven by

LRRK2 is a contributing factor.

A sex-dependent effect of the *R1441C* mutation was reported here, with females exhibiting an age-acquired

immune cell exhaustion phenotype which could only be elicited in macrophages from males upon a

secondary, chronic inflammatory insult. Sex-dependent effects of *LRRK2* mutations have previously been

reported. A protective effect of the *G2019S* mutation against *S. typhimurium* has been observed in mice,

with female mutant mice showing increased survival relative to B6 controls which was not observed in

males (*59*). Such observations are also consistent with the sexual dimorphism reported in subjects with PD

who also carry a heterozygous p.*G2019S* mutation in *LRRK2*, with female sex being more common amongst

carriers than non-carriers (*71-73*). This sexual dimorphism has also been reported for the *R1441C/G/H*

mutations (*72*). We observed an *R1441C*-dependent phenotype in macrophages from male mice only when

a secondary, chronic inflammatory insult was applied. It is particularly interesting that genes that are

significantly upregulated with age in *R1441C* macrophages from male mice are associated with pathways

which are heavily implicated in the phenotypes we observed in macrophages from young *R1441C* females.

An increase in gene expression in pathways associated with cytokine production is consistent with our

previous observation of increased IL6, IL10 and TNF release in aged male pMacs relative to B6 controls.

Further research is therefore required to determine the extent to which this sexual dimorphism is driven by

altered immune system susceptibility to *LRRK2* mutations in males and females.

We observed alterations in genes enriched in the KEGG Parkinson's Disease pathway in *R1441C* female

macrophages as a function of age, with these genes being mostly associated with mitochondrial respiration

and oxidative stress. Not surprisingly, mitochondrial dysfunction has emerged as one of the key

mechanisms underlying the pathogenesis of both sporadic PD and familial Parkinsonism (*74, 75*). LRRK2

has been heavily implicated in mitochondrial function and mitophagy, with *LRRK2* mutations causing

reduced mitochondria potential and decreased ATP production, mitochondrial DNA damage, mitochondrial

fragmentation and altered rates of mitophagy in various cell types and pre-clinical models (*76-79*). But

importantly, our findings herein reveal that pathways once thought to be primarily important to neuronal

demise (mitochondrial and oxidative stress) are also disrupted in peripheral immune cells by PD-associated

mutations in a sex- and age-dependent manner and have important implications for monitoring risk, target

engagement, and disease progression of PD and perhaps other NDs by assessing peripheral immune cell

types *ex vivo*. It is known that, in T cells, mitochondrial stress induced by chronic antigen stimulation drives

immune exhaustion (*80*). Further research is therefore required to fully understand the role of mitochondria

and mitophagy, and therefore lysosomal function, in the *R1441C*-mediated immune cell exhaustion

phenotypes observed here.

It has previously been shown that LRRK2 exerts differential effects on inflammatory responses in a tissue-

specific manner, and it has been suggested that there is a differential role of LRRK2 in the periphery versus

the central nervous system (CNS) (*59, 81*). It therefore remains to be determined how the data reported here

impacts our understanding of LRRK2 in myeloid cells in both the periphery and the CNS. However,

alterations in type II interferon responses have been observed in both induced pluripotent stem cell (iPSC)-

derived microglia and macrophages from patients harboring *G2019S* mutations (*82*), and similar effects of

LRRK2 mutations on lysosomal function have also been reported in patient iPSC-derived microglia and

macrophages (*83*). However, it has been demonstrated that expression of *R1441C*-Lrrk2 in peripheral

lymphocytes alone is sufficient to induce dopaminergic neuronal cell loss in the midbrains of LPS-treated

mice (*33*), suggesting that the periphery plays a pertinent and perhaps primary role in *LRRK2*-PD, and

future research into how immune exhaustion in the periphery impacts cross-talk with the CNS is needed to

inform future immunomodulatory therapeutic interventions for PD.

We successfully demonstrated that, like macrophages from aged *R1441C* mice, myeloid cells from *R1441C*

and *Y1699C* patients exhibited hypophagocytosis and increased expression of immune exhaustion markers

relative to those from HCs and *R1441C*-NMCs. However, it is unclear if the same hyper-responsive

phenotype that was observed in young *R1441C* female mice is exhibited in myeloid cells in *R1441C* carriers

at a younger age or prior to disease onset. Indeed, we did show that *R1441C*-NMCs did not exhibit these

immune exhaustion phenotypes, and it is noteworthy that the average age of the 3 *R1441C*-NMCs in this

study was 30.5 years (see supplementary material), whilst the average age of the *R1441C* and *Y1699C*-PD

subjects was 42 and 63.5, respectively. Further investigation is therefore required to determine if PBMCs

from NMCs, upon follow-up after further aging or disease onset, exhibit immune exhaustion. Furthermore,

further research is required to understand if these phenotypes are exclusive to *LRRK2*-pathogenic variants

located in the GTPase domain or if they are seen in other *LRRK2* mutant immune cells as well as in

idiopathic PD or sporadic cases.

LRRK2-targeting therapies, including antisense oligonucleotides (ASOs) and kinase inhibitors, have been

suggested as a potential therapeutic for *LRRK2*-PD, with many hypothesizing that targeting the increased

LRRK2 levels and kinase activity will be beneficial to CNS-related pathology. Our findings suggest that

more research is required to understand the location and optimal timing for the administration of these

LRRK2-targeting therapeutics. Specifically, if LRRK2 plays a critical role in pathogen and infection control

in a pathogen or tissue-specific manner, it may mean that LRRK2-targeting therapies may need to be limited

to carriers of *LRRK2* pathogenic variants and/or may need to be delivered in a CNS compartment-targeted

manner as opposed to systemically to avoid deleterious collateral damage to immune function. Indeed, the

LRRK2 ASO currently being tested in the clinic is being delivered via intrathecal injection directly into the

cisterna magna, thus limiting systemic exposure (NCT03976349). Lastly, perhaps somewhat surprisingly,

we found that while LRRK2 kinase inhibition could rescue some of the effector functions of aged mutant-

LRRK2 mice, it had no beneficial effects on the immune-cell exhaustion phenotypes observed here in

macrophages from aged mice; however, it did prove to be protective in macrophages from young mice to

prevent the development of an immune-cell exhaustive phenotype. Such observations suggest that LRRK2-

targeting therapies would elicit the most beneficial effects during the prodromal phases of disease, and

further research on PBMCs from prodromal patients with pre-motor features at risk for development of

motor PD due to a family history will be required to investigate this possibility and inform future clinical

trials.

MATERIALS AND METHODS

Study design

The primary objective of this research was to investigate the effects of the *R1441C LRRK2* pathogenic variant on myeloid cell effector function in a sex- and age-dependent manner. We hypothesized that a biphasic, age-dependent effect of the *R1441C* pathogenic variant would be observed, affecting antigen presentation, cytokine release, phagocytosis and lysosomal function in macrophages. To explore our hypothesis, we used *R1441C* knock-in mice and B6 controls and assessed myeloid cell effector function both *ex vivo* and *in vivo,* in males and females at 2-3 months and 18-21 months. All studies were performed by investigators blinded to genotype and treatment groups at the University of Florida, USA. Power analyses based on the labs pilot and published data indicated that n = 12 per group for flow-cytometry and cytokine release assays, and n = 6 for imaging based immunophenotyping and immunoblotting would be needed to yield 80% power with a α set to 0.05. The study used human PBMCs received from collaborators at University Hospital Schleswig-Holstein, University of Luebeck, Germany and the University Hospital Eppendorf, Hamburg, Germany. Samples were received at the University of Florida and investigators were unblinded after completion of analysis. Deidentified data provided included sex, age at collection, age of diagnosis, clinical diagnosis and genetic status.

Animals

R1441C knock-in (Strain #009346; B6.Cg-Lrrk2tm1.1Shn/J) mice have previously been characterized (*84*) and were maintained in the McKnight Brain Institute vivarium (University of Florida) at 22°C at 60-70% humidity with a 12-hour light/dark cycle and *ad libitum* access to standard rodent chow and water. C57BL/6 (Strain #000664) controls were used for all studies. All animal procedures were approved by the University of Florida Institutional Animal Care and Use Committee and were in accordance with the National Institute of Health *Guide for the Care and Use of Laboratory Animals* (NIH Publications No. 80-23) revised 1996. Female and male mice were aged to 2-3 or 18-21-months-old and sacrificed via cervical dislocation or rapid decapitation.

Human subjects

This study was reviewed and approved by the University of Lubeck Institutional Review Board (AZ16-039). Participants provided written informed consent to participate. During recruitment, a family history was used to assess history of disease and patients were screened for *LRRK2* variants. See Table S3 for patient demographics.

Statistics and data analysis

Data and statistical analyses were performed using IBM SPSS statistics 27 or GraphPad Prism 9. For assessing differences between groups, data were analyzed by either 1-way or 2-way analysis of variance (ANOVA), or by t-test. In instances when data did not fit parametric assumptions, Kruskal-Wallis non-parametric ANOVA was used. *Post hoc* tests following ANOVAs were conducted using Tukey HSD or Bonferroni correction. Two-tailed levels of significance were used and $p < 0.05$ was considered statistically significant. Graphs are depicted by means +/- standard error of the mean (SEM).

References

1. Von Campenhausen, S., B. Bornschein, R. Wick, K. Bötzel, C. Sampaio, W. Poewe, W. Oertel, U. Siebert, K. Berger, and R. Dodel, Prevalence and incidence of Parkinson's disease in Europe. **15**(4): p. 473-490 (2005).
2. Marras, C., J.C. Beck, J.H. Bower, E. Roberts, B. Ritz, G.W. Ross, R.D. Abbott, R. Savica, S.K. Van Den Eeden, A.W. Willis, C.M. Tanner, and P.G. Parkinson's Foundation, Prevalence of Parkinson's disease across North America. *NPJ Parkinsons Dis*. **4**: p. 21 (2018).
3. Yang, W., J.L. Hamilton, C. Kopil, J.C. Beck, C.M. Tanner, R.L. Albin, E. Ray Dorsey, N. Dahodwala, I. Cintina, P. Hogan, and T. Thompson, Current and projected future economic burden of Parkinson's disease in the U.S. *npj Parkinson's Disease*. **6**(1) (2020).
4. Bloem, B.R., M.S. Okun, and C. Klein, Parkinson's disease. *Lancet*. **397**(10291): p. 2284-2303 (2021).
5. Tansey, M.G., Wallings, R. L., Houser, M. C., Herrick, M. K., Keating, C. E. & Joers, V., The contributions of inflammation and immune dysfunction to Parkinson disease. *Nature Reviews Immunology*. (2022).
6. Blauwendraat, C., M.A. Nalls, and A.B. Singleton, The genetic architecture of Parkinson's disease. *Lancet Neurol*. **19**(2): p. 170-178 (2020).
7. West, A.B., D.J. Moore, S. Biskup, A. Bugayenko, W.W. Smith, C.A. Ross, V.L. Dawson, and T.M. Dawson, Parkinson's disease-associated mutations in leucine-rich repeat kinase 2 augment kinase activity. *Proc Natl Acad Sci U S A*. **102**(46): p. 16842-7 (2005).

8.   West, A.B., D.J. Moore, C. Choi, S.A. Andrabi, X. Li, D. Dikeman, S. Biskup, Z. Zhang, K.L. Lim, V.L.
Dawson, and T.M. Dawson, Parkinson's disease-associated mutations in LRRK2 link enhanced GTP-binding
and kinase activities to neuronal toxicity. *Hum Mol Genet*. **16**(2): p. 223-32 (2007).

9.   Funayama, M., K. Hasegawa, H. Kowa, M. Saito, S. Tsuji, and F. Obata, A new locus for Parkinson's disease
(PARK8) maps to chromosome 12p11.2-q13.1. *Ann Neurol*. **51**(3): p. 296-301 (2002).

10.   Funayama, M., Y. Li, H. Tomiyama, H. Yoshino, Y. Imamichi, M. Yamamoto, M. Murata, T. Toda, Y.
Mizuno, and N. Hattori, Leucine-rich repeat kinase 2 G2385R variant is a risk factor for Parkinson disease
in Asian population. *Neuroreport*. **18**(3): p. 273-5 (2007).

11.   Hatano, T., M. Funayama, S.I. Kubo, I.F. Mata, Y. Oji, A. Mori, C.P. Zabetian, S.M. Waldherr, H. Yoshino,
G. Oyama, Y. Shimo, K.I. Fujimoto, H. Oshima, Y. Kunii, H. Yabe, Y. Mizuno, and N. Hattori, Identification
of a Japanese family with LRRK2 p.R1441G-related Parkinson's disease. *Neurobiol Aging*. **35**(11): p. 2656
e17-2656 e23 (2014).

12.   Ross, O.A., Y.R. Wu, M.C. Lee, M. Funayama, M.L. Chen, A.I. Soto, I.F. Mata, G.J. Lee-Chen, C.M. Chen,
M. Tang, Y. Zhao, N. Hattori, M.J. Farrer, E.K. Tan, and R.M. Wu, Analysis of Lrrk2 R1628P as a risk factor
for Parkinson's disease. *Ann Neurol*. **64**(1): p. 88-92 (2008).

13.   Haugarvoll, K. and Z.K. Wszolek, Clinical features of LRRK2 parkinsonism. **15**: p. S205-S208 (2009).

14.   Satake, W., Y. Nakabayashi, I. Mizuta, Y. Hirota, C. Ito, M. Kubo, T. Kawaguchi, T. Tsunoda, M. Watanabe,
A. Takeda, H. Tomiyama, K. Nakashima, K. Hasegawa, F. Obata, T. Yoshikawa, H. Kawakami, S. Sakoda,
M. Yamamoto, N. Hattori, M. Murata, Y. Nakamura, and T. Toda, Genome-wide association study identifies
common variants at four loci as genetic risk factors for Parkinson's disease. *Nat Genet*. **41**(12): p. 1303-7
(2009).

15.   Gaig, C., M. Ezquerra, M.J. Marti, E. Muñoz, F. Valldeoriola, and E. Tolosa, LRRK2 Mutations in Spanish
Patients With Parkinson Disease. *Archives of Neurology*. **63**(3): p. 377 (2006).

16.   Steger, M., F. Tonelli, G. Ito, P. Davies, M. Trost, M. Vetter, S. Wachter, E. Lorentzen, G. Duddy, S. Wilson,
M.A. Baptista, B.K. Fiske, M.J. Fell, J.A. Morrow, A.D. Reith, D.R. Alessi, and M. Mann,
Phosphoproteomics reveals that Parkinson's disease kinase LRRK2 regulates a subset of Rab GTPases. *Elife*.
5 (2016).

17.   Lewis, P.A., E. Greggio, A. Beilina, S. Jain, A. Baker, and M.R. Cookson, The R1441C mutation of LRRK2
disrupts GTP hydrolysis. **357**(3): p. 668-671 (2007).

18.   Cook, D.A., G.T. Kannarkat, A.F. Cintron, L.M. Butkovich, K.B. Fraser, J. Chang, N. Grigoryan, S.A.
Factor, A.B. West, J.M. Boss, and M.G. Tansey, LRRK2 levels in immune cells are increased in Parkinson's
disease. *NPJ Parkinsons Dis*. **3**: p. 11 (2017).

19.   Gardet, A., Y. Benita, C. Li, B.E. Sands, I. Ballester, C. Stevens, J.R. Korzenik, J.D. Rioux, M.J. Daly, R.J.
Xavier, and D.K. Podolsky, LRRK2 Is Involved in the IFN- Response and Host Response to Pathogens. *The*
Journal of Immunology. **185**(9): p. 5577-5585 (2010).

20.   Atashrazm, F., D. Hammond, G. Perera, M.F. Bolliger, E. Matar, G.M. Halliday, B. Schule, S.J.G. Lewis,
R.J. Nichols, and N. Dzamko, LRRK2-mediated Rab10 phosphorylation in immune cells from Parkinson's
disease patients. *Mov Disord*. **34**(3): p. 406-415 (2019).

21.   Wallings, R.L., L.P. Hughes, H.A. Staley, Z.D. Simon, N.R. McFarland, R.N. Alcalay, A. Garrido, M.J.
Marti, E.T. Sarro, N. Dzamko, and M.G. Tansey, WHOPPA Enables Parallel Assessment of Leucine-Rich
Repeat Kinase 2 and Glucocerebrosidase Enzymatic Activity in Parkinson's Disease Monocytes. *Front Cell*
Neurosci. **16**: p. 892899 (2022).

22.   Kozina, E., S. Sadasivan, Y. Jiao, Y. Dou, Z. Ma, H. Tan, K. Kodali, T. Shaw, J. Peng, and R.J. Smeyne,
Mutant LRRK2 mediates peripheral and central immune responses leading to neurodegeneration in vivo.
Brain. **141**(6): p. 1753-1769 (2018).

23.   Rideout, H.J. and D.B. Re, LRRK2 and the "LRRKtosome" at the Crossroads of Programmed Cell Death:
Clues from RIP Kinase Relatives. *Adv Neurobiol*. **14**: p. 193-208 (2017).

24.   Wang, Z., S. Arat, M. Magid-Slav, and J.R. Brown, Meta-analysis of human gene expression in response to
Mycobacterium tuberculosis infection reveals potential therapeutic targets. *BMC Systems Biology*. **12**(1)
(2018).

25.   Hartlova, A., S. Herbst, J. Peltier, A. Rodgers, O. Bilkei-Gorzo, A. Fearns, B.D. Dill, H. Lee, R. Flynn, S.A.
Cowley, P. Davies, P.A. Lewis, I.G. Ganley, J. Martinez, D.R. Alessi, A.D. Reith, M. Trost, and M.G.
Gutierrez, LRRK2 is a negative regulator of Mycobacterium tuberculosis phagosome maturation in
macrophages. *EMBO J*. **37**(12) (2018).

26. Liu, W., X.N. Liu, Y. Li, J. Zhao, Z. Liu, Z. Hu, Y. Wang, Y. Yao, A.W. Miller, B. Su, M.R. Cookson, X.
Li, and Z. Kang, LRRK2 promotes the activation of NLRC4 inflammasome during Salmonella Typhimurium
infection. *The Journal of Experimental Medicine*. **214**(10): p. 3051-3066 (2017).

27. Zhang, Q., Y. Pan, R. Yan, B. Zeng, H. Wang, X. Zhang, W. Li, H. Wei, and Z. Liu, Commensal bacteria
direct selective cargo sorting to promote symbiosis. **16**(9): p. 918-926 (2015).

28. Fava, V.M., J. Manry, A. Cobat, M. Orlova, N. Van Thuc, N.N. Ba, V.H. Thai, L. Abel, A. Alcaïs, and E.
Schurr, A Missense LRRK2 Variant Is a Risk Factor for Excessive Inflammatory Responses in Leprosy.
PLOS Neglected Tropical Diseases. **10**(2): p. e0004412 (2016).

29. Witoelar, A., I.E. Jansen, Y. Wang, R.S. Desikan, J.R. Gibbs, C. Blauwendraat, W.K. Thompson, D.G.
Hernandez, S. Djurovic, A.J. Schork, F. Bettella, D. Ellinghaus, A. Franke, B.A. Lie, L.K. McEvoy, T.H.
Karlsen, S. Lesage, H.R. Morris, A. Brice, N.W. Wood, P. Heutink, J. Hardy, A.B. Singleton, A.M. Dale, T.
Gasser, O.A. Andreassen, and M. Sharma, Genome-wide Pleiotropy Between Parkinson Disease and
Autoimmune Diseases. *JAMA Neurology*. **74**(7): p. 780 (2017).

30. Villumsen, M., S. Aznar, B. Pakkenberg, T. Jess, and T. Brudek, Inflammatory bowel disease increases the
risk of Parkinson's disease: a Danish nationwide cohort study 1977–2014. *Gut*. **68**(1): p. 18-24 (2019).

31. Skaper, S.D., L. Facci, M. Zusso, and P. Giusti, An Inflammation-Centric View of Neurological Disease:
Beyond the Neuron. *Frontiers in Cellular Neuroscience*. **12** (2018).

32. Gelders, G., V. Baekelandt, and A. Van der Perren, Linking Neuroinflammation and Neurodegeneration in
Parkinson's Disease. *J Immunol Res*. **2018**: p. 4784268 (2018).

33. Kozina, E., M. Byrne, and R.J. Smeyne, Mutant LRRK2 in lymphocytes regulates neurodegeneration via IL-
6 in an inflammatory model of Parkinson's disease. *npj Parkinson's Disease*. **8**(1) (2022).

34. Kim, K.S., P.C. Marcogliese, J. Yang, S.M. Callaghan, V. Resende, E. Abdel-Messih, C. Marras, N.P.
Visanji, J. Huang, M.G. Schlossmacher, L. Trinkle-Mulcahy, R.S. Slack, A.E. Lang, and D.S. Park,
Regulation of myeloid cell phagocytosis by LRRK2 via WAVE2 complex stabilization is altered in
Parkinson's disease. *Proceedings of the National Academy of Sciences*. **115**(22): p. E5164-E5173 (2018).

35. Liu, Z., E. Xu, H.T. Zhao, T. Cole, and A.B. West, LRRK2 and Rab10 coordinate macropinocytosis to
mediate immunological responses in phagocytes. *EMBO J*. **39**(20): p. e104862 (2020).

36. Tong, Y., E. Giaime, H. Yamaguchi, T. Ichimura, Y. Liu, H. Si, H. Cai, J.V. Bonventre, and J. Shen, Loss of
leucine-rich repeat kinase 2 causes age-dependent bi-phasic alterations of the autophagy pathway. **7**(1): p. 2
(2012).

37. Cassado, A.D.A., M.R. D'Império Lima, and K.R. Bortoluci, Revisiting Mouse Peritoneal
Macrophages: Heterogeneity, Development, and Function. *Frontiers in Immunology*. **6** (2015).

38. Schapansky, J., J.D. Nardozzi, F. Felizia, and M.J. Lavoie, Membrane recruitment of endogenous LRRK2
precedes its potent regulation of autophagy. **23**(16): p. 4201-4214 (2014).

39. Hakimi, M., T. Selvanantham, E. Swinton, R.F. Padmore, Y. Tong, G. Kabbach, K. Venderova, S.E.
Girardin, D.E. Bulman, C.R. Scherzer, M.J. LaVoie, D. Gris, D.S. Park, J.B. Angel, J. Shen, D.J. Philpott,
and M.G. Schlossmacher, Parkinson's disease-linked LRRK2 is expressed in circulating and tissue immune
cells and upregulated following recognition of microbial structures. *J Neural Transm (Vienna)*. **118**(5): p.
795-808 (2011).

40. Moehle, M.S., J.P.L. Daher, T.D. Hull, R. Boddu, H.A. Abdelmotilib, J. Mobley, G.T. Kannarkat, M.G.
Tansey, and A.B. West, The G2019S LRRK2 mutation increases myeloid cell chemotactic responses and
enhances LRRK2 binding to actin-regulatory proteins. **24**(15): p. 4250-4267 (2015).

41. Kaminski, H., M. Lemoine, and T. Pradeu, Immunological exhaustion: How to make a disparate concept
operational? *PLOS Pathogens*. **17**(9): p. e1009892 (2021).

42. Pradhan, K., Z. Yi, S. Geng, and L. Li, Development of Exhausted Memory Monocytes and Underlying
Mechanisms. *Front Immunol*. **12**: p. 778830 (2021).

43. Strader, S. and A.B. West, The interplay between monocytes, alpha-synuclein and LRRK2 in Parkinson's
disease. *Biochem Soc Trans*. **51**(2): p. 747-758 (2023).

44. Jin, J., H. Zhang, C.M. Weyand, and J.J. Goronzy, Lysosomes in T Cell Immunity and Aging. *Front Aging*.
2: p. 809539 (2021).

45. Hennies, C.M., M.A. Lehn, and E.M. Janssen, Quantitating MHC class II trafficking in primary dendritic
cells using imaging flow cytometry. *J Immunol Methods*. **423**: p. 18-28 (2015).

46. Watts, C., The endosome-lysosome pathway and information generation in the immune system. *Biochim
Biophys Acta*. **1824**(1): p. 14-21 (2012).

47. Boomer, J.S., J.M. Green, and R.S. Hotchkiss, The changing immune system in sepsis. *Virulence*. **5**(1): p.
45-56 (2014).

48.   Jubel, J.M., Z.R. Barbati, C. Burger, D.C. Wirtz, and F.A. Schildberg, The Role of PD-1 in Acute and Chronic
791   Infection. *Front Immunol*. **11**: p. 487 (2020).
792   49.   Weinstock, A., K. Rahman, O. Yaacov, H. Nishi, P. Menon, C.A. Nikain, M.L. Garabedian, S. Pena, N.
793   Akbar, B.E. Sansbury, S.P. Heffron, J. Liu, G. Marecki, D. Fernandez, E.J. Brown, K.V. Ruggles, S.A.
794   Ramsey, C. Giannarelli, M. Spite, R.P. Choudhury, P. Loke, and E.A. Fisher, Wnt signaling enhances
795   macrophage responses to IL-4 and promotes resolution of atherosclerosis. *Elife*. **10** (2021).
796   50.   Morris, R., N.J. Kershaw, and J.J. Babon, The molecular details of cytokine signaling via the JAK/STAT
797   pathway. *Protein Science*. **27**(12): p. 1984-2009 (2018).
798   51.   West, A.B., D.J. Moore, S. Biskup, A. Bugayenko, W.W. Smith, C.A. Ross, V.L. Dawson, and T.M. Dawson,
799   From The Cover: Parkinson's disease-associated mutations in leucine-rich repeat kinase 2 augment kinase
800   activity. **102**(46): p. 16842-16847 (2005).
801   52.   Daniëls, V., R. Vancraenenbroeck, B.M.H. Law, E. Greggio, E. Lobbestael, F. Gao, M. De Maeyer, M.R.
802   Cookson, K. Harvey, V. Baekelandt, and J.-M. Taymans, Insight into the mode of action of the LRRK2
803   Y1699C pathogenic mutant. **116**(2): p. 304-315 (2011).
804   53.   Biosa, A., A. Trancikova, L. Civiero, L. Glauser, L. Bubacco, E. Greggio, and D.J. Moore, GTPase activity
805   regulates kinase activity and cellular phenotypes of Parkinson's disease-associated LRRK2. **22**(6): p. 1140-
806   1156 (2013).
807   54.   Nguyen, A.P. and D.J. Moore, Understanding the GTPase Activity of LRRK2: Regulation, Function, and
808   Neurotoxicity. *Adv Neurobiol*. **14**: p. 71-88 (2017).
809   55.   Nazish, I., C. Arber, T.M. Piers, T.T. Warner, J.A. Hardy, P.A. Lewis, J.M. Pocock, and R. Bandopadhyay,
810   Abrogation of LRRK2 dependent Rab10 phosphorylation with TLR4 activation and alterations in evoked
811   cytokine release in immune cells. *Neurochem Int*. **147**: p. 105070 (2021).
812   56.   Sanin, D.E., C.T. Prendergast, and A.P. Mountford, IL-10 Production in Macrophages Is Regulated by a
813   TLR-Driven CREB-Mediated Mechanism That Is Linked to Genes Involved in Cell Metabolism. *J Immunol*.
814   **195**(3): p. 1218-32 (2015).
815   57.   Tanaka, T., M. Narazaki, and T. Kishimoto, IL-6 in inflammation, immunity, and disease. *Cold Spring Harb
816   Perspect Biol*. **6**(10): p. a016295 (2014).
817   58.   Li, X., H. Korner, and X. Liu, Susceptibility to Intracellular Infections: Contributions of TNF to Immune
818   Defense. *Front Microbiol*. **11**: p. 1643 (2020).
819   59.   Shutinoski, B., M. Hakimi, I.E. Harmsen, M. Lunn, J. Rocha, N. Lengacher, Y.Y. Zhou, J. Khan, A. Nguyen,
820   Q. Hake-Volling, D. El-Kodsi, J. Li, A. Alikashani, C. Beauchamp, J. Majithia, K. Coombs, D. Shimshek,
821   P.C. Marcogliese, D.S. Park, J.D. Rioux, D.J. Philpott, J.M. Woulfe, S. Hayley, S. Sad, J.J. Tomlinson, E.G.
822   Brown, and M.G. Schlossmacher, Lrrk2 alleles modulate inflammation during microbial infection of mice in
823   a sex-dependent manner. *Science Translational Medicine*. **11**(511): p. eaas9292 (2019).
824   60.   Herrick, M.K. and M.G. Tansey, Is LRRK2 the missing link between inflammatory bowel disease and
825   Parkinson's disease? *npj Parkinson's Disease*. **7**(1) (2021).
826   61.   Daseke, M.J., 2nd, M.A.A. Tenkorang-Impraim, Y. Ma, U. Chalise, S.R. Konfrst, M.R. Garrett, K.Y.
827   DeLeon-Pennell, and M.L. Lindsey, Exogenous IL-4 shuts off pro-inflammation in neutrophils while
828   stimulating anti-inflammation in macrophages to induce neutrophil phagocytosis following myocardial
829   infarction. *J Mol Cell Cardiol*. **145**: p. 112-121 (2020).
830   62.   Fava, V., M. Orlova, A. Cobat, A. Alcaïs, M. Mira, and E. Schurr, Genetics of leprosy reactions: an overview.
831   **107**: p. 132-142 (2012).
832   63.   Yi, J.S., M.A. Cox, and A.J. Zajac, T-cell exhaustion: characteristics, causes and conversion. *Immunology*.
833   **129**(4): p. 474-81 (2010).
834   64.   Jiang, S., H. Gao, Q. Luo, P. Wang, and X. Yang, The correlation of lymphocyte subsets, natural killer cell,
835   and Parkinson's disease: a meta-analysis. *Neurol Sci*. **38**(8): p. 1373-1380 (2017).
836   65.   Saunders, J.A., K.A. Estes, L.M. Kosloski, H.E. Allen, K.M. Dempsey, D.R. Torres-Russotto, J.L. Meza,
837   P.M. Santamaria, J.M. Bertoni, D.L. Murman, H.H. Ali, D.G. Standaert, R.L. Mosley, and H.E. Gendelman,
838   CD4+ regulatory and effector/memory T cell subsets profile motor dysfunction in Parkinson's disease. *J
839   Neuroimmune Pharmacol*. **7**(4): p. 927-38 (2012).
840   66.   Fiszer, U., E. Mix, S. Fredrikson, V. Kostulas, and H. Link, Parkinson's disease and immunological
841   abnormalities: increase of HLA-DR expression on monocytes in cerebrospinal fluid and of CD45RO+ T cells
842   in peripheral blood. *Acta Neurol Scand*. **90**(3): p. 160-6 (1994).
843   67.   Kustrimovic, N., C. Comi, L. Magistrelli, E. Rasini, M. Legnaro, R. Bombelli, I. Aleksic, F. Blandini, B.
844   Minafra, G. Riboldazzi, A. Sturchio, M. Mauri, G. Bono, F. Marino, and M. Cosentino, Parkinson's disease

patients have a complex phenotypic and functional Th1 bias: cross-sectional studies of CD4+ Th1/Th2/T17
and Treg in drug-naive and drug-treated patients. *J Neuroinflammation*. **15**(1): p. 205 (2018).

68. Rosenkranz, D., S. Weyer, E. Tolosa, A. Gaenslen, D. Berg, T. Leyhe, T. Gasser, and L. Stoltze, Higher
frequency of regulatory T cells in the elderly and increased suppressive activity in neurodegeneration. *J
Neuroimmunol*. **188**(1-2): p. 117-27 (2007).

69. McRitchie, B.R. and B. Akkaya, Exhaust the exhausters: Targeting regulatory T cells in the tumor
microenvironment. *Front Immunol*. **13**: p. 940052 (2022).

70. Tiemessen, M.M., A.L. Jagger, H.G. Evans, M.J. van Herwijnen, S. John, and L.S. Taams,
CD4+CD25+Foxp3+ regulatory T cells induce alternative activation of human monocytes/macrophages.
Proc Natl Acad Sci U S A. **104**(49): p. 19446-51 (2007).

71. Marder, K., Y. Wang, R.N. Alcalay, H. Mejia-Santana, M.X. Tang, A. Lee, D. Raymond, A. Mirelman, R.
Saunders-Pullman, L. Clark, L. Ozelius, A. Orr-Urtreger, N. Giladi, S. Bressman, and L.A.J. Consortium,
Age-specific penetrance of LRRK2 G2019S in the Michael J. Fox Ashkenazi Jewish LRRK2 Consortium.
Neurology. **85**(1): p. 89-95 (2015).

72. Cilia, R., C. Siri, D. Rusconi, R. Allegra, A. Ghiglietti, G. Sacilotto, M. Zini, A.L. Zecchinelli, R. Asselta, S.
Duga, A.M. Paganoni, G. Pezzoli, M. Seia, and S. Goldwurm, LRRK2 mutations in Parkinson's disease:
confirmation of a gender effect in the Italian population. *Parkinsonism Relat Disord*. **20**(8): p. 911-4 (2014).

73. Trinh, J., I. Guella, and M.J. Farrer, Disease penetrance of late-onset parkinsonism: a meta-analysis. *JAMA
Neurol*. **71**(12): p. 1535-9 (2014).

74. Giannoccaro, M.P., C. La Morgia, G. Rizzo, and V. Carelli, Mitochondrial DNA and primary mitochondrial
dysfunction in Parkinson's disease. *Mov Disord*. **32**(3): p. 346-363 (2017).

75. Larsen, S.B., Z. Hanss, and R. Kruger, The genetic architecture of mitochondrial dysfunction in Parkinson's
disease. *Cell Tissue Res*. **373**(1): p. 21-37 (2018).

76. Williamson, M.G., M. Madureira, W. McGuinness, R. Heon-Roberts, E.D. Mock, K. Naidoo, K.M.L. Cramb,
M.C. Caiazza, A.B. Malpartida, M. Lavelle, K. Savory, S.W. Humble, R. Patterson, J.B. Davis, N. Connor-
Robson, B.J. Ryan, and R. Wade-Martins, Mitochondrial dysfunction and mitophagy defects in LRRK2-
R1441C Parkinson's disease models. *Hum Mol Genet*. **32**(18): p. 2808-2821 (2023).

77. Mortiboys, H., K.K. Johansen, J.O. Aasly, and O. Bandmann, Mitochondrial impairment in patients with
Parkinson disease with the G2019S mutation in LRRK2. *Neurology*. **75**(22): p. 2017-20 (2010).

78. Gonzalez-Hunt, C.P., E.A. Thacker, C.M. Toste, S. Boularand, S. Deprets, L. Dubois, and L.H. Sanders,
Mitochondrial DNA damage as a potential biomarker of LRRK2 kinase activity in LRRK2 Parkinson's
disease. *Scientific Reports*. **10**(1) (2020).

79. Sanders, L.H., J. Laganière, O. Cooper, S.K. Mak, B.J. Vu, Y.A. Huang, D.E. Paschon, M. Vangipuram, R.
Sundararajan, F.D. Urnov, J.W. Langston, P.D. Gregory, H.S. Zhang, J.T. Greenamyre, O. Isacson, and B.
Schüle, LRRK2 mutations cause mitochondrial DNA damage in iPSC-derived neural cells from Parkinson's
disease patients: Reversal by gene correction. *Neurobiology of Disease*. **62**: p. 381-386 (2014).

80. Scharping, N.E., D.B. Rivadeneira, A.V. Menk, P.D.A. Vignali, B.R. Ford, N.L. Rittenhouse, R. Peralta, Y.
Wang, Y. Wang, K. DePeaux, A.C. Poholek, and G.M. Delgoffe, Mitochondrial stress induced by continuous
stimulation under hypoxia rapidly drives T cell exhaustion. *Nat Immunol*. **22**(2): p. 205-215 (2021).

81. Wallings, R.L., M.K. Herrick, and M.G. Tansey, LRRK2 at the Interface Between Peripheral and Central
Immune Function in Parkinson's. *Frontiers in Neuroscience*. **14** (2020).

82. Panagiotakopoulou, V., D. Ivanyuk, S. De Cicco, W. Haq, A. Arsic, C. Yu, D. Messelodi, M. Oldrati, D.C.
Schondorf, M.J. Perez, R.P. Cassatella, M. Jakobi, N. Schneiderhan-Marra, T. Gasser, I. Nikic-Spiegel, and
M. Deleidi, Interferon-gamma signaling synergizes with LRRK2 in neurons and microglia derived from
human induced pluripotent stem cells. *Nat Commun*. **11**(1): p. 5163 (2020).

83. Yadavalli, N. and S.M. Ferguson, LRRK2 suppresses lysosome degradative activity in macrophages and
microglia through MiT-TFE transcription factor inhibition. *Proceedings of the National Academy of
Sciences*. **120**(31) (2023).

84. Tong, Y., A. Pisani, G. Martella, M. Karouani, H. Yamaguchi, E.N. Pothos, and J. Shen, R1441C mutation
in LRRK2 impairs dopaminergic neurotransmission in mice. *Proc Natl Acad Sci U S A*. **106**(34): p. 14622-7
(2009).

Acknowledgments: We thank members of the Tansey lab for useful discussions and edits of the manuscript. We thank the UF Interdisciplinary Center for Biotechnology Research (UF | ICBR) for use of flow cytometry and sequencing facilities and advice.

Funding: Partial funding for this work was derived from awards from the Bright Focus Fellowship A2021017F (RLW), NIH/NINDS grant RF1NS128800 (MGT), NIH NIA 1RF1AG057247, and the joint efforts of The Michael J. Fox Foundation for Parkinson's Research (MJFF) and the Aligning Science Across Parkinson's (ASAP) initiative. MJFF administers the grant ASAP-020527 on behalf of ASAP and itself. For the purpose of open access, the author has applied a CC-BY public copyright license to the Author Accepted Manuscript (AAM) version arising from this submission.

Author contributions:

Conceptualization: RLW, MGT

Methodology: RLW, HAS, KMF, SS, NB, SZ, TU, CK, ES

Investigation: RLW, KMF

Visualization: RLW, MGT

Funding acquisition: RLW, MGT

Project administration: MGT

Supervision: MGT

Writing – original draft: RLW, HAS, KMF, SS, NB, SZ, TU, CK, ES, MGT

Competing interests: All authors hold no competing interests

Data and materials availability: All data are available in the main text or supplementary files. Correspondence and requests for materials should be addressed to MGT.

Figure 1. *R1441C* mutation leads to age dependent biphasic alteration in antigen presentation. pMacs

from 2-3-month and 18-21-month-old female or male, B6 or *R1441C* mice were stimulated with 100U IFNγ

for 18-hours +/- 100nM PF360. (A, B) Lrrk2 protein levels were assessed and normalized to β-Actin levels

and quantified. Representative western blots shown. (C, D, E, F) Surface MHC-II median fluorescence

intensity (MFI) was quantified on LPMs and via flow cytometry. (G-J) pMacs were stained for

intracellular and extracellular MHC-II MFI and Ex:IcMHC-II ratio quantified. Scale bars, 100μM. Bars

represent mean +/- SEM (N = 6-12). One/two-way ANOVA, Bonferroni *post hoc*, groups sharing the same

letters are not significantly different (p > 0.05) whilst groups displaying different letters are significantly

different (p < 0.05).

Figure 2. *R1441C* mutation leads to age dependent biphasic alteration in cytokine release ex vivo and circulating cytokines in vivo. (A) pMacs from 2-3-month and 18-21-month-old female or male, B6 or *R1441C* mice were stimulated with 100U IFNγ for 18-hours and media collected. (B, C, D, E) Cytokine levels of IL10, IL6 and TNF were quantified and normalized to live cell count. Bars represent mean +/- SEM (N = 8-10). Two-way ANOVA, Bonferroni *post hoc*. (F) Trunk blood was collected via trunk bleed and plasma was isolated. (G, H) Cytokine levels of IL10, IL6 and TNF were quantified. Bars represent mean +/- SEM (N = 8-10). Student's T-test. Groups sharing the same letters are not significantly different (p > 0.05) whilst groups displaying different letters are significantly different (p < 0.05).

Figure 3. *R1441C* **mutation leads to age dependent biphasic alteration in lysosomal activity.** pMacs from 2-3-month and 18-21-month-old female or male, B6 or *R1441C* mice were stimulated with 100U IFNγ for 18-hours. (A) Schematic of YAe assay. (B-E) Cells were co-treated with 100nM PF360 or 50nM Bafilomycin A1 (baf) and YAe MFI was quantified on LPMs via flow-cytometry. (F-H) DQ-BSA and BMV109 MFI was quantified in LPMs via flow cytometry. (J-O) DQ-BSA and BMV109 MFI was quantified in pMacs via microscopy. Scale bars, 100μM. Bars represent mean +/- SEM (N = 6-12). Two - way ANOVA, Bonferroni *post hoc*, or Student's T-test. Groups sharing the same letters are not significantly different (p > 0.05) whilst groups displaying different letters are significantly different (p < 0.05).

Figure 4. *R1441C* **Lrrk2 mutation alters pathogen uptake and pathogen-mediated DNA damage in age- and sex-dependent manner**. pMacs from 2-3-month and 18-21-month-old female, B6 or *R1441C* mice were incubated with 100μg of pHrodo *E. coli*. +/- 100nM PF360 (A-F) pMacs were imaged every 20 minutes for 5 hours, and GFP+ cells (% of total) and GFP MFI was measured. Scale bars, 100μM. Three-way ANOVA, Bonferroni *post hoc*. Main effects of groups are reported in each graph key. Groups sharing the same letters are not significantly different (p > 0.05) whilst groups displaying different letters are significantly different (p < 0.05). (G, H) TUNEL MFI was quantified in LPMs via flow cytometry. Bars represent mean +/- SEM (N = 6-12). One-way ANOVA, Bonferroni *post hoc*. Groups sharing the same letters are not significantly different (p > 0.05) whilst groups displaying different letters are significantly different (p < 0.05).

Figure 5. *R1441C* **Lrrk2 mutation induces immune exhaustion in a kinase-dependent manner**. pMacs

from 2-3-month and 18-21-month-old female, B6 or *R1441C* mice were incubated with 100ng/mL of LPS

for 5 days +/- 100nM PF360. (A) Schematic of study design. (B, C) Lrrk2 protein levels were assessed and

normalized to β-Actin levels and quantified. Representative western blots shown. (D, E) PD1+ LPM count

was quantified via flow cytometry. (F) After 5-day LPS or vehicle treatment, pMacs were stimulated with

100nG/mL of IFNγ for 18-hours and MHC-II MFI expression was quantified via flow cytometry. Bars

represent mean +/- SEM (N = 6-12). Two -way ANOVA, Bonferroni *post hoc*. (G, H, I) After 5-day LPS

or vehicle treatment, pMacs were incubated with 100μg of pHrodo *E. coli*. +/- 100nM PF360. pMacs were

imaged every 20 minutes for 5 hours, and GFP+ cells (% of total) and GFP MFI was measured. Scale bars,

100μM. Three-way ANOVA, Bonferroni *post hoc*. Main effects of groups are reported in each graph key.

Groups sharing the same letters are not significantly different (p > 0.05) whilst groups displaying different

letters are significantly different (p < 0.05).

Figure 6. *R1441C* Lrrk2 mutation alters transcriptional regulation of mitochondrial and oxidative stress genes as a function of age in female macrophages. Transcriptomic analysis from B6 or *R1441C* vehicle and IFNγ-treated pMacs, harvested from 2-3-month or18-21-month-old. (A, B, C). Significant Old:Young DEGs in IFNγ-treated pMacs were counted and compared across B6 and *R1441C*. Volcano plots show genes with log₂-transformed fold change >0.5 and an adjusted p-value ≤ 0.05. (D) KEGG pathway and GO BP enrichment analysis identifies pathways in Old:Young DEGs seen only in *R1441C* samples. (F). Heat maps show KEGG: Parkinson's Disease pathway with significant Old:Young DEGs seen only in *R1441C* samples.

Figure 7. *R1441C* **and** *Y1699C-LRRK2* **monocyte-derived patient macrophages exhibit hypophagocytosis and increased markers for exhaustion.** (A) Schematic of study design. (B, C) Cryopreserved PBMCs were cryorecovered and plated in the presence of hM-CSF to differentiate myeloid cells to macrophages. Macrophages were incubated with 100μg of pHrodo *E. coli* and imaged every 20 minutes for 5 hours, and GFP+ cells (% of total) quantified. . Scale bars, 100μM. Three-way ANOVA, Bonferroni *post hoc*. Main effects of groups are reported in each graph key. Groups sharing the same letters are not significantly different (p > 0.05) whilst groups displaying different letters are significantly different (p < 0.05). (D, E, F) PD1+ cell count and PD1 expression was quantified in total PBMCs via flow cytometry. Bars represent mean +/- SEM (N = 6-12). One-way ANOVA, Bonferroni *post hoc*. Groups sharing the same letters are not significantly different (p > 0.05) whilst groups displaying different letters are significantly different (p < 0.05). (G) Monocyte-derived macrophages were stained for PD1 and microscopy images captured. Scale bars, 100μM.

1037   **References** (*85-88*)

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059