Title: The *R1441C-LRRK2* mutation induces myeloid immune cell exhaustion

in an age- and sex-dependent manner

Authors: Rebecca L., Wallings[1,2,6], Karen McFarland[2,3], Hannah A. Staley[1,2], Noelle

Neighbarger[1,2], Susen Schaake[4], Norbert Brüggemann[4,5], Simone Zittel[5], Tatiana Usnich[4,5],

Christine Klein[4,6], Esther M Sammler[6,7,8] and Malú Gámez Tansey[1, 2, 3,6]*

Affiliations:

[1] Department of Neuroscience, [2] Center for Translational Research in Neurodegenerative Disease,

University of Florida, College of Medicine, McKnight Brain Institute, Gainesville, Florida, USA

[3] Department of Neurology and Fixel Institute for Neurological Diseases, University of Florida

Health, Gainesville, Florida, USA

[4] Institute of Neurogenetics, University of Lübeck, Lübeck, Germany.

[5] Department of Neurology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

[6] Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase,

MD, USA

[7] MRC Protein Phosphorylation and Ubiquitylation Unit, School of Life Sciences, University of

Dundee, Dundee DD1 5EH, U.K. d.r.alessi@dundee.ac.uk e.m.sammler@dundee.ac.uk.

[8] Department of Neurology, School of Medicine, Ninewells Hospital, Ninewells Drive, Dundee

DD1 9SY, U.K.

*Corresponding author. Email: mgtansey@ufl.edu

Abstract: Considering age is the greatest risk factor for many neurodegenerative diseases, aging, in particular aging of the immune system, is the most underappreciated and understudied contributing factor in the neurodegeneration field. Genetic variation around the *LRRK2* gene affects risk of both familial and sporadic Parkinson's disease (PD). The leucine-rich repeat kinase 2 (LRRK2) protein has been implicated in peripheral immune signaling, however, the effects of an aging immune system on LRRK2 function have been neglected to be considered. We demonstrate here that the *R1441C* mutation induces a hyper-responsive phenotype in macrophages from young female mice, characterized by increased effector functions, including stimulation-dependent antigen presentation, cytokine release, phagocytosis, and lysosomal function. This is followed by age-acquired immune cell exhaustion in a Lrrk2-kinase-dependent manner. Immune-exhausted macrophages exhibit suppressed antigen presentation and hypophagocytosis, which is also demonstrated in myeloid cells from *R1441C* and *Y1699C*-PD patients. Our novel findings that *LRRK2* mutations confer immunological advantage at a young age but may predispose the carrier to age-acquired immune cell exhaustion have significant implications for LRRK2 biology and therapeutic development. Indeed, LRRK2 has become an appealing target in PD, but our findings suggest that more research is required to understand the cell-type specific consequences and optimal timing of LRRK2-targeting therapeutics.

One Sentence Summary: The *R1441C-LRRK2* mutation causes an age-acquired immune cell exhaustion in macrophages in a sex-dependent manner

Main Text:

INTRODUCTION

Parkinson's disease (PD) is a common, progressive neurodegenerative disease, affecting around 1-2% of the population over the age of 65 (*1*). PD prevalence will likely increase two-fold by the year 2030 (*2*). In addition, the projected total economic burden will surpass $79 billion by 2037 (*3*), underscoring the need to reduce PD incidence and delay disease progression. The fact that there are currently no disease-modifying drugs for people with PD indicates that knowledge gaps still need to be closed to identify ways to cure or prevent this disease. Although the etiology of PD is largely unknown, it is thought to involve a complex interaction between various genetic and environmental factors (*4*). Although typically thought of as a disease limited to the central nervous system (CNS), evidence suggests a crucial and fundamental role of inflammation in both the CNS and the periphery in PD pathogenesis (*5*). Numerous studies of peripheral blood and cerebrospinal fluid from patients with PD suggest alterations in markers of inflammation and immune cell populations that could initiate or exacerbate neuroinflammation and perpetuate the neurodegenerative process (*6-9*). Furthermore, a number of disease genes and risk factors have been identified as modulators of immune function in PD (*10-12*).

Pathogenic variants in the gene encoding leucine-rich repeat kinase 2 (LRRK2) account for the majority of familial PD patients (*13*). LRRK2 is a large, complex protein containing six functional domains, two of which have known enzymatic activities; a ROC domain (GTPase activity) and a MAPKK domain (kinase activity). There are numerous single-nucleotide substitutions in each of these domains that have been shown to be pathogenic (*14-21*). The second most prevalent mutation, *R1441C*, located in the GTPase ROC domain, is present in about 46% of familial PD that has an origin in the Basque region in Spain (*22*) and leads to decreased LRRK2 GTPase hydrolysis and increased LRRK2 protein kinase activity (*23, 24*).

LRRK2 is expressed in innate and adaptive immune cells and its expression is tightly regulated by immune stimulation (*9, 25-28*). Furthermore, LRRK2 is a member of the receptor interacting protein (RIP) kinase family, which is a group of proteins that detect and respond to cellular stress by regulating cell death and

activation of the immune system (*29*). LRRK2 function has also been implicated in pathogen control based

on observations that LRRK2 variants confer differential susceptibility to a number of bacterial infections,

including *Mycobacterium tuberculosis* (*30, 31*), *Salmonella Typhimurium* (*32*), and *Listeria monocytogenes*

(33), as well as an increased risk of developing leprosy (*34*) and inflammatory bowel disease such as

Crohn's (*35*) and ulcerative colitis (*36*). Collectively such data strongly suggest a generalized role of

LRRK2 in immune system regulation.

Although research has focused primarily on neuroinflammation in PD over the last two decades, it is

becoming increasingly clear that peripheral inflammatory responses may play important roles in PD

pathogenesis (*37, 38*). For example, immune signaling plays a role in the regulation of neurodegeneration

in a *LRRK*2-PD pre-clinical model, with the *R1441C* mutation altering the type II interferon response in

peripheral immune cells (*28*). Furthermore, neutralization of IL-6 over-production in the periphery of these

mice prevents substantia nigra *pars compacta* (SNpc) dopaminergic (DA) neuron loss in the presence of

peripheral lipopolysaccharide (LPS) exposure (*39*). Importantly, immune dysfunction in the periphery takes

primacy over the CNS, as the presence of *Lrrk2* mutations in lymphocytes alone is sufficient to induce DA

neuron loss in the midbrain of LPS-treated mice (*39*). Such data suggest a pertinent role of the peripheral

immune system in *LRRK2*-PD, with immune dysfunction driving the development of the disease.

As well as a role in immune response, a vast amount of research has focused on the role of LRRK2 at the

lysosome. In immune cells, specifically, LRRK2 has been shown to mediate a number of lysosome-

dependent immune functions such as phagocytosis (*40*) and antigen presentation (*9, 27, 41*). Intriguingly,

LRRK2 has been shown to alter lysosomal function and autophagy in kidneys in an age-dependent, biphasic

manner, with enhanced autophagy levels in young *Lrrk2*-null mice, that are reduced and drop below control

levels once the mice reach 20 months of age (*42*). Despite this intriguing observation, few studies have

investigated the effects of *LRRK2* mutations on lysosomal processes in the context of aging. Considering

age is the greatest risk factor for many neurodegenerative diseases, aging, in particular aging of the immune

system (*5*), is the most underappreciated and understudied contributing factor to neurodegeneration in the

field.

To close this important gap, we assessed the effects of *LRRK2* mutations on immune responses and

lysosomal function in peripheral immune cells in young versus old mice. We used *R1441C-Lrrk2* mutant

knock-in (KI) mice based on previous findings demonstrating an effect of the *R1441C/G* mutation on

peripheral immune responses (*28, 39*) whilst evading limitations of an overexpression model and

investigating the effects of the *R1441C* mutation in physiologically relevant levels. Peritoneal macrophages

(pMacs) were isolated from these mice to examine the effects of the *R1441C* mutation on macrophage cell

function at different ages. pMacs are not a homogenous population of cells, but rather a mix of small and

large pMacs (SPMs and LPMs, respectively). LPMs are resident to the peritoneal cavity and are

traditionally thought of as anti-inflammatory, phagocytic and responsible for the presentation of exogenous

antigens (*43*). SPMs, on the other hand, are generated from bone-marrow-derived myeloid precursors which

migrate to the peritoneal cavity in response to infection, inflammatory stimuli, or thioglycolate, and present

a pro-inflammatory functional profile (*43*). We observed a bi-phasic, age-dependent effect of the *R1441C-*

Lrrk2 KI mice, with increased antigen presentation, production of anti-inflammatory cytokines, lysosomal

activity, and pathogen uptake in peripheral macrophages from young (2-3 months) *R1441C-Lrrk2* KI mice

examined *ex vivo*. Conversely, peripheral macrophages from aged (18-21 months) *R1441C-Lrrk2* KI mice

exhibited a diminished capacity to present antigens in the presence of inflammatory insults, decreased

lysosomal function and pathogen uptake, with concomitant increases in DNA fragmentation in the presence

of pathogens. Importantly, we demonstrate that this is a sex-dependent phenotype, with male *R1441C-Lrrk2*

KI failing to present this age-dependent phenotype. Furthermore, we demonstrate that the age-dependent

hypophagocytosis and decreased antigen presentation in the *R1441C-LRRK2* KI females is due to an

acquired immune cell exhaustion driven by increased Lrrk2 protein kinase activity and immune responses

whilst young. These phenotypes were also observed in human peripheral myeloid cells, with monocyte-

derived macrophages from *R1441C* and *Y1699C* patients exhibiting decreased pathogen uptake and

increased programmed cell death protein ligand 1 (PDL1) expression, consistent with immune cell

exhaustion.

RESULTS

LRRK2 R1441C **mutant immune cells display age-dependent biphasic alteration in antigen**

presentation

LRRK2 protein expression has been shown to increase in immune cells from both murine pre-clinical

models and patient cells in response to inflammatory stimuli (*9, 27, 28, 44-46*). To determine whether this

is age-dependent, Lrrk2 protein expression were assessed in pMacs from young and aged mice treated with

100U of IFNγ for 18h. In pMacs from young females, IFNγ treatment significantly increased total Lrrk2

protein in both B6 and *R1441C* pMacs (Fig. 1A, B; $p < 0.01$). Relative to pMacs from young female mice,

pMacs from aged females expressed increased total Lrrk2 protein under vehicle stimulation, with aged

R1441C pMacs expressing significantly less Lrrk2 protein relative to B6 aged pMacs ($p < 0.05$).

Furthermore, pMacs from aged females did not exhibit a significant increase in Lrrk2 protein expression

with IFNγ treatment, with aged *R1441C* pMacs still expressing significantly less Lrrk2 protein relative to

B6 aged pMacs in IFNγ-treatment ($p<0.05$). In pMacs from young male mice, the same observations were

made as in the females (Fig. S1A, B). In aged male mice, although a lack of IFNγ-mediated Lrrk2 increase

was observed as seen in females, the effect of the *R1441C* mutation was not observed. These observations

indicate that Lrrk2 protein increase in macrophages as a function of age in both males and females, however,

the *R1441C* mutation limits this increase in a sex-dependent manner.

To decipher whether these alterations in Lrrk2 protein expression was accompanied by increases in Lrrk2

kinase activity, phosphorylation of Rab10 at T73 (pT73), a *bona fide* kinase-substrate of Lrrk2 (*23*), was

quantified. In pMacs from young female mice, a significant ($p<0.01$) increase in pT73 Rab10 expression

was observed in response to IFNγ in *R1441C-Lrrk2* cells, whilst this was not observed in pMacs from young

B6 females (Fig. 1C, D). Interestingly, this phenotype was reversed in pMacs from aged female mice, with

a significant increase (p<0.01) in pT73 Rab10 expression observed in response to IFNγ in B6 but not

R1441C-Lrrk2 cells. Of note, the pT73 Rab10 expression in pMacs from aged B6 females in response to

IFNγ was significantly (p<0.05) elevated relative to all other groups, suggesting that, in B6 females, Lrrk2

kinase activity increases with age. In pMacs from young male mice, however, no significant effect of the

R1441C mutation was observed in vehicle or IFNγ conditions (Fig. SB, C), although a significant (p<0.01)

increase in pT73 Rab10 expression was observed in response to IFNγ. Interestingly, a significant increase

in pT73 Rab10 expression was observed in pMacs from aged male mice relative to those from young males,

suggesting that Lrrk2 kinase activity increases with age in pMacs from male mice. Furthermore, upon IFNγ

treatment, a significant increase in pT73 Rab10 expression was observed in pMacs from aged *R1441C-*

Lrrk2 males relative to all other groups, suggesting an additive effect of the *R1441C* mutation on Lrrk2

kinase activity in an ageing male immune system.

As pMacs are not a homogenous population of cells, as previously discussed, we used multi-color flow

cytometry-based methods to immunophenotype LPMs and SPMs, as the two populations can be

distinguished based on CD11b expression (Fig. S1D) (*43*). No differences were seen in LPM or SPM count

between genotypes in males or females at either young or aged (Fig. S1E-H).

To begin to investigate the effects of the *R1441C* mutation on pMac function at different ages, surface

MHC-II expression was assessed on both LPMs and SPMs via multi-color flow cytometry as a measure of

antigen presentation in both vehicle and IFNγ treated pMacs. In pMacs from young females, the *R1441C*

mutation increases surface MHC-II expression on LPMs in IFNγ treatments (Fig 1D). Interestingly,

although no effect of age was observed in vehicle conditions, surface MHC-II expression was significantly

decreased in aged pMacs from both B6 and *R1441C* when treated with IFNγ relative to LPM from young

mice (p<0.001). Furthermore, a significant decrease in surface MHC-II expression was observed in LPM

from aged *R1441C* females relative to aged B6 LPM (p<0.01), indicative of decreased ability to engage in

antigen presentation in these cells. This effect of the *R1441C* mutation was not observed in LPMs from

male mice (Fig. 1E). However, as seen in LPM from female mice, an effect of age was also observed in

LPM from male mice, with surface MHC-II expression significantly decreased in LPM from aged mice, in

both vehicle and IFNγ conditions (p<0.001). Regarding surface MHC-II expression on SPMs, the same

effect of age was observed in both male and female pMacs, with SPM from aged mice expressing

significantly less surface MHC-II relative to SPM from young mice (p<0.001) (Fig. S1I, J).

With alterations in surface MHC-II expression, there is likely increased and decreased transport of MHC-

II-complexes to the cell surface in pMacs from young and aged *R1441C* females, respectively. To test this,

using pMacs treated with vehicle or IFNγ, we co-stained for extracellular vs intracellular MHC-II

(ExMHCII vs IcMHCII) and monitored expression via fluorescence microscopy. pMacs were co-treated

with 100nM PF360 to determine the effects of Lrrk2 kinase activity on surface MHC-II expression.

Treatment with the Lrrk2 kinase inhibitor, PF360 (PF-06685360)PF360 reduced expression of Lrrk2

phosphorylated at S935 (pS935) Lrrk2 in all groups (Fig. S2A, B). Although no significant effects of

genotype or age were observed in vehicle treated pMacs from female mice, upon IFNγ stimulation, a

significant increase in Ex:IcMHCII ratio was observed in pMacs from young *R1441C* mice relative to other

groups (p<0.05) (Fig. 1F, Fig. S2C). Furthermore, a significant decrease in Ex:IcMHCII ratio was observed

in pMacs from aged *R1441C* females relative to aged B6 pMacs and young pMacs (p<0.01) (Fig. 1G). The

increased Ex:IcMHCII ratio in pMacs from young *R1441C* females was ameliorated upon Lrrk2 kinase

inhibition (Fig. S2C, D), with this treatment also decreasing Ex:IcMHCII ratio in pMacs from young and

aged B6 females. However, no effect of Lrrk2 kinase inhibition was seen in pMacs from aged *R1441C*

females. Such data suggests that Lrrk2-kinase inhibition reduces the transport of IcMHCII to the cell surface

to engage in antigen presentation, but this may occur in an age- and genotype-dependent manner. No effect

of the *R1441C* mutation was observed in males (Fig. 1H, I, Fig. S2D). It seems therefore that the *R1441C*

mutation exhibits a bi-phasic, age- and sex-dependent effect on antigen presentation in macrophages. The

effect of Lrrk2 protein kinase inhibition on antigen presentation, however, does not appear to be bi-phasic,

with Lrrk2 kinase inhibition decreasing antigen presentation in pMacs from both young and aged mice, at

least in B6 mice.

LRRK2 R1441C **mutant immune cells display age-dependent biphasic alteration in cytokine release**

ex vivo **and circulating cytokines** *in vivo*

To investigate the extent to which alterations in antigen presentation were accompanied by changes in

cytokine release, media from vehicle- and IFNγ treated cells were collected and cytokine release was

quantified on a multiplexed immunoassay panel (Fig. 2A). In pMacs from young *R1441C* females, increases

in IL6 relative to B6 controls was observed in vehicle treated conditions, which was ameliorated upon

ageing (Fig. 2B). Interestingly, a bi-phasic, age-dependent pattern was observed regarding IL10 release,

with increased IL10 observed in pMacs from young *R1441C* females relative to young B6 controls, whilst,

conversely, decreased IL10 was observed in pMacs from aged *R1441C* females relative to aged B6 controls

(Fig. 2B). As well, TNF was significantly elevated in IFNγ-treated pMacs from R1441C aged females

relative to aged B6 controls and both young B6 and young *R1441C* females (p<0.05) (Fig. 2B). With a

decrease in the release of the anti-inflammatory cytokine, IL10, and an increase in the release of the pro-

inflammatory cytokine, TNF, in females, these findings indicate that pMacs from aged *R1441C* female

mice promote a pro-inflammatory microenvironment.

In pMacs from *R1441C* males, IL6 release was significantly increased in IFNγ conditions relative to B6

controls in both young and aged mice (p<0.05) (Fig. 2C). Similarly, IL10 release was significantly increased

in pMacs from both aged and young *R1441C* males relative to B6 controls (p<0.05) (Fig.2C). Regarding

TNF release from pMacs from male mice, no genotype effects were observed in pMacs from young mice,

whilst an increase in TNF was observed in aged R1441C males relative to aged B6 controls, although this

did not significantly differ from pMacs from neither young B6 or *R1441C* males (Fig.2C). It seems

therefore that, in pMacs from aged male mice, the *R1441C*-associated increase in IL6 and IL10 seen in

young mice is retained with age but *R1441C* macrophages secrete more TNF than B6 controls after

stimulation.

To determine if these alterations in cytokine release associated with the *R1441C* mutation *ex vivo* was also

detectable *in vivo*, plasma from whole blood was collected and cytokine release was quantified (Fig. 2D).

A significant increase in IL10, IL6, and TNF was observed in plasma from young *R1441C* female mice

relative to young B6 controls (p<0.05) (Fig. 2E). Interestingly, an increase in these cytokines were observed

with age, although IL6 and IL10 release was significantly reduced in aged *R1441C* females relative to B6

controls (p<0.01), confirming the biphasic age-dependent effects observed *ex vivo*. In young male mice, a

significant increase in both IL6, IL10 and TNF were also observed in *R1441C* males relative to B6 controls

(p<0.01) (Fig. 2F). However, unlike in females, an age-dependent increase in cytokine release across

genotypes was not observed, with the only genotype effect being observed in aged male mice seen with

IL10, with a decrease in IL10 observed in aged *R1441C* males relative to aged B6 controls.

235   The R1441C Lrrk2 mutation is associated with alterations of monocyte populations and microglia

236   activation in aged female mice

The observation of a biphasic immune response, with an initial hyperinflammation state proceeded by a

subsequent downregulation of immune responses, prompted us to hypothesize that what was being observed

in aged *R1441C* female immune cells was immune cell exhaustion. Immune cell exhaustion arises from

chronic stimulation of immune cells, leading to a downregulation, or even absence, of an expected effector

response (e.g. antigen presentation, cytokine release or phagocytosis) (*47*). It is known that monocytes

pushed to exhaustion shift to express higher Ly6C levels and a reduction in Ly6C$^-$ monocytes under resting

conditions (*48*). We therefore assessed monocyte sub-populations in the blood of young and aged mice to

determine if a shift to an exhausted state could be observed. Indeed, we observed a detectable increase in

Ly6C$^+$:Ly6C$^-$ in the peripheral blood mononuclear cells isolated from blood of aged *R1441C* female mice

relative to aged B6 controls and young B6 and *R1441C* female mice (Fig. S3A) with no significant change

observed in male mice (Fig. S3B). The fact that no effect of genotype was observed in young females and

only aged females, suggests the development of an age-acquired immune cell exhaustion phenotype in the

R1441C females. This is further supported by the observation that in aged *R1441C* female mice, but not

males, an increase in PDL1 expression, an immune checkpoint molecule, was upregulated on LyC+

monocytes from blood relative to B6 controls (Fig. S3C, D).

It has been suggested that circulating monocytes, disrupted in PD, may infiltrate the brain parenchyma and

affect local glial and adaptive immune responses (*49*). To determine whether exhausted monocytes could

be infiltrating the brain parenchyma of aged *R1441C* female mice, immune cells were isolated from whole

brain and assessed via multi-color flow cytometry (Fig. S3E). Indeed, in aged *R1441C* females, an increase

in overall Ly6C$^+$ monocytes in the brain parenchyma was observed relative to B6 controls, whilst MHC-

II/Ly6C$^+$ were decreased (Fig. S3F, G), suggesting that whilst there is an increase in infiltrating Ly6C$^+$

monocytes in the brains of aged *R1441C* females, they exhibit the same decrease in antigen presenting

capabilities to those observed in pMacs *ex vivo*. Importantly, alongside this increase in infiltrating

monocytes into the brain parenchyma, a concomitant increase in MHC-II expression on microglia was

observed in aged *R1441C* females relative to B6 controls (Fig. S3H) with no changes in overall microglia

number (Fig. S3I). No changes were observed in the brains of male mice (Fig. S3J-M).

LRRK2 R1441C mutant immune cells display age-dependent biphasic alterations in lysosomal activity

As previously discussed, loss of *Lrrk2* leads to biphasic alterations in lysosomal function in kidneys (*42*).

Interestingly, lysosomal dysfunction has been implicated in immune cell exhaustion, with decreased

autophagy levels and lysosomal function being associated with increased T cell exhaustion (*50*). We

therefore hypothesized that the biphasic immune response observed in *R1441C* females would be associated

with concomitant biphasic lysosomal function.

The Eα: YAe model is used to monitor the antigen presentation capabilities of cells by incubating an

endogenous peptide (Eα 52-68) which is subsequently phagocytosed, transported to the lysosome, loaded

onto an MHC-II complex at the lysosome and transported back to the plasma membrane for antigen

presentation (Fig. 3A). This Eα peptide-loaded MHC-II can subsequently be detected via flow cytometry

using the YAe antibody (*51*); DOI: 10.1016/j.jim.2015.04.023). We observed that YAe median

fluorescence intensity (MFI) was significantly increased in IFNγ-treated LPMs from young *R1441C*

females relative to age-matched B6 controls and pMacs from aged females ($p<0.05$) (Fig. 3B). Furthermore,

IFNγ-treated LPMs from aged *R1441C* females exhibited decreased YAe MFI relative to other groups. Co-

treatment with PF360 ameliorated the increased YAe MFI observed in LPMs from young *R1441C* females.

Antigen presentation and pathogen sensing require protease action and sufficient lysosomal function in

order to occur (*52*), which is why lysosomotropic agents have been shown to decrease peptide-loaded-

MHC-II surface expression in antigen presenting cells (*51*). Indeed, when co-treated with the vacuolar H+

ATPase (V-ATPase) inhibitor, Bafilomycin A1, YAe MFI significantly decreased in both *R1441C* and B6

LPMs from both young and aged mice ($p<0.01$). Although no effect of genotype was observed in male mice

at either ages, the same treatment effects were observed as seen in females, with PF360 and Bafilomycin

A1 co-treatment significant decreasing YAe MFI relative to IFNγ conditions ($p<0.01$) (Fig. 3C). Due to

this observation and the crucial role of the lysosome in antigen presentation, we next sought to probe the

effects of the *R1441C* mutation on lysosomal function in these pMacs.

pMacs were plated and incubated with the lysosomal degradation probe, DQ Red BSA, and the pan-

cathepsin probe, BMV109, and collected for flow cytometry to quantify lysosomal activity. It was observed

that LPMs from young *R1441C* females exhibited increased DQ-BSA and BMV109 MFI relative to age-

matched controls and pMacs from aged B6 and *R1441C* females (Fig. 3D, E), indicative of increased

lysosomal protein degradation and pan-cathepsin activity, respectively. Conversely, LPMs from aged

R1441C females exhibited decreased DQ-BSA and BMV109 MFI relative to age-matched controls and

pMacs from young B6 and *R1441C* females. Interestingly, LPM from both aged B6 and *R1441C* females

exhibited decreased DQ-BSA MFI relative to young mice, reflecting an overall decrease in lysosomal

degradative capacity with age. No effects of genotype were observed in LPMs from male mice (Fig. S4A,

B). These observations made via flow cytometry were confirmed with fluorescence microscopy (Fig. 3F-I;

Fig. S4C-F). As was observed with YAe measurements, Lrrk2 kinase inhibition via treatment with PF360

significantly reduced BMV109 MFI observed via microscopy (p<0.05) in all groups except aged B6 and

R1441C females, which exhibited decreased measurement of pan cathepsin activity which were unaffected

by Lrrk2 kinase inhibition. However, Lrrk2 kinase inhibition was able to decrease DQ-BSA MFI in all

groups.

_R1441C_ Lrrk2 mutation is association with age-acquired hypo-phagocytosis

Lysosomal function in macrophages is crucial for a number of other immune-related functions in myeloid

cells, including phagocytosis. Therefore, we posited that the observed biphasic effects of the *R1441C*

mutation on lysosomal function would have downstream consequences on phagocytosis in pMacs. To

explore this hypothesis, pMacs were plated and incubated with pHrodo™ Green *E. coli* BioParticles™,

after which uptake and fluorescence intensity were monitored over 5 hours. An increase in GFP MFI was

observed in pMac cultures from young *R1441C* females relative to age-matched B6 controls and pMac

cultures from aged B6 and *R1441C* females (Fig. 4A, B). As well, a significant reduction in GFP MFI was

observed in pMacs from aged *R1441C* females relative to age-matched B6 controls and young B6 and

R1441C females (p<0.01). Lrrk2 kinase inhibition with PF360 significantly reduced GFP MFI in young

*R1441C fe*males (p<0.01), with no effects of Lrrk2 kinase inhibition on GFP MFI in other groups. In male

mice, pMacs from young *R1441C* males did exhibit increased GFP MFI relative to age-matched B6 controls

and aged B6 and *R1441C* mice which could be significantly reduced with PF360 co-treatment (p<0.05)

(Fig. S5A-B). Interestingly, an overall effect of age was observed in pMac cultures from male mice, with

aged B6 and *R1441C* pMacs exhibiting decreased GFP MFI relative to pMacs cultures from young male

mice. It was still observed, however, that, despite this decrease in phagocytosis with age, pMacs from aged

R1441C males exhibited increased GFP MFI relative to age-matched B6 controls. Collectively, it seems

that pMacs from *R1441C* females exhibit increased phagocytic uptake at a young age and develop an age-

dependent hypophagocytic phenotype. pMacs from *R1441C* males, however, exhibit this same increased

phagocytic uptake at a young age but retain this increase in old ages.

Hypophagocytic activity in *R1441C* macrophages is associated with aberrant pathogen-mediated

DNA damage

Our data thus far suggests a potential age-acquired immune cell exhaustion in macrophages from *R1441C*

female mice. In other disease states associated with cellular exhaustion, apoptotic depletion of immune cells

is also observed (*53*). To investigate the degree of pathogen-induced apoptosis in pMacs from *R1441C*

mice, pMacs were plated in the presence of *E. coli* for 12 hours, harvested and stained using terminal

deoxynucleotidyl transferase dUTP nick end labeling (TUNEL), which detects DNA breaks formed during

the final phase of apoptosis, and TUNEL fluorescence was assessed via flow cytometry. LPMs from young

R1441C females exhibited decreased TUNEL MFI in response to *E. Coli* relative to age-matched B6

controls and LPMs from aged mice, whilst an increase was observed in LPMs from aged *R1441C* females

relative to all other groups (Fig. 4C). Reduced apoptosis in young *R1441C* mice could lead to prolonged

presence of activated immune cells which should otherwise be eliminated, resulting in an increased and

prolonged inflammatory response and, if sustained for prolonged times, could lead to immune cell

exhaustion. Indeed, in response to *E. Coli*, no effect of genotype was observed in aged males (Fig. S5C).

These novel findings support the conclusion that *R1441C* males may evade the age-acquired immune cell

exhaustion that is seen in females by increasing clearance of activated immune cells.

Immune cell exhaustion in *R1441C* Lrrk2 mutant immune cells is dependent on Lrrk2 kinase activity

Our data thus far suggest a potential age-acquired immune cell exhaustion in macrophages from *R1441C*

female mice. We hypothesized that this age-acquired immune cell exhaustion was driven by the initial

hyperinflammatory phase when these mice are young, characterized by increased cytokine release,

increased antigen presentation, and reduced apoptosis of activated immune cells. We therefore

hypothesized that pMacs from young *R1441C* female mice could be pushed to acquire this age-dependent

immune cell exhaustion phenotype if subjected to chronic overstimulation *ex vivo.* To test this, pMacs were

isolated and plated in the presence of a vehicle or 100ng/mL LPS for a 5-day chronic treatment (Fig. 5A).

We first sought to understand how Lrrk2 expression is altered by LPS-induced exhaustion and how age and

the *R1441C* affects this. In pMacs from young females, an increase in Lrrk2 expression was observed in

response to LPS-induced exhaustion (Fig. 5B, C). Importantly, there was no significant effect of treatment

in pMacs from aged female mice, with aged *R1441C* pMacs expressing significantly less Lrrk2 relative to

B6 controls (p<0.01). Interestingly, these same observations were made in pMacs from aged male mice

(Fig. S6A, B) suggesting that, with an additional chronic inflammatory stress, aged *R1441C* males can

exhibit some of the phenotypes observed in females.

A number of negative costimulatory markers that are associated with immune cell exhaustion have been

identified, including the immune checkpoint molecule, PDL1 (*54*). PDL1 expression was quantified via

multi-color flow cytometry in pMacs from young and aged mice to determine if pMacs from aged *R1441C*

females were exhausted as hypothesized. Indeed, an increase in PDL1$^+$ cells was observed in both vehicle

and LPS-exhausted aged *R1441C* LPMs relative to age-matched B6 controls as well as young mice (Fig.

5D). As this was seen in both vehicle and LPS conditions, these exciting findings suggest that pMacs from

R1441C females are already exhibiting immune cell exhaustion markers, which is then exacerbated by

chronic LPS-exhaustion. Interestingly, a significant increase in PDL1$^+$ cell count was observed in pMacs

from young *R1441C* females in both vehicle and LPS-exhausted conditions relative to age-matched B6

controls (p<0.01) (Fig. 5D). Interestingly, when cells were pushed to exhaustion with Lrrk2 kinase inhibitor

co-treatment, a significant reduction in PDL1$^+$ cell count was observed (p<0.05). These data suggest that

R1441C macrophages from female mice may be susceptible to immune cell exhaustion in a Lrrk2-kinase dependent manner.

LPMs from aged *R1441C* male mice, although no significant effect of genotype was observed in vehicle conditions, when pushed to exhaustion with LPS, an increase in PDL1+ LPMs relative to age-matched B6 controls, as well as pMacs from young mice, was observed (Fig. S6C). The lack of differences in PDL1+ cells between genotypes or ages in male mice suggests there is no effect of the *R1441C* mutation on immune cell exhaustion in naïve animals. However, based on the increased PDL1+ cell counts observed in aged *R1441C* males when treated with LPS for 5 days, we conclude that a secondary, chronic inflammatory insult can push *R1441C* pMacs from male mice to acquire immune cell exhaustion like that seen in female mice.

We next sought to determine if LPS-induced exhausted macrophages from young female *R1441C* mice also exhibited the diminished antigen presentation capabilities that we observe in the aged *R1441C* macrophages. After a 5-day treatment with vehicle or LPS +/- PF360, macrophages were treated with IFNγ and MHC-II expression was quantified via multi-color flow cytometry. In LPMs subjected to a 5-day vehicle pre-treatment, the same increased MHC-II expression in *R1441C* macrophages relative to B6 controls was observed (Fig. 5E). As hypothesized, in LPMs pushed to exhaustion with a 5-day LPS pre-treatment, a decreased response to IFNγ was elicited in both genotypes; however, *R1441C* LPMs exhibited decreased MHC-II relative to B6 controls in response to IFNγ, phenocopying what was observed in macrophages with aged mice. Importantly, this could be ameliorated by co-treatment with PF360 during the 5-day pre-treatment, suggesting that the push to immune cell exhaustion and vulnerability of *R1441C* macrophages is dependent on Lrrk2 protein kinase activity. Furthermore, we also observed that pMacs from young *R1441C* females pushed to exhaustion with a 5-day pre-treatment of LPS exhibited a significant reduction in phagocytosis as measured by uptake of pHrodo™ Green *E. coli* relative to B6 controls detected by GFP+ cell percentage and GFP MFI which was ameliorated with PF360 co-treatment ($p<0.01$) (Fig. 5F-H).

Based on the observation that the PDL1$^+$ cell count was elevated in LPMs from aged *R1441C* males relative

to B6 controls, we hypothesized that a 5-day LPS treatment would induce immune cell exhaustion in pMacs

from aged *R1441C* males similar to that seen in female mice. After a 5-day treatment with vehicle or LPS

+/- PF30, macrophages from aged male mice were treated with IFNγ and MHC-II expression quantified via

multi-color flow cytometry. LPMs from aged males pushed to exhaustion with 5-day LPS pre-treatment,

R1441C LPMs exhibited decreased MHC-II relative to B6 controls in response to IFNγ, phenocopying what

was observed in macrophages from aged female *R1441C* mice (Fig. S6E). Additionally, the same

hypophagocytic phenotype observed in aged *R1441C* female pMacs was observed in males once pushed to

exhaustion with a 5-day LPS treatment (Fig. S6F-H). Collectively such data support the hypothesis that,

with a secondary hit in the form of a chronic, inflammatory stimulus, male *R1441C* mice exhibit increased

immune cell exhaustion relative to B6 controls, similar to that seen in naïve *R1441C* female macrophages.

Transcriptomic profiling reveals age- and sex-specific alterations in critical effector function

pathways in *R1441C* macrophages

To explore the age and sexually dimorphic effects of the *R1441C* mutation on macrophage function, RNA

was harvested from pMacs post-vehicle or IFNγ treatment and processed for RNA sequencing (RNAseq).

A principal component analysis (PCA) was conducted from the resulting dataset which revealed that groups

in our dataset primarily cluster based on age and treatment with vehicle or IFNγ treatment (Fig. S7A).

We first sought to identify key cellular pathways altered under different treatment conditions in *R1441C*

pMacs relative to B6 controls. In pMacs from young females treated with IFNγ, Gene Ontology (GO)

enrichment analysis revealed that upregulated differentially expressed genes (DEGs) in *R1441C*

macrophages were enriched in pathways such as regulation of phagocytosis, T cell differentiation, cytokine

production and defense response to virus (Fig. S7B, C). In pMacs from aged females treated with IFNγ,

GO enrichment analysis showed downregulated DEGs in *R1441C* macrophages were enriched in pathways

such as detection of external biotic stimulus, acute inflammatory response, and regulation of MAP kinase

activity (Fig. S7D, E). Collectively, such observations support our previous data that shows an effect of

R1441C Lrrk2 on functional readouts such as increased phagocytosis and cytokine production in response

to IFNγ in young females relative to B6 controls, whilst in aged mice, effector functions responsible for

defense responses are significantly downregulated relative to B6 controls.

Overall, fewer genes were dysregulated in *R1441C* pMacs from young females treated with vehicle, relative

to pMacs treated with IFNγ (Fig. S8A), although these DEGs were enriched in similar pathways (Fig. S8B).

Similar observations were made with pMacs treated with vehicle from aged females (Fig. S8C). As Lrrk2

protein increases in response to IFNγ, it follows that DEGs driven by the *R1441C* mutation would have a

more significant biological impact in these conditions where Lrrk2 protein expression are increased.

Furthermore, given that the majority of phenotypes we observe in pMacs from *R1441C* females are seen in

response to an inflammatory insult or immunological challenge, such as IFNγ or *E. Coli,* it is logical that

more *R1441C*:B6 DEGs would be observed in these conditions relative to vehicle treatments. Few

R1441C:B6 DEGs were observed in pMacs from male mice at both ages and in both treatments (Fig. S8D-

I). This is unsurprising, given that many phenotypes were not observed in *R1441C* male mice. pMacs from

aged *R1441C* only begin to exhibit phenotypes similar to *R1441C* female mice when pushed to immune

cell exhaustion and re-challenged with IFNγ. It is therefore likely that to identify key cellular pathways that

are altered by the *R1441C* mutation in male mice, a second, chronic immune insult is needed to elicit those

changes.

R1441C Lrrk2 mutation alters transcriptional regulation of mitochondrial and oxidative stress genes

in macrophages in a sex- and age-dependent manner

Based on the observation that the *R1441C* mutation alters gene expression in pMacs from aged female

mice, we next sought to determine the key cellular pathways that this mutation altered in the aging process.

When comparing DEGs in pMacs from both B6 and *R1441C* aged mice treated with IFNγ relative to those

from young pMacs, there was a large degree of overlap between the DEGs identified in the two genotypes

(Fig. 6A-C). However, 2012 genes were identified to be differentially expressed upon aging and novel to

R1441C pMacs. KEGG pathway enrichment analysis revealed that these 2012 genes unique to aged

R1441C pMacs were significantly enriched for pathways associated with neurodegenerative diseases, such

as Parkinson's disease, Alzheimer's disease and Huntington's disease ($p<0.05$) (Fig. 6D). Similarly, GO

biological pathway enrichment analysis of these DEGs showed a significant enrichment ($p<0.05$) in

pathways related to mitochondria, oxidative stress and metabolism, including mitochondrial respiratory

chain complex I assembly, glucose and pyruvate metabolic process and cellular respiration, which are all

known key lysosomal and mitochondrial functions. Many DEGs observed in the Parkinson's Disease

KEGG pathway were present in the GO biological pathways associated with mitochondrial function and

oxidative stress, including many *Nduf* and *Cox* genes*, Park7, Pink1* and *Prkn* (Fig. 6E). Furthermore, GO

and KEGG pathway enrichment analysis of DEGs identified as upregulated in pMacs from aged *R1441C*

females relative to aged *R1441C* males showed a significant enrichment ($p<0.05$) in these same pathways,

including Parkinson's Disease, cellular respiration and ribonucleotide biosynthetic process (Fig. 6F). Given

that pMacs from aged *R1441C* females are susceptible to the age-dependent, bi-phasic immunophenotype

described here, whilst pMacs from aged *R1441C* males seem to be protected, altered mitochondrial function

as a function of age in the females, which is not observed in the males, may underlie this observation.

When comparing DEGs in aged pMacs from both B6 and *R1441C* female mice treated with vehicle from,

unsurprisingly, different DEGs were identified relative to what was seen in IFNγ conditions (Fig. S9A-D).

As opposed to pathways related to mitochondria and oxidative induced by IFNγ conditions, DEGs novel to

R1441C in vehicle conditions were significantly enriched ($p<0.05$) in pathways such as wound healing

responses, and Wnt, ERK and PI3K signaling. It seems, therefore, that at rest, or under vehicle-treatment

conditions, the *R1441C* mutation exerts its effects on aging through pathways crucial for metabolism, cell

survival, and cell growth. The Wnt signaling cascade is crucial in macrophages and serves as a regulator of

macrophage inflammation-resolving state (*55*). However, during an inflammatory insult, the *R1441C*

mutation exerts its effects on aging through pathways crucial for oxidative stress and cellular respiration.

Interestingly, when assessing DEGs in pMacs from aged male mice treated with IFNγ, 540 genes were

identified as novel and upregulated in *R1441C* macrophages that were not significantly affected by aging

in B6 controls (Fig. S9E-G). Pathway enrichment analysis revealed that these 540 upregulated genes

significantly enriched pathways (p<0.05) related to type II interferon production, cytokine production, and

NFkB and JNK signaling (Fig. S9H). We also observed that, with the 5-day LPS stimulation, pMacs from

aged *R1441C* male mice could be pushed to exhibit phenotypes that paralleled those seen in aged female

R1441C macrophages. It has previously been reported that sustained and chronic STAT1 signaling, known

to be stimulated by cytokine signaling (*56*), is necessary to induce macrophage exhaustion (*48*).

Collectively, we interpret our findings that the upregulated DEGs, including driving increased cytokine

release in *R1441C* aged male macrophages, may predispose these macrophages to become exhausted upon

LPS-stimulation.

R1441C- and Y1699C-LRRK2 monocyte-derived patient macrophages exhibit hypophagocytosis and

increased markers for exhaustion

To assess the effects of PD-associated mutations on immune cell exhaustion in humans carrying *LRRK2*

mutations, PBMCs were collected from 2 PD-*Y1699C* and 1 PD-*R1441C* patient, 9 healthy controls, and 3

non-manifesting carriers (NMC) of the *R1441C* pathogenic variant (Fig. 7A). *R1441C* and *Y1699C* variant

are located in the GTPase domain of the LRRK2 protein, and both have been shown to perturb LRRK2's

GTPase activity and have been reported to elevate its kinase activity (*15, 23, 24, 57, 58*). In the cohorts

available for this study, although healthy controls were balanced for sex (5 male and 4 female), all PD

patients and NMCs were male (Table S3).

Pathogen phagocytosis was quantified in monocyte-derived macrophages of these individuals and a

hypophagocytic phenotype was observed in monocyte-derived macrophages from PD-*R1441C* and *Y1699C*

relative to healthy controls and *R1441C* NMCs, as indicated by a reduced number of GFP$^+$ cells (Fig. 7B,

C), with no differences in GFP MFI observed between groups (Fig. S10A). Next, to assess whether

increased markers of immune cell exhaustion accompanied reduced rates of phagocytosis, PDL1 expression

was assessed in peripheral blood mononuclear cells (PBMCs). Consistent with our mouse data, we observed

that PBMCs from both PD-*Y1699C* and PD-*R1441C* patients exhibited increased PDL1$^+$ cell count and

MFI (Fig. 7D, E). When quantifying PDL1 expression in specific immune-cell subsets (Fig. S10B), it was

determined that this overall increase in PDL1 expression was driven by expression in classical monocytes,

with increased PDL1 expression observed in PD-*Y1699C* and *R1441C* classical monocytes relative to

healthy controls and *R1441C* NMCs (Fig. 7F), with no significant differences observed in other cell types

(Fig. S10C, D). Similarly, increased PDL1 expression was evident in monocyte-derived macrophages from

PD-*Y1699C* and *R1441C* patients relative to healthy controls and *R1441C* NMCs (Fig. 7G). Collectively,

the immune cell exhaustion phenotype observed in aged *R1441C* female mice is also associated with an

exhaustion phenotype in cells from patients harboring the *LRRK2-Y1699C* or *R1441C* mutation.

DISCUSSION

Considering age is the greatest risk factor for many neurodegenerative diseases, aging, in particular aging

of the immune system, is the most underappreciated and understudied contributing factor in the

neurodegeneration field. Our novel findings reveal a bi-phasic age- and sex-dependent effect of the *R1441C*

mutation in peripheral immune cells, with increased cytokine release, antigen presentation, phagocytosis

and lysosomal activity in macrophages from young female *R1441C* mice, with many of these phenotypes

ameliorated by Lrrk2 protein kinase inhibition. Our findings further reveal an age-acquired immune cell

exhaustion in macrophages from *R1441C* females associated with hypophagocytosis, decreased antigen

presentation, and reduced lysosomal pan-cathepsin activation and protein degradation, increased pathogen-

mediated DNA breakage, and increased PDL1 expression indicative of immune cell exhaustion.

Importantly, this immune cell exhaustion phenotype was also observed in monocytes and macrophages

from human subjects with PD and harboring the *R1441C* and the *Y1699C* mutation. Furthermore, it was

demonstrated that, although macrophages from male mice did not exhibit this bi-phasic age-dependent

phenotype, they could be pushed to exhaustion with chronic LPS stimulation, suggesting that, along with

age, a secondary inflammatory insult is required to observe immune cell exhaustion phenotypes in

macrophages from *R1441C* males. Finally, it was demonstrated that there is an age-acquired dysregulation

in the expression of genes associated with PD risk, mitochondrial function, and oxidative stress in

macrophages from female *R1441C* mice.

Consistent with findings in *ex vivo* studies in macrophages, we consistently observed elevated expression

of both IL10 and IL6 in circulating plasma from young *R1441C* mice. It has recently been shown that

RAW264.7 cells expressing *T1348N-LRRK2*, an artificial P-Loop null mutation that disrupts GTP binding

(*59, 60*), produce significantly less IL10 relative to wild-type cells in response to LPS and Zymosan (*61*).

IL10 is produced by macrophages and is critical in limiting immune-mediated pathology (*62*), and it was

therefore suggested by Nazish *et al* that there may be a neuroprotective role of LRRK2 in immune signaling

through regulation of IL10 secretion. In addition, we also observed elevations in IL6, which is known to be

promptly and transiently produced in response to infections and tissue injuries during host defense through

the stimulation of immune reactions (*63*). These observations implicate a protective role of Lrrk2 in young

R1441C mice. The pro-inflammatory cytokine, TNF, was also increased in circulating plasma of young

R1441C mice, a compensatory increase that may also contribute to organisms ability to fight infection given

that mice lacking TNF exhibit increased susceptibility to infection with increased bacterial load (*64*). In

support of the protective role of LRRK2 in immune signaling, LRRK2 has been demonstrated to provide

increased pathogen control of *Salmonella typhimurium (32, 65)* and *Listeria monocytogenes* (*33*).

Consistent with these reports, we observed increased *E. coli* uptake and decreased *E. coli*-mediated DNA

damage in macrophages from young *R1441C* female mice, providing further support for a protective role

of LRRK2 in immune signaling and defense against pathogens, at least in a young organism. Given that

LRRK2 mutations are associated with not only PD, but also Crohn's Disease (CD) (*2, 12, 66*) and leprosy

(*67*), it is difficult to understand why these mutations would have persisted in the human population. Given

the evidence discussed and the data presented here, it is plausible to speculate that these mutations may

have persisted due to the immunological advantage they confer to young organisms, including protection

against certain infections.

For the first time, we have reported here an age-dependent, bi-phasic effect of LRRK2 on immune cell

function, with the *R1441C* mutation inducing myeloid cell immune cell exhaustion in a sex-dependent

manner. The concept of immune cell exhaustion was first described as the clonal deletion of virus-specific

CD8$^+$ T-cells that occurs during high-grade chronic infections, and in T-cells is characterized by the

stepwise and progressive loss of T-cell functions (*68*). Regarding exhausted myeloid cells, pathogenic

inflammation and immunosuppression, characterized by decreased antigen presentation, hypophagocytosis

and altered cytokine release, have been described more recently (*48*). The immune system is composed of

numerous cell types with varying states of activation, and cell-to-cell communication during immune

responses is crucial for host defense and resolution of inflammation. The observations made here in innate

immune cells, therefore, likely have downstream effects on myeloid cell communication with adaptive

immune cell types in the periphery. Indeed, alterations in the adaptive immune system have been reported

in PD patients. For example, a meta-analysis of 943 cases of PD identified a reduction in circulating CD4$^+$

T cells in PD-patients (*69, 70*). More specifically, HLA-DR$^+$ T cells and CD45RO$^+$ memory T cells have

been shown to be increased in people with PD relative to healthy age- and sex-matched controls, while

naive CD4$^+$ T cells are reduced (*70-72*). CD4$^+$FOXP3$^+$ Treg cells have increased suppressive activity in

patients with PD (*73*). Interestingly, one characteristic of immune cell exhaustion is the dysregulated cross-

talk between CD4$^+$ and CD8$^+$ T-cells due to the suppressive action of Tregs (*74*). Furthermore, Tregs

directly inhibit the antigen-presenting function of monocytes and macrophages (*75*). Further research is

required to understand the role of cell-to-cell communication in these myeloid cell phenotypes, and if

immune cell exhaustion driven by LRRK2 is a contributing factor.

A sex-dependent effect of the *R1441C* mutation was reported here, with females exhibiting an age-acquired

immune cell exhaustion phenotype which could only be elicited in macrophages from males upon a

secondary, chronic inflammatory insult. Sex-dependent effects of *LRRK2* mutations have previously been

reported. A protective effect of the *G2019S* mutation against *S. typhimurium* has been observed in mice,

with female mutant mice showing increased survival relative to B6 controls which was not observed in

males (*65*). Such observations are also consistent with the sexual dimorphism reported in subjects with PD

who also carry a heterozygous p.*G2019S* mutation in *LRRK2*, with female sex being more common amongst

carriers than non-carriers (*76-78*). This sexual dimorphism has also been reported for the *R1441C/G/H*

mutations (*77*). We observed an *R1441C*-dependent phenotype in macrophages from male mice only when

a secondary, chronic inflammatory insult was applied. It is particularly interesting that genes that are

significantly upregulated with age in *R1441C* macrophages from male mice are associated with pathways

which are heavily implicated in the phenotypes we observed in macrophages from young *R1441C* females.

An increase in gene expression in pathways associated with cytokine production is consistent with our

previous observation of increased IL6, IL10 and TNF release in aged male pMacs relative to B6 controls.

Further research is therefore required to determine the extent to which this sexual dimorphism is driven by

altered immune system susceptibility to *LRRK2* mutations in males and females.

We observed alterations in genes enriched in the KEGG Parkinson's Disease pathway in *R1441C* female

macrophages as a function of age, with these genes being mostly associated with mitochondrial respiration

and oxidative stress. Not surprisingly, mitochondrial dysfunction has emerged as one of the key

mechanisms underlying the pathogenesis of both sporadic PD and familial Parkinsonism (*79, 80*). LRRK2

has been heavily implicated in mitochondrial function and mitophagy, with *LRRK2* mutations causing

reduced mitochondria potential and decreased ATP production, mitochondrial DNA damage, mitochondrial

fragmentation and altered rates of mitophagy in various cell types and pre-clinical models (*81-84*). But

importantly, our findings herein reveal that pathways once thought to be primarily important to neuronal

demise (mitochondrial and oxidative stress) are also disrupted in peripheral immune cells by PD-associated

mutations in a sex- and age-dependent manner and have important implications for monitoring risk, target

engagement, and disease progression of PD and perhaps other NDs by assessing peripheral immune cell

types *ex vivo*. It is known that, in T cells, mitochondrial stress induced by chronic antigen stimulation drives

immune cell exhaustion (*85*). Further research is therefore required to fully understand the role of

mitochondria and mitophagy, and therefore lysosomal function, in the *R1441C*-mediated immune cell

exhaustion phenotypes observed here.

GO and KEGG pathway enrichment analysis of DEGs identified as upregulated in pMacs from aged

R1441C females relative to aged *R1441C* males showed a significant enrichment in pathways related to

mitochondrial function and cellular respiration. Interestingly, estrogen increases expression of respiratory

complexes, antioxidant molecules, and anti-apoptotic factors that directly impact mitochondrial structure

and function, with aging in women being associated with a reduction in estrogen formation and the

subsequent development of mitochondrial dysfunction (*86*). Given that pMacs from aged *R1441C* females

are susceptible to the age-dependent, bi-phasic immunophenotype described here, whilst pMacs from aged

R1441C males seem to be protected, altered mitochondrial function as a consequence of aging may underlie

this observation. It was also observed that, with chronic-LPS stimulation, pMacs from aged *R1441C* males

elicited a phenotype comparable to that seen in pMacs from aged *R1441C* females, suggesting a secondary,

chronic inflammatory insult is needed in conjunction with aging in *R1441C* males for immune cell

exhaustion to occur. It has previously been reported that upon LPS stimulation, decreased mitochondrial

membrane potential decreases and ROS levels increase in macrophages (*87*). Further research is therefore

required to decipher if altered mitochondrial function underlies the synergizing effects of aging and chronic

inflammation in *R1441C*-induced immune cell exhaustion in macrophages from males.

It has previously been shown that LRRK2 exerts differential effects on inflammatory responses in a tissue-

specific manner, and it has been suggested that there is a differential role of LRRK2 in the periphery versus

the CNS (*65, 88*). It therefore remains to be determined how the data reported here impacts our

understanding of LRRK2 in myeloid cells in both the periphery and the CNS. Indeed, we observed a

decrease in MHC-II expression in peripheral myeloid cells of aged *R1441C* females, whilst microglia

expressed increased MHC-II in these mice relative to B6 controls, implying a differential effect of LRRK2

in the periphery vs. the CNS. Alterations in type II interferon responses have been observed in both induced

pluripotent stem cell (iPSC)-derived microglia and macrophages from patients harboring *G2019S* mutations

(*89*), and similar effects of *LRRK2* mutations on lysosomal function have also been reported in patient

iPSC-derived microglia and macrophages (*90*). However, it has been demonstrated that expression of

R1441C-Lrrk2 in peripheral lymphocytes alone is sufficient to induce dopaminergic neuronal cell loss in

the midbrains of LPS-treated mice (*39*), suggesting that the periphery plays a pertinent and perhaps primary

role in *LRRK2*-PD, and future research into how immune cell exhaustion in the periphery impacts cross-

talk with the CNS is needed to inform future immunomodulatory therapeutic interventions for PD.

We successfully demonstrated that, like macrophages from aged *R1441C* mice, myeloid cells from *R1441C*

and *Y1699C* patients exhibited hypophagocytosis and increased expression of immune cell exhaustion

markers relative to those from HCs and *R1441C*-NMCs. However, it is unclear if the same hyper-responsive

phenotype that was observed in young *R1441C* female mice is exhibited in myeloid cells in *R1441C* carriers

at a younger age or prior to disease onset. Indeed, we did show that *R1441C*-NMCs did not exhibit these

immune cell exhaustion phenotypes, and it is noteworthy that the average age of the 3 *R1441C*-NMCs in

this study was 30.5 years (Table S3), whilst the average age of the *R1441C* and *Y1699C*-PD subjects was

42 and 63.5, respectively. Further investigation is therefore required to determine if PBMCs from NMCs,

upon follow-up after further aging or disease onset, exhibit immune cell exhaustion. As well, it is interesting

to note that, despite the immune cell exhaustion phenotype being observed in female *R1441C* KI mice, all

the *R1441C* and *Y1699C*-PD patients recruited here were male (Table S3), and PBMCs from these male

patients exhibited similar immune cell exhaustion phenotypes observed in female *R1441C* mice. We do,

however, observe these immune cell exhaustion phenotypes in macrophages from aged *R1441C* males upon

a secondary, chronic inflammatory insult. Such observations support the hypothesis that complex gene-by-

environment interactions combine with an ageing immune system to create the 'perfect storm' that enables

the development and progression of PD (*5*). Further research is therefore required to determine if sex plays

a role in this 'perfect storm'. Furthermore, further research is required to understand if these phenotypes

are exclusive to *LRRK2*-pathogenic variants located in the GTPase domain or if they are seen in other

LRRK2 mutant immune cells, such as the more common *G2019S* mutation, as well as in idiopathic PD or

sporadic cases.

LRRK2-targeting therapies, including antisense oligonucleotides (ASOs) and kinase inhibitors, have been

suggested as a potential therapeutic for *LRRK2*-PD, with many hypothesizing that targeting the increased

LRRK2 levels and kinase activity will be beneficial to CNS-related pathology. Our findings suggest that

more research is required to understand the location and optimal timing for the administration of these

LRRK2-targeting therapeutics. Specifically, if LRRK2 plays a critical role in pathogen and infection control

in a pathogen or tissue-specific manner, it may mean that LRRK2-targeting therapies may need to be limited

to carriers of *LRRK2* pathogenic variants and/or may need to be delivered in a CNS compartment-targeted

manner as opposed to systemically to avoid deleterious collateral damage to immune function. Indeed, the

LRRK2 ASO currently being tested in the clinic is being delivered via intrathecal injection directly into the

cisterna magna, thus limiting systemic exposure (NCT03976349). Lastly, perhaps somewhat surprisingly,

we found that, while LRRK2 kinase inhibition could rescue some of the effector functions of young mutant-

LRRK2 mice, it had no beneficial effects on the immune-cell exhaustion phenotypes observed here in

macrophages from aged mice; however, it did prove to be protective in macrophages from young mice to

prevent the development of an immune-cell exhaustive phenotype. Such observations suggest that LRRK2-

targeting therapies would elicit the most beneficial effects during the prodromal phases of disease, and

further research on PBMCs from prodromal patients with pre-motor features at risk for development of

motor PD due to a family history will be required to investigate this possibility and inform future clinical

trials.

Our study is not without limitations, and further steps are needed to translate our findings into clinical

samples further. Firstly, although we provide evidence of immune cell exhaustion in patient immune cells,

this was carried out in a small population due to the availability of cells to us at the time. It is hard to

conclude here whether *R1441C*-NMC did not exhibit immune cell exhaustion phenotypes due to the fact

that these phenotypes only present themselves upon disease onset, or, whether immune cell exhaustion

phenotypes are not present amongst all *R1441C* carriers. Furthermore, due to the fact that penetrance of

these mutations is low, it may be that the NMCs recruited for this study will not develop PD and this

explains their lack of immune cell exhaustion phenotypes. Larger, longitudinal studies are therefore

required in order to decipher between these possibilities. Secondly, due to the lack of a longitudinal study

design and limited sample numbers, it is hard to conclude whether age or chronic immune insult are the

primary driver of immune cell exhaustion in our patient samples, and whether immune cell exhaustion is a

key driver of PD onset and/or progression. Thirdly, although the *G2019S-LRRK2* mutation is the largest

contributor to familial forms of PD, and LRRK2 has been implicated in sporadic forms of the disease, the

R1441C mutation is comparably rarer, with prevalence estimates between 0.03-2% (*91*). The focus on

R1441C, therefore, may be limited, and further studies are required to determine if these phenotypes are

mutation-specific and if they are seen in sporadic and idiopathic-PD cases. Lastly, although increased PDL1

expression and hypophagocytosis were observed in patient samples, there are other effector functions

altered due to immune cell exhaustion, and in other cell types, such as cytokine release, T cell proliferation

and differentiation. Further studies in clinical and pre-clinical samples are therefore required to understand

the effects of LRRK2 mutations on other immune cell effector functions and how immune cell exhaustion

may be implicated in this.

MATERIALS AND METHODS

Study design

The primary objective of this research was to investigate the effects of the *R1441C LRRK2* pathogenic

variant on myeloid cell effector function in a sex- and age-dependent manner. We hypothesized that a

biphasic, age-dependent effect of the *R1441C* pathogenic variant would be observed, affecting antigen

presentation, cytokine release, phagocytosis and lysosomal function in macrophages. To explore our

hypothesis, we used *R1441C* KI mice and B6 controls and assessed myeloid cell effector function both *ex*

vivo and *in vivo,* in males and females at 2-3 months and 18-21 months. All studies were performed by

investigators blinded to genotype and treatment groups at the University of Florida, USA. Power analyses

based on the labs pilot and published data indicated that n = 8-12 per group for flow-cytometry and cytokine

release assays, and n = 4-6 for imaging based immunophenotyping and immunoblotting would be needed

to yield 80% power with a α set to 0.05. Exclusion criteria for experiments involving mice included mice

who reached endpoint before time of experiment (due to health-related issues), mice who were found to

have tumors whilst collecting tissue, and datapoints that fell more than 3SD +/- of the average of that group.

Multiple cohorts of mice (each cohort consisting of 6-12 mice per group depending on the assay) were used

to generate this data set. As 2 sexes, 2 genotypes and 2 ages were used to generate this data, each cohort

consisted of 8 groups, equating to 48-96 mice. Each cohort was separated into batches and experiments

carried out, and data acquired and analyzed for each batch, with data from batches merged post-analysis to

form the overall data set for that cohort. As well, entirely independent cohorts of mice were used to produce

data using orthogonal assays (e.g. 1 cohort used for flow cytometry, 1 cohort used for microscopy and

western blotting analysis) to ensure the phenotypes we were observing were not assay- or cohort-dependent.

By generating this data across batches of animals and using multiple orthogonal assays to confirm

phenotypes in separate cohorts of mice, we ensure that our data is robust and reproducible. A schematic of

this study design is shown in Fig.S11.

The study used human PBMCs received from collaborators at University Hospital Schleswig-Holstein,

University of Luebeck, Germany and the University Hospital Eppendorf, Hamburg, Germany. Samples

were received at the University of Florida and investigators were unblinded after completion of analysis.

Deidentified data provided included sex, age at collection, age of diagnosis, clinical diagnosis and genetic

status.

Animals

R1441C knock-in (Strain #009346; B6.Cg-Lrrk2tm1.1Shn/J) mice have previously been characterized (*92*)

and were maintained in the McKnight Brain Institute vivarium (University of Florida) at 22°C at 60-70%

humidity with a 12-hour light/dark cycle and *ad libitum* access to standard rodent chow and water. C57BL/6

(Strain #000664) controls were used for all studies. All animal procedures were approved by the University

of Florida Institutional Animal Care and Use Committee and were in accordance with the National Institute

of Health *Guide for the Care and Use of Laboratory Animals* (NIH Publications No. 80-23) revised 1996.

Female and male mice were aged to 2-3 or 18-21-months-old and sacrificed via cervical dislocation or rapid

decapitation.

Human subjects

This study was reviewed and approved by the University of Lubeck Institutional Review Board (AZ16-

039). Participants provided written informed consent to participate. During recruitment, a family history

was used to assess history of disease and patients were screened for *LRRK2* variants. See Table S3 for

patient demographics.

Statistics and data analysis

All individual level data can be found in data file S1 and is also available on Zenodo

(10.5281/zenodo.11099669). Data and statistical analyses were performed using IBM SPSS statistics 27 or

GraphPad Prism 9. For assessing differences between groups, data were analyzed by either 1-way or 2-way

analysis of variance (ANOVA), or by t-test. In instances when data did not fit parametric assumptions,

Kruskal-Wallis non-parametric ANOVA was used. *Post hoc* tests following ANOVAs were conducted

using Tukey HSD or Bonferroni correction. Two-tailed levels of significance were used and $p < 0.05$ was

considered statistically significant. Graphs are depicted by means +/- standard error of the mean (SEM).

List of Supplementary Materials:

Materials and Methods

Supplementary figures:

Figure S1. Characterization of *ex vivo* pMac cultures.

Figure S2. *R1441C* mutation leads to age dependent biphasic alteration in antigen presentation.

Figure S3. The *R1441C* Lrrk2 mutation causes alterations of monocyte population and microglia activation

in aged female mice.

Figure S4. *R1441C* mutation leads to age-dependent biphasic alteration in lysosomal activity.

Figure S5. *R1441C* Lrrk2 mutation alters pathogen uptake and pathogen-mediated DNA damage in age-

and sex-dependent manner.

Figure S6. *R1441C* Lrrk2 mutation induces immune cell exhaustion in a kinase-dependent manner.

Figure S7. Transcriptomic profiling reveals alterations in critical effector function pathways in *R1441C*

macrophages in age- and sex- dependent manner.

Figure S8. *R1441C* Lrrk2 mutation alters biological processes in macrophages in a sex- and age-dependent

manner.

Figure S9. *R1441C* Lrrk2 mutation alters transcriptional changes associated with age in sex dependent

manner.

Figure S10. *R1441C*- and *Y1699C*-LRRK2 monocyte derived macrophages exhibit hypo phagocytosis and

increased markers for immune cell exhaustion.

Figure S11. Schematic of study design with mice.

Supplementary tables:

Table S1. Flow cytometry marker antibody panel (mouse)

Table S2. Antibodies for immunoblotting

Table S3. Patient demographics

Table S4. Flow cytometry marker antibody panel (human)

References (*93-98*)

References

1. Von Campenhausen, S., B. Bornschein, R. Wick, K. Bötzel, C. Sampaio, W. Poewe, W. Oertel, U. Siebert,
K. Berger, and R. Dodel, Prevalence and incidence of Parkinson's disease in Europe. **15**(4): p. 473-490
(2005).
2. Marras, C., J.C. Beck, J.H. Bower, E. Roberts, B. Ritz, G.W. Ross, R.D. Abbott, R. Savica, S.K. Van Den
Eeden, A.W. Willis, C.M. Tanner, and P.G. Parkinson's Foundation, Prevalence of Parkinson's disease across
North America. *NPJ Parkinsons Dis*. **4**: p. 21 (2018).
3. Yang, W., J.L. Hamilton, C. Kopil, J.C. Beck, C.M. Tanner, R.L. Albin, E. Ray Dorsey, N. Dahodwala, I.
Cintina, P. Hogan, and T. Thompson, Current and projected future economic burden of Parkinson's disease
in the U.S. *npj Parkinson's Disease*. **6**(1) (2020).
4. Bloem, B.R., M.S. Okun, and C. Klein, Parkinson's disease. *Lancet*. **397**(10291): p. 2284-2303 (2021).
5. Tansey, M.G., Wallings, R. L., Houser, M. C., Herrick, M. K., Keating, C. E. & Joers, V., The contributions
of inflammation and immune dysfunction to Parkinson disease. *Nature Reviews Immunology*. (2022).
6. Bliederhaeuser, C., L. Zondler, V. Grozdanov, W.P. Ruf, D. Brenner, H.L. Melrose, P. Bauer, A.C. Ludolph,
F. Gillardon, J. Kassubek, J.H. Weishaupt, and K.M. Danzer, LRRK2 contributes to monocyte dysregulation
in Parkinson's disease. *Acta Neuropathol Commun*. **4**(1): p. 123 (2016).
7. Grozdanov, V., C. Bliederhaeuser, W.P. Ruf, V. Roth, K. Fundel-Clemens, L. Zondler, D. Brenner, A.
Martin-Villalba, B. Hengerer, J. Kassubek, A.C. Ludolph, J.H. Weishaupt, and K.M. Danzer, Inflammatory
dysregulation of blood monocytes in Parkinson's disease patients. **128**(5): p. 651-663 (2014).
8. Grozdanov, V., L. Bousset, M. Hoffmeister, C. Bliederhaeuser, C. Meier, K. Madiona, L. Pieri, M. Kiechle,
P.J. McLean, J. Kassubek, C. Behrends, A.C. Ludolph, J.H. Weishaupt, R. Melki, and K.M. Danzer,
Increased Immune Activation by Pathologic alpha-Synuclein in Parkinson's Disease. *Ann Neurol*. **86**(4): p.
593-606 (2019).
9. Cook, D.A., G.T. Kannarkat, A.F. Cintron, L.M. Butkovich, K.B. Fraser, J. Chang, N. Grigoryan, S.A.
Factor, A.B. West, J.M. Boss, and M.G. Tansey, LRRK2 levels in immune cells are increased in Parkinson's
disease. *NPJ Parkinsons Dis*. **3**: p. 11 (2017).
10. Kline, E.M., M.C. Houser, M.K. Herrick, P. Seibler, C. Klein, A.B. West, and T. M.G., Genetic and
Environmental Factors in Parkinson's Disease Converge on Immune Function and Inflammation. *Movement
Disorders*. **36**(1): p. 25-36 (2021).
11. Houser, M.C. and M.G. Tansey, The gut-brain axis: is intestinal inflammation a silent driver of Parkinson's
disease pathogenesis? *NPJ Parkinsons Dis*. **3**: p. 3 (2017).
12. Herrick, M.K. and M.G. Tansey, Is LRRK2 the missing link between inflammatory bowel disease and
Parkinson's disease? *npj Parkinson's Disease*. **7**(1) (2021).
13. Blauwendraat, C., M.A. Nalls, and A.B. Singleton, The genetic architecture of Parkinson's disease. *Lancet
Neurol*. **19**(2): p. 170-178 (2020).
14. West, A.B., D.J. Moore, S. Biskup, A. Bugayenko, W.W. Smith, C.A. Ross, V.L. Dawson, and T.M. Dawson,
Parkinson's disease-associated mutations in leucine-rich repeat kinase 2 augment kinase activity. *Proc Natl
Acad Sci U S A*. **102**(46): p. 16842-7 (2005).
15. West, A.B., D.J. Moore, C. Choi, S.A. Andrabi, X. Li, D. Dikeman, S. Biskup, Z. Zhang, K.L. Lim, V.L.
Dawson, and T.M. Dawson, Parkinson's disease-associated mutations in LRRK2 link enhanced GTP-binding
and kinase activities to neuronal toxicity. *Hum Mol Genet*. **16**(2): p. 223-32 (2007).
16. Funayama, M., K. Hasegawa, H. Kowa, M. Saito, S. Tsuji, and F. Obata, A new locus for Parkinson's disease
(PARK8) maps to chromosome 12p11.2-q13.1. *Ann Neurol*. **51**(3): p. 296-301 (2002).
17. Funayama, M., Y. Li, H. Tomiyama, H. Yoshino, Y. Imamichi, M. Yamamoto, M. Murata, T. Toda, Y.
Mizuno, and N. Hattori, Leucine-rich repeat kinase 2 G2385R variant is a risk factor for Parkinson disease
in Asian population. *Neuroreport*. **18**(3): p. 273-5 (2007).
18. Hatano, T., M. Funayama, S.I. Kubo, I.F. Mata, Y. Oji, A. Mori, C.P. Zabetian, S.M. Waldherr, H. Yoshino,
G. Oyama, Y. Shimo, K.I. Fujimoto, H. Oshima, Y. Kunii, H. Yabe, Y. Mizuno, and N. Hattori, Identification
of a Japanese family with LRRK2 p.R1441G-related Parkinson's disease. *Neurobiol Aging*. **35**(11): p. 2656
e17-2656 e23 (2014).

19.   Ross, O.A., Y.R. Wu, M.C. Lee, M. Funayama, M.L. Chen, A.I. Soto, I.F. Mata, G.J. Lee-Chen, C.M. Chen,
M. Tang, Y. Zhao, N. Hattori, M.J. Farrer, E.K. Tan, and R.M. Wu, Analysis of Lrrk2 R1628P as a risk factor
for Parkinson's disease. *Ann Neurol*. **64**(1): p. 88-92 (2008).
20.   Haugarvoll, K. and Z.K. Wszolek, Clinical features of LRRK2 parkinsonism. **15**: p. S205-S208 (2009).
21.   Satake, W., Y. Nakabayashi, I. Mizuta, Y. Hirota, C. Ito, M. Kubo, T. Kawaguchi, T. Tsunoda, M. Watanabe,
A. Takeda, H. Tomiyama, K. Nakashima, K. Hasegawa, F. Obata, T. Yoshikawa, H. Kawakami, S. Sakoda,
M. Yamamoto, N. Hattori, M. Murata, Y. Nakamura, and T. Toda, Genome-wide association study identifies
common variants at four loci as genetic risk factors for Parkinson's disease. *Nat Genet*. **41**(12): p. 1303-7
(2009).
22.   Gaig, C., M. Ezquerra, M.J. Marti, E. Muñoz, F. Valldeoriola, and E. Tolosa, LRRK2 Mutations in Spanish
Patients With Parkinson Disease. *Archives of Neurology*. **63**(3): p. 377 (2006).
23.   Steger, M., F. Tonelli, G. Ito, P. Davies, M. Trost, M. Vetter, S. Wachter, E. Lorentzen, G. Duddy, S. Wilson,
M.A. Baptista, B.K. Fiske, M.J. Fell, J.A. Morrow, A.D. Reith, D.R. Alessi, and M. Mann,
Phosphoproteomics reveals that Parkinson's disease kinase LRRK2 regulates a subset of Rab GTPases. *Elife*.
5 (2016).
24.   Lewis, P.A., E. Greggio, A. Beilina, S. Jain, A. Baker, and M.R. Cookson, The R1441C mutation of LRRK2
disrupts GTP hydrolysis. **357**(3): p. 668-671 (2007).
25.   Gardet, A., Y. Benita, C. Li, B.E. Sands, I. Ballester, C. Stevens, J.R. Korzenik, J.D. Rioux, M.J. Daly, R.J.
Xavier, and D.K. Podolsky, LRRK2 Is Involved in the IFN- Response and Host Response to Pathogens. *The
Journal of Immunology*. **185**(9): p. 5577-5585 (2010).
26.   Atashrazm, F., D. Hammond, G. Perera, M.F. Bolliger, E. Matar, G.M. Halliday, B. Schule, S.J.G. Lewis,
R.J. Nichols, and N. Dzamko, LRRK2-mediated Rab10 phosphorylation in immune cells from Parkinson's
disease patients. *Mov Disord*. **34**(3): p. 406-415 (2019).
27.   Wallings, R.L., L.P. Hughes, H.A. Staley, Z.D. Simon, N.R. McFarland, R.N. Alcalay, A. Garrido, M.J.
Marti, E.T. Sarro, N. Dzamko, and M.G. Tansey, WHOPPA Enables Parallel Assessment of Leucine-Rich
Repeat Kinase 2 and Glucocerebrosidase Enzymatic Activity in Parkinson's Disease Monocytes. *Front Cell
Neurosci*. **16**: p. 892899 (2022).
28.   Kozina, E., S. Sadasivan, Y. Jiao, Y. Dou, Z. Ma, H. Tan, K. Kodali, T. Shaw, J. Peng, and R.J. Smeyne,
Mutant LRRK2 mediates peripheral and central immune responses leading to neurodegeneration in vivo.
Brain. **141**(6): p. 1753-1769 (2018).
29.   Rideout, H.J. and D.B. Re, LRRK2 and the "LRRKtosome" at the Crossroads of Programmed Cell Death:
Clues from RIP Kinase Relatives. *Adv Neurobiol*. **14**: p. 193-208 (2017).
30.   Wang, Z., S. Arat, M. Magid-Slav, and J.R. Brown, Meta-analysis of human gene expression in response to
Mycobacterium tuberculosis infection reveals potential therapeutic targets. *BMC Systems Biology*. **12**(1)
(2018).
31.   Hartlova, A., S. Herbst, J. Peltier, A. Rodgers, O. Bilkei-Gorzo, A. Fearns, B.D. Dill, H. Lee, R. Flynn, S.A.
Cowley, P. Davies, P.A. Lewis, I.G. Ganley, J. Martinez, D.R. Alessi, A.D. Reith, M. Trost, and M.G.
Gutierrez, LRRK2 is a negative regulator of Mycobacterium tuberculosis phagosome maturation in
macrophages. *EMBO J*. **37**(12) (2018).
32.   Liu, W., X.N. Liu, Y. Li, J. Zhao, Z. Liu, Z. Hu, Y. Wang, Y. Yao, A.W. Miller, B. Su, M.R. Cookson, X.
Li, and Z. Kang, LRRK2 promotes the activation of NLRC4 inflammasome during Salmonella Typhimurium
infection. *The Journal of Experimental Medicine*. **214**(10): p. 3051-3066 (2017).
33.   Zhang, Q., Y. Pan, R. Yan, B. Zeng, H. Wang, X. Zhang, W. Li, H. Wei, and Z. Liu, Commensal bacteria
direct selective cargo sorting to promote symbiosis. **16**(9): p. 918-926 (2015).
34.   Fava, V.M., J. Manry, A. Cobat, M. Orlova, N. Van Thuc, N.N. Ba, V.H. Thai, L. Abel, A. Alcaïs, and E.
Schurr, A Missense LRRK2 Variant Is a Risk Factor for Excessive Inflammatory Responses in Leprosy.
PLOS Neglected Tropical Diseases. **10**(2): p. e0004412 (2016).
35.   Witoelar, A., I.E. Jansen, Y. Wang, R.S. Desikan, J.R. Gibbs, C. Blauwendraat, W.K. Thompson, D.G.
Hernandez, S. Djurovic, A.J. Schork, F. Bettella, D. Ellinghaus, A. Franke, B.A. Lie, L.K. McEvoy, T.H.
Karlsen, S. Lesage, H.R. Morris, A. Brice, N.W. Wood, P. Heutink, J. Hardy, A.B. Singleton, A.M. Dale, T.
Gasser, O.A. Andreassen, and M. Sharma, Genome-wide Pleiotropy Between Parkinson Disease and
Autoimmune Diseases. *JAMA Neurology*. **74**(7): p. 780 (2017).
36.   Villumsen, M., S. Aznar, B. Pakkenberg, T. Jess, and T. Brudek, Inflammatory bowel disease increases the
risk of Parkinson's disease: a Danish nationwide cohort study 1977–2014. *Gut*. **68**(1): p. 18-24 (2019).
37.   Skaper, S.D., L. Facci, M. Zusso, and P. Giusti, An Inflammation-Centric View of Neurological Disease:
Beyond the Neuron. *Frontiers in Cellular Neuroscience*. **12** (2018).

38. Gelders, G., V. Baekelandt, and A. Van der Perren, Linking Neuroinflammation and Neurodegeneration in
Parkinson's Disease. *J Immunol Res*. **2018**: p. 4784268 (2018).
39. Kozina, E., M. Byrne, and R.J. Smeyne, Mutant LRRK2 in lymphocytes regulates neurodegeneration via IL-
6 in an inflammatory model of Parkinson's disease. *npj Parkinson's Disease*. **8**(1) (2022).
40. Kim, K.S., P.C. Marcogliese, J. Yang, S.M. Callaghan, V. Resende, E. Abdel-Messih, C. Marras, N.P.
Visanji, J. Huang, M.G. Schlossmacher, L. Trinkle-Mulcahy, R.S. Slack, A.E. Lang, and D.S. Park,
Regulation of myeloid cell phagocytosis by LRRK2 via WAVE2 complex stabilization is altered in
Parkinson's disease. *Proceedings of the National Academy of Sciences*. **115**(22): p. E5164-E5173 (2018).
41. Liu, Z., E. Xu, H.T. Zhao, T. Cole, and A.B. West, LRRK2 and Rab10 coordinate macropinocytosis to
mediate immunological responses in phagocytes. *EMBO J*. **39**(20): p. e104862 (2020).
42. Tong, Y., E. Giaime, H. Yamaguchi, T. Ichimura, Y. Liu, H. Si, H. Cai, J.V. Bonventre, and J. Shen, Loss of
leucine-rich repeat kinase 2 causes age-dependent bi-phasic alterations of the autophagy pathway. **7**(1): p. 2
(2012).
43. Cassado, A.D.A., M.R. D'Império Lima, and K.R. Bortoluci, Revisiting Mouse Peritoneal
Macrophages: Heterogeneity, Development, and Function. *Frontiers in Immunology*. **6** (2015).
44. Schapansky, J., J.D. Nardozzi, F. Felizia, and M.J. Lavoie, Membrane recruitment of endogenous LRRK2
precedes its potent regulation of autophagy. **23**(16): p. 4201-4214 (2014).
45. Hakimi, M., T. Selvanantham, E. Swinton, R.F. Padmore, Y. Tong, G. Kabbach, K. Venderova, S.E.
Girardin, D.E. Bulman, C.R. Scherzer, M.J. LaVoie, D. Gris, D.S. Park, J.B. Angel, J. Shen, D.J. Philpott,
and M.G. Schlossmacher, Parkinson's disease-linked LRRK2 is expressed in circulating and tissue immune
cells and upregulated following recognition of microbial structures. *J Neural Transm (Vienna)*. **118**(5): p.
795-808 (2011).
46. Moehle, M.S., J.P.L. Daher, T.D. Hull, R. Boddu, H.A. Abdelmotilib, J. Mobley, G.T. Kannarkat, M.G.
Tansey, and A.B. West, The G2019S LRRK2 mutation increases myeloid cell chemotactic responses and
enhances LRRK2 binding to actin-regulatory proteins. **24**(15): p. 4250-4267 (2015).
47. Kaminski, H., M. Lemoine, and T. Pradeu, Immunological exhaustion: How to make a disparate concept
operational? *PLOS Pathogens*. **17**(9): p. e1009892 (2021).
48. Pradhan, K., Z. Yi, S. Geng, and L. Li, Development of Exhausted Memory Monocytes and Underlying
Mechanisms. *Front Immunol*. **12**: p. 778830 (2021).
49. Strader, S. and A.B. West, The interplay between monocytes, alpha-synuclein and LRRK2 in Parkinson's
disease. *Biochem Soc Trans*. **51**(2): p. 747-758 (2023).
50. Jin, J., H. Zhang, C.M. Weyand, and J.J. Goronzy, Lysosomes in T Cell Immunity and Aging. *Front Aging*.
2: p. 809539 (2021).
51. Hennies, C.M., M.A. Lehn, and E.M. Janssen, Quantitating MHC class II trafficking in primary dendritic
cells using imaging flow cytometry. *J Immunol Methods*. **423**: p. 18-28 (2015).
52. Watts, C., The endosome-lysosome pathway and information generation in the immune system. *Biochim
Biophys Acta*. **1824**(1): p. 14-21 (2012).
53. Boomer, J.S., J.M. Green, and R.S. Hotchkiss, The changing immune system in sepsis. *Virulence*. **5**(1): p.
45-56 (2014).
54. Jubel, J.M., Z.R. Barbati, C. Burger, D.C. Wirtz, and F.A. Schildberg, The Role of PD-1 in Acute and Chronic
Infection. *Front Immunol*. **11**: p. 487 (2020).
55. Weinstock, A., K. Rahman, O. Yaacov, H. Nishi, P. Menon, C.A. Nikain, M.L. Garabedian, S. Pena, N.
Akbar, B.E. Sansbury, S.P. Heffron, J. Liu, G. Marecki, D. Fernandez, E.J. Brown, K.V. Ruggles, S.A.
Ramsey, C. Giannarelli, M. Spite, R.P. Choudhury, P. Loke, and E.A. Fisher, Wnt signaling enhances
macrophage responses to IL-4 and promotes resolution of atherosclerosis. *Elife*. **10** (2021).
56. Morris, R., N.J. Kershaw, and J.J. Babon, The molecular details of cytokine signaling via the JAK/STAT
pathway. *Protein Science*. **27**(12): p. 1984-2009 (2018).
57. West, A.B., D.J. Moore, S. Biskup, A. Bugayenko, W.W. Smith, C.A. Ross, V.L. Dawson, and T.M. Dawson,
From The Cover: Parkinson's disease-associated mutations in leucine-rich repeat kinase 2 augment kinase
activity. **102**(46): p. 16842-16847 (2005).
58. Daniëls, V., R. Vancraenenbroeck, B.M.H. Law, E. Greggio, E. Lobbestael, F. Gao, M. De Maeyer, M.R.
Cookson, K. Harvey, V. Baekelandt, and J.-M. Taymans, Insight into the mode of action of the LRRK2
Y1699C pathogenic mutant. **116**(2): p. 304-315 (2011).
59. Biosa, A., A. Trancikova, L. Civiero, L. Glauser, L. Bubacco, E. Greggio, and D.J. Moore, GTPase activity
regulates kinase activity and cellular phenotypes of Parkinson's disease-associated LRRK2. **22**(6): p. 1140-
1156 (2013).

60. Nguyen, A.P. and D.J. Moore, Understanding the GTPase Activity of LRRK2: Regulation, Function, and
Neurotoxicity. *Adv Neurobiol*. **14**: p. 71-88 (2017).
61. Nazish, I., C. Arber, T.M. Piers, T.T. Warner, J.A. Hardy, P.A. Lewis, J.M. Pocock, and R. Bandopadhyay,
Abrogation of LRRK2 dependent Rab10 phosphorylation with TLR4 activation and alterations in evoked
cytokine release in immune cells. *Neurochem Int*. **147**: p. 105070 (2021).
62. Sanin, D.E., C.T. Prendergast, and A.P. Mountford, IL-10 Production in Macrophages Is Regulated by a
TLR-Driven CREB-Mediated Mechanism That Is Linked to Genes Involved in Cell Metabolism. *J Immunol*.
195(3): p. 1218-32 (2015).
63. Tanaka, T., M. Narazaki, and T. Kishimoto, IL-6 in inflammation, immunity, and disease. *Cold Spring Harb*
Perspect Biol. **6**(10): p. a016295 (2014).
64. Li, X., H. Korner, and X. Liu, Susceptibility to Intracellular Infections: Contributions of TNF to Immune
Defense. *Front Microbiol*. **11**: p. 1643 (2020).
65. Shutinoski, B., M. Hakimi, I.E. Harmsen, M. Lunn, J. Rocha, N. Lengacher, Y.Y. Zhou, J. Khan, A. Nguyen,
Q. Hake-Volling, D. El-Kodsi, J. Li, A. Alikashani, C. Beauchamp, J. Majithia, K. Coombs, D. Shimshek,
P.C. Marcogliese, D.S. Park, J.D. Rioux, D.J. Philpott, J.M. Woulfe, S. Hayley, S. Sad, J.J. Tomlinson, E.G.
Brown, and M.G. Schlossmacher, Lrrk2 alleles modulate inflammation during microbial infection of mice in
a sex-dependent manner. *Science Translational Medicine*. **11**(511): p. eaas9292 (2019).
66. Daseke, M.J., 2nd, M.A.A. Tenkorang-Impraim, Y. Ma, U. Chalise, S.R. Konfrst, M.R. Garrett, K.Y.
DeLeon-Pennell, and M.L. Lindsey, Exogenous IL-4 shuts off pro-inflammation in neutrophils while
stimulating anti-inflammation in macrophages to induce neutrophil phagocytosis following myocardial
infarction. *J Mol Cell Cardiol*. **145**: p. 112-121 (2020).
67. Fava, V., M. Orlova, A. Cobat, A. Alcaïs, M. Mira, and E. Schurr, Genetics of leprosy reactions: an overview.
107: p. 132-142 (2012).
68. Yi, J.S., M.A. Cox, and A.J. Zajac, T-cell exhaustion: characteristics, causes and conversion. *Immunology*.
129(4): p. 474-81 (2010).
69. Jiang, S., H. Gao, Q. Luo, P. Wang, and X. Yang, The correlation of lymphocyte subsets, natural killer cell,
and Parkinson's disease: a meta-analysis. *Neurol Sci*. **38**(8): p. 1373-1380 (2017).
70. Saunders, J.A., K.A. Estes, L.M. Kosloski, H.E. Allen, K.M. Dempsey, D.R. Torres-Russotto, J.L. Meza,
P.M. Santamaria, J.M. Bertoni, D.L. Murman, H.H. Ali, D.G. Standaert, R.L. Mosley, and H.E. Gendelman,
CD4+ regulatory and effector/memory T cell subsets profile motor dysfunction in Parkinson's disease. *J*
Neuroimmune Pharmacol. **7**(4): p. 927-38 (2012).
71. Fiszer, U., E. Mix, S. Fredrikson, V. Kostulas, and H. Link, Parkinson's disease and immunological
abnormalities: increase of HLA-DR expression on monocytes in cerebrospinal fluid and of CD45RO+ T cells
in peripheral blood. *Acta Neurol Scand*. **90**(3): p. 160-6 (1994).
72. Kustrimovic, N., C. Comi, L. Magistrelli, E. Rasini, M. Legnaro, R. Bombelli, I. Aleksic, F. Blandini, B.
Minafra, G. Riboldazzi, A. Sturchio, M. Mauri, G. Bono, F. Marino, and M. Cosentino, Parkinson's disease
patients have a complex phenotypic and functional Th1 bias: cross-sectional studies of CD4+ Th1/Th2/T17
and Treg in drug-naive and drug-treated patients. *J Neuroinflammation*. **15**(1): p. 205 (2018).
73. Rosenkranz, D., S. Weyer, E. Tolosa, A. Gaenslen, D. Berg, T. Leyhe, T. Gasser, and L. Stoltze, Higher
frequency of regulatory T cells in the elderly and increased suppressive activity in neurodegeneration. *J*
Neuroimmunol. **188**(1-2): p. 117-27 (2007).
74. McRitchie, B.R. and B. Akkaya, Exhaust the exhausters: Targeting regulatory T cells in the tumor
microenvironment. *Front Immunol*. **13**: p. 940052 (2022).
75. Tiemessen, M.M., A.L. Jagger, H.G. Evans, M.J. van Herwijnen, S. John, and L.S. Taams,
CD4+CD25+Foxp3+ regulatory T cells induce alternative activation of human monocytes/macrophages.
Proc Natl Acad Sci U S A. **104**(49): p. 19446-51 (2007).
76. Marder, K., Y. Wang, R.N. Alcalay, H. Mejia-Santana, M.X. Tang, A. Lee, D. Raymond, A. Mirelman, R.
Saunders-Pullman, L. Clark, L. Ozelius, A. Orr-Urtreger, N. Giladi, S. Bressman, and L.A.J. Consortium,
Age-specific penetrance of LRRK2 G2019S in the Michael J. Fox Ashkenazi Jewish LRRK2 Consortium.
Neurology. **85**(1): p. 89-95 (2015).
77. Cilia, R., C. Siri, D. Rusconi, R. Allegra, A. Ghiglietti, G. Sacilotto, M. Zini, A.L. Zecchinelli, R. Asselta, S.
Duga, A.M. Paganoni, G. Pezzoli, M. Seia, and S. Goldwurm, LRRK2 mutations in Parkinson's disease:
confirmation of a gender effect in the Italian population. *Parkinsonism Relat Disord*. **20**(8): p. 911-4 (2014).
78. Trinh, J., I. Guella, and M.J. Farrer, Disease penetrance of late-onset parkinsonism: a meta-analysis. *JAMA*
Neurol. **71**(12): p. 1535-9 (2014).

79. Giannoccaro, M.P., C. La Morgia, G. Rizzo, and V. Carelli, Mitochondrial DNA and primary mitochondrial
dysfunction in Parkinson's disease. *Mov Disord*. **32**(3): p. 346-363 (2017).
80. Larsen, S.B., Z. Hanss, and R. Kruger, The genetic architecture of mitochondrial dysfunction in Parkinson's
disease. *Cell Tissue Res*. **373**(1): p. 21-37 (2018).
81. Williamson, M.G., M. Madureira, W. McGuinness, R. Heon-Roberts, E.D. Mock, K. Naidoo, K.M.L. Cramb,
M.C. Caiazza, A.B. Malpartida, M. Lavelle, K. Savory, S.W. Humble, R. Patterson, J.B. Davis, N. Connor-
Robson, B.J. Ryan, and R. Wade-Martins, Mitochondrial dysfunction and mitophagy defects in LRRK2-
R1441C Parkinson's disease models. *Hum Mol Genet*. **32**(18): p. 2808-2821 (2023).
82. Mortiboys, H., K.K. Johansen, J.O. Aasly, and O. Bandmann, Mitochondrial impairment in patients with
Parkinson disease with the G2019S mutation in LRRK2. *Neurology*. **75**(22): p. 2017-20 (2010).
83. Gonzalez-Hunt, C.P., E.A. Thacker, C.M. Toste, S. Boularand, S. Deprets, L. Dubois, and L.H. Sanders,
Mitochondrial DNA damage as a potential biomarker of LRRK2 kinase activity in LRRK2 Parkinson's
disease. *Scientific Reports*. **10**(1) (2020).
84. Sanders, L.H., J. Laganière, O. Cooper, S.K. Mak, B.J. Vu, Y.A. Huang, D.E. Paschon, M. Vangipuram, R.
Sundararajan, F.D. Urnov, J.W. Langston, P.D. Gregory, H.S. Zhang, J.T. Greenamyre, O. Isacson, and B.
Schüle, LRRK2 mutations cause mitochondrial DNA damage in iPSC-derived neural cells from Parkinson's
disease patients: Reversal by gene correction. *Neurobiology of Disease*. **62**: p. 381-386 (2014).
85. Scharping, N.E., D.B. Rivadeneira, A.V. Menk, P.D.A. Vignali, B.R. Ford, N.L. Rittenhouse, R. Peralta, Y.
Wang, Y. Wang, K. DePeaux, A.C. Poholek, and G.M. Delgoffe, Mitochondrial stress induced by continuous
stimulation under hypoxia rapidly drives T cell exhaustion. *Nat Immunol*. **22**(2): p. 205-215 (2021).
86. Klinge, C.M., Estrogenic control of mitochondrial function. *Redox Biol*. **31**: p. 101435 (2020).
87. Ji, D., J.-Y. Yin, D.-F. Li, C.-T. Zhu, J.-P. Ye, and Y.-Q. Pan, Effects of inflammatory and anti-inflammatory
environments on the macrophage mitochondrial function. *Scientific Reports*. **10**(1) (2020).
88. Wallings, R.L., M.K. Herrick, and M.G. Tansey, LRRK2 at the Interface Between Peripheral and Central
Immune Function in Parkinson's. *Frontiers in Neuroscience*. **14** (2020).
89. Panagiotakopoulou, V., D. Ivanyuk, S. De Cicco, W. Haq, A. Arsic, C. Yu, D. Messelodi, M. Oldrati, D.C.
Schondorf, M.J. Perez, R.P. Cassatella, M. Jakobi, N. Schneiderhan-Marra, T. Gasser, I. Nikic-Spiegel, and
M. Deleidi, Interferon-gamma signaling synergizes with LRRK2 in neurons and microglia derived from
human induced pluripotent stem cells. *Nat Commun*. **11**(1): p. 5163 (2020).
90. Yadavalli, N. and S.M. Ferguson, LRRK2 suppresses lysosome degradative activity in macrophages and
microglia through MiT-TFE transcription factor inhibition. *Proceedings of the National Academy of
Sciences*. **120**(31) (2023).
91. Simpson, C., L. Vinikoor-Imler, F.L. Nassan, J. Shirvan, C. Lally, T. Dam, and N. Maserejian, Prevalence
of ten LRRK2 variants in Parkinson's disease: A comprehensive review. *Parkinsonism Relat Disord*. **98**: p.
103-113 (2022).
92. Tong, Y., A. Pisani, G. Martella, M. Karouani, H. Yamaguchi, E.N. Pothos, and J. Shen, R1441C mutation
in LRRK2 impairs dopaminergic neurotransmission in mice. *Proc Natl Acad Sci U S A*. **106**(34): p. 14622-7
(2009).
1022

Acknowledgments: We thank members of the Tansey lab for useful discussions and edits of the

manuscript. We thank the UF Interdisciplinary Center for Biotechnology Research (UF | ICBR) for use of

flow cytometry and sequencing facilities and advice. Partial funding for this work was derived from awards

from the Bright Focus Fellowship A2021017F (RLW), NIH/NINDS grant RF1NS128800 (MGT), NIH

NIA 1RF1AG057247, and the joint efforts of The Michael J. Fox Foundation for Parkinson's Research

(MJFF) and the Aligning Science Across Parkinson's (ASAP) initiative. MJFF administers the grant ASAP-

020527 on behalf of ASAP and itself. For the purpose of open access, the author has applied a CC-BY

public copyright license to the Author Accepted Manuscript (AAM) version arising from this submission.

N.B. was supported by the DFG (BR4328.2-1, GRK1957), the Michael J Fox Foundation, the Collaborative

Center for X-linked Dystonia-Parkinsonism and the EU Joint Programme - Neurodegenerative Disease

Research (JPND)

1034

1035

Author contributions:

Conceptualization: RLW, MGT

Methodology: RLW, HAS, KMF, SS, NB, SZ, TU, CK, ES

Investigation: RLW, KMF

Visualization: RLW, MGT

Funding acquisition: RLW, MGT

Project administration: MGT

Supervision: MGT

Writing – original draft: RLW, HAS, KMF, SS, NB, SZ, TU, CK, ES, MGT

All authors reviewed and approved the final manuscript.

Competing interests: MGT is a current advisor/consultant for INmune Bio, Merck, Forward Therapeutics,

Weston Foundation, Alzheimer's Association, Bright Focus Foundation, New Horizons Research,

SciNeuro, NysnoBio, Longevity, iMetabolic Pharma, Novo Nordisk, and iMMvention and an AE for

Science Advances. N.B. received honaria from Abbott, Abbvie, Biogen, Biomarin, Bridgebio, Centogene,

Esteve, Ipsen, Merz, Teva, Zambon. CK is medical advisor to Centogene, Retromer Therapeutics, and

Takeda and Speakers Honoraria from Desitin and Bial.

All other authors hold no competing interests.

1053

1054   **Data and materials availability:** All data are available in the main text or supplementary files, including

1055   links to data repositories and individual level data in Data File S1. Correspondence and requests for

1056   materials should be addressed to MGT.

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072

1073

1074

Figures

Figure 1. *R1441C* mutation leads to age dependent biphasic alteration in antigen presentation. pMacs

from 2-3-month and 18-21-month-old female or male, B6 or *R1441C* mice were stimulated with 100U IFNγ

for 18-hours. (A, B) Total Lrrk2 protein expression was assessed and normalized to β-Actin and quantified.

Representative western blots shown. (C, D) T73 Rab10 protein expression was assessed and normalized to

total Rab10 and quantified. Representative western blots shown. (E, F) Surface MHC-II median

fluorescence intensity (MFI) was quantified on LPMs and via flow cytometry. (G-J) pMacs were stained

for intracellular and extracellular MHC-II MFI and Ex:IcMHC-II ratio quantified. Scale bars, 100μM. Bars

represent mean +/- SEM (N = 6-12 mice). Three-way ANOVA, Bonferroni *post hoc*, groups sharing the

same letters are not significantly different (p > 0.05) whilst groups displaying different letters are

significantly different (p < 0.05).

1087

Figure 2. ***R1441C* mutation leads to age dependent biphasic alteration in cytokine release *ex vivo* and**

circulating cytokines *in vivo*. (A) pMacs from 2-3-month and 18-21-month-old female or male, B6 or

R1441C mice were stimulated with 100U IFNγ for 18-hours and media collected. (B, C) IL10, IL6 and

TNF were quantified and normalized to live cell count. (D) Trunk blood was collected via trunk bleed and

plasma was isolated. (E, F) IL10, IL6 and TNF were quantified. Bars represent mean +/- SEM (N = 8-10

mice). Three-way ANOVA, Bonferroni *post hoc*.

1094

Figure 3. *R1441C* mutation leads to age dependent biphasic alteration in lysosomal activity. pMacs from 2-3-month and 18-21-month-old female or male, B6 or *R1441C* mice were stimulated with 100U IFNγ for 18-hours. (A) Schematic of YAe assay. (B, C) Cells were co-treated with 100nM PF360 or 50nM Bafilomycin A1 (baf) and YAe MFI was quantified on LPMs via flow-cytometry. (D, E) DQ-BSA and BMV109 MFI was quantified in LPMs via flow cytometry. (F-I) DQ-BSA and BMV109 MFI was quantified in pMacs via microscopy. Scale bars, 100μM. Bars represent mean +/- SEM (N = 6-12 mice). Two -way ANOVA, Bonferroni *post hoc*, or Student's T-test. Groups sharing the same letters are not significantly different ($p > 0.05$) whilst groups displaying different letters are significantly different ($p < 0.05$).

Figure 4. *R1441C* **Lrrk2 mutation alters pathogen uptake and pathogen-mediated DNA damage in age- and sex-dependent manner**. pMacs from 2-3-month and 18-21-month-old female, B6 or *R1441C* mice were incubated with 100μg of pHrodo *E. coli*. +/- 100nM PF360. (A, B) pMacs were imaged every 20 minutes for 5 hours, and GFP fluorescence was measured. Scale bars, 100μM. Four-way ANOVA, Bonferroni *post hoc*. Main effects of groups are reported in the graph key. Groups sharing the same letters are not significantly different ($p > 0.05$) whilst groups displaying different letters are significantly different ($p < 0.05$). (C) TUNEL MFI was quantified in LPMs via flow cytometry. Bars represent mean +/- SEM (N = 4-12 mice). One-way ANOVA, Bonferroni *post hoc*. Groups sharing the same letters are not significantly different ($p > 0.05$) whilst groups displaying different letters are significantly different ($p < 0.05$).

1123

Figure 5. *R1441C* **Lrrk2 mutation induces immune cell exhaustion in a kinase-dependent manner**.

pMacs from 2-3-month and 18-21-month-old female, B6 or *R1441C* mice were incubated with 100ng/mL

of LPS for 5 days +/- 100nM PF360. (A) Schematic of study design. (B, C) Lrrk2 protein expression was

assessed and normalized to β-Actin and quantified. Representative western blots shown. (D) PDL1+ LPM

count was quantified via flow cytometry. (E) After 5-day LPS or vehicle treatment, pMacs were stimulated

with 100nG/mL of IFNγ for 18-hours and MHC-II MFI expression was quantified via flow cytometry. Bars

represent mean +/- SEM (N = 6-12 mice). Three -way ANOVA, Bonferroni *post hoc*. (F-H) After 5-day

LPS or vehicle treatment, pMacs were incubated with 100μg of pHrodo *E. coli*. +/- 100nM PF360. pMacs

were imaged every 20 minutes for 5 hours, and GFP+ cells (% of total) and GFP MFI was measured. Scale

bars, 100μM. Four-way ANOVA, Bonferroni *post hoc*. Main effects of groups are reported in each graph

key. Groups sharing the same letters are not significantly different (p > 0.05) whilst groups displaying

different letters are significantly different (p < 0.05).

1136

Figure 6. *R1441C* Lrrk2 mutation alters transcriptional regulation of mitochondrial and oxidative stress genes as a function of age in macrophages from mutant females. Transcriptomic analysis from B6 or *R1441C* vehicle and IFNγ-treated pMacs, harvested from 2-3-month or18-21-month-old. (A, B, C). Significant Old:Young DEGs in IFNγ-treated pMacs were counted and compared across B6 and *R1441C*. Volcano plots show genes with log$_2$-transformed fold change > 0.5 and an adjusted p-value ≤ 0.05. (D) KEGG pathway and GO BP enrichment analysis identifies pathways in Old:Young DEGs seen only in *R1441C* samples. (E). Heat maps show KEGG: Parkinson's Disease pathway with significant Old:Young DEGs seen only in *R1441C* samples. (F) KEGG pathway and GO BP enrichment analysis identifies pathways in Aged R1441C F:M upregulated DEGs. pMac-derived RNA was used from 4-6 mice per sex, genotype, treatment and age in transcriptomic studies.

1147

Figure 7. *R1441C* **and** *Y1699C-LRRK2* **monocyte-derived patient macrophages exhibit hypophagocytosis and increased markers for immune cell exhaustion.** (A) Schematic of study design. (B, C) Cryopreserved PBMCs were cryorecovered and plated in the presence of hM-CSF to differentiate myeloid cells to macrophages. Macrophages were incubated with 100μg of pHrodo *E. coli* and imaged every 20 minutes for 5 hours, and GFP+ cells (% of total) quantified. . Scale bars, 100μM. Three-way ANOVA, Bonferroni *post hoc*. Main effects of groups are reported in each graph key. Groups sharing the same letters are not significantly different ($p > 0.05$) whilst groups displaying different letters are significantly different ($p < 0.05$). (D, E, F) PDL1+ cell count and PDL1 expression was quantified in total PBMCs via flow cytometry. Bars represent mean +/- SEM (N = 3-9 participants). One-way ANOVA, Bonferroni *post hoc*. Groups sharing the same letters are not significantly different ($p > 0.05$) whilst groups displaying different letters are significantly different ($p < 0.05$). (G) Monocyte-derived macrophages were stained for PDL1 and microscopy images captured. Scale bars, 100μM.

1166

1167

1168

1169

1170

1171

1172

1173

1174

1175

1176

1177

1178

1179

1180

1181

1182

1183

1184