Assessing the quality of *de novo* parasite genomes assembled using only Oxford Nanopore Technologies MinION data

Kaylee S. Herzog[1], Rachel Wu[1], John M. Hawdon[2], Peter Nejsum[3], and Joseph R. Fauver[1]*

[1]Department of Epidemiology, University of Nebraska Medical Center, Omaha, NE 68198

[2]Department of Microbiology, Immunology, and Tropical Medicine, The George Washington University, Washington, D.C. 20037

[3]Department of Clinical Medicine, Aarhus University, 8200 Aarhus, Denmark

*Lead Contact/Corresponding Author: Joseph R. Fauver, jfauver@unmc.edu

Summary

Parasitic nematode infections represent a significant burden of disease in impoverished populations. Genomic studies of parasitic nematodes have revealed novel drug and vaccine targets and provided unprecedented insights into parasite biology. A key component of these studies is the availability of high-quality reference genomes (i.e., genomes that are contiguous, complete, and accurate) that capture the biological diversity contained within species. However, relatively few genomic resources exist for parasitic nematodes and few species are represented by more than a single reference genome. Streamlined laboratory and computational workflows to generate high-quality reference genomes from individual specimens using a single data source have the potential to increase the availability of genomic data from parasitic nematodes. The Oxford Nanopore Technologies (ONT) MinION is an accessible sequencing platform capable of generating ultra-long read data ideal for assembling genomes. However, lower read-level accuracy of ONT data has previously required assemblies to be error-corrected with more accurate short-read data. In this study, we assessed the quality of *de novo* genome assemblies for three species of parasitic nematodes (*Brugia malayi, Trichuris trichiura,* and *Ancylostoma caninum*) generated using only ONT MinION data. Assemblies were benchmarked against current reference genomes and against additional assemblies that were supplemented with short-read Illumina data through polishing or hybrid assembly approaches. For each species, assemblies generated using only MinION data had similar or superior measures of contiguity, completeness, and gene content. In terms of gene composition, depending on the species, between 88.9–97.6% of complete coding sequences predicted in MinION data only assemblies were identical to those predicted in assemblies polished with Illumina data. Polishing

41 MinION data only assemblies with Illumina data therefore improved gene-level accuracy
42 to a degree. Furthermore, modified DNA extraction and library preparation protocols
43 produced sufficient genomic DNA from *B. malayi* and *T. trichiura* to generate *de novo*
44 assemblies from individual specimens.
45
Data Availability Statement
Quality-controlled MinION and Illumina data for each species are deposited in the
NCBI Sequence Read Archive (SRA) under the BioProject accession ID
PRJNA1074771 under BioSample accession nos. SAMN39888962 (*Brugia malayi*),
SAMN39888963 (*Trichuris trichiura*) and SAMN39888964 (*Ancylostoma caninum*). Final
assemblies for each species are publicly available on GenBank.
52
Introduction
Parasitic nematodes represent an enormous burden of disease. It is estimated
that 1.5 billion people are infected with soil-transmitted helminths, while approximately
120 million and 15 million people suffer from filariasis and onchocerciasis, respectively[1].
Helminths are responsible for nearly 12 million Disability-Adjusted Life Years (DALYs)
lost annually[1]. In addition to the morbidity caused by nematodes themselves, infection
also exacerbates the disease burden of other existing conditions such as malaria, HIV,
and tuberculosis, and can reduce immunological response to important vaccines[2-6].
Reference genomes for parasitic nematodes have proven to be an invaluable resource
for increasing our understanding of helminth biology, treatment, and control. For
example, the landmark comparative genomic study by the International Helminth
Genomes Consortium[7] analyzed dozens of complete genomes representing all major
helminth lineages. This extensive dataset allowed for the identification of multiple gene
family expansions in various groups that are targets for novel drug and vaccine
development. Additional more focused comparative genomic work has revealed
differences between helminth and host metabolic pathways and extracellular vesicle
protein content that represent potential druggable targets, evidence for positive
selection in gene families that are uniquely expanded in flatworms and implicated in the
biology of endoparasitism, conserved immunomodulatory proteins that are potential
vaccine targets for soil-transmitted helminths, and patterns of coevolution between
parasitic roundworms and their mammalian hosts[8-14]. These advances further highlight
helminth reference genomes as indispensable tools for comparative biological and
biomedical research.
Reductions in the cost and improved ease of access of Next-Generation
Sequencing (NGS) has made the assembly of whole genomes from helminths more
feasible[15]. As a result, reference genomes have now been generated for the majority of
species of medical and veterinary importance[16, 17]. Most of these species, however, are
still represented by only a single reference genome. These solitary references are often

generated from laboratory models that have been maintained for decades and therefore
cannot represent the biological diversity observed in natural helminth populations[18, 19].
Generating multiple reference genomes per species from contemporary and
geographically disparate populations would begin to capture the genomic diversity of
helminths, and allow for characterization of differences in, for example, genome
organization, gene copy number and structural variants, and adaptation to local
selective pressures. When generating a genome from a natural population, using
genomic DNA from one specimen is ideal to avoid complications in the assembly
process. Therefore, streamlined laboratory and computational workflows that allow high-
quality assemblies to be generated from an individual specimen using a single data
source would be invaluable. The Oxford Nanopore Technologies (ONT) MinION is a
single molecule sequencing platform capable of generating long reads ideal for *de novo*
genome assembly. Lower read-level accuracy of ONT data has previously required
long-read assemblies to be error-corrected with more accurate short-read data. Recent
updates to ONT chemistries, however, have decreased the error rate of MinION reads,
allowing for the potential to generate helminth reference genomes using only long-read
data.
This study aims to assess the contiguity, completeness, gene content, and gene
composition of whole genome *de novo* assemblies of parasitic nematodes using
MinION data only. Three species that collectively span the breadth of parasitic
nematode diversity were utilized for sequencing: the filarial worm *Brugia malayi*
(Spirurina), the whipworm *Trichuris trichiura* (Trichinellida), and the dog hookworm
Ancylostoma caninum (Rhabditina). Assemblies generated using only ONT MinION
data were compared to reference genomes for each species, as well as to assemblies
supplemented with short-read Illumina data through polishing or hybrid assembly
approaches that were generated as a part of this study. This work highlights a
straightforward approach for generating high-quality *de novo* genome assemblies from
parasitic nematodes using only data generated from the ONT MinION.
109
Results
<u>Data generation</u>
Total gDNA was successfully extracted from a single adult worm for each of *B.*
malayi and *T. trichiura*, and from a pool of L3 larvae for *A. caninum*. For each species,
both MinION and Illumina sequencing libraries were prepared from the same source of
gDNA. The bioinformatic pipeline used to generate each assembliy type is outlined in
Figure 1. Total amounts sequence data generated for each species prior to basecalling
and/or quality control were as follows: ~15.9 Gb (MinION) and ~7.9 Gb (Illumina) for *B.*
malayi; ~26.8 Gb (MinION) and ~4.5 Gb (Illumina) for *T. trichiura*; and ~33.1 Gb
(MinION) and ~20.8 Gb (Illumina) for *A. caninum* (Supplemental Table 1). For all three
species, a median quality score value of >Q16 was achieved for MinION reads, and for

MinION reads aligned to final MinION data only assemblies (Supplemental Figure 1).
Values for average depth of coverage of quality-controlled reads mapped to species-
specific reference assemblies were as follows: 124.85× (MinION) and 82.95× (Illumina)
for *B. malayi*; 249.91× (MinION) and 48.58× (Illumina) for for *T. trichiura*; and 49.42×
(MinION) 38.64× (Illumina) for *A. caninum*. For all species and assembly types,
complete mitochondrial genomes were concurrently generated. For *B. malayi*, the
complete genome for the *Wolbachia* endosymbiont was also assembled in a single
circular genome.
129
Assembly contiguity
All assemblies generated in this study were shorter in terms of total length
compared to current reference genomes, with the exception of the MinION data only
assembly for *B. malayi* (Table 1). Of the assemblies generated in this study, hybrid
assemblies were consistently the most contiguous as represented by higher N50 values
(Table 1). The most improvement in terms of contiguity was observed in *A. caninum*,
where each assembly approach resulted in a shorter genome and more contiguous
assembly compared to the current reference genome. Within a species, all assemblies
had similar levels of GC content.
139
Assembly completeness
miniBUSCO scores were nearly identical among the three assembly types
generated for all three species and matched or exceeded completeness scores of
current reference assemblies (Table 2). For *A. caninum*, miniBUSCO scores indicate
that the assemblies generated here are more complete than the existing International
Helminth Genomes Consortium[7] reference assembly. All assemblies compared for *T.*
trichiura, including both available references, had high proportions of missing BUSCOs
(i.e., ~40.8–42.4%; see Table 2). According to the assessment of contiguity versus
completeness, the assemblies generated here for *B. malayi* and *A. caninum* can be
classified as "tier 1" genomes (Figure 2). For *T. trichiura*, the contiguity of assemblies
generated here is sufficient to qualify them each as "tier 1", but, as mentioned above,
high proportions of BUSCO missingness prevented any trichurid assembly assessed
from achieving "tier 1" status (see Figure 2).
153
Gene content and composition
For organelle genomes, within a species, the mitochondrial and *Wolbachia*
genomes generated were nearly identical (i.e., >99.9% pairwise nucleotide identity)
across assembly types. For gene datasets produced by GeMoMa, the number of
predicted genes was roughly equal across assembly types for each species (Figure 3).
Additionally, the majority of these genes were shared between assembly types. Within a
species, genes predicted across assembly types were similar or identical in mean gene

length, mean length of introns, exons, and coding sequences, and mean number of
exons per coding sequence (Table 3). Pairwise nucleotide comparisons of genes
shared between MinION only data assemblies and MinION assemblies polished with
Illumina data showed 88–98% of these genes to be identical at the nucleotide level
(Figure 4A). The majority of differences in gene composition were the result of single
SNPs or single indels, with few shared genes demonstrating greater than ten
mismatches (Figure 4B).
168
Discussion
<u>Sample processing and sequencing</u>
The DNA extraction and MinION library preparation protocols described here
were optimized for low input gDNA extracted from individual parasitic nematodes,
allowing us to retain the majority of input gDNA through library preparation. Genomic
DNA from a single individual is the ideal input for *de novo* whole genome assembly for
diploid organisms. This allows assembly pipelines to contend with only two potential
haplotypes, leading to more accurate assemblies with less haplotypic duplication. An
alternative method would be to sequence multiple individuals from a highly inbred
homozygous line. Many species cannot be maintained in a laboratory setting, however,
making the establishment of inbred lines for these taxa challenging or impossible[20].
Additionally, sampling from natural, rather than laboratory-maintained, populations is
preferred for many genomic studies, further highlighting the advantage of whole-
genome sequencing from single individuals[21]. For *B. malayi* and *T. trichiura*, which both
have relatively large adult stages (~3–5 cm in length) and tractable genome sizes (<100
Mb), optimized protocols allowed for the generation of both short- and long-read data
from a single adult worm. Adults of *A. caninum*, however, tend not to exceed ~1.5 cm in
length, and have the largest genome size of the three focal species in this study (i.e.,
~400 Mb). Thus, for *A. caninum*, even if adult worms had been accessible, these
protocols would likely still not have allowed for generation of sufficient genomic data
from a single adult. Despite using a pooled sample, however, copy number spectrum
plots suggest that incorporation of a purging step produces results similar to those for
single individuals (see Supplemental Figure 2). For nematodes where gDNA from single
individuals does not meet the requisite quantity for sequencing on the ONT MinION,
additional strategies such as whole genome amplification can be pursued.
There was an association between the average Q score of MinION read datasets
and the total amount of data generated by a flow cell: More efficient MinION sequencing
runs generated data with higher average Q scores (see Supplemental Table 1;
Supplemental Figure 1). For example, MinION libraries for both *B. malayi* and *T.
trichiura* generated ~10.8–11.8 Gb of data per flow cell post-basecalling with
comparably high Q scores. Conversely, libraries for *A. caninum* generated ~7.7–8.5 Gb
of data per flow cell post-basecalling, and these data had the lowest average Q score of

the three species sequenced (see Supplemental Table 1; Supplemental Figure 1). We
observed that flow cells that were used to sequence the *A. caninum* libraries
experienced the most rapid decline in pore availability. This observation highlights the
variability in data generation and quality depending on the sample type when using the
latest ONT chemistries.
206
<u>Estimation of genome size and heterozygosity</u>
There were appreciable differences in overall size of assemblies produced in this
study compared to reference genomes (see Table 1). These differences were notable
for both *T. trichiura* and *A. caninum*, but most dramatic for *A. caninum*, where
assemblies produced here were >100 Mb smaller than the existing reference. This is
likely due to the fact that a purging step to remove haplotypic duplication was included
in the bioinformatic pipeline for all assembly types generated in this study (see Figure
1). This was not the case for the reference genome for *A. caninum*[7], and thus it likely
includes haplotypic duplication that artificially inflates genome size. Three lines of
evidence support that the genomes assembled here more accurately represent the true
haplotypic genome size for both *T. trichiura* and *A. caninum*. First, the lengths of these
assemblies more closely match k-mer based genome size estimates than do those of
the existing reference genomes (see Table 1). Second, miniBUSCO scores were largely
indistinguishable across all assembly types (including existing reference assemblies),
indicating that all are comparably complete despite ranging in size. In fact, the Doyle et
al. reference assembly for *T. trichiura*[22] and the International Helminth Genomes
Consortium reference assembly for *A. caninum*[7] had higher proportions of duplicated
BUSCOs as compared to the assemblies generated herein (see Table 2). Finally, for
hybrid assemblies, copy number spectrum plots suggest they were not "over-purged"
and thus missing genomic content (see Supplemental Figure 2). As previously
advocated by other authors, purging haplotypic duplication is an important step in
genome curation regardless of assembly approach[23, 24]. Unsurprisingly, purging
becomes increasingly important for highly heterozygous sample types, as observed
here for *A. caninum*.
Results from GenomeScope (see Supplemental Table 1) and copy number
spectrum plots (see Supplemental Figure 2) indicate low heterozygosity for *B. malayi*
and elevated heterozygosity for *T. trichiura* and *A. caninum*. These results are
congruent with the fact that the single individual of *B. malayi* sequenced came from a
long-maintained inbred laboratory strain while the individual of *T. trichiura* sequenced
came from a naturally infected human host, and sequence data for *A. caninum* were
generated from a pool of individuals. When sequencing adult females that are
potentially gravid, as was the case for *B. malayi*, analyses like GenomeScope and KAT
that utilize k-mer spectra can be altered by the presence of paternal haplotypes in eggs.
Given that the female of *B. malayi* sequenced came from an inbred strain, however,

maternal and paternal haplotypes are likely to be identical, or highly similar, ameliorating this concern. It is worth noting that the k-mer based approaches used to estimate genome size and heterozygosity and to assess the effectiveness of purging rely on short-read data as input, and therefore cannot be used to evaluate long-read only assemblies at this time.

Assembly contiguity

The reference genome assemblies for both *B. malayi* and *T. trichuris* are highly contiguous and nearing chromosome scale. For *B. malayi* and *T. trichuris*, the N50 values for the assemblies generated in this study were substantially lower compared to those of the highly contiguous reference genome. This is due in part to the additional data sources used to generate these references, which for *B. malayi* included optical mapping[25] and for *T. trichiura* included unspecified updates that resulted in improvements to contiguity[22]. For *A. caninum*, however, the contiguity of the assemblies generated here represent marked improvements as compared to the reference genome. Hybrid assemblies were in all cases more contiguous than MinION data assemblies (see Table 1). This result was somewhat unexpected given that including short-read data in whole genome assembly has traditionally been touted to increase accuracy rather than contiguity[26-28]. Increased contiguity of hybrid assemblies over long read-only assemblies has been observed in bacteria whole genome sequencing[29, 30] but does not appear to have been reported previously for eukaryotes. It is worth noting that the long reads generated for each species here did not approach the read lengths that MinION devices are capable of sequencing. Across species, read length N50 values ranged from ~2.7–8.6 kb (see Supplemental Table 1), which are relatively small. This may be due to the optimized MinION library preparation approach used here, which relied on washing final libraries with a titration of Short and Long Fragment Buffer to retain some short fragments of gDNA in addition to long fragments (see Methods). Rigorous size selection which could have improved read lengths was therefore sacrificed to enable sequencing from low quantities of input gDNA. Had it been possible, generation of ultra-long reads would have likely lessened the gap in contiguity metrics between MinION data and hybrid assemblies. Additionally, a single assembly algorithm was chosen to generate long-read and hybrid assemblies (i.e., Canu and MaSuRCA, respectively). Given the same read-level data, different assembly algorithms may have produced assemblies with different contiguity metrics.

Assembly completeness

In general, completeness values as measured by miniBUSCO are indistinguishable between assembly types for *B. malayi* and *T. trichiura*. Assemblies generated here for *A. caninum* showed an improvement in the number of single copy genes identified and a reduction in the number of duplicated and fragmented genes

identified as compared to the reference genome. Results from miniBUSCO indicate a
high proportion of missingness for all trichurid assemblies (see Table 2; Figure 2).
Underrepresentation in the BUSCO "nematoda_odb10" dataset has been noted to result
in biases and underestimation in ortholog detection for some species of nematodes[31].
This presents a challenge for using traditional completeness metrics like miniBUSCO to
assess trichurid genome assemblies.
287
Gene content and composition
In terms of gene content, within a species, similar numbers of genes were
predicted by GeMoMA for each of the three assembly types generated (see Figure 3).
Though not all predicted genes were identified in all three assemblies, large proportions
were found to be shared among them (i.e., ~93–98%, depending on the species). In
particular, MinION only assemblies and MinION assemblies polished with Illumina data
were highly similar in terms of gene content, with ~98.7–99.7% of predicted genes
shared between them within a species. Furthermore, predicted genes sets had, by
species, identical or nearly identical mean genes lengths, mean intron, exon, and
coding sequence
lengths, and mean numbers of exons per coding sequence (see Table 3), providing
further evidence for a high degree of similarity in gene content across assembly types.
In terms of gene composition, a major goal of this study was to assess the
potential of Illumina data to correct errors that may be present in the final assembly as a
result of less accurate MinION data. To evaluate this, gene datasets predicted by
GeMoMa for MinION data only assemblies and for MinION assemblies polished with
Illumina data were pairwise aligned and assessed for mismatches. Comparisons
between MinION data only assemblies and hybrid assemblies are not tenable because
long-read assemblers like Canu and hybrid assemblers like MaSuRCA utilize assembly
algorithms that handle read-level data dissimilarly[32, 33]. Given that gene predictions for
each assembly were based on existing reference genomes and their corresponding
annotations (see Methods), plus these reference genomes were generated from
different biological samples, comparison to reference genomes was also not salient.
The most direct comparison to determine whether short reads improve gene
accuracy in final assemblies is the MinION data only assembly versus that same
assembly polished with Illumina short reads. If Illumina polishing substantially improves
accuracy, homology-based gene prediction would be expected to result in a low
proportion of genes shared between these two assembly types, and or/the majority of
shared genes to differ at the nucleotide level. This was not the case as all three species
showed high proportions of predicted genes shared between assembly types, and the
majority of the genes shared between the two assembly types were identical (see
Figures 3, 4A). For those that were not identical, most differed by only a single SNP or
indel. These single differences are presumably corrections made during Illumina

polishing (Figure 4B). Despite this, a substantial number of gene comparisons still
contained 10–100+ mismatches (Figure 4B). It is unlikely these differences result from
polishing errors in MinION data only assemblies; rather, they are more likely the result
of homology-based gene prediction models identifying incongruent genes between
assembly types in a small number of cases. In summary, depending on the species,
correcting with short reads did not change nucleotide calls in coding sequences for
~88–98% of the genes compared.
328
Conclusions
For this study, *de novo* whole genome assemblies were generated for three
species of parasitic nematodes (*Brugia malayi*, *Trichuris trichiura*, and *Ancylostoma*
caninum) using only MinION long-read data, using MinION data polished with Illumina
short reads, and using a combined hybrid approach. For *B. malayi* and *T. trichiura*,
optimized gDNA extraction and library preparation protocols allowed for the generation
of complete genomes from individual adult worms. For all species sequenced, MinION
data only assemblies had similar, or superior, measures of contiguity and completeness
as compared to existing reference genomes. The most substantial improvements in
quality metrics were observed for *A. caninum*, which was the only one of the three focal
species for which the existing reference genome is not a near-chromosome scale
assembly. Among the three assembly types generated, predicted gene content was
nearly identical with a species, and the vast majority of predicted genes shared between
MinION data only assemblies and MinION assemblies polished with short reads were
identical at the nucleotide level. For some genes, however, polishing did result in the
correction of single SNPs or indels. Although additional data types beyond MinION long
reads are needed to produce reference-quality, chromosome-scale assemblies, the
results of this study demonstrate that MinION data alone can be used to generate highly
contiguous and complete *de novo* whole genomes from parasitic helminths.
348
Limitations of the Study
This study has multiple limitations. For two of the three focal species, gDNA was
generated from adult worms. For most species, and particularly those that infect
humans, adult nematodes are difficult or impossible to obtain, somewhat limiting the
utility of this approach in practice. Furthermore, the assemblies generated here remain
incomplete in terms of the modern reference genome standards. They are not
chromosome-scale, they are not annotated, and additional data types are required to
complete them. Generating chromosome-scale annotated assemblies was not the goal
of this study. Rather, it was to evaluate the feasibility and quality of whole genome
assemblies generated from an accessible NGS data type. For *B. malayi* and *T. trichiura*,
existing reference genomes were already exceptionally high-quality "tier 1"-status
assemblies *sensu* the International Helminth Genomes Consortium[7], and the

assemblies generated here using a single data source approached their quality in contiguity and matched their quality in completeness. The comparison in which the added value of long-read MinION data is most evident is that of *A. caninum*. Using gDNA from the same strain and sample type on which the existing reference assembly was based, the genomic resources available for this species were improved considerably.

Author Contributions

KSH and JRF designed the study. JMH provided the specimens of *Ancylostoma caninum* sequenced. PN provided the specimen of *Trichuris trichiura* sequenced. KSH extracted genomic DNA and prepared and sequenced MinION libraries. KSH and JRF determined and refined bioinformatic pipelines. KSH performed bioinformatic analyses. RW aided KSH in gene-level comparison analyses. KSH and JRF interpreted data and generated figures and tables. KSH and JRF drafted the manuscript. All authors contributed to revising and approving the final version of the manuscript.

Acknowledgements

The authors thank the BEI Resources NIH/NIAID Filariasis Research Reagent Resource Center for providing specimens of *Brugia malayi* for sequencing. Thanks are also due to Dr. Michael Wiley, Dr. Shaun Cross, Kristen Bernhard, and Mahmood Al-Haehm (University of Nebraska Medical Center) for providing access and support for use of select laboratory equipment. The authors additionally thank Christopher Castaldi and Irina Tikhonova (Yale Center for Genome Analysis, Yale School of Medicine) for generation of, and correspondence regarding, Illumina sequence data. Bioinformatic analysis was enabled by the University of Nebraska-Lincoln Holland Computing Center computing cluster.

Declaration of Interests

The authors declare no competing interests.

Figure Titles and Legends

Figure 1. Bioinformatic pipelines used to generate *de novo* whole genome assemblies. The MinION data only pipeline is provided at left and the hybrid pipeline is provided at right.

Figure 2. Plot of single copy miniBUSCO score versus the log value of N50 length divided by contig (or scaffold) count for the assemblies generated as part of this study, and for all genomes of nematodes that parasitize animals as adults available from WormBase ParaSite. Dotted lines indicate the cutoff for "tier 1" genome

status *sensu* the International Helminth Genomes Consortium (i.e., >85% single copy BUSCO score and >1.6 log value for the contiguity metric). A key to symbolic designations is provided at top left.

Figure 3. Venn diagrams of the number of genes predicted by GeMoMa that were shared among the assembly types generated. (A) *Brugia malayi*, (B) *Trichuris trichiura* and (C) *Ancylostoma caninum*.

Figure 4. Summary of differences in gene composition identified in predicted genes shared between MinION data assemblies and MinION assemblies polished with Illumina data. (A) Proportions of shared genes identified at the nucleotide level as identical, differing by SNPs only, differing by indels only, or differing by both SNPS and indels for *Brugia malayi* (left), *Trichuris trichiura* (center), and *Ancylostoma caninum* (right). **(B)** Number of differences identified between non-identical shared genes for *Brugia malayi* (left), *Trichuris trichiura* (center), and *Ancylostoma caninum* (right). A key to bar plot colors is provided at bottom center.

Materials and Methods

Specimen acquisition

The specimens sequenced were obtained from the same strains previously used to generate existing reference genomes for each species. Adults of *B. malayi* (FR3 strain) were acquired through the NIH/NIAID Filariasis Research Reagent Resource Center via BEI Resources (Michalski et al., 2011). Worms were received frozen and stored at -80°C. Prior to sequencing, worms were thawed at 4°C and subsequently preserved in DNA/RNA Shield (Zymo Research, Irvine, CA, USA). Adults of *T. trichiura* were obtained from a Danish patient after anthelmintic treatment for an infection acquired in Uganda. Worms were received preserved in 70% ethanol and were stored at 4°C prior to sequencing. Pooled third-stage (L3) larvae of *A. caninum* (Baltimore strain) were obtained from experimental infection in laboratory-maintained canines. Pooled L3s were received frozen, stored at -80°C, and thawed at 4°C prior to sequencing.

DNA extraction

Total nucleic acid was extracted separately from each of a single adult female (*B. malayi*), a single adult male (*T. trichuris*), and a pool of L3 larvae (*A. caninum*) using a *Quick*-DNAÔ HMW MagBead Kit (Zymo Research, Irvine, CA, USA) and modified Solid Tissue extraction protocol (Supplemental Table 1). Prior to extraction, adult worms were placed in clean 1.5 mL DNA LoBind® tubes (Eppendorf® North America, Enfield, CT, USA) using sterilized forceps, while pooled L3s were centrifuged at 6,540 *g* for 3 min to allow for the removal of excess supernatant. All samples were then homogenized

mechanically using a Monarch® Pestle Set single-use microtube pestle (New England
Biolabs® Inc., Ipswich, MA, USA). Homogenization was performed both prior to, and
immediately following, the addition of Proteinase K. Samples were then digested in a
digital dry bath at 55°C for 24 hr with occasional flick mixing. The protocol for DNA
purification was then followed, with these modifications: After the addition of 33 µL of
MagBinding Beads, samples were placed on a benchtop rotating mixer for 40 min to 1.5
hr at RT; and after the addition of 52 µL DNA Elution Buffer, samples were incubated in
a digital dry bath at 37°C for 2 hr with occasional flick mixing, then incubated at RT
overnight prior to collecting eluted DNA. To avoid DNA fragmentation, samples were
mixed by flicking rather than pipette mixing wherever possible. The concentration of
each extraction was measured using a Qubit4Ô 1X dsDNA High Sensitivity (HS) Assay
(ThermoFisher Scientific, Waltham, MA, USA) and fragment size distribution was
determined using an Agilent 2200 TapeStation System and associated protocol for
genomic DNA ScreenTape analysis (Agilent, Santa Clara, CA, USA). For *T. trichiura*,
TapeStation analysis was not performed.

456   For *A. caninum*, initial attempts to sequence ONT libraries resulted in low
sequencing efficiency, poor data quality, and rapid pore loss. To ameliorate concerns of
protein contamination, extracted genomic DNA (gDNA) was purified via bead-based
cleanup prior to final ONT library preparation. This cleanup was performed as follows:
Two aliquots of extracted gDNA were each bound to a 0.5✗ volume of AMPure XP
beads (Beckman Coulter, Brea, CA, USA) on a benchtop rotator mixer for 1 hr at RT.
Beads were then washed twice with 80% EtOH, air-dried, and gDNA was eluted in 52
µL Zymo DNA Elution Buffer for 2 hr at 37°C with occasional flick mixing, followed by
overnight elution at RT. This cleanup process was repeated a second time for both
aliquots. Purified extractions were then quantified using a Qubit4[TM] 1X dsDNA High HS
Assay and the Agilent 2200 TapeStation System.

MinION library preparation

469   Aliquots of gDNA extracted for each species were used to prepare one or more
libraries for whole genome sequencing on an ONT MinION desktop sequencer (see
Supplemental Table 1). Libraries were prepared using a combination of the DNA Clean
& Concentrator MagBead (Zymo Research, Irvine, CA, USA) and ONT SQK-LSK114
Ligation Sequencing (Oxford Nanopore Technologies, Oxford, United Kingdom) Kits,
and a modified hybrid protocol. First, gDNA was incubated with ONT DNA repair and
end preparation reagents for 10 min at RT followed by 10 min at 65°C. End preparation
reactions were then bound to 20 µL Zymo MagBinding Beads in 4✗ volumes of Zymo
DNA MagBinding Buffer on a benchtop rotating mixer for 30 min to 2 hr. Beads were
then washed twice with Zymo DNA Wash Buffer, air-dried for 10 min, eluted in 51 µL
Zymo DNA Elution Buffer via manual agitation for 10 min, and quantified via Qubit 1X
dsDNA HS Assay. Sequencing adaptors were then ligated via a 15 min incubation at

RT, and libraries were bound to a 0.4× volumes of AMPure XP beads on a benchtop
rotator mixer at RT for 1 hr. Beads were then washed twice with either ONT Long
Fragment Buffer (*A. caninum*) or a titrated wash mix of 1:3 ONT Short Fragment
Buffer:ONT Long Fragment Buffer (*B. malayi* and *T. trichiura*). After washing, beads
were air-dried, then libraries were eluted in 15–17 µL of ONT Elution Buffer for 2 hr at
37°C with occasional flick mixing followed by overnight elution at RT. Final libraries
were quantified using a Qubit4Ô 1X dsDNA High HS Assay and the Agilent 2200
TapeStation System. For *T. trichiura*, TapeStation analysis was not performed.
Additionally, for *T. trichiura*, a portion of ONT library reserved after sequencing on one
flow cell was re-washed prior to sequencing on a second flow cell (see Supplemental
Table 1). This aliquot of library was bound to a 0.4× volume of AMPure XP beads on a
benchtop rotator mixer for 1 hr, washed twice with ONT Long Fragment Buffer, air-dried,
then eluted in 17 µL of ONT Elution Buffer for 2 hr at 37°C with occasional flick mixing
followed by overnight elution at RT.
495
MinION sequencing, basecalling, and read data quality control
Portions of ONT libraries were sequenced for ~62–80 hr each on one (*B. malayi*),
two (*T. trichiura*), or three (*A. caninum*) ONT MinION R10.4.1 flow cells. The amount of
library loaded onto each flow cell and the total number of pores available at the start of
sequencing are provided in Supplemental Table 1. MinKNOW software (ONT) v.
22.10.7 (*B. malayi*), v. 22.10.10 (*T. trichiura*), or v. 22.12.7 (*A. caninum*) was used to run
each flow cell with pore scans every 1.5 hr. After sequencing, signal data (i.e., fast5
files) from each flow cell were basecalled using Guppy v. 6.3.4 (ONT). The bioinformatic
pipeline used to generate MinION data assemblies is outlined in Figure 1. Sequencing
adaptors were simultaneously removed by specifying the "⬜⬜trim_adapters" flag. The
results of each Guppy run were summarized using NanoPlot v. 1.40.2[34]. Fastq files that
passed basecalling were input to Mash v. 2.2.2[35] to confirm there was no significant
contamination in the read-level data prior to assembly. Estimated genome sizes
provided to Mash were based on the lengths of the existing reference assemblies
available for each species, and were thus set as 88 Mb for *B. malayi*[36], 465 Mb for *A.*
caninum[7]   and   75   Mb   for   *T.   trichiura*[22]
(https://github.com/stephenrdoyle/ancient_trichuris/tree/master/02_data). Average depth
of coverage of reads across the species-specific reference genome and proportion of
reads mapped were estimated using Minimap2 v. 2.16[37] and SAMtools v. 1.9[38].
515
Illumina library preparation, sequencing, and read data quality control
Aliquots of gDNA extracted for each species were sent to the Yale Center for
Genome Analysis (YCGA) for Illumina whole genome library preparation and
sequencing (see Supplemental Table 1). Libraries were prepared using an xGenTM
cfDNA & FFPE DNA Library Prep Kit with unique dual indexing (Integrated DNA

Technologies, Newark, NJ, USA) and standard protocol. Libraries were subjected to 5–
7 cycles of PCR to increase concentration (see Supplemental Table 1) and multiplexed
for 2×150 paired-end sequencing on an Illumina NovaSeq 6000 targeting 50× depth of
coverage genome-wide for each species. Reads were demultiplexed by the YCGA and
subsequently filtered for remaining adaptor contamination, quality, and length using
fastp[39]. The bioinformatic pipeline used to process Illumina data and generate hybrid
assemblies is outlined in Figure 1. Fastq files that passed fastp were input to Mash to
confirm there was no significant contamination in the read-level data prior to assembly.
Estimated genome sizes provided to Mash were based on the lengths of the existing
reference assemblies available for each species (see above). Average depth of
coverage across the species-specific reference genome and proportion of reads
mapped were estimated using BWA v. 0.7.17 and SAMtools v. 1.9[38]. A second round of
fastp was run on the raw demultiplexed Illumina data to remove sequencing adapters,
only, by specifying "￭￭detect_adapter_for_pe", "￭￭disable_length_filtering", and
"￭￭disable_quality_filtering". These data were the used for hybrid genome assembly, as
MaSuRCA utilizes built-in error correction and cleaning.
537
<u>Estimation of genome size and heterozygosity</u>
Quality-controlled Ilumina data were used to estimate genome size and
heterozygosity for each species. First, Jellyfish v. 2.3.0[40] was used to generate k-mer
spectra, specifying a k-mer size of 21 and an initial hash size of 1,000,000,000.
Histograms from Jellyfish were then provided to the GenomeScope web-based
graphical user interface[41] to visualize k-mer spectra and estimate heterozygosity.
544
<u>MinION data genome assembly</u>
FASTQ files that passed basecalling by Guppy were concatenated and used as
input to Canu v. 2.1[32] (ONT) to generate MinION data assemblies. For *T. trichiura* and
A. caninum, a 3 kb minimum read length requirement was specified, and for *B. malayi*, a
5 kb minimum read length requirement was specified. Estimated genome sizes provided
to Canu were based on the lengths of the existing reference assemblies available for
each species (see above). Contigs in the resulting Canu assemblies that were indicated
as potential alternative alleles (i.e., with FASTA headers including
"suggestBubble=yes") were removed using a combination of SeqKit v 0.10.1[42] and the
filterbyname.sh script of BBMap v.38.84[43]. Resulting "popped" assemblies were used as
input to purge_dups v. 1.2.5[44] to remove false duplications. These "popped and purged"
assemblies were then polished with one round of Racon v. 1.5.0[45] followed by one
round of Medaka v. 1.7.2 (https://github.com/nanoporetech/medaka) to generate
"popped, purged, and polished" assemblies. Copies of these final assemblies were also
separately polished with Illumina data using Pilon v. 1.24[46] and BWA. Three iterations of
Pilon were iteratively run specifying the "￭￭diploid" flag.

561
<u>Hybrid genome assembly</u>
FASTQ files that passes basecalling by Guppy and Illumina data from which only
sequencing adaptors were removed by fastp were used as input to MaSuRCA v. 4.1.0
(Zimin et al., 2013) to generate hybrid assemblies. Default settings were used except for
setting "LHE_COVERAGE" to "35" and "cgwErrorRate=0.15", and setting "JF_SIZE" to a
value ten times the genome size of the species-specific reference assembly.
Assemblies output by MaSuRCA were input to purge_dups v. 1.2.5[44] to remove false
duplications.
570
<u>Quality control and refinement of final assemblies</u>
To ensure "purged" MaSuRCA hybrid assemblies were not over- or under-
purged, copy number spectrum plots were generated using the K-mer Analysis Toolkit
(KAT) v. 2.4.0[47] specifying a k-mer size of 31. "Popped, purged, and polished" Canu
MinION data assemblies and "purged" MaSuRCA hybrid assemblies were input to
BlobTools v. 1.1.1[48] to identify potential contaminants. First, the reads used to generate
each assembly were mapped to the assembly using Minimap2 (MinION data) and/or
BWA (Illumina data), and SAMtools was used to generate sorted BAM files for each
mapping. The "blastn" function in BLAST v. 2.14.0[49] was then used to compare all
contigs to reference sequences available in the National Center for Biotechnology
Information (NCBI)'s nucleotide databases, specifying "evalue 1e-25",
"max_target_seqs 10" and "max_hsps 1". BLAST outputs and sorted BAM files were
then used as input to BlobTools. Contigs that could be confirmed as host, human, or
bacterial contamination were excluded from final assemblies. Quality scores (i.e., Q
scores) of basecalled MinION reads, and Q scores of reads when mapped to the final
"popped, purged, and polished" MinION data assemblies, were measured using
NanoPlot and plotted using the "geom_density" function in the ggplot2 package[50] in R v.
4.3.0[51] in RStudio v. 2023.03.1[52].
589
<u>Mitogenome and *Wolbachia* genome assembly</u>
For *B. malayi* and *A. caninum* contigs containing the mitogenomes—and, for *B.*
malayi, the genome of its *Wolbachia* endosymbiont—were extracted from "popped"
MinION and unpurged MaSuRCA assemblies. For *T. trichiura*, mitogenomes were
extracted from the original "unpopped" Canu assembly and the unpurged MaSuRCA
assembly. These organelle genomes contigs were then mapped to species-specific
reference organelle genomes using Geneious Prime v. 2022.0.1[53] and manually
reconfigured to match the linear orientation of the reference. For *B. malayi* and *T.*
trichiura, reference organelle genomes were extracted from the reference genome
assemblies for each species (see above). For *A. caninum*, for which a mitochondrial
contig is not labeled in the reference assembly, a reference mitogenome was

downloaded from GenBank (accession no. MN215971)[54]. Contigs containing complete
organelle genomes were then manually added to the appropriate final version of each
assembly.
604
<u>Assessing assembly contiguity and completeness</u>
Contiguity and completeness were assessed for each assembly using QUAST v.
5.0.2[55] and miniBUSCO v. 0.2[56], respectively. For miniBUSCO, the Nematoda ortholog
database (nematoda_odb10) was specified. For comparison, miniBUSCO was also run
on the aforementioned reference genomes for each species, as well as for all genomes
of nematodes that parasitize animals as adults available from WormBase ParaSite[16, 17].
For each assembly, the proportion of single copy BUSCOs present was plotted against
the log value of the assembly's N50 length divided by its scaffold or contig count using
R v. 4.2.3 via RStudio v. 2023.06.1. Cutoff values of >85% (for proportion of single copy
BUSCOs present) and >1.6 (for the log value of the contiguity metric) were used to
assess whether each assembly could be classified as a "tier 1" genome following the
standard for helminth genomes established by the International Helminth Genomes
Consortium[7].
618
<u>Assessing gene content and composition</u>
Homology-based gene prediction was performed for each assembly generated
here using GeMoMA[57, 58]. Predictions were made using default settings and the
species-specific reference genome and annotation as input. Resulting GFF files were
summarized using AGAT[59]. For each species, the predicted genes shared between the
MinION data assembly and the MinION assembly polished with Illumina data were
pairwise aligned using the "pairwiseAlignment" function in Biostrings package v. 2.66.0[60]
in R v. 4.2.3 via RStudio v. 2023.06.1 specifying the alignment type as "global".
Biostrings was then used to characterize each pairwise alignment in terms of percent
identity, alignment length, gene lengths, and number of single nucleotide
polymorphisms (i.e., SNPS) and insertions/deletions (i.e., indels). Summary figures for
all comparisons were generated in R v. 4.3.0 via RStudio v. 2023.03.1 using the ggplot2
package.
632
633 **Supplemental Figure Titles and Legends**
634
635 **Supplemental Figure 1.** Histograms of average read quality scores (i.e., Q scores) for
636 unaligned basecalled MinION read data, and for basecalled MinION read data aligned
637 to the final MinION data only genome assembly. (A) *Brugia malayi*, (B) *Trichuris*
638 *trichiura*, and (C) *Ancylostoma caninum*.
639

Supplemental Figure 2. K-mer multiplicity plots generated from purged hybrid assemblies. (A) *Brugia malayi*, (B) *Trichuris trichiura*, and (C) *Anyclostoma caninum*.

References

1. 2020. The Global Burden of Disease Study 2019. The Lancet 396(10258): 1129–1306.

2. Biraro, I.A., M. Egesa, F. Toulza, J. Levin, S. Cose, M. Joloba, S. Smith, H.M. Dockrell, A. Katamba, and A.M. Elliott. 2014. Impact of co-infections and BCG immunisation on immune responses among household contacts of tuberculosis patients in a Ugandan cohort. PLoS One 9(11): e111517. DOI: 10.1371/journal.pone.0111517.

3. Kizito, D., R. Tweyongyere, A. Namatovu, E.L. Webb, L. Muhangi, S.A. Lule, H. Bukenya, S. Cose, and A.M. Elliott. 2013. Factors affecting the infant antibody response to measles immunisation in Entebbe-Uganda. BMC Public Health 13(619): 1–9. DOI: 10.1186/1471-2458-13-619.

4. Morawski, B.M., M. Yunus, E. Kerukadho, G. Turyasingura, L. Barbra, A.M. Ojok, A.R. DiNardo, S. Sowinski, D.R. Boulware, and R. Mejia. 2017. Hookworm infection is associated with decreased CD4+ T cell counts in HIV-infected adult Ugandans. PLoS Neglected Tropical Diseases 11(5): e0005634. DOI: 10.1371/journal.pntd.0005634.

5. Nash, S., A.J. Mentzer, S.A. Lule, D. Kizito, G. Smits, F.R. van der Klis, and A.M. Elliott. 2017. The impact of prenatal exposure to parasitic infections and to anthelminthic treatment on antibody responses to routine immunisations given in infancy: Secondary analysis of a randomised controlled trial. PLoS Neglected Tropical Diseases 11(2): e0005213. DOI: 10.1371/journal.pntd.0005213.

6. Ndyomugyenyi, R., N. Kabatereine, A. Olsen, and P. Magnussen. 2008. Malaria and hookworm infections in relation to haemoglobin and serum ferritin levels in pregnancy in Masindi district, western Uganda. Transactions of the Royal Society of Tropical Medicine and Hygiene 102(2): 130–136. DOI: 10.1016/j.trstmh.2007.09.015.

7. Consortium, I.H.G. 2019. Comparative genomics of the major parasitic worms. Nature Genetics 51(1): 163–174. DOI: 10.1038/s41588-018-0262-1.

8. Bennett, A.P., E. de la Torre-Escudero, and M.W. Robinson. 2020. Helminth genome analysis reveals conservation of extracellular vesicle biogenesis pathways but divergence of RNA loading machinery between phyla. International Journal for Parasitology 50(9): 655–661. DOI: 10.1016/j.ijpara.2020.04.004.

9. Collington, E., B. Lobb, N.A. Mazen, A.C. Doxey, and D.M. Glerum. 2023. Phylogenomic analysis of 155 helminth species reveals widespread absence of oxygen metabolic Capacity. Genome Biology and Evolution 15(8): evad135. DOI: 10.1093/gbe/evad135.

10. Hu, Y., L. Yu, H. Fan, G. Huang, Q. Wu, Y. Nie, S. Liu, L. Yan, and F. Wei. 2021. Genomic signatures of coevolution between nonmodel mammals and parasitic roundworms. Molecular Biology and Evolution 38(2): 531–544. DOI: 10.1093/molbev/msaa243.

684   11.   Luo, G., R. Gong, P. Li, Q. Li, and X. Wei. 2022. Comparative genomic analysis
685   of *Echinococcus multilocularis* with other tapeworms. Biologia 77(9): 2743-2750.
686   DOI: 10.1007/s11756-022-01120-9.

687   12.   Montaño, K.J., C. Cuéllar, and J. Sotillo. 2021. Rodent models for the study of
688   soil-transmitted helminths: A proteomics approach. Frontiers in Cellular and
689   Infection Microbiology 11: 639573. DOI: 10.3389/fcimb.2021.639573.

690   13.   Rosa, B.A., Y.-J. Choi, S.N. McNulty, H. Jung, J. Martin, T. Agatsuma, H.
691   Sugiyama, T.H. Le, P.N. Doanh, and W. Maleewong. 2020. Comparative
692   genomics and transcriptomics of 4 *Paragonimus* species provide insights into
693   lung fluke parasitism and pathogenesis. GigaScience 9(7): giaa073. DOI:
694   10.1093/gigascience/giaa073.

695   14.   Wang, J. 2021. Genomics of the parasitic nematode *Ascaris* and its relatives.
696   Genes 12(4). DOI: 10.3390/genes12040493.

697   15.   Doyle, S.R. 2022. Improving helminth genome resources in the post-genomic
698   era. Trends in Parasitology 38(10): 831–840. DOI: 10.1016/j.pt.2022.06.002.

699   16.   Howe, K.L., B.J. Bolt, S. Cain, J. Chan, W.J. Chen, P. Davis, J. Done, T. Down,
700   S. Gao, and C. Grove. 2016. WormBase 2016: expanding to enable helminth
701   genomic research. Nucleic Acids Research 44(D1): D774–D780. DOI:
702   10.1093/nar/gkv1217.

703   17.   Howe, K.L., B.J. Bolt, M. Shafie, P. Kersey, and M. Berriman. 2017. WormBase
704   ParaSite– a comprehensive resource for helminth genomics. Molecular and
705   Biochemical Parasitology 215: 2–10. DOI: 10.1016/j.molbiopara.2016.11.005.

706   18.   Valiente-Mullor, C., B. Beamud, I. Ansari, C. Francés-Cuesta, N. García-
707   González, L. Mejía, P. Ruiz-Hueso, and F. González-Candelas. 2021. One is not
708   enough: on the effects of reference genome for the mapping and subsequent
709   analyses of short-reads. PLoS Computational Biology 17(1): e1008678. DOI:
710   10.1371/journal.pcbi.1008678.

711   19.   Yang, X., W.-P. Lee, K. Ye, and C. Lee. 2019. One reference genome is not
712   enough. Genome Biology 20(104): 1–3. DOI: 10.1186/s13059-019-1717-0.

713   20.   Solares, E.A., Y. Tao, A.D. Long, and B.S. Gaut. 2021. HapSolo: an optimization
714   approach for removing secondary haplotigs during diploid genome assembly and
715   scaffolding. BMC Bioinformatics 22(1): 1–17. DOI: 10.1186/s12859-020-03939-y.

716   21.   Adams, M., J. McBroome, N. Maurer, E. Pepper-Tunick, N.F. Saremi, R.E.
717   Green, C. Vollmers, and R.B. Corbett-Detig. 2020. One fly–one genome:
718   chromosome-scale genome assembly of a single outbred *Drosophila*
719   *melanogaster.* Nucleic Acids Research 48(13): e75. DOI: 10.1093/nar/gkaa450.

720   22.   Doyle, S.R., M.J. Søe, P. Nejsum, M. Betson, P.J. Cooper, L. Peng, X.-Q. Zhu,
721   A. Sanchez, G. Matamoros, and G.A.F. Sandoval. 2022. Population genomics of
722   ancient and modern *Trichuris trichiura.* Nature Communications 13(1): 3888.
723   DOI: 10.1038/s41467-022-31487-x.

724   23.   Howe, K., W. Chow, J. Collins, S. Pelan, D.-L. Pointon, Y. Sims, J. Torrance, A.
725   Tracey, and J. Wood. 2021. Significantly improving the quality of genome
726   assemblies through curation. Gigascience 10(1): giaa153. DOI:
727   10.1093/gigascience/giaa153.

728   24.   Rhie, A., S.A. McCarthy, O. Fedrigo, J. Damas, G. Formenti, S. Koren, M.
729   Uliano-Silva, W. Chow, A. Fungtammasan, and J. Kim. 2021. Towards complete

and error-free genome assemblies of all vertebrate species. Nature 592(7856):
737–746. DOI: 10.1038/s41586-021-03451-0.

25. Tracey, A., J.M. Foster, M. Paulini, A. Grote, J. Mattick, Y.-C. Tsai, M. Chung,
J.A. Cotton, T.A. Clark, and A. Geber. 2020. Nearly complete genome sequence
of *Brugia malayi* strain FR3. Microbiology Resource Announcements 9(24). DOI:
10.1128/mra. 00154-20.

26. Bashir, A., A.A. Klammer, W.P. Robins, C.-S. Chin, D. Webster, E. Paxinos, D.
Hsu, M. Ashby, S. Wang, and P. Peluso. 2012. A hybrid approach for the
automated finishing of bacterial genomes. Nature Biotechnology 30(7): 701–707.
DOI: 10.1038/nbt.2288.

27. English, A.C., S. Richards, Y. Han, M. Wang, V. Vee, J. Qu, X. Qin, D.M. Muzny,
J.G. Reid, and K.C. Worley. 2012. Mind the gap: upgrading genomes with Pacific
Biosciences RS long-read sequencing technology. PloS One 7(11): e47768. DOI:
10.1371/journal.pone.0047768.

28. Koren, S., M.C. Schatz, B.P. Walenz, J. Martin, J.T. Howard, G. Ganapathy, Z.
Wang, D.A. Rasko, W.R. McCombie, and E.D. Jarvis. 2012. Hybrid error
correction and *de novo* assembly of single-molecule sequencing reads. Nature
Biotechnology 30(7): 693–700. DOI: 10.1038/nbt.2280.

29. George, S., L. Pankhurst, A. Hubbard, A. Votintseva, N. Stoesser, A.E.
Sheppard, A. Mathers, R. Norris, I. Navickaite, and C. Eaton. 2017. Resolving
plasmid structures in Enterobacteriaceae using the MinION nanopore sequencer:
assessment of MinION and MinION/Illumina hybrid data assembly approaches.
Microbial Genomics 3(8): e000118. DOI: 10.1099/mgen.0.000118.

30. Neal-McKinney, J.M., K.C. Liu, C.M. Lock, W.-H. Wu, and J. Hu. 2021.
Comparison of MiSeq, MinION, and hybrid genome sequencing for analysis of
Campylobacter jejuni. Scientific Reports 11(1): 5676. DOI: 10.1038/s41598-021-
84956-6.

31. Rödelsperger, C. 2021. The community-curated *Pristionchus pacificus* genome
facilitates automated gene annotation improvement in related nematodes. BMC
Genomics 22(216). DOI: 10.1186/s12864-021-07529-x.

32. Koren, S., B.P. Walenz, K. Berlin, J.R. Miller, N.H. Bergman, and A.M. Phillippy.
2017. Canu: scalable and accurate long-read assembly via adaptive k-mer
weighting and repeat separation. Genome Research 27(5): 722–736. DOI:
10.1101/gr.215087.116.

33. Zimin, A.V., G. Marçais, D. Puiu, M. Roberts, S.L. Salzberg, and J.A. Yorke.
2013. The MaSuRCA genome assembler. Bioinformatics 29(21): 2669–2677.
DOI: 10.1093/bioinformatics/btt476.

34. De Coster, W. and R. Rademakers. 2023. NanoPack2: population-scale
evaluation of long-read sequencing data. Bioinformatics 39(5): btad311. DOI:
10.1093/bioinformatics/btad311.

35. Ondov, B.D., T.J. Treangen, P. Melsted, A.B. Mallonee, N.H. Bergman, S. Koren,
and A.M. Phillippy. 2016. Mash: fast genome and metagenome distance
estimation using MinHash. Genome Biology 17(132): 1–14. DOI:
10.1186/s13059-016-0997-x.

36. Ghedin, E., S. Wang, D. Spiro, E. Caler, Q. Zhao, J. Crabtree, J.E. Allen, A.L.
Delcher, D.B. Guiliano, and D. Miranda-Saavedra. 2007. Draft genome of the

filarial nematode parasite *Brugia malayi.* Science 317(5845): 1756–1760. DOI:
10.1126/science.1145406.

37. Li, H. 2018. Minimap2: pairwise alignment for nucleotide sequences.
Bioinformatics 34(18): 3094–3100. DOI: 10.1093/bioinformatics/bty191.

38. Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G.
Abecasis, and R. Durbin. 2009. The sequence alignment/map format and
SAMtools. Bioinformatics 25(16): 2078–2079. DOI:
10.1093/bioinformatics/btp352.

39. Chen, S., Y. Zhou, Y. Chen, and J. Gu. 2018. fastp: an ultra-fast all-in-one
FASTQ preprocessor. Bioinformatics 34(17): i884–i890. DOI:
10.1093/bioinformatics/bty560.

40. Marçais, G. and C. Kingsford. 2011. A fast, lock-free approach for efficient
parallel counting of occurrences of k-mers. Bioinformatics 27(6): 764–770. DOI:
10.1093/bioinformatics/btr011.

41. Vurture, G.W., F.J. Sedlazeck, M. Nattestad, C.J. Underwood, H. Fang, J.
Gurtowski, and M.C. Schatz. 2017. GenomeScope: fast reference-free genome
profiling from short reads. Bioinformatics 33(14): 2202–2204. DOI:
10.1093/bioinformatics/btx153.

42. Shen, W., S. Le, Y. Li, and F. Hu. 2016. SeqKit: a cross-platform and ultrafast
toolkit for FASTA/Q file manipulation. PLoS One 11(10): e0163962. DOI:
10.1371/journal.pone.0163962.

43. Bushnell, B., *BBMap: a fast, accurate, splice-aware aligner*. 2014, Lawrence
Berkeley National Lab.(LBNL), Berkeley, CA (United States).

44. Guan, D., S.A. McCarthy, J. Wood, K. Howe, Y. Wang, and R. Durbin. 2020.
Identifying and removing haplotypic duplication in primary genome assemblies.
Bioinformatics 36(9): 2896–2898. DOI: 10.1093/bioinformatics/btaa025.

45. Vaser, R., I. Sović, N. Nagarajan, and M. Šikić. 2017. Fast and accurate de novo
genome assembly from long uncorrected reads. Genome Research 27(5): 737–
746. DOI: 10.1101/gr.214270.116.

46. Walker, B.J., T. Abeel, T. Shea, M. Priest, A. Abouelliel, S. Sakthikumar, C.A.
Cuomo, Q. Zeng, J. Wortman, and S.K. Young. 2014. Pilon: an integrated tool for
comprehensive microbial variant detection and genome assembly improvement.
PLoS one 9(11): e112963. DOI: 10.1371/journal.pone.0112963.

47. Mapleson, D., G. Garcia Accinelli, G. Kettleborough, J. Wright, and B.J. Clavijo.
2017. KAT: a K-mer analysis toolkit to quality control NGS datasets and genome
assemblies. Bioinformatics 33(4): 574–576. DOI: 10.1093/bioinformatics/btw663.

48. Laetsch, D.R. and M.L. Blaxter. 2017. BlobTools: Interrogation of genome
assemblies. F1000Research 6(1287). DOI: 10.12688/f1000research.12232.1.

49. Altschul, S.F., W. Gish, W. Miller, E.W. Myers, and D.J. Lipman. 1990. Basic
local alignment search tool. Journal of Molecular Biology 215(3): 403–410.

50. Wickham, H., *ggplot2: Elegant Graphics for Data Analysis*. 978-3-319-24277-4,
https://ggplot2.tidyverse.org ed. 2016: Springer-Verlag New York. 189.

51. Team, R.C. 2023. A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

52. Team, R. 2020. RStudio: Integrated Development for R. http://www.rstudio.com/.
PBC, Boston, MA, USA.

53.   Geneious Prime v. 2022.0.1. https://www.geneious.com.

54.   Xie, Y., Z. Xu, Y. Zheng, Y. Li, Y. Liu, L. Wang, X. Zhou, Z. Zuo, X. Gu, and G. Yang. 2019. The mitochondrial genome of the dog hookworm *Ancylostoma caninum* (Nematoda, Ancylostomatidae) from Southwest China. Mitochondrial DNA Part B 4(2): 3002–3004. DOI: 10.1080/23802359.2019.1666048.

55.   Gurevich, A., V. Saveliev, N. Vyahhi, and G. Tesler. 2013. QUAST: quality assessment tool for genome assemblies. Bioinformatics 29(8): 1072–1075. DOI: 10.1093/bioinformatics/btt086.

56.   Huang, N. and H. Li. 2023. miniBUSCO: a faster and more accurate reimplementation of BUSCO. bioRxiv: 2023.06. 03.543588. DOI: 10.1101/2023.06.03.543588.

57.   Keilwagen, J., F. Hartung, M. Paulini, S.O. Twardziok, and J. Grau. 2018. Combining RNA-seq data and homology-based gene prediction for plants, animals and fungi. BMC Bioinformatics 19(189). DOI: 10.1186/s12859-018-2203-5.

58.   Keilwagen, J., M. Wenk, J.L. Erickson, M.H. Schattat, J. Grau, and F. Hartung. 2016. Using intron position conservation for homology-based gene prediction. Nucleic Acids Research 44(9): e89. DOI: 10.1093/nar/gkw092.

59.   Dainat, J., D. Hereñú, and P. Pucholt. 2020. AGAT: Another Gff Analysis Toolkit to handle annotations in any GTF/GFF format (Version v0.7.0). DOI: 10.5281/zenodo.3552717.

60.   Pages, H., P. Aboyoun, R. Gentleman, S. DebRoy, M.H. Pages, D. DataImport, S. BSgenome, R. XStringSet-class, R. MaskedXString-class, and R. XStringSet-io. 2013. Package 'Biostrings'. Bioconductor: 18129.

Table 1. Comparative quality metrics output by QUAST for the assemblies generated as part of this study and the reference assemblies available for each species. Abbreviations: bp=base pairs.

		Length (bp)	No. contigs	N50 (bp)	GC content	N content
Brugia malayi	Ghedin et al. (2007) reference assembly	88,235,797	197	14,214,749	28.5%	0.30%
	MinION data only assembly	88,513,498	88	4,666,641	28.43%	0%
	MinION assembly polished with Illumina data	88,428,546	88	4,667,161	28.45%	0%
	Hybrid assembly	84,528,932	49	3,943,761	28.57%	<0.001%
Trichuris trichiura	Foth et al. (2014) reference assembly	75,474,068	4,086	70,602	42.3%	0.30%
	Doyle et al. (2022) reference assembly	80,573,711	113	11,299,416	42.3%	0.31%
	MinION data only assembly	73,094,541	174	806,679	42.22%	0%
	MinION assembly polished with Illumina data	72,959,001	174	805,117	42.23%	0%
	Hybrid assembly	72,705,678	63	2,773,302	41.15%	0%
Ancylostoma caninum	International Helminth Genomes Consortium (2019) reference assembly	465,750,606	25,339	256,700	42.8%	13.50%
	MinION data only assembly	358,159,610	1,578	638,901	43.36%	0%
	MinION assembly polished with Illumina data	357,986,713	1,578	637,973	43.38%	0%
	Hybrid assembly	348,034,555	652	1,095,227	43.24%	<0.001%

Table 2. Scores from miniBUSCO for the assemblies generated as part of this study and the reference assemblies available for each species. Scores are presented as number of BUSCOs recovered in each assembly followed in parentheses by proportion of the total number of nematode orthologs assessed by miniBUSCO (n=3,131).

		Single copy	Duplicated	Fragmented	Missing
Brugia malayi	Ghedin et al. (2007) reference assembly	3,100 (99.01%)	15 (0.48%)	8 (0.26%)	8 (0.26%)
	MinION data only assembly	3,097 (98.91%)	17 (0.54%)	9 (0.29%)	8 (0.26%)
	MinION assembly polished with Illumina data	3,097 (98.91%)	17 (0.54%)	9 (0.29%)	8 (0.26%)
	Hybrid assembly	3,096 (98.88%)	17 (0.54%)	8 (0.26%)	10 (0.32%)
Trichuris trichiura	Foth et al. (2014) reference assembly	1,676 (53.53%)	15 (0.48%)	112 (3.58%)	1,328 (42.41%)
	Doyle et al. (2022) reference assembly	1,702 (54.36%)	46 (1.47%)	102 (3.26%)	1,281 (40.91%)
	MinION data only assembly	1,717 (54.84%)	20 (0.64%)	99 (3.16%)	1,295 (41.36%)
	MinION assembly polished with Illumina data	1,716 (54.81%)	20 (0.64%)	100 (3.19%)	1,295 (41.36%)
	Hybrid assembly	1,725 (55.09%)	22 (0.70%)	104 (3.32%)	1,280 (40.88%)
Ancylostoma caninum	International Helminth Genomes Consortium (2019) reference assembly	2,654 (84.77%)	239 (7.63%)	143 (4.57%)	95 (3.03%)
	MinION data only assembly	2,865 (91.50%)	100 (3.19%)	66 (2.11%)	100 (3.19%)
	MinION assembly polished with Illumina data	2,867 (91.57%)	99 (3.16%)	65 (0.08%)	100 (3.19%)
	Hybrid assembly	2,883 (92.08%)	111 (3.55%)	57 (1.82%)	80 (2.56%)

Table 3. Comparative summary statistics output by AGAT for genes predicted by GeMoMa in assembly types for each species.
Mean lengths are presented as number of base pairs. Abbreviations: CDS=coding sequence.

		Mean gene length	Mean exon length	Mean intron length	Mean CDS length	Mean no. exons per CDS
Brugia malayi	MinION data assembly	4,299	148	346	1,394	9.4
	MinION assembly polished with Illumina data	4,291	148	346	1,394	9.4
	Hybrid assembly	4,267	148	344	1,389	9.4
Trichuris trichiura	MinION data assembly	2,916	212	325	1,280	6.0
	MinION assembly polished with Illumina data	2,909	212	325	1,279	6.0
	Hybrid assembly	2,918	212	324	1,283	6.0
Ancylostoma caninum	MinION data assembly	2,663	125	332	822	6.5
	MinION assembly polished with Illumina data	2,655	125	330	821	6.5
	Hybrid assembly	2,681	125	332	825	6.6

	MinION fast5 data	**Illumina fastq data**

Raw data quality control

MinION fast5 data
- Use *guppy* specifying a super high accuracy basecalling model to trim adapters and basecall raw fast5 data
- Run *nanoplot* to assess quality of basecalled data
- Run *mash* to visualize distances between reads, ensure they come from the same source organism, and identify potential contaminants
- Identify read-to-read overlaps and estimate average depth of coverage to a reference genome with *minimap2* and *samtools*
- *Optional:* run *nanofilt* to specify length and quality cut-offs for read dataset prior to assembly (alternatively, specify a read length minimum to during assembly)

Illumina fastq data
- Use *fastp* to trim adapters and assess data quality simultaneously
- Run *mash* to visualize distances between reads, ensure they come from the same source organism, and identify potential contaminants
- Identify estimate average depth of coverage to a reference genome with *BWA* and *samtools*
- Generate and analyse a k-mer spectrum using *jellyfish* or *kmc* and then the *GenomeScope online GUI* to estimate genome size, heterozygosity, error & repeat rates, etc.
- *Optional:* run *smudgeplot* and/or *enquire* to confirm ploidy and further explore k-mer based results
- Run *fastp* again, only removing sequencing adaptors, to generate input for *MaSuRCA*

MinION-only assembly
- Use *canu* to generate a *de novo* long-read MinION data-only assembly
- Remove contigs that are suspected bubbles using *SeqKit* and the *BBMap filterbyname.sh script*
- Use *purge_dups* to remove duplicated regions of the assembly resulting from failure of the assembler to recognize allelic contigs
- Separately and manually assemble organelle genomes from the unpurged assembly using *minimap2, samtools* and *Geneious*

Hybrid assembly
- Use *MaSuRCA* to generate a *de novo* hybrid assembly
- Use *purge_dups* to remove duplicated regions of the assembly resulting from failure of the assembler to recognize allelic contigs
- Separately and manually assemble organelle genomes from the unpurged assembly using *minimap2, BWA, samtools* and *Geneious*

Polishing

MinION fast5 data
- Polish the MinION data-only assembly with raw, basecalled MinION long-read data using *racon* (facilitated by *minimap2*) followed by *medaka*

Illumina fastq data
- *Optional: use quality-controlled Illumina data and pilon to polish a copy of the popped, purged, and long read-polished MinION data-only assembly*

Refinement

MinION fast5 data
- Identify, visualize, and remove contaminants from final assembly using *BlobTools*

Illumina fastq data
- Detect false duplication and assess completeness of hybrid assembly (i.e., whether assembly contains all k-mers present in the reads) and whether heterozygosity has been correctly collapsed using the "assembly spectra copy number plots" function in the *Kmer Analysis Toolkit (KAT)*
- Identify, visualize, and remove common contaminants from final assembly using *BlobTools*

Assessment

MinION fast5 data
- Use *QUAST* to compare the final assembly to a reference genome
- Use the "nucmer", "dnadiff" and "mummerplot" functions of *MuMMer4* to compare the final assemblies to the reference
- Use *miniBUSCO* to identify the proportion of expected single copy orthologs present in the final assembly

Illumina fastq data
- Use *QUAST* to compare the final assembly to a reference genome
- Use the "nucmer", "dnadiff" and "mummerplot" functions of *MuMMer4* to compare the final assemblies to the reference
- Use *miniBUSCO* to identify the proportion of expected single copy orthologs present in the final assembly