Generating reference-quality *de novo* genome assemblies from parasitic

nematodes using the Oxford Nanopore Technologies MinION sequencing platform

Kaylee S. Herzog[1], Rachel Wu[1], John M. Hawdon[2], Peter Nejsum[3], and Joseph R.

Fauver[1]*

[1]Department of Epidemiology, University of Nebraska Medical Center, Omaha, NE 68198

[2]Department of Microbiology, Immunology, and Tropical Medicine, The George

Washington University, Washington, D.C. 20037

[3]Department of Clinical Medicine, Aarhus University, 8200 Aarhus, Denmark

*Author for Correspondence: Joseph R. Fauver, Department of Epidemiology, University

of Nebraska Medical Center, Omaha, NE 68198, (402) 552-3397, (402) 559-4961,

jfauver@unmc.edu

Abstract

Parasitic nematode infections represent a significant burden of disease in
impoverished populations. High prevalence of infection coupled with widespread
geographic distributions means populations of parasitic nematodes likely harbor high
levels of genetic diversity that are not reflected in currently available genomic resources.
Comparative genomic studies of parasitic nematodes have identified novel drug and
vaccine targets and yielded unprecedented insights into helminth biology, highlighting the
potential of high-quality reference genomes. Few genomic resources exist for parasitic
nematodes, however, and most species are represented by only a single genome. This
is partly due to challenges associated with generating genomic data for these parasites.
Many species are small in size and yet have relatively large and complex genomes,
making their genomes difficult and expensive to sequence and assemble. Short-read
sequence data has primarily been used to generate genomic data for parasitic
nematodes, but whole genome assemblies based on short-read data are highly
discontiguous. Recent improvements in the accuracy of Oxford Nanopore Technologies
(ONT) long-read data makes it an appealing approach for improving existing reference
genomes and generating novel genome assemblies from uncharacterized species. This
study targets three species of parasitic nematodes to compare the current gold standard
hybrid *de novo* genome assembly approach to one that relies on long-read ONT MinION
data only. Hybrid assemblies were found to be more contiguous, but nonetheless, MinION
data alone were used successfully to generate highly contiguous, complete, and accurate
assemblies for all species quickly and inexpensively.

Key Words

Parasitic nematodes, whole genome sequencing, genome assembly, Oxford Nanopore
Technologies, MinION sequencing, BUSCO

Significant Statement

High quality helminth genome assemblies are invaluable tools for identifying novel drug
and vaccine targets and monitoring the effectiveness of control campaigns, but
generating regionally relevant reference-quality genomes from a broad diversity of
populations and species can be challenging when relying on traditional approaches. This
study demonstrates that Oxford Nanopore Technologies MinION long-read data alone
can be used to quickly and inexpensively generate highly contiguous, complete, and
accurate whole genome assemblies for parasitic nematodes. These findings highlight an
accessible method for increasing the resources available for comparative helminth
genomics and a path toward incorporating genomic epidemiology as a routine component
of assessing control campaign effectiveness.

Data Availability Statement

Quality-controlled MinION and Illumina data for each species are deposited in the NCBI
Sequence Read Archive (SRA) under the BioProject accession number XXXX. Final
assemblies for each species are publicly available on WormBase ParaSite.

Introduction

Parasitic nematodes inflict an enormous burden of human disease. It is estimated that 1.5 billion people alone are infected with soil-transmitted helminths, while approximately 120 million and 15 million people suffer from filariasis and onchocerciasis, respectively (Global Burden of Disease Study, 2020). In addition to the morbidity and mortality caused by nematodes themselves, infection with these parasites also exacerbates the disease burden of other existing conditions such as malaria, HIV, and tuberculosis, and can reduce immunological response to important vaccines (Ndyomugyenyi et al., 2008; Kizito et al., 2013; Biraro et al., 2014; Morawski et al., 2017; Nash et al., 2017). Accordingly, the World Health Organization (WHO) has established large-scale global control programs to reduce the incidence of nematode infections (Casulli, 2021; International Helminth Genomes Consortium, 2021). These programs are centered around broad distribution of anthelmintic drugs to all individuals in vulnerable communities regardless of their infection status. Because a limited number of inexpensive drugs can be used to treat multiple parasitic infections, these mass drug administration campaigns (i.e., MDAs) are touted as an inexpensive, broadly effective control measure for a variety of nematode parasites. As programmatic monitoring focuses almost exclusively on metrics of infected individuals themselves, however, the question of the impact of repeated exposure to anthelmintic drugs on nematode populations remains unanswered. Recent advances in next generation sequencing (NGS) technologies now allow for the monitoring of parasite population structure over the course of long-term treatment with anthelminthics, thus enabling the application of genomic epidemiological approaches to assess control program efficacy (Berger et al., 2021).

Genomic epidemiology requires high quality and regionally specific reference

genomes for the species of parasite(s) being treated and monitored (Formenti et al.,

2022). This is due to the fact that most genomic epidemiological approaches for

eukaryotes rely on population-level datasets in which individuals are genotyped based on

mapping reads for loci of interest to a conspecific reference-quality genome (Sims et al.,

2014; Hoban et al., 2016). The advent of NGS—and specifically, reductions in its cost

and improved ease of access—has recently made the generation of genomic data from

many populations of parasites feasible (Doyle 2022). While this has resulted in an

increased number of population-level studies for parasites that are based on genomic

data (e.g., Brabec et al., 2016; Sweet & Johnson, 2018; Berger et al., 2021; Fraija-

Fernández et al., 2021; Doyle et al., 2022; Virrueta Herrera et al., 2022), there has not

been an equivalent increase in the generation of reference genomes representative of

these populations.

The reference genomes that do exist for parasitic nematodes have proven to be

an invaluable resource for increasing our understanding of parasite biology, treatment,

and control. For example, the landmark comparative genomic study by the International

Helminth Genomes Consortium analyzed tens of complete genomes representing all

major helminth lineages, including nematodes. This expansive dataset allowed for

identification of multiple gene family expansions in various groups that are targets for

novel drug and vaccine development (International Helminth Genomes Consortium,

2019). Studies such as these highlight the importance of parasite genomes for not only

population-level genomic epidemiological approaches, but also as indispensable tools for

comparative biomedical research.

Historically, short-read sequences like those produced by Illumina NGS platforms

have been the data type of choice for assembling parasite genomes. Though highly

accurate, short-read data alone are unable to produce contiguous reference genomes for

most species (Bashir et al., 2012; English et al., 2012; Koren et al., 2012). The Oxford

Nanopore Technologies (ONT) MinION NGS platform offers an appealing complimentary

approach. Its benefits include reduced equipment costs and infrastructure requirements

as compared to traditional Illumina platforms, ease of use and portability, and, importantly,

the ability to generate the long-read sequence data needed to produce contiguous

reference-quality genomes. Recent updates to ONT chemistries put the error rates of

MinION sequence data on par with those of Illumina platforms, allowing for the potential

to generate parasite reference genomes using long-read data, only.

This study aims to compare whole genomes of parasitic nematodes assembled *de*

novo using the current gold standard hybrid approach of combining both short- and long-

read data during the assembly process versus using Q20+ ONT MinION long-read data,

only. We utilize three species for sequencing which collectively span the breadth of

nematode diversity: the filarial worm *Brugia malayi* (Spirurina), the whipworm *Trichuris*

trichiura (Trichinellida), and the dog hookworm *Ancylostoma caninum* (Rhabditina).

Ultimately, the goal of this study is to evaluate whether the improved quality of Q20+

MinION long reads make these data alone sufficient for generating reference-quality

assemblies, highlighting a quick, inexpensive, and accessible method for generating

parasite genomes.

Results

128 <u>DNA extraction, library preparation, and sequencing</u>

129   The total amount of gDNA extracted from each sample type and the mean

130 fragment lengths of extractions are provided in Supplementary Table 1.

131   *MinION library preparation and sequencing:* Figure 1 provides a summary of the

132 bioinformatic pipeline used to process MinION sequence data. The concentrations and

133 mean fragment sizes of MinION libraries are provided in Supplementary Table 1. The

134 amount of data generated from each flow cell and the amount of data successfully

135 basecalled by Guppy are also provided in Supplementary Table 1. Select NanoPlot

136 outputs summarizing each Guppy run are provided for each species separately in

137 Supplementary Figures 1–3. For *T. trichiura* and *A. caninum*, for which more than one

138 flow cell's worth of data were generated, NanoPlot plots are based on input from two (*T.*

139 *trichiura*) or three (*A. caninum*) sequencing summary files. Values for average depth of

140 coverage and proportion of basecalled MinION reads mapped to species-specific

141 reference genomes as estimated using Minimap2 and SAMtools were 124.85× (98.77%

142 of reads mapped) for *B. malayi*, 249.91× (99.91% of reads mapped) for *T. trichiura*, and

143 49.42× (99.85% of reads mapped) for *A. caninum.* For *B. malayi*, percent identity and

144 median multiplicity values output by Mash indicate the majority of MinION reads were

145 classified as *B. malayi* or its *Wolbachia* endosymbiont (see Supplementary Table 2). For

146 *T. trichiura* and *A. caninum*, Mash outputs were less straightforward. Neither species was

147 identified within the MinION read sets analyzed; rather, various lineages of virus, bacteria,

148 and fungi were assigned, largely with relatively low percent identity and median

149 multiplicity values (see Supplementary Table 2).

MinION fast5 data

Illumina fastq data

Raw data quality control

- Use *guppy* specifying a super high accuracy basecalling model to trim adapters and basecall raw fast5 data
- Run *nanoplot* to assess quality of basecalled data
- Run *mash* to visualize distances between reads, ensure they come from the same source organism, and identify potential contaminants
- Identify read-to-read overlaps and estimate average depth of coverage to a reference genome with *minimap2* and *samtools*
- *Optional:* run *nanofilt* to specify length and quality cut-offs for read dataset prior to assembly (alternatively, specify a read length minimum to during assembly)

- Use *fastp* to trim adapters and assess data quality simultaneously
- Run *mash* to visualize distances between reads, ensure they come from the same source organism, and identify potential contaminants
- Identify estimate average depth of coverage to a reference genome with *BWA* and *samtools*
- Generate and analyse a k-mer spectrum using *jellyfish* or *kmc* and then the *GenomeScope online GUI* to estimate genome size, heterozygosity, error & repeat rates, etc.
- *Optional:* run *smudgeplot* and/or *enquire* to confirm ploidy and further explore k-mer based results
- Run *fastp* again, only removing sequencing adaptors, to generate input for *MaSuRCA*

MinION-only assembly

- Use *canu* to generate a *de novo* long-read MinION data-only assembly
- Remove contigs that are suspected bubbles using *SeqKit* and the *BBMap filterbyname.sh script*
- Use *purge_dups* to remove duplicated regions of the assembly resulting from failure of the assembler to recognize allelic contigs
- Separately and manually assemble organelle genomes from the unpurged assembly using *minimap2, samtools* and *Geneious*

Hybrid assembly

- Use *MaSuRCA* to generate a *de novo* hybrid assembly
- Use *purge_dups* to remove duplicated regions of the assembly resulting from failure of the assembler to recognize allelic contigs
- Separately and manually assemble organelle genomes from the unpurged assembly using *minimap2, BWA, samtools* and *Geneious*

Polishing

- Polish the MinION data-only assembly with raw, basecalled MinION long-read data using *racon* (facilitated by *minimap2*) followed by *medaka*

- *Optional: use quality-controlled Illumina data and pilon to polish a copy of the popped, purged, and long read-polished MinION data-only assembly*

Refinement

- Identify, visualize, and remove contaminants from final assembly using *BlobTools*

- Detect false duplication and assess completeness of hybrid assembly (i.e., whether assembly contains all k-mers present in the reads) and whether heterozygosity has been correctly collapsed using the "assembly spectra copy number plots" function in the *Kmer Analysis Toolkit (KAT)*
- Identify, visualize, and remove common contaminants from final assembly using *BlobTools*

Assessment

- Use *QUAST* to compare the final assembly to a reference genome
- Use the "nucmer", "dnadiff" and "mummerplot" functions of *MuMMer4* to compare the final assemblies to the reference
- Use *miniBUSCO* to identify the proportion of expected single copy orthologs present in the final assembly

- Use *QUAST* to compare the final assembly to a reference genome
- Use the "nucmer", "dnadiff" and "mummerplot" functions of *MuMMer4* to compare the final assemblies to the reference
- Use *miniBUSCO* to identify the proportion of expected single copy orthologs present in the final assembly

Figure 1. Bioinformatic pipeline used to generate MinION data (left) and hybrid (right) genome assemblies.

Illumina library preparation and sequencing: Figure 1 provides a summary of the

bioinformatic pipeline used to process Illumina sequence data. The amount of raw

sequencing data generated for each Illumina library, and the amount of data retained

following the first iteration of filtration with fastp, are provided in Supplementary Table 1.

Values for average depth of coverage and proportion of quality-controlled Illumina reads

mapped across species-specific reference genomes as estimated using BWA and

SAMtools were 82.95× (95.74% of reads mapped) for *B. malayi*, 48.58× (90.71% of reads

mapped) for *T. trichiura*, and 38.64× (90.31% of reads mapped) for *A. caninum*. As with

MinION data, Mash results indicated that Illumina read datasets for *B. malayi* were

overwhelmingly identified as *B. malayi* or its *Wolbachia* endosymbiont but indicated no

species-appropriate hits for *T. trichiura* or *A. caninum* (see Supplementary Table 2).

<u>Estimation of genome size, heterozygosity, and ploidy</u>

GenomeScope plots for each species are provided in Supplementary Figure 4.

Genome size (and heterozygosity) estimates from Jellyfish and GenomeScope were

85,917,606 bp (0.275%) for *B. malayi*, 68,538,097 bp (1.35%) for *T. trichiura,* and

329,957,709 (1.09%) for *A. caninum*. Genome size estimates for all three species were

lower than the current reference assembly lengths of ~88 Mb for *B. malayi* (Ghedin et al.,

2007),   ~75–80   Mb   for   *T.   trichiura*   (Foth   et   al.,   2014)

(https://github.com/stephenrdoyle/ancient_trichuris/tree/master/02_data;   Doyle   et   al.,

2022), and ~465 Mb for *A. caninum* (International Helminth Genomes Consortium, 2019).

Model fit values estimated by GenomeScope ranged from ~93.4–94.5% (*B. malayi*),

~91.6–93.3% (*T. trichiura*) and ~94.1–96.2% (*A. caninum*). For *T. trichiura*, based on log-

likelihood scores, nQuire preferred a triploid model over diploid and tetraploid models for

both Illumina and MinION read datasets (see Supplementary Table 3).

<u>MinION data assembly and refinement</u>

Histograms of Q scores for MinION reads versus Q scores for MinION reads

aligned to final MinION data assemblies are provided for each species in Supplementary

Figure 5. For all three species, a median Q score value of >Q16 was achieved in both

categories. *Brugia malayi* had the highest median Q score, approaching Q20 in both

categories, followed closely by *T. trichiura*. The BlobPlots and summary of by-contig

BLAST results produced by BlobTools for MinION data assemblies are provided in

Supplementary Figures 6–8 and Supplementary Table 4, respectively. For *B. malayi* and

A. caninum, four contigs totaling 49,030 bp and four contigs totaling 116,599 bp,

respectively, were confidently identified as potential contamination and were removed

from final assemblies (see Supplementary Table 4); however, contigs from the *T. trichiura*

and *A. caninum* assemblies returned largely off-target BLAST results (see Supplementary

Table 4; Supplementary Figures 7, 8).

<u>Hybrid assembly and refinement</u>

K-mer multiplicity plots output by KAT for purged hybrid assemblies are provided

for all three species in Supplementary Figure 9. A single peak in the plot for *B. malayi*

indicates low heterozygosity whereas two distinct peaks in the plots for *T. trichiura* and *A.*

caninum indicates higher levels of heterozygosity. For all three species, KAT plots

suggest that errors and—in the case of *T. trichiura* and *A. caninum*, heterozygosity—were

successfully purged from final hybrid assemblies. The BlobPlots and summary of by-

contig BLAST results produced by BlobTools for hybrid assemblies are provided in

Supplementary Figures 10–12 and Supplementary Table 4, respectively. Only a single

contig was confidently identified as contamination and eliminated from a hybrid assembly

(i.e., for *A. caninum*; see Supplementary Table 4). As with MinION data assemblies, *T.*

trichiura and *A. caninum* assemblies returned high proportions off-target BLAST results

(see Supplementary Table 4; Supplementary Figures 11, 12).

<u>Assembly contiguity, accuracy, and completeness</u>

Assembly length, contig number, N50 value, and percent GC and N content as

output by QUAST are provided for MinION data, hybrid, and reference assemblies in

Table 1. The number of gSNPs and gINdels identified by MuMmer4 between MinION data

and hybrid assemblies, and the nucleotide-level percent identity where these two

assemblies align, are also provided in Table 1. Graphs of contig number by genome size

are provided for each species in Figure 2. With the exception of the MinION data

assembly for *B. malayi*, all assemblies generated here were shorter in terms of total length

than the current reference assemblies available for each species (Table 1; Figure 2). In

comparison to the genomes available on WormBase ParaSite for *T. trichiura* (Foth et al.,

2014), and *A. caninum* (International Helminth Genomes Consortium, 2019), both MinION

data and hybrid assemblies were more contiguous (as measured by a smaller contig

number and a larger N50 value) than the existing reference assemblies, and hybrid

assemblies were in all cases more contiguous than MinION data assemblies (Table 1;

Figure 2). By species, all assemblies had similar levels of GC content within, and MinION

Table 1. Comparative quality metrics for the assemblies generated as part of this study and the reference assemblies available for each focal species.

Species	Assembly	QUAST					MuMmer4		
		Length (bp)	No. contigs	N50 (bp)	GC content	N content	Nucleotide-level genome identity where aligned	No. gSNPS	No. gIndels
Brugia malayi	WormBase ParaSite reference	88,235,797	197	14,214,749	28.5%	0.30%			
	MinION data-only assembly	88,513,498	88	4,666,641	28.43%	0%	99.34%	150,681	9,434
	Hybrid assembly	84,528,932	49	3,943,761	28.57%	<0.001%	98.68%		
Trichuris trichiura	WormBase ParaSite reference	75,474,068	4,086	70,602	42.3%	0.30%			
	Doyle et al. (2022) reference	80,573,711	113	112,99,416	42.3%	0.31%			
	MinION data-only assembly	73,094,541	174	806,679	42.22%	0%	99.32%	35,115	36,140
	Hybrid assembly	72,705,678	63	27,73,302	41.15%	0%	99.60%		
Ancylostoma caninum	WormBase ParaSite Reference	465,750,606	25,339	256,700	42.8%	13.50%			
	MinION data-only assembly	358,159,610	1,578	638,901	43.36%	0%	97.33%	299,510	1,258,880
	Hybrid assembly	348,034,555	652	1,095,227	43.24%	<0.001%	96.16%		

219   data and hybrid assemblies were comprised of 0%, or essentially 0%, ambiguous

220   nucleotide calls, respectively (Table 1). Nucleotide-level alignment identity between

221   MinION data and hybrid assemblies as assessed by MuMmer4 ranged from ~96.2–99.6%

222   (Table 1).

Figure 2. Comparison of contiguity of the assemblies generated as part of this study and existing reference assemblies for (A) *Brugia malayi***, (B)** *Trichuris trichiura***, and (C)** *Ancylostoma caninum***.** Numbers in parenthesis within keys at lower right are the total number of contigs in each assembly; asterisks indicate a scaffold count rather than a contig count.

Scores from miniBUSCO for all assembly types are summarized for each species

in Table 2. For all three species, scores for each category are nearly identical among the

three assembly types generated herein (i.e., MinION data, MinION data polished with

Illumina data, and hybrid). For *A. caninum*, miniBUSCO scores indicate MinION and

hybrid assemblies are more complete as compared to the existing reference assembly.

All assemblies compared for *T. trichiura*, including both available references, had high

proportions of missing BUSCOs (i.e., ~40.8–42.4%; see Table 2). A summary of the

assessment of "tier 1" helminth genome status *sensu* the International Helminth Genomes

Consortium (International Helminth Genomes Consortium, 2019) for the assemblies

generated here, and assemblies of all nematodes that parasitize animal hosts available

from WormBase ParaSite, are presented in Figure 3. Both the MinION data and hybrid

assemblies for *B. malayi* and *A. caninum* are classified as "tier 1" genomes. For *T.*

trichiura, the contiguity of both the MinION data and hybrid assembly is sufficient to qualify

them each as "tier 1", but, as mentioned above, all assemblies for trichurids appear to be

missing a significant proportion of nematode BUSCOs, preventing any trichurid assembly

assessed from achieving "tier 1" status (see Figure 3).

In terms of assembly accuracy, one of the most salient questions is how different

MinION data assemblies are from assemblies that include both MinION and Illumina data.

Table 2. Scores from miniBUSCO for MinION data-only, MinION data-only polished with Illumina data, hybrid, and reference assemblies for each focal species. Scores are presented as number of BUSCOs recovered in each assembly followed in parentheses by proportion of the total number of nematode orthologs assessed by miniBUSCO (n=3,131).

		Single copy	Duplicated	Fragmented	Missing
Brugia malayi	WormBase ParaSite reference	3,100 (99.01%)	15 (0.48%)	8 (0.26%)	8 (0.26%)
	MinION data-only assembly	3,097 (98.91%)	17 (0.54%)	9 (0.29%)	8 (0.26%)
	MinION assembly + Illumina polishing	3,097 (98.91%)	17 (0.54%)	9 (0.29%)	8 (0.26%)
	Hybrid assembly	3,096 (98.88%)	17 (0.54%)	8 (0.26%)	10 (0.32%)
Trichuris trichiura	WormBase ParaSite reference	1,676 (53.53%)	15 (0.48%)	112 (3.58%)	1,328 (42.41%)
	Doyle et al. (2022) Reference	1,702 (54.36%)	46 (1.47%)	102 (3.26%)	1,281 (40.91%)
	MinION data-only assembly	1,717 (54.84%)	20 (0.64%)	99 (3.16%)	1,295 (41.36%)
	MinION assembly + Illumina polishing	1,716 (54.81%)	20 (0.64%)	100 (3.19%)	1,295 (41.36%)
	Hybrid assembly	1,725 (55.09%)	22 (0.70%)	104 (3.32%)	1,280 (40.88%)
Ancylostoma caninum	WormBase ParaSite reference	2,654 (84.77%)	239 (7.63%)	143 (4.57%)	95 (3.03%)
	MinION data-only assembly	2,865 (91.50%)	100 (3.19%)	66 (2.11%)	100 (3.19%)
	MinION assembly + Illumina polishing	2,867 (91.57%)	99 (3.16%)	65 (0.08%)	100 (3.19%)
	Hybrid assembly	2,883 (92.08%)	111 (3.55%)	57 (1.82%)	80 (2.56%)

To address this question, we utilized BUSCO to objectively generate comparative

gene datasets. BUSCO outputs the DNA sequence for each single copy ortholog it

identifies in an assembly. We compiled these sequences for MinION data assemblies,

MinION data assemblies polished with Illumina data (i.e., "polished" assemblies) and

hybrid assemblies, and for each species, we identified which sequences were shared

between MinION data and polished, and MinION data and hybrid, assemblies. We then

generated pairwise alignments for each shared gene and used comparative identity

metrics to assess nucleotide-level differences. The goal of generating these comparisons

Figure 3. Plot of single copy miniBUSCO score versus the log value of N50 length divided by scaffold (or contig) count for the assemblies generated as part of this study, and for all genomes of nematodes that parasitize animals as adults available from WormBase ParaSite. Dotted lines indicate the cutoff for "tier 1" genome status *sensu* the International Helminth Genome Consortium (i.e., >85% single copy BUSCO score and >1.6 log value for the contiguity metric). A key to symbolic designations is provided at top left.

was to assess whether the accuracy of assembled protein coding sequences is

systematically affected by data type.

A summary of pairwise alignment designations and percent identities for genes

shared between assembly types for each species is provided in Table 3. For all species,

a greater proportion of identical genes were shared between MinION data and polished

assemblies (2,893 for *B. malayi*, 1,485 for *T. trichiura*, and 1,801 for *A. caninum*) as

Table 3. Comparison of pairwise alignments for genes shared between assembly types generated as part of this study. Columns for each species represent values for the MinION data assembly compared to either the hybrid assembly (left) or the polished assembly (right). The total number of complete single copy BUSCOs shared between assemblies is out of the total number of nematode orthologs assessed by BUSCO (n=3,131).

	Brugia malayi		*Trichuris trichiura*		*Ancylostoma caninum*	
	Versus hybrid assembly	Versus polished assembly	Versus hybrid assembly	Versus polished assembly	Versus hybrid assembly	Versus polished assembly
Total no. complete single copy genes shared between assemblies	3,047	3,071	1,750	1,799	2,656	2,812
No. (percent) of identical alignments	1,959 (64.29%)	2,893 (94.20%)	622 (35.54%)	1,485 (82.55%)	1,207 (45.44%)	1,801 (64.05%)
No. (percent) of alignments that differ by SNPs only	411 (13.49%)	10 (0.36%)	420 (24.00%)	90 (5.00%)	316 (11.90%)	222 (7.89%)
No. (percent) of alignments that differ by SNPs and Indels	477 (15.65%)	93 (3.03%)	664 (37.94%)	177 (9.83%)	947 (35.66%)	600 (21.34%)
No. (percent) of alignments that differ by Indels only	200 (6.56%)	75 (2.44%)	44 (2.51%)	47 (2.61%)	186 (7.00%)	189 (6.72%)
Average percent identity | Average percent identity where aligned	98.08% | 99.74%	99.59% | 99.95%	95.80% | 98.74%	99.02% | 99.75%	95.19% | 98.82%	97.45% | 99.52%

compared to MinION data and hybrid assemblies (1,959 for *B. malayi*, 622 for *T. trichiura*,

and 1,207 for *A. caninum*) (see Table 3). Additionally, across species, average percent

identity, and average percent identity where aligned, were each greater for pairwise

alignments from MinION data versus polished assemblies as compared to hybrid

assemblies (see Table 3).

Dot plots of the lengths of genes shared between assembly types that differed by

length variants (i.e., Indels only, or SNPs and Indels), are provided for each species in

Figure 4A–F. (These alignments were of particular interest for determining whether one

assembly method systematically over- or under-represents gene length.) For all species,

a greater proportion of shared genes were found to differ by length variants in MinION

data versus hybrid assemblies as compared to polished assemblies. For each species

and comparison, no substantial difference was identified between the number of genes

identified as longer in MinION data versus polished or hybrid assemblies (Figure 4).

For all species, assembly comparisons, and pairwise alignment designations,

violin plots of percent identity and percent identity where aligned for alignments that

scored >90% in identity are provided in Figure 4G and 4F, respectively. For MinION data

versus hybrid assemblies, 88.65% and 98.36% of alignments had >90% identity and

>90% identity where aligned, respectively. For MinION data versus polished assemblies,

95.54% and 99.37% of alignments had >90% identity and >90% identity where aligned,

respectively. Across all species and comparisons, most pairwise alignments generated

for were found to be identical or nearly identical, with comparatively few alignments

scoring <97% in identity (see Figure 4G, F). Raw data characterizing each pairwise

Figure 4. Comparisons of genes that differ by length variants shared between assembly types. Dot plots of lengths of shared genes for (A–B) *Brugia malayi*, (C–D) *Trichuris trichiura*, and (E–F) *Ancylostoma caninum*. Dotted lines indicate equal lengths. Violin plots of (G) percent identity and (H) percent identity where aligned for all pairwise alignments generated for shared genes for each of the three focal species.

alignment generated for each assembly comparison are presented in Supplementary

Table 5.

Discussion

<u>Sample processing and sequencing</u>

Genomic DNA extracted from a single individual is the ideal input for *de novo* whole

genome assembly. This is in part because, for diploid organisms, assembly pipelines

need only contend with two potential haplotypes, leading to more accurate assemblies

with less haplotypic duplication. An alternative method would be to sequence multiple

individuals from a highly inbred homozygous line. Many species cannot be maintained in

a laboratory setting, however, making the establishment of inbred lines for these taxa

challenging or impossible (Solares et al., 2021). It is also now well-recognized that

inbreeding can significantly reduce genome size (Price, 1976; Fierst et al., 2015; Roessler

et al., 2019), making these lines poor representatives of the species as whole.

Furthermore, sampling from natural, rather than laboratory-maintained, populations is a

requirement for most genomic studies (including genomic epidemiology), further

highlighting the advantage of whole-genome sequencing from single individuals.

Considering these advantages, the DNA extraction and MinION library preparation

protocols described here were optimized for low input gDNA extracted from individual

parasitic nematodes. For *B. malayi* and *T. trichiura*, which both have relatively large adult

stages (i.e., ~3–5 cm in length) and tractable genome sizes (i.e., <100 Mb), these

protocols allowed for the generation of both short- and long-read data from a single adult

worm. Adults of *A. caninum*, however, tend not to exceed ~1.5 cm in length, and yet have

the largest genome size of the three focal species in this study (i.e., ~400 Mb). Thus, for

A. caninum, even if we had had access to adult worms, we expect that using these

optimized protocols would still not have allowed for generation of genomic data from a

single adult. Moving forward, the approaches outlined here can be adapted for use with

additional species of parasitic nematodes.

There was an association between the average Q score of MinION read datasets

and the total amount of data generated by a flow cell: More efficient MinION sequencing

runs generated data with higher average Q scores (see Supplementary Table 1). For

example, MinION libraries for both *B. malayi* and *T. trichiura* were sequenced for ~63–72

hr each and generated ~10.8–11.8 Gb of data per flow cell post-basecalling with

comparably high Q scores. Conversely, libraries for *A. caninum*, which were sequenced

for ~68–80 hr each, only generated ~7.7–8.5 Gb of data per flow cell post-basecalling,

and these data had the lowest average Q score of the three species sequenced (see

Supplementary Table 1; Supplementary Figure 5). Differences in pore occupancy during

sequencing could potentially explain this correlation. According to MinKNOW run reports,

the flow cell used to sequence the *B. malayi* library had the greatest proportion of

available and actively sequencing pores throughout the sequencing run while the flow

cells that were used to sequence the three *A. caninum* libraries experiences the most

rapid decline in pore availability. Whatever the reason for this association, it is clear that

sequencing efficiency should be taken into consideration when attempting to achieve

Q20+ data using the latest ONT chemistries.

Estimating contamination in read-level data

In terms of detecting potential contamination in read-level data, Mash worked well

for *B. malayi*, but did not return species-appropriate hits for *T. trichiura* or *A. caninum* (see

Supplementary Table 2). Given that high proportions (~90.3–99.9%) of both MinION and

Illumina reads mapped to the reference genomes for all species, however, it is unlikely

that these results indicate significant read-level contamination for the later two species.

Of the three species, however, *B. malayi* is the least well-represented in the NCBI RefSeq

archive (the databased utilized by Mash) in terms of total number of records, so why Mash

performed most effectively for *B. malayi* is unclear. It is worth noting that read N50 lengths

for *B. malayi* greater than those for *T. trichiura* or *A. caninum* (Supplementary Table 1),

and so perhaps longer sequences allowed the NCBI RefSeq archive to be queried more

accurately by Mash.

<u>Estimation of genome size, heterozygosity, and ploidy</u>

For *B. malayi*, the MinION data assembly was slightly larger than the current

reference assembly, while the hybrid assembly was appreciably smaller (see Table 1).

For each of *T. trichiura* and *A. caninum*, both assemblies generated as part of this study

were considerably smaller than existing reference genomes (see Table 1). For *A.*

caninum, this different was most dramatic, on the order of >100 Mb. Assemblies

generated here are likely smaller than current references due to that fact that a purging

step to remove haplotypic duplication was included in the bioinformatic pipeline for both

assembly types (see Figure 1). This was not the case in the generation of existing

reference genomes (Ghedin et al., 2007; Foth et al., 2014; Consortium, 2019), and so

they likely include haplotypic duplication that artificially inflates genome size—particularly

for *T. trichiura* and *A. caninum*, which were sequenced from heterozygous sample types.

Three lines of evidence support that the genomes assembled here more accurately

represent the true genome size for each species: (1) The lengths of these assemblies

more closely match k-mer based genome size estimates than do those of the existing

reference genomes (see Table 1, Supplementary Figure 4); (2) miniBUSCO scores are

largely indistinguishable across all assembly types (including existing reference

assemblies), indicating that all are comparably complete despite ranging in size; in fact,

the Doyle et al. (2022) reference assembly for *T. trichiura* and the WormBase ParaSite

reference assembly for *A. caninum* had higher proportions of duplicated BUSCOs as

compared to the assemblies generated herein (see Table 2); and (3) for hybrid

assemblies, KAT plots suggest they are not "over-purged" and thus missing genomic

content (see Supplementary Figure 9). As advocated previously, purging haplotypic

duplication is an important step in genome curation regardless of assembly approach

(Howe et al., 2021; Rhie et al., 2021), and, unsurprisingly, becomes increasingly

important for highly heterozygous sample types such as that sequenced for *A. caninum*.

Results from GenomeScope (Supplementary Figure 4) and the k-mer multiplicity

plots output by KAT (Supplementary Figure 9) indicate low heterozygosity for *B. malayi*

and elevated heterozygosity for *T. trichiura* and *A. caninum*. These results are congruent

with the fact that the specimen of *B. malayi* sequenced came from a long-maintained

inbred laboratory strain while *T. trichiura* and *A. caninum* were sequenced from

specimens from infections maintained for shorter periods of time.

Preliminary data from nQuire suggest that *T. trichiura* may be triploid (see

Supplementary Table 3). This result was unexpected, given that the well-studied model

species for trichuriasis, *Trichuris muris*, is diploid (Špakulová et al., 1994). An early study

by Goswami (1978) on the karyotypes of trichurids reports that *T. trichiura* is also diploid.

Goswami does report supernumerary chromosomes (i.e., small abnormal chromosomes

with few or no banding landmarks; Spinner et al., 2013) in *Trichuris ovis* and *Trichuris*

globulosa, but makes no mention of supernumerary chromosomes in *T. trichiura*. Despite

the publication of Goswami's work, more recent literature perplexingly states that the

karyotype of *T. trichiura* remains unknown (Foth et al., 2014; Yousuf et al., 2022). The

estimation of ploidy for *T. trichiura* made here was based purely on a bioinformatic

approach and should therefore not be considered a definitive assessment. Moving

forward, it seems a worthwhile pursuit to confirm the chromosome number and copy

number for this medically important species using a modern karyotyping approach.

<u>Assembly refinement</u>

As with Mash, BlobTools was a more effective tool for assessing potential

contamination in final assemblies for *B. malayi* than for *T. trichiura* and *A. caninum*. For

example, for *B. malayi*, BlobTools identified four small contigs in the MinION data

assembly that represented sequence from the jird host, and contigs in both assembly

types that represented sequence from the *Wolbachia* endosymbiont, with all remaining

contigs identified as belonging to the phylum Nematoda (see Supplementary Tables 3

and 4). For *T. trichiura* and *A. caninum*, however, while many contigs could be identified

as belonging to the phylum Nematoda, a large proportion of contigs in both assembly

types returned off-target results from other phyla, or could not be assigned to a taxon at

all (see Supplementary Table 4). All three species are represented in the NIH BLAST

database on which BlobTools is based, but the assemblies generated for *B. malayi* were

more contiguous than those generated for *T. trichiura* and *A. caninum*. As noted above

for Mash, perhaps longer contig lengths allowed for the return of more on-target BLAST

results for *B. malayi*.

<u>Assembly, contiguity, accuracy, and completeness</u>

Hybrid assemblies were in all cases more contiguous than MinION data

assemblies (see Table 1, Figure 2). (As polishing with Illumina data improves accuracy

rather than contiguity, MinION data assemblies polished with Illumina data were no

included in contiguity comparisons.) This result was somewhat unexpected given that

including short-read data in whole genome assembly has traditionally been touted to

increase accuracy rather than contiguity (Bashir et al., 2012; English et al., 2012; Koren

et al., 2012). While this was indeed the case when utilizing hybrid approaches with older

ONT chemistries, the results of the comparisons made here demonstrate that the

improved read-level quality of Q20+ MinION data may lessen the value of short-read data

for improving assembly accuracy. For example, alignment identity metrics indicate a high

degree of identity (i.e., ~96–99%) between MinION data and hybrid assemblies for each

species (see MuMmer4 results in Table 1).

Though often reported (e.g., Fu et al., 2020; Wang et al., 2020; Yi et al., 2020;

Ludington & Sanders, 2021; Voelker et al., 2021; Okenmor et al., 2022), genome-wide

alignment identity metrics like those produced by MuMmer4 are a broad and imperfect

measure of assembly accuracy. Therefore, we endeavored to conduct a more in-depth

analysis of accuracy by characterizing pairwise alignments of BUSCO sequences shared

between assembly types. The long-read and hybrid assemblers utilized here are based

on different assembly algorithms that handle read-level data dissimilarly. Though both are

built on a Celera Assembler foundation, Canu's MinHash Alignment Process (MHAP) is

broadly an overlap layout consensus-based method (Koren et al., 2017) while MaSuRCA

relies on a "super-read" algorithm that combines aspects of both overlap layout

consensus and de Bruijn graph-based approaches (Zimin et al., 2013). For assessing the

ability of short-read data to improve long-read assembly accuracy, comparing MinION

data assemblies to MinION data assemblies polished with Illumina data is a more salient

comparison than comparing to hybrid assemblies. If polishing with Illumina data

substantially improved the accuracy of MinION data assemblies, we would expect the

majority of genes found in both assembly types to differ at the sequence level. This was

not the case, as for all three species, most genes shared between these two assembly

types were identical or nearly identical at the nucleotide level (see Table 3, Figure 4,

Supplementary Figures 13–15). The biggest differences were attributable to pairwise

alignments containing dramatic length variants, as evidenced by high average percent

identities where aligned (i.e., >99.5% across species; Table 3) and comparably lower

average percent identities (i.e., >97.4% across species; Table 3), which take into account

difference caused by Indels. Given that most alignments were identical or nearly identical,

the dramatic length variants observed between MinION data only and Illumina-polished

assemblies are more than likely the result of inconsistencies in the way BUSCO identifies

genes.

Assembly accuracy is a complex metric that is challenging to assess for large

eukaryotic genomes, and none of the methods utilized here to measure accuracy are

alone sufficient assessments. In combination, however, the results of these methods

suggest that when working with Q20+ MinION data for whole genome *de novo* assembly,

the added value of short-read data comes more measurably from improvements to

contiguity rather than accuracy. Increased contiguity of hybrid assemblies over long read-

only assemblies has been observed in bacteria whole genome sequencing (George et

al., 2017; Neal-McKinney et al., 2021), but to our knowledge has not been previously

reported for eukaryotes. It is worth noting that the long reads generated for each species

here did not approach the read lengths that MinION devices are capable of sequencing.

Across species, read length N50 values ranged from ~2.7–8.6 kb (Supplementary Table

1), which are relatively small values. This is likely in part due to the fact that generating

data from low-input samples, as was done here for single or larval worms, precludes

rigorous size selection. It is possible that the generation of ultra-long reads would have

lessened the gap in contiguity metrics between MinION data and hybrid assemblies.It

should also be noted that generating short-read data in addition to long-read data also

has the added benefit of allowing for the use of k-mer based approaches to estimate

genome size and heterozygosity and to assess the effectiveness of purging haplotypic

duplication.

Results from miniBUSCO indicate a high proportion of missingness (i.e., >40%)

for all trichurid assemblies (see Table 2; Figure 3). Why trichurids appear to be missing

nearly half of nematode BUSCOs is not well understood. Completeness of the existing

WormBase ParaSite reference assemblies for *T. trichiura* and *T. suis* were previously

assessed using the Core Eukaryotic Genes Mapping Approach (i.e., CEGMA) (Parra et

al., 2007), which indicated much higher rates of completeness for these assemblies (i.e.,

~94–96%) (Foth et al., 2014; Jex et al., 2014). However, the creators of CEGMA have

since acknowledged BUSCO as a superior tool for assessing assembly completeness

(Bradnam, 2015). Its origins aside, this presumed genome reduction clearly presents a

challenge for assessing trichurid genome assemblies when using traditional

completeness metrics.

Conclusions

This study represents the first comprehensive comparison of Q20+ MinION data

versus hybrid assembly approaches for generating complete genomes from parasitic

nematodes, and from eukaryotes more broadly. Traditionally, hybrid approaches—

including both incorporating short- and long-read in assembly and polishing long-read

assemblies with short-read data—have been shown to improve assembly-level accuracy.

With the advent of improved read-level quality of Q20+ MinION data, however, it appears

that, at least in this study, where the generation of ultra-long reads was not possible, the

more evident benefit of a hybrid approach was improved contiguity rather than improved

assembly accuracy. Our pairwise alignment analysis of shared genes suggests that most

differences observed between assembly types result from differences in assembly

algorithms or BUSCO implementation rather than significant errors in read-level MinION

data that systematically affect final genome assemblies.

This study was of course met with limitations. For two of the three focal species,

gDNA was generated from adult worms. For most species, and particularly those that

infect humans, adult nematodes are difficult or impossible to obtain, somewhat limiting

the utility of this approach in practice. And although we highlight the potential utility of

Q20+ MinION data for population-level analyses, it was outside the scope of this study to

benchmark MinION data for population genomics in eukaryotes. Finally, the assemblies

generated here remain admittedly incomplete in terms of the modern reference genome

standards. They do not approach chromosome-scale, and they are not annotated, and

additional data types are required to complete them. Generating complete reference

assemblies was not the goal of this study, however; rather, we sought to evaluate the

feasibility and quality of whole genome assemblies generated from an accessible and

inexpensive sequence data type. For *B. malayi* and *T. trichiura*, existing reference

genomes were already exceptionally high-quality, "tier 1"-status assemblies *sensu* the

International Helminth Genomes Consortium Consortium (2019), and the MinION data

and hybrid assemblies generated in general matched them in terms of comparative

quality metrics. The comparison in which the added value of Q20+ MinION data is most

evident is that of *A. caninum*. Using gDNA from the same strain and sample type on which

the existing reference assembly was based, we were able to considerably improve the

genomic resources available for this species relatively easily and inexpensively. These

results highlight the generation of Q20+ MinION data as a simple and cost-effective way

to both shift existing reference genomes for parasitic nematodes into "tier 1" status, as

well to generate *de novo* whole genome assemblies.

In sum, we demonstrated that it is reasonably straightforward to generate high-

quality "tier 1" genomes for a variety of species of parasitic nematodes using Q20+

MinION data alone. Additionally, for species of sufficient size and tractable genome

length, the optimized DNA extraction and library preparation protocols outlined here allow

for the generation of complete genomes from individual specimens. More broadly, the

quality and accessibility of Q20+ MinION data have the potential to make generating real-

time genomic data a routine part of control program monitoring and surveillance,

ultimately accelerating progress toward disease elimination.

Materials and Methods

<u>Specimen acquisition</u>

The specimens sequenced were obtained from the same strains previously used

to generate existing reference genomes for each species. Adults of *B. malayi* (FR3 strain)

were acquired through the NIH/NIAID Filariasis Research Reagent Resource Center via

BEI Resources (Michalski et al., 2011). Worms were received frozen and stored at -80°C.

Prior to sequencing, worms were thawed at 4°C and subsequently preserved in DNA/RNA

Shield (Zymo Research, Irvine, CA, USA). Adults of *T. trichiura* were obtained from a

Danish patient after anthelmintic treatment for an infection acquired in Uganda. Worms

were received preserved in 70% ethanol and were stored at 4°C prior to sequencing.

Pooled third stage (L3) larvae of *A. caninum* (Baltimore strain) were obtained from

experimental infection in laboratory-maintained canines. Pooled L3s were received

frozen, stored at -80°C, and thawed at 4°C prior to sequencing.

<u>DNA extraction</u>

Total nucleic acid was extracted separately from each of a single adult female (*B.*

malayi), a single adult male (*T. trichuris*), and a pool of L3 larvae (*A. caninum*) using a

Quick-DNA™ HMW MagBead Kit (Zymo Research, Irvine, CA, USA) and modified Solid

Tissue extraction protocol. Prior to extraction, adult worms were placed in clean 1.5 mL

DNA LoBind® tubes (Eppendorf® North America, Enfield, CT, USA) using sterilized

forceps, while pooled L3s were centrifuged at 6,540 *g* for 3 min to allow for the removal

of excess supernatant. All samples were then homogenized mechanically using a

Monarch® Pestle Set single-use microtube pestle (New England Biolabs® Inc., Ipswich,

MA, USA). Homogenization was performed both prior to, and immediately following, the

addition of Proteinase K. Samples were then digested in a digital dry bath at 55°C for 24

hr with occasional flick mixing. The protocol for DNA purification was then followed, with

these modifications: After the addition of 33 µL of MagBinding Beads, samples were

placed on a benchtop rotating mixer for 40 min to 1.5 hr at RT; and after the addition of

52 µL DNA Elution Buffer, samples were incubated in a digital dry bath at 37°C for 2 hr

with occasional flick mixing, then incubated at RT overnight prior to collecting eluted DNA.

To avoid DNA fragmentation, samples were mixed by flicking rather than pipette mixing

wherever possible. The concentration of each extraction was measured using a Qubit4™

1X dsDNA High Sensitivity (HS) Assay (ThermoFisher Scientific, Waltham, MA, USA) and

fragment size distribution was determined using an Agilent 2200 TapeStation System and

associated protocol for genomic DNA ScreenTape analysis (Agilent, Santa Clara, CA,

USA). For *T. trichiura*, TapeStation analysis was not performed.

For *A. caninum*, initial attempts to sequence ONT libraries resulted in low

sequencing efficiency, poor data quality, and rapid pore loss. To ameliorate concerns of

protein contamination, extracted gDNA was purified via bead-based cleanup prior to final

ONT library preparation. This cleanup was performed as follows: Two aliquots of

extracted gDNA were each bound to a 0.5× volume of AMPure XP beads (Beckman

Coulter, Brea, CA, USA) on a benchtop rotator mixer for 1 hr at RT. Beads were then

washed twice with 80% EtOH, air-dried, and gDNA was eluted in 52 µL Zymo DNA Elution

Buffer for 2 hr at 37°C with occasional flick mixing, followed by overnight elution at RT.

This cleanup process was repeated a second time for both aliquots. Purified extractions

were then quantified using a Qubit4™ 1X dsDNA High HS Assay and the Agilent 2200

TapeStation System.

<u>MinION library preparation</u>

Aliquots of gDNA extracted for each species were used to prepare one or more

libraries for whole genome sequencing on an ONT MinION desktop sequencer (see

Supplementary Table 1). Libraries were prepared using a combination of the DNA Clean

& Concentrator MagBead (Zymo Research, Irvine, CA, USA) and ONT SQK-LSK114

Ligation Sequencing (Oxford Nanopore Technologies, Oxford, United Kingdom) Kits, and

a modified hybrid protocol. First, gDNA was incubated with ONT DNA repair and end

preparation reagents for 10 min at RT followed by 10 min at 65°C. End preparation

reactions were then bound to 20 µL Zymo MagBinding Beads in 4× volumes of Zymo

DNA MagBinding Buffer on a benchtop rotating mixer for 30 min to 2 hr. Beads were then

washed twice with Zymo DNA Wash Buffer, air-dried for 10 min, eluted in 51 µL Zymo

DNA Elution Buffer via manual agitation for 10 min, and quantified via Qubit4™ 1X dsDNA

HS Assay. Sequencing adaptors were then ligated via a 15 min incubation at RT, and

libraries were bound to a 0.4× volumes of AMPure XP beads on a benchtop rotator mixer

at RT for 1 hr. Beads were then washed twice with either ONT Long Fragment Buffer (*A.*

caninum) or a titrated wash mix of 1:3 ONT Short Fragment Buffer:ONT Long Fragment

Buffer (*B. malayi* and *T. trichiura*). After washing, beads were air-dried, then libraries were

eluted in 15–17 µL of ONT Elution Buffer for 2 hr at 37°C with occasional flick mixing

followed by overnight elution at RT. Final libraries were quantified using a Qubit4™ 1X

dsDNA High HS Assay and the Agilent 2200 TapeStation System. For *T. trichiura*,

TapeStation analysis was not performed. Additionally for *T. trichiura*, a portion of ONT

library reserved after sequencing on one flow cell was re-washed prior to sequencing on

a second flow cell (see Supplementary Table 1). This aliquot of library was bound to a

0.4× volume of AMPure XP beads on a benchtop rotator mixer for 1 hr, washed twice with

ONT Long Fragment Buffer, air-dried, then eluted in 17 µL of ONT Elution Buffer for 2 hr

at 37°C with occasional flick mixing followed by overnight elution at RT.

<u>MinION sequencing, basecalling, and read data quality control</u>

Portions of ONT libraries were sequenced for ~62–80 hr each on one (*B. malayi*),

two (*T. trichiura*), or three (*A. caninum*) ONT MinION R10.4.1 flow cells. The amount of

library loaded onto each flow cell and the total number of pores available at the start of

sequencing are provided in Supplementary Table 1. MinKNOW software (ONT) v. 22.10.7

(*B. malayi*), v. 22.10.10 (*T. trichiura*), or v. 22.12.7 (*A. caninum*) was used to run each

flow cell with pore scans every 1.5 hr. After sequencing, signal data (i.e., fast5 files) from

each flow cell were basecalled using Guppy v. 6.3.4 (ONT). Sequencing adaptors were

simultaneously removed by specifying the "--trim_adapters" flag. The results of each

Guppy run were summarized using NanoPlot v. 1.40.2 (De Coster & Rademakers, 2023).

Fastq files that passed basecalling were input to Mash v. 2.2.2 (Ondov et al., 2016) to

confirm there was no significant contamination in the read-level data prior to assembly.

Estimated genome sizes provided to Mash were based on the lengths of the existing

reference assemblies available for each species, and were thus set as 88 Mb for *B. malayi*

(Ghedin et al., 2007), 465 Mb for *A. caninum* (Consortium, 2019) and 75 Mb for *T. trichiura*

(https://github.com/stephenrdoyle/ancient_trichuris/tree/master/02_data) (Doyle et al.,

2022). Average depth of coverage of reads across the species-specific reference genome

and proportion of reads mapped were estimated using Minimap2 v. 2.16 (Li, 2018) and

SAMtools v. 1.9 (Li et al., 2009).

<u>Illumina library preparation, sequencing, and read data quality control</u>

Aliquots of gDNA extracted for each species were sent to the Yale Center for

Genome Analysis (YCGA) for Illumina whole genome library preparation and sequencing

(see Supplementary Table 1). Libraries were prepared using an xGenTM cfDNA & FFPE

DNA Library Prep Kit with unique dual indexing (Integrated DNA Technologies, Newark,

NJ, USA) and standard protocol. Libraries were subjected to 5–7 cycles of PCR to

increase concentration (see Supplementary Table 1) and multiplexed for 2×150 paired-

end sequencing on an Illumina NovaSeq 6000 targeting 50× depth of coverage genome-

wide for each species. Reads were demultiplexed by the YCGA and subsequently filtered

for remaining adaptor contamination, quality, and length using fastp (Chen et al., 2018).

Fastq files that passed fastp were input to Mash to confirm there was no significant

contamination in the read-level data prior to assembly. Estimated genome sizes provided

to Mash were based on the lengths of the existing reference assemblies available for

each species (see above). Average depth of coverage across the species-specific

reference genome and proportion of reads mapped were estimated using BWA v. 0.7.17

and SAMtools v. 1.9 (Li et al., 2009). A second round of fastp was run on the raw

demultiplexed Illumina data to remove sequencing adapters, only, by specifying

"--detect_adapter_for_pe", "--disable_length_filtering", and "--disable_quality_filtering".

These data were the used for hybrid genome assembly, as MaSuRCA utilizes built-in

error correction and cleaning.

<u>Estimation of genome size, heterozygosity, and ploidy</u>

Quality-controlled Ilumina data were used to estimate genome size and

heterozygosity for each species. First, Jellyfish v. 2.3.0. (Marçais & Kingsford, 2011) was

used to generate k-mer spectra, specifying a k-mer size of 21 and an initial hash size of

1,000,000,000. Histograms from Jellyfish were then provided to the GenomeScope web-

based graphical user interface (Vurture et al., 2017) to visualize k-mer spectra and

estimate heterozygosity. For *T. trichiura*, ploidy was estimated bioinformatically using

quality-controlled Ilumina data and basecalled MinION data in conjunction with nQuire

(Weiß et al., 2018).

<u>MinION data genome assembly</u>

Fastq files that passed basecalling via Guppy were concatenated and used as

input to Canu v. 2.1 (ONT) (Koren et al., 2017) to generate MinION data assemblies. For

T. trichiura and *A. caninum*, a 3 kb minimum read length requirement was specified, and

for *B. malayi*, a 5 kb minimum read length requirement was specified. Estimated genome

sizes provided to Canu were based on the lengths of the existing reference assemblies

available for each species (see above). Contigs in the resulting Canu assemblies that

were indicated as potential alternative alleles (i.e., with FASTA headers including

"suggestBubble=yes") were removed using a combination of SeqKit v 0.10.1 (Shen et al.,

2016) and the filterbyname.sh script of BBMap v.38.84 (Bushnell, 2014). Resulting

"popped" assemblies were used as input to purge_dups v. 1.2.5 (Guan et al., 2020) to

remove false duplications. These "popped and purged" assemblies were then polished

with one round of Racon v. 1.5.0 (Vaser et al., 2017) followed by one round of Medaka v.

1.7.2 (https://github.com/nanoporetech/medaka) to generate "popped, purged, and

polished" assemblies.

Hybrid genome assembly

FASTQ files that passes basecalling by Guppy and Illumina data from which only

sequencing adaptors were removed by fastp were used as input to MaSuRCA v. 4.1.0

(Zimin et al., 2013) to generate hybrid assemblies. Default settings were used except for

setting "LHE_COVERAGE" to "35" and "cgwErrorRate=0.15", and setting "JF_SIZE" to a

value ten times the genome size of the species-specific reference assembly. Assemblies

output by MaSuRCA were input to purge_dups v. 1.2.5 (Guan et al., 2020) to remove

false duplications.

Quality control and refinement of final assemblies

To ensure "purged" MaSuRCA hybrid assemblies were not over- or under-purged,

copy number spectrum plots were generated using the K-mer Analysis Toolkit (KAT)

v. 2.4.0 (Mapleson et al., 2017) specifying a k-mer size of 31. "Popped, purged, and

polished" Canu MinION data assemblies and "purged" MaSuRCA hybrid assemblies were

input to BlobTools v. 1.1.1 (Laetsch & Blaxter, 2017) to identify potential contaminants.

First, the reads used to generate each assembly were mapped to the assembly using

Minimap2 (MinION data) and/or BWA (Illumina data), and SAMtools was used to generate

sorted BAM files for each mapping. The "blastn" function in BLAST v. 2.14.0 (Altschul et

al., 1990) was then used to compare all contigs to reference sequences available in the

National Center for Biotechnology Information (NCBI)'s nucleotide databases, specifying

"-evalue 1e-25", "-max_target_seqs 10" and "-max_hsps 1". BLAST outputs and sorted

BAM files were then used as input to BlobTools. Contigs that could be confirmed as host,

human, or bacterial contamination were excluded from final assemblies. Quality scores

(i.e., Q scores) of basecalled MinION reads, and Q scores of reads when mapped to the

final "popped, purged, and polished" MinION data assemblies, were measured using

NanoPlot and plotted using the "geom_density" function in the ggplot2 package

(Wickham, 2016) in R v. 4.3.0 (Team, 2023) in RStudio v. 2023.03.1 (Team, 2020).

<u>Mitogenome and *Wolbachia* genome assembly</u>

For *B. malayi* and *A. caninum* contigs containing the mitogenomes—and, for *B.*

malayi, the genome of its *Wolbachia* endosymbiont—were extracted from "popped"

MinION and unpurged MaSuRCA assemblies. For *T. trichiura*, mitogenomes were

extracted from the original "unpopped" Canu assembly and the unpurged MaSuRCA

assembly. These organelle genomes contigs were then mapped to species-specific

reference organelle genomes using Geneious v. 2022.0.1 ("Geneious Prime v.

2022.0.1,") and manually reconfigured to match the linear orientation of the reference.

For *B. malayi* and *T. trichiura*, reference organelle genomes were extracted from the

reference genome assemblies for each species (see above). For *A. caninum*, for which a

mitochondrial contig is not labeled in the reference assembly, a reference mitogenome

was downloaded from GenBank (accession no. MN215971) (Xie et al., 2019). Contigs

containing complete organelle genomes were then manually added to the appropriate

final version of each assembly.

<u>BUSCO comparison to assess assembly accuracy</u>

Copies of final "popped, purged and polished" MinION data assemblies generated for

each species (i.e., assemblies from which potential contaminants identified by BlobTools

had been removed, and to which organelle genomes had been added) were separately

polished with Illumina data using Pilon v. 1.24 (Walker et al., 2014) and BWA. Three

iterations of Pilon were iteratively run specifying the "--diploid" flag. BUSCO v. 4.0.6

(Seppey et al., 2019) was then run for MinION data, polished, and hybrid assemblies for

each   species   specifying   comparison   to   the   Nematoda   ortholog   database

(nematoda_odb10). Two single copy ortholog datasets were generated per species from

the resulting BUSCO output: one containing sequences found in both the MinION data

and the polished assembly, and one containing sequences found in both the MinION data

and the hybrid assembly. For each dataset and species, the Genes shared between the

two assembly types were pairwise aligned using the "pairwiseAlignment" function in

Biostrings package v. 2.66.0 (Pages et al., 2013) in R v. 4.2.3 via RStudio v. 2023.06.1

specifying the alignment type as "global". Biostrings and custom R code were then used

to retain only pairwise alignments in which Genes from both assemblies started and

ended with eukaryotic start and stop codons, respectively. Biostrings was then used to

characterize each pairwise alignment in terms of percent identity, alignment length,

BUSCO lengths, and number of single nucleotide polymorphisms (i.e., SNPS) and

insertions/deletions (i.e., Indels). Summary figures for all comparisons were generated in

R v. 4.3.0 via RStudio v. 2023.03.1 using the ggplot2 package.

<u>Assessing assembly contiguity, accuracy, and completeness</u>

Contiguity and accuracy were respectively assessed by comparing the MinION

data assembly to hybrid and reference assemblies using QUAST v. 5.0.2 (Gurevich et

al., 2013), and to the hybrid assembly using the "nucmer" and "dnadiff" modules of

MuMmer4 v. 4.0.0beta2 (Marçais et al., 2018). To assess assembly completeness,

miniBUSCO v. 0.2 (Huang & Li, 2023) was run on all three final assemblies (i.e., MinION

data, MinION data polished with Illumina data, and hybrid) generated for each species,

specifying comparison to the Nematoda ortholog database (nematoda_odb10). For

comparison, miniBUSCO was also run on the aforementioned reference genomes for

each species, as well as for all genomes of nematodes that parasitize animal as adults

available from WormBase ParaSite (Howe et al., 2016; Howe et al., 2017). For each

assembly, the proportion of single copy BUSCOs present was plotted against the log

value of the assembly's N50 length divided by its scaffold or contig count using R v. 4.2.3

via RStudio v. 2023.06.1. Cutoff values of >85% (for proportion of single copy BUSCOs

present) and >1.6 (for the log value of the contiguity metric) were used to assess whether

each assembly could be classified as a "tier 1" genome following the standard for helminth

genomes established by the International Helminth Genomes Consortium (Consortium,

2019).

Acknowledgements

The authors thank the BEI Resources NIH/NIAID Filariasis Research Reagent Resource Center for providing specimens of *Brugia malayi* for sequencing. Thanks are also due to Dr. Michael Wiley, Dr. Shaun Cross, Kristen Bernhard, and Mahmood Al-Haehm (University of Nebraska Medical Center) for providing access and support for use of select laboratory equipment. The authors additionally thank Christopher Castaldi and Irina Tikhonova (Yale Center for Genome Analysis, Yale School of Medicine) for generation of, and correspondence regarding, Illumina sequence data. Bioinformatic analysis was enabled by the University of Nebraska-Lincoln Holland Computing Center computing cluster.

Author Contributions

KSH and JRF designed the study. JMH provided the specimens of *Ancylostoma caninum* sequenced. PN provided the specimen of *Trichuris trichiura* sequenced. KSH extracted genomic DNA and prepared and sequenced MinION libraries. KSH and JRF determined and refined bioinformatic pipelines. KSH performed bioinformatic analyses. RW aided KSH in the generation of comparative BUSCO data. KSH and JRF interpreted data and generated figures and tables. KSH and JRF drafted the manuscript. All authors contributed to revising and approving the final version of the manuscript.

Literature Cited

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. 1990. Basic local alignment

search tool. Journal of Molecular Biology, 215(3), 403–410.

Bashir, A., Klammer, A. A., Robins, W. P., Chin, C.-S., Webster, D., Paxinos, E., Hsu, D., Ashby, M.,

Wang, S., & Peluso, P. 2012. A hybrid approach for the automated finishing of bacterial

genomes. Nature Biotechnology, 30(7), 701–707. DOI: 10.1038/nbt.2288.

Berger, D. J., Crellen, T., Lamberton, P. H., Allan, F., Tracey, A., Noonan, J. D., Kabatereine, N. B.,

Tukahebwa, E. M., Adriko, M., & Holroyd, N. 2021. Whole-genome sequencing of

Schistosoma mansoni reveals extensive diversity with limited selection despite mass

drug administration. Nature Communications, 12(1), 4776. DOI: 10.1038/s41467-021-

24958-0

Biraro, I. A., Egesa, M., Toulza, F., Levin, J., Cose, S., Joloba, M., Smith, S., Dockrell, H. M.,

Katamba, A., & Elliott, A. M. 2014. Impact of co-infections and BCG immunisation on

immune responses among household contacts of tuberculosis patients in a Ugandan

cohort. PLOS One, 9(11), e111517. DOI: 10.1371/journal.pone.0111517.

Brabec, J., Kuchta, R., Scholz, T., & Littlewood, D. T. J. 2016. Paralogues of nuclear ribosomal

genes conceal phylogenetic signals within the invasive Asian fish tapeworm lineage:

Evidence from next generation sequencing data. International Journal for Parasitology,

46(9), 555–562. DOI: 10.1016/j.ijpara.2016.03.009.

Bradnam, K. (2015). Goodbye CEGMA, hello BUSCO!

Bushnell, B. 2014. *BBMap: a fast, accurate, splice-aware aligner*. Lawrence Berkeley National

Lab.(LBNL), Berkeley, CA (United States). Retrieved from.

Casulli, A. 2021. New global targets for NTDs in the WHO roadmap 2021–2030. PLOS Neglected

Tropical Diseases, 15(5), e0009373. DOI: 10.1371/journal.pntd.0009373.

Chen, S., Zhou, Y., Chen, Y., & Gu, J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics, 34(17), i884–i890. DOI: 10.1093/bioinformatics/bty560.

Consortium, I. H. G. 2019. Comparative genomics of the major parasitic worms. Nature

Genetics, 51(1), 163–174. DOI: 10.1038/s41588-018-0262-1.

De Coster, W., & Rademakers, R. 2023. NanoPack2: population-scale evaluation of long-read

sequencing data. Bioinformatics, 39(5), btad311. DOI: 10.1093/bioinformatics/btad311.

Doyle, S. Improving helminth genome resources in the post-genomic era. Trends in

Parasitology, 38(10), 831–840. DOI: 10.1016/j.pt.2022.06.002.

Doyle, S. R., Søe, M. J., Nejsum, P., Betson, M., Cooper, P. J., Peng, L., Zhu, X.-Q., Sanchez, A.,

Matamoros, G., & Sandoval, G. A. F. 2022. Population genomics of ancient and modern

Trichuris trichiura. Nature Communications, 13(1), 3888. DOI: 10.1038/s41467-022-

31487-x.

English, A. C., Richards, S., Han, Y., Wang, M., Vee, V., Qu, J., Qin, X., Muzny, D. M., Reid, J. G., &

Worley, K. C. 2012. Mind the gap: upgrading genomes with Pacific Biosciences RS long-

read sequencing technology. PLOS One, 7(11), e47768. DOI:

10.1371/journal.pone.0047768.

Fierst, J. L., Willis, J. H., Thomas, C. G., Wang, W., Reynolds, R. M., Ahearne, T. E., Cutter, A. D.,

& Phillips, P. C. 2015. Reproductive mode and the evolution of genome size and

structure in *Caenorhabditis* nematodes. PLoS Genetics, 11(6), e1005323. DOI:

10.1371/journal.pgen.1005323.

Formenti, G., Theissinger, K., Fernandes, C., Bista, I., Bombarely, A., Bleidorn, C., Ciofi, C.,

Crottini, A., Godoy, J. A., & Höglund, J. 2022. The era of reference genomes in

conservation genomics. Trends in Ecology & Evolution, 37(3), 197–202. DOI:

10.1016/j.tree.2021.11.008.

Foth, B. J., Tsai, I. J., Reid, A. J., Bancroft, A. J., Nichol, S., Tracey, A., Holroyd, N., Cotton, J. A.,

Stanley, E. J., & Zarowiecki, M. 2014. Whipworm genome and dual-species

transcriptome analyses provide molecular insights into an intimate host-parasite

interaction. Nature Genetics, 46(7), 693–700. DOI: 10.1038/ng.3010.

Fraija-Fernández, N., Waeschenbach, A., Briscoe, A. G., Hocking, S., Kuchta, R., Nyman, T., &

Littlewood, D. T. J. 2021. Evolutionary transitions in broad tapeworms (Cestoda:

Diphyllobothriidea) revealed by mitogenome and nuclear ribosomal operon

phylogenetics. Molecular Phylogenetics and Evolution, 163(107262), 11 pp. DOI:

10.1016/j.ympev.2021.107262.

Fu, Z., Li, Y., Elling, A. A., & Snyder, W. E. 2020. A draft genome of a field-collected *Steinernema*

feltiae strain NW. Journal of Nematology, 52(1), 1–7. DOI: 10.21307/jofnem-2020-003.

Geneious Prime v. 2022.0.1. https://www.geneious.com.

George, S., Pankhurst, L., Hubbard, A., Votintseva, A., Stoesser, N., Sheppard, A. E., Mathers, A.,

Norris, R., Navickaite, I., & Eaton, C. 2017. Resolving plasmid structures in

Enterobacteriaceae using the MinION nanopore sequencer: assessment of MinION and

MinION/Illumina hybrid data assembly approaches. Microbial Genomics, 3(8), e000118.

DOI: 10.1099/mgen.0.000118.

Ghedin, E., Wang, S., Spiro, D., Caler, E., Zhao, Q., Crabtree, J., Allen, J. E., Delcher, A. L.,

Guiliano, D. B., & Miranda-Saavedra, D. 2007. Draft genome of the filarial nematode

parasite *Brugia malayi*. Science, 317(5845), 1756–1760. DOI: 10.1126/science.1145406.

The Global Burden of Disease Study 2019. 2020. The Lancet, 396(10258), 1129–1306.

Goswami, U. 1978. Some cytogenetical aspects of genus *Trichuris* (Nematoda). Current Science,

47(11), 368–371.

Guan, D., McCarthy, S. A., Wood, J., Howe, K., Wang, Y., & Durbin, R. 2020. Identifying and

removing haplotypic duplication in primary genome assemblies. Bioinformatics, 36(9),

2896–2898. DOI: 10.1093/bioinformatics/btaa025.

Gurevich, A., Saveliev, V., Vyahhi, N., & Tesler, G. 2013. QUAST: quality assessment tool for

genome assemblies. Bioinformatics, 29(8), 1072–1075. DOI:

10.1093/bioinformatics/btt086.

Hoban, S., Kelley, J. L., Lotterhos, K. E., Antolin, M. F., Bradburd, G., Lowry, D. B., Poss, M. L.,

Reed, L. K., Storfer, A., & Whitlock, M. C. 2016. Finding the genomic basis of local

adaptation: pitfalls, practical solutions, and future directions. The American Naturalist,

188(4), 379–397. DOI: 10.1086/688018.

Howe, K., Chow, W., Collins, J., Pelan, S., Pointon, D.-L., Sims, Y., Torrance, J., Tracey, A., &

Wood, J. 2021. Significantly improving the quality of genome assemblies through

curation. Gigascience, 10(1), giaa153. DOI: 10.1093/gigascience/giaa153.

Howe, K. L., Bolt, B. J., Cain, S., Chan, J., Chen, W. J., Davis, P., Done, J., Down, T., Gao, S., &

Grove, C. 2016. WormBase 2016: expanding to enable helminth genomic research.

Nucleic Acids Research, 44(D1), D774–D780. DOI: 10.1093/nar/gkv1217.

Howe, K. L., Bolt, B. J., Shafie, M., Kersey, P., & Berriman, M. 2017. WormBase ParaSite– a

comprehensive resource for helminth genomics. Molecular and Biochemical

Parasitology, 215, 2–10. DOI: 10.1016/j.molbiopara.2016.11.005.

Huang, N., & Li, H. 2023. miniBUSCO: a faster and more accurate reimplementation of BUSCO.

bioRxiv, 2023.2006. 2003.543588. DOI: 10.1101/2023.06.03.543588.

Jex, A. R., Nejsum, P., Schwarz, E. M., Hu, L., Young, N. D., Hall, R. S., Korhonen, P. K., Liao, S.,

Thamsborg, S., & Xia, J. 2014. Genome and transcriptome of the porcine whipworm

Trichuris suis. Nature Genetics, 46(7), 701–706. DOI: 10.1038/ng.3012.

Kizito, D., Tweyongyere, R., Namatovu, A., Webb, E. L., Muhangi, L., Lule, S. A., Bukenya, H.,

Cose, S., & Elliott, A. M. 2013. Factors affecting the infant antibody response to measles

immunisation in Entebbe-Uganda. BMC Public Health, 13(619), 1–9. DOI: 10.1186/1471-

2458-13-619.

Koren, S., Schatz, M. C., Walenz, B. P., Martin, J., Howard, J. T., Ganapathy, G., Wang, Z., Rasko,

D. A., McCombie, W. R., & Jarvis, E. D. 2012. Hybrid error correction and *de novo*

assembly of single-molecule sequencing reads. Nature Biotechnology, 30(7), 693–700.

DOI: 10.1038/nbt.2280.

Koren, S., Walenz, B. P., Berlin, K., Miller, J. R., Bergman, N. H., & Phillippy, A. M. 2017. Canu:

scalable and accurate long-read assembly via adaptive k-mer weighting and repeat

separation. Genome Research, 27(5), 722–736. DOI: 10.1101/gr.215087.116.

Laetsch, D. R., & Blaxter, M. L. 2017. BlobTools: Interrogation of genome assemblies.

F1000Research, 6(1287). DOI: 10.12688/f1000research.12232.1.

Li, H. 2018. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics, 34(18),

3094–3100. DOI: 10.1093/bioinformatics/bty191.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,

Durbin, R., & Subgroup, G. P. D. P. 2009. The sequence alignment/map format and

SAMtools. Bioinformatics, 25(16), 2078-2079. DOI: 10.1093/bioinformatics/btp352.

Ludington, A. J., & Sanders, K. L. 2021. Demographic analyses of marine and terrestrial snakes

(Elapidae) using whole genome sequences. Molecular Ecology, 30(2), 545–554. DOI:

10.1111/mec.15726.

Mapleson, D., Garcia Accinelli, G., Kettleborough, G., Wright, J., & Clavijo, B. J. 2017. KAT: a K-

mer analysis toolkit to quality control NGS datasets and genome assemblies.

Bioinformatics, 33(4), 574–576. DOI: 10.1093/bioinformatics/btw663.

Marçais, G., Delcher, A. L., Phillippy, A. M., Coston, R., Salzberg, S. L., & Zimin, A. 2018.

MUMmer4: A fast and versatile genome alignment system. PLoS Computational Biology,

14(1), e1005944. DOI: 10.1371/journal.pcbi.1005944.

Marçais, G., & Kingsford, C. 2011. A fast, lock-free approach for efficient parallel counting of

occurrences of k-mers. Bioinformatics, 27(6), 764–770. DOI:

10.1093/bioinformatics/btr011.

Michalski, M. L., Griffiths, K. G., Williams, S. A., Kaplan, R. M., & Moorhead, A. R. 2011. The NIH-

NIAID filariasis research reagent resource center. PLOS Neglected Tropical Diseases,

5(11), e1261. DOI: 10.1371/journal.pntd.0001261.

Morawski, B. M., Yunus, M., Kerukadho, E., Turyasingura, G., Barbra, L., Ojok, A. M., DiNardo, A.

R., Sowinski, S., Boulware, D. R., & Mejia, R. 2017. Hookworm infection is associated

with decreased CD4+ T cell counts in HIV-infected adult Ugandans. PLOS Neglected

Tropical Diseases, 11(5), e0005634. DOI: 10.1371/journal.pntd.0005634.

Nash, S., Mentzer, A. J., Lule, S. A., Kizito, D., Smits, G., van der Klis, F. R., & Elliott, A. M. 2017.

The impact of prenatal exposure to parasitic infections and to anthelminthic treatment

on antibody responses to routine immunisations given in infancy: Secondary analysis of

a randomised controlled trial. PLOS Neglected Tropical Diseases, 11(2), e0005213. DOI:

10.1371/journal.pntd.0005213.

Ndyomugyenyi, R., Kabatereine, N., Olsen, A., & Magnussen, P. 2008. Malaria and hookworm

infections in relation to haemoglobin and serum ferritin levels in pregnancy in Masindi

district, western Uganda. Transactions of the Royal Society of Tropical Medicine and

Hygiene, 102(2), 130–136. DOI: 10.1016/j.trstmh.2007.09.015.

Neal-McKinney, J. M., Liu, K. C., Lock, C. M., Wu, W.-H., & Hu, J. 2021. Comparison of MiSeq,

MinION, and hybrid genome sequencing for analysis of *Campylobacter jejuni*. Scientific

Reports, 11(1), 5676. DOI: 10.1038/s41598-021-84956-6.

Okenmor, G. A., Kutshik, R. J., & Longdet, I. Y. 2022. Draft genome sequence of *Lysinibacillus* sp.

strain BPa_S21, isolated from petroleum-contaminated soil in Delta State, Nigeria.

Microbiology Resource Announcements, 11(8), e00456-00422. DOI:

10.1128/mra.00456-22.

Ondov, B. D., Treangen, T. J., Melsted, P., Mallonee, A. B., Bergman, N. H., Koren, S., & Phillippy,

A. M. 2016. Mash: fast genome and metagenome distance estimation using MinHash.

Genome Biology, 17(132), 1–14. DOI: 10.1186/s13059-016-0997-x.

Organization, W. H. 2021. Ending the neglect to attain the Sustainable Development Goals: A

road map for neglected tropical diseases 2021–2030 [Internet].

Pages, H., Aboyoun, P., Gentleman, R., DebRoy, S., Pages, M. H., DataImport, D., BSgenome, S.,

XStringSet-class, R., MaskedXString-class, R., & XStringSet-io, R. 2013. Package

'Biostrings'. Bioconductor, 18129.

Parra, G., Bradnam, K., & Korf, I. 2007. CEGMA: a pipeline to accurately annotate core genes in

eukaryotic genomes. Bioinformatics, 23(9), 1061–1067. DOI:

10.1093/bioinformatics/btm071.

Price, H. J. 1976. Evolution of DNA content in higher plants. The Botanical Review, 42, 27–52.

DOI: 10.1007/BF02860861.

Rhie, A., McCarthy, S. A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., Uliano-Silva, M., Chow,

W., Fungtammasan, A., & Kim, J. 2021. Towards complete and error-free genome

assemblies of all vertebrate species. Nature, 592(7856), 737–746. DOI: 10.1038/s41586-

021-03451-0.

Roessler, K., Muyle, A., Diez, C. M., Gaut, G. R., Bousios, A., Stitzer, M. C., Seymour, D. K.,

Doebley, J. F., Liu, Q., & Gaut, B. S. 2019. The genome-wide dynamics of purging during

selfing in maize. Nature Plants, 5(9), 980–990. DOI: 10.1038/s41477-019-0508-7.

Seppey, M., Manni, M., & Zdobnov, E. M. 2019. BUSCO: assessing genome assembly and

annotation completeness. Gene prediction: methods and protocols(33), 227–245. DOI:

10.1093/bioinformatics/btx153.

Shen, W., Le, S., Li, Y., & Hu, F. 2016. SeqKit: a cross-platform and ultrafast toolkit for FASTA/Q

file manipulation. PLOS One, 11(10), e0163962. DOI: 10.1371/journal.pone.0163962.

Sims, D., Sudbery, I., Ilott, N. E., Heger, A., & Ponting, C. P. 2014. Sequencing depth and

coverage: key considerations in genomic analyses. Nature Reviews Genetics, 15(2), 121–

132. DOI: 10.1038/nrg3642.

Solares, E. A., Tao, Y., Long, A. D., & Gaut, B. S. 2021. HapSolo: an optimization approach for

removing secondary haplotigs during diploid genome assembly and scaffolding. BMC

Bioinformatics, 22(1), 1–17. DOI: 10.1186/s12859-020-03939-y.

Špakulová, M., Kralova, I., & Cutillas, C. 1994. Studies on the karyotype and gametogenesis in

Trichuris muris. Journal of Helminthology, 68(1), 67–72. DOI:

10.1017/S0022149X0001350X.

Spinner, N. B., Conlin, L. K., Mulchandani, S., & Emanuel, B. S. 2013. Deletions and other

structural abnormalities of the autosomes. *In* Emery and Rimoin's Essential Medical

Genetics. D. L. Rimoin, R. E. Pyeritz, & B. Korf (Eds.). Vol. 161. Oxford, UK: Elsevier.

Sweet, A. D., & Johnson, K. P. 2018. The role of parasite dispersal in shaping a host–parasite

system at multiple evolutionary scales. Molecular Ecology, 27(24), 5104–5119. DOI:

10.1111/mec.14937.

Team, R. 2020. RStudio: Integrated Development for R. http://www.rstudio.com/. PBC, Boston,

MA, USA.

Team, R. C. 2023. A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Vaser, R., Sović, I., Nagarajan, N., & Šikić, M. 2017. Fast and accurate de novo genome assembly

from long uncorrected reads. Genome Research, 27(5), 737–746. DOI:

10.1101/gr.214270.116.

Virrueta Herrera, S., Johnson, K. P., Sweet, A. D., Ylinen, E., Kunnasranta, M., & Nyman, T. 2022.

High levels of inbreeding with spatial and host-associated structure in lice of an

endangered freshwater seal. Molecular Ecology, 31(18), 4593–4606. DOI:

10.1111/mec.16569.

Voelker, J., Shepherd, M., & Mauleon, R. 2021. A high-quality draft genome for *Melaleuca*

alternifolia (tea tree): a new platform for evolutionary genomics of myrtaceous terpene-

rich species. GigaByte(gigabyte28). DOI: 0.46471/gigabyte.28.

Vurture, G. W., Sedlazeck, F. J., Nattestad, M., Underwood, C. J., Fang, H., Gurtowski, J., &

Schatz, M. C. 2017. GenomeScope: fast reference-free genome profiling from short

reads. Bioinformatics, 33(14), 2202–2204. DOI: 10.1093/bioinformatics/btx153.

Walker, B. J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., Cuomo, C. A., Zeng,

Q., Wortman, J., & Young, S. K. 2014. Pilon: an integrated tool for comprehensive

microbial variant detection and genome assembly improvement. PLOS One, 9(11),

e112963. DOI: 10.1371/journal.pone.0112963.

Wang, W., Das, A., Kainer, D., Schalamun, M., Morales-Suarez, A., Schwessinger, B., & Lanfear,

R. 2020. The draft nuclear genome assembly of *Eucalyptus pauciflora*: a pipeline for

comparing de novo assemblies. Gigascience, 9(1), giz160. DOI:

10.1093/gigascience/giz160.

Weiß, C. L., Pais, M., Cano, L. M., Kamoun, S., & Burbano, H. A. 2018. nQuire: a statistical

framework for ploidy estimation using next generation sequencing. BMC Bioinformatics,

19(122). DOI: 10.1186/s12859-018-2128-z.

Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis (978-3-319-24277-4,

https://ggplot2.tidyverse.org ed.). Springer-Verlag New York. pp.

Xie, Y., Xu, Z., Zheng, Y., Li, Y., Liu, Y., Wang, L., Zhou, X., Zuo, Z., Gu, X., & Yang, G. 2019. The

mitochondrial genome of the dog hookworm *Ancylostoma caninum* (Nematoda,

Ancylostomatidae) from Southwest China. Mitochondrial DNA Part B, 4(2), 3002–3004.

DOI: 10.1080/23802359.2019.1666048.

Yi, L., Dalai, M., Su, R., Lin, W., Erdenedalai, M., Luvsantseren, B., Chimedtseren, C., Wang, Z., &

Hasi, S. 2020. Whole-genome sequencing of wild Siberian musk deer (*Moschus*

moschiferus) provides insights into its genetic features. BMC Genomics, 21(108), 1–13.

DOI: 10.1186/s12864-020-6495-2.

Yousuf, P., Parvaiz, S., Razzak, S., & Zehbi, N. (Eds.). (2022). *Trichuris: A critical review*:

IntechOpen.

Zimin, A. V., Marçais, G., Puiu, D., Roberts, M., Salzberg, S. L., & Yorke, J. A. 2013. The MaSuRCA

genome assembler. Bioinformatics, 29(21), 2669–2677. DOI:

10.1093/bioinformatics/btt476.