Assessing rapid adaptation through epigenetic inheritance: a new

experimental approach

Alexandra Chávez[1,2,] Meret Huber[1,2*,]

1 Institute of Organismic and Molecular Evolution, University of Mainz, Johann-

Joachim-Becher-Weg 7, 55128 Mainz, Germany

2 Institute of Plant Biology and Biotechnology, University of Münster, Schlossplatz 7,

48143 Münster, Germany

*Correspondence: Meret Huber, Institute of Organismic and Molecular Evolution,

Johann-Joachim-Becher-Weg 7, 55128 Mainz, Germany; meret.huber@uni-

mainz.de

Running headline: Adaption through epigenetic inheritance

Abstract

1. Epigenetic inheritance is hypothesized to mediate rapid adaptation to stresses via two fundamentally different routes: first, through spontaneous epimutations that arise in a largely stochastic manner in the presence or absence of stress; if these spontaneous epimutations are heritable and beneficial, they may be selected upon ("stochastic route"); and second, through environment-induced epialleles that arise uniformly among individuals; if heritable, these epialleles may lead to stress adaptation even in the absence of selection ("deterministic route"). Testing and teasing apart these two routes is challenging, largely because a suitable experimental approach is lacking.

2. Here, we propose an experimental approach that allows to simultaneously assess the contribution of the stochastic and deterministic route. The essence of the approach is to manipulate the efficacy of selection through the population size and thereby to test whether selection is required for adaptation (stochastic route). To this end, genetically uniform populations are grown under different environments across multiple generations ("pre-treatment") at two different population sizes: in large populations, in which selection is effective; and in small populations, in which drift overcomes the effect of selection. If the deterministic route contributes to adaptation, variation in fitness, phenotypes or epigenetic marks will arise between the small populations of the different pre-treatments. If the stochastic route contributes to adaptation, variation will arise between the small and large population within each pre-treatment. As a proof-of-principle, we tested whether small

and large monoclonal populations of the aquatic duckweed *Spirodela*

polyrhiza may adapt to copper excess outdoors.

3. After five to seven generations of pre-treatment and a subsequent multi-

generational growth under control conditions, large populations outperformed

small populations under copper excess. Furthermore, small populations pre-

treated with copper excess tended to have higher fitness under copper excess

than small populations pre-treated under control conditions. These data

suggest that both the stochastic and deterministic route may alter plant fitness

under recurring stress.

4. The proposed approach will allow to experimentally evaluate whether species

may adapt to stresses through either stochastic and deterministic epigenetic

changes, which is fundamental to understand whether and how epigenetic

inheritance may lead to rapid stress adaptation.

Keywords

Epigenetic and non-genetic inheritance, rapid adaptation to stress, transgenerational

inheritance, transgenerational plasticity, deterministic and stochastic epigenetic

variation, experimental evolution, epimutation, duckweed

INTRODUCTION

The hallmark of evolutionary biology states that species adapt to stresses through selection of phenotypic variants that are arise from DNA sequence variation. Yet, increasing evidence shows that DNA sequence variation is not the sole source of heritable phenotypic variation: for instance, flowers of natural toadflax mutants have radial instead of bilateral symmetry because the floral symmetry gene *Lcyc* became hypermethylated (Cubas et al., 1999). Similarly, homeotic floral phenotypes in the oil palm arose during tissue culture because the homeotic gene DEFICIENS is alternatively spliced upon spontaneous hypomethylation of an intron located transposon (Ong-Abdullah et al., 2015). Furthermore, fruit ripening in a spontenous tomato mutant is hampered because the *Cnr* locus was hypermerthylated and thereby silenced (Manning et al., 2006). These examples highlight that mechanisms other than DNA sequence variation may lead to heritable, variable and fitness-relevant phenotypes, and as such they open up an exciting question: what role do mechanisms that generate heritable phenotypic variation in the absence of DNA sequence variation play in the adaptation to environmental stresses? Considering the on-going loss of intra-specific genetic diversity and the rapid pace of environmental change (Sage, 2020; Des Roches et al., 2021), answering this question is becoming an increasingly pressing issue.

Heritable phenotypic variation that is not based on DNA sequence variation may arise through so-called "epigenetic" or "non-genetic" inheritance (Bonduriansky & Day, 2009; O'Dea et al., 2016). Epigenetic inheritance often refers to mechanisms that alter gene expression across generations through genome-associated mechanisms such as DNA methylation, histone modifications and small RNAs, whereas non-genetic inheritance includes any other mechanism such as the vertical

transfer of microbes, nutrients or hormones (Bonduriansky & Day, 2009). To improve

readibility, we here use the term "epigenetic" to refer to both epigenetic and non-

genetic mechanisms; differences among these mechanisms are, however,

discussed.

Many epigenetic marks share two feasures: first, epigenetic marks may

change spontaneously over generations in a stochastic manner (Becker et al., 2011;

Schmitz et al., 2011), and second, epigenetic marks may be induced by the

environment and thereby lead to predictable phenotypic variation (Wibowo et al.,

2016; Zhang et al., 2018). Consequently, epigenetic inheritance may lead to stress

adaptation through two fundamentally different routes: the "stochastic" and the

"deterministic" route (Fig. 1a) (Baugh & Day, 2020).

The "stochastic route" is comparable to the adaptive process that is based on

DNA sequence variation. Here, spontaneous changes ("epimutations") arise either in

the presence or absence of stress in a largely random manner. For instance, in

Arabidopsis thaliana, variations in CG and to a much lower extent also CHG and

CHH methylation (H=A, T, C) steadily accumulate across generations (Becker et al.,

2011; Schmitz et al., 2011; Denkena et al., 2021), and both the frequency and the

spectrum of these spontaneous epimutations may be altered by stress (Jiang et al.,

2014; Johannes & Schmitz, 2019). These spontaneous epimutations are semi-

stable, meaning that many epimutations are inherited across several generations but

also frequently reverse (Johannes & Schmitz, 2019). If epimutations convey fitness

benefits under stress, these variants can be selected upon, leading to stress

adaptation.

The "deterministic route", in contrast, is not comparable to the adaptive

process that is based on DNA sequence variation. In the deterministic route,

environmental stresses induce epigenetic variation and thereby lead to predictable

epigenetic variations that are shared among individuals ("deterministic epialleles").

For instance, in *A. thaliana,* environmental stresses may remodel CHG and CHH

methylation in specific repeat sequences, which is associated with altered

expression of stress-related genes nearby (Luna et al., 2012; Wibowo et al., 2016;

Annacondia et al., 2021). Such stress-induced epialleles are usually considered to

have limited heritability, as they often vanish after maximally three offspring

generation (Luna et al., 2012; Wibowo et al., 2016), and thus may occur due to direct

stress exposure during the early development of the offspring rather than being truly

inherited. Recent evidence, however, indicates that deterministic epialleles may be

substantially more stable than anticipated: in the duckweed *Lemna minor*, heat

stress remodelled CHG-methylation, and part of this variation was still observed after

at least three clonal generations under control conditions (Van Antro et al., 2023).

Similarly, in the closely related duckweed *S. polyrhiza*, copper excess induced

variation in phenotypes and fitness that persisted for up to 10 clonal generations

under control conditions (Huber et al., 2021). If deterministic epialleles are heritable

across multiple generations, as these studies suggest, they may lead to rapid stress

adaption even in the absence of selection.

METHODS

The problem: teasing apart the stochastic and deterministic routes

While the stochastic route relies on random variation being selected upon, the

deterministic route may allow populations to adapt in the absence of selection. Does

epigenetic inheritance thereby lead to a fundamentally different adaptive process in

which adaptation is, to a certain extent, predictable and deterministic, or does

adaptation trough epigenetic inheritance proceeds like genetic adaptation through

selection of phenotypic variants that are randomly generated? Answering this

question is fundamental to understand whether and how epigenetic inheritance may

lead to rapid adaption. Yet, assessing and teasing apart the stochastic and

deterministic route is experimentally challenging.

Experiments that aim to test the deterministic route are relatively common,

e.g. (Luna et al., 2012; Wibowo et al., 2016; Baugh & Day, 2020; Huber et al., 2021).

Usually, genetically uniform organisms are grown for multiple generations under

different environments ("pre-treatment") as single descendants - thereby

confounding genetic effects as well as selection of beneficial variants are almost

completely avoided (Baugh & Day, 2020). Subsequently, individuals are grown for at

least three generations under control conditions to avoid that the organisms were

directly exposed to stress during their early development (Heard & Martienssen,

2014). Subsequently, fitness and epigenetic variation between the pre-treatments

are assessed. Ideally this experiment is accompanied with genetic or chemical

manipulation of the epigenetic machinery and/or genes that are transgenerationally

regulated to test the molecular mechanisms. While this approach is powerful to

assess the role of deterministic epialleles in stress adaptation, it does not assess

whether selection of stochastic variants may lead to stress adaptation.

Experiments that test whether stochastic epigenetic variants may mediate

rapid stress adaption through natural selection are rather rare, but see (Kronholm et

al., 2017; Schmid et al., 2018; Heckwolf et al., 2020; Huber et al., 2021), likely

because of the experimental challenges. The most common approach is to grow

large, genetically uniform populations for many generations under different

environments ("pre-treatment"), subsequently grow individuals for a few generations

under a shared control environment to erase environment-induced effects, and then

compare fitness and epigenetic variation between the populations of the different

pre-treatments. The limitation of this approach is that during the pre-treatment,

populations simultaneously experience both selection of stochastic variants as well

as the induction of deterministic epialleles – and if the deterministic epialleles are

heritable for multiple generations, and show unexpected temporal inheritance

patterns (Bell & Hellmann, 2019; Huber et al., 2021) – variation between the

populations could be either due to selection of stochastic variants or heritable

deterministic epialleles. To assess whether stochastic epigenetic variation may

mediate rapid adaption through natural selection, we thus need an approach in

which selection of stochastic variants can take place while teasing apart the

contribution of deterministic epialleles.

The solution: manipulating the efficacy of selection through the population

size

Here, we propose an experimental approach that unifies both of the above-

mentioned setups and thereby allows to simultaneously assess whether rapid stress

adaptation may be mediated by selection of stochastic variants or the formation of

deterministic epialleles. One of the key differences between the stochastic and

deterministic route is whether selection is needed – and as such, to differentiate

among these two routes, one should manipulate the efficacy of selection. This can

be achieved through the population size, as selection is most effective in large

populations (Wahl et al., 2002; McDonald, 2019). We therefore propose to subject

populations for several generations to different environments ("pre-treatments") at

two different population sizes: on the one hand, in population sizes sufficiently large

that selection is effective. These populations may undergo both the stochastic and

deterministic route. On the other hand, in population sizes sufficiently small that

selection is not effective, i.e. that drift overcomes the effect of selection. This is

achieved once the effective population size $N_e < 1/s$ (s = selection coefficient)

(Nielsen & Slatkin, 2013; McDonald, 2019); the smallest possible effective population

size is achieved in single descendant lineages. Thus, the small population will evolve

almost exclusively through drift, and thereby only undergo the deterministic route.

Importantly, the large and small populations must be grown side by side in the very

same environment to ensure that the deterministic route is equally induced in the

small and large populations.

After several generations, one can assess whether populations adapted to

stress through the stochastic and/or deterministic route. To this end, individuals of

the small and large populations of the different pre-treatments are grown in a shared

control environment for at least three generations to ensure that any variation is truly

heritable. Subsequently, phenotypes and fitness under the different environments

are compared. This allows differentiating whether selection of epigenetic variants

and/or deterministic epialleles lead to rapid adaptation: if deterministic epialleles

confer resistance, variation in resistance and epigenetic variation will arise between

the small populations of the different pre-treatments. If selection of epigenetic

variants contributes to resistance, variation in resistance and in epigenetic variation

will establish between the small and large population within each pre-treatment (Fig.

1b).

(a) Routes of epigenetic adaptation

(b) Experimental approach

***Figure 1. The two routes of rapid adaptation through epigenetic inheritance.** (a)*

Epigenetic inheritance may facilitate adaptation either via deterministic epialleles

(deterministic route) or selection of stochastic epigenetic variants (stochastic route).

Each circle and triangle indicates the DNA methylation level of a specific locus of

different individuals. (b) To differentiate among the deterministic and stochastic route,

organisms are grown in two different environments (pre-treatments) in large and small

populations; after a multi-generational recovery phase in a shared control environment,

fitness, phenotypes and epigenetic marks in the presence and absence of recurring

stress are compared between populations sizes and pre-treatments.

The requirements: is my experimental system suitable for the approach?

While the proposed experiment is in principle applicable to species across the tree of

life, the species under investigation should have following characteristics:

   1. *Rapid reproduction and small body size.* As with all species used for

      experimental evolution, rapid reproduction and small body size are highly

      desirable.

2. *Low genetic variation.* Teasing apart genetic from epigenetic effects is

notoriously difficult, see discussion below. We thus suggest minimizing

genetic variation by using clonally reproducing organisms or organisms that

are highly inbred; ideally, genetic mutation rates should be small to reduce

genetic variation that arise during the experiment.

3. *Accurate assessments of fitness and epigenetic variation.* Measuring

Darwinian fitness is critical when assessing adaptation – thus, direct fitness

assessments (e.g. number of offspring) is preferred over indirect assessments

such as phenotypes or altered performance of interacting species; such

assessments are nevertheless valuable. Furthermore, the ability to measure

epigenetic variation (e.g. DNA methylation, histone modification, small RNAs)

in a largely unbiased manner and to link this variation to specific genes is

highly desirable. Although these measurements often require a reference

genome, new developments in high-throughput sequencing will likely

overcome some of the limitations of non-model species that do not have a

high-quality reference genome or whose genome is prohibitively large

(Amarasinghe et al., 2020; Gawehns et al., 2022).

Proof-of-concept: applying the method to assess rapid adaptation in

monoclonal duckweed populations

One biological system that is ideal to test adaptation through the deterministic or

stochastic route is the giant duckweed, *Spirodela polyrhiza,* one of the smallest and

fastest growing flowering plants. *Spirodela polyrhiza* reproduces almost exclusively

asexually by budding with a maximal doubling time of two to three days (Landolt

1957; Ziegler et al., 2015). The species has a very low mutation rate with

approximately one point mutation every eighty generations (Xu et al., 2019). In

nature, *S. polyrhiza* is exposed to diverse environmental stresses including copper

excess, which is a common pollutant in aquatic ecosystems (Greger, 1999; Li &

Xiong, 2004).

To test whether *S. polyrhiza* acquires stress resistance through the stochastic

or deterministic route, we grew monoclonal *S. polyrhiza* populations in the presence

and absence of copper excess ("pre-treatment") in both small and large populations

("selection regimen") for approximately seven generations in ponds outdoors and

assessed plant resistance to copper excess after five generations of growth in a

shared control environment. The small populations consisted of single descendant

lineages – the smallest possible population size – which were floating freely inside

the large populations (Fig. 2a).

On May 2021, we set up 20 ponds (80 x 60 x 32cm, AuerPackaging,

Amerang, Germany) inside an open field site in Münster, Germany (51°57'54.0'N,

7°36'22.4'E). Ponds were buried to the rim with soil and filled with 10 L of pond soil

(Floragard, Oldenburg, Germany), 40 mL of organic fruit and vegetable fertilizer

(organic NPK 3,1+0,5+4,1; COMPO, Münster, Germany) and tap water with a total

volume of 100 L. The boxes had a draining scape and lids consisting of a stainless-

steel mesh (mesh opening 0.63 mm, wire diameter, 0.224 mm; Haver&Boeker,

Oelde, Germany). On May 26th, each box received around 3000 duckweeds of the

genotype 7498 that were pre-cultivated inside 10% N-medium (Appenroth et al.,

1996) for one month in indoor growth cabinets (GroBank, CLF PlantClimatics,

Wertingen, Germany) operating at 26°C, 135 µmol photons m^{-2} s^{-1} supplied through

LEDs with 16:8 h light:dark cycles. After one week of acclimation, we added 20 µM

$CuSO_4$ to half of the ponds (n=10) to initiate the stress pre-treatment (Fig. 2a).

Across the entire experiments, we measured copper concentration weekly and after

each rainfall and added $CuSO_4$ if needed to maintain 20 µM $CuSO_4$. After adding

$CuSO_4$ for the first time, we randomly chose four plants per pond as the founders of

the single descendant lineages. To trace the single descendant lineages, we marked

the plants with a dot using a permanent marker (Stabilo OHPen universal) (Fig. 2a).

The remaining plants reproduced freely and covered the surface after 12 days, which

corresponds to four to five generations of growth and a maximal population size of

25,000 individuals. To avoid crowding, we randomly removed 30% of the plants.

Subsequently, we also marked four plants from the populations and followed them

for two generations as single descendant lineages to equalize the marking treatment

and the plant age between the single descendants and large populations. To grow

the plants from the different pre-treatments and selection regimens in a uniform

environment ("recovery phase"), we moved the third marked generation of the

population and the sixth generation of the single descendants (approx. 24 days of

pre-treatment) into separate cavities of polyvinyl chloride disks (7 cm diameter) that

were floating inside the control ponds. Due to algae bloom, these plants did not

reproduce any offspring, so we moved the same plants back to the initial indoor

conditions after 38 days of growth outdoors. To establish a shared environment

among plants from the different pre-treatments and selection regimens, we placed

the plants from the single descendant lineages and the large populations from each

control and copper pre-treatment pond pair together inside a 250mL polypropylene

beakers (Plastikbecher GmbH, Giengen an der Brenz, Germany) filled with 150

mL10% N-medium and covered with perforated and transparent lids (four groups per

beaker). Plants from the four different groups were marked with specific colour

patterns to follow the individual lineages. To reduce algal growth, we changed the

medium every two to three days and wrapped the sides of the beakers with three

layers of black plastic foil to decrease light incidence. Inside the plastic beakers, we

propagated each lineage for five generations using single descendant propagation.

To assess plant fitness and resistance to copper excess, we moved the first and

second daughter plant of generation five into separate 250 ml polypropylene beakers

containing control conditions (10% N-medium) or recurrent copper excess (20 µM

$CuSO_4$ in 10% N-medium); the first and second offspring were randomly assigned to

the treatment groups. After setting up the experiment, we took a picture for surface

area analysis using a camera installation with a webcam (HD Pro Webcam C270,

Logitech, Lausanne, Switzerland; webcam software 2.12.8). To reduce algae growth,

we exchanged the medium after four days. After eight days of growth, we took a

second picture and then gently dried the plants with a tissue paper before weighting

the plants. Fresh mass of the plant that initiated the fitness assay was assessed

separately from the fresh mass of the offspring to measure growth rates based on

biomass, as described below.

Statistical analyses

We calculated growth rates based on frond surface coverage and fresh weight

[(End value)-(Initial value)]/days. Additionally, we estimated resistance to copper

excess as the ratio in plant surface area of plant grown under copper excess)

relative to the mean value of plants under control treatment). To test whether plant

resistance and growth rates under control conditions and copper excess are affected

by the selection regimen and the pre-treatment, we fitted a linear mixed effects

model using selection regimen and pre-treatment as fixed effects, and the pair of a

control and a copper pre-treatment ponds ("pair") and blocks of ponds (groups of

three to four ponds, "block") regarding their physical position in the field as random

effects (~Selection*Pre-treatment+(1|pair)+(1|block)). Models were run in R v4.3.1

using the package lme4 v1.1.34 (Bates et al., 2015), and *P*-values of the fixed

effects were estimated with Type III Analysis of Variance with the Satterthwaite´s

method using anova of the package stats (R Core Team 2023). We used DHARMa

v0.4.6 (Hartig, 2022) to verify whether the scaled residuals of our fitted models

correlated to the simulated values from the same model. When analyzing plant

resistance and fitness under copper excess, we found an influential sample in the

copper pre-treated large population under copper environment, which showed

Cook´s distance higher than 0.5, low correlation between the standardized residuals

and theoretical quantiles (Q-Q plot) and whose sibling was missing in the control

environment. Therefore, we excluded this outlier from the analysis. Plots were

displayed with ggplot2 v3.4.3 (Wickham, 2016).

RESULTS

Proof-of-concept: duckweeds possibly adapt through both the stochastic and

deterministic epigenetic routes

To assess whether *S. polyrhiza* acquires resistance to copper excess through either

the deterministic or stochastic route, we measured plant resistance of single

descendants and large populations grown under control and copper pre-treatment,

the latter reduced plant growth rates outdoors by 30%. Both the presence of

selection and copper pre-treatment affected plant resistance: regardless of the pre-

treatment, plants from the large populations tended to have higher resistance to

copper excess than plants from the single descendant lineages (Fig. 1b; Selection:

$F_{(1,3.33)} = 7.24$, $P = 0.066$, linear mixed effects model). Furthermore, pre-treatment in

copper stress tended to increase copper resistance in the single descendants but not

in the populations (Fig. 1b; Selection x pre-treatment: $F_{(1,12.78)} = 7.48$, $P = 0.017$,

linear mixed effects model). Similar results were obtained when resistance was

calculated based on biomass (Supporting Information Appendix Fig. S1; Selection x

pre-treatment: $F_{(1,13.01)} = 5.32$, $P = 0.038$, linear mixed effects model) instead of

surface coverage, as described above.

To differentiate whether the increase in resistance was due to lower growth

under control conditions or higher growth under copper excess, we analysed plant

fitness under both treatment environments separately. Under control conditions,

neither the population size nor the pre-treatment altered plant fitness (Fig. 2c;

$F_{(1,10.33)} = 0.50$, $P > 0.1$, linear mixed effects models). In contrast, under copper

excess, plants from the large populations outperformed plants from the single

descendant lineages regardless of the pre-treatment. (Fig. 2d; Selection: $F_{(1,3.68)} =$

8.34, $P = 0.049$, mixed effects models). Furthermore, pre-treatment in copper stress

tended to increase plant fitness under copper excess in the single descendant

lineages but not in the populations (Selection x pre-treatment: $F_{(1,12.79)} = 3.86$, $P =$

0.071, linear mixed effects models). Therefore, variations in copper resistance were

mainly due to variations in growth rates under copper excess and not under control

conditions.

(a) Experimental setup

(b) Resistance

(c) Control environment

(d) Copper environment

Figure 2. Testing the experimental approach by growing large and small monoclonal populations of Spirodela polyrhiza in the presence and absence of copper excess in outdoor mesocosms. *(a) Overview of the experimental setup. (b) Plant resistance to copper excess (growth rates under copper excess compared to the group´s mean growth rate under control conditions) depended on the population size and the pre-treatment. (c) Growth rates under control environment was not affected by the population size and the pre-treatment environment. (d) Growth rates under copper excess were affected by the population size and the pre-treatment. For b-c:* P*-values refer to linear mixed effects models. The black circles mark the means, and error bars show the standard errors.* $N_{(control\ pre\text{-}treat\ *\ Pop\ regimen)}$ = 5; $N_{(control\ pre\text{-}treat\ *\ Sgl\ regimen)}$ = 3-5; $N_{(copper\ pre\text{-}treat\ *\ Pop\ regimen)}$ = 3; $N_{(copper\ pre\text{-}treat\ *\ Sgl\ regimen)}$ = 4-5. *Pre-treat = pre-treatment; Sel = selection; Pop = large population; Sgl = single descendants.*

DISCUSSION

We here presented an experimental approach to test a long-standing controversy:

whether epigenetic inheritance can lead to rapid adaptation to environmental

stresses through stochastic or deterministic epigenetic variation. Our proof-of-

concept experiments suggests that both the deterministic and stochastic route may

alter duckweed fitness under recurring stress.

Proof-of-concept experiment

In our experiment, offspring from large populations outperformed offspring

from the single descendants regardless of the pre-treatment. The higher fitness of

the large populations compared to single descendants may have been caused by

two factors: first, variations among single descendants and large populations may

have been confounded by our method to identify the single descendants during the

pre-treatment phase. To identify the single descendants, we marked the single

descendants throughout the pre-treatment phase. Although we also marked plants

from the large populations in the last two generations of the pre-treatment, we

cannot fully rule out that the prolonged marking of the single descendants affected

plant fitness even after the five generations of growth under a shared control

conditions prior fitness assay. In this case, the difference between the large

populations and single descendants is not caused by selection (stochastic route) but

is in fact a pre-treatment effect (deterministic route) that persisted across five

generations. Second, the higher fitness of the large populations than of single

descendants may reflect true differences caused by selection. Our experiment was

limited to five generations. While such short durations are sufficient for natural

populations to adapt through standing genetic variation (Züst et al., 2012), it is

unclear whether monoclonal populations may acquire sufficient variation through

epigenetic inheritance for such rapid adaptation taking place. Future experiments

that assess variation in plant fitness among single descendant lineages, and long-

term experiments that ensure that small and large populations experience the very

same environment during the pre-treatment phase will be needed to differentiate

among these two possibilities.

While the largest differences in fitness was caused by the different population

size, we also obtained evidence for deterministic effects (interaction of population

size and pre-treatment on plant resistance). This is in line with recent studies

suggesting that the previous environment alters plant fitness and epigenetic marks

across multiple clonal generations (González et al., 2018; Li et al., 2018; Benson et

al., 2021; Colinas et al., 2023; Sammarco et al., 2023). Future manipulative studies

are needed that test for specific molecular mechanism of the deterministic route to

confirm the presence of this phenomenon and to understand whether the observed

phenomena are truly caused by epigenetic inheritance *sensu stricto* (genome-

associated mechanisms) or by other non-genetic inheritance mechanisms.

While the proposed experiment is not designed to tease apart the specific

epigenetic or non-genetic mechanism, the approach can – if carried out carefully –

resolve the contribution of the deterministic and stochastic route in a wide range of

organisms. In addition, the approach can also be used to assess fitness burdens of

genetic mutations, as proposed previously (Ho et al., 2021). We would like to

highlight a few points to make maximal use of this experimental approach:

Inferring the stochastic and deterministic route

The above-described experiment with duckweeds reveals a critical issue: the pre-

treatment environment of the small and large population must be equal, otherwise

one cannot differentiate between the stochastic and deterministic route. In our

experiment, this could be achieved by caging small populations inside the large

population – this would alleviate the potential bias of the marking, however, one must

ensure that the small populations still experience the same environment as the large

populations despite the caging. In duckweeds, this can be achieved by separating

the small populations from the large populations using floating disks while keeping

the density of the small and large populations equal. It is of utmost importance to

carefully design of the experiments considering the specific biology of each

organism.

So far, we treated the stochastic and deterministic route as separate, non-

overlapping entities. This may be an unrealistic assumption. For instance,

deterministic epialleles may not be induced in an entirely deterministic manner;

environmental stresses may simply increase the likelihood of an epiallele to appear

but not cause all individuals changing their epigenetic status. Selection could thus

still act on environment-induced "deterministic" epialleles. Similar accounts for the

stochastic epimutations: environmental stresses may increase the likelihood that

certain epimutations appear (Zheng et al., 2017; Hazarika et al., 2022) and thus

epimutations may not be entirely stochastic. Are the stochastic and deterministic

route therefore two extremes of the same process? One argument holds against this:

in *A. thaliana*, spontaneous epimutations mostly accumulate in the CG context

whereas environment-induced epialleles are most prominent in the CHH and CHG

context (Lin et al., 2022; Yadav et al., 2022). This supports the idea that the

stochastic and deterministic route are, at least in the context of DNA methylation in

plants, two fundamentally different processes.

Inferring the molecular mechanisms

Another critical issue is the interpretation of the results: is the underlying

molecular mechanism of genetic or epigenetic nature? Even when using highly

inbred or clonally reproducing species, genetic mutations may accumulate and

account for variation in phenotypes or fitness, particularly in the large populations.

We suggest following experiments and analyses to tease apart genetic from

epigenetic effects: first, unbiased methods (e.g. high-throughput sequencing) should

be deployed to screen for epigenetic variations. Second, transcriptome and

proteome data should be generated to link fitness to phenotypes to genes to the

epigenome. Third, differentially regulated genes or epigenetic machineries should be

manipulated both genetically as well as chemically to test whether the identified

genes or epigenetic machinery affect stress resistance or the process of stress

adaptation. Fourth, phenotypes or fitness should be continuously assessed for

multiple generations after stress release – if variations between groups have an

epigenetic basis, the variation should diminish or at least change in some lineages

over time. Fifth, high-throughput sequencing should be deployed to screen for

genetic variation in identified genomic regions. While in isolation none of these

approaches will provide a clear answer whether genetic or epigenetic mechanisms

are at play, combining these approaches will be powerful to disentangle genetic and

epigenetic factors.

CONCLUSIONS

Taken together, we here provide an experimental framework to assess the relative

contribution of deterministic epialleles and selection of stochastic epigenetic variants

to rapid stress adaptation. Through this approach, carefully designed experiments

can provide novel insights whether and how epigenetic inheritance may lead to rapid

stress adaptation. Considering the on-going loss of intra-specific genetic diversity

and the rapid pace of environmental change, these insights may become

increasingly relevant when assessing species potential to resist global change.

ACKNOWLEDGEMENTS

We would like to thank Pauline Prüsener, Anke Grosse-Brinkhaus, Maite Görtz

Lizarraga and the Botanical Garden of the University of Münster for supporting the

outdoor experiment and Shuqing Xu for providing facilities. This study profited from

interactions within the GenEvo Research Training Group 407023052/GRK2526/1

funded by the German Research Foundation (DFG). The work was funded by the

Volkswagen Foundation (Grant Nr 97236) to MH and the German Research

Foundation to MH (grant Nr. 512079118). The project was supported by the

University of Münster and the University of Mainz.

CONFLICT OF INTEREST

480   The authors declare no conflict of interest.

AUTHOR CONTRIBUTION

483   MH conceived the ideas and designed methodology, AC collected the data, AC and

484   MH analysed the data, MH and AC drafted the manuscript. Both authors contributed

485   critically to the drafts and gave final approval for publication.

DATA AVAILABILITY

488   The raw data of the outdoor experimental evolution experiment will be archived on

489   Dryad upon acceptance.

REFERENCES

Amarasinghe, S.L., Su, S., Dong, X., Zappia, L., Ritchie, M.E. & Gouil, Q. (2020)
Opportunities and challenges in long-read sequencing data analysis. *Genome*
Biology, **21,** 30.

Annacondia, M.L., Markovic, D., Reig-Valiente, J.L., Scaltsoyiannes, V., Pieterse,
C.M.J., Ninkovic, V., Slotkin, R.K. & Martinez, G. (2021) Aphid feeding

induces the relaxation of epigenetic control and the associated regulation of

the defense response in *Arabidopsis. New Phytologist,* **230,** 1185-1200.

Appenroth, K.J., Teller, S. & Horn, M. (1996) Photophysiology of turion formation and

germination in *Spirodela polyrhiza. Biologia Plantarum,* **38,** 95.

Baugh, L.R. & Day, T. (2020) Nongenetic inheritance and multigenerational plasticity

in the nematode *C. elegans. eLife,* **9,** e58498.

Becker, C., Hagmann, J., Müller, J., Koenig, D., Stegle, O., Borgwardt, K. & Weigel,

D. (2011) Spontaneous epigenetic variation in the *Arabidopsis thaliana*

methylome. *Nature,* **480,** 245-249.

Bell, A.M. & Hellmann, J.K. (2019) An integrative framework for understanding the

mechanisms and multigenerational consequences of transgenerational

plasticity. *Annual Review of Ecology, Evolution and Systematics,* **50,** 97-118.

Benson, C.W., Mao, Q. & Huff, D.R. (2021) Global DNA methylation predicts

epigenetic reprogramming and transgenerational plasticity in Poa annua L.

Crop Science, **61,** 3011-3022.

Bonduriansky, R. & Day, T. (2009) Nongenetic inheritance and its evolutionary

implications. *Annual Review of Ecology, Evolution and Systematics,* **40,** 103-

125.

Colinas, N., Montero-Pau, J., Carmona, M.J., Sabatino, R., Di Cesare, A., Eckert,

E.M. & García-Roger, E.M. (2023) Transgenerational expression profiles of a

sex related and an epigenetic control gene in the rotifer Brachionus plicatilis in

relation to environmental predictability. *Hydrobiologia.*

Cubas, P., Vincent, C. & Coen, E. (1999) An epigenetic mutation responsible for

natural variation in floral symmetry. *Nature,* **401,** 157-161.

Denkena, J., Johannes, F. & Colomé-Tatché, M. (2021) Region-level epimutation

rates in *Arabidopsis thaliana*. *Heredity,* **127,** 190-202.

Des Roches, S., Pendleton, L.H., Shapiro, B. & Palkovacs, E.P. (2021) Conserving

intraspecific variation for nature's contributions to people. *Nature Ecology &*

Evolution, **5,** 574-582.

Gawehns, F., Postuma, M., van Antro, M., Nunn, A., Sepers, B., Fatma, S., van

Gurp, T.P., Wagemaker, N.C.A.M., Mateman, A.C., Milanovic-Ivanovic, S.,

Groβe, I., van Oers, K., Vergeer, P. & Verhoeven, K.J.F. (2022) epiGBS2:

Improvements and evaluation of highly multiplexed, epiGBS-based reduced

representation bisulfite sequencing. *Molecular Ecology Resources,* **22,** 2087-

2104.

González, A.P.R., Preite, V., Verhoeven, K.J.F. & Latzel, V. (2018)

Transgenerational effects and epigenetic memory in the clonal plant *Trifolium*

repens. Frontiers in Plant Science, **9,** 1677-1677.

Greger, M. (1999) Metal Availability and Bioconcentration in Plants. *Heavy Metal*

Stress in Plants: From Molecules to Ecosystems (eds M.N.V. Prasad & J.

Hagemeyer), pp. 1-27. Springer Berlin Heidelberg, Berlin, Heidelberg.

Hartig, F. (2022) DHARMa: Residual Diagnostics for Hierarchical (Multi-Level /

Mixed) Regression Models.

Hazarika, R.R., Serra, M., Zhang, Z., Zhang, Y., Schmitz, R.J. & Johannes, F. (2022)

Molecular properties of epimutation hotspots. *Nature Plants,* **8,** 146-156.

Heard, E. & Martienssen, R.A. (2014) Transgenerational epigenetic inheritance:

myths and mechanisms. *Cell,* **157,** 95-109.

Heckwolf, M.J., Meyer, B.S., Häsler, R., Höppner, M.P., Eizaguirre, C. & Reusch,

T.B.H. (2020) Two different epigenetic information channels in wild three-

spined sticklebacks are involved in salinity adaptation. *Science advances,* **6,**

eaaz1138-eaaz1138.

Ho, E.K.H., Bellis, E.S., Calkins, J., Adrion, J.R., Latta Iv, L.C. & Schaack, S. (2021)

Engines of change: Transposable element mutation rates are high and

variable within Daphnia magna. *PLOS Genetics,* **17,** e1009827.

Huber, M., Gablenz, S. & Höfer, M. (2021) Transgenerational non-genetic

inheritance has fitness costs and benefits under recurring stress in the clonal

duckweed *Spirodela polyrhiza*. *Proceedings of the Royal Society of London,*

Series B: Biological Sciences, **288**.

Jiang, C., Mithani, A., Belfield, E.J., Mott, R., Hurst, L.D. & Harberd, N.P. (2014)

Environmentally responsive genome-wide accumulation of de novo

Arabidopsis thaliana mutations and epimutations. *Genome Research,* **24,**

1821-1829.

Johannes, F. & Schmitz, R.J. (2019) Spontaneous epimutations in plants. *New*

Phytologist, **221,** 1253-1259.

Kronholm, I., Bassett, A., Baulcombe, D. & Collins, S. (2017) Epigenetic and genetic

contributions to adaptation in Chlamydomonas. *Molecular Biology and*

Evolution, **34,** 2285-2306.

Landolt, E. (1957) Physiologische und ökologische Untersuchungen an Lemnaceen.

Bericht der Schweizerischen Botanischen Gesellschaft, **67,** 271–410.

Li, K., Chen, J., Wei, Q., Li, Q. & Lei, N. (2018) Effects of Transgenerational

Plasticity on Morphological and Physiological Properties of Stoloniferous Herb

Centella asiatica Subjected to High/Low Light. *Frontiers in Plant Science,* **9**.

Li, T. & Xiong, Z. (2004) A novel response of wild-type duckweed (Lemna

paucicostata Hegelm.) to heavy metals. *Environmental Toxicology,* **19,** 95-

102.

Lin, X., Zhou, M., Yao, J., Li, Q.Q. & Zhang, Y.-Y. (2022) Phenotypic and Methylome

Responses to Salt Stress in Arabidopsis thaliana Natural Accessions.

Frontiers in Plant Science, **13**.

Luna, E., Bruce, T.J.A., Roberts, M.R., Flors, V. & Ton, J. (2012) Next-generation

systemic acquired resistance. *Plant Physiology,* **158,** 844-853.

Manning, K., Tör, M., Poole, M., Hong, Y., Thompson, A.J., King, G.J., Giovannoni,

J.J. & Seymour, G.B. (2006) A naturally occurring epigenetic mutation in a

gene encoding an SBP-box transcription factor inhibits tomato fruit ripening.

Nat Genet, **38,** 948-952.

McDonald, M.J. (2019) Microbial experimental evolution - a proving ground for

evolutionary theory and a tool for discovery. *EMBO reports,* **20,** e46992-

e46992.

Nielsen, R. & Slatkin, M. (2013) *An introduction to population genetics: theory and*

applications. Sinauer.

O'Dea, R.E., Noble, D.W.A., Johnson, S.L., Hesselson, D. & Nakagawa, S. (2016)

The role of non-genetic inheritance in evolutionary rescue: epigenetic

buffering, heritable bet hedging and epigenetic traps. *Environmental*

Epigenetics, **2**.

Ong-Abdullah, M., Ordway, J., Jiang, N., Ooi, S., Kok, S.Y., Sarpan, N., Azimi, N.,

Hashim, A., Ishak, Z., Rosli, S., Ahmad Malike, F., Abu Bakar, N., Marjuni, M.,

Abdullah, N., Yaakub, Z., Din, A., Rajanaidu, N., Singh, R., Low, L. &

593   Martienssen, R. (2015) Loss of Karma transposon methylation underlies the

594   mantled somaclonal variant of oil palm. *Nature,* **525,** 533-537.

595 Sage, R.F. (2020) Global change biology: A primer. *Global Change Biology,* **26,** 3-

596   30.

597 Sammarco, I., Díez Rodríguez, B., Galanti, D., Nunn, A., Becker, C., Bossdorf, O.,

598   Münzbergová, Z. & Latzel, V. (2023) DNA methylation in the wild: epigenetic

599   transgenerational inheritance can mediate adaptation in clones of wild

600   strawberry (Fragaria vesca). *Research Square.*

601 Schmid, M.W., Heichinger, C., Coman Schmid, D., Guthörl, D., Gagliardini, V.,

602   Bruggmann, R., Aluri, S., Aquino, C., Schmid, B., Turnbull, L.A. &

603   Grossniklaus, U. (2018) Contribution of epigenetic variation to adaptation in

604   *Arabidopsis. Nature Communications,* **9,** 4446.

605 Schmitz, R.J., Schultz, M.D., Lewsey, M.G., O'Malley, R.C., Urich, M.A., Libiger, O.,

606   Schork, N.J. & Ecker, J.R. (2011) Transgenerational epigenetic instability is a

607   source of novel methylation variants. *Science,* **334,** 369-373.

608 Team, R.C. (2023) R: A Language and Environment for Statistical Computing. R

609   Foundation for Statistical Computing, Vienna, Austria.

610 Van Antro, M., Prelovsek, S., Ivanovic, S., Gawehns, F., Wagemaker, N.C.A.M.,

611   Mysara, M., Horemans, N., Vergeer, P. & Verhoeven, K.J.F. (2023) Stability of

612   temperature-induced DNA methylation changes across clonal generation in

613   the common Duckweed *Lemna minor* L. *Molecular Ecology.*

614 Wahl, L.M., Gerrish, P.J. & Saika-Voivod, I. (2002) Evaluating the impact of

615   population bottlenecks in experimental evolution. *Genetics,* **162,** 961-971.

616 Wibowo, A., Becker, C., Marconi, G., Durr, J., Price, J., Hagmann, J., Papareddy, R.,

617   Putra, H., Kageyama, J., Becker, J., Weigel, D. & Gutierrez-Marcos, J. (2016)

Hyperosmotic stress memory in Arabidopsis is mediated by distinct

epigenetically labile sites in the genome and is restricted in the male germline

by DNA glycosylase activity. *eLife,* **5,** e13546.

Wickham, H. (2016) *ggplot2: Elegant Graphics for Data Analysis,* 2 edn. Use R!

Xu, S., Stapley, J., Gablenz, S., Boyer, J., Appenroth, K.J., Sree, K.S., Gershenzon,

J., Widmer, A. & Huber, M. (2019) Low genetic variation is associated with

low mutation rate in the giant duckweed. *Nature Communications,* **10,** 1243.

Yadav, N.S., Titov, V., Ayemere, I., Byeon, B., Ilnytskyy, Y. & Kovalchuk, I. (2022)

Multigenerational Exposure to Heat Stress Induces Phenotypic Resilience,

and Genetic and Epigenetic Variations in Arabidopsis thaliana Offspring.

Frontiers in Plant Science, **13**.

Zhang, H., Lang, Z. & Zhu, J.-K. (2018) Dynamics and function of DNA methylation

in plants. *Nature Reviews: Molecular Cell Biology,* **19,** 489-506.

Zheng, X., Chen, L., Xia, H., Wei, H., Lou, Q., Li, M., Li, T. & Luo, L. (2017)

Transgenerational epimutations induced by multi-generation drought

imposition mediate rice plant's adaptation to drought condition. *Scientific*

Reports, **7,** 39843.

Ziegler, P., Adelmann, K., Zimmer, S., Schmidt, C. & Appenroth, K.J. (2015) Relative

in vitro growth rates of duckweeds (Lemnaceae) - the most rapidly growing

higher plants. *Plant Biol (Stuttg),* **17 Suppl 1,** 33-41.

Züst, T., Heichinger, C., Grossniklaus, U., Harrington, R., Kliebenstein, D.J. &

Turnbull, L.A. (2012) Natural enemies drive geographic variation in plant

defenses. *Science,* **338,** 116-119.