Assessing rapid adaptation through epigenetic inheritance: a new

experimental approach

Alexandra Chávez[1,2,] Meret Huber[1,2*,]

[1] Institute of Organismic and Molecular Evolution, University of Mainz, Johann-

Joachim-Becher-Weg 7, 55128 Mainz, Germany

[2] Institute of Plant Biology and Biotechnology, University of Münster, Schlossplatz 7,

48143 Münster, Germany

*Correspondence: Meret Huber, Institute of Organismic and Molecular Evolution,

Johann-Joachim-Becher-Weg 7, 55128 Mainz, Germany; meret.huber@uni-

mainz.de

Running headline: Adaption through epigenetic inheritance

Abstract

1. Epigenetic inheritance is hypothesized to mediate rapid adaptation to stresses

via two fundamentally different routes: first, through spontaneous

epimutations that arise in a largely stochastic manner in the presence or

absence of stress; if these spontaneous epimutations are heritable and

beneficial, they may be selected upon ("stochastic route"); and second,

through environment-induced epialleles that arise uniformly among

individuals; if heritable, these epialleles may lead to stress adaptation even in

the absence of selection ("deterministic route"). Testing and teasing apart

these two routes is challenging, largely because a suitable experimental

approach is lacking.

2. Here, we propose an experimental approach that allows to simultaneously

assess the contribution of the stochastic and deterministic route. The essence

of the approach is to manipulate the efficacy of selection through the

population size and thereby to test whether selection is required for

adaptation (stochastic route). To this end, genetically uniform populations are

grown under different environments across multiple generations ("pre-

treatment") at two different population sizes: in large populations, in which

selection is effective; and in small populations, in which drift overcomes the

effect of selection. If the deterministic route contributes to adaptation, variation

in fitness, phenotypes or epigenetic marks will arise between the small

populations of the different pre-treatments. If the stochastic route contributes

to adaptation, variation will arise between the small and large population

within each pre-treatment. As a proof-of-principle, we tested whether small

and large monoclonal populations of the aquatic duckweed *Spirodela*

polyrhiza may adapt to copper excess outdoors.

3. After five to seven generations of pre-treatment and a subsequent multi-

generational growth under control conditions, large populations outperformed

small populations under copper excess. Furthermore, small populations pre-

treated with copper excess tended to have higher fitness under copper excess

than small populations pre-treated under control conditions. These data

suggest that both the stochastic and deterministic route may alter plant fitness

under recurring stress.

4. The proposed approach will allow to experimentally evaluate whether species

may adapt to stresses through either stochastic and deterministic epigenetic

changes, which is fundamental to understand whether and how epigenetic

inheritance may lead to rapid stress adaptation.

Keywords

Epigenetic inheritance, non-genetic inheritance, transgenerational plasticity, rapid

adaptation to stress, deterministic and stochastic epigenetic variation, experimental

evolution, epimutation, duckweed

INTRODUCTION

The hallmark of evolutionary biology states that species adapt to stresses through

selection of phenotypic variants that are arise from DNA sequence variation. Yet,

increasing evidence shows that DNA sequence variation is not the sole source of

heritable phenotypic variation: for instance, flowers of natural toadflax mutants

develop radial instead of bilateral symmetry because a floral symmetry gene became

hypermethylated (Cubas et al., 1999). Similarly, homeotic floral phenotypes in the oil

palm arose during tissue culture because an homeotic gene is alternatively spliced

upon spontaneous hypomethylation of an intron located transposon (Ong-Abdullah

et al., 2015). Furthermore, fruit ripening in a spontenous tomato mutant is hampered

because a gene of the SBP-box family of trascription factors became

hypermerthylated and was thereby silenced (Manning et al., 2006). These examples

highlight that mechanisms other than DNA sequence variation may lead to heritable,

variable and fitness-relevant phenotypes, and as such they open up an exciting

question: which role do mechanisms that generate heritable phenotypic variation in

the absence of DNA sequence variation play in the adaptation to environmental

stresses? Considering the on-going loss of intra-specific genetic diversity and the

rapid pace of environmental change (Des Roches et al., 2021; Sage, 2020),

answering this question is becoming an increasingly relevant.

Heritable phenotypic variation that is not caused by DNA sequence changes

may arise through so-called "non-genetic" or "epigenetic" inheritance (Bonduriansky

& Day, 2009; O'Dea et al., 2016). Epigenetic inheritance often refers to mechanisms

that alter gene expression across generations through genome-associated

mechanisms such as DNA methylation, histone modifications and small RNAs,

whereas non-genetic inheritance includes any other mechanism such as the vertical

transfer of microbes, nutrients or hormones (Bonduriansky & Day, 2009). In this

manuscript, we use the term "epigenetic" to refer to both epigenetic and other non-

genetic mechanisms in order to improve readibily.

Many epigenetic marks share two features: first, epigenetic marks may

change spontaneously over generations in a stochastic manner (Becker et al., 2011;

Schmitz et al., 2011), and second, epigenetic marks may be induced by the

environment and thereby lead to predictable phenotypic variation (Wibowo et al.,

2016; Zhang et al., 2018). Consequently, epigenetic inheritance may lead to stress

adaptation through two fundamentally different routes: the "stochastic" and the

"deterministic" route (Fig. 1a) (Baugh & Day, 2020).

The stochastic route is comparable to the adaptive process that is based on

DNA sequence variation. Here, spontaneous changes ("epimutations") arise either in

the presence or absence of stress in a largely random manner. For instance, in

Arabidopsis thaliana, variations in CG and to a much lower extent also CHG and

CHH methylation (H=A, T, C) steadily accumulate across generations (Becker et al.,

2011; Denkena et al., 2021; Schmitz et al., 2011), and both the frequency and the

spectrum of these spontaneous epimutations may be altered by stress (Jiang et al.,

2014; Johannes & Schmitz, 2019). In plants, these spontaneous epimutations are

semi-stable, meaning that many epimutations are inherited across several

generations but also frequently reverse (Johannes & Schmitz, 2019). Spontaneous

epimutations increase the phenotypic diversity in genetically uniform individuals upon

which natural selection can act.

The deterministic route, in contrast, is not comparable to the adaptive process

that is based on DNA sequence variation. In the deterministic route, also described

as transgenerational plasticity, reviewed in Anastasiadi et al. (2021), environmental

stresses induce epigenetic variation and thereby lead to predictable epigenetic

variations that are shared among individuals ("deterministic epialleles"). For instance,

in *A. thaliana,* environmental stresses may remodel CHG and CHH methylation in

specific repeat sequences, which is associated with altered expression of stress-

related genes nearby (Annacondia et al., 2021; Luna et al., 2012; Wibowo et al.,

2016). Such stress-induced epialleles are usually considered to have limited

heritability, as they often vanish after maximally three offspring generation (Luna et

al., 2012; Wibowo et al., 2016), and thus may occur due to direct stress exposure

during the early development of the offspring rather than being truly inherited.

Recent evidence, however, indicates that deterministic epialleles may be

substantially more stable than anticipated: in the duckweed *Lemna minor*, heat

stress remodelled CHG-methylation, and part of this variation was still observed after

at least three clonal generations under control conditions (Van Antro et al., 2023).

Similarly, in the closely related duckweed *S. polyrhiza*, copper excess induced

variation in phenotypes and fitness that persisted for up to 10 clonal generations

under control conditions (Huber et al., 2021). If deterministic epialleles are heritable

across multiple generations, as these studies suggest, they may lead to rapid stress

adaption even in the absence of selection.

METHODS

The problem: teasing apart the stochastic and deterministic routes

While the stochastic route relies on random variation being selected upon, the

deterministic route may allow populations to adapt in the absence of selection. Does

epigenetic inheritance thereby lead to a fundamentally different adaptive process in

which adaptation is, to a certain extent, predictable and deterministic, or does

adaptation through epigenetic inheritance proceeds similar to genetic adaptation

through selection of phenotypic variants that are generated through random

epimutations? Answering this question is fundamental to understand whether and

how epigenetic inheritance may lead to rapid adaption. Yet, assessing and teasing

apart the stochastic and deterministic route is experimentally challenging.

Experiments that aim to test the deterministic route are relatively common,

e.g. Baugh & Day (2020), Huber et al. (2021), Luna et al. (2012), Wibowo et al.

(2016). Usually, genetically uniform organisms are grown for multiple generations

under different environments ("pre-treatment") as single descendants - thereby

confounding genetic effects as well as selection of beneficial (epi-)genetic variants

are almost completely avoided (Baugh & Day, 2020). Subsequently, individuals are

grown for at least three generations under control conditions to avoid that the

organisms were directly exposed to stress during their early development (Heard &

Martienssen, 2014). Subsequently, fitness and epigenetic variation between the pre-

treatments are assessed. Ideally this experiment is accompanied with genetic or

chemical manipulation of the epigenetic machinery and/or genes that are

transgenerationally regulated to test the molecular mechanisms. While this approach

is powerful to test the role of deterministic epialleles in stress adaptation, it does not

allow to assess whether selection of stochastic variants may lead to stress

adaptation.

Experiments that test whether stochastic epigenetic variants may mediate

rapid stress adaption through natural selection are rather rare, but see Heckwolf et

al. (2020); Huber et al. (2021), Kronholm et al. (2017), Schmid et al. (2018), likely

because of the experimental challenges. The most common approach is to grow

large, genetically uniform populations for many generations under different

environments ("pre-treatment"), subsequently grow individuals for a few generations

under a shared control environment to erase environment-induced effects, and then

compare fitness and epigenetic variation between the populations of the different

pre-treatments. The limitation of this approach is that during the pre-treatment,

populations simultaneously experience both selection of stochastic variants as well

as the induction of deterministic epialleles – and if the deterministic epialleles are

heritable for multiple generations, and show unexpected temporal inheritance

patterns (Bell & Hellmann, 2019; Huber et al., 2021) – variation between the

populations could be either due to selection of stochastic variants or heritable

deterministic epialleles. To assess whether stochastic epigenetic variation may

mediate rapid adaption through natural selection, we thus need an approach in

which selection of stochastic variants can take place while teasing apart the

contribution of deterministic epialleles.

The solution: manipulating the efficacy of selection through the population

size

Here, we propose an experimental approach that unifies both of the above-

mentioned setups and thereby allows to simultaneously assess whether selection of

stochastic variants and/or the formation of deterministic epialleles mediate rapid

stress adaptation. One of the key differences between the stochastic and

deterministic route is whether selection is needed – and as such, to differentiate

among these two routes, one should manipulate the efficacy of selection. This can

be achieved through the population size, as selection is most effective in large

populations (McDonald, 2019; Wahl et al., 2002). We therefore propose to subject

populations for several generations to different environments ("pre-treatments") at

two different population sizes: on the one hand, in population sizes sufficiently large

that selection is effective. These populations may undergo both the stochastic and

deterministic route. On the other hand, in population sizes sufficiently small that

selection is not effective, i.e. that drift overcomes the effect of selection. This is

achieved once the effective population size $N_e < 1/s$ (s = selection coefficient)

(McDonald, 2019; Nielsen & Slatkin, 2013); the smallest possible effective population

size is achieved in single descendant lineages. Thus, the small population will evolve

almost exclusively through drift, and thereby only undergo the deterministic route.

Importantly, the large and small populations must be grown side by side in the very

same environment to ensure that the deterministic route is equally induced in the

small and large populations.

After several generations of pre-treatment, one can assess whether

populations adapted to stress through the stochastic and/or deterministic route. To

this end, individuals of the small and large populations of the different pre-treatments

are grown in a shared control environment for at least three generations to ensure

that any variation is truly heritable. Subsequently, phenotypes and fitness under the

different environments are compared. This allows differentiating whether selection of

epigenetic variants and/or deterministic epialleles lead to rapid adaptation: if

deterministic epialleles confer resistance, variation in resistance and epigenetic

variation will arise between the small populations of the different pre-treatments. If

selection of epigenetic variants contributes to resistance, variation in resistance and

in epigenetic variation will establish between the small and large population within

each pre-treatment (Fig. 1b).

(a) Routes of epigenetic adaptation

(b) Experimental approach

Figure 1. The two routes of rapid adaptation through epigenetic inheritance. (a)
Epigenetic inheritance may facilitate adaptation either via deterministic epialleles
(deterministic route) or selection of stochastic epigenetic variants (stochastic route).
Each circle and triangle indicate the DNA methylation level of a specific locus of
different individuals. (b) To differentiate among the deterministic and stochastic route,
organisms are grown in two different environments (pre-treatments) in large and small
populations; after a multi-generational recovery phase in a shared control environment,
fitness, phenotypes and epigenetic marks in the presence and absence of recurring
stress are compared between populations sizes and pre-treatments.

The requirements: is my experimental system suitable for the approach?

While the proposed experiment is in principle applicable to species across the tree of

life, the species under investigation should have following characteristics:

1. *Rapid reproduction and small body size.* As with all species used for

experimental evolution, rapid reproduction and small body size are highly

desirable.

2. *Low genetic variation.* Teasing apart genetic from epigenetic effects is

notoriously difficult, see discussion below. We thus suggest minimizing

genetic variation by using clonally reproducing organisms or organisms that

are highly inbred; ideally, genetic mutation rates should be small to reduce

genetic variation that arise during the experiment.

3. *Accurate assessments of fitness and epigenetic variation.* Measuring

Darwinian fitness is critical when assessing adaptation – thus, direct fitness

assessments (e.g. number of offspring) is preferred over indirect assessments

such as phenotypes or altered performance of interacting species; such

assessments are nevertheless valuable. Furthermore, the ability to measure

epigenetic variation (e.g. DNA methylation, histone modification, small RNAs)

in a largely unbiased manner and to link these variations to specific genes is

highly desirable. Although these analyses often require a reference genome,

new developments in high-throughput sequencing will likely overcome some

of the limitations of non-model species that do not have a high-quality

reference genome or whose genome is prohibitively large (Amarasinghe et

al., 2020; Gawehns et al., 2022).

Proof-of-concept: applying the method to assess rapid adaptation in

monoclonal duckweed populations

A species that is ideal to test adaptation through the deterministic or stochastic route

is the giant duckweed, *Spirodela polyrhiza,* one of the smallest and fastest growing

flowering plants. *Spirodela polyrhiza* reproduces almost exclusively asexually by

budding with a maximal doubling time of two to three days (Landolt, 1957; Ziegler et

al., 2015). The plant consists of a flat, thallus-like shoot, the so-called frond, which

allows to adequately estimate plant fitness by estimating growth rates based on

surface area. The species has a very low mutation rate with approximately one point

mutation every eighty generations (Xu et al., 2019). In nature, *S. polyrhiza* is

exposed to diverse environmental stresses including copper excess, which is a

common pollutant in aquatic ecosystems (Greger, 1999; Li & Xiong, 2004).

To test whether *S. polyrhiza* acquires stress resistance through the stochastic or

deterministic route, we grew monoclonal *S. polyrhiza* populations in the presence

and absence of copper excess ("pre-treatment") in both small and large populations

("selection regimen") for approximately seven generations in ponds outdoors and

assessed plant resistance to copper excess after five generations of growth in a

shared control environment. The small populations consisted of single descendant

lineages – the smallest possible population size – which were floating freely inside

the large populations (Fig 2A).

On May 2021, we set up 20 ponds (80 x 60 x 32cm, AuerPackaging,

Amerang, Germany) inside an open field site in Münster, Germany (51°57'54.0'N,

7°36'22.4'E). Ponds were buried to the rim with soil and filled with 10 L of pond soil

(Floragard, Oldenburg, Germany), 40 mL of organic fruit and vegetable fertilizer

(organic NPK 3,1+0,5+4,1; COMPO, Münster, Germany) and tap water with a total

volume of 100 L. The boxes had a draining scape and lids consisting of a stainless-

steel mesh (mesh opening 0.63 mm, wire diameter, 0.224 mm; Haver&Boeker,

Oelde, Germany). On May 26th, each box received around 3000 duckweed fronds

of the genotype 7498 that were pre-cultivated inside 10% N-medium (Appenroth et

al., 1996) for one month in indoor growth cabinets (GroBank, CLF PlantClimatics,

Wertingen, Germany) operating at 26°C, 135 μmol photons m^{-2} s^{-1} supplied through

LEDs with 16:8 h light:dark cycles. After one week of outdoor acclimation, we added

20 µM CuSO$_4$ to half of the ponds (n=10) to initiate the stress pre-treatment (Fig.

2a). Across the entire experiments, we measured copper concentration weekly and

after each rainfall; if needed, CuSO$_4$ was added to maintain 20 µM CuSO$_4$. After

adding CuSO$_4$ for the first time, we randomly chose four plants per pond as the

founders of the single descendant lineages. To trace the single descendant lineages,

we marked the plants with a dot using a permanent marker (Stabilo OHPen

universal) (Fig. 2a). The remaining plants reproduced freely and covered the surface

after 12 days, which corresponds to four to five generations of growth and a maximal

population size of 25,000 individuals. To avoid crowding, we randomly removed 30%

of the plants. Subsequently, we also marked four plants from the populations and

followed them for two generations as single descendant lineages to equalize the

marking treatment and the plant age between the single descendants and large

populations. To grow the plants from the different pre-treatments and selection

regimens in a uniform environment ("recovery phase"), we moved the third marked

generation of the population and the sixth generation of the single descendants

(approx. 24 days of pre-treatment) into separate cavities of polyvinyl chloride disks (7

cm diameter) that were floating inside the control ponds. Due to algae bloom, these

plants did not reproduce any offspring. We thus transferred the very same plants

back to the initial indoor conditions after 38 days of growth outdoors. To establish a

shared environment among plants from the different pre-treatments and selection

regimens, we placed the plants from the single descendant lineages and the large

populations from each control and copper pre-treatment pond pair together inside a

250mL polypropylene beakers (Plastikbecher GmbH, Giengen an der Brenz,

Germany) filled with 150 mL10% N-medium and covered with perforated and

transparent lids (four groups per beaker). Plants from the four different groups were

marked with specific colors to follow the individual lineages; colors were randomized

among treatments. To reduce algal growth, we changed the medium every two to

three days and wrapped the sides of the beakers with three layers of black plastic foil

to decrease light incidence. Inside the plastic beakers, we propagated each lineage

for five generations using single descendant propagation. To assess plant fitness

and resistance to copper excess, we moved the first and second daughter plant of

generation five into separate 250 ml polypropylene beakers containing control

conditions (10% N-medium) or recurrent copper excess (20 µM $CuSO_4$ in 10% N-

medium); the first and second offspring were randomly assigned to the treatment

groups. After setting up the experiment, we took a picture for surface area analysis

using a camera installation with a webcam (HD Pro Webcam C270, Logitech,

Lausanne, Switzerland; webcam software 2.12.8). To reduce algae growth, we

exchanged the medium after four days, and we took a second picture after eight

days of growth.

Statistical analyses

We calculated growth rates based on frond surface area [(End value)-(Initial

value)]/days. Additionally, we estimated resistance to copper excess as the ratio in

the surface area of plants grown under copper excess relative to the mean value of

plants grown under control treatment). To test whether plant resistance and growth

rates under control conditions and copper excess are affected by the selection

regimen and the pre-treatment, we fitted a linear mixed effects model using selection

regimen and pre-treatment as fixed effects, and as random effects the pair of a

control and a copper pre-treatment ponds ("pair") and blocks of ponds (groups of

three to four ponds, "block") regarding their physical position in the field

(~Selection*Pre-treatment+(1|pair)+(1|block)). Models were run in R v4.3.1 using the

package lme4 v1.1.34 (Bates et al., 2015), and *P*-values of the fixed effects were

estimated with Type III Analysis of Variance with the Satterthwaite´s method using

anova of the package stats (R Core Team, 2023). We used the packages DHARMa

v0.4.6 (Hartig, 2022) to verify whether the scaled residuals of our fitted models

correlated to the simulated values from the same model, and effects v4.2.2 (Fox &

Weisberg, 2018) to predict the effects of the fitted model. When analyzing plant

resistance and fitness under copper excess, we found an influential sample in the

copper pre-treated large population under copper environment, which showed

Cook´s distance higher than 0.5, low correlation between the standardized residuals

and theoretical quantiles (Q-Q plot) and whose corresponding sibling in the control

environment was lost. Therefore, we excluded this outlier from the analysis. Plots

were displayed with ggplot2 v3.4.3 (Wickham, 2016) and data was organized with

readxl v1.4.2 (Wickham & Bryan, 2023), data.table v1.14.8 (Dowle & Srinivasan,

2023), tidytext v0.4.1 (Silge & Robinson, 2016) and dplyr v1.1.2 (Wickham et al.,

2023). Figures were edited with Inkscape (Inkscape Project, 2018).

RESULTS

Proof-of-concept: duckweeds possibly adapt through both the stochastic and

deterministic epigenetic routes

To assess whether *S. polyrhiza* acquires resistance to copper excess through either

the deterministic or stochastic route, we measured plant resistance of single

descendants and large populations grown under control and copper pre-treatment,

the latter reduced plant growth rates outdoors by 16 % (Supporting Information Fig.

S1). Both the presence of selection and copper pre-treatment affected plant

resistance: regardless of the pre-treatment, plants from the large populations tended

to have higher resistance to copper excess than plants from the single descendant

lineages (Fig. 1b; selection: $F_{(1,3.33)} = 7.24$, $P = 0.066$, linear mixed effects model).

Furthermore, pre-treatment in copper stress tended to increase copper resistance in

the single descendants but not in the populations (Fig. 1b; selection x pre-treatment:

$F_{(1,12.78)} = 7.48$, $P = 0.017$, linear mixed effects model).

347   To differentiate whether the increase in resistance was due to lower growth

under control conditions or higher growth under copper excess, we analysed plant

fitness under both treatment environments separately. Under control conditions,

neither the population size nor the pre-treatment altered plant fitness (Fig. 2c;

$F_{(1,10.33)} = 0.50$, $P > 0.1$, linear mixed effects models). In contrast, under copper

excess, plants from the large populations outperformed plants from the single

descendant lineages regardless of the pre-treatment (Fig 2d; selection: $F_{(1,3.68)} =$

8.34, $P = 0.049$, mixed effects models). Furthermore, pre-treatment in copper stress

tended to increase plant fitness under copper excess in the single descendant

lineages but not in the populations (selection x pre-treatment: $F_{(1,12.79)} = 3.86$, $P =$

0.071, linear mixed effects models). Therefore, variations in copper resistance were

mainly due to variations in growth rates under copper excess and not under control

conditions.

(a) Experimental setup

(b) Resistance

(c) Control environment

(d) Copper environment

Figure 2. Testing the experimental approach by growing large and small monoclonal populations of Spirodela polyrhiza *in the presence and absence of copper excess in outdoor mesocosms. (a) Overview of the experimental setup. (b) Plant resistance to copper excess (growth rates under copper excess compared to the group´s mean growth rate under control conditions) depended on the population size and the pre-treatment. (c) Growth rates under control environment was not affected by the population size and the pre-treatment environment. (d) Growth rates under copper excess were affected by the population size and the pre-treatment. For b-c: P-values refer to linear mixed effects models. The black circles show the means, and the error bars the standard errors. N = 3-6. Pre-treat = pre-treatment; Sel = selection; Pop = large population; Sgl = single descendants.*

DISCUSSION

Here we presented an experimental approach to test a long-standing controversy:

whether epigenetic inheritance can lead to rapid adaptation to environmental

stresses through stochastic or deterministic epigenetic variation. Our proof-of-

concept experiments suggests that both the deterministic and stochastic routes may

alter duckweed fitness under recurring stress.

Proof-of-concept experiment

Our experiment suggests that duckweed populations may acquire stress

resistance through the stochastic route, as, regardless of the pre-treatment, offspring

from large populations outperformed offspring from the single descendants. The

higher fitness of the large populations compared to single descendants may have

been caused by two factors: first, variations among single descendants and large

populations may have been confounded by our method to identify the single

descendants during the pre-treatment phase. To identify the single descendants, we

marked the single descendants throughout the pre-treatment phase. Although we

also marked plants from the large populations in the last two generations of the pre-

treatment, we cannot fully rule out that the prolonged marking of the single

descendants affected plant fitness even after the five generations of growth under

shared control conditions prior fitness assay. In this case, the difference between the

large populations and single descendants is not caused by selection (stochastic

route) but by the marking, a pre-treatment effect (deterministic route) that would

have persisted across five generations. Second, the higher fitness of the large

populations than of the single descendants may have been caused by selection of

stochastic variants. Our experiment was limited to five generations. While such short

durations are sufficient for natural populations to become adapted through selection

of standing genetic variations (Züst et al., 2012), it is unclear whether monoclonal

populations may acquire sufficient epigenetic variation for such rapid adaptation

taking place. Future experiments that assess variation in plant fitness among single

descendant lineages, and long-term experiments that ensure that small and large

populations experience the very same environment during the pre-treatment phase

will be needed to differentiate among these two possibilities.

While the largest differences in fitness was caused by the different population

size, we also obtained evidence for deterministic effects (interaction of population

size and pre-treatment on plant resistance): copper pre-treatment increased fitness

of stress single descendants and decreased fitness of the population plants. This is

in line with recent studies suggesting that the previous environment alters plant

fitness and epigenetic marks across multiple clonal generations (Anastasiadi et al.,

2021; Benson et al., 2021; Colinas et al., 2023; González et al., 2018; Li et al., 2018;

Sammarco et al., 2023). Future manipulative studies are needed to test for specific

mechanisms of the deterministic route to confirm the presence of this phenomenon

and to understand whether the observed routes are truly caused by epigenetic

inheritance *sensu stricto* (genome-associated mechanisms) or by other non-genetic

inheritance mechanisms.

While the proposed experiment is not designed to tease apart the specific

epigenetic mechanism, the approach can – if carried out carefully – resolve the

contribution of the deterministic and stochastic route in a wide range of organisms. In

addition, the approach can also be used to assess the burdens of genetic mutation

accumulation, as proposed previously (Ho et al., 2021; Ho et al., 2020). We would

like to highlight a few points to make maximal use of this experimental approach:

Inferring the stochastic and deterministic route

The above-described experiment with duckweeds reveals a critical issue: the pre-

treatment environment of the small and large population must be equal, otherwise

one cannot differentiate between the stochastic and deterministic route. In our

experiment, this could be achieved by caging small populations inside the large

population – this would alleviate the potential bias of the marking. However, one

must ensure that the small populations still experience the same environment as the

large populations despite the caging. In duckweeds, this can be achieved by

separating the small populations from the large populations using floating disks while

keeping the density of the small and large populations equal. It is of utmost

importance to carefully design of the experiments considering the specific biology of

each organism.

So far, we treated the stochastic and deterministic route as separate, non-

overlapping entities. This may be an unrealistic assumption. For instance,

deterministic epialleles may not be induced in an entirely deterministic manner;

environmental stresses may simply increase the likelihood of an epiallele to appear

but not cause all individuals changing their epigenetic status. Selection could thus

still act on environment-induced "deterministic" epialleles. Similar accounts for the

stochastic epimutations: environmental stresses may increase the likelihood that

certain epimutations appear, associated to the presence of epigenetic hotspots

(Hazarika et al., 2022; Zheng et al., 2017), and thus epimutations may not be entirely

stochastic. Are the stochastic and deterministic route therefore two extremes of the

same process? One argument holds against this: in *A. thaliana*, spontaneous

epimutations mostly accumulate in the CG context (Becker et al., 2011; Denkena et

al., 2021; Schmitz et al., 2011; van der Graaf et al., 2015) whereas environment-

induced epialleles are most prominent in the CHG and/or CHH context (Annacondia

et al., 2021; Lin et al., 2022; Wibowo et al., 2016; Yadav et al., 2022; Zhou et al.,

2019). This supports the idea that the stochastic and deterministic route are, at least

in the context of DNA methylation in plants, two fundamentally different processes.

Inferring the molecular mechanisms

Another critical issue is the interpretation of the results: is the underlying

molecular mechanism of genetic or epigenetic nature? Even when using highly

inbred or clonally reproducing species, genetic mutations may accumulate and

account for variation in phenotypes or fitness, particularly in the large populations.

We suggest following experiments and analyses to tease apart genetic from

epigenetic effects: first, unbiased methods (e.g. high-throughput sequencing) should

be deployed to screen for epigenetic variations. Second, transcriptome and

proteome data should be generated to link fitness to phenotypes to gene expression

and epigenetic variation. Third, differentially regulated genes or epigenetic

machineries should be manipulated both genetically as well as chemically to test

whether the identified genes or epigenetic machinery affect stress resistance or the

process of stress adaptation. Fourth, phenotypes or fitness should be continuously

assessed for multiple generations after stress release – if variations between groups

have an epigenetic basis, the variation should diminish or at least change in some

lineages over time. Fifth, high-throughput sequencing should be deployed to screen

for genetic variation in identified genomic regions. While in isolation none of these

approaches will provide a clear answer whether genetic or epigenetic mechanisms

are at play, combining these approaches will be powerful to disentangle genetic and

epigenetic factors.

CONCLUSIONS

Taken together, we here provide an experimental framework to assess the relative

contribution of deterministic epialleles and selection of stochastic epigenetic variants

to rapid stress adaptation. Through this approach, carefully designed experiments

can provide novel insights whether and how epigenetic inheritance may lead to rapid

stress adaptation. Considering the on-going loss of intra-specific genetic diversity

and the rapid pace of environmental change, these insights may become

increasingly relevant when assessing species potential to resist global change.

ACKNOWLEDGEMENTS

We would like to thank Pauline Prüsener, Anke Grosse-Brinkhaus, Maite Görtz

Lizarraga and the Botanical Garden of the University of Münster for supporting the

outdoor experiment and Shuqing Xu for providing facilities. This study profited from

interactions within the GenEvo Research Training Group 407023052/GRK2526/1

funded by the German Research Foundation (DFG). The work was funded by the

Volkswagen Foundation (Grant Nr 97236) to MH and the German Research

Foundation to MH (grant Nr. 512079118). The project was supported by the

University of Münster and the University of Mainz.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

AUTHOR CONTRIBUTION

MH conceived the ideas and designed methodology, AC collected the data, AC and

MH analysed the data, MH and AC drafted the manuscript. Both authors contributed

critically to the drafts and gave final approval for publication.

STATEMENT ON INCLUSION

This study was discussed and carried out by a group of scientists originating from

different continents and by technicians from the local region. All authors were

engaged in the design of the experiments to ensure that a diverse set of

perspectives were included. Efforts were made to consider relevant work published

in the local language.

DATA AVAILABILITY

The raw data of the outdoor experimental evolution experiment will be archived on Dryad upon acceptance.

REFERENCES

Amarasinghe, S.L., Su, S., Dong, X., Zappia, L., Ritchie, M.E. & Gouil, Q. (2020). Opportunities and challenges in long-read sequencing data analysis. *Genome Biology,* **21,** 30.

Anastasiadi, D., Venney, C.J., Bernatchez, L. & Wellenreuther, M. (2021). Epigenetic inheritance and reproductive mode in plants and animals. *Trends in Ecology & Evolution,* **36,** 1124-1140.

Annacondia, M.L., Markovic, D., Reig-Valiente, J.L., Scaltsoyiannes, V., Pieterse, C.M.J., Ninkovic, V., Slotkin, R.K. & Martinez, G. (2021). Aphid feeding induces the relaxation of epigenetic control and the associated regulation of the defense response in *Arabidopsis. New Phytologist,* **230,** 1185-1200.

Appenroth, K.J., Teller, S. & Horn, M. (1996). Photophysiology of turion formation and germination in *Spirodela polyrhiza. Biologia Plantarum,* **38,** 95.

Baugh, L.R. & Day, T. (2020). Nongenetic inheritance and multigenerational plasticity in the nematode *C. elegans. eLife,* **9,** e58498.

Becker, C., Hagmann, J., Müller, J., Koenig, D., Stegle, O., Borgwardt, K. & Weigel, D. (2011). Spontaneous epigenetic variation in the *Arabidopsis thaliana* methylome. *Nature,* **480,** 245-249.

Bell, A.M. & Hellmann, J.K. (2019). An integrative framework for understanding the

mechanisms and multigenerational consequences of transgenerational

plasticity. *Annual Review of Ecology, Evolution and Systematics,* **50,** 97-118.

Benson, C.W., Mao, Q. & Huff, D.R. (2021). Global DNA methylation predicts

epigenetic reprogramming and transgenerational plasticity in Poa annua L.

Crop Science, **61,** 3011-3022.

Bonduriansky, R. & Day, T. (2009). Nongenetic inheritance and its evolutionary

implications. *Annual Review of Ecology, Evolution and Systematics,* **40,** 103-

125.

Colinas, N., Montero-Pau, J., Carmona, M.J., Sabatino, R., Di Cesare, A., Eckert,

E.M. & García-Roger, E.M. (2023). Transgenerational expression profiles of a

sex related and an epigenetic control gene in the rotifer Brachionus plicatilis in

relation to environmental predictability. *Hydrobiologia.*

Cubas, P., Vincent, C. & Coen, E. (1999). An epigenetic mutation responsible for

natural variation in floral symmetry. *Nature,* **401,** 157-161.

Denkena, J., Johannes, F. & Colomé-Tatché, M. (2021). Region-level epimutation

rates in *Arabidopsis thaliana. Heredity,* **127,** 190-202.

Des Roches, S., Pendleton, L.H., Shapiro, B. & Palkovacs, E.P. (2021). Conserving

intraspecific variation for nature's contributions to people. *Nature Ecology &*

Evolution, **5,** 574-582.

Dowle, M. & Srinivasan, A. (2023). data.table: Extension of `data.frame`.

Fox, J. & Weisberg, S. (2018). Visualizing Fit and Lack of Fit in Complex Regression

Models with Predictor Effect Plots and Partial Residuals. *Journal of Statistical*

Software, **87,** 1 - 27.

Gawehns, F., Postuma, M., van Antro, M., Nunn, A., Sepers, B., Fatma, S., van

Gurp, T.P., Wagemaker, N.C.A.M., Mateman, A.C., Milanovic-Ivanovic, S.,

Groβe, I., van Oers, K., Vergeer, P. & Verhoeven, K.J.F. (2022). epiGBS2:

Improvements and evaluation of highly multiplexed, epiGBS-based reduced

representation bisulfite sequencing. *Molecular Ecology Resources,* **22,** 2087-

2104.

González, A.P.R., Preite, V., Verhoeven, K.J.F. & Latzel, V. (2018).

Transgenerational effects and epigenetic memory in the clonal plant *Trifolium*

repens. Frontiers in Plant Science, **9,** 1677-1677.

Greger, M. (1999). Metal Availability and Bioconcentration in Plants. *Heavy Metal*

Stress in Plants: From Molecules to Ecosystems (eds M.N.V. Prasad & J.

Hagemeyer), pp. 1-27. Springer Berlin Heidelberg, Berlin, Heidelberg.

Hartig, F. (2022). DHARMa: Residual Diagnostics for Hierarchical (Multi-Level /

Mixed) Regression Models.

Hazarika, R.R., Serra, M., Zhang, Z., Zhang, Y., Schmitz, R.J. & Johannes, F.

(2022). Molecular properties of epimutation hotspots. *Nature Plants,* **8,** 146-

156.

Heard, E. & Martienssen, R.A. (2014). Transgenerational epigenetic inheritance:

myths and mechanisms. *Cell,* **157,** 95-109.

Heckwolf, M.J., Meyer, B.S., Häsler, R., Höppner, M.P., Eizaguirre, C. & Reusch,

T.B.H. (2020). Two different epigenetic information channels in wild three-

spined sticklebacks are involved in salinity adaptation. *Science advances,* **6,**

eaaz1138-eaaz1138.

Ho, E.K.H., Bellis, E.S., Calkins, J., Adrion, J.R., Latta Iv, L.C. & Schaack, S. (2021).

Engines of change: Transposable element mutation rates are high and

variable within Daphnia magna. *PLOS Genetics,* **17,** e1009827.

Ho, E.K.H., Macrae, F., Latta, L.C., 4th, McIlroy, P., Ebert, D., Fields, P.D., Benner,

M.J. & Schaack, S. (2020). High and Highly Variable Spontaneous Mutation

Rates in Daphnia. *Molecular Biology and Evolution,* **37,** 3258-3266.

Huber, M., Gablenz, S. & Höfer, M. (2021). Transgenerational non-genetic

inheritance has fitness costs and benefits under recurring stress in the clonal

duckweed *Spirodela polyrhiza. Proceedings of the Royal Society of London,*

Series B: Biological Sciences, **288**.

Inkscape Project. (2018). Inkscape.

Jiang, C., Mithani, A., Belfield, E.J., Mott, R., Hurst, L.D. & Harberd, N.P. (2014)

Environmentally responsive genome-wide accumulation of de novo

Arabidopsis thaliana mutations and epimutations. *Genome Research,* **24,**

1821-1829.

Johannes, F. & Schmitz, R.J. (2019). Spontaneous epimutations in plants. *New*

Phytologist, **221,** 1253-1259.

Kronholm, I., Bassett, A., Baulcombe, D. & Collins, S. (2017). Epigenetic and genetic

contributions to adaptation in Chlamydomonas. *Molecular Biology and*

Evolution, **34,** 2285-2306.

Landolt, E. (1957). Physiologische und ökologische Untersuchungen an Lemnaceen.

Bericht der Schweizerischen Botanischen Gesellschaft, **67,** 271–410.

Li, K., Chen, J., Wei, Q., Li, Q. & Lei, N. (2018). Effects of Transgenerational

Plasticity on Morphological and Physiological Properties of Stoloniferous Herb

Centella asiatica Subjected to High/Low Light. *Frontiers in Plant Science,* **9**.

Li, T. & Xiong, Z. (2004). A novel response of wild-type duckweed (Lemna

593   paucicostata Hegelm.) to heavy metals. *Environmental Toxicology,* **19,** 95-

594   102.

595   Lin, X., Zhou, M., Yao, J., Li, Q.Q. & Zhang, Y.-Y. (2022). Phenotypic and

596   Methylome Responses to Salt Stress in Arabidopsis thaliana Natural

597   Accessions. *Frontiers in Plant Science,* **13**.

598   Luna, E., Bruce, T.J.A., Roberts, M.R., Flors, V. & Ton, J. (2012). Next-generation

599   systemic acquired resistance. *Plant Physiology,* **158,** 844-853.

600   Manning, K., Tör, M., Poole, M., Hong, Y., Thompson, A.J., King, G.J., Giovannoni,

601   J.J. & Seymour, G.B. (2006). A naturally occurring epigenetic mutation in a

602   gene encoding an SBP-box transcription factor inhibits tomato fruit ripening.

603   *Nat Genet,* **38,** 948-952.

604   McDonald, M.J. (2019). Microbial experimental evolution - a proving ground for

605   evolutionary theory and a tool for discovery. *EMBO reports,* **20,** e46992-

606   e46992.

607   Nielsen, R. & Slatkin, M. (2013). *An introduction to population genetics: theory and*

608   *applications.* Sinauer.

609   O'Dea, R.E., Noble, D.W.A., Johnson, S.L., Hesselson, D. & Nakagawa, S. (2016).

610   The role of non-genetic inheritance in evolutionary rescue: epigenetic

611   buffering, heritable bet hedging and epigenetic traps. *Environmental*

612   *Epigenetics,* **2**.

613   Ong-Abdullah, M., Ordway, J., Jiang, N., Ooi, S., Kok, S.Y., Sarpan, N., Azimi, N.,

614   Hashim, A., Ishak, Z., Rosli, S., Ahmad Malike, F., Abu Bakar, N., Marjuni, M.,

615   Abdullah, N., Yaakub, Z., Din, A., Rajanaidu, N., Singh, R., Low, L. &

Martienssen, R. (2015) Loss of Karma transposon methylation underlies the mantled somaclonal variant of oil palm. *Nature,* **525,** 533-537.

Sage, R.F. (2020). Global change biology: A primer. *Global Change Biology,* **26,** 3-30.

Sammarco, I., Díez Rodríguez, B., Galanti, D., Nunn, A., Becker, C., Bossdorf, O., Münzbergová, Z. & Latzel, V. (2023). DNA methylation in the wild: epigenetic transgenerational inheritance can mediate adaptation in clones of wild strawberry (Fragaria vesca). *Research Square.*

Schmid, M.W., Heichinger, C., Coman Schmid, D., Guthörl, D., Gagliardini, V., Bruggmann, R., Aluri, S., Aquino, C., Schmid, B., Turnbull, L.A. & Grossniklaus, U. (2018). Contribution of epigenetic variation to adaptation in *Arabidopsis. Nature Communications,* **9,** 4446.

Schmitz, R.J., Schultz, M.D., Lewsey, M.G., O'Malley, R.C., Urich, M.A., Libiger, O., Schork, N.J. & Ecker, J.R. (2011). Transgenerational epigenetic instability is a source of novel methylation variants. *Science,* **334,** 369-373.

Silge, J. & Robinson, D. (2016). tidytext: Text Mining and Analysis Using Tidy Data Principles in R. *Journal of Open Source Software,* **1,** 37.

R Core Team. (2023). R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.

Van Antro, M., Prelovsek, S., Ivanovic, S., Gawehns, F., Wagemaker, N.C.A.M., Mysara, M., Horemans, N., Vergeer, P. & Verhoeven, K.J.F. (2023). Stability of temperature-induced DNA methylation changes across clonal generation in the common Duckweed *Lemna minor* L. *Molecular Ecology.*

van der Graaf, A., Wardenaar, R., Neumann, D.A., Taudt, A., Shaw, R.G., Jansen, R.C., Schmitz, R.J., Colome-Tatche, M. & Johannes, F. (2015). Rate,

spectrum, and evolutionary dynamics of spontaneous epimutations.

Proceedings of the National Academy of Sciences, USA, **112,** 6676-6681.

Wahl, L.M., Gerrish, P.J. & Saika-Voivod, I. (2002). Evaluating the impact of

population bottlenecks in experimental evolution. *Genetics,* **162,** 961-971.

Wibowo, A., Becker, C., Marconi, G., Durr, J., Price, J., Hagmann, J., Papareddy, R.,

Putra, H., Kageyama, J., Becker, J., Weigel, D. & Gutierrez-Marcos, J. (2016).

Hyperosmotic stress memory in Arabidopsis is mediated by distinct

epigenetically labile sites in the genome and is restricted in the male germline

by DNA glycosylase activity. *eLife,* **5,** e13546.

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis,* 2 edn. Use R!

Wickham, H. & Bryan, J. (2023) readxl: Read Excel Files.

Wickham, H., François, R., Henry, L., Müller, K. & Vaughan, D. (2023) dplyr: A

Grammar of Data Manipulation.

Xu, S., Stapley, J., Gablenz, S., Boyer, J., Appenroth, K.J., Sree, K.S., Gershenzon,

J., Widmer, A. & Huber, M. (2019). Low genetic variation is associated with

low mutation rate in the giant duckweed. *Nature Communications,* **10,** 1243.

Yadav, N.S., Titov, V., Ayemere, I., Byeon, B., Ilnytskyy, Y. & Kovalchuk, I. (2022).

Multigenerational Exposure to Heat Stress Induces Phenotypic Resilience,

and Genetic and Epigenetic Variations in Arabidopsis thaliana Offspring.

Frontiers in Plant Science, **13**.

Zhang, H., Lang, Z. & Zhu, J.-K. (2018). Dynamics and function of DNA methylation

in plants. *Nature Reviews: Molecular Cell Biology,* **19,** 489-506.

Zheng, X., Chen, L., Xia, H., Wei, H., Lou, Q., Li, M., Li, T. & Luo, L. (2017).

Transgenerational epimutations induced by multi-generation drought

imposition mediate rice plant's adaptation to drought condition. *Scientific*

Reports, **7,** 39843.

Zhou, M., Sng, N.J., LeFrois, C.E., Paul, A.-L. & Ferl, R.J. (2019). Epigenomics in an

extraterrestrial environment: organ-specific alteration of DNA methylation and

gene expression elicited by spaceflight in Arabidopsis thaliana. *BMC*

Genomics, **20,** 205.

Ziegler, P., Adelmann, K., Zimmer, S., Schmidt, C. & Appenroth, K.J. (2015).

Relative in vitro growth rates of duckweeds (Lemnaceae) - the most rapidly

growing higher plants. *Plant Biol (Stuttg),* **17 Suppl 1,** 33-41.

Züst, T., Heichinger, C., Grossniklaus, U., Harrington, R., Kliebenstein, D.J. &

Turnbull, L.A. (2012). Natural enemies drive geographic variation in plant

defenses. *Science,* **338,** 116-119.