<u>**Prior heat stress increases pathogen susceptibility in the model cnidarian *Exaptasia diaphana***</u>

Sofia C. Diaz de Villegas[1], Erin. M. Borbee[1], Peyton Y. Abdelbaki[1], Lauren E. Fuess[1*]

[1]Texas State University, San Marcos, TX, USA
[*]**Corresponding Author:** lfuess@txstate.edu

Abstract

Anthropogenic climate change has significantly altered terrestrial and marine ecosystems globally, often in the form of climate-related events such as thermal anomalies and disease outbreaks. Although the isolated effects of these stressors have been well documented, a growing body of literature suggests that stressors often interact, resulting in complex effects on ecosystems. Recent declines in coral cover from a variety of climate-related stressors make reef ecosystems well-suited to investigate sequential associations between climate impacts, particularly heat stress and disease. Here we used the model cnidarian *Exaiptasia diaphana* to investigate mechanisms linking heat stress to increased disease susceptibility. We examined anemone pathogen susceptibility and physiology (symbiosis, immunity, and energetics) following recovery from heat stress. We observed significantly increased pathogen susceptibility in anemones exposed to heat stress, likely linked to observed suppression of multiple immunological parameters. Furthermore, our data suggest that changes in resource allocation might drive observed reductions in immunity following thermal stress. These findings are an important first step towards understanding temporal associations between two of the most significant climate-associated stressors, heat and disease.

Keywords
cnidarian, heat stress, symbiosis, disease, multiple stressors, immunity

Introduction

Anthropogenic pressure has led to environmental and ecological changes in terrestrial and aquatic ecosystems across the world, often in the form of climate-related events. (Hansen & Stone, 2015; Rosenzweig et al., 2008). Marine ecosystems, in particular, have been significantly affected by multiple climate-induced stressors, such as ocean warming and disease outbreaks (Doney et al., 2011; Hoegh-Guldberg & Bruno, 2010; Scavia et al., 2002). Both of these stressors directly affect organismal physiology through alteration of metabolic function, and indirectly affect a variety of ecological processes, such as oxygen availability, nutrient cycling, and ocean circulation (Doney et al., 2011; Smith et al., 2023). As such, ocean warming, specifically marine heatwaves, are a leading cause of mass mortality in marine organisms (Genin et al., 2020; Rivetti et al., 2014; Smale, 2020; Wild et al., 2019). Similarly, increases in the frequency and severity of disease outbreaks pose significant threats to marine ecosystems. Marine taxa are affected by a large diversity of pathogens ranging from viruses and bacteria to fungi and metazoans (Bateman et al., 2020), many of which have triggered significant mortality events in a variety of marine ecosystems (Hamilton et al., 2021; Patterson, 1996; Petton et al., 2015; Sanderson & Alexander, 2020; Short et al., 1987).

Although the isolated effects of ocean warming, disease outbreaks, and other climate-associated stressors have been well documented within marine ecosystems (Doney et al., 2011; Hoegh-Guldberg & Bruno, 2010), a growing body of literature suggests that these stressors often coincide. Depending on the circumstances and stressors in question, interactions may range from cumulative to synergistic, or even antagonistic in nature (Adams, 2005; Crain et al., 2008). Associations between heat stress and disease outbreaks are particularly prevalent, and have often contributed to extensive marine mortality (Harvell et al., 2002). Consequently, studies have attempted to improve understanding of associations between heat stress and disease in marine ecosystems, but most have focused only on simultaneous synergy between the two stressors (Ban et al., 2014; Harvell et al.; Olsen & Duarte, 2015). However, frequent increases in disease outbreaks in the weeks to months following marine heatwaves suggest that these two stressors may more commonly accumulate in a sequential, rather than simultaneous, manner, (Case et al., 2011; Godwin et al., 2020; Marcogliese, 2008; Vezzulli et al., 2013). Despite these observations, the mechanisms driving sequential temporal patterns between heat stress and disease are poorly understood and necessitate further investigation.

Coral reef ecosystems present a unique opportunity to investigate sequential associations
between heat stress and disease. These ecosystems are commonly faced with a variety of climate
stressors (Smith & Buddemeier, 1992), which have resulted in unprecedented declines in coral
cover over the past several decades (Bruno & Selig, 2007; Gardner et al., 2003). Increasing sea
surface temperature (SST), in particular, has exacerbated the frequency and severity of coral
decline. Two major drivers of these declines are heat-induced bleaching—loss of endosymbiotic
algae (Brown, 1997)—and disease (Harvell et al., 2007). Although isolated heat-induced
bleaching (Johnston et al., 2019; McClanahan et al., 2007; Winter et al., 1998) and disease are
common (Richardson et al., 1998; Subhan et al., 2020), these events have also been frequently
observed in close spatial and temporal associations (Brandt & McManus, 2009; Burke et al.,
2023; Harvell et al., 2001; Howells et al., 2020). In many cases, coral disease follows bleaching
(Croquer & Weil, 2009), with extensive disease-related mortality often reported in the weeks to
months after heat stress (Jones et al., 2021; Miller et al., 2009; Miller et al., 2006). Despite these
observations, nearly all experimental studies have focused on the synergistic effects of
simultaneous heat stress and disease, specifically documenting increases in disease susceptibility
when these stressors occur at the same time (Palmer, 2018; Zaragoza et al., 2014). Consequently,
many questions remain regarding the mechanisms linking sequential associations between heat
stress and disease outbreaks. The increasing prevalence of heat-induced bleaching events and
disease outbreaks highlight the need for an improved understanding of how biotic and abiotic
stressors interact through time within the context of changing coral reef environments.

While tropical scleractinian corals are exceptionally important from an ecological standpoint,
their precarious species statuses and inherent experimental limitations complicate the laboratory
investigation of complex biological phenomenon in these species. Consequently, there has been a
rise in the development of closely related model species with traits better suited to experimental
manipulation. Of these cnidarian models, *Exaiptasia diaphana*, commonly known as Aiptasia, is
a species of anemone from the same class as scleractinian corals (Hexacorallia). Aiptasia is
widely regarded for its ecological abundance and generation of genetically identical clonal lines,
allowing for control of genetic variation in experimentation (Baumgarten et al., 2015).
Furthermore, like scleractinian corals, *E. diaphana* hosts endosymbiotic algae of the family
Symbiodiniaceae (LaJeunesse et al., 2018). We leveraged the *E. diaphana* model system to
investigate the mechanisms linking heat stress and disease susceptibility in cnidarians.

Specifically, we examined host immune and energetic responses to pathogen exposure following

heat-stress in two genetically distinct clonal lines of *E. diaphana*. Here we document a

significant association between heat stress and increased pathogen susceptibility and highlight

the potential physiological mechanisms of this association. The results described here are an

important first step towards understanding temporal links between two of the most significant

climate-associated stressors faced by cnidarians.

Methods

Experimental Design

Anemones were obtained from laboratory-maintained clonal lines of *E. diaphana* (VWB9

and H2; V. Weis). Anemones from each clonal line were randomly assigned to one of four

possible treatment combinations: control or heat stress and placebo or live bacteria. Individuals

were transferred accordingly into sterile 6-well culture plates filled with 10 mL of filter-sterile

artificial sea water (FASW). Control plates were maintained in a temperature-controlled Percival

incubator at 27°C for the duration of the experiment. Plates assigned to heat treatment were

randomly divided between two separate temperature-controlled incubators designated for heat

stress. Prior to the start of the experiment, all individuals were acclimated to well plates for three

days at 27°C in their respective incubators. Throughout the entire experiment, anemones in all

three incubators were maintained under the same light conditions (12hr:12hr light: dark cycles

with approximately 40-60 µmol·m^{-2} PAR) and received bi-weekly feeding and water changes

until the point of infection, Plate locations were randomized within and between incubators at the

time of water changes, as appropriate.

Experimental Heat Stress & Immune Challenge

Heat stress consisted of a seven-day ramping period where temperatures were raised 1°C per

day until they reached 34°C. Temperatures were then maintained at 34°C for five days based on

data from pilot studies **(Supplemental Table 1).** During the experiment, bleaching was

confirmed by visual assessment of anemone color and was found to be consistent with the

expected timeline. Following heat stress, anemones were immediately returned to 27°C and

allowed to recover for two weeks. At the conclusion of this recovery period, surviving anemones

from both control and heat treatments were inoculated with either a pathogen or a placebo

(FASW) treatment according to *a priori* assignments. Pathogen exposure was conducted using

the known coral pathogen *V. coralliilyticus* BAA 450 (Ben-Haim et al., 2003) at a predetermined

LD50 dosage of 1 x 10^8 CFU·mL^{-1} (Zaragoza et al., 2014). Final bacterial concentrations were

confirmed with serial dilutions and plating. Following exposure, a subset of anemones was

monitored for survivorship (**Supplemental Table 2**): mortality data was collected every two

hours for the first 36 hours, and then daily for the remainder of the experiment (96 hours post-

infection) at which point 89% of exposed anemones had died. Individuals were considered dead

when they exhibited more than 75% tissue degradation or had 75% tissue degradation for more

than 24 hours.

Sample Collection and Processing

Twelve hours after inoculation, a second subset of randomly selected anemones was sampled

for the assessment of symbiont density, immune activity, and energetic reserves (**Supplemental**

Table 2). Anemones were anesthetized in 10 mL of $MgCl_2$ (0.37 M) prior to sampling. First, we

excised a single tentacle for estimates of symbiont density. Subsequent tentacle processing was

modified from existing protocols (Tortorelli et al., 2020). Following tentacle excision, the

remaining anemone was halved using a sterile razor blade; one half was flash frozen for the

immune and energetic assays described herein. Frozen tissue samples were homogenized with 3

mm glass beads in 1.5 mL of 100 mM Tris + 0.05 M DTT buffer (pH 7.8) for 1 minute at 5 m·s^{-1}

(Tim Bateman, *pers. comm.*) using a bead mill (Fisherbrand). For each sample, a 200 µL aliquot

of homogenate was transferred into a separate microcentrifuge tube and stored at -20°C for

melanin assays. The remaining homogenate for each sample was then transferred into new 2 mL

microcentrifuge tubes and centrifuged at 2250xg at 4°C for five minutes. Resultant sample

supernatants were aliquoted accordingly for each set of assays (protein, lipid, carbohydrates, ash-

free dry weight) and stored at -80°C (protein fraction) or -20°C (all other fractions) prior to

downstream analyses.

Symbiont Density Estimation

Mounted tentacles were imaged using brightfield and fluorescence microscopy on the

Cytation1 Cell Imaging Multimode Reader (Biotek). Approximate tentacle area was determined

via brightfield microscopy. Symbiont autofluorescence was measured using a chlorophyll a

fluorescent filter (Agilent; EX 445/45 nm, EM 685/40 nm). Brightfield and fluorescent images of

each tentacle were combined to standardize fluorescence by tentacle area. The average pixel
intensity of the chlorophyll a fluorescence within each standardized tentacle area was used as a
proxy for symbiont density.

Immune Assays

Cnidarian immunity was measured using a suite of standardized immune assays originally
developed for scleractinian corals (Mydlarz & Palmer, 2011; Pinzón C et al., 2014) and adapted
for use in *E. diaphana* (**Supplemental File 1**). We measured responses related to three major
innate immune pathways: phenoloxidase, antibacterial, and antioxidant pathways. The
phenoloxidase pathway is an enzymatic cascade resulting in the synthesis of melanin – an
effector molecule involved in wound healing and pathogen encapsulation (Mydlarz et al., 2016).
Here we quantify the phenoloxidase cascade using both total phenoloxidase activity (enzymatic
activity of prophenoloxidase + phenoloxidase) and melanin concentration. An additional
essential component of coral immunity is the production of antimicrobial peptides and
metabolites that can neutralize a wide variety of pathogenic bacteria (Mydlarz et al., 2016;
Palmer & Traylor-Knowles, 2018). Antibacterial activity here was quantified by measuring
bacterial growth rates of *Vibrio coralliilyticus* in the presence of protein extracts generated from
host tissue. Finally, we also measured activity of the important antioxidant catalase, using
established methods (Pinzón et al., 2014). Cnidarians produce antioxidant enzymes as a
mechanism to protect against reactive oxygen species generated by both hosts and pathogens
during infection and associated immune responses (Parisi et al., 2020)

All immune assays were run in triplicate on separate 96-well plates and measured using the
Cytation1 Cell Imaging Multimode Reader and Gen5 Software (Biotek). 100mM Tris + 0.05
mM DTT buffer was used as a negative control for all assays. All enzymatic immune assays
were standardized by protein concentration using Red660 colorimetric assays, while melanin
concentration was standardized by desiccated tissue weight.

Energetic Assays

Total lipids and total carbohydrates assays were used to determine approximate host
energetic budget. Lipids and carbohydrates are commonly used proxies for energetic budgets in
multiple cnidarian species (Lesser, 2013). Lipid and carbohydrate concentrations were measured
and standardized by ash-free dry weight following modifications to previously established

protocols (Cheng et al., 2011; Fitt et al., 2000; Masuko et al., 2005). Total lipid and total

carbohydrate assays were run in triplicate and duplicate, respectively, and measured in separate

96-well plates using the Cytation1 Cell Imaging Multimode Reader and Gen5 Software (Biotek).

Tris + DTT buffer was used as a negative control for both assays.

Statistical Analyses

All statistical analyses were performed using R Version 4.3.0 (2023-04-21; R Core Team

2023). Prior to hypothesis testing and modeling sample outliers were identified using the Rosner

test from the EnvStats package (v2.7.0; Millard, SP., 2013) and removed. Shapiro-Wilke and

Bartlett tests were used to confirm the assumptions of normal distribution and equal variances,

respectively. Several assays failed to meet assumptions necessary for parametric testing, even

following transformation, necessitating the use of non-parametric analyses.

Mortality

Survivorship data from our first subset of anemones were used to assess the effects of

pathogen exposure, heat stress and genotype (clonal line) on mortality. First, the effects of

pathogen exposure on mortality were assessed using a Cox proportional hazard's model

(Survival ~ immune challenge; n = 188; survival package v3.5.5; Therneau, T, 2023). We

then used a secondary Cox model to investigate the effects of heat stress and genotype on

mortality within the subset of pathogen-exposed anemones only (Survival ~ treatment +

clonal line; n = 90).

Univariate hypothesis testing

A second subset of anemones reserved for assessment of symbiont density, energetic

reserves, and immune activity, was used to investigate hypotheses related to mechanisms of

increased mortality. To start, we tested for differences in symbiont density (mean chlorophyll

fluorescence intensity) or energetic reserves (total lipids and total carbohydrates) between

treatment groups (control vs. heat) and clonal lines. The impacts of heat stress or clonal line

on symbiont density, lipids, and carbohydrates were assessed independently with separate

Welch's two-sample t-tests (symbiont density and lipids) or non-parametric Wilcoxon Rank

Sum tests (carbohydrates).

Redundancy analysis

To investigate the impact of our independent variables (heat stress, pathogen exposure,
and clonal line) and covariates (symbiont density, total lipids, and total carbohydrates) on
host immune activity, we ran a redundancy analysis (RDA) using the vegan package (v2.6.4;
Oksanen, J. et al., 2022). Before starting with this multivariate approach, we tested for and
confirmed no collinearity among our variables. All continuous variables were centered and
scaled before separation into predictor (i.e., symbiont density, lipids, carbohydrates) and
response (i.e., catalase, total phenoloxidase, and antibacterial activity, and melanin
concentration) data frames. An RDA was used to identify which variables best predicted
variation in immunity across our samples. We started by running an RDA with all possible
predictor variables (i.e. clonal line, temperature treatment, pathogen exposure, lipids, and
carbohydrates). From there, we used a forward selection to identify which predictor variables
explained significant variation in our dataset. After identifying significant predictor variables,
we ran a final RDA using the best-fit model determined by forward selection (~ Treatment +
Clonal Line + Immune Challenge + Lipids).

Linear modeling

Finally, we used linear modeling to assess variation in individual immune parameters
with respect to the statistically relevant factors determined via RDA. For normally distributed
immune parameters (catalase and antibacterial activity) we used general linear models
(GLMs). Global models with all possible additive and interactive variables were analyzed for
catalase and antibacterial activity independently. All models were dredged and ranked via
AICc using the MuMIn package (v1.47.5; Barton, K., 2023), where applicable. Models with
delta AICc values less than or equal to 2 were averaged to determine the best-fit models for
each immune parameter. For non-normally distributed parameters, we used a similar
approach with modifications to account for unequal variances and zero-inflated data. To
account for unequal variances within total phenoloxidase data we used the nlme package
(v3.1.162; Pinheiro, J. et al., 2023) to run generalized least squares (GLS) modeling with the
same model dredging and averaging approach as described above. Finally, for melanin, we
adopted a two-tiered strategy that was motivated by the high number of zero measurements
(see results). First, we used a probit model (i.e. a generalized linear model with binomial

error structure) to predict the presence of melanin among samples using heat treatment,
clonal line, immune challenge and lipid concentration as predictors. We then removed
samples in which melanin was not detected and used multiple regression to predict melanin
concentrations employing the same predictors.

Results

Mortality

Anemone mortality was significantly greater in anemones exposed to pathogens (Cox
proportional hazards model; hazard ratio = 103; 95% CI [25 – 421]; $p < 0.001$; **Supplemental**
Fig. 1). Furthermore, within pathogen exposed anemones, mortality rate was significantly greater
in the heat stressed anemones relative to controls (hazard ratio = 2.15; 95% CI [1.34 – 3.46]; $p =$
0.002) and among H2 anemones relative to VWB9 (hazard ratio = 0.26; 95% CI [0.16-0.42]; $p <$
0.001; **Fig. 1**).

Symbiont Density & Energetic Reserves

Symbiont density, estimated by mean chlorophyll a fluorescence pixel intensity, was
marginally, but not significantly, higher in heat stressed anemones compared to controls
(Welch's two sample t-test; t = -1.93, df = 68.7, $p = 0.0579$. 95% CI [-3740 – 63.3]; **Fig. 2a**).
However, symbiont density did significantly vary across clonal lines (Welch's two sample t test;
t = 2.04, df = 49.4, $p = 0.0472$; 95% CI [26.5 – 4100]); H2 had significantly greater symbiont
density than VWB9 anemones **(Fig. 2b).** For energetic reserves, total carbohydrates varied
between treatment groups. Carbohydrates were significantly lower in heat treated anemones
compared to control anemones (Wilcoxon Rank Sum test; W = 1460, $p < 0.01$; **Fig. 2c**) and in
the VWB9 clonal line compared to the H2 clonal line (Wilcoxon Rank Sum test; W = 733, p =
0.0116; **Fig. 2d**). Total lipids did not significantly differ between either temperature treatment
(Welch's two sample t-test; t = -0.945, df = 77.3, $p = 0.348$, 95% CI [0.0774 – 0.0846], **Fig 2e**)
or clonal lines (Welch's two sample t-test; t = -0.436, df = 84.0, $p = 0.664$, 95% CI [-0.0184 –
0.0117], **Fig 2f**).

Immune Responses

The RDA explained significant variation along the first three axes of the ordination. These
axes explained 14.32% (RDA1), 7.84% (RDA2), and 7.57% (RDA3) for a cumulative proportion

explained of 29.74% (**Fig. 3**). Treatment (ANOVA, $p = 0.001$), clonal line (ANOVA, $p = 0.001$),

pathogen exposure (ANOVA, $p = 0.003$), and lipid concentration (ANOVA, $p = 0.032$) all

explained significant variation in immune responses along these axes. Temperature treatment and

lipid concentration explained variation mostly along the first RDA axis, while clonal line and

pathogen exposure explained variation along both the first and second RDA axes. When

considering immune responses, the variables that drove the variation along the first axis (RDA1,

14.32%) were catalase activity and melanin concentration while PPO activity drove the variation

on the second axis (RDA2, 7.84%).

Linear modeling of the effects of each of our significant predictors on individual immune

metrics revealed metric specific effects (**Table 1**). Both catalase activity and melanin

concentration were significantly lower in heat treated anemones relative to control temperatures

(Catalase GLM, $p < 0.001$; Melanin GLM, $p < 0.001$; **Fig. 4).** Catalase activity was also

significantly different between clonal lines (GLM, $p < 0.001$); VWB9 anemones had

significantly greater catalase activity than H2 (**Supplemental Fig. 2**). Both antibacterial activity

and melanin concentration were significantly correlated with lipid concentration (**Fig 5**).

Notably, antibacterial activity was significantly different as a result of the interaction between

heat treatment and lipid concentration (GLM, p < 0.01; **Fig. 5a**). Heat stressed anemones had a

significant negative association between lipids and antibacterial activity (LM; $R^2_{adj} = 0.103$, F(1,

25) = 4.66 , $p = 0.0396$). However, control anemones maintained a positive, but insignificant,

association between lipids and antibacterial activity (LM; $R^2_{adj} = 0.0557$, F(1, 24) = 3.196 , $p =$

0.0865). While heat treatment predicted the presence or absence of melanin (GLM, $p < 0.001$),

melanin concentration, when present, was significantly positively associated with lipid

concentration independent of other predictors (0s removed GLM, $p < 0.001$; **Fig. 5b**). Total

phenoloxidase was the only metric to vary as a result of pathogen exposure; activity was reduced

in anemones following pathogen exposure (GLS, $p < 0.01$; **Supplemental Fig. 3**).

Discussion

As anthropogenic climate change worsens, marine ecosystems are faced with a multitude of

stressors that may form complex associations with each other (Doney et al., 2011; Hoegh-

Guldberg & Bruno, 2010). However, empirical study of interactive stressors, particularly those

occurring asynchronously, has been limited, leaving many questions regarding the nature and

mechanisms of associations amongst stressors (Pirotta et al., 2022). Cnidarians, which face a
diversity of climate related stressors(Ban et al., 2014; Maor-Landaw & Levy, 2016) , provide a
relevant model to investigate the effects of multiple stressors on marine species. Here we
conducted one of the first experimental studies investigating the consequences of heat stress on
cnidarian pathogen susceptibility. Our results support a long held theory (Muller et al., 2018) that
previous heat stress increases disease susceptibility and provide important new insight regarding
the mechanisms of these patterns. Specifically, we highlight changes in host cnidarian immunity
following heat stress which are likely driven by shifting resource allocation which may
contribute to observed patterns of differential mortality.

Mortality differs across temperature treatments and clonal lines

Susceptibility to pathogen exposure was significantly higher among those anemones exposed
to heat stress. This result is in agreement with documented increases in disease and disease-
related mortality following coral bleaching in the wild (Harvell et al., 2001; Matsuda et al., 2020;
Miller et al., 2009; Precht et al., 2016; Sakai et al., 2019; Wright et al., 2017). Notably, while
previous laboratory experiments have documented impacts of simultaneous heat stress and
pathogen exposure (Muller et al., 2018; Zaragoza et al., 2014) ours is the first to document
extended consequences of heat stress on pathogen susceptibility, more accurately representing
observed environmental patterns in nature. Future studies combining our approach with emergent
experimental disease techniques (Huntley et al., 2022; Traylor-Knowles et al., 2021) will allow
for more broadscale applications to complex cnidarian and marine disease systems.

We also observed significant genotypic differences in mortality. Anemones from the VWB9
clonal line were less susceptible to pathogen-induced mortality than H2. Both clonal lines
originate from similar geographic regions and host the same species of algal endosymbiont
(*Brevolium minutum*) (Thornhill et al., 2013), suggesting that fine scale genetic variation likely
contributes to these differences. While intraspecific variation in cnidarian disease resistance has
been linked to genetic variation in the wild (Miller et al., 2019; Vollmer & Kline, 2008), there
are a lack of studies investigating intraspecific variation in laboratory settings (Wright et al.,
2017) . Similarly, few experimental studies investigating disease dynamics in Aiptasia
incorporate genetic variation (Krediet et al., 2014; Zaragoza et al., 2014)**.** Our findings suggest
that inclusion of genetic variation in cnidarian studies, including those involving *E. diaphana*,
may provide increased insight regarding the roles of genetic variability in contributing to

resistance or susceptibility to stress.

Mechanisms of differential mortality: roles of symbiont density and energetic reserves

Beyond documenting differences in mortality as a result of environmental and genetic

factors, we employed an integrative approach to investigate the underlying mechanisms driving

these differences. First, we tested for differences in symbiont density and energetic reserves

between heat stressed and control anemones. Both symbiont densities and energetic reserves may

be affected during heat stress events (Rodrigues & Grottoli, 2007). Furthermore, changes in

symbiont densities or energetic reserves may impact host immunity (Changsut et al., 2022;

Valadez-Ingersoll et al., 2023), and consequently pathogen susceptibility (Palmer et al., 2010).

We failed to detect any differences in symbiont density after two weeks of recovery, likely due

to *E. diaphana*'s capacity for exceptionally rapid recovery from dysbiosis (Tortorelli et al.,

2020). Thus, the two weeks prior to pathogen exposure in which anemones were allowed to

recover from heat stress was likely ample time for symbiont densities to partially recover in our

heat stress group. In contrast, we did observe differences in carbohydrate concentration as a

result of heat stress: anemones from heat stress treatments had significantly lower carbohydrate

concentration than anemones from control treatments. Thus, while symbiont density recovers

quickly, carbohydrate concentration appears to recover more slowly, potentially contributing to

increased pathogen susceptibility. More fine-scale approaches leveraging metabolomic or

carbon-flux modeling may be useful in determining the complex effects of heat stress on host

energetic budget, and subsequent effects on pathogen susceptibility.

Mechanisms of differential mortality: roles of host immunity

Next, we focused on the potential roles of shifts in immunological activity which might

contribute to observed patterns of mortality. When considering immunity holistically using

multivariate statistics, we saw significant effects of temperature, clonal line, immune challenge,

and lipid concentration on immune response. Here we discuss the impacts of temperature and

lipid concentration on immunity as these factors had the most prevalent impacts on holistic

immunity and individual immune parameters. Examination of the effects of heat stress on

individual immune parameters revealed strong suppression of both catalase activity and melanin

concentration. While numerous studies have evidenced the short term effects of heat stress (days

– weeks) on cnidarian host immunity (Krueger et al., 2015; Palmer, 2018; Caroline V. Palmer et
al., 2011; Wall et al., 2018) the long-term effects of heat stress (months – years) on immune
activity are not well documented (Mydlarz et al., 2009; Wall et al., 2021). Generally, heat stress
is believed to induce activation of a variety of immune parameters, including antioxidant (e.g.,
catalase) and phenoloxidase enzymes (Krueger et al., 2015; Palmer, 2018). However, here we
document pronounced suppression of immunity (catalase, antibacterial activity, melanin) after
two weeks of recovery from heat stress, suggesting that immune activation is dampened during
recovery. Our approach provides new insight suggesting suppression manifests quickly after heat
stress and highlights the need for frequent, repeated sampling during bleaching and recovery to
better understand the prolonged effects of increasing SSTs on cnidarian immunity.

Investigating the role of symbiotic immunosuppression
While immune suppression following thermal stress may be the result a number of
mechanisms, one of the most commonly proposed mechanisms involves dynamic changes in
symbiosis. Suppression of host immunity by symbionts has been hypothesized as an important
mechanism required for the successful establishment and maintenance of symbiotic relationships
(Mansfield & Gilmore, 2019; Merselis et al., 2018). Previous laboratory studies have confirmed
negative associations between symbiosis and immunity across several cnidarians (Detournay et
al., 2012; Fuess et al., 2020a; Fuess, et al., 2020b; Jacobovitz et al., 2021; Mansfield et al., 2017;
Merselis et al., 2018). Nonetheless, few studies have examined the dynamics of immune-
symbiosis interplay during the breakdown and re-establishment of symbiosis. It is possible that
the rapid repopulation of endosymbionts following heat-induced dysbiosis exacerbates symbiotic
immunosuppression, potentially explaining, in part, naturally observed patterns of disease after
thermal bleaching events. While symbiont density neither significantly differed between
temperature treatments, nor significantly predicted holistic immunity, it is possible that indirect
or time lagged effects of changing symbiont densities may be influencing observed patterns of
differential mortality. Specifically, rapid and excessive symbiont proliferation (evidenced by
marginally higher symbiont densities in heat treated anemones) may have had significant
prolonged effects on host immunity. Future studies incorporating time-series approaches to
sampling may be used to test this hypothesis.

Investigating the role of energetic immunosuppression and the importance of resource

allocation

A second potential hypothesis to explain observed associations between heat stress and
immunity is that changes in resource allocation as a result of resource limitations during
dysbiosis may result in immune suppression. According to resource allocation theory, organisms
must distribute a finite amount of resources amongst diverse biological processes—growth,
reproduction, metabolism, immunity, etc. (Ng'oma et al., 2017). This may involve trade-offs
where resources are shifted between competing needs in order to maximize fitness (Garland et
al., 2022; Zera & Harshman, 2001). Consequently, reduction in the total amount of resources
may alter the proportions in which resources are distributed, potentially resulting in the
suppression of one or more biological processes. As such, immune suppression is one of the
most commonly documented energetic trade-offs associated with resource limitation (Norris &
Evans, 2000; Rantala & Roff, 2005; Rauw, 2012; Sadd & Schmid-Hempel, 2009; van der Most
et al., 2011). In cnidarians, the loss of symbionts during a heat induced bleaching event may
severely limit the number of energetic resources available to the host. These limitations likely
exacerbate resource tradeoffs and result in changes to resource allocation, including allocation to
host immune responses. Alternatively, immune-energetic trade-offs may also occur in the
absence of resource limitation in response to biotic or abiotic pressures: organisms may
dynamically shift resources between processes in response to pressing needs caused by
environmental stress or other changes in biological condition (McNamara & Buchanan, 2005).

In congruence with this hypothesis, the relationship between antibacterial activity and lipids
differed as a result of temperature treatment potentially due to differences in resource allocation.
We observed a negative association between lipids and antibacterial activity only following heat
stress. Increased investment of lipids towards antibacterial activity under control conditions is
likely indicative of immunological "front-loading"; increased constitutive activity of potent
preventative immune metrics which confer increased pathogen resistance (Mydlarz et al., 2016).
However, under heat stress, lipids may be directed away from preventative mechanisms (i.e.
pathogen defense) in favor of more pressing stress response pathways. The most frequently
measured response to heat stress is production of antioxidant enzymes which mitigate oxidative
damage to host tissues (Krueger et al., 2015; Yakovleva et al., 2005). While we did not detect
significant associations between antioxidant enzymes and lipids, it is possible that resources
were allocated to other, unmeasured stress-related mechanisms such as fluorescent pigments,

mycosporine-like amino acids, or heat-shock proteins (Baird et al., 2009). In contrast, melanin

concentration was significantly positively associated with lipid concentration regardless of

temperature treatment, likely due to its multi-faceted roles in cnidarian host fitness (C. V. Palmer

et al., 2011; Palmer & Traylor-Knowles, 2018).

Other factors contributing to variation in immune activity

Finally, both genotype and pathogen exposure were also significant predictors of immune

activity. Specifically, catalase activity was significantly different between clonal lines: VWB9

anemones had higher catalase activity than H2 anemones. Given that antioxidant activity is an

essential component of coral immunity (Parisi et al., 2020), these differences may be related to

differential clonal line mortality. Additionally, total phenoloxidase activity was lower in

anemones exposed to pathogens. Rapid changes in immune activity, and in particular the

phenoloxidase cascade, have been observed following pathogen exposure in cnidarians (Caroline

V. Palmer et al., 2011). Notably, the phenoloxidase cascade results in the production of melanin

which has been linked to pathogen defense and wound healing (Mydlarz et al., 2010). Thus,

observed declines in phenoloxidase could indicate increased investment in melanin synthesis in

anemones exposed to pathogenic challenge. While we did not observe significant differences in

melanin concentrations between placebo and infected anemones, it is possible that sampling

timing (12 hours post infection) coincided with intermediate stages of the signaling cascade.

Conclusion & Future Directions

Multiple stressors are a common consequence of anthropogenic climate change and can have

pervasive effects on a variety of marine ecosystems (Adams, 2005; Crain et al., 2008). As such,

an improved understanding of organismal responses to climate-induced stressors is imperative to

the conservation of these ecosystems. Here we detail one of the first studies to investigate the

mechanistic links between prior heat stress and increased disease susceptibility in a marine

species. Our study establishes the ability to reproduce heat stress and disease associations under

laboratory conditions using a tractable model system, allowing for more nuanced and robust

investigations of ecologically relevant phenomena. Using this experimental approach, we were

able to demonstrate prolonged effects of prior heat stress on host immunity, which may

contribute to observed pathogen-induced mortality. Furthermore, our results implicate changes in

resource allocation as a potential mechanism of heat-induced immunosuppression. Further

studies incorporating time series approaches will be necessary to fully disentangle the observed
patterns and associated mechanisms and assess their broad applicability to marine systems.
Specifically, future work should aim to sample across multiple timepoints from the onset of
bleaching through acute and long-term recovery to determine the effects of dynamic changes in
symbiont density and energetic reserves on host immunity and pathogen susceptibility.
Additional comparative experimentation using scleractinian corals and non-cnidarian species
will also help determine the applicability of our results to other systems. In summary, our results
provide an important starting point for understanding the mechanistic links between heat stress
and disease that will aid in the conservation of marine species, including cnidarians, in the face
of a changing climate. Most importantly, they generate testable hypotheses that will be
invaluable in better understanding the effects of multiple stressors on diverse species and
ecosystem functions.
515

Acknowledgements
We would like to thank Sam Bedgood and Virginia Weis for providing anemone stock
populations as well as Tim Bateman for sharing Aiptasia protocols. Additionally, we would like
to thank Dr. Benjamin Martin and the Rhodes Lab at Texas State University for equipment
usage. Finally, we would like to acknowledge the Symbiommunity Lab members and Texas
State EEB discussion group for experimental assistance and data/manuscript feedback,
respectively. In particular, we thank Chris Nice and Jim Fordyce for their contributions to the
statistical analyses presented. Funding was provided by Texas State University (Startup to LEF)
and the National Academies' Gulf Research Program (Early Career Fellowship to LEF).
525

526   **Conflict of Interest**
527   The authors declare they have no competing interests.
528

Author Contributions
SCD was responsible for experimental design, sample collection, data analysis and
interpretation, and drafting/revising the manuscript. LEF developed the study concept,
contributed to experimental design and data analysis, conducted critical revision of the

manuscript and provided funding for the study. EMB supported statistical analyses, graphical

generation, and manuscript revision. PYA assisted with data collection.

Literature Cited

Adams, S. M. (2005). Assessing cause and effect of multiple stressors on marine systems. *Marine Pollution Bulletin*, *51*(8), 649-657. https://doi.org/https://doi.org/10.1016/j.marpolbul.2004.11.040

Baird, A. H., Bhagooli, R., Ralph, P. J., & Takahashi, S. (2009). Coral bleaching: the role of the host. *Trends in Ecology & Evolution*, *24*(1), 16-20. https://doi.org/https://doi.org/10.1016/j.tree.2008.09.005

Ban, S. S., Graham, N. A., & Connolly, S. R. (2014). Evidence for multiple stressor interactions and effects on coral reefs. *Glob Chang Biol*, *20*(3), 681-697. https://doi.org/10.1111/gcb.12453

Bateman, K. S., Feist, S. W., Bignell, J. P., Bass, D., & Stentiford, G. D. (2020). Marine pathogen diversity and disease outcomes. In *Marine Disease Ecology* (pp. 0). Oxford University Press. https://doi.org/10.1093/oso/9780198821632.003.0001

Baumgarten, S., Simakov, O., Esherick, L. Y., Liew, Y. J., Lehnert, E. M., Michell, C. T., Li, Y., Hambleton, E. A., Guse, A., Oates, M. E., Gough, J., Weis, V. M., Aranda, M., Pringle, J. R., & Voolstra, C. R. (2015). The genome of Aiptasia, a sea anemone model for coral symbiosis. *Proceedings of the National Academy of Sciences*, *112*(38), 11893-11898. https://doi.org/10.1073/pnas.1513318112

Ben-Haim, Y., Thompson, F., Thompson, C., Cnockaert, M., Hoste, B., Swings, J., & Rosenberg, E. (2003). Vibrio coralliilyticus sp. nov., a temperature-dependent pathogen of the coral Pocillopora damicornis. *International journal of systematic and evolutionary microbiology*, *53*(1), 309-315.

Brandt, M. E., & McManus, J. W. (2009). Disease incidence is related to bleaching extent in reef-building corals. *Ecology*, *90*(10), 2859-2867. https://doi.org/10.1890/08-0445.1

Brown, B. E. (1997). Coral bleaching: causes and consequences. *Coral Reefs*, *16*(1), S129-S138. https://doi.org/10.1007/s003380050249

Bruno, J. F., & Selig, E. R. (2007). Regional Decline of Coral Cover in the Indo-Pacific: Timing, Extent, and Subregional Comparisons. *PLOS ONE*, *2*(8), e711. https://doi.org/10.1371/journal.pone.0000711

Burke, S., Pottier, P., Lagisz, M., Macartney, E. L., Ainsworth, T., Drobniak, S. M., & Nakagawa, S. (2023). The impact of rising temperatures on the prevalence of coral diseases and its predictability: A global meta-analysis. *Ecology Letters*, *26*(8), 1466-1481. https://doi.org/https://doi.org/10.1111/ele.14266

Case, R. J., Longford, S. R., Campbell, A. H., Low, A., Tujula, N., Steinberg, P. D., & Kjelleberg, S. (2011). Temperature induced bacterial virulence and bleaching disease in a chemically defended marine macroalga. *Environ Microbiol*, *13*(2), 529-537. https://doi.org/10.1111/j.1462-2920.2010.02356.x

Changsut, I., Womack, H. R., Shickle, A., Sharp, K. H., & Fuess, L. E. (2022). Variation in symbiont density is linked to changes in constitutive immunity in the facultatively symbiotic coral, *Astrangia poculata*. *Biology Letters*, *18*(11), 20220273. https://doi.org/doi:10.1098/rsbl.2022.0273

Cheng, Y.-S., Zheng, Y., & VanderGheynst, J. S. (2011). Rapid quantitative analysis of lipids using a colorimetric method in a microplate format. *Lipids*, *46*(1), 95-103. https://doi.org/10.1007/s11745-010-3494-0

Crain, C. M., Kroeker, K., & Halpern, B. S. (2008). Interactive and cumulative effects of multiple human stressors in marine systems. *Ecol Lett*, *11*(12), 1304-1315.

584   https://doi.org/10.1111/j.1461-0248.2008.01253.x

585 Croquer, A., & Weil, E. (2009). Changes in Caribbean coral disease prevalence after the 2005
586   bleaching event. *Dis Aquat Organ*, *87*(1-2), 33-43. https://doi.org/10.3354/dao02164

587 Doney, S. C., Ruckelshaus, M., Emmett Duffy, J., Barry, J. P., Chan, F., English, C. A., Galindo,
588   H. M., Grebmeier, J. M., Hollowed, A. B., Knowlton, N., Polovina, J., Rabalais, N. N.,
589   Sydeman, W. J., & Talley, L. D. (2011). Climate Change Impacts on Marine Ecosystems.
590   *Annual Review of Marine Science*, *4*(1), 11-37. https://doi.org/10.1146/annurev-marine-
591   041911-111611

592 Fitt, W. K., McFarland, F. K., Warner, M. E., & Chilcoat, G. C. (2000). Seasonal patterns of
593   tissue biomass and densities of symbiotic dinoflagellates in reef corals and relation to
594   coral bleaching. *Limnology and Oceanography*, *45*(3), 677-685.
595   https://doi.org/https://doi.org/10.4319/lo.2000.45.3.0677

596 Gardner, T. A., Côté, I. M., Gill, J. A., Grant, A., & Watkinson, A. R. (2003). Long-Term
597   Region-Wide Declines in Caribbean Corals. *Science*, *301*(5635), 958-960.
598   https://doi.org/10.1126/science.1086050

599 Genin, A., Levy, L., Sharon, G., Raitsos, D. E., & Diamant, A. (2020). Rapid onsets of warming
600   events trigger mass mortality of coral reef fish. *Proc Natl Acad Sci U S A*, *117*(41),
601   25378-25385. https://doi.org/10.1073/pnas.2009748117

602 Godwin, S. C., Fast, M. D., Kuparinen, A., Medcalf, K. E., & Hutchings, J. A. (2020). Increasing
603   temperatures accentuate negative fitness consequences of a marine parasite. *Scientific
604   Reports*, *10*(1), 18467. https://doi.org/10.1038/s41598-020-74948-3

605 Hamilton, S. L., Saccomanno, V. R., Heady, W. N., Gehman, A. L., Lonhart, S. I., Beas-Luna,
606   R., Francis, F. T., Lee, L., Rogers-Bennett, L., Salomon, A. K., & Gravem, S. A. (2021).
607   Disease-driven mass mortality event leads to widespread extirpation and variable
608   recovery potential of a marine predator across the eastern Pacific. *Proc Biol Sci*,
609   *288*(1957), 20211195. https://doi.org/10.1098/rspb.2021.1195

610 Hansen, G., & Stone, D. (2015). Assessing the observed impact of anthropogenic climate
611   change. *Nature Climate Change*, *6*(5), 532-537. https://doi.org/10.1038/nclimate2896

612 Harvell, C. D., Mitchell, C. E., Ward, J. R., Altizer, S., Dobson, A. P., Ostfeld, R. S., & Samuel,
613   M. D. (2002). Climate Warming and Disease Risks for Terrestrial and Marine Biota.
614   *Science*, *296*(5576), 2158-2162. https://doi.org/10.1126/science.1063699

615 Harvell, C. D., Montecino-Latorre, D., Caldwell, J. M., Burt, J. M., Bosley, K., Keller, A.,
616   Heron, S. F., Salomon, A. K., Lee, L., Pontier, O., Pattengill-Semmens, C., & Gaydos, J.
617   K. Disease epidemic and a marine heat wave are associated with the continental-scale
618   collapse of a pivotal predator (Pycnopodia helianthoides). *Science Advances*, *5*(1),
619   eaau7042. https://doi.org/10.1126/sciadv.aau7042

620 Harvell, D., Jordán-Dahlgren, E., Merkel, S., Rosenberg, E., Raymundo, L., Smith, G., Weil, E.,
621   & Willis, B. (2007). Coral disease, environmental drivers, and the balance between coral
622   and microbial associates. *Oceanography*, *20*, 172-195.

623 Harvell, D., Kim, K., Quirolo, C., Weir, J., & Smith, G. (2001). Coral bleaching and disease:
624   contributors to 1998 mass mortality in Briareum asbestinum (Octocorallia, Gorgonacea).
625   *Hydrobiologia*, *460*(1), 97-104. https://doi.org/10.1023/A:1013169331913

626 Hoegh-Guldberg, O., & Bruno, J. F. (2010). The impact of climate change on the world's marine
627   ecosystems. *Science*, *328*(5985), 1523-1528. https://doi.org/10.1126/science.1189930

628 Howells, E. J., Vaughan, G. O., Work, T. M., Burt, J. A., & Abrego, D. (2020). Annual
629   outbreaks of coral disease coincide with extreme seasonal warming. *Coral Reefs*, *39*(3),

771-781. https://doi.org/10.1007/s00338-020-01946-2

Huntley, N., Brandt, M. E., Becker, C. C., Miller, C. A., Meiling, S. S., Correa, A. M. S., Holstein, D. M., Muller, E. M., Mydlarz, L. D., Smith, T. B., & Apprill, A. (2022). Experimental transmission of Stony Coral Tissue Loss Disease results in differential microbial responses within coral mucus and tissue. *ISME Communications*, *2*(1), 46. https://doi.org/10.1038/s43705-022-00126-3

Johnston, M. A., Hickerson, E. L., Nuttall, M. F., Blakeway, R. D., Sterne, T. K., Eckert, R. J., & Schmahl, G. P. (2019). Coral bleaching and recovery from 2016 to 2017 at East and West Flower Garden Banks, Gulf of Mexico. *Coral Reefs*, *38*(4), 787-799. https://doi.org/10.1007/s00338-019-01788-7

Jones, N. P., Kabay, L., Semon Lunz, K., & Gilliam, D. S. (2021). Temperature stress and disease drives the extirpation of the threatened pillar coral, Dendrogyra cylindrus, in southeast Florida. *Scientific Reports*, *11*(1), 14113. https://doi.org/10.1038/s41598-021-93111-0

Krediet, C. J., Meyer, J. L., Gimbrone, N., Yanong, R., Berzins, I., Alagely, A., Castro, H., Ritchie, K. B., Paul, V. J., & Teplitski, M. (2014). Interactions between the tropical sea anemone Aiptasia pallida and Serratia marcescens, an opportunistic pathogen of corals. *Environ Microbiol Rep*, *6*(3), 287-292. https://doi.org/10.1111/1758-2229.12151

Krueger, T., Hawkins, T. D., Becker, S., Pontasch, S., Dove, S., Hoegh-Guldberg, O., Leggat, W., Fisher, P. L., & Davy, S. K. (2015). Differential coral bleaching—Contrasting the activity and response of enzymatic antioxidants in symbiotic partners under thermal stress. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, *190*, 15-25. https://doi.org/https://doi.org/10.1016/j.cbpa.2015.08.012

LaJeunesse, T. C., Parkinson, J. E., Gabrielson, P. W., Jeong, H. J., Reimer, J. D., Voolstra, C. R., & Santos, S. R. (2018). Systematic Revision of Symbiodiniaceae Highlights the Antiquity and Diversity of Coral Endosymbionts. *Current biology: CB*, *28*(16), 2570-2580.e2576. https://doi.org/10.1016/j.cub.2018.07.008

Lesser, M. P. (2013). Using energetic budgets to assess the effects of environmental stress on corals: are we measuring the right things? *Coral Reefs*, *32*(1), 25-33. https://doi.org/10.1007/s00338-012-0993-x

Maor-Landaw, K., & Levy, O. (2016). Survey of Cnidarian Gene Expression Profiles in Response to Environmental Stressors: Summarizing 20 Years of Research, What Are We Heading for? In S. Goffredo & Z. Dubinsky (Eds.), *The cnidaria, past, present and future: the world of medusa and her sisters* (pp. 523-543). Springer International Publishing. https://doi.org/10.1007/978-3-319-31305-4_32

Marcogliese, D. (2008). The impact of climate change on the parasites and infectious diseases of aquatic animals. *Revue scientifique et technique (International Office of Epizootics)*, *27*, 467-484.

Masuko, T., Minami, A., Iwasaki, N., Majima, T., Nishimura, S., & Lee, Y. C. (2005). Carbohydrate analysis by a phenol-sulfuric acid method in microplate format. *Anal Biochem*, *339*(1), 69-72. https://doi.org/10.1016/j.ab.2004.12.001

Matsuda, S. B., Huffmyer, A. S., Lenz, E. A., Davidson, J. M., Hancock, J. R., Przybylowski, A., Innis, T., Gates, R. D., & Barott, K. L. (2020). Coral Bleaching Susceptibility Is Predictive of Subsequent Mortality Within but Not Between Coral Species. *Frontiers in Ecology and Evolution*, *8*. https://www.frontiersin.org/articles/10.3389/fevo.2020.00178

McClanahan, T. R., Ateweberhan, M., Muhando, C. A., Maina, J., & Mohammed, M. S. (2007).

Effects of Climate and Seawater Temperature Variation on Coral Bleaching and Mortality. *Ecological Monographs*, *77*(4), 503-525. https://doi.org/10.1890/06-1182.1

McNamara, J. M., & Buchanan, K. L. (2005). Stress, resource allocation, and mortality. *Behavioral Ecology*, *16*(6), 1008-1017. https://doi.org/10.1093/beheco/ari087

Miller, J., Muller, E., Rogers, C., Waara, R., Atkinson, A., Whelan, K. R. T., Patterson, M., & Witcher, B. (2009). Coral disease following massive bleaching in 2005 causes 60% decline in coral cover on reefs in the US Virgin Islands. *Coral Reefs*, *28*(4), 925. https://doi.org/10.1007/s00338-009-0531-7

Miller, J., Waara, R., Muller, E., & Rogers, C. (2006). Coral bleaching and disease combine to cause extensive mortality on reefs in US Virgin Islands. *Coral Reefs*, *25*(3), 418-418. https://doi.org/10.1007/s00338-006-0125-6

Miller, M. W., Colburn, P. J., Pontes, E., Williams, D. E., Bright, A. J., Serrano, X. M., & Peters, E. C. (2019). Genotypic variation in disease susceptibility among cultured stocks of elkhorn and staghorn corals. *PeerJ*, *7*, e6751. https://doi.org/10.7717/peerj.6751

Muller, E. M., Bartels, E., & Baums, I. B. (2018). Bleaching causes loss of disease resistance within the threatened coral species Acropora cervicornis. *eLife*, *7*, e35066. https://doi.org/10.7554/eLife.35066

Mydlarz, L. D., Couch, C. S., Weil, E., Smith, G., & Harvell, C. D. (2009). Immune defenses of healthy, bleached and diseased Montastraea faveolata during a natural bleaching event. *Diseases of Aquatic Organisms*, *87*(1-2), 67-78. https://doi.org/10.3354/dao02088

Mydlarz, L. D., Fuess, L., Mann, W., Pinzón, J. H., & Gochfeld, D. J. (2016). Cnidarian immunity: from genomes to phenomes. *The cnidaria, past, present and future: the world of medusa and her sisters*, 441-466.

Mydlarz, L. D., McGinty, E. S., & Harvell, C. D. (2010). What are the physiological and immunological responses of coral to climate warming and disease? *Journal of Experimental Biology*, *213*(6), 934-945. https://doi.org/10.1242/jeb.037580

Mydlarz, L. D., & Palmer, C. V. (2011). The presence of multiple phenoloxidases in Caribbean reef-building corals. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, *159*(4), 372-378. https://doi.org/https://doi.org/10.1016/j.cbpa.2011.03.029

Olsen, Y. S., & Duarte, C. M. (2015). Combined effect of warming and infection by Labyrinthula sp. on the Mediterranean seagrass Cymodocea nodosa. *Marine Ecology Progress Series*, *532*, 101-109. https://doi.org/10.3354/meps11343

Palmer, C. V. (2018). Warmer Water Affects Immunity of a Tolerant Reef Coral. *Frontiers in Marine Science*, *5*. https://www.frontiersin.org/articles/10.3389/fmars.2018.00253

Palmer, C. V., Bythell, J. C., & Willis, B. L. (2010). Levels of immunity parameters underpin bleaching and disease susceptibility of reef corals. *The FASEB Journal*, *24*(6), 1935-1946. https://doi.org/10.1096/fj.09-152447

Palmer, C. V., Bythell, J. C., & Willis, B. L. (2011). A comparative study of phenoloxidase activity in diseased and bleached colonies of the coral Acropora millepora. *Developmental & Comparative Immunology*, *35*(10), 1098-1101. https://doi.org/10.1016/j.dci.2011.04.001

Palmer, C. V., McGinty, E. S., Cummings, D. J., Smith, S. M., Bartels, E., & Mydlarz, L. D. (2011). Patterns of coral ecological immunology: variation in the responses of Caribbean corals to elevated temperature and a pathogen elicitor. *Journal of Experimental Biology*, *214*(24), 4240-4249. https://doi.org/10.1242/jeb.061267

Palmer, C. V., & Traylor-Knowles, N. G. (2018). Cnidaria: Anthozoans in the Hot Seat. In E. L.
Cooper & E. L. Cooper (Eds.), *Advances in comparative immunology* (pp. 51-93).
https://link.springer.com/10.1007/978-3-319-76768-0_3
Parisi, M. G., Parrinello, D., Stabili, L., & Cammarata, M. (2020). Cnidarian Immunity and the
Repertoire of Defense Mechanisms in Anthozoans. *Biology*, *9*(9), 283.
https://doi.org/10.3390/biology9090283
Patterson, K. R. (1996). Modelling the impact of disease-induced mortality in an exploited
population: the outbreak of the fungal parasite (Ichthyophonus)(hoferi) in the North Sea
herring (Clupea harengus). *Canadian Journal of Fisheries and Aquatic Sciences*, *53*(12),
2870-2887. https://doi.org/10.1139/f96-234
Petton, B., Boudry, P., Alunno-Bruscia, M., & Pernet, F. (2015). Factors influencing disease-
induced mortality of Pacific oysters Crassostrea gigas. *Aquaculture Environment*
Interactions, *6*(3), 205-222. https://doi.org/10.3354/aei00125
Pinzón C, J. H., Beach-Letendre, J., Weil, E., & Mydlarz, L. D. (2014). Relationship between
Phylogeny and Immunity Suggests Older Caribbean Coral Lineages Are More Resistant
to Disease. *PLOS ONE*, *9*(8), e104787. https://doi.org/10.1371/journal.pone.0104787
Pinzón, C. J., Dornberger, L., Beach-Letendre, J., Weil, E., & Mydlarz, L. D. (2014). The link
between immunity and life history traits in scleractinian corals. *PeerJ*, *2*, e628.
https://doi.org/10.7717/peerj.628
Pirotta, E., Thomas, L., Costa, D. P., Hall, A. J., Harris, C. M., Harwood, J., Kraus, S. D., Miller,
P. J. O., Moore, M. J., Photopoulou, T., Rolland, R. M., Schwacke, L., Simmons, S. E.,
Southall, B. L., & Tyack, P. L. (2022). Understanding the combined effects of multiple
stressors: A new perspective on a longstanding challenge. *Sci Total Environ*, *821*,
153322. https://doi.org/10.1016/j.scitotenv.2022.153322
Precht, W. F., Gintert, B. E., Robbart, M. L., Fura, R., & van Woesik, R. (2016). Unprecedented
Disease-Related Coral Mortality in Southeastern Florida. *Scientific Reports*, *6*(1), 31374.
https://doi.org/10.1038/srep31374
Richardson, L. L., Goldberg, W. M., Carlton, R. G., & Halas, J. C. (1998). Coral disease
outbreak in the Florida Keys: plague type II. *Revista de Biología tropical*, *46*(S5), 187-
198.
Rivetti, I., Fraschetti, S., Lionello, P., Zambianchi, E., & Boero, F. (2014). Global warming and
mass mortalities of benthic invertebrates in the Mediterranean Sea. *PLOS ONE*, *9*(12),
e115655. https://doi.org/10.1371/journal.pone.0115655
Rodrigues, L. J., & Grottoli, A. G. (2007). Energy reserves and metabolism as indicators of coral
recovery from bleaching. *Limnology and Oceanography*, *52*(5), 1874-1882.
https://doi.org/10.4319/lo.2007.52.5.1874
Rosenzweig, C., Karoly, D., Vicarelli, M., Neofotis, P., Wu, Q., Casassa, G., Menzel, A., Root,
T. L., Estrella, N., Seguin, B., Tryjanowski, P., Liu, C., Rawlins, S., & Imeson, A.
(2008). Attributing physical and biological impacts to anthropogenic climate change.
Nature, *453*(7193), 353-357. https://doi.org/10.1038/nature06937
Sakai, K., Singh, T., & Iguchi, A. (2019). Bleaching and post-bleaching mortality of Acropora
corals on a heat-susceptible reef in 2016. *PeerJ*, *7*, e8138.
https://doi.org/10.7717/peerj.8138
Sanderson, C. E., & Alexander, K. A. (2020). Unchartered waters: Climate change likely to
intensify infectious disease outbreaks causing mass mortality events in marine mammals.
Glob Chang Biol, *26*(8), 4284-4301. https://doi.org/10.1111/gcb.15163

Scavia, D., Field, J. C., Boesch, D. F., Buddemeier, R. W., Burkett, V., Cayan, D. R., Fogarty, M., Harwell, M. A., Howarth, R. W., Mason, C., Reed, D. J., Royer, T. C., Sallenger, A. H., & Titus, J. G. (2002). Climate change impacts on U.S. Coastal and Marine Ecosystems. *Estuaries*, *25*(2), 149-164. https://doi.org/10.1007/BF02691304

Short, F. T., Muehlstein, L. K., & Porter, D. (1987). EELGRASS WASTING DISEASE: CAUSE AND RECURRENCE OF A MARINE EPIDEMIC. *The Biological Bulletin*, *173*(3), 557-562. https://doi.org/10.2307/1541701

Smale, D. A. (2020). Impacts of ocean warming on kelp forest ecosystems. *New Phytol*, *225*(4), 1447-1454. https://doi.org/10.1111/nph.16107

Smith, K. E., Burrows, M. T., Hobday, A. J., King, N. G., Moore, P. J., Sen Gupta, A., Thomsen, M. S., Wernberg, T., & Smale, D. A. (2023). Biological Impacts of Marine Heatwaves. *Annual Review of Marine Science*, *15*(1), 119-145. https://doi.org/10.1146/annurev-marine-032122-121437

Smith, S. V., & Buddemeier, R. W. (1992). Global Change and Coral Reef Ecosystems. *Annual Review of Ecology and Systematics*, *23*, 89-118. http://www.jstor.org/stable/2097283

Subhan, B., Arafat, D., Rahmawati, F., Dasmasela, Y. H., Royhan, Q. M., Madduppa, H., Santoso, P., & Prabowo, B. (2020). Coral disease at Mansuar Island, Raja Ampat, Indonesia. *IOP Conference Series: Earth and Environmental Science*, *429*(1). https://doi.org/10.1088/1755-1315/429/1/012027

Thornhill, D. J., Xiang, Y., Pettay, D. T., Zhong, M., & Santos, S. R. (2013). Population genetic data of a model symbiotic cnidarian system reveal remarkable symbiotic specificity and vectored introductions across ocean basins. *Mol Ecol*, *22*(17), 4499-4515. https://doi.org/10.1111/mec.12416

Tortorelli, G., Belderok, R., Davy, S. K., McFadden, G. I., & van Oppen, M. J. H. (2020). Host Genotypic Effect on Algal Symbiosis Establishment in the Coral Model, the Anemone Exaiptasia diaphana, From the Great Barrier Reef. *Frontiers in Marine Science*, *6*. https://www.frontiersin.org/articles/10.3389/fmars.2019.00833

Traylor-Knowles, N., Connelly, M. T., Young, B. D., Eaton, K., Muller, E. M., Paul, V. J., Ushijima, B., DeMerlis, A., Drown, M. K., Goncalves, A., Kron, N., Snyder, G. A., Martin, C., & Rodriguez, K. (2021). Gene Expression Response to Stony Coral Tissue Loss Disease Transmission in M. cavernosa and O. faveolata From Florida [Original Research]. *Frontiers in Marine Science*, *8*. https://doi.org/10.3389/fmars.2021.681563

Valadez-Ingersoll, M., Aguirre Carrion, P. J., Bodnar, C. A., Desai, N. A., Gilmore, T. D., & Davies, S. W. (2023). Nutrient deprivation differentially affects gene expression, immunity, and pathogen susceptibility across symbiotic states in a model cnidarian. *bioRxiv*, 2023.2007. 2030.551141.

Vezzulli, L., Colwell, R. R., & Pruzzo, C. (2013). Ocean Warming and Spread of Pathogenic Vibrios in the Aquatic Environment. *Microbial Ecology*, *65*(4), 817-825. https://doi.org/10.1007/s00248-012-0163-2

Vollmer, S. V., & Kline, D. I. (2008). Natural Disease Resistance in Threatened Staghorn Corals. *PLOS ONE*, *3*(11), e3718. https://doi.org/10.1371/journal.pone.0003718

Wall, C. B., Ricci, C. A., Foulds, G. E., Mydlarz, L. D., Gates, R. D., & Putnam, H. M. (2018). The effects of environmental history and thermal stress on coral physiology and immunity. *Marine Biology*, *165*(3), 56. https://doi.org/10.1007/s00227-018-3317-z

Wall, C. B., Ricci, C. A., Wen, A. D., Ledbetter, B. E., Klinger, D. E., Mydlarz, L. D., Gates, R. D., & Putnam, H. M. (2021). Shifting baselines: Physiological legacies contribute to the

response of reef corals to frequent heatwaves. *Functional Ecology*, *35*(6), 1366-1378.
https://doi.org/10.1111/1365-2435.13795

Wild, S., Krutzen, M., Rankin, R. W., Hoppitt, W. J. E., Gerber, L., & Allen, S. J. (2019). Long-
term decline in survival and reproduction of dolphins following a marine heatwave.
Current biology: CB, *29*(7), R239-R240. https://doi.org/10.1016/j.cub.2019.02.047

Winter, A., Appeldoorn, R., Bruckner, A., Williams Jr, E., & Goenaga, C. (1998). Sea surface
temperatures and coral reef bleaching off La Parguera, Puerto Rico (northeastern
Caribbean Sea). *Coral Reefs*, *17*, 377-382.

Wright, R. M., Kenkel, C. D., Dunn, C. E., Shilling, E. N., Bay, L. K., & Matz, M. V. (2017).
Intraspecific differences in molecular stress responses and coral pathobiome contribute to
mortality under bacterial challenge in Acropora millepora | Scientific Reports. *Scientific*
Reports, *7*(2609). https://www.nature.com/articles/s41598-017-02685-1

Yakovleva, I., Bhagooli, R., Takemura, A., & Hidaka, M. (2005). Differential susceptibility to
oxidative stress of two scleractinian corals: Antioxidant functioning of mycosporine-
glycine. *Comparative biochemistry and physiology. Part B, Biochemistry & molecular*
biology, *139*, 721-730. https://doi.org/10.1016/j.cbpc.2004.08.016

Zaragoza, W. J., Krediet, C. J., Meyer, J. L., Canas, G., Ritchie, K. B., & Teplitski, M. (2014).
Outcomes of infections of sea anemone Aiptasia pallida with Vibrio spp. pathogenic to
corals. *Microbial Ecology*, *68*(2), 388-396. https://doi.org/10.1007/s00248-014-0397-2

Tables and Figures

Table 1: Best-fit linear models for each immune parameter when including temperature treatment, clonal line, immune challenge, and lipids as predictors. † indicate models that did not undergo model dredging and averaging. Asterisks (*) represent significant p-value ($\alpha =$ 0.05).

Catalase					
Predictors	**Estimates**	**SE**	**SEadj**	**z**	**p value**
(Intercept)	336	29.3	29.7	11.3	**< 0.001 ***
Treatment	-126	31.8	32.2	3.90	**< 0.001 ***
Clonal Line	62.8	29.0	29.4	2.13	**0.0329 ***
Immune Challenge	43.6	37.8	38.1	1.14	0.253
Treatment: Immune	-14.9	36.5	36.7	0.404	0.686
Clonal: Immune	7.33	26.8	27.0	0.271	0.786
Antibacterial Activity					
Predictors	**Estimates**	**SE**	**SEadj**	**z**	**p value**
(Intercept)	1.86	0.112	0.114	16.3	**< 0.001 ***
Treatment	0.327	0.157	0.160	2.04	**0.0411 ***
Immune Challenge	0.00205	0.103	0.104	0.020	0.984
Lipids	2.32	1.21	1.23	1.88	0.0605
Treatment: Immune	-0.0487	0.0884	0.0891	0.547	0.584
Treatment: Lipids	-4.66	1.63	1.65	2.82	**< 0.01***
Immune: Lipids	-0.369	1.03	1.04	0.355	0.723
Total Phenoloxidase					
Predictors	**Estimates**	**SE**	**SEadj**	**z**	**p value**
(Intercept)	0.0233	0.00136	0.00138	16.9	**< 0.001 ***
Treatment	-3.04E-02	1.10E-02	1.12E-02	2.71	**< 0.01 ****
Clonal Line	-0.00156	0.00272	0.00275	0.569	0.569
Immune Challenge	0.00478	0.00176	0.00178	2.68	**< 0.01 ****
Treatment: Clonal	-0.00356	0.00413	0.00415	0.857	0.391
Treatment: Immune	0.000337	0.00137	0.00138	0.244	0.807
Clonal: Immune	-0.00133	0.00268	0.00270	.494	0.621

Melanin†				
Predictors	**Estimates**	**SE**	**t**	**p value**
(Intercept)	2.31	1.36	1.70	0.089
Treatment	-2.45	0.675	-3.62	**< 0.001****
Clonal Line	1.12	0.635	1.77	0.077
Immune Challenge	0.494	0.624	0.791	0.429
Lipids	-11.4	12.5	-0.916	0.360
0s removed Melanin†				
Predictors	**Estimates**	**SE**	**t**	**p value**
(Intercept)	-1.49E-04	1.01E-04	-1.47	0.149
Treatment	1.62E-05	5.19E-05	0.311	0.757
Clonal Line	-1.24E-06	4.96E-05	-0.025	0.980
Immune Challenge	5.35E-05	4.83E-05	1.11	0.275
Lipids	4.02E-03	9.34E-04	4.31	**<0.001 ***

Figure 1: (a-b) Kaplan-Meier survivorship curves for pathogen exposed anemones split based on comparisons across **(a)** temperature treatment and **(b)** genotype (clonal line). **(c)** Cox proportional hazards models of pathogen exposed anemones compared across temperature treatment and genotype. Vertical dotted line at 1 indicates the reference to which the second group is compared. The distance from 1 indicates the probability of mortality (i.e. heat stress is about twice as likely to die from immune challenge as heat). Both treatment ($p < 0.01$) and genotype ($p < 0.001$) had significant impacts on survival. Hazards ratios are displayed for both predictors with 95% confidence intervals.

Figure 2: Box plots displaying symbiont density (**a-b**), carbohydrate concentration (**c-d**), and lipid concentration (**e-f**0 across temperature treatments and clonal lines. (**a**) Symbiont density was non-significantly different between temperature treatments (Welch's two sample t-test; $p = 0.058$); **b**) However, symbiont density was significantly higher in H2 anemones compared to VWB9 (Welch's two sample t-test; $p = 0.046$). Carbohydrate concentrations, on the other hand, were (**c**) significantly lower in heat-stressed anemones compared to control (Wilcoxon Rank Sum; $p < 0.01$) and (**d**) higher in the VWB9 clonal line compared to H2 (Wilcoxon Rank Sum; $p < 0.01$). Lipid concentrations did not differ across either (**e**) temperature treatments (Welch's two sample t-test; $p = 0.348$) or (**f**) clonal lines (Welch's two sample t-test; $p = 0.664$).

Figure 3. RDA plot showing significant predictors for explaining variation in holistic immunity. Data points represent individual samples. Ordination was constructed from response variables which in this case were immune parameters. Color indicates temperature treatment, shape indicates clonal line, and open shape/filled shape indicates immune challenge. Black boxes represent predictor variables and white boxes represent response variables. Arrows represent strength and directionality of each variable in multi-variate space.

Figure 4: Differences in individual immune parameters as a result of temperature treatment. Prior heat treatment significantly affected **(a)** catalase activity (GLM, $p < 0.001$) and **(b)** melanin concentration (GLM, $p < 0.001$; 0s removed GLM, $p < 0.001$).

Figure 5: Best-fit linear models of the association between lipids and **(a)** antibacterial activity and **(b)** melanin concentration. **(a)** Heat treatment significantly affected associations between standardized lipid concentration and antibacterial activity (GLM, $p < 0.01$). **b)** However, melanin concentration was consistently associated with lipid concentration regardless of treatment group (GLM, $p < 0.001$; 0s removed GLM, $p < 0.001$). Points represent individual samples color coded by treatment group. Red trendline represents linear model of temperature- dependent association. Grey trendlines represent linear models of associations independent of temperature.