Unlocking the Mycorrhizal Nitrogen Pathway Puzzle : Metabolic Modelling and multi-omics unveil Pyrimidines' Role in Maize Nutrition via Arbuscular Mycorrhizal Fungi Amidst Nitrogen Scarcity

Authors :

Bérengère Decouard[1], Niaz Bahar Chowdhury[2], Aurélien Saou[3], Martine Rigault[1], Isabelle Quilleré[1], Thomas Sapir[1], Anne Marmagne[1], Christine Paysant le Roux[4], Alexandra Launay-Avon[4], Florence Guerard[4], Caroline Mauve[4], Bertrand Gakière[4], Céline Lévy-Leduc[3], Pierre Barbillon[3], Pierre-Emmanuel Courty[5], Daniel Wipf[5], Bertrand Hirel[1], Rajib Saha [2]*, Alia Dellagi[1]*

1 : Université Paris-Saclay, INRAE, AgroParisTech, Institut Jean-Pierre Bourgin (IJPB), 78000, Versailles, France.

2 : Chemical and Biomolecular Engineering, University of Nebraska-Lincoln, Lincoln, NE, United States of America

3 : Université Paris-Saclay, AgroParisTech, INRAE, UMR MIA Paris-Saclay, 91120 Palaiseau, France

4 : Université Paris-Saclay, CNRS, INRAE, Université Evry, Institute of Plant Sciences Paris-Saclay (IPS2), 91190, Gif sur Yvette, France. Université Paris Cité, CNRS, INRAE, Institute of Plant Sciences Paris-Saclay (IPS2), 91190, Gif sur Yvette, France

5 : Agroécologie, AgroSup Dijon, Centre National de la Recherche Scientifique (CNRS), Université de Bourgogne, Institut National de Recherche pour l'Agriculture, l'Alimentation, et l'Environnement (INRAE), Université Bourgogne Franche-Comté, Dijon, France.

* Corresponding authors

Alia Dellagi

Address: Université Paris-Saclay, INRAE, AgroParisTech, Institut Jean-Pierre Bourgin (IJPB),

78000, Versailles, France.

Phone: +33 1 30 83 32 28

E-mail: alia.dellagi@agroparistech.fr

Rajib Saha

Address: Chemical and Biomolecular Engineering, University of Nebraska-Lincoln, Lincoln,

NE, USA

Phone: +1 402 472 7531

E-mail: rsaha2@unl.edu

Key words: N use efficiency; Nitrogen uptake; arbuscular mycorrhizal fungi, Maize, metabolic

modelling.

Abstract 332

Maize is currently the most productive cereal crop in the world (www.faostat.org). Maize can

form a symbiotic relationship with the Arbuscular Mycorrhizal Fungus (AMF), *Rhizophagus*

irregularis. In this relationship, the fungus provides the plant with additional water and

mineral nutrients, while the plant supplies carbon compounds to the fungus. Little is known

about the N metabolism disruption during symbiosis in both partners. To address this issue,

two genetically distant maize lines were studied in terms of physiological and molecular

responses to AMF inoculation by dual RNA-seq, metabolomics and phenotyping.

Interestingly, the beneficial effects of the AMF were observed mainly under conditions of

limited N fertilization. Under such conditions, the AMF helped maintain plant biomass

production. The availability of nitrogen was found to be a crucial factor influencing all the

traits studied showing that the level of N supply plays a pivotal role in determining how maize

plants interact with the AMF. Despite the two maize lines showing different transcriptomic

and metabolomic responses to *R. irregularis*, their agro-physiological traits remained similar.

Both the plant and fungal transcriptomes were more significantly influenced by the level of N

nutrition rather than the specific maize genotype. This suggests that N availability has a more

profound impact on gene expression in both organisms than the genetic makeup of the

maize plant. To understand the metabolic implications of this symbiotic relationship, we

integrated transcriptomic data into our recently built multi-organ Genome-scale metabolic

model (GSM) called iZMA6517. Remarkably, this modelling approach was supported by

metabolomics profiling, in particular increased leaf pyrimidine levels in response to AMF

inoculation under limiting N supply. Consistently, fungal genes involved in pyrimidine de novo

synthesis and salvage were found to be expressed in symbiotic roots. Our work highlights

nucleotide and ureides metabolism as previously unrecognized factors contributing to the

symbiotic N nutrition facilitated by *R. irregularis*, thereby enhancing maize growth. This study

demonstrates the effectiveness of integrating multi-omics approaches with mathematical

modelling to uncover novel metabolic mechanisms associated with AM symbiosis, without a

priori.

Introduction 908 words

Nitrogen (N) is the fourth most abundant component of plant biomass, which explains its

pivotal role in agriculture [1,2]. Indeed, large inputs of N fertilizers are necessary to ensure

optimal crop productivity [1,3]. However, only 30 to 50% of the N available in the soil is taken

up by most crops, resulting in important N losses which can cause damage to the environment such as water pollution and increased greenhouse effect [4,5]. Ensuring crop productivity while minimizing N supply can be monitored by measuring nitrogen use efficiency (NUE). A better understanding of how to enhance NUE is a major challenge in the context of agroecology as it represents the capacity of the plant to produce grains or vegetative biomass per unit of available N [3,6,7]. Direct uptake of minerals including N and P from the soil by plants is ensured by transporters known to drive the so called "direct transport". Interestingly, to fulfil their N-associated metabolic needs, plants can establish symbiosis with beneficial soil microbes which help to improve their N nutrition by different mechanisms, some of which are well characterized. Exploiting the capacity of a crop to benefit from beneficial microbes in terms of N nutrition in order to maintain yields while reducing chemically synthesized N input is challenging in the context of a more sustainable agriculture [8–13]. Nitrogen can be transferred from the soil to the roots when the plant is colonized by Arbuscular Mycorrhizal Fungi (AMFs), a symbiotic process occurring in 80% of land plants including cereals [8,12–14]. The early steps of AMF symbiosis involve signal exchange between the host roots of the host plant and the fungus. This signalling process leads to establishment of a physical contact between the partners during which the fungus develops structures like a hyphopodium on the root epidermal cell followed by its growth inside these cells up to the cortical cells where it differentiates into highly branched structures called "arbuscules". Around these arbuscules, the cortical cells undergo strong reorganization leading to the formation of a structure called "periarbuscular membrane" (PAM). Dual nutritional exchange occurs at the interface of this membrane where the transfer of carbon containing molecules derived from photosynthesis such as hexoses and lipids feed the fungus, which in turn provides minerals such as N and P [8,11,15–17]. The extra-radical mycelium, which represents an extension of the arbuscular structures, allows the fungus to explore the surrounding rhizosphere and to take up minerals further translocated to the periarbuscular interface to be finally delivered to the host plant.

It is well established that phosphorus (P) deficiency is the main driver for the establishment

of mycorrhizal symbiotic associations [18–22] . In contrast, the benefit of a mycorrhizal symbiosis

in terms of N uptake and utilization by the host plant remains to be fully characterized [11–

14].

Maize is currently the most productive cereal crop in the world (www.faostat.org) with a

global yield exceeding that of rice [23]. Different maize genotypes are able to develop a

symbiotic association with AMF [24,25]. Several studies have shown that AMF can supply

substantial amount of N to maize [26,27]. However, physiological and molecular characterization

of maize's interaction with AMF under different N supply levels has never been undertaken.

A Genome-Scale Model (GSM) encompasses the majority of known metabolic reactions

within a biological system. It employs techniques such as flux balance analysis (FBA) [28], flux

variability analysis (FVA) [29], and parsimonious FBA [30], to predict the flux of reactions

accurately. Genome-scale metabolic modelling (GSM) is a computational tool to model plant

metabolism and applied to plant system can highlight potential key metabolic pathways in

plants in response to environmental cues [31]. Initial efforts on maize metabolic models were

limited to its leaf [32] and root [33] GSMs, which showed agreement with existing literature

regarding phenotypical predictions. Most recently, a multi-organ GSM of maize line B73,

iZMA6517 [34], is proposed which includes full scale GSMs of root, stalk, kernel, and leaf;

connected via phloem vascular tissue. iZMA6517 is the largest maize multi-organ GSM of

maize till date and captured accurate plant phenotype under heat and cold stress conditions.

In the present study, we have investigated in an integrated manner the impact of N nutrition

on the interaction between two maize lines and the AMF *Rhizophagus irregularis* both at the

physiological and molecular levels. Using phenotyping, dual RNA-seq transcriptomics and

metabolomics approaches, we generated a set of paired data (same plants used for all

analysis). We have characterized such interactions using two maize lines. Both maize lines

displayed improvement of agronomic performance in response to inoculation with *R.*

irregularis under limited N supply. Thereby, apart from experimental works, we used

*i*ZMA6517 to predict plant agronomic performance and organ-specific metabolic bottlenecks

accurately to explore the symbiotic relationship with *R. irregularis*. *i*ZMA6517 GSM identified

pyrimidine biosynthetic pathway as a key physiological process involved in the efficiency of N

utilization in the symbiotic association when N is limiting. Interestingly, this prediction was

supported by monitoring pyrimidine levels in leaves of symbiotic plants. Thus, modelling

approach highlighted nucleotide and ureides metabolism as previously unrecognized factors

contributing to the symbiotic N nutrition facilitated by *R. irregularis*, thereby enhancing maize

growth. This research highlights how the combination of multi-omics strategies and

mathematical modelling can reveal new metabolic processes associated with AM symbiosis

without preconceived notions. Our work highlights nucleotide and ureides metabolism as

previously unrecognized factors contributing to the symbiotic N nutrition facilitated by *R.*

irregularis, thereby enhancing maize growth.

Material and methods 2518 words

Plant material, growth conditions and inoculation

The maize inbred lines used in this study B73 and Lo3 (Supplementary table 1). The reference maize genotype B73, for which the genome was fully sequenced, [35]. The fungal strain used is the highest quality of purity strain *Rhizophagus irregularis* DAOM197198 (Agronutrition, Toulouse, France). Inoculum was prepared in distilled water and the spore suspension was added next to the germinated seed or plantlet with a pipette.

In all experiments, seeds were surface-sterilized as follows. Seeds were first incubated in ethanol for 5 minutes at 28°C followed by rinsing with distilled water. Then they were incubated for 45 minutes in a 15% commercial bleach solution with 0.01% Triton x100 then rinsed in distilled water. They were placed on wet sterile Whatman paper in Petri dishes closed with parafilm and incubated in the dark for 48-72h at 20°C. Germinated seeds were selected and then sown in pots of 4 litres containing a mix of sterilized unfertilized peat (40% Baltic blond peat 0-10 mm, 60% Brown peat 0-10 mm, 75 g oligo elements, pH $CaCl_2$ 5.3 – 5.8, Stender, France) and dried clay beads (reference: 17831050 BILLE ARGILE LIA 8/16, Soufflet, France) 1:1/v:v. For experimental design and the different steps of growing and harvesting of the experiment in 2020 and 2021 see Supplementary Figure 1 and 2 respectively. For meteorological conditions within the greenhouse during 2020 see Supplementary Figure 3. For each condition, 9-10 plants were used. Three inoculations with spores of *R. irregularis* were performed as follow: The first inoculation with 500 spores/plant at sowing, second inoculation 1 week after sowing with 200 spores/plant and a third inoculation 2 weeks after sowing with 200 spores/plant. Control plants were treated with distilled water. Plants were grown under three different nitrogen regimes: low N (LN, 1mM NO_3^-), medium N (MN, 5 mM NO_3^-) and high N (HN, 10 mM NO_3^-). The specific composition of these regimes is provided in Supplementary Table 2. For each N regime, plants were irrigated for the first three weeks with the solution corresponding to the chosen N regime with low P (named P-) containing 10 µM phosphate in order to trigger AMF colonization. Then, at

dates indicated in Supplementary Figure 1 and 2, irrigation was changed to the same N level

but 300 µM P (named P+, Supplementary Table 2). Increasing P level in the irrigation

solution was necessary to allow plants reach the maturity stage. To determine ^{15}N uptake,

^{15}N labelling was performed at indicated dates (Supplementary Figure 1). 100 mL of 45 mM

KNO_3 solution enriched at 2% ^{15}N was applied to each plant in order to provide 1,25mg of

^{15}N.

Harvesting procedures

Plants grown in 2020 were used for the phenotyping, metabolomics, RNA-seq and RT-qPCR

experiments. Plants grown in 2021 were used to monitor fungal colonization and RT-qPCR

experiments. Depending on the experiment and the biological question asked, one, two or

three developmental stages were chosen for tissue harvesting. The first one is the vegetative

stage (VS) corresponding to the 7- to 8-leaf stage [36] is an indicator of a highly metabolically

and photosynthetically active leaf tissue. The second stage corresponds to 15 days after the

tasselling (VT+15) and plant is in the early phase of silking (data not shown). This stage was

chosen because it allows root harvesting at a reproductive stage without compromising

kernel production. The third stage is the maturity stage (MS) and corresponds to plants

displaying mature ears. Dates of harvesting for VS, VT+15 stage and MS differed depending

on the condition and the year of experiment (Supplementary Figure 1 and 2). For leaf

tissues, harvesting consisted in cutting a piece of 1 cm x 10 cm of leaf mesophyll along the

midrib from the 6th fully emerged leaf for the VS. For VT+15 stage and MS, a piece of 1 cm x

10 cm leaf surface from the leaf below the ear was harvested. Roots were harvested at VS,

VT+15 stage or MS depending on the type of experiment. For each treatment/condition, roots

were collected from 5 plants. Thin secondary roots were harvested from the surrounding root

system, close to the pot borders. They were washed with distilled water, then frozen in liquid

nitrogen or stored in distilled water. Following root harvesting, the plants grown in 2020 grew

until maturity.

Quantification of yield, biomass, chlorophyll, C, N, ^{15}N and P

Plant organs were harvested then oven dried at 80°C for 3 to 5 days then ground to fine

powder. Grain, biomass of the aerial parts including stalks, leaves, and cobs were weighed

after drying. The upper regions of the root system was harvested. The percentage of C and

N were quantified using the FLASH 2000 Organic Elemental Analyzer (Thermo Fisher

Scientific Villebon, France) and the ^{15}N enrichment using the Delta V Advantage isotope ratio

mass spectrometer (Thermo Fisher Scientific, France) as described in [37]. Supplementary

Table 3 indicates the calculations of different traits with C and N content. A% (Atom percent)

represents the ^{15}N enrichment in the sample [^{15}N/(total N)]. Since the natural ^{15}N abundance

in N unlabelled samples was 0.3663 (A% control), specific enrichments due to the ^{15}N uptake

were calculated as E% = (A% - 0.3363).The P concentration was measured using the

molybdate blue method on a Shimadzu UV-160 spectrophotometer (Shimadzu Biotech) after

acid digestion [38].

210   Quantification of root colonization level by AMF structures

Thin secondary roots were harvested as indicated above and carefully washed in distilled

water. They were incubated in 10% KOH overnight at room temperature. They were then

rinsed in distilled water and incubated in 5% black ink (Sheaffer Skrip Bottled Ink, France)

solution for 10 min at 70°C. Washing was performed by incubating roots in 8% acetic acid for

25 min. For the quantification of % of AM colonization (total, arbuscules, vesicles) 50 pieces

of 1 cm length each were mounted on a slide for observation of AM structures under light

microscope using the slide method [39]. Quantifications were performed on roots collected from

5 plants for each condition.

219   RNA extraction for RT qPCR and RNAseq

220   RNA extraction was performed with TRIzol (Thermo Fisher Scientific, USA) protocol. Fresh

221   tissue of leaves or roots (100 mg) were frozen in liquid nitrogen then ground to fine powder

222   that was homogenized in 1 mL of TRIzol. RNA-extraction was performed as described in [34]

223   and was used for RT-qPCR and RNA-seq.

cDNA synthesis and RT-PCR

It was performed as indicated in Decouard et al. [37]. Primer sequences are described in Supplementary table 4.

RNA-seq and Bioinformatic Analysis

RNA-seq libraries were constructed by the POPS platform (IPS2) using the Illumina Stranded mRNA Prep kit (Illumina®, California, U.S.A.) according to the supplier's instructions. Libraries were sequenced by the Genoscope (Évry-Courcouronnes, France) on a Novaseq 6000 instrument (Illumina) with 150 base paired-end sequences. RNA-seq raw data cleaning was made as described in [34]. Statistical RNA-seq analyses are described in Supplementary File 1. Filtered reads were then mapped and counted using STAR (v2.7.3a, [40]) with the following parameters --alignIntronMin 5 --alignIntronMax 60000 --outSAMprimaryFlag AllBestScore --outFilterMultimapScoreRange 0 --outFilterMultimapNmax 20 on a combined reference genome made by concatenating the files of the *Zea mays* B73 (V4.49) and *Rhizophagus irregularis* (V2) reference genomes and the associated GTF annotation files (also combined). Between 75.3 and 96.1 % of the reads were uniquely mapped (median = 91.9%). Between 68 and 88 % of the reads were associated to annotated genes (median = 85.3%).

Metabolomic analyses

All extraction steps were performed in 2 mL Safelock Eppendorf tubes. The ground dried samples (5 mg DW) were resuspended in 1 mL of frozen (-20°C) Water:Acetonitrile:Isopropanol (2:3:3) containing Ribitol at 4 mg/L and extracted for 10 min at 4°C with shaking at 1500 rpm in an Eppendorf Thermomixer. Insoluble material was removed by centrifugation at 13500 rpm for 10 min. 700 µL were collected and 70 µL of myristic acid d27 at 0.3 g/L was added as an internal standard for retention time locking. After centrifugation, aliquots of each extract (100 µL) were dried for 4 h at 35 °C in a Speed-Vac and stored at -80°C. Two Blank tubes underwent the same steps as the samples.

Sample derivatization and analysis : All steps of GC-MS analyses were done as described in

Fiehn et al.[41,42]. Agilent literature PUB: 5989-8310EN (Agilent G1676AA Agilent Fiehn

GC/MS Metabolomics RTL Library User Guide. June 2008. Agilent P/N: G1676-90000,

Agilent Fiehn GC/MS Metabolomics RTL Library Product Note. May 2008). Samples were

taken out of -80°C, warmed 15 min before opening and speed-vac dried again for 1 hour at

35 °C before adding 10 µl of 20 mg/mL methoxyamine in pyridine to the samples and the

reaction was performed for 90 min at 30°C under continuous shaking in an Eppendorf

thermomixer. 90 µL of N-methyl-N-trimethylsilyl-trifluoroacetamide (MSTFA) (REGIS

TECHNOLOGIES 1-270590-200 -10x1g) were then added and the reaction continued for 30

min at 37°C. After cooling, 90 µL were transferred to an Agilent vial for injection.

4 hours after derivatization 1 µL of sample was injected in splitless mode on an Agilent

7890B gas chromatograph coupled to an Agilent 5977A mass spectrometer. The column was

an ZORBAX DB5-MS from Agilent (30 m with 10 m integraguard column - ref 122-5532G).

An injection in split mode with a ratio of 1:30 was systematically performed for saturated

quantification compounds. Oven temperature ramp was 60 °C for 1 min then 10 °C/min to

325 °C for 10 min. Helium constant flow was 1.1 mL/min. Temperatures were as follows:

injector: 250°C, transfer line: 290°C, source: 230 °C and quadripole 150 °C. The quadrupole

mass spectrometer was switched on after a 5.90 min solvent delay time, scanning from 50-

600 µ. Absolute retention times were locked to the internal standard d27-myristic acid using

the RTL system provided in Agilent's Masshunter software. Retention time locking reduces

run-to-run retention time variation. Samples were randomized. A fatty acid methyl esters mix

(C8, C9, C10, C12, C14, C16, C18, C20, C22, C24, C26, C28, C30) was injected in the

middle of the queue for external RI calibration.

Data processing and quantification: Raw Agilent datafiles were analyzed with AMDIS

http://chemdata.nist.gov/mass-spc/amdis/. The Agilent Fiehn GC/MS Metabolomics RTL

Library (version June 2008) was employed for metabolite identifications. This library is the

most comprehensive library of metabolite GC/MS spectra that is commercially available. It

contains searchable GC/MS EI spectra and retention time indexes from approximately 700

common metabolites. Peak areas determined with the Masshunter Quantitative Analysis

(Agilent) in splitless and split 30 modes. Because automated peak integration was

occasionally erroneous, integration was verified manually for each compound in all analyses.

Resulting areas were compiled in one single Excel file for comparison. Peak areas were

normalized to Ribitol and Dry Weight. Metabolite contents are expressed in arbitrary units

(semi-quantitative determination).

Amino acid quantification by OPA-HPLC

Extraction procedure was identical that one described for metabolomics. Here 200µL of the

extraction solution was taken and dried in a Speed-Vac evaporator for 2 h at 30°C before

adding 1,3 mL of H_2O milliQ. The o-phtaldialdéhyde (OPA) reagent was made 48 h before

first use by dissolving OPA at 12mg/mL in 200 µL of methanol and adding 1.8 mL of sodium

borate 0.5M (pH 9.5) and 40 µL of 2-mercaptoethanol. The reagent was filtered into an

autosampler vial and used for up to 3 days. Precolumn derivatization was performed in the

injection loop by automated mixing of 10 µL sample and 10 µL OPA reagent, followed by a

delay of 2 min prior to injection. The HPLC (Alliance Waters 2695, Waters Corporation,

Massachusetts, USA) fitted with a fluo detector (Multi λ Fluorescence Detector 2475) was

used. Compounds were measured at λ excitation of 340 nm and λ emission of 455 nm. The

chromatographic separation was performed with Symmetry C18 column (3.5µm, 4.6*150

mm) by gradient elution at 40 °C using buffer A (20% methanol, 80% sodium acetate, 1%

tetrahydrofuran, pH 5.9) and buffer B (80% methanol, 20% sodium acetate, pH 5.9). Buffer

flow rate was 0.8 mL/min throughout and total run time per injection was 42 min. The

chromatography data were analysed by Empower software. Peak identity was confirmed by

co-elution with authentic standards. Concentration was calculated with calibration curve

using peak area of compound of interest.

Nucleotide quantification LC–TOF.

Nucleotide quantification as performed on frozen leaves of the same maize plants used for

metabolome and RNAseq. Extraction and quantification were performed as described in [43].

Statistical analyses

All statistical analyses were performed with RStudio 2023.06.0+421 "Mountain Hydrangea"

under the free software environment R version 4.3.1 (Beagle Scouts). All Anova were

implemented with the "aov" R package. All heatmaps were generated with the estimated

marginal means that were calculated using the R package "emmeans". Heatmap for physio-

agronomical traits was generated with 'pheatmap' 1.0.12 package (Date 2019-01-04. Author

Raivo Kolde) by implementing the ward.D2 hierarchical clustering method [44]. Heatmap for

physio-agronomical traits was generated with the 'pheatmap' R package 1.0.12 Date 2019-

01-04 (Author Raivo Kolde). Heatmaps for metabolomics, including the interactive ones,

were generated with "heatmaply" R package [45].

Metabolomics data were analyzed with the R package MultiVarSel [46] which is the

implementation of the variable selection approach in the multivariate linear model proposed

in [47] theoretically analysed in [46] and applied in several contexts of omics data analyses such

as [48,49]. The list of R packages used in this study *sbm: Stochastic Blockmodels*. R package

version 0.4.5-9000, <https://grosssbm.github.io/sbm/>. Kolde R (2019). _pheatmap: Pretty

Heatmaps_. R package version 1.0.12, <https://CRAN.R-project.org/package=pheatmap>.

Lenth R (2023). _emmeans: Estimated Marginal Means, aka Least-Squares Means_. R

package version 1.8.6, <https://CRAN.R-project.org/package=emmeans>. Perrot-Dockès M,

Lévy-Leduc C, Chiquet J. R package MultiVarSel available from the Comprehensive R

Archive Network, 2017. <https://CRAN.R-project.org/package=MultiVarSel>.

Genome-Scale Metabolic Modelling (GSM) and Transcriptomics Data Integration

To investigate the metabolism of maize from systems level, we used *i*ZMA6517 as the

template model [34]. The model is specific to B73 genotype and consists of organ-specific

GSMs of root, stalk, leaf, and kernel. All these individual organ GSMs are connected through

a common phloem pool. The model has been described in our earlier work [34]. The model was

contextualized with HN Control, HN Rhizo, LN Control, and LN Rhizo using EXTREAM

algorithm [34]. The upper and lower limits of the reaction fluxes for the mentioned conditions

are presented in the Supplementary Table 5. The General Algebraic Modeling System

(GAMS) version 24.7.4 with IBM CPLEX solver was used to run FBA and FVA, E-Flux, and

the FSA algorithm on the model. Each of the algorithms was scripted in GAMS and then run

on a Linux-based high-performance cluster computing system at the University of Nebraska-

Lincoln. Instructions for running files in GAMS through NEOS server have been published [50].

All the necessary files for COBRApy users can also be accessed in the GitHub depository

(https://github.com/ssbio/iZMA6517).

Metabolic Bottleneck Analysis (MBA)

To determine the metabolic bottlenecks in each of HN Control, HN Rhizo, LN Control, and

LN Rhizo conditions, we used Metabolic Bottleneck Analysis (MBA) [34].. MBA expands the

flux space of each reaction separately and assesses its impact on the whole plant biomass

growth rate. Further details on MBA can be accessed in the original publications. MBA

pinpoints plant-wide metabolic bottlenecks, which were further analyzed for biological

insights.

Results 2547 words

The impact of *Rhizophagus irregularis* on maize growth and yield is strongly

dependent on the level of N nutrition

To investigate the diversity of the molecular and physiological mechanisms involved in maize

interaction with AMF in terms of N nutrition we used two genetically distant AMF-responsive

maize lines, B73 and Lo3. B73 is known be responsive to AMF, resulting in increased

biomass [25,51]. Lo3 genotype was selected due to its enhanced biomass response following

inoculation with *R. irregularis* in two rounds of selection (data not shown). In order to find at

which level of N fertilization the two lines benefit most from the fungal inoculation, these were

grown in a greenhouse under low N (LN), medium N (MN) or high N (HN) corresponding to

1 mM NO_3^-, 5 mM NO_3^- and 10 mM NO_3^- in the nutrient solution respectively. The impact of

R. irregularis inoculation on a panel of maize physio-agronomic traits was monitored

(Supplementary table 3). *R. irregularis* inoculation resulted in an increase in the shoot

biomass production (Aerial Parts, AP) and grain yield under LN in both lines (Figure 1A, 1B),

and under MN for B73. A negative impact of *R. irregularis* inoculation was observed on B73

grain yield when plants were grown under HN (Figure 1B). Interestingly, yield of mycorrhizal

plants grown under LN was similar to yield of non-inoculated plants fed with 5 times more

NO_3^- (MN). NUE increased in the AP only in LN conditions for both inoculated lines

compared to controls (Figure 1C). An increase in NUE for grain was observed under LN and

MN fertilization conditions only in B73 (Figure 1D). Fungal inoculation resulted in a significant

increase in the N content of aerial parts under MN for B73 and under HN for Lo3 indicating

that these lines are able to store higher levels of N in the vegetative parts upon association

with the AMF even under ample N (Supplementary Figure 4A, B, C). To determine whether N

uptake is modified during the symbiotic interaction, plants were labelled with a pulse of ^{15}N

provided at the vegetative stage. The total quantity of ^{15}N in aerial parts and grains measured

at maturity remained unchanged whatever the treatment (Supplementary Figure 5D and E),

except for the AP of Lo3 under HN. In inoculated plants, an increase in the total phosphorus

content was observed mainly under HN conditions (Supplementary Figure 4E, F, G).

Hierarchical clustering analysis (HCA) was performed using the physio-agronomic traits

listed in Supplementary Table 3 (Figure 1 E). The HCA shows that the N supply is the major

driver of all traits. All the LN plants were grouped in a cluster where values for the traits

related to AP and roots were higher. On the contrary, in cluster B, traits related mainly to

grains were higher. In cluster B, genotypes were distributed into two subgroups B1 and B2.

Interestingly, in cluster A, traits related to grains exhibited higher values in the inoculated

plants compared to the control indicating that *R. irregularis* allowed plants to behave more

similarly to group B plants, albeit still under N stress deficiency. These data show that *R.*

irregularis inoculation can mitigate the impact of N limitation on physio-agronomic traits.

Marker genes for N uptake and assimilation are differentially regulated in response to

AMF inoculation and N supply

The finding that several NUE traits, notably those related to NO_3^- uptake were modified by

AMF treatment prompted us to investigate further which molecular components of symbiosis

and N transport were involved at the root level. Roots were harvested at the VT+15 stage,

corresponding to 15 days after tasselling (see materials and methods) that was identified as

the most relevant stage to investigate maize-AMF interaction (Supplementary Figure 6). By

monitoring the level of *R. irregularis* constitutive transcript *RiLSU,* a constitutive *R. irregularis*

transcript*,* we found that high N input led to increased fungal biomass in roots (Figure 2A).

Then, the level of transcripts for the plant genes *ZmAMT3.1*, *ZmPHT6.1; ZmHA1* and

ZmNPF4.5 markers of the symbiotic association and representative of NH_4^+, phosphorus and

NO_3^- transport respectively was quantified by qRT-PCR [13,27,52,53].

The expression level of the four well-known symbiotic markers mentioned previously,

ZmPHT1.6, *ZmNPF4.5*, *ZmHA1*, a proton pump, and *ZmAMT3.1* is up-regulated in response

to the fungus in both genotypes and under all N supply conditions (Figure 2B). It is

interesting to notice that the higher the N nutrition level, the higher the expression of these

markers (Figure 2B). To know more about direct N transport genes in those conditions, we

investigated expression levels of two maker genes of N starvation stress *(ZmNRT2.1a, and*

ZmNRT2.2) encoding high affinity transporters, known to be upregulated under N deficiency

[2,54–56] (Figure 2B). As expected, the level of N supply had a drastic effect in both maize

genotypes on transcripts levels for the nitrate transporters *ZmNRT2.1a* and *ZmNRT2.2* which

were more highly expressed in LN plants compared to MN and HN plants. Interestingly, the

inoculation with *R. irregularis* on LN plants resulted in decreased levels of the *ZmNRT2.1a*

and *ZmNRT2.2* transcripts. These data indicate that the inoculation of N starved plants with

the fungus mimics an additional N supply.

Leaf and root transcriptome reprogramming in response to *R. irregularis* colonization

integrates a genotype and N-availability effects

To further characterize the biological processes involved in maize response to *R. irregularis*

colonization as a function of N fertilization, a whole genome transcriptomics approach was

conducted in leaves and a dual RNA-seq in roots of plants grown under HN and LN

conditions. The MN condition was considered as equivalent to HN and was not further

investigated. The leaf below the ear was chosen for providing a good proxy of the sink-to-

source transition during grain filling [57]. First, we defined the differentially expressed genes

(DEGs) as those exhibiting a fold change of the transcript levels in inoculated plants

compared to non-inoculated plants that was higher than 1.5 or lower than -1.5 (*p*<0.05).

Details for the numbers of DEGs depending on organ, N nutrition level and genotype are

shown in Figure 3. GO enrichment analysis revealed that for those common up-regulated

genes in roots, the biological processes overrepresented were those related to proteolysis,

lipid metabolism, ethylene response and membrane transport (Supplementary Figure 7). The

common down regulated genes were represented by five main functional categories

including cell wall biogenesis, metal homeostasis, phosphate transport, stress responses and

lipid metabolism (Supplementary Figure 7). Specific features of B73 root transcriptome

consist mainly on up-regulation of genes related to immunity (both under LN and HN).

Specific features of the root Lo3 transcriptome resides mainly on up-regulation of autophagy

related genes (under LN) and terpenoid metabolism genes, and down regulation of genes

related to DNA metabolism and replication (both under LN and HN) (Supplementary Figure

7). We performed the same data analysis in leaves (Figure 3, Supplementary Figure 7).

Specific features of B73 leaf transcriptome consist mainly on translation and auxin signalling

genes (under LN), and redox reactions, carbohydrate transport and defence related genes

(under HN). Specific features of the leaf Lo3 transcriptome reside in photosynthesis, cell wall

remodelling, cytoskeleton remodelling, stress response, signalling, while in the down-

regulated genes are related to gamma-amino butyric acid (GABA) metabolism and

pyrimidine biosynthesis (under both LN and HN). Taken together, these data show that leaf

and root transcriptomic responses to the AMF differ depending on maize genotype and N

supply.

Leaf and root metabolomics profiles in response to *R. irregularis* are shaped by N

nutrition

To further characterize the metabolic responses of the two maize lines to *R. irregularis*

inoculation*,* a primary metabolome profiling was performed on the same leaf and root

samples as those studied in the previous section. Again, plant physio-agronomical traits and

metabolomics are paired data. Metabolic profiles showed that plant organ and N supply had

the strongest impact on leaf and root metabolite composition (Supplementary Figures 8A and

B). AMF inoculation and maize genotype had globally weaker impacts on metabolite profiles

(Figures 4 A and B and supplementary Figure 8C and D). In leaf samples, 132 water soluble

metabolites were detected of which 103 were significantly different in at least on condition

after ANOVA statistical analysis ($P < 0.05$) followed by a Bonferroni post-hoc test and

correction. In roots, there were 120 water-soluble metabolites of which 117 were found to be

significantly different in at least one condition. The identity of metabolites can be found on the

interactive heatmaps (Supplementary interactive Files 1 and 2).

In roots, fungal inoculation triggered accumulation of lipids such as trans-13-octadecenoic

acid and palmitoleic in both lines Supplementary Figure 9 A. The accumulation of these two

lipids was much higher under HN than under LN. We observed that in inoculated roots, there

was a global decrease in sucrose, glucose and fructose levels Supplementary Figure 9 B.

Several metabolites linked to N metabolism were differentially accumulated in inoculated

roots Supplementary Figure 9 D. For example, Glu, Arg, citrulline, Asn. Some metabolites

related to immunity were accumulated differentially in response to the fungus. For instance,

shikimic acid was accumulated in LN roots of inoculated B73 plants compared to controls

Supplementary Figure 9 C. In leaves, sucrose showed a significant decrease in HN in both

maize lines (Supplementary Figure 10). Gln and Asn accumulated in Lo3 leaves under HN in

response to *R. irregularis*.

To identify metabolites representative of the plant response to *R. irregularis* under each

growth condition, metabolites for which abundance is significantly correlated with specific

treatment conditions were identified by using a variable selection approach in a multivariate

linear model conducted with the MultiVarSel approach (see materials and methods, Figure 4

C and D). For example, we found that under LN, xylulose, was correlated with leaves of Lo3

inoculated plants (Figure 4C). In roots, benzoic acid was accumulated in inoculated B73

roots of plants grown under LN (Figure 4D). These data indicate that specific metabolites

accumulation is correlated with particular conditions depending on the plant genotype and

the N supply in response to *R. irregularis*.

Multi-organ metabolic modelling reveals key metabolic processes triggered by *R.

irregularis allowing maize to mitigate N limitation stress**

To identify key metabolic reactions involved in yield and biomass reshaping in response to *R.*

irregularis symbiosis, we integrated the transcriptomic data in a recently reconstructed multi-

organ genome scale modelling (GSM) iZMA6517 using EXTREAM [34]. This was followed by a

metabolic bottleneck analysis (MBA) [34] to pinpoint metabolic bottlenecks for biomass growth

rate. First, the B73 leaf and root transcriptomic data were used as input to EXTREAM to

contextualize iZMA6517. EXTREAM equally distributes transcript levels for each gene based

on the number of cognate biochemical reactions and contextualize the GSM. The in silico

analysis predicted that growth rates were increased under LN in response to *R. irregularis* inoculation compared to non-inoculated plants, which is expected to lead to grain yield and biomass increase. While under HN, a negative impact on maize growth rate was predicted (Figure 5A). In particular, the model predicted absence of grain production in control of LN B73 plants. Strikingly, these model predictions agree with the observed yield and biomass experimental data in Figure 1, confirming the predictive accuracy of the iZMA6517. When applied to the Lo3 genotype leaf and root transcriptomic data, iZMA6517 predicted a slight increase in grain yield and leaf biomass under LN in Lo3 inoculated plants. The model was better fitting the experimental B73 phenotype in terms of biomass compared to the Lo3 experimental data.

This result prompted us to determine the metabolic processes that could explain the improved yield under LN B73 plants and LN Lo3 in response to *R. irregularis* inoculation by identifying the potential metabolic bottlenecks relieved by the AMF under limiting N using MBA (see materials and methods). MBA expands the flux space of each reaction individually and assesses the impact of the expanded flux space on the biomass growth rate. For this purpose, we compared the bottlenecks identified in LN control plants to the bottlenecks identified in inoculated LN plants (Figure 5B; Table 1 and 2). Four metabolic bottlenecks relieved by *R. irregularis* inoculation were identified (Table 1). Interestingly 13 new bottlenecks were emerged following *R. irregularis* inoculation. On the other hand, we found that four metabolic bottlenecks that were present in LN control plants were relieved by N supply in the HN plants. There were three metabolic bottlenecks relieved by both HN supply conditions and *R. irregularis* inoculated condition (photosynthesis, purine and pyrimidine metabolism, aromatic acid metabolism). The same analysis was conducted in Lo3 maize genotype. There were 6 metabolic bottlenecks relieved by *R. irregularis* inoculation under LN conditions (Table 2). These bottlenecks are involved in Pyrimidine metabolism, Purine metabolism, and Calvin-Benson-Bassham cycle. These data suggest that in both maize lines, *R. irregularis* inoculation triggers modifications in pyrimidine metabolism. More

specifically, the model predicts that the limiting reactions that are relieved by symbiosis

should result in an accumulation of UMP, UDP and UTP in leaves under LN. Thus, the levels

of UMP, UDP, UDP-glc and UTP were monitored in leaves under LN and HN (Figure 5D and

Supplementary Figure 12). Remarkably, we found an increase in UMP, UDP and UDP-glc

levels in leaves of plants inoculated with the AMF under LN only in the Lo3 genotype, thus

confirming MBA predictions. Together these data indicate that the metabolic modelling was a

good predictor of biomass in the context of maize-*R. irregularis* interaction especially for B73

maize genotype. The MBA analysis was able to predict that pyrimidine biosynthesis is crucial

in this interaction, in agreement with experimental observation of pyrimidine accumulation in

leaves of symbiotic Lo3 plants under LN.

Fungal transcriptome of in inoculated maize roots is driven by N availability

We then investigated the metabolic cross talk between *R. irregularis* and maize from th

fungal side. For this purpose, we determined the fungal genes displaying a higher expression

in LN roots compared to HN roots and vice versa. This selection was performed for B73 and

Lo3 separately. The Venn diagrams in Figure 6A and 6B show the numbers of fungal genes

up-regulated under LN compared to HN or vice versa in each maize line. In both maize lines,

the functional categories overrepresented in *R. irregularis* DEG were similar in each N

nutrition condition Figure 6C. Among the DEG functional categories, several genes were

involved in amino acid and nucleotide metabolism and transport as well as in inorganic

transport. We focused on the expression profiles of nucleotide and amino acid metabolism

genes and transporter genes as well as phosphate and ammonium acid transport genes in

the annotation database of JGI (https://genome.jgi.doe.gov/portal/). Supplementary Table 6

shows the expression pattern of these fungal genes. Interestingly, fungal ammonia

transporters are more highly expressed under LN than under HN. The fungal amino acid

biosynthetic genes up-regulated are mainly involved in asparagine, arginine, glutamine,

glutamate and serine pathways. Several fungal amino acid transporter genes were found to

be up-regulated either in LN or HN roots. The fungal genes related to pyrimidine de novo

biosynthesis and ureide metabolism (Figure 6D) are expressed in both LN and HN roots.

Interestingly, although expressed under all nutritional conditions they are globally up-

regulated in HN roots.   These data show that the fungal transcriptome is strongly affected by

the nutritional N status of the host and the soil and that fungal amino acid and pyrimidine

metabolism genes are expressed in symbiotic roots.

 Discussion_1335 **words**

 In our study, we showed that a symbiotic association with *R. irregularis* was able to
compensate for decreased maize plant performance when N fertilization is reduced. In the
literature, depending on both the N supply level and the form of N supplied, different effects
on yield and/or biomass production have been reported, including an increase, no effect, or a
reduction in productivity when the plant develops a symbiotic association with an AMF
[26,58,59]. In the present investigation, we estimated that kernel yield can be maintained with a 2-
to 5-fold reduction in N input when two maize lines were inoculated with *R. irregularis*. Such
a beneficial impact was previously observed for maize plant biomass production under low N
fertilizer inputs [26]. One can, therefore, conclude that the positive impact of AMF on plant
performance can be variable depending on the AMF and the host genotype as well as the
level and type of available N source [9,13,58].

Our transcriptome study allowed us to provide an overview of biological processes putatively
involved during the establishment of a symbiotic association between the AMF *R. irregularis*
and maize. The DEGs that were either up or down-regulated in roots were enriched in
transcripts encoding proteins involved in a variety of biological functions : defence, hormone
signalling N and C metabolism, cell wall and chromatin remodelling, proteolysis, metal
homeostasis and lipid metabolism. Similar results were previously obtained in several other
model and crop species [27,60–67]. In particular, the most highly up-regulated genes in roots of
B73 and Lo3 under both N nutrition conditions in response to *R. irregularis* inoculation are
encoding peptidases, peptidase inhibitors, chitinases and senescence genes
(Supplementary Table 7a and b). Interestingly, autophagy genes are overrepresented in Lo3
genes up-regulated in roots under LN following inoculation with *R. irregularis* (Supplementary
Table 8, Supplementary Figure 7). These data suggest that important proteolysis processes
are taking place in in inoculated roots that may involve autophagy especially in the Lo3
maize line. Autophagy is a process known to recycle N during senescence, and is known to
provide N assimilates necessary for maize kernel filling [68,69].

Changes in the level of expression of genes involved in sugar, amino acid, and lipid

metabolism were confirmed by metabolomics analyses (Supplementary Figure 9 and 10).

Globally, we observed lipid accumulation in roots colonized by AMFs and a drop in soluble

sugars in both roots and shoots, a phenomenon already described [16,70]. This is considered a

way of fuelling the symbiotic fungus with a carbon source as lipids. In inoculated plants,

arginine (Arg), known to be a key molecule involved in the transfer of N from the fungus to

the plant [11,13,66,71] accumulates under HN conditions not only in the roots of line B73 HN but

also in the leaves.

To delve deeper than a mere data overview, the incorporation of transcriptomics data into the

iZMA6517 metabolic model is especially potent, as we observed concurrence between

iZMA6517 predictions and actual experimental results regarding biomass increase triggered

by the AMF. Our previous efforts of maize metabolic modelling resulted in excellent

phenotypical predictions for including the photosynthetic behaviour of maize leaf [32],

predicting the root phenotype under nitrogen-optimal and nitrogen deficient conditions[33] and

whole plant phenotype for heat and cold stress conditions [34]. Thereby, our whole-plant model

of maize, iZMA6517 [34] was an ideal candidate to identify metabolic underpinnings of maize

for different nitrogen availability. Interestingly, iZMA6517 allowed the identification of

pyrimidine metabolism as a crucial player in this symbiosis. Indeed, our transcriptomic

analysis demonstrates that, regardless of nitrogen feeding conditions, pyrimidine

biosynthesis processes are consistently downregulated in leaves of inoculated plants

(Supplementary Figure 7, Supplementary Table 10). Under LN conditions, several key genes

encoding enzymes involved in nucleotide metabolism, including putative Orotidine-5'-

phosphate decarboxylase, thiaminase, Uridine 5'-monophosphate synthase, Cytidine

triphosphate synthase, are downregulated in leaves (Supplementary Table 10). A critical

gene in this context is *Zm00001d023833*, encoding a putative nucleoside diphosphate kinase

(NDPK), which participates in pyrimidine and purine nucleotide salvage [72]. This enzyme was

identified as a metabolic bottleneck relieved by AMF inoculation (Figure 5, Table 1 and 2)

and its expression in the context of symbiosis was predicted to lead to accumulation of

pyrimidine in leaves. Remarkably, this accumulation was confirmed experimentally by the

accumulation of UMP and UDP in Lo3 leaves of LN symbiotic plants (Figure 5D).

In line with these data, fungal transcriptomic analysis by RNAseq shows that pyrimidine and

ureide metabolism are expressed by *R. irregularis*. The integration of maize and fungal RNA-

seq and metabolome analyses has allowed us to construct a simplified model depicting the

various metabolic N related events occurring in the symbiotic association considering both

lines (Figure 7, Supplementary Figure 11). Under LN, fungal argininosuccinate lyase,

carbamoyl-phosphate synthase and arginase genes are upregulated, potentially leading to

urea accumulation. However, urea level drops in LN inoculated roots (Supplementary Figure

11), a fact that is consistent with higher expression of *R. irregularis* urease under LN (Figure

6, Supplementary Table 6). In parallel, fungal genes involved in pyrimidine de novo synthesis

and salvage are expressed in both Lo3 and B73 symbiotic roots under LN and HN (Figure 6,

Supplementary Table 6). Our hypothesis is that pyrimidines are transferred to the leaves via

the phloem sap, and that they are provided by the fungus especially under LN. This is

consistent with the finding that in Lo3 leaves of symbiotic plants, UMP, UDP and UDP-glc

levels were higher compared to control plants. The fact that in B73 leaves, and in HN leaves,

we did not observe an increase of these metabolites in response to AMF inoculation

indicates that their accumulation is tightly controlled. In our experiment the level of

pyrimidines was globally higher in HN leaves than LN leaves but no impact of AMF

inoculation was observed in HN plants (Supplementary Figure 12) indicating that the steady

state of these compounds may vary over time. On the other hand, allantoin accumulates

exclusively in HN roots and its level remains unaffected by inoculation (Supplementary

Figure 11). This metabolite can be converted into allantoic acid, which may be released into

the apoplast [73]. A fungal allantoicase is upregulated in HN roots, indicating possible

conversion of allantoic acid into urea under HN conditions. This transfer of ureides is akin to

the process observed in symbiotic associations between tropical legumes and Rhizobia

species [73]. Previous studies have also identified upregulation of genes involved in allantoin

synthesis in roots of AMF-inoculated plants [74]. Consequently, it is plausible that ureide

transfer mirrors that occurring in legume nodules.

Fungal arbuscules are known to undergo controlled senescence [17,22]. Our data suggest that

autophagy and proteolysis in general are upregulated in mycorrhizal maize roots, with higher

expression in Lo3 LN roots (Figure 3 and Supplementary Table 8). Autophagy may

contributes to the degradation of fungal cellular products, potentially releasing nucleosides.

The fungal nucleotide salvage process can then recycle nucleosides that are probably

transported to the leaves thus repressing de novo plant pyrimidine biosynthesis pathway

(Supplementary Figure 7, Supplementary Table 9) [72,73]. This implies that AMF likely supplies

the host plant with nucleotides and nucleosides, potentially pyrimidines, to circumvent the

deficiency in NDPK activity, thereby alleviating the associated metabolic bottleneck. This

observation is consistent with the known role of NDPK in plant survival with respect to

pyrimidine salvage [72,73]. It is noteworthy that pyrimidine biosynthesis can be an essential

process for plant N management [75,76]. Pyrimidine metabolism holds significance not only in

crop performance but also in nitrogen use efficiency (NUE) [73,77,78]. Recent findings

demonstrate a correlation between maize leaf nitrogen accumulation and NDPK protein

levels [79], further supporting the hypothesis that *R. irregularis* stimulates the reprogramming of

plant pyrimidine metabolism to enhance maize performance under nitrogen-limiting

conditions.

Physiological profiling combined with metabolic modelling revealed pyrimidine and ureide

metabolism as previously unrecognized pivotal components in this interaction, warranting

additional investigation [80,81]. Additionally, the contrasting strategies adopted by our two

maize lines to capitalize on the fungus highlight the significance of harnessing maize's broad

genetic diversity for a holistic comprehension of this symbiotic relationship [24,82,83].

Data Availability

RNAseq project is deposited (Gene Expression Omnibus,Edgard R. et al. 2002): https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE235654 ; accession no. GSE235654. All steps of the experiment, from growth conditions to bioinformatic analyses, were detailed in CATdb [84]: http://tools.ips2.u-psud.fr.fr/CATdb/; Project: NGS2021_19_Rhizophagus according to the MINSEQE 'minimum information about a high throughput sequencing experiment'.

Acknowledgments:

BD acknowledges Université Paris Saclay for PhD fellowship. AD acknowledges the benefits received from France 2030 program (Saclay Plant Sciences, reference n° ANR-17-EUR-0007, EUR SPS-GSR integrated into France 2030 (reference n° ANR-11-IDEX-0003-02), INRAE Grant IB BAP 2021 MYCORN, and IJPB's Plant Observatory technological platforms. RS gratefully acknowledges funding support from National Science Foundation (NSF) CAREER grant (1943310) and Nebraska Collaboration Initiative Grant (21-1106-6011). We thank Michel Lebrusq, Lilan Diahuron, Patrick Grillot, and Christian Jeudy for taking care of the plants in the greenhouse. We thank the POPS platform for the RNAseq experiment and data. We also thank Etienne Delannoy from POPS platform for fruitful discussions. We are grateful to Cyril Bauland and Carine Palaffre from the INRAE for providing seeds from the Centre de Ressources Biologiques of Saint Martin de Hinx (France). We acknowledge Pr. Mario MOTTO from ISC, Bergamo (Italy) for providing Lo3 seeds.

675 **Table 1: List of metabolic bottlenecks relieved and created by *R. irregularis* or N supply in B73.**
676 Metabolic bottlenecks were identified by MBA that are relieved by either R. irregularis treatment or HN
677 treatment compared to LN control conditions are indicated as "debottlnecked". Those that were newly
678 created following R. irregularis or HN treatment were compared to LN control conditions are indicated
679 as "new metabolic" bottelnecks. M : mesophyll, R : root, p : plastid, t : thylakoid, c : cytosol, x : vacuole.

Reactions Debottlenecked by *R. irregularis* treatment under LN compared to LN CTRL	
R03817[M,t]	Photosynthesis light reactions
R00124[R,p]	Purine and Pyrimidine metabolism
R00692[R,p]	Phenylalanine, tyrosine, and tryptophan metabolism
R04241_mod[R,m]	Folate metabolism
Reactions Debottlenecked by HN treatment compared to LN CTRL	
R03817[M,t]	Photosynthesis light reactions
R00124[R,p]	Purine and Pyrimidine metabolism
R00692[R,p]	Phenylalanine, tyrosine, and tryptophan metabolism
R00095[R,x]	Ascorbate and aldarate metabolism
New bottlenecks with LN RHIZO compared to LN CTRL	
R03346[M,c]	Nicotinate and nicotinamide metabolism
R03346[B,c]	Nicotinate and nicotinamide metabolism
R00086[M,pm]	Purine metabolism
R00986[R,p]	Purine metabolism
MR00968[B,c]	Benzoate metabolism
R00086[B,pm]	Purine metabolism
R00341[R,c]	Pyruvate metabolism
R00216[B,p]	Pyruvate metabolism
R01058[B,c]	Glycolysis
R02678[R,m]	Phenylalanine, tyrosine, and tryptophan metabolism
R00710[R,m]	Pyruvate metabolism
R01067[R,c]	Calvin-Benson-Bassham cycle
R01067[R,p]	Calvin-Benson-Bassham cycle
New bottlenecks in HN CTRL compared to LN CTRL	
R00341[R,c]	Pyruvate metabolism
R00709_mod[R,m]	TCA cycle
R00713[R,m]	Alanine, aspartate, and glutamate metabolism
R01067[R,c]	Calvin-Benson-Bassham cycle
R01067[R,p]	Calvin-Benson-Bassham cycle
R01667[R,p]	Pyrimidine metabolism
R00235[B,x]	Pyruvate metabolism
R00885[B,c]	Fructose and mannose metabolism
R01058[B,c]	Glycolysis
R02080[M,c]	Phenylalanine, tyrosine, and tryptophan metabolism
R00214[M,m]	Pyruvate metabolism
R00124[B,p]	Purine metabolism
R00267_mod[M,c]	Alanine, aspartate, and glutamate metabolism
R00235[R,p]	Pyruvate metabolism

681 **Table 2: List of metabolic bottlenecks relieved and created by *R. irregularis* or N supply on Lo3.**
682 Metabolic bottlenecks were identified by MBA that are relieved by either *R. irregularis* treatment or HN
683 treatment compared to LN control conditions are indicated as "debottlenecked". Those that were newly
684 created following *R. irregularis* or HN treatment were compared to LN control conditions are indicated
685 as "new metabolic" bottlenecks. M : mesophyll, R : root, p : plastid, t : thylakoid, c : cytosol, x : vacuole.

Reactions Debottlenecked by *R. irregularis* treatment under LN compared to LN CTRL	
R01080[R,p]	Pyrimidine metabolism
R02143[R,p]	Purine metabolism
R01667[R,p]	Pyrimidine metabolism
R01067[R,c]	Calvin-Benson-Bassham cycle
R01067[R,p]	Calvin-Benson-Bassham cycle
R00281[R,p]	Non pathway information
Reactions Debottlenecked by HN treatment compared to LN CTRL	
R04241_mod[R,m]	Folate metabolism
R05395[R,p]	DDT degradation
New bottlenecks with LN RHIZO compared to LN CTRL	
R00709_mod[R,m]	TCA cycle
R00713[R,m]	Alanine, aspartate, and glutamate metabolism
MR00886[R,m]	Acyl carrier protein metabolism
New bottlenecks in HN CTRL compared to LN CTRL	
R00709_mod[R,m]	TCA cycle
R00713[R,m]	Alanine, aspartate, and glutamate metabolism
R00885[B,c]	Fructose and mannose metabolism

References:

1. Hirel, B., Tétu, T., Lea, P. J. & Dubois, F. Improving Nitrogen Use Efficiency in Crops for Sustainable Agriculture. *Sustainability* vol. 3 (2011).

2. Hirel, B. & Krapp, A. B. T.-R. M. in L. S. Nitrogen Utilization in Plants I Biological and Agronomic Importance. in (Elsevier, 2020). doi:https://doi.org/10.1016/B978-0-12-809633-8.21265-X.

3. Hirel, B., Le Gouis, J., Ney, B. & Gallais, A. The challenge of improving nitrogen use efficiency in crop plants: Towards a more central role for genetic variability and quantitative genetics within integrated approaches. in *Journal of Experimental Botany* (2007). doi:10.1093/jxb/erm097.

4. Schlesinger, W. H. On the fate of anthropogenic nitrogen. *Proc. Natl. Acad. Sci.* **106**, 203 LP – 208 (2009).

5. Lassaletta, L. *et al.* Nitrogen use in the global food system: past trends and future trajectories of agronomic performance, pollution, trade, and dietary demand. *Environ. Res. Lett.* **11**, 095007 (2016).

6. Lea, P. J. & Miflin, B. J. Nitrogen Assimilation and its Relevance to Crop Improvement. *Annual Plant Reviews online* 1–40 (2018) doi:doi:10.1002/9781119312994.apr0448.

7. Liu, Q. *et al.* Improving Crop Nitrogen Use Efficiency Toward Sustainable Green Revolution. *Annu. Rev. Plant Biol.* **73**, 523–551 (2022).

8. Gianinazzi, S. *et al.* Agroecology: the key role of arbuscular mycorrhizas in ecosystem services. *Mycorrhiza* **20**, 519–530 (2010).

9. Verzeaux, J., Hirel, B., Dubois, F., Lea, P. J. & Tétu, T. Agricultural practices to improve nitrogen use efficiency through the use of arbuscular mycorrhizae: Basic and agronomic aspects. *Plant Sci.* **264**, 48–56 (2017).

10. Wipf, D., Krajinski, F., van Tuinen, D., Recorbet, G. & Courty, P.-E. Trading on the arbuscular mycorrhiza market: from arbuscules to common mycorrhizal networks. *New Phytol.* **223**, 1127–1142 (2019).

11. Dellagi, A., Quillere, I. & Hirel, B. Beneficial soil-borne bacteria and fungi: a promising way to improve plant nitrogen acquisition. *J. Exp. Bot.* **71**, 4469–4479 (2020).

12. Gojon, A. Nitrogen acquisition in arbuscular mycorrhizal symbioses: A step into the real world. *J. Plant Physiol.* **271**, 153646 (2022).

13. Xie, K. *et al.* Plant nitrogen nutrition: The roles of arbuscular mycorrhizal fungi. *J. Plant Physiol.* **269**, 153591 (2022).

14. Chen, E. C. H. *et al.* High intraspecific genome diversity in the model arbuscular mycorrhizal symbiont Rhizophagus irregularis. *New Phytol.* **220**, 1161–1171 (2018).

15. Smith, S. E. & Read, D. J. Colonization of roots and anatomy of VA mycorrhizas. *Mycorrhizal Symbiosis* 33–II (2002) doi:10.1016/b978-012652840-4/50003-6.

16. MacLean, A. M., Bravo, A. & Harrison, M. J. Plant Signaling and Metabolic Pathways Enabling Arbuscular Mycorrhizal Symbiosis. *Plant Cell* **29**, 2319–2335 (2017).

17. Choi, J., Summers, W. & Paszkowski, U. Mechanisms Underlying Establishment of Arbuscular Mycorrhizal Symbioses. *Annu. Rev. Phytopathol.* **56**, 135–160 (2018).

18. Breuillin, F. *et al.* Phosphate systemically inhibits development of arbuscular mycorrhiza in Petunia hybrida and represses genes involved in mycorrhizal

functioning. *Plant J.* **64**, 1002–1017 (2010).

19.   Smith, S. E., Jakobsen, I., Grønlund, M. & Smith, F. A. Roles of Arbuscular
Mycorrhizas in Plant Phosphorus Nutrition: Interactions between Pathways of
Phosphorus Uptake in Arbuscular Mycorrhizal Roots Have Important Implications for
Understanding and Manipulating Plant Phosphorus Acquisition. *Plant Physiol.* **156**,
1050–1057 (2011).

20.   Balzergue, C., Chabaud, M., Barker, D., Bécard, G. & Rochange, S. High phosphate
reduces host ability to develop arbuscular mycorrhizal symbiosis without affecting root
calcium spiking responses to the fungus . *Frontiers in Plant Science* vol. 4 (2013).

21.   Wang, W. *et al.* Nutrient Exchange and Regulation in Arbuscular Mycorrhizal
Symbiosis. *Mol. Plant* **10**, 1147–1158 (2017).

22.   Paries, M. & Gutjahr, C. The good, the bad, and the phosphate: regulation of
beneficial and detrimental plant–microbe interactions by the plant phosphate status.
New Phytol. **239**, 29–46 (2023).

23.   Liu, J., Fernie, A. R. & Yan, J. The Past, Present, and Future of Maize Improvement:
Domestication, Genomics, and Functional Genomic Routes toward Crop
Enhancement. *Plant Commun.* **1**, 100010 (2020).

24.   Sawers, R. J. H. H., Ramírez-Flores, M. R., Olalde-Portugal, V. & Paszkowski, U. The
impact of domestication and crop improvement on arbuscular mycorrhizal symbiosis in
cereals: insights from genetics and genomics. *New Phytol.* **220**, 1135–1140 (2018).

25.   Berger, F. & Gutjahr, C. Factors affecting plant responsiveness to arbuscular
mycorrhiza. *Curr. Opin. Plant Biol.* **59**, 101994 (2021).

26.   Quiroga, G., Erice, G., Aroca, R., Delgado-Huertas, A. & Ruiz-Lozano, J. M.
Elucidating the Possible Involvement of Maize Aquaporins and Arbuscular Mycorrhizal
Symbiosis in the Plant Ammonium and Urea Transport under Drought Stress
Conditions. *Plants* vol. 9 (2020).

27.   Hui, J. *et al.* The mycorrhiza-specific ammonium transporter ZmAMT3;1 mediates
mycorrhiza-dependent nitrogen uptake in maize roots. *Plant Cell* **34**, 4066–4087
(2022).

28.   Orth, J. D., Thiele, I. & Palsson, B. Ø. What is flux balance analysis? *Nat. Biotechnol.*
28, 245–248 (2010).

29.   Mahadevan, R. & Schilling, C. H. The effects of alternate optimal solutions in
constraint-based genome-scale metabolic models. *Metab. Eng.* **5**, 264–276 (2003).

30.   Lewis, N. E. *et al.* Omic data from evolved *E. coli* are consistent with computed
optimal growth from genome‐scale models. *Mol. Syst. Biol.* **6**, 390 (2010).

31.   Morris, D. S., Chowdhury, N. B., Kathol, M. & Saha, R. Opening the door to nitrogen
fixation in oxygenic phototrophs. *Trends Biotechnol.* **41**, 283–285 (2023).

32.   Simons, M. *et al.* Assessing the Metabolic Impact of Nitrogen Availability Using a
Compartmentalized Maize Leaf Genome-Scale Model   . *Plant Physiol.* **166**, 1659–
1674 (2014).

33.   Chowdhury, N. B. *et al.* Dissecting the metabolic reprogramming of maize root under
nitrogen-deficient stress conditions. *J. Exp. Bot.* **73**, 275–291 (2022).

34.   Chowdhury, N. B. *et al.* A multi-organ maize metabolic model connects temperature
stress with energy production and reducing power generation. *iScience* **26**, 108400
(2023).

35. Schnable, P. S. *et al.* The B73 Maize Genome: Complexity, Diversity, and Dynamics.
Science (80-.). **326**, 1112–1115 (2009).

36. Cañas, R. A. *et al.* Exploiting the Genetic Diversity of Maize Using a Combined
Metabolomic, Enzyme Activity Profiling, and Metabolic Modeling Approach to Link Leaf
Physiology to Kernel Yield. *Plant Cell* **29**, 919–943 (2017).

37. Decouard, B. *et al.* Genotypic Variation of Nitrogen Use Efficiency and Amino Acid
Metabolism in Barley   . *Frontiers in Plant Science*   vol. 12 (2022).

38. Murphy, J. & Riley, J. P. A modified single solution method for the determination of
phosphate in natural waters. *Anal. Chim. Acta* **27**, 31–36 (1962).

39. Giovannetti, M. & Mosse, B. An Evaluation of Techniques for Measuring Vesicular
Arbuscular Mycorrhizal Infection in Roots. *New Phytol.* **84**, 489–500 (1980).

40. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21
(2013).

41. Fiehn, O. *et al.* Quality control for plant metabolomics: reporting MSI-compliant
studies. *Plant J.* **53**, 691–704 (2008).

42. Fiehn, O. Metabolite Profiling in Arabidopsis BT - Arabidopsis Protocols. in (eds.
Salinas, J. & Sanchez-Serrano, J. J.) 439–447 (Humana Press, 2006). doi:10.1385/1-
59745-003-0:439.

43. Guérard, F., Pétriacq, P., Gakière, B. & Tcherkez, G. Liquid chromatography/time-of-
flight mass spectrometry for the analysis of plant samples: A method for simultaneous
screening of common cofactors or nucleotides and application to an engineered plant
line. *Plant Physiol. Biochem.* **49**, 1117–1125 (2011).

44. Murtagh, F. & Legendre, P. Ward's Hierarchical Agglomerative Clustering Method:
Which Algorithms Implement Ward's Criterion? *J. Classif.* **31**, 274–295 (2014).

45. Galili, T., O'Callaghan, A., Sidi, J. & Sievert, C. heatmaply: an R package for creating
interactive cluster heatmaps for online publishing. *Bioinformatics* **34**, 1600–1602
(2018).

46. Perrot-Dockès, M., Lévy-Leduc, C., Sansonnet, L. & Chiquet, J. Variable selection in
multivariate linear models with high-dimensional covariance matrix estimation. *J.*
Multivar. Anal. **166**, 78–97 (2018).

47. Perrot-Dockès, M. *et al.* A variable selection approach in the multivariate linear model:
an application to LC-MS metabolomics data. **17**, (2018).

48. Grandclaudon, M. *et al.* A Quantitative Multivariate Model of Human Dendritic Cell-T
Helper Cell Communication. *Cell* **179**, 432-447.e21 (2019).

49. Perrot-Dockès, M., Lévy-Leduc, C. & Rajjou, L. Estimation of Large Block Structured
Covariance Matrices: Application to 'Multi-Omic' Approaches to Study Seed Quality. *J.*
R. Stat. Soc. Ser. C Appl. Stat. **71**, 119–147 (2022).

50. Chowdhury, N. B., Alsiyabi, A. & Saha, R. Characterizing the Interplay of Rubisco and
Nitrogenase Enzymes in Anaerobic-Photoheterotrophically Grown
Rhodopseudomonas palustris CGA009 through a Genome-Scale Metabolic and
Expression Model. *Microbiol. Spectr.* **10**, e01463-22 (2022).

51. Ramírez-Flores, M. R., Bello-Bello, E., Rellán-Álvarez, R., Sawers, R. J. H. & Olalde-
Portugal, V. Inoculation with the mycorrhizal fungus Rhizophagus irregularis
modulates the relationship between root growth and nutrient content in maize (Zea
mays ssp. mays L.). *Plant Direct* **3**, e00192 (2019).

52. Wang, E. *et al.* A H+-ATPase That Energizes Nutrient Uptake during Mycorrhizal
Symbioses in Rice and Medicago truncatula . *Plant Cell* **26**, 1818–1830 (2014).

53. Sawers, R. J. H. *et al.* Phosphorus acquisition efficiency in arbuscular mycorrhizal
maize is correlated with the abundance of root-external hyphae and the accumulation
of transcripts encoding PHT1 phosphate transporters. *New Phytol.* **214**, 632–643
(2017).

54. Krapp, A. *et al.* Nitrate transport and signalling in Arabidopsis. *Journal of Experimental*
Botany (2014) doi:10.1093/jxb/eru001.

55. Wang, Y.-Y., Cheng, Y.-H., Chen, K.-E. & Tsay, Y.-F. Nitrate Transport, Signaling, and
Use Efficiency. *Annu. Rev. Plant Biol.* **69**, 85–122 (2018).

56. Dechorgnat, J. *et al.* Tissue and nitrogen-linked expression profiles of ammonium and
nitrate transporters in maize. *BMC Plant Biol.* **19**, 206 (2019).

57. Prioul, J.-L. & Schwebel-Dugué, N. Source-Sink Manipulations and Carbohydrate
Metabolism in Maize. *Crop Sci.* **32**, cropsci1992.0011183X003200030035x (1992).

58. Ingraffia, R. *et al.* Nitrogen Type and Availability Drive Mycorrhizal Effects on Wheat
Performance, Nitrogen Uptake and Recovery, and Production Sustainability .
Frontiers in Plant Science vol. 11 760 (2020).

59. Yooyongwech, S. *et al.* Matching of Nitrogen Enhancement and Photosynthetic
Efficiency by Arbuscular Mycorrhiza in Maize (Zea mays L.) in Relation to Organic
Fertilizer Type. *Plants* vol. 11 (2022).

60. Fiorilli, V. *et al.* Global and cell-type gene expression profiles in tomato plants
colonized by an arbuscular mycorrhizal fungus. *New Phytol.* **184**, 975–987 (2009).

61. Gallou, A., Declerck, S. & Cranenbrouck, S. Transcriptional regulation of defence
genes and involvement of the WRKY transcription factor in arbuscular mycorrhizal
potato root colonization. *Funct. Integr. Genomics* **12**, 183–198 (2012).

62. Schaarschmidt, S., Gresshoff, P. M. & Hause, B. Analyzing the soybean transcriptome
during autoregulation of mycorrhization identifies the transcription factors GmNF-
YA1a/b as positive regulators of arbuscular mycorrhization. *Genome Biol.* **14**, R62
(2013).

63. Meng, L. *et al.* Arbuscular mycorrhizal fungi and rhizobium facilitate nitrogen uptake
and transfer in soybean/maize intercropping system . *Frontiers in Plant Science* vol.
6 (2015).

64. Handa, Y. *et al.* RNA-seq Transcriptional Profiling of an Arbuscular Mycorrhiza
Provides Insights into Regulated and Coordinated Gene Expression in Lotus japonicus
and Rhizophagus irregularis. *Plant Cell Physiol.* **56**, 1490–1511 (2015).

65. Vangelisti, A. *et al.* Transcriptome changes induced by arbuscular mycorrhizal fungi in
sunflower (Helianthus annuus L.) roots. *Sci. Rep.* **8**, 4 (2018).

66. Kaur, S. & Suseela, V. Unraveling Arbuscular Mycorrhiza-Induced Changes in Plant
Primary and Secondary Metabolome. *Metabolites* vol. 10 (2020).

67. Wang, S. *et al.* Functional analysis of the OsNPF4.5 nitrate transporter reveals a
conserved mycorrhizal pathway of nitrogen acquisition in plants. *Proc. Natl. Acad. Sci.*
U. S. A. **117**, 16649–16659 (2020).

68. Li, F. *et al.* Autophagic Recycling Plays a Central Role in Maize Nitrogen
Remobilization. *Plant Cell* **27**, 1389–1408 (2015).

69. Masclaux-Daubresse, C., Chen, Q. & Havé, M. Regulation of nutrient recycling via
autophagy. *Curr. Opin. Plant Biol.* **39**, 8–17 (2017).

70. Keymer, A. *et al.* Lipid transfer from plants to arbuscular mycorrhiza fungi. *Elife* **6**,
e29107 (2017).

71. Chen, A., Gu, M., Wang, S., Chen, J. & Xu, G. Transport properties and regulatory
roles of nitrogen in arbuscular mycorrhizal symbiosis. *Semin. Cell Dev. Biol.* **74**, 80–88
(2018).

72. Witte, C. P. & Herde, M. Nucleotide metabolism in plants. *Plant Physiol.* **182**, 63–78
(2020).

73. Tegeder, M. Transporters involved in source to sink partitioning of amino acids and
ureides: opportunities for crop improvement. *J. Exp. Bot.* **65**, 1865–1878 (2014).

74. Nanjareddy, K., Arthikala, M.-K., Gómez, B.-M., Blanco, L. & Lara, M. Differentially
expressed genes in mycorrhized and nodulated roots of common bean are associated
with defense, cell wall architecture, N metabolism, and P metabolism. *PLoS One* **12**,
e0182328 (2017).

75. Zrenner, R., Stitt, M., Sonnewald, U. & Boldt, R. PYRIMIDINE AND PURINE
BIOSYNTHESIS AND DEGRADATION IN PLANTS. *Annu. Rev. Plant Biol.* **57**, 805–
836 (2006).

76. Fernie, A. R. *et al.* Synchronization of developmental, molecular and metabolic
aspects of source–sink interactions. *Nat. Plants* **6**, 55–66 (2020).

77. Lee, D.-K. *et al.* A Nitrogen Molecular Sensing System, Comprised of the
ALLANTOINASE and UREIDE PERMEASE 1 Genes, Can Be Used to Monitor N
Status in Rice   . *Frontiers in Plant Science*   vol. 9 (2018).

78. Melino, V. J., Tester, M. A. & Okamoto, M. Strategies for engineering improved
nitrogen use efficiency in crop plants via redistribution and recycling of organic
nitrogen. *Curr. Opin. Biotechnol.* **73**, 263–269 (2022).

79. Urrutia, M. *et al.* Identification of metabolic and protein markers representative of the
impact of mild nitrogen deficit on agronomic performance of maize hybrids. (2023).

80. BARROSO, J., CASIMIRO, A., CARRAPIÇO, F., SLOMÉ, M. & PAIS, S. Localization
of uricase in mycorrhizas of Ophrys lutea Cav. *New Phytol.* **108**, 335–340 (1988).

81. Thakur, K. & Garg, N. Differential Role of Silicon and Rhizoglomus intraradices in
Modulating Amide and Ureide Metabolism of Seasonally Different Legume Species
Subjected to Nickel Toxicity. *Silicon* (2023) doi:10.1007/s12633-023-02599-w.

82. Schmidt, J. E., Bowles, T. M. & Gaudin, A. C. M. Using Ancient Traits to Convert Soil
Health into Crop Yield: Impact of Selection on Maize Root and Rhizosphere Function
. *Frontiers in Plant Science*   vol. 7 (2016).

83. Liu, H., Brettell, L. E. & Singh, B. Linking the Phyllosphere Microbiome to Plant Health.
Trends Plant Sci. **25**, 841–844 (2020).

84. Gagnot, S. *et al.* CATdb: a public access to Arabidopsis transcriptome data from the
URGV-CATMA platform. *Nucleic Acids Res.* **36**, D986–D990 (2008).

Figure 1: Impact of N supply on maize responses to *R. irregularis* for biomass and N content shows more pronounced beneficial AMF effects under LN. A and, B : grain yield and aerial part (AP) biomass. C and D : Nitrogen Use Efficiency (NUE) in grains and in AP. Ctrl: Mock non inoculated plants, Rhizo: inoculated with *R. irregularis*. Boxplots depict the interquartile range, with the median demarcated by a horizontal line in the middle, upper, and lower part of the box, representing the first and third quartiles and the range of data distribution. Dots overlaying the boxplots represent independent plants, each corresponding to a single observation in the dataset (N=7 to 10 plants). Different letters indicate significant differences between groups identified through one-way ANOVA testing followed by Tukey's Honestly Significant Difference (HSD) post hoc analysis (p < 0.05). An asterisk indicates a significant difference between control and inoculated plants, determined by a t-test (p < 0.05). E: Hierarchical Clustering Analysis was performed following Ward method of clustering. Indicated traits are measured in Aerial parts (AP), roots (R) or grains (G). Cpercent: Percentage of C. Npercent: percentage of N. CN: C/N ratio. 15NEpercent: percentage of ¹⁵N. Ppercent: percentage of P. Q15Nmg: quantity of 15N in organ per plant. NUE: Nitrogen Use Efficiency. GY: Grain Yield per plant. The color scale indicates if the trait is higher (orange) or lower (blue) in the indicated conditions with respect to the others. Details for the formulas used to calculate these traits are indicated in Supplementary Table 3.

Figure 2: Impact of N supply on expression of genes involved in N and P transport in roots suggests that AM symbiosis alleviates N deficiency stress. Plants were grown and harvested in 2020 as indicated in materials and methods under low nitrate (LN, 1mM NO_3^-), moderate nitrate (MN, 5mM NO_3^-) or high nitrate (HN, 10mM NO_3^-). Roots were harvested at VT+15 stage. A: Quantification of *R. irregularis* in roots by qRT-PCR amplification of the *RiLSU* constitutive fungal transcript normalized with maize actin transcripts. B: Normalized transcript levels with actin as a reference for indicated genes monitored by qRT-PCR. Ctrl: Mock non inoculated plants, Rhizo: inoculated with *R. irregularis*. Bar plots represent the means, accompanied by dots for individual observations overlaid on the bars, with each dot corresponding to a single observation in the dataset (N=5). Different letters indicate significant differences in gene expression levels identified through one-way ANOVA analysis followed by Tukey's honestly significant difference (HSD) test. The black bars represent standard errors. An asterisk indicates a statistically significant difference between mock and *R. irregularis*-inoculated plants, determined by a two-sample t-test (p < 0.05).

Figure 3 : RNAseq data showing the overlap between maize DEGs and functional GO enrichment categories. A: Genes were selected so that their fold change (inoculated versus control) was > to 1.5 and statistically significant in the 3 replicates. B73 LN and B73 HN represent DEG genes up or down regulated in inoculated plants compared to control plants in LN or HN respectively; LO3 LN and LO3 HN represent DEG genes up or down regulated n inoculated plants compared to control plants in LN or HN respectively. **B:** main GO functional categories in DEGs down in inoculated versus control (on the right with blue arrows) and DEG down (on the left with orange arrows) in inoculated versus control.

Figure 4: Metabolomic analyses of leaves and roots reveals key metabolites specific to symbiosis under
LN or HN nutrition. Plants were grown and harvested in 2020 as indicated in materials and methods under low
nitrate (LN, 1mM NO$_3^-$) or high nitrate (HN, 10mM NO$_3^-$). Leaves and roots were harvested at VT+15 stage. Ctrl:
Mock non inoculated control, Rhizo: inoculated with *R. irregularis,* Metabolites were monitored by GS/MS. **A** and

B : HCA heatmap showing metabolomic profiles of leaves and roots of inoculated or control plants under LN or
HN in leaves and roots respectively. **C** and **D**: Analyses of metabolomics data conducted with the R package
MultiVarSel with a threshold of 0.884 showing the metabolites on which the genotype, the N nutrition of the *R.*
irregularis inoculation have an impact. Leaves, **C;** roots, **D.** The color scale indicates if the metabolites are
accumulated (red) or down accumulated (blue) in the indicated conditions with respect to the others.
919

Figure 5: Genome based metabolic modeling reveals essential impact of AMF on pyrimidine metabolism. A: RNAseq data were used to predict different organ growth rates in the B73 and Lo3 maize lines. EXTREAM algorithm predicted higher yield under LN in B73 and Lo3. B : Venn diagrams showing the metabolic bottlenecks under indicated conditions. The debottlenecked reactions from the LN CTRL condition are those on the left yellow circle outside the intersection. LN CTRL (yellow circle). HN CTRL (blue circle). LN RHIZO (red circle). The list of the metabolic bottlenecks is on Tables 1 and 2. C: Metabolic reactions related to the pyrimidine pathway that are predicted to be debottlenecked by *R. irregularis* inoculation and by HN treatment shown in B. All the reactions shown here are in the plastid compartment. D: Quantification of pyrimidines in leaves of the same plants from the 2020 experiment, used for RNAseq and metabolomics. The nucleotides were extracted then quantified by LC–TOF according to materials and methods. N=3 pools of leaves harvested from 2 to 3 plants. CTRL Mock non inoculated, RHIZO: inoculated with *R. irregularis*. Different letters indicate significant differences in nucleotide levels identified through one-way ANOVA analysis followed by Tukey's honestly significant difference (HSD) test. The black bars represent standard errors. One or two asterisks indicates a statistically significant difference between mock and *R. irregularis*-inoculated plants, determined by a two-sample t-test p < 0.05 or p<0.01 respectively.

Figure 6 : N supply strongly impacts *Rhizophagus irregularis* transcriptome and pyrimidine biosynthetic genes in maize roots. Fungal DEGs were selected so that they have a fold change (FC >1.5). **A** : Up in LN indicates *R. irregularis* genes that are more highly expressed in inoculated LN roots compared to inoculated HN roots. **B**: Up in HN indicates *R. irregularis* genes that are more highly expressed in HN compared to LN. **C**: Gene ontology of functional categories enrichment was based on KOG annotations of encoded *R. irregularis* proteins (see materials and methods). Each bar represents the functional category enriched in the given condition. p-value of enrichment was < 0.05; only categories enriched more than 2 fold and displaying more than 2 genes are represented and used to build the heatmap. The color scale indicates the over-representation or the down-representation in red and blue respectively. **D**: read counts of *R. irregularis* transcripts related to pyrimidine de novo biosynthesis and salvage (from Supplementary Table 6).

Figure 7: Hypothetical working model for maize and *R. irregularis* N nutrition under LN and HN.
To build the model, the metabolomic data from B73 and Lo3 maize lines were pooled for simplification (Supplementary Figure 10) and we used genes in Supplementary Tables 6 and 8. In some cases genes with multigene families such as Glutamine Synthetase encoding genes have different

expression profiles. For simplification purposes, these cases were not considered in the hypothetical
mechanistic model below.
Genes are indicated in italics. Metabolites are in boxes. Surrounded by orange boxes: fungal
metabolites. Surrounded by green boxes: plant metabolites. Written in green or orange are plant or
fungal genes respectively. If no box or circle is indicated, this means that there are no clear data bout
the corresponding gene or metabolite. AA: Amino acids. NucB: Nucleobases. Pur: Purine. Pyr:
Pyrimidine. IRM: Intraradical mycelium. ECM: Extraradical mycelium. .Alla: Allantoic Acid. Alln :
Allantoin. Alln-ase: Allantoinase. Arg: arginine. Asn: Asparagine. Asp : Aspartate. b-Ala :b-alanine. Citr
: Citrulline. Glu: Glutamate. Gln: Glutamine. Orn : Ornithine. GOGAT: Glutamine Oxo-Glutarate
Aminotransferase. Oxa : Oxalic Acid. NR : Nitrate reductase. NiR : Nitrite reductase. AAT : Amino
acid transporter. AMT : Ammonium transporter. DUR : Urea transporter. NPF : plant Nitrate
transporter. NRT : *R. irregularis* nitrate transporter. CPS : Carbamoyl-phosphate synthase. ASL :
Argininosuccinate lyase. Pyr de novo: Pyrimidine de novo biosynthetic pathway. Pyr degradation:
Pyrimidine degradation. Pur de novo: Pur de novo biosynthetic pathway.
942

A

DEGs Up regulated in leaves

DEGs Down regulated in leaves

B73 HN   LO3 LN
B73 LN   LO3 HN
        786   1534
    52   49   107
        29   11
1892   325
    1136   2   49
        57   4
            42

DEGs Up regulated in roots

DEGs Down regulated in roots

B73 HN   LO3 LN
B73 LN   LO3 HN
        991   3119
    244   772   599
        196   315
1621   503
    263   59
        54   38   112
            56

B

A

Pyrimidine metabolism

Up in Lo3 LN vs HN

Up in B73 LN vs HN

793 | 548 | 699

Up in Lo3 HN vs LN

Up in B73 HN vs LN

987 | 609 | 750

Enriched GO categories

Nuclear structure
Intracellular trafficking, secretion, and vesicular transport
Translation, ribosomal structure and biogenesis
Transcription
Cytoskeleton
RNA processing and modification
Defense mechanisms
Function unknown
Amino acid transport and metabolism
Posttranslational modification, protein turnover, chaperones
Chromatin structure and dynamics
Carbohydrate transport and metabolism
General function prediction only
Inorganic ion transport and metabolism
Cell cycle control, cell division, chromosome partitioning
Replication, recombination and repair
Lipid transport and metabolism
Energy production and conversion
Nucleotide transport and metabolism
Coenzyme transport and metabolism
Cell wall/membrane/envelope biogenesis
Signal transduction mechanisms
Secondary metabolites biosynthesis, transport and catabolism

LO3 | B73 | LO3 | B73

Up in LN | Up in HN

2 | 1 | 0 | -1 | -2

D

- LO3_LN
- LO3_HN
- B73_LN
- B73_HN

Urate oxidase
Urease
Uracil phosphoribosyltransferase
Orotate phosphoribosyltransferase
Cytosine deaminase FCY1
Deoxycytidylate deaminase
Allantoicase
Uridine 5'- monophosphate synthase
Uridine 5'- monophosphate synthase
dUTPase

0.00 | 200.00 | 400.00

read counts