1

Cortical circuits for cross-modal generalization

Maëlle Guyoton[1]*, Giulio Matteucci[1]*, Charlie G. Foucher[1], Matthew P. Getz[2],

Julijana Gjorgjieva[2] & Sami El-Boustani[1]

[1] Department of Basic Neurosciences, Faculty of Medicine, University of Geneva, 1

Rue Michel-Servet, 1206 Geneva, Switzerland.

[2] School of Life Sciences, Technical University of Munich, 85354 Freising, Germany.

* These authors contributed equally to this work.

Summary

Adapting goal-directed behaviors to changing sensory conditions is a fundamental aspect of intelligence. The brain uses abstract representations of the environment to generalize learned associations across sensory modalities. The circuit organization that mediates such cross-modal generalizations remains, however, unknown. Here, we demonstrate that mice can bidirectionally generalize sensorimotor task rules between touch and vision by using abstract representations of peri-personal space within the cortex. Using large-scale mapping in the dorsal cortex at single-cell resolution, we discovered multimodal neurons with congruent spatial representations within multiple associative areas of the dorsal and ventral stream. Optogenetic sensory substitution and systematic silencing of these associative areas revealed that a single area in the dorsal stream is necessary and sufficient for cross-modal generalization. Our results identify and comprehensively describe a cortical circuit organization that underlies an essential cognitive function, providing a structural and functional basis for abstract reasoning in the mammalian brain.

Introduction

Objects possess unique physical properties that are detected by different sensory organs. The brain seamlessly integrates these sensory inputs to create unified percepts and abstract representations of the environment, which are essential for generalizing behaviors to unfamiliar situations[1]. This phenomenon is especially pronounced in the peri-personal space, where visual and tactile systems converge[2]. Indeed, an object's position and identity, initially discerned through touch in darkness, can immediately be recognized by sight in light. Cross-modal generalization —also called cross-modal transfer learning— describes the process by which recognition in one sensory modality enables the generalization of learned associations to others, a capability observed across diverse species[2–5], suggesting a foundational circuit organization yet to be elucidated. While neuronal correlates of increasingly abstract representations have been reported[6–9], the cortical architecture enabling the use of such representations for generalized learning across sensory modalities is not yet understood. Mice have proven to be a valuable model for dissecting circuits responsible for multisensory integration and their role in goal-directed behavior[10–13]. Mice rely on visuo-tactile inputs for behaviors such as gap crossing[14], navigation[15], and object recognition[6]. Because whiskers occupy a substantial portion of the visual field, both somatosensory and visual systems frequently receive correlated inputs (Fig. 1a). In the superior colliculus, multimodal representations of whisker and visual information possess topographically co-aligned functional maps[16,17], potentially enhancing reflexive behaviors like gaze and head orientation by increasing the salience of spatially congruent multisensory events[18]. Cortical circuits, critical for conscious perception and goal-directed behaviors, also display multimodal visuo-tactile responses. In particular, associative areas within the posterior parietal cortex

receive inputs from both the primary visual cortex (V1) and the primary somatosensory

cortex (S1)[19]. However, the precise nature of the functional and anatomical

organization of cortical areas dedicated to visuo-tactile processing, as well as their

potential role in cross-modal generalization of previously learned sensorimotor

associations, remains unknown. Here we investigate these circuits in mice performing

goal-directed behaviors.

Results

Cross-modal generalization in mice

We designed a behavioral paradigm to test the ability of mice to generalize

sensorimotor associations learned through whisker sensations to the visual modality.

Given that both sensory modalities share a common spatial organization within the

peri-personal space (Fig. 1a), the task includes visual and tactile stimuli that originate

from locations that are spatially congruent (Fig. 1b). In the dark, head-fixed and water-

restricted mice were first trained on a Go/No go tactile discrimination task, where they

had to discriminate between two whiskers vertically arranged in the same column of

the whisker pad. Mice were rewarded with a drop of water if they licked a spout upon

stimulation of the top whisker (B2) whereas they were punished with a 10-second-long

timeout if they licked for the bottom whisker (C2). Once mice became expert at the

task and performed stably with a high percentage of correct trials over at least 3

consecutive sessions, we switched the task to a Go/No go visual task. In this condition,

we replaced the top and bottom whisker stimulations with "top" and "bottom" visual

stimuli. These stimuli consisted of black squares on a gray background drifting along

the same rostro-caudal direction than the whisker deflections. The screen was

oriented to be centered and parallel to the right retina on the same side as the

previously stimulated whiskers. The location of the moving squares along the vertical

axis was chosen to roughly match the location of the whiskers within the visual field of

the animals (see Methods).

Fig. 1. Cross-modal generalization in mice.

a, Illustration of the common organization of peri-personal space for visual and whisker tactile
inputs in mice. **b,** Schematic of the behavioral Go/No go paradigm for studying cross-modal
generalization of spatial information in mice. **c,** Two types of tactile-to-visual modality switch:
"rule-preserving", wherein the spatial location of rewarded stimuli is preserved, and "rule-
reversing", wherein the location of rewarded stimuli is reversed. **d,** Left: example of a session
with the tactile task the day before modality switch. Conditional lick probabilities over trials are
shown for the top whisker (blue), the bottom whisker (red) and in absence of stimuli (pink).
Task performance computed as the percentage of correct discrimination trials (see Methods)
is shown in green with chance level as gray dashed line. Traces were computed on a sliding
window of 60 trials. Right: same as left for the first visual session following a rule-preserving
modality switch. **e,** Left: task performance and conditional lick probabilities averaged across
mice (N=5 mice) over consecutive sessions before and after a rule-preserving switch (vertical
dashed line). Shaded area: S.E.M. Color code as in **d**. Right: detection (purple) and
discrimination (green) performance distribution for the session before and after the switch
(two-sided paired t-test comparing days, *p<0.05, **p<0.01, ***p<0.001, N.S. Not significant).

To test whether mice can use previous associations learned during the tactile task to infer the reward contingency (i.e. the rule that determines when a reward will be delivered following a sensorimotor response) in a new visual task, we considered two scenarios: a "rule-preserving" and a "rule-reversing" modality switch. The cohort of mice undergoing the rule-preserving modality switch could obtain reward by licking for stimuli presented at the same spatial location after the switch. For mice undergoing the rule-reversing switch, the reward contingency was spatially reversed following the switch to the new modality (Fig. 1c). Following rule-preserving modality switches, we observed rapid generalization of the learned association to the new modality. Mice seamlessly performed the task with a comparable level of high performance as observed in the session preceding the switch (Fig. 1d,e). In contrast, task performance was strongly affected following a rule-reversing modality switch, consistently falling at chance level. Mice typically attempted to lick in response to both stimuli during the early part of the session before disengaging and ceasing any licking behavior (Fig. 1f). In some cases, mice attempted to lick first for visual stimuli spatially congruent with the previously rewarded whisker, leading performance to briefly drop below chance level. Following rule-reversing switches, mice displayed a strong resistance to engaging with the task for several consecutive sessions (Fig. 1g). Such behavior persisted despite strong thirst-driven motivation, as indicated by the prolonged licking bouts observed when water drops were manually delivered to maintain task engagement[20]. This implies that the mouse's failure to perform after a rule-reversing

modality switch stems from conflicting prior knowledge of the task rule, rather than

from a lack of motivational drive. We verified that this result was not caused by a

preference for the top whisker or for the top visual stimulus by repeating the same

experiments with cohorts of mice trained to lick for bottom whisker stimulations. Mice

maintained high task performance after switches that preserved the spatial rule,

regardless of whether they were initially trained on one whisker or another, but

performance dropped when the rule was reversed (Extended Data Fig. 1a-c). We

further confirmed that mice could also generalize the task rule from the visual to the

tactile modality, demonstrating bidirectional cross-modal generalization of learned

associations (Extended Data Fig. 1g-l).

Despite the limited visibility of the capillary glass tubes used for whisker

stimulations in the dark, we tested whether mice could potentially rely on tubes moving

as visual cues to perform the tactile task, thereby generalizing within the visual domain

instead of across sensory modalities. We carried out control experiments where mice

proficient in the tactile task underwent sessions with whiskers temporarily removed

from the tubes, and subsequently reintroduced. We observed that both detection and

discrimination performance dropped to chance level immediately after the whiskers

were removed from the capillary tubes, but recovered to expert levels once the

whiskers were reinserted into the piezo stimulators (Extended Data Fig. 2). This

demonstrates that mice were not using visual cues to perform the whisker

discrimination task in the dark.

Besides generalizing task rule through common spatial organization, mice

could also potentially generalize the abstract Go/No go structure common to the two

tasks (i.e. "acting upon one stimulation, the Go stimulus, leads to reward while acting

upon the other, the No go stimulus, leads to a timeout") to rapidly increase

performance after the switch. To investigate this possibility, we trained mice on an

auditory Go/No go discrimination task, which produced similar levels of motivation

without prior knowledge of spatial rules related to reward contingency. Mice were then

switched to the spatial visual task (Extended Data Fig. 1d). The stimuli used for the

auditory task were two pure tones played from the same speaker, bearing no clear

spatial relationship to the visual stimuli introduced after the modality switch. In absence

of spatial prior, performance dropped to chance level after the switch but steadily

recovered to expert level over the next few sessions (Extended Data Fig. 1e,f). We

concluded that mice could learn the task after modality switch significantly faster in

absence of the conflicting spatial prior observed for mice experiencing rule-reversing

modality switches (Fig. 1h).

Finally, we compared our results for rule-reversing modality switches with

conditions in which reward contingencies were switched within the tactile modality, as

previously studied[21,22]. After mice reached expert level in the tactile task, we reversed

the reward contingencies between the two whiskers. This led to a stark drop in

discrimination performance below chance level, indicating that mice rigidly persisted

in performing the task following the original rule (Extended Data Fig. 3). Following the

switch, performance increased but remained below or at chance level for at least three

consecutive sessions. Thus, mice behave differently when reward contingencies are

switched within the same sensory modality as they continue to inflexibly produce the

same sensorimotor transformation, likely reflecting ingrained habitual behaviors.

Co-aligned visuo-tactile spatial maps in the dorsal cortex

Our behavioral results suggest that mice can generalize previously learned

sensorimotor associations across sensory modalities (Extended Data Fig. 4), based

on a common representation of the peri-personal space. Cortical circuits are

necessary for conscious perception and are believed to mediate flexible goal-directed

behaviors such as cross-modal generalization[2,23]. To pinpoint the cortical regions

responsible for visuo-tactile generalization, we mapped the topographic

representation of vertical space for both modalities in the dorsal cortex of transgenic

mice expressing the calcium indicator GCaMP6f in cortical layers 2/3 (Fig. 2a, see

Methods). We first used established retinotopic and somatotopic mapping

protocols[20,24] to identify whisker-responding and retinotopically organized cortical

areas through a 5 mm diameter cranial window over the posterior part of the left dorsal

cortex (Extended Data Fig. 5a). Whisker response patterns and retinotopic sign maps

enabled us to precisely fit a projection of the Allen Mouse Brain Atlas to the cranial

window of each mouse (see Methods). We used this atlas to register all functional

maps into a common reference frame (Extended Data Fig. 5b-f).

Fig. 2. Co-aligned visual and tactile functional maps in the dorsal cortex.

195   **a,** Top: schematic of the visuo-tactile sparse noise protocol used during wide-field imaging of
196   GCaMP6f through a cranial window. Bottom: matrix of all combinations of visual and whisker
197   stimuli (green: multisensory, orange: visual, magenta: tactile). **b,** Somatotopic map of vertical

space computed from whisker stimuli averaged across mice (N=29 mice for all panels of the figure), with transparency defined by response significance in each pixel (see Methods). A projection of the Allen Mouse Brain Atlas is overlaid on top with areas names and orientation. **c,** Average retinotopic map of vertical space. **d,** Average modality preference map between visual and tactile responses. **e,** Average spatial coherence map between visual and tactile representations (see Methods). **f,** Average multisensory modulation map comparing visuo-tactile responses and combination of unisensory responses (see Methods). **g,** Multisensory modulation index for pixels belonging in regions of high spatial coherence compared to regions without spatial coherence (n=26,286 pixels versus n=28,394 pixels, unpaired two-sided Wilcoxon test, ***$p<10^{-300}$). **h,** Multisensory modulation index for pixels belonging to unimodal or multimodal regions (n=10,264 pixels versus n=15,921 pixels, unpaired two-sided t-test, ***$p<10^{-116}$).

To characterize the cortical representation of the vertical space for both unisensory and multisensory stimulations, we designed a visuo-tactile sparse noise protocol (Fig. 2a, see Methods). This protocol was used in task-naïve mice to obtain retinotopic and somatotopic maps for vertical space by computing the preferred spatial position for each pixel. In response to whisker stimuli, we identified the well-established somatotopic arrangement of the primary and secondary whisker somatosensory cortices, S1 and S2, which exhibited a topographic inversion at their boundary (Fig. 2b). Importantly, we observed that whisker stimulations also evoked organized somatotopic maps in several visually responsive areas including the anterior (A), rostro-lateral (RL), antero-lateral (AL) and latero-intermediate (LI) areas. This suggested that whisker representations might be present in a more extended cortical network than previously reported[25]. Strikingly, the maps obtained with visual stimuli displayed a very similar organization to the tactile ones in these associative areas as well as in S1 and S2 (Fig. 2c). The extended spatial representations evoked by visual or tactile stimuli were found consistently across mice (Extended Data Fig. 5g). This suggests that spatially localized stimuli, regardless of their visual or tactile nature, might share a common topographic representation that facilitates mapping between sensory modalities, as previously observed in the superior colliculus[16,17].

Spatial representations evoked by these two modalities displayed an angular offset

that we estimated around 30 degrees by comparing the angle difference between

gradient vectors obtained from these maps (see Methods). This might reflect the

mouse's internal model of how whisker sensations align with their corresponding

locations in the visual field[13].

We further investigated the functional properties of these representations by

first computing a modality preference index to assess what sensory modality

dominates each area (Fig. 2d). As expected, S1 and S2 were dominated by tactile

inputs whereas V1 was dominated by visual inputs. RL and the region at the border

between AL and LI displayed a more balanced preference for both modalities. In

addition, we measured the spatial coherence between maps of vertical retinotopy and

somatotopy indicating local co-alignment (Fig. 2e, see Methods). This confirmed a

widespread spatial co-alignment across most associative areas in the belt between

V1 and S1. We further computed a multisensory modulation index, comparing

multisensory responses triggered by visuo-tactile stimuli to the maximal unisensory

response on a pixel-by-pixel basis (see Methods). The resulting map revealed a strong

multisensory enhancement in visuo-tactile associative areas and in S1 (Fig. 2f).

Multisensory responses were comparable under visuo-tactile conditions where

whisker stimuli were synchronous or delayed by 0.15 seconds to ensure simultaneity

of evoked responses in the cortex (Extended Data Fig. 6), with a tendency to show

stronger enhancement in the latter case, as previously documented[19]. Moreover, we

found that multisensory enhancement was more pronounced in regions with higher

coherence between spatial maps and with strong bimodal representations (Fig. 2g,h).

Anatomical origin of spatial maps in the dorsal cortex

Functional maps measured with wide-field calcium imaging could result from direct inputs from visual and tactile primary cortical areas, from evoked top-down inputs[26], or even be the result of highly stereotypical uninstructed movements evoked by sensory stimuli[27]. To investigate the synaptic origin of these maps, we performed anatomical experiments to map feedforward and feedback projections between primary sensory areas and associative areas displaying visuo-tactile representations (Fig. 3a). We obtained visual and tactile functional maps for representation of vertical space in wild-type mice using intrinsic optical signal imaging under low isoflurane anesthesia (Extended Data Fig. 7a,b,d,e, see Methods). These maps were then used to identify two cortical locations representing distinct iso-horizontal vertical positions in V1 or to target B2 and C2 barrels in S1. We then opened the cranial window and injected two adeno-associated viral vectors to induce expression of tracer proteins GFP and tdTomato in the respective locations (Extended Data Fig. 7c,f). After 10-15 days, transcardial perfusions were performed and brains were extracted, flattened, and sliced (see Methods). Enriched M2 subtype muscarinic acetylcholine receptors (M2 AChR) in V1 and S1 barrel field were used as landmarks to fit the Allen Mouse Brain Atlas on the reconstructed stack, confirming the location of injection sites along the vertical representation of V1 and S1 (Fig. 3b,c). Axonal projections from primary sensory areas were found in associative cortical regions where visuo-tactile responses were measured with the same spatial organization. Variability in injection sites allowed us to create an anatomical map that described axonal preferences for top and bottom location (Fig. 3b,c), aligning closely with wide-field imaging results (Fig. 2b,c). This

confirmed that the functional maps are inherited, at least in part, from direct

feedforward projections from primary cortical areas.

Fig. 3. Co-aligned visual and tactile anatomical projection maps in the dorsal cortex.
a, Schematic of the data collection and analysis pipeline for topographic anatomical tracing.
b, Anterograde labeling of S1 projections using viral vectors injections. Left: injection sites in
B2 and C2 barrels in whisker S1. Right: conserved somatotopic organization of projections in
associative areas for the cortex shown in the left (top). Bottom: Reconstructed whisker
preference averaged across mice using all injection sites depicted by circles (see Methods,
N=9 mice and n=18 injections, Pearson coefficient of correlation between anatomical and
wide-field map: 0.8311, $p<10^{-300}$). **c,** Same as **b** but for anterograde labeling of V1 projections
with two injection sites along the iso-horizontal axis (N=7 mice and n=14 injections, Pearson
coefficient of correlation: 0.5121, $p<10^{-300}$). **d**, Retrograde labeling of S1-projecting neurons
using CTB-555 and CTB-647. Top: Examples of CTB-labelled neurons spatially organized in
associative areas. Bottom: Reconstructed map of preferred whisker in projecting neurons over
the dorsal cortex combining all injection sites (N=5 mice and n=8 injections, Pearson
coefficient of correlation: 0.663, $p<10^{-300}$).

While visual stimuli could evoke organized responses in S1 (Fig. 2c), no direct

projections were found between V1 and S1[28] suggesting the existence of spatially

organized feedback projections from associative areas to S1. Previous work has

shown that feedback projections from higher visual areas (including A, RL, AL and LI)

to V1 are spatially organized along the vertical dimension[29] but it is not clear if this

holds true for feedback projections to S1. Feedback projections from associative areas

to S1 were characterized using the same strategy, but with injections of Cholera Toxin

Subunit B (CTB) conjugated either with Alexa555 or Alexa 647 (Fig. 3d). S1-projecting

neurons for top and bottom locations aligned with wide-field imaging (Fig. 2b) and

anterograde tracing (Fig. 3b). We additionally confirmed that the same was true for V1

(Extended Data Fig. 7g) as previously reported[29]. Thus, we observed a shared spatial

organization of feedforward and feedback projections between primary and

associative areas.

Single-neuron visuo-tactile functional properties

Spatially organized feedforward and feedback projections could facilitate

generalized sensorimotor learning through transfer of spatial information across

sensory modalities. However, functional maps obtained with wide-field imaging do not

reveal precise computations performed at single-cell level and could still be prone to

artifacts produced by neuronal processes originating from other brain structures. To

overcome this limitation and assess if neurons in associative areas can mediate cross-

modal generalization, we performed two-photon calcium imaging in a subset of mice

(Fig. 4a). Single neuron somatic GCaMP6f signal was extracted during the visuo-

tactile sparse noise protocol in fields-of-view covering different cortical areas identified

with the atlas (Extended Data Fig. 8a-f). Given the observed fluorescence response

patterns to various unimodal and multimodal stimuli (Fig. 4b), we could map specific

functional properties of single neurons onto the reference atlas, pooling data across

mice, for comparison with corresponding wide-field regions. Many recordings were

performed across a large portion of the dorsal cortex to extensively cover responsive

visuo-tactile areas (Extended Data Fig. 8g-h). We reconstructed the somatotopic map

of vertical space across the cranial window, which closely aligned with the wide-field

map (Fig. 4c). Neurons with whisker tactile responses were found across the belt of

associative areas following the somatotopic organization. This further confirmed the

existence of an extended network of whisker responsive and visuo-tactile cortical

regions[25]. The same was true for neurons responding to visual stimuli which were

found across most visual and tactile areas, even extending to S1 and S2 (Fig. 4d).

Reconstructed population maps based on single-neuron properties consistently

matched with functional maps measured with wide-field calcium imaging (Extended

Data Fig. 9).

Fig. 4. Visuo-tactile representation of peri-personal space in single neurons.

a, Example cranial window and two-photon field-of-view overlapping with the medial part of

area RL. **b,** Response pattern of GCaMP6f fluorescence for the example neuron highlighted

with yellow circle in **a**. Unisensory responses (z-scored) are shown on the side for visual

(orange) or whisker (magenta) stimuli. Predicted multisensory responses are shown for visuo-

tactile stimuli (black) together with measured responses (green). Error bars: S.E.M. **c,**

Population of single neurons imaged with two-photon microscopy to measure significant

responses to whisker stimuli (n=567 neurons). Each spot represents a single neuron, and the

color code indicates preference for top (blue) or bottom (red) stimuli. The reconstructed wide-

Among all neurons imaged, we identified four functional cell types. Unisensory visual or tactile neurons only responded to their respective modality, gated neurons responded only when both visual and tactile stimuli were presented together, and bimodal visuo-tactile neurons responded to both unisensory visual and whisker stimuli (Extended Data Fig. 10a-b). Importantly, bimodal visuo-tactile neurons localized into two distinct clusters (Fig. 4e), corresponding to the multimodal domains identified by wide-field imaging (Fig. 2d), and consistent with anatomical projection patterns (Fig. 3b,c). As these clusters coincided with areas associated with the dorsal (A, RL) and ventral (AL, LI) streams, we used these designations moving forward. Due to their distinct functional properties, visuo-tactile neurons may facilitate cross-modal generalization. A Bayesian decoder trained to discriminate whisker stimuli based on individual neuronal responses was evaluated for generalization to visual stimuli (see

Methods). Visuo-tactile neurons in associative areas uniquely enabled effective

generalization across sensory modalities (see Fig. 4f). Other neuron types

demonstrated significantly lower decoding accuracy when we tested their ability to

generalize the tactile discrimination to the visual modality, particularly with larger

populations (see Extended Data Fig. 10c,d).

We further characterized the response properties of single neurons for visuo-

tactile stimuli in comparison with their responses predicted from unisensory responses

(Fig. 4b, see Methods). We found that bimodal neurons were typically tuned to

spatially congruent stimuli across both modalities. The example neuron in Figure 4b

responded preferentially to the bottom part of the visual field and to the bottom whisker.

Using unisensory responses, we predicted the response pattern to visuo-tactile stimuli

as the maximum response between the two modalities for each combination. When

comparing the predicted response with the measured one, we observed suppression

in incongruent combinations and enhancement in congruent ones. This nonlinear

modulation profile enables neurons to maintain their spatial selectivity independently

of the stimulated modality (visual, tactile, or visuo-tactile). This response property is

reminiscent of the supramodal encoding of object orientation reported in visuo-tactile

neurons of the rat posterior parietal cortex[6]. Population analysis confirmed that spatial

congruence in multisensory selectivity is prevalent among populations of multimodal

neurons in both the ventral and dorsal stream (Fig. 4g). Additionally, the two neuronal

populations exhibited sharper tuning for whisker and visual positions than that

predicted by their unisensory responses alone (Fig. 4h). Neurons from the ventral

domain displayed sharper tuning for particular visual locations, whereas neurons in

the dorsal domain exhibited greater tuning to individual whiskers (Fig. 4i,j), consistent

with modality preferences observed in wide-field imaging data (Fig. 2d). These results

showed that bimodal neurons in visuo-tactile associative areas are indeed specifically

responding to spatially congruent visuo-tactile stimuli and could mediate cross-modal

generalization during goal-directed tasks.

Cross-modal generalization loss-of-function

Cross-modal decodability of stimulus location observed at single-cell level (Fig.

4f), alongside the anatomical connectivity between visual and whisker somatosensory

cortices (Fig. 3), suggest that associative visuo-tactile cortical areas could mediate

cross-modal generalization of goal-directed behaviors. We performed loss-of-function

experiments to assess the necessity of these areas for visuo-tactile generalization.

Given that task-naïve mice (those not yet exposed to any behavioral tasks) exhibited

co-aligned and anatomically connected spatial maps, we reasoned that the

reverberation of evoked responses across this extensive visuo-tactile network could

potentially drive learning processes beyond the initially stimulated modality. This could

facilitate cross-modal generalization before any direct exposure to the second task

with the other modality. To prevent this possibility, we decided to chronically silence

associative cortical areas prior to any sensorimotor learning. We virally expressed the

tetanus toxin light chain (TeNT-P2A-GFP) in ventral and dorsal areas, thereby

preventing neurotransmitter vesicle release in transfected neurons (Fig. 5a, see

Methods). Neurons expressing TeNT also co-expressed GFP, allowing comparison of

the expression pattern with the fitted atlas (Fig. 5b). After subtracting blood vessel

patterns and comparing with the cranial window before injection (see Methods), we

characterized the extent of GFP expression and overlap with different visuo-tactile

areas (Extended Data Fig. 11a-d). To ensure strong expression of TeNT to effectively

suppress vesicle release in the transfected neurons before beginning the behavioral

training, we waited at least four weeks after viral injections[30]. Mice normally learned

the whisker discrimination task and reached expert performances comparable to those

observed in mice trained without TeNT expression (Fig. 5c, Extended Data Fig. 11a-

d). However, when switching to the visual task in the rule-preserving condition,

performance typically dropped to chance level (Fig. 5d,k).

To identify which visuo-tactile areas are necessary for cross-modal

generalization, we expressed TeNT-P2A-GFP in different cortical locations across

mice (see Methods). We were able to cover all visuo-tactile areas with varying degrees

of overlap with the expression of TeNT (Extended Data Fig. 11e). Taking advantage

of this variability, we calculated for each cortical area the extent to which the overlap

of its surface with TeNT expression correlated with any decline in performance

following a rule-preserving switch (Extended Data Fig. 11f). This analysis revealed

that generalization impairment was significantly correlated with TeNT expression in

area RL only (Fig. 5e). To confirm this result, we also performed a reverse correlation

analysis mapping the average TeNT coverage that evoked a complete impairment of

cross-modal generalization (see Methods). Here again, RL silencing was found to

consistently prevent cross-modal generalization, revealing that this associative area

is necessary for the transfer to occur (Fig. 5f). Grouping mice depending on whether

TeNT expression pattern was located in the dorsal or ventral stream (Fig. 5g,i), we

confirmed that dorsal stream silencing severely affected generalization performance

(Fig. 5h), leading to slower re-learning after rule-preserving switches.

Fig. 5. Area RL is necessary for cross-modal generalization.

a, Schematic of the experimental timeline for loss-of-function experiments. **b,** Cranial window with expression of TeNT-P2A-GFP where viral vectors were injected. Atlas overlaid for reference. **c,** Example session before a rule-preserving modality switch from a tactile to a visual task. Color code as in Fig. 1d. **d,** Session following the modality switch. **e,** Area-based correlation between TeNT-P2A-GFP expression overlap and performance drop following modality switch. Color map indicates Pearson coefficients of correlation ρ. Areas with p<0.05 are indicated with a thick border (RL with p=0.047). **f,** Average TeNT-P2A-GFP coverage for mice with impaired cross-modal generalization (see Methods). The map is displayed after subtraction of the average coverage across all injected mice. **g,** Average TeNT-P2A-GFP coverage of all mice where only dorsal neurons were silenced (N=8 mice). Dots indicate center-of-mass location for each mouse. **h,** Left: Average task performance and conditional lick probabilities across sessions for mice in **g** with rule-preserving modality switch (vertical dashed line). Shaded area: S.E.M. Color code as in **c**. Right: Detection (purple) and discrimination (green) performance distribution the day before and after the switch (paired two-sided t-test comparing days, *p<0.05, **p<0.01, ***p<0.001, N.S. Not significant).

In contrast, silencing areas in the ventral stream did not prevent most mice from performing above chance level following a rule-preserving switch (Fig. 5j). Although mice expressing TeNT in RL initially failed to detect or discriminate visual stimuli after task switch, they regained performance in the visual task after several training sessions (Fig. 5h). This indicates that despite a comparable capacity to perform the visual or tactile task, these mice were not able to generalize learning across sensory modalities. This suggests a very selective effect of RL silencing in impairing cross-modal generalization abilities but not unisensory skills.

We further assessed how silencing area RL with TeNT affects task performance following rule-reversing modality switch (Extended Data Fig. 12a-b). Following the switch, their performance fell to chance level but recovered faster than what we observed for control mice (Extended Data Fig. 12c). To fully characterize the learning rate following modality switches in TeNT-expressing mice, we focused on all mice that exhibited complete generalization impairment (i.e. displaying performance near chance level on the first session following the switch regardless of whether the condition was rule-reversing or rule-preserving). For these mice, we quantified the re-learning rate over several sessions following the modality switch and observed that the distribution was comparable to that seen during non-spatial auditory-to-visual

switches (Fig. 5l). This implies that silencing area RL prevents mice from applying prior

knowledge about the spatial task rule, whether it is conflicting or congruent with the

new task rule, resulting in the mice learning the visual task without prior spatial

knowledge.

Optogenetic sensory substitution for cross-modal generalization

To determine whether cortical area RL is not only necessary, but also sufficient

for visuo-tactile generalization, we conducted gain-of-function experiments by

substituting visual stimuli after modality switch with direct optogenetic stimulations of

RL "top" and "bottom" encoding subregions (Fig. 6a). Projections from primary sensory

areas to the ventral and dorsal visuo-tactile stream being largely distinct (Extended

Data Fig. 7h-j), we assumed that optogenetic activations of RL would remain restricted

mostly to the dorsal stream. Functional mapping was first performed to identify

domains of RL encoding top and bottom visual stimuli in Ai32 transgenic mice

expressing Cre-dependent ChR2-eYFP. Viral vectors AAV1.CaMKIIa.Cre were then

injected in this area to broadly express ChR2-eYFP in RL (Fig. 6b). We employed a

projector-based microscope to precisely shape blue light patterns to selectively excite

subregions of RL encoding top and bottom stimuli. This approach allowed us to

substitute visual stimuli in the upper or lower visual field with precise spatio-temporal

optostimulations following the modality switch (see Methods).

Fig. 6. Area RL is sufficient for cross-modal generalization.

a, Schematic of the behavioral experiment with direct ChR2-mediated optostimulations of RL subregions following modality switch. **b,** Example cranial window of an Ai32 mouse expressing ChR2-eYFP after injection of AAV1.CaMKIIa.Cre viral vector in area RL. Atlas overlaid for reference. Subregions of RL encoding top or bottom stimuli are indicated in blue and red, respectively. Blue light patterns are shaped to match these subregions (see Methods). **c,** Left: Average task performance and conditional lick probabilities across sessions for mice experiencing a rule-preserving modality switch from the whisker task to the optogenetic task (N=7 mice). Right: detection (purple) and discrimination (green) performance distribution the session before and after the switch (paired two-sided t-test comparing days, *p<0.05, **p<0.01, ***p<0.001, N.S. Not significant). Performances are also tested against chance level (two-sided t-test, *p<0.05, **p<0.01, ***p<0.001, Blank: Not significant). **d,** Same as **c** but for mice undergoing a rule-reversing switch (N=6 mice, paired two-sided t-test comparing days, *p<0.05, **p<0.01, ***p<0.001, N.S. Not significant).

After mice began responding to optogenetic blue light stimulations, we assessed their capacity to transfer previously learned sensorimotor associations from the whisker task to the optogenetic task, both when the spatial rule was preserved or reversed. Following rule-preserving modality switch, mice maintain a comparable level of discrimination performance (Fig. 6c). In contrast, mice undergoing the rule-reversing modality switch performed at chance level after the switch and gradually relearned the task in the following sessions (Fig. 6d). This suggests that activity in RL alone is sufficient to induce cross-modal generalization.

Network model for generalized sensorimotor learning

Based on our experimental findings on the functional and anatomical organization of visuo-tactile cortical circuits, as well as the effect of RL silencing on cross-modal generalization, we developed a network model to recapitulate the behavior we observed in mice. Our model included areas V1, S1, and RL, each comprising two recurrent networks of excitatory and inhibitory neurons encoding "top" and "bottom" spatial location, respectively. Each of these areas projected to reward-modulated decision variables representing a downstream motor area (Fig. 7a). Training the model on the whisker discrimination task involved external inputs to S1 (representing thalamic inputs) and reward-modulated learning via Hebbian plasticity of the sensorimotor connections (see Methods). Model simulations showed that after switching inputs to V1, mimicking a rule-preserving switch to the visual modality, performance remained stable (Fig. 7b). However, if the spatial rule of the task was reversed upon modality switch, performance dropped below chance level and then slowly recovered (Fig. 7c, Extended Data Fig. 13). This delay in performance recovery was due to the model weights needing to override the previously learned structure before learning the new association. Mimicking the loss-of-function experiments, we silenced the area RL in the model during training, causing the neural network to learn the visual task from scratch after the rule-preserving switch without any spatial prior in the weight structure (Fig. 7d). The absence of direct connections between the modeled sensory areas prevented ongoing learning in the non-stimulated area. The model suggests that the activity reverberation mediated by bidirectional connections induces a consistent weight structure across connected areas. These weights, however,

needed to be reorganized following a rule-reversing modality switch, leading to

delayed relearning time (Extended Data Fig. 13c-d).

Fig.7. Neural network architecture for cross-modal generalization.

a, Schematic of the neural network model for cross-modal generalization. Synapses projecting
to the decision-computing area undergo synaptic plasticity during sensorimotor learning. **b,**
Action probability conditional on inputs to S1 before and inputs to V1 after, a rule-preserving
modality switch (vertical dashed line). Red: rewarded bottom stimulations, blue: non-rewarded
top stimulations. Discrimination performance is shown in green. Shaded areas: standard
deviation with n=50 simulations. **c,** Same as **b** after a rule-reversing modality switch. **d,** Same
as **b** with silenced RL. **e,** Number of steps necessary to cross a performance threshold of 75%
in the full network model (left) or in the model with silenced RL (right) after rule-preserving
("Preser.") or rule-reversing ("Rever.") modality switches (n=50 simulations in all except
"Normal Rever." for which n=49). Error bars indicate median, first and third quartiles. **f,** Left:
Number of sessions needed to cross a performance threshold of 65% following the modality
switch. Training was stopped after 12 sessions (dashed line) and mice that did not reach the
criteria before this session are plotted on the dashed line. Mice numbers are indicated for each
group at the bottom of the bar. Error bars: S.E.M. Right: Surprise matrix computed from
pairwise unpaired two-sided t-test between conditions. A hierarchical clustering based on
cosine similarity was used to group conditions based on surprise values.

To quantify the effect of different task conditions in the learning trajectory

following modality switch in the model, we computed the time needed in simulation

steps to recover expert level. Following a rule-preserving modality switch, the time to

relearn was near-instantaneous whereas it was much longer following rule-reversing

switches (Fig. 7e). Interestingly, when RL was silenced, the time to relearn was in-

between these two distributions reflecting that the model learned the task from scratch

(Fig. 7e). We compared these results to the time needed to relearn the task after

modality switch in all conditions used with our mice (Fig. 7f). In line with the model,

mice rapidly recovered high-performance levels after a rule-preserving modality switch.

In contrast, time to relearn the task was dramatically prolonged following rule-

reversing switches. This was true for both visual stimuli and optostimulations following

the switch. In TeNT-expressing mice, however, the time to relearn the task was

moderate and similar for both rule-preserving and rule-reversing modality switches.

Importantly, the time to relearn in these mice was comparable to the time needed to

reach expert level in the auditory task in absence of any spatial prior, reminiscent of

the "RL silenced" case in the model. The distribution of peak performance observed

for our mice was mostly determined by the nature of the sensory inputs after the switch

rather than the switch type (Extended Data Fig. 14).

Discussion

Our results elucidate the cortical architecture underlying the abstract

representation of peri-personal space. We identified the specific cortical circuit that

enables rapid generalization of sensorimotor associations learned from one modality

to another, forming the neural substrate for a key component of flexible behavior.

Cross-modal generalization has been reported across species including apes[3],

rodents[2] and, more recently, bumblebees[4]. Early research into the mammalian brain

circuits involved in cross-modal generalization underscores the necessity of different

brain structures to process different types of information such as spatial, temporal, and

object-related features[23,31,32]. The posterior parietal cortex (PPC) plays an important

role in processing spatial information[33,34]. Our results point out that area RL, a

subregion of PPC, is crucial for cross-modal generalization based on abstract

representations of peri-personal space. Area RL displays visuo-tactile neuronal

responses[19] and is implicated in whisking movements control[35]. It is strategically

positioned between the primary visual cortex and the primary whisker somatosensory

cortex, facilitating bidirectional information transfer between these sensory systems.

Area RL possesses a retinotopic map biased towards the lower part of the visual field[24]

where most whiskers are visible, and its neurons are tuned to high binocular disparity,

aligning with objects in close proximity, potentially within whiskers' reach[36]. It was also

implicated in optic flow processing[37] while its contribution to high-order motion

computations remains debated[38,39]. Our findings unveil a fundamental functional

specialization of area RL, confirming its unique role in facilitating visuo-tactile

coordination and sensory abstraction in the peri-personal space. Future research

could elucidate how this area may contribute to sensorimotor integration during active

multimodal exploration of nearby objects, similar to multimodal limb posture encoding

observed in the primate PPC[40].

Other forms of cross-modal generalization, such as object recognition, may

engage different cortical regions. Indeed, areas of the dorsal stream are classically

associated with spatial, attentional, and motion processing or motor guidance, while

areas of the ventral stream are involved in object recognition. Previous studies have

reported visuo-tactile multimodal responses in both the ventral and dorsal pathway in

primates[41–43], including areas initially thought to be exclusive to visual processing like

V4[44] and MT/V5[45]. Our study uncovered visuo-tactile areas belonging to the ventral or

dorsal stream in mice[46,47]. Specifically, RL and A are typically associated to the dorsal

stream while LI, with its significant projection to the postrhinal (POR) cortex, is

associated with the ventral stream[46]. Lesions in the perirhinal cortex in rats, downstream to POR, obstruct cross-modal object recognition during spontaneous exploration[31]. Moreover, areas LI and AL have been functionally characterized as specialized for shape processing and object recognition[48,49], suggesting that they could represent a rodent homologue of the visuo-haptic subregion of the lateral occipital complex (LOtv) reported in human fMRI studies[41]. These reports suggest that the ventral pathway might contribute specifically to cross-modal object recognition. Future research focusing on objects discrimination could provide further insight into the specific role of visuo-tactile areas in the ventral stream.

Wide-field imaging and anatomical mapping experiments revealed a more extensive and organized network of visuo-tactile associative cortical areas than previously identified[19,46,50]. Many associative areas exhibited a shared representation of vertical space between visual and whisker tactile inputs. These spatial maps extended even to primary sensory cortices through associative areas, with strong responses observed in S1 following visual stimulations despite no direct projections between V1 and S1[28]. Although recent findings advise caution in interpreting cross-modal signals between primary cortical areas due to potential confusion with signals evoked by uninstructed movements[27], our results suggest that certain fundamental features, such as spatial location, indeed have a shared representations across various cortical areas. These representations possibly support object or event-oriented abstract encoding[51] for cross-modal generalized learning. Organized feedback projections from associative areas to primary sensory cortices appear to play an unsuspected role for learning, in addition to their known role in shaping functional properties with contextual information[52]. This raises the question of whether the same or different feedback projections subserve these functions.

Consistent with the literature on multisensory integration[53], we observed multisensory modulations that enhanced responses to spatially congruent visuo-tactile stimuli and suppressed responses to incongruent stimuli. This response pattern could potentially arise from surround suppression for incongruent inputs, mediated by local parvalbumin-positive interneurons as observed for conflicting visuo-auditory stimuli[12]. By comparing congruent to incongruent combinations, we uncovered multisensory modulation rules that shape a supramodal representation of peri-personal space. Supramodal encoding of gratings orientation was reported in neurons of the rat posterior parietal cortex (PPC) in response to visuo-tactile stimuli[6] despite a lack of organized functional maps for orientation selectivity in rodents. Further investigations at the single-neuron level could help elucidate the mechanisms of synaptic plasticity underlying the emergence of these supramodal representations during development.

Information generalization across modalities is crucial not only for biological brains but also for artificial systems. With the advent of multimodal large language models such as GPT-4o and Gemini, it is widely recognized that multimodality is fundamental for general artificial intelligence[55,56] and can lead to the emergence of abstract conceptual representations in machine learning systems[57]. We believe that the supramodal encoding of peri-personal space by multisensory neurons we reported could represent an early instance of the same computational mechanism leading to the emergence of conceptual representations[9] that form the building blocks of flexible mental representations known as cognitive maps[1]. In this light, our work represents a major step toward understanding the circuit architecture of one of the most fundamental cognitive computations: sensory abstraction.

References

1. Behrens, T. E. J. *et al.* What Is a Cognitive Map? Organizing Knowledge for Flexible Behavior. *Neuron* **100**, 490–509 (2018).

2. Cloke, J. M., Jacklin, D. L. & Winters, B. D. The neural bases of crossmodal object recognition in non-human primates and rodents: A review. *Behavioural Brain Research* **285**, 118–130 (2015).

3. Davenport, R. K. & Rogers, C. M. Intermodal Equivalence of Stimuli in Apes. *Science (1979)* **168**, 279–280 (1970).

4. Solvi, C., Gutierrez Al-Khudhairy, S. & Chittka, L. Bumble bees display cross-modal object recognition between visual and tactile senses. *Science (1979)* **367**, 910–912 (2020).

5. Bruck, J. N. & Pack, A. A. Understanding across the senses: cross-modal studies of cognition in cetaceans. *Anim Cogn* **25**, 1059–1075 (2022).

6. Nikbakht, N., Tafreshiha, A., Zoccolan, D. & Diamond, M. E. Supralinear and Supramodal Integration of Visual and Tactile Signals in Rats: Psychophysics and Neuronal Mechanisms. *Neuron* **97**, 626-639.e8 (2018).

7. Samborska, V., Butler, J. L., Walton, M. E., Behrens, T. E. J. & Akam, T. Complementary task representations in hippocampus and prefrontal cortex for generalizing the structure of problems. *Nat Neurosci* **25**, 1314–1326 (2022).

8. Miller, E. K., Nieder, A., Freedman, D. J. & Wallis, J. D. Neural correlates of categories and concepts. *Curr Opin Neurobiol* **13**, 198–203 (2003).

9.   Quiroga, R. Q., Reddy, L., Kreiman, G., Koch, C. & Fried, I. Invariant visual

representation by single neurons in the human brain. *Nature* **435**, 1102–1107

(2005).

10.   Garner, A. R. & Keller, G. B. A cortical circuit for audio-visual predictions. *Nat*

Neurosci **25**, 98–105 (2022).

11.   Godenzini, L. *et al.* Auditory input enhances somatosensory encoding and tactile

goal-directed behavior. *Nat Commun* **12**, (2021).

12.   Song, Y. H. *et al.* A Neural Circuit for Auditory Dominance over Visual Perception.

Neuron **93**, 940-954.e6 (2017).

13.   Weiler, S. *et al.* A primary sensory cortical interareal feedforward inhibitory

circuit for tacto-visual integration. *Nat Commun* **15**, 3081 (2024).

14.   Jenkinson, E. W. & Glickstein, M. Whiskers, barrels, and cortical efferent pathways

in gap crossing by rats. *J Neurophysiol* **84**, 1781–1789 (2000).

15.   Sofroniew, N. J., Cohen, J. D., Lee, A. K. & Svoboda, K. Natural whisker-guided

behavior by head-fixed mice in tactile virtual reality. *Journal of Neuroscience* **34**,

9537–9550 (2014).

16.   Drager, U. C. & Hubel, D. H. Physiology of visual cells in mouse superior colliculus

and correlation with somatosensory and auditory input. *Nature* **253**, 203–204

(1975).

17.   Drager, U. C. & Hubel, D. H. Topography of visual and somatosensory projections

to mouse superior colliculus. *J Neurophysiol* **39**, 91–101 (1976).

18.   Stein, B. E., Stanford, T. R. & Rowland, B. A. Development of multisensory

integration from the perspective of the individual neuron. *Nature Reviews*

Neuroscience vol. 15 520–535 Preprint at https://doi.org/10.1038/nrn3742

(2014).

19.   Olcese, U., Iurilli, G. & Medini, P. Cellular and synaptic architecture of

multisensory integration in the mouse neocortex. *Neuron* **79**, 579–593 (2013).

20.   Matteucci, G. *et al.* Cortical sensory processing across motivational states during

goal-directed behavior. *Neuron* **110**, 4176-4193.e10 (2022).

21.   Chéreau, R. *et al.* Dynamic perceptual feature selectivity in primary

somatosensory cortex upon reversal learning. *Nat Commun* **11**, 3245 (2020).

22.   Banerjee, A. *et al.* Value-guided remapping of sensory cortex by lateral

orbitofrontal cortex. *Nature* **585**, 245–250 (2020).

23.   Tees, R. The effects of posterior parietal and posterior temporal cortical lesions

on multimodal spatial and nonspatial competencies in rats. *Behavioural Brain*

Research **106**, 55–73 (1999).

24.   Zhuang, J. *et al.* An extended retinotopic map of mouse cortex. *Elife* **6**, 1–29

(2017).

25.   Yamashita, T. *et al.* Diverse Long-Range Axonal Projections of Excitatory Layer

2/3 Neurons in Mouse Barrel Cortex. *Front Neuroanat* **12**, (2018).

26.   Choi, I., Lee, J.-Y. & Lee, S.-H. Bottom-up and top-down modulation of

multisensory integration. *Curr Opin Neurobiol* **52**, 115–122 (2018).

27.   Bimbard, C. *et al.* Behavioral origin of sound-evoked activity in mouse visual

751   cortex. *Nat Neurosci* **26**, 251–258 (2023).

752   28.   Wang, Q. & Burkhalter, A. Area map of mouse visual cortex. *Journal of*

753   *Comparative Neurology* **502**, 339–357 (2007).

754   29.   Morimoto, M. M., Uchishiba, E. & Saleem, A. B. Organization of feedback

755   projections to mouse primary visual cortex. *iScience* **24**, (2021).

756   30.   Zingg, B., Peng, B., Huang, J., Tao, H. W. & Zhang, L. I. Synaptic Specificity and

757   Application of Anterograde Transsynaptic AAV for Probing Neural Circuitry. *The*

758   *Journal of Neuroscience* **40**, 3250–3267 (2020).

759   31.   Winters, B. D. & Reid, J. M. A Distributed Cortical Representation Underlies

760   Crossmodal Object Recognition in Rats. *The Journal of Neuroscience* **30**, 6253–

761   6261 (2010).

762   32.   Yeterian, E. H. The effects of visual or auditory cortical lesions on specific cross-

763   modal transfer in the rat. *Neuropsychologia* **15**, 517–527 (1977).

764   33.   Whitlock, J. R. Posterior parietal cortex. *Current Biology* **27**, R691–R695 (2017).

765   34.   Pasalar, S., Ro, T. & Beauchamp, M. S. TMS of posterior parietal cortex disrupts

766   visual tactile multisensory integration. *European Journal of Neuroscience* **31**,

767   1783–1790 (2010).

768   35.   Mohan, H. *et al.* Functional Architecture and Encoding of Tactile Sensorimotor

769   Behavior in Rat Posterior Parietal Cortex. *The Journal of Neuroscience* **39**, 7332–

770   7343 (2019).

36.   La Chioma, A., Bonhoeffer, T. & Hübener, M. Area-Specific Mapping of Binocular

Disparity across Mouse Visual Cortex. *Current Biology* **29**, 2954-2960.e5 (2019).

37.   Rasmussen, R. N., Matsumoto, A., Arvin, S. & Yonehara, K. Binocular integration of

retinal motion information underlies optic flow processing by the cortex. *Current*

Biology **31**, 1165-1174.e6 (2021).

38.   Matteucci, G., Bellacosa Marotti, R., Zattera, B. & Zoccolan, D. Truly pattern:

Nonlinear integration of motion signals is required to account for the responses

of pattern cells in rat visual cortex. *Sci Adv* **9**, (2023).

39.   Sit, K. K. & Goard, M. J. Distributed and retinotopically asymmetric processing of

coherent motion in mouse visual cortex. *Nat Commun* **11**, 3565 (2020).

40.   Graziano, M. S. A., Cooke, D. F. & Taylor, C. S. R. Coding the Location of the Arm by

Sight. *Science (1979)* **290**, 1782–1786 (2000).

41.   Amedi, A., Malach, R., Hendler, T., Peled, S. & Zohary, E. Visuo-haptic object-

related activation in the ventral visual pathway. *Nat Neurosci* **4**, 324–330 (2001).

42.   James, T. W. & Kim, S. Dorsal and Ventral Cortical Pathways for Visuo-haptic

Shape Integration Revealed Using fMRI. in *Multisensory Object Perception in the*

Primate Brain 231–250 (Springer New York, New York, NY, 2010).

doi:10.1007/978-1-4419-5615-6_13.

43.   Lacey, S. & Sathian, K. Visuo-haptic multisensory object recognition,

categorization, and representation. *Front Psychol* **5**, (2014).

44.   Maunsell, J. H. R., Sclar, G., Nealey, T. A. & DePriest, D. D. Extraretinal

792   representations in area V4 in the macaque monkey. *Vis Neurosci* **7**, 561–573

793   (1991).

794   45.   Hagen, M. C. *et al.* Tactile motion activates the human middle temporal/V5

795   (MT/V5) complex. *European Journal of Neuroscience* **16**, 957–964 (2002).

796   46.   Wang, Q., Sporns, O. & Burkhalter, A. Network analysis of corticocortical

797   connections reveals ventral and dorsal processing streams in mouse visual

798   cortex. *Journal of Neuroscience* **32**, 4386–4399 (2012).

799   47.   Marshel, J. H., Garrett, M. E., Nauhaus, I. & Callaway, E. M. Functional specialization

800   of seven mouse visual cortical areas. *Neuron* **72**, 1040–1054 (2011).

801   48.   Tafazoli, S. *et al.* Emergence of transformation-tolerant representations of visual

802   objects in rat lateral extrastriate cortex. *Elife* **6**, (2017).

803   49.   Froudarakis, E. *et al.* Object manifold geometry across the mouse cortical visual

804   hierarchy. doi:10.1101/2020.08.20.258798.

805   50.   Caron-Guyon, J. *et al.* Neuronal encoding of multisensory motion features in the

806   rat associative parietal cortex. *Cerebral Cortex* **30**, 5372–5386 (2020).

807   51.   Ghazanfar, A. A. & Schroeder, C. E. Is neocortex essentially multisensory? *Trends*

808   *in Cognitive Sciences* vol. 10 278–285 Preprint at

809   https://doi.org/10.1016/j.tics.2006.04.008 (2006).

810   52.   Keller, A. J., Roth, M. M. & Scanziani, M. Feedback generates a second receptive

811   field in neurons of the visual cortex. *Nature* **582**, 545–549 (2020).

812   53.   Stein, B. E. & Stanford, T. R. Multisensory integration: Current issues from the

813   perspective of the single neuron. *Nat Rev Neurosci* **9**, 255–266 (2008).

814   54.   Gemini Team *et al.* Gemini: A Family of Highly Capable Multimodal Models.

815   (2023).

816   55.   Fu, L. *et al.* A Touch, Vision, and Language Dataset for Multimodal Alignment.

817   (2024).

818   56.   Wu, J., Gan, W., Chen, Z., Wan, S. & Yu, P. S. Multimodal Large Language Models: A

819   Survey. (2023).

820   57.   Goh, G. *et al.* Multimodal Neurons in Artificial Neural Networks. *Distill* **6**, (2021).

821   58.   Lein, E. S. *et al.* Genome-wide atlas of gene expression in the adult mouse brain.

822   *Nature* **445**, 168–176 (2007).

823   59.   Pachitariu, M. *et al.* Suite2p: beyond 10,000 neurons with standard two-photon

824   microscopy. *bioRxiv* 061507 (2016) doi:10.1101/061507.

825   60.   Wallace, M. T., Meredith, M. A. & Stein, B. E. Multisensory Integration in the

826   Superior Colliculus of the Alert Cat. *J Neurophysiol* **80**, 1006–1010 (1998).

827   61.   Jazayeri, M. & Movshon, J. A. Optimal representation of sensory information by

828   neural populations. *Nat Neurosci* **9**, 690–696 (2006).

829   62.   Ma, W. J., Beck, J. M., Latham, P. E. & Pouget, A. Bayesian inference with

830   probabilistic population codes. *Nat Neurosci* **9**, 1432–1438 (2006).

831   63.   Boyden, E. S., Zhang, F., Bamberg, E., Nagel, G. & Deisseroth, K. Millisecond-

832   timescale, genetically targeted optical control of neural activity. *Nat Neurosci* **8**,

833   1263–1268 (2005).

834   64.   Efron, B. & Tibshirani, R. J. *An Introduction to the Bootstrap*. (Chapman and

835   Hall/CRC, 1994). doi:10.1201/9780429246593.

Methods

Methods

<u>Animals</u>

All experiments were carried out in accordance with the Institutional Animal Care and

Use Committee of the University of Geneva and with permission of the Geneva

cantonal authorities (GE258B). C57BL/6J wild-type mice and transgenic mouse lines

were housed 2–6 mice per cage under a 12/12-h non-inverted light/dark cycle with ad

libitum access to food and water. Transgenic mice used for calcium imaging were

obtained as a crossing between Ai148-D mice (Jackson Laboratories, stock number

030328) and Rasgrf2-2A-dCre mice (Jackson Laboratories, stock number 022864).

Ai148-D mouse line is a Cre-dependent reporter line containing a gene encoding the

calcium indicator GCaMP6f at the Igs7 locus. Exposure to Cre recombinase through

viral vector injections or crossing with Cre-expressing mice resulted in expression of

GCaMP6f. Rasgrf2-2A-dCre mouse line expressed a trimethoprim-inducible Cre

recombinase directed by endogenous Rasgrf2 promoter/enhancer elements. When

induced, Cre recombinase activity is observed in cortical layers 2/3 and other scattered

cells of the cortex, hypothalamus, thalamus, and midbrain. Trimethoprim (TMP) i.p.

injections were performed for 5 consecutive days at least two weeks before any

surgical intervention (0.25 mg/g of body weight diluted in DMSO and 0.9% NaCl). As

a result, TMP injected crossed Ai148D x Rasgrf2-2A-dCre mice expressed GCaMP6f

in cortical layer 2/3. For optogenetic experiments, we used Ai32 transgenic mice that

express Channelrhodopsin-2 (ChR2) fused to enhanced Yellow Fluorescent Protein

(eYFP) in a Cre-dependent manner, enabling fast neuronal activation in vivo through

blue light illumination (Jackson Laboratories, stock number 012569). Males and

females aged 2–5 months and weighing approximately 20–25 g were used for the

experiments.

Viral vectors and markers

For anterograde labeling, AAV2.CAG.GFP (Addgene, #37825-AAV2, titer: 7×10^{12} vg/mL) and AAV2.CAG.tdTomato (Addgene, #59462-AAV2, titer: 4×10^{12} vg/mL) were used. The volume of viral vectors injected was around 50-75nL in each site at a depth of approximately 400-500 μm from the surface of the cortex. Brains were collected 3 weeks after the injections. For retrograde labeling, Cholera Toxin subunit B conjugated to Alexa Fluor 555 (Invitrogen, reference number C22843) or Alexa Fluor 647 (Invitrogen, reference number C34778) diluted in PBS were injected (~50nL in each site, 400-500 μm below the surface). Brains were collected 10 days after the injections. For silencing experiments, AAV-DJ.CMV.eGFP-2A-TeNT viral vectors were injected in the region of interest (Wu Tsai Neurosciences Institute, reference number GVVC-AAV-70) at 3 different depths along the cortical column (800 μm, 500 μm and 200 μm respectively). Behavioral protocols started at least one month after the injections were done. For optogenetic experiments, Ai32 mice expressing Cre-dependent ChR2-eYFP were injected with AAV1.CamKII0.4.Cre.SV40 (Addgene, #105558-AAV1, titer: 3.5×10^{12} vg/mL) in three different locations along RL and two different depths (250 μm and 500 μm) to express ChR2-eYFP in most of this area.

Stereotaxic surgeries

Pain management was first performed by administrating the opioid Buprenorphine subcutaneously (0.1 mg/kg) before starting the surgery. Mice were anesthetized inside an induction chamber with 3% isoflurane mixed with oxygen and then fixed on the stereotaxic apparatus (Model 940, Kopf). A custom-made nose-clamp has been adapted to the apparatus to maintain the position of the animal, allowing head rotation.

Body temperature was constantly monitored through a thermic probe and adjusted to

~37°C via a heating pad placed below the mouse (DC Temperature Controller, FHC).

Breathing rate was regularly monitored by visual inspection. Ophthalmic gel (Vitamin

A, Bausch Lomb) was applied on both eyes to ensure protection from light and prevent

them from drying out. A local anesthetic was injected under the skin of the head before

the surgical incision (mix Lidocaine/Bupivacaine, 6mg/kg and 2.5 mg/kg respectively).

During the surgery, mice were anesthetized with a constant isoflurane level lowered

around 2%. At the end of the surgery, an anti-inflammatory was also administered

subcutaneously (Carprofen, 7.5mg/kg) and animals were warmed with a heating lamp

for at least 15 minutes until recovery from anesthesia. A second anti-inflammatory

(Ibuprofen, Algifor) was added to drinking water for 3-days post-op and the weight was

checked daily to ensure that weight was kept above 15% of the original weight prior to

the surgery. All animals were implanted with a cranial implant. After removing the skin

and tissues on top of the head, the skull was cleaned, dried and thinned. The mouse

head was tilted approximately at a 30° angle, ensuring better access to the left

hemisphere. A custom-made metallic implant was placed on the top of the skull using

a custom-made holder. It was then fixed with a layer of glue (Loctite 401, Henkel) and

additional layers of dental acrylic (Pala, Kulzer) to ensure the implant is securely

attached. Dental acrylic was covered with black nail polish to prevent light

contamination from visual stimuli during imaging experiments. After at least 3 days of

recovery, mice could undergo additional procedures. Animals that underwent imaging

sessions were implanted with a glass window composed of a top round cover slip of 7

mm diameter and two superimposed 5 mm diameter cover slips (CS-7R and CS-5R,

Multi Channel Systems). The three concentric cover slips were glued together with UV

glue (Optical Adhesive n°68, Norland). The craniotomy was the size of the smaller

window and was drilled above the posterior part of the dorsal cortex. Before removing

skullcap, we used a custom-made perfusion chamber with saline solution (NaCl 0.9%)

to rinse the craniotomy continuously and reduce bleeding stains. A custom-made

holder with air-suction was used to hold and position the window. The cranial window

was gently brought down until it was in contact with the brain. The window was then

fixed with UV glue, super glue and dental acrylic. For stereotaxic injections, glass

pipettes (5-000-2005, Drummond) were pulled (Model P-97, Sutter Instrument Co) and

broken to obtain a tip of ~10-15µm inner diameter. Pipettes were further beveled to

create a sharp tip to avoid cortical damage during insertion. Injection sites were

determined using functional mapping. Injections were done using a single-axis oil

hydraulic micromanipulator (R.MO-10, Narishige). The pipette was slowly inserted

inside the cortex until reaching the desired depth. Viral vectors or other reagents were

injected at a speed of ~2nL/s. When the whole volume was injected, we waited 5

minutes with the pipette in the same position before gently removing it.

<u>Behavioral training</u>

A Matlab custom-made graphical user interface (GUI) was developed from the Matlab

App Designer to control the behavioral tasks and monitor performance. The GUI

allowed real time visualization of the animal's performance and online modification of

the parameters (e.g. stimuli parameters, punishment duration, stimuli proportion).

Animals underwent water restriction 2 to 4 days before the training started and were

handled every day by the experimenter for at least 10 minutes. During the pre-training

phase, mice were habituated to head-fixation and placed on the setup. In the first

session, only Go stimuli (i.e., stimuli for which licking responses are rewarded) were

presented and rewards were delivered automatically regardless of mouse actions. Go

trials proportion was progressively reduced as No go trials (i.e., trials featuring the

stimulus for which licking responses were punished) were added in the following

sessions and mice were required to lick to obtain rewards during Go trials. Mice were

trained once a day, every day at the same hour. Eight different behavioral tasks (two

tactile tasks, two visual tasks, two auditory tasks and two optogenetic tasks) were used

for our experiments (Fig. 1, Extended Data Fig. 1, Fig. 5 and Fig. 6), all following a

Go/No go discrimination paradigm. In the tactile tasks, two whiskers ("top" whisker B2

and "bottom" whisker C2) were inserted inside glass capillaries each attached to a

piezo actuator that could create a small deflection of about 1 mm along the rostro-

caudal axis. In one version of the task, B2 whisker stimulations were associated to a

reward (Go trials). In the other version, C2 whisker stimulations were the Go trials. In

the visual tasks, two drifting squares (a "top" square and a "bottom" square, relative to

the midline in the mouse visual field) were presented on the screen. In one version of

the task, top square stimulations were the Go trials while in the other version of the

task, bottom squares stimulations were the Go trials. The auditory task followed the

same structure as the other tasks but using two short pure tones of 6 kHz and 12 kHz

as Go and No go stimuli, respectively. These tones were delivered from a speaker

located next to the mouse on the same side as the visual and tactile stimuli. The

optogenetic task was performed by directly stimulating subregions of RL

corresponding to top or bottom multimodal representation using shaped blue light

patterns. In one version of the task, stimulating the top region was associated with a

reward, while in the other version, it was the bottom region. The total trial duration of

a single trial was 4 s: after a 2 s quiet window (during which licking resulted in trial

abortion) the stimulus was presented, and the mouse was allowed to lick during a 2 s

response window. During Go trials, the mouse could obtain a water reward upon

licking the spout following the stimulus (Hit trials). Failure to lick would result in a Miss

trial. During No go trials, the mouse had to refrain from licking (Correct rejection trials,

CR) or it was punished with a time-out of 10 s (False Alarm trials, FA). Some trials

were presented without any stimulus (catch trials). If the mouse licked during catch

trials, no time-out was applied. If mice were too compulsive (i.e. licking during the quiet

window was too frequent), a 10 s time-out early lick punishment could also be applied.

The proportion of each trial type was the following: Go trials = 30%, No go trials = 50%,

Catch trials = 20%. All tactile stimuli were generated through Matlab data acquisition

toolbox controlling a piezo actuator (Bimorph bendor piezo actuator PB4NB2S,

Thorlabs) through a National Instrument card. All visual stimuli were generated using

Matlab and PsychToolBox. Stimuli were presented on a gray background through a

LCD monitor (20x15cm, pixels, 60Hz refresh rate, Pi-shop) positioned 10 cm from the

eye, with a 30° angle to the right of the midline. The screen was also tilted with a 30°

angle along the horizontal plane to match the mouse head angle with the intent of

roughly aligning the bottom and top parts of the screen to the resting position of the

C2 and B2 whiskers in the mouse visual field. Stimulations during behavior consisted

of black squares moving through the screen in the rostro-caudal direction (bar width

was 15°, stimulus duration was 175ms, speed was 500°/sec), on a gray background.

Wide-field microscopy

We used a custom-made wide-field epifluorescence microscope setup[39] including a

sCMOS camera (ORCA-Flash4.0 V3, Hamamatsu). Magnification was achieved

through a 0.63X C-mount camera adapter for Olympus Microscopes. The field-of-view

size was 5.6 mm x 5.6 mm. The camera and adapters were mounted on a base

allowing vertical movement with manual focus. LED white illumination (740mW,

1225mA, Thorlabs) could be controlled via a T-Cube LED driver (LEDD1B, Thorlabs).

Filter cubes could be changed for different types of imaging. For imaging GCaMP6f,

GFP excitation, emission and dichroic filters were used. For intrinsic optical signals

imaging, Cy3/5 excitation, emission and dichroic filters were used. An objective (MVX

Plan Apochromat with 2x, Olympus) was attached to the microscope base. The

somatotopic mapping protocol consisted of repetitive rostro-caudal pulsatile

deflections (~1 mm amplitude) of either B2 or C2 whiskers for 50-80 trials each

followed by a quiet window. The retinotopic mapping protocol consisted of drifting bars.

A contrast reversing checkerboard was presented within the bar to better drive neural

activity (0.04 cycles/° of spatial frequency and 2 Hz of temporal frequency). In each

trial the bar was swept in the four cardinal directions: left to right, right to left, bottom

to top, and top to bottom. Single trials were repeated 20 to 40 times. For anatomical

experiments, only C57BL/6J mice were used. To image the intrinsic optical signal, we

used longer tactile stimulations and slower visual stimulations. Mice were fixed on the

platform and anesthetized during the procedure with isoflurane level lowered than 1%.

Body temperature was monitored with a probe and adjusted to 37°C using a heating

pad (DC Temperature Controller, FHC). For all the other experiments, we used Ai148-

D x Rasgrf2-2A-dCre mice and calcium imaging. The visuo-tactile sparse noise

protocol consisted in combinations of visuo-tactile stimuli: three whisker conditions (C2

whisker, B2 whisker and no whisker), eight vertical positions of the moving square.

The ninth position corresponded to a no visual stimulus condition (i.e. blank screen).

Visual and tactile stimulus onsets were either synchronous or delayed (0.15 s delay,

visual leading tactile stimulus). In total, 54 different combinations were presented in

pseudo-random order with a 1 s inter-stimulus interval. The full sequence was

repeated 60 times. Total sparse noise protocol duration was approximately 1 hour. At

the beginning of each recording, we took a picture of the cranial window with blood

vessels pattern on focus as reference image. The focus was then set at ~300μm below

the surface to maximize signal collection. Light was adjusted to prevent saturation.

Before each imaging session, the window was cleaned with 70% ethanol and eyes

hydrated with mineral oil.

1018

<u>Two-photon microscopy</u>

The two-photon microscope was custom-made (INSS Company). It consisted of a

femtosecond laser with wavelength range 690-1040nm (Tunable Ti:Sapphire with

dispersion compensation MaiTai DeepSee, Spectra Physics) which beam was

directed with Resonant/Galvo scan mirrors and the emitted signals were detected by

2 GaAsP amplified PMTs (PMT2101/M, Thorlabs). Imaging was performed through

ScanImage (Vidrio Technologies). Images were acquired at approximately 30 frames

per second. Two-photon calcium imaging during the visuo-tactile sparse noise

protocol described in the wide-field section was performed on Ai148-D x Rasgrf2-2A-

dCre mice. After cleaning the cranial window with 70% ethanol, a hydrophobic

chamber was made between the head plate and the imaging platform using liquid

plastic (Smooth-Cast 325, Smooth-On) and the objective was immersed in distilled

water. We ensured no polluting light could reach the objective by covering it with a

dark protective sleeve and by turning off the light in the room. At the beginning of each

recording, an anatomical picture of the field-of-view was taken using a CCD camera.

When switching the microscope in two-photon mode, we took an image of the surface

blood vessels at magnification x1 and x1.5 for further realignment. For each location,

we typically imaged at three different depths between 100 µm and 300 µm below the

surface. Each mouse underwent no more than two imaging protocols per day.

1038

1039

Histology

Mice were anesthetized with 3% isoflurane and euthanized with i.p. injection of Pentobarbital (Eskornarkon, 150 mg/kg) and Buprenorphine (0.1 mg/kg). They were then transcardiacally perfused using a peristaltic pump (ISM829, Cole-Parmer) with 0.01 M PBS, pH 7.4 for 2 min, and then 4% paraformaldehyde (PFA) in phosphate buffer (PB; pH 7.4) for 3 min. The brains were post-fixed at 4°C in PFA for 48 h and then transferred into PBS. For anatomical experiments of Fig. 3, the brains were post-fixed at 4°C in PFA for 2 h and washed 3 times for 15 min in PBS. The left cerebral hemisphere was separated from the right hemisphere and subcortical parts were removed with a spatula. The left hemisphere was flattened between glass slides and kept flattened at 4°C in PFA for 12 h. After PBS washing, flattened hemispheres were embedded in agar gel 4% before cutting slices of 60µm thickness with a vibratome (VT 1000 S, Leica).

Immunohistochemistry

Brain sections were incubated with slight agitation (40 rotations/min) for 2 h at room temperature in a saturation/permeabilization solution containing a mix of 5% BSA and NGS, 0.3% triton X-100 and PBS. Brain sections were then incubated with slight agitation (40 rotations/min) overnight at 4°C with a rat anti-muscarinic acetylcholine receptor M2 (M2 AChR) primary antibody (1:500, Sigma-Aldrich, reference number MAB367) in the same blocking solution. Sections were washed 3 times for 15 min in PBS before a 2 h incubation with either a donkey anti-rat secondary antibody conjugated with Alexa Fluor 488 (dilution 1:500, Invitrogen, #A21208) or with a goat anti-rat secondary antibody conjugated with Cy5 (dilution 1:500, Invitrogen, #A10525)

1064 in blocking solution at room temperature without agitation. Hoechst solution (dilution

1065 1:1000, Invitrogen, # 33342) was used for fluorescent nuclear counterstaining. Slices

1066 were incubated for 10 min in that solution with slight agitation then washed 2 times for

1067 5 min in PBS.

1068

1069 <u>Histological imaging</u>

1070 Slices were mounted on Superfrost microscope slides (Epredia) with mounting

1071 medium (Fluoromount) and covered with 24x50 mm coverslips (Menzel-Gläser). All

1072 photomicrographs were taken using a Zeiss Axio Scan.Z1 or a Leica Stellaris 5

1073 microscope at the bioimaging platform of the University of Geneva.

1074

<u>Wide-field calcium imaging analysis: Retinotopic and whisker mapping protocols</u>

Responses to drifting checkerboard stimuli were averaged across trials for each

condition and converted in df/f using a pre-stimulus time window as baseline. Azimuth

and elevation preference maps were then computed[24].To segment visual areas,

azimuth and elevation maps were combined to generate a visual field sign map. The

sign map was computed as the sine of the difference between the vertical and

horizontal retinotopic gradients for each pixel. Somatotopic maps were obtained by

averaging the response df/f across B2 or C2 stimulation trials over a short time window

of 200 ms.

1084

1085

 Wide-field calcium imaging analysis: Sparse noise protocol

After an initial down sampling to a resolution of 100 x 100 pixels by bicubic

interpolation, 50 Hz framerate videos acquired during the sparse noise protocol

underwent pixelwise notch filtering (f_0=12 Hz, f_w=6 Hz) and discrete wavelet transform

(DWT) detrending (setting lowest frequency approximation coefficients to zero in a 6-

level decomposition) to remove artifacts and low-frequency drifts. After these initial

preprocessing steps, videos were z-scored using as reference distribution the

ensemble of all pixel values corresponding to blank stimuli (i.e. trials with no visual

and no tactile stimulation). Next, trials with a high baseline activity (>=0.5 z-score on

average, corresponding to a strong positive fluctuation of spontaneous activity

preceding the stimulus) were discarded and z-score and baseline subtraction were

computed again including only remaining trials. This was meant to better separate

local responses to sensory stimulations from ongoing spontaneous activity. Starting

from the z-scored data obtained in this way, response surprises (i.e. -log10($p_{t\text{-}test}$))

across trials were computed for each time point and each stimulus to define spatial

responsivity masks. Average visual or tactile top or bottom response movies were

obtained by averaging the median responses across trials to the relevant stimuli. For

visual "top" all stimulation conditions (either unimodal or multimodal) in which the

visual stimulus was present in grid positions 7 or 8 were used whereas for visual

"bottom" all visual conditions including positions 3 or 4 were used. Retinotopic and

somatotopic maps of vertical space (Fig. 2b,c and Extended Data Fig. 5g) were

obtained by taking the difference between average visual or tactile top and bottom

response maps. Modality preference maps (Fig. 2d and Extended Data Fig. 5g) were

obtained by computing pixel-by-pixel the difference between the maximum response

for visual and tactile stimuli. Multisensory enhancement maps (Fig. 2f and Extended

Data Fig. 5g) were obtained by computing pixel-by-pixel the difference between the

maximum unisensory response between visual and tactile stimuli (predicted response)

and the multisensory response (measured responses). For all the difference maps

mentioned above the input maps were independently normalized to the [0,1] interval

before taking the difference (ensuring their range to be bound in the [-1,1] interval).

Retinotopy-somatotopy spatial coherence maps (Fig. 2e and Extended Data Fig. 5g)

were obtained by convolutionally (stride = 1 patch size = 8 pixels) computing the

correlation-based similarity of matching patches of the two different maps. The result

was then scaled by the average peak-to-peak range in that patch (to diminish

contribution from pixel with very low response magnitude). Comparison of grand

average maps at single pixel level (Fig. 2g,h) was performed on responsive pixels only

(exceeding a surprise threshold corresponding to p_{t-test} <= 0.1 for either visual or tactile

responses). We compared pixels with low and high modality preference (i.e. below or

above 85% quantile of the absolute value distribution within the responsive region) or

with low and high coherence (i.e. below or above 15% quantile of the absolute value

distribution within the responsive region). Grand average maps displayed in Fig. 2b,c,f

are obtained by averaging frames of the corresponding grand-average movie

(obtained by realigning single mouse movies to the common reference atlas and then

averaging) over different time windows: frames 18 to 20 for the somatotopy (i.e. 0.16

s to 0.20 s after stimulus onset); frames 25 to 27 for the retinotopy (i.e. 0.30 s to 0.34

s after stimulus onset); and frames 28 to 30 for multisensory modulation (i.e. 0.36 s to

0.4 s after stimulus onset). These windows were chosen to be roughly centered on the

peak of these signals. Grand average modality preference and spatial coherence

maps displayed in Fig. 2d,e were computed on maps obtained above (average z-score

maps or position selectivity maps, respectively). Signal time courses displayed in

Extended Data Fig. 6 were obtained by integrating grand-average movies across time-

varying regions of interest (ROIs). These ROIs were specifically defined to track the activity 'bump' evoked by stimulation in each area. To prevent bias in estimating the response —particularly in areas with higher cortical magnification factors, which could lower the average by including many non-responsive pixels— these masks were created by intersecting a static mask, which selects each area based on the fitted reference atlas, with a dynamic responsivity mask. This dynamic mask selects, for each movie frame, only those pixels that across mice show an average maximum response surprise, exceeding a threshold (corresponding to a p-value ≤ 0.01 across conditions). To measure angular mismatch between retinotopic and somatotopic maps we computed vertical retinotopy gradient vectors from the grand-average version of these maps (using Matlab "gradient" function with a scale of 4 pixels followed by a Gaussian smoothing of gradient components). Then a region of interest overlapping with RL was defined and we computed the average angular difference between the two gradient fields.

Atlas fitting and registration

The reference Allen Mouse Brain atlas[58] was first tilted sideways at 30° and then projected to match the skull tilt in our experiments and therefore the layout of areas in the dorsal cortex. The resulting atlas was manually fitted to the wide-field imaging field-of-view for each mouse by visually aligning the atlas boundary lines to reproducible landmarks from functional maps. Both the maps obtained from the sparse noise protocol (i.e. vertical retinotopy and somatotopy difference maps) and the maps obtained from the whisker and retinotopic mapping protocols were used to register the atlas. Landmarks used include the outlines of the sign map regions, the position of C2 and B2 whisker activity bumps in S1 and S2, the reversal of vertical retinotopy at the

boundary of each visual area (see Extended Data Fig. 5 for example). To relate

microscale and macroscale functional properties measured respectively by two-

photon and wide-field imaging experiments, we also manually reconstructed the

position of each two-photon field-of-view (FOV) in the frame of reference with the atlas

fitted to the wide-field maps. To do so, we aligned the blood vessels pattern visible in

each two-photon FOV with the one visible in the wide-field cortical image (as shown

in Extended Data Fig. 8f). To compare data across mice and compute grand averages,

we developed a Matlab pipeline to robustly realign atlases to one another. This

pipeline was based on iterative application of the image registration algorithm

implemented by Matlab function "imregtform" (considering "rigid" transformations in

"monomodal" mode) to stack images depicting atlas boundaries. This pipeline enabled

us to obtain the rotation and shift required to register all maps and neuron positions to

the frame of reference of a common atlas. For the anatomical tracing experiments (Fig.

3b-d), a similar processing was performed with ImageJ through registration of atlases

fitted using the M2 AChR staining. Brain from all mice with similar injections could then

be averaged together. To compare the somatotopy and retinotopy of axonal

projections or projecting neurons with the functional topographic maps obtained in

wide-field we brought them in a common reference frame by registering a crop of the

M2 AChR-fitted atlas to the common wide-field/two-photon atlas (by finding the

registering affine transformation with Matlab functions "cpselect" and "fitgeotrans").

1182

<u>Two-photon calcium imaging analysis: Pre-processing and functional cell-types</u>

To extract time-varying somatic GCaMP6f calcium signals, we used the Suite2p

toolbox[59]. Neuropil contamination was corrected by subtracting the fluorescent signal

from a surrounding ring $F_{Surround}(t)$ from somatic fluorescence: $F(t) = F_{Soma}(t)$ -

$\alpha * F_{Surround}(t)$ with $\alpha=0.7$. Neuropil-corrected fluorescence signals $F(t)$ were then

converted in z-score by subtracting from each trace the mean value and dividing by

the standard deviation of $F(t)$ over the last 0.2 s of the baseline window preceding the

stimulus onset (pooling across all trials). Using these z-scored fluorescence traces we

computed response surprises (i.e. $-\log10(p_{z\text{-test}})$) across trials for each stimulus and

for each time sample. Similarly, we computed the coefficient of variation (CV) across

trials quantifying response variability. A neuron was considered reliably responsive if

its response exceeded a stringent surprise threshold $Surprise_{th} = 8$ (i.e. $p_{z\text{-test}} <= 10^{-8}$)

while remaining below a coefficient of variation threshold value $CV_{th} = 4$ for at least 4

consecutive time bins in any stimulus condition. This responsivity criterion was

requested for each neuron to be included in subsequent analyses. For each neuron

included, we used a more inclusive criteria to characterize the extent of their tuning

properties. Indeed, neurons exceeding a less stringent threshold $Surprise_{th} = 2$ (i.e. $p_{t\text{-}}$

$_{test} <= 0.01$) while remaining below the same coefficient of variation threshold at any

time bin of a visual, tactile or visuo-tactile condition were considered "responsive" to

these conditions. The responsivity pattern of each neuron was used to classify them

in one of four functional classes: 1) neurons responsive in visual but not tactile

conditions were termed "visual neuron"; 2) neurons responsive in tactile but not visual

conditions were termed "tactile neuron"; 3) neurons responsive in both tactile and

visual conditions were termed "visuo-tactile neuron"; 4) neurons responsive only in

visuo-tactile conditions (both visual and tactile stimuli presented together) were termed

"gated neuron". Only visuo-tactile neurons displaying significant responses in both

modalities were used for the analyses displayed in Fig. 4g-j. Measured multisensory

responses were compared to a "max model" of multisensory interaction[60] (as in Fig.

4b and Extended Data Fig. 10b). Predicted visuo-tactile z-scored traces for each

multisensory stimulus condition were computed trial-wise by taking the max (for each

time bin) between the response to the same visual and tactile stimulus presented

alone (i.e. the corresponding unimodal conditions) and subsequently averaging across

trials. Average responses (predicted or measured) were computed over a response

window spanning from 0.2 s to ~0.733 s following stimulus onset for each stimulus

and each responsive neuron.

1218

<u>Two-photon calcium imaging analysis: Computation of indices</u>

We computed the "multisensory modulation index" (MI) as the normalized difference

between predicted and measured visuo-tactile responses at every time bin for every

stimulus condition. To summarize the intensity of multisensory modulation of each

neuron across stimulus conditions over time we computed the min-max range of the

integral of MI values over the same time window mentioned above (i.e. from 0.2 s to

0.733 s) across all valid conditions. Valid stimulus conditions comprised significant

responses as well as all multisensory conditions containing at least one unisensory

stimulus to which the neuron is responsive and displaying significant multisensory

modulations (i.e. with p<0.05 of a bootstrap-t test for MI value for at least one time bin,

see "quantification and statistical analysis" section below). Preferred vertical position

in the tactile or visual space of each neuron was quantified as the center-of-mass of

these average responses along the corresponding dimension of the stimulus grid on

visuo-tactile (measured or predicted) conditions. This center-of-mass was converted

into a "position preference index" ranging from +1 for neurons tuned to the top

visual/B2 whisker stimulus to -1 for neurons tuned to the bottom visual/C2 whisker

stimulus (as in Fig. 4c,d,g and Extended Data Fig. 9a-b). The sharpness of positional

tuning of neurons was computed as the inverse of the best sigma parameter of a

Gaussian fit of the average response to each visual position or as the absolute value

of the tactile position selectivity index described above. This selectivity index was

computed on the visuo-tactile condition of max response (both in the measured or

predicted case). These indices ranged from 0 for neurons responding equally to all

positions (i.e. completely positionally untuned) to 1 for neurons responding to only one

position (i.e. maximally positionally tuned). The values of positional preference and

selectivity indices shown in Fig. 4g,h,i,j are averaged over both delay conditions (i.e.

with or without delay between tactile and visual stimuli, see Extended Data Fig. 6).

Neurons were labeled as ventral or dorsal by running a k-means clustering algorithm

(with $n_{centroids}$=2) on the spatial distribution of visuo-tactile neurons over the surface of

the dorsal cortex (Fig. 4e).

1248

<u>Two-photon calcium imaging analysis: Correspondence with wide-field</u>

To quantify the correspondence between single-neuron properties distributed across

the cortex and the wide-field maps, we created smoothed spatial maps from 2D

histograms that depict the dominant response properties at each cortical location. We

generated retinotopic and somatotopic maps from two-photon data (Fig. 4c,d and

Extended Data Fig. 9a,b) by smoothing a 2D normalized histogram of neurons with

positional preferences (top-preference index≥0.05, bottom-preference index≤0.05)

using a Gaussian kernel (σ~3% of total wide-field FOV) and then calculated the

difference between these maps. For modality preference maps reconstructed using

two-photon data (Extended Data Fig. 9c), we applied a Gaussian smoothing (σ~3% of

FOV) to 2D normalized histograms that describe the distribution of unimodal visual

and tactile neurons, calculating the normalized difference between them. To map

multisensory modulation (Extended Data Fig. 9d), we applied Gaussian smoothing

(σ~3% of FOV) to a MI-range-weighted histogram showing the spatial distribution of

multisensory modulations, then normalized this against the unweighted distribution

that includes unresponsive neurons. We included unresponsive neurons to properly

capture the relative density of multisensory modulation in the local population. To

quantitatively assess the match between two-photon reconstructed maps and the

wide-field maps, we computed a Pearson correlation coefficient restricted to the pixels

with good two-photon coverage (thresholding the map shown in Extended Data Fig.

8g at a threshold of 2,000 neurons/mm2).

1270

<u>Two-photon calcium imaging analysis: Decoding analysis</u>

To test whether the neuronal representation observed across the dorsal cortex of

naïve mice could support cross-modal generalization observed in our behavioral

experiments, we adopted a decoding approach. We trained a Bayesian classifier to

infer stimulus category (top or bottom) from single-neuron responses in the tactile

modality and then tested the accuracy on the visual modality. This decoder was

inspired by original works introducing neuronal population probabilistic codes[61,62] and

is similar to the approach used in previous work[20]. We built feature vectors

representing the response of each neuron to each stimulus category (i.e. "top" and

"bottom") in each modality. The feature vectors were defined, for each neuron, as the

z-score response integrated over a 0.2 s - 0.733 s time window in every trial of the

most responsive stimulus position pertaining to each category (visual top = positions

5 to 8, visual bottom = positions 1 to 4, tactile top = B2, tactile bottom = C2). The

response feature vectors were used in a trial-wise fashion as predictors to infer the

"top" or "bottom" labels corresponding to the location of the stimulus delivered to the

animal. We incorporated 4-fold cross-validation to mitigate overfitting and assess the

classifier's accuracy outside the training dataset. Trials were partitioned into four

batches, three of them for training and one for testing. The decoding was assessed

both at the single neuron level (n_{pool} = 1 and averaged over all neurons of a given class

to get the values displayed in Extended Data Fig. 10d) as well as on pools of neurons

of increasing size (n_{pool} = [1,50] to get the curves displayed in Extended Data Fig. 10c).

During population decoding, we repeated the procedure 40 times, each time randomly

resampling the neurons in the decoding pool without replacement to ensure that the

results were not dependent on a specific selection and partitioning of neurons.

Decoding accuracy (defined as the fraction of trials with correct predictions) was

averaged across resampled data and cross-validation splits. Prior to classification,

each neuron's response features within the population vector were individually

normalized to a 0-1 range across trials. Additionally, trials were randomly reordered,

maintaining their corresponding correct labels, to eliminate noise-related correlations.

In this probabilistic framework the "training" consists simply in computing the average

tuning curves (i.e. stimulus-triggered average) using all trials included in the training

set and taking the logarithm of this vector. The subsequent "testing" of the decoder

requires to generate a prediction. Predictions were made by calculating each test trial's

feature vector log-likelihood for the 'top' or 'bottom' stimulus, then assigning the label

with the highest log-likelihood as the predicted outcome. The log-likelihood can be

obtained by taking a matrix product between the matrix storing the log of tuning curves

computed at the "training" stage and the current population feature vector (after

subtracting a corrective term proportional to the sum of all tuning curves to account for

any representational bias in the decoded population). Tactile decoding performances

represented in Extended Data Fig. 10c,d corresponded to the average cross-validated

test accuracies on the same modality on which the Bayesian classifier was trained (i.e.

tactile).

1313

Behavioral analysis

To quantify mouse behavior over time within single sessions we isolated different stimulation conditions (i.e. trials with top stimuli, trials with bottom stimuli and Catch trials) to compute a smoothed conditional lick probability over a sliding window of 60 trials. We calculated the conditional probabilities of licking by dividing the number of trials with a mouse lick response to a specific stimulus by the total presentations of that stimulus within the moving time window. These curves are the ones displayed throughout the paper (Fig. 1,5,6 and Extended Data Fig. 1,2,3,11,12) and labelled "Top", "Bottom" and "Spont". From Go and No go lick probabilities (defined by the task rule to coincide with top or bottom), we also computed a "Discrimination performance" metric as the percentage of correct trials for discrimination. Specifically, we defined behavioral performance as the average between the rate of correct choices upon delivery of the Go stimulus (i.e. Hit rate = conditional lick probability for Go trials) and the rate of correct rejects upon delivery of the No go stimulus (i.e. Correct Reject rate = 1 — False Alarm rate where False Alarm rate = conditional lick probability for No go trials). This way of computing the performance makes it insensitive to the proportion of Go and No go trials. Following a similar logic, we defined a "Detection performance" as the maximum between the Go detection rate (i.e. average Hit rate and Catch Correct Reject rate) and the No go detection rate (i.e. average False Alarm rate and Catch Correct Reject rate). Behavioral states were classified as previously described[20]. Conditional lick probabilities and performance per session were computed only during engaged trials of each session (i.e. discarding the part of the session during which conditional lick probabilities dropped due to task disengagement). Switch-aligned performance trajectories shown throughout the paper (Fig. 1,5,6 and Extended Data

Fig. 1,3,12) were obtained realigning the curve of average session performance and

conditional lick probabilities around the session of task switch. Accompanying bar

plots display the difference of discrimination and detection performance between the

last session before the switch and the first session after the switch. In Extended Data

Fig. 2, we plotted the average conditional lick probabilities and the performance time

course across mice that underwent whisker removal control sessions. These average

curves were obtained by dividing each session in three chunks: "before", "during" and

"after" whisker removal and averaging them separately after a stretching interpolation

to match their duration. Accompanying bar plots display the average discrimination

and detection performance in these three chunks (excluding boundary timepoints to

avoid contamination due to the temporal smoothing induced by the sliding window we

used to compute the traces). Learning rates following the switch (Fig. 1h and Fig. 5l)

were obtained measuring the slope of the best fitted line for 3 consecutive sessions

after switch for all mice in each group. We excluded mice that did not show a drop of

performance to chance level after the switch as they successfully generalized from

one modality to the other. The relationship between the TeNT-P2A-GFP expression

overlap with each cortical area and the cross-modal generalization performance

impairment in rule-preserving conditions was assessed using Pearson coefficient of

correlation. Specifically, for each mouse we calculated the Pearson coefficient of

correlation between two vectors: 1) the vector representing the fractional overlap of

TeNT expression binary masks (displayed in Extended Data Fig. 11) and each atlas-

defined cortical area, and 2) the vector representing the change in performance

following modality switch (performance averaged over 3 consecutive sessions after

switch). This analysis was performed for each cortical area (examples for areas RL

and AL are shown in Extended Data Fig. 11f) and displayed as a correlation map in

Fig. 5e. For the reverse correlation approach, we computed the difference between

the average of all TeNT-injected mice GFP fluorescence binary masks and the one of

mice displaying impairment of cross-modal generalization ability (defined as mice

performing below 57.5% of correct trials on average in the 3 sessions following a rule-

preserving modality switch). This difference took null or negative values in areas where

TeNT expression did not affect the behavior and positive values in regions producing

behavioral impairment (i.e. regions where TeNT expression is enriched in the

impairment-triggered ensemble compared to the ensemble including all TeNT-

expressing mice).

1372

<u>Quantification of TeNT expression and ventral/dorsal grouping</u>

Wide-field fluorescence images of dorsal cortex displaying the pattern of expression

of TeNT-P2A-GFP in each injected mouse were first pre-processed to remove the

blood vessels from the surface. This was necessary to obtain a smooth, non-occluded

estimate of the spatial profile of GFP expression across the surface of the cortex. To

do so we first segmented the blood vessels with adaptive thresholding (using Matlab

function "adaptthresh") and then inpaint them by local interpolation (using Matlab

function "regionfill") (Fig. 5g,i and Extended Data Fig. 11a-d and 12b). Next, we

computed a median-subtracted map of df/f calculated with respect to a manually

annotated region of interest on the border of each window (selected to correspond to

a fluorescence profile stable around baseline level, usually at the farthest end of the

window with respect to the injection site). To compare images across different mice

and account for variations in imaging conditions, we normalized saturation by clipping

the values in each map to the minimum peak value observed across all mice. After

normalizing these resaturated df/f maps to 1 we computed their 0.90 contour (i.e. the

expression "patch") to define the binary TeNT-P2A-GFP expression mask to be used

in subsequent analyses to evaluate the spatial extent of toxin expression. The overlay

of these masks for all mice is displayed in Extended Data Fig. 11e. Next, TeNT-

expressing mice undergoing rule-preserving modality switch were subdivided in

"dorsal" and "ventral" subgroups by computing a d-prime measure (distance

normalized by patch size) with respect to the two multimodal visuo-tactile neuron

clusters found by k-means in the two-photon analysis of Fig. 4e. Neurons outside the

region of low d-prime with patch centroid falling respectively within AL/LI or A/RL

region were categorized as ventral and dorsal, respectively (Fig. 5g,i showed the

overlay of the binary masks together with expression centers as dots over the common

atlas). The same procedure was followed to validate the targeting of TeNT injections

to the dorsal region in mice undergoing rule-reversing modality switch (Extended Data

Fig. 12b).

1401

<u>Anatomical tracing analysis</u>

For each brain, M2 AChR immunostaining and tracer fluorescence images of each

slice were realigned by matching the radial blood vessel pattern. To do so, we applied

the "Normalize local contrast" function in ImageJ and then inverted the image to crop

low intensity pixels and isolate blood vessel holes. We then iteratively registered these

images across the stack in ImageJ using the "MultiStackReg" function with affine

transformation. The transformation could then be applied to the original stack. Slices

with good fluorescence signals were then averaged together. For anterograde tracing

data, we first applied an "Unsharp mask" function (radius: 20, mask weight: 0.8) and

pixel values were log-transformed to enhance axonal projections with respect to the

strong signal at the injection site where cell bodies are labelled. For retrograde tracing

data, a top hat filter (radius: 12) was applied to highlight the signal carried by CTB-

positive cells followed by the "Enhance contrast" function (Saturated pixels: 0.55%).

Realigned average slices for each mouse were then manually fitted to a projection of

the Allen mouse brain atlas using M2 AChR landmarks[58]. Individual mouse atlases

were then registered to common reference atlas. Centroid positions for injection

locations were determined by calculating the center of mass of the fluorescence

signals within manually outlined regions surrounding the injection sites. For

anterograde tracing data, fluorescence images corresponding to different injections

were normalized to the unit range per channel across brains and median-equalized

across channels within brains. For retrograde tracing data, fluorescence images

corresponding to different injections were rescaled to have the same peak value

across channels and brains. Center-of-mass coordinates for each injection in each

brain were projected on a line corresponding to the vertical meridian in the somatotopic

or retinotopic map at the targeted site. This yielded, for each injection, a vertical

coordinate as the center location. Lastly, we obtained an average projection map by

weighting the coordinate of each injection center with its normalized pixel intensity at

every pixel across the stack of fluorescence images (Fig. 3b,c,d and Extended Data

Fig. 7g). The maps obtained in this way represent for every pixel the preferred

projection origin (anterograde) or projection target (retrograde) location along the

vertical line, allowing for an understanding of the topographic mapping of inputs and

outputs associated with various regions. We also established a statistical metric to

assess the significance of projections at each location. This was accomplished by

comparing the median difference between two distributions: the local distribution of

normalized pixel intensities at each location and the distribution of normalized pixel

intensities from a specific frontal-lateral region of the slice. The latter region was

selected based on its lack of fluorescence signal, as determined by an initial visual

examination of the raw images. The center of the color scale used to plot the projection

maps corresponds to the median projection coordinate of the injection ensemble.

Average projection maps were compared to wide-field retinotopic and somatotopic

maps. This was achieved by aligning a selected portion (matching the cranial window

field-of-view) of the projection maps with the shared wide-field/two-photon atlas.

Subsequently, we calculated the Pearson coefficient of correlation between the two

images as a measure of similarity. Pixels exhibiting non-significant projection signal

were discarded for this analysis.

1447

<u>Optogenetic substitution experiments</u>

To test if the dorsal cortical area RL is sufficient for cross-modal generalization, we

used area-specific optogenetic activation as a substitute for visually evoked responses.

We expressed Channelrhodopsin-2 in the cortex by injecting

AAV1.CamKII0.4.Cre.SV40 into the whole RL area of Ai32 mice. This approach

induced a widespread expression of Channelrhodopsin-2 (ChR2) fused with

Enhanced Yellow Fluorescent Protein (eYFP). ChR2, a light-gated cation-selective

channel, generates depolarizing photocurrents in response to blue illumination,

thereby allowing for the excitation of cortical neurons[63]. To illuminate the cortical

surface of head-fixed mice during behavioral tasks, we built a behavioral rig equipped

with a Digital Light Processing (DLP) module (STAR 3.0 EVM Monochrome LED

Projection Modules, DLP6500, ViALUX GmbH). This module features a 1024 × 768

Texas Instruments micro-mirror chip and a 460 nm blue LED, integrated with an

inverted tandem-lens configuration macroscope. The optical tower assembly

comprised two lenses configured in an L-shaped pathway, separated by a dichroic

mirror. Light from the macroscope enters through the projection lens (L1; 150 mm smc

Pentax-A 645 lens) at the rear, reaching a dichroic mirror (640 nm cutoff wavelength;

60 × 60 mm; Semrock) positioned at a 45° angle. This mirror reflects the light towards

the front end of the objective lens (L2; 50 mm Nikon NIKKOR lens), which projects the

image onto the mouse's cortex. Stimulation intensity was adjusted, for each mouse,

to ensure detection while minimizing heat dissipation in the cortex: this was achieved

by operating the stimulation LED within a power range that produced an irradiance

between 3.5mW/mm^2 and 11.5mW/mm^2. The refresh rate was set at 100Hz with a

projected pixel size measuring approximately 6.5 μm. To prevent mice from directly

perceiving the blue light used for optogenetic stimulations, we enclosed the optical

column and the mouse skull with an opaque sleeve. To optogenetically stimulate the

cortex with task relevant information concerning the vertical position of the stimulus,

we shaped the blue light projection in every trial to match the top- or bottom- encoding

subregions of the RL map. Light patterns were obtained by thresholding the average

retinotopy movies within RL boundaries as defined by the fitted atlas. To effectively

drive neuronal activity with blue light, we pulsed illumination at 30 Hz[63]. We used a

pulsing pattern over the top or bottom encoding part of RL lasting one second within

the response window. Before switching to the full optogenetic substitution task, mice

that had achieved stable expert performance for three sessions in a row in the original

tactile task first underwent 1 to 3 optogenetic habituation sessions. During these

sessions, they were exposed solely to the Go light pattern (with a distribution of 80%

Go trials and 20% Catch trials with light stimulations outside the cranial window). As

soon as mice displayed a consistent licking response to optogenetic stimulations they

were transitioned to the full task (with a distribution of 30% Go, 50% No go trials and

20% Catch). Real-time projection control was achieved integrating ALP Vialux API

with the custom-made Matlab graphical user interface (GUI) used to run the behavioral

tasks and monitor mouse performance. Analysis of task performance during

optogenetic substitution sessions was carried out exactly as described in the

"behavioral analysis" section above.

1492

<u>Network model</u>

Code will be made available at: https://github.com/comp-neural-circuits. We

constructed three recurrent excitatory (E) / inhibitory (I) rate networks representative

of the areas V1, S1, and RL (Fig. 7a). We abstracted the decision process to a pair of

readout units which in turn drove an integrator through a drift-diffusion process.

Recurrent rates of the units in each region evolved according to

$$\tau_i \dot{r}_i = -r_i + f\left(W_{ij}r_j + I(t) + \sigma_r x_i(t)\right)$$

where W_{ij} are the connection weights from unit j to unit i, $I(t)$ is a source of external

input to V1 and S1 cells only, and x_i is noise independent to each unit, evolving

according to $dx_i = -x_i dt + \sigma_x dW$ where dW is a Wiener process with scaling factor σ_x.

We used a saturating transfer function, given by:

$$f(x) = \beta \left\lfloor \frac{1}{1 + \exp(-(x - x_0)/\rho)} - \frac{1}{1 + \exp(x_0/\rho)} \right\rfloor_+ .$$

where $\lfloor \cdot \rfloor_+$ denotes a threshold at zero. Each region was divided into two subnetworks

(corresponding to top- and bottom-preferring cells) of N_E excitatory neurons and $N_I =$

N_E inhibitory neurons each. These subnetworks were anatomically segregated such

that weights were partitioned into top- and bottom-preferring units which did not

interact (i.e. $W_{ij} = 0$ for i top and j bottom, or i bottom and j top; see Fig. 7). Within a

subnetwork, neurons were all-to-all connected. Projections between regions

respected subnetwork boundaries and were limited to E neurons. Weights of

feedforward projection connections (i.e. V1 or S1 to RL) were scaled slightly up relative

to the within-region weights, and feedback projections (i.e. RL to V1 or S1) were scaled

slightly down relative to within-region weights (see Table). To model experiments with

TeNT-based silencing of RL we removed all incoming and outgoing connections to the

model RL units. Simulations were subdivided into multiple trials of length 0.4 s, defined

as a period of no external stimulation t_{isi} = 0.2 s, i.e. $I(t)$ = 0, together with presentation

of a stimulus $I(t)$ > 0 to the V1 or S1 network (mimicking the visual or tactile tasks,

respectively) for time t_{stim}= 0.2 s which was exclusive to either top- or bottom-preferring

units in one of the two sensory regions (V1 or S1). The stimulus was modeled as a

step function, i.e. $I(t)$ = c_s, where c_s is a constant. The stimulus order was randomized

for each simulation. In order to model rewards in our framework we considered two

abstract readout units, z_1 and z_2, corresponding to Go and No go responses,

respectively. The activity of these units was a weighted linear sum of activity from E

neurons within the recurrent network $z_i(t) = \alpha W_{ij} r_j$ where α is a scaling constant. The

network's decision was determined by producing a decision variable δ_i for each z_i by

integrating z_i from stimulus onset in the presence of independent noise:

$d\delta_i = z_i dt + \sigma dW$

where dW is a Wiener process with scaling factor σ. Decisions were instantiated once

a fixed threshold θ_0 was reached by either z_1 or z_2 resulting in the specific decision

encoded by that readout unit; if neither z_1 nor z_2 crossed the threshold before the end

of the stimulus period, the action was considered No go. Once a decision was reached

the response z_i was modified according to $\tilde{z}_i(t) = z_i(t) + c_r$, until the end of the

stimulus interval. If the decision was correct relative to the presented stimulus, then

$c_r > 0$, otherwise $c_r < 0$. $|c_r|$ was defined as the difference between the current z_i

and the last value of z_i before the stimulus period. Only projection weights from each

region (V1, S1, and RL) to the output variables z_i were plastic. Weights evolved

according to a Hebbian rule:

$\dot{W}_{ij} = \gamma z_i r_j$

where $i \in \{1, 2\}$, $j \in \{1, ..., N_E\}$, and γ is the learning rate. Weights were bounded

between 0 and $W_{max} = 1$ and normalized after each update to the initialized mean $\overline{W_z}$

within a region. Weights were initialized as a Gaussian centered at mean $\overline{W_z}$ = 0.5 with

standard deviation $\sigma_z = 0.1$. All model simulations were run for 100 trials before and

after a switch. To compare model performance with the animal behavior we recorded

the trial outcome for each stimulus presentation. We then computed a performance

metric and "action" probabilities conditioned on the stimulus in a sliding window of 15

trials. In particular, if we define s_1 as input to the bottom-preferring units and s_2 as input

to the top-preferring units, within each window we calculated P($z_1|s_1$) and P($z_1|s_2$) as

the conditional action probabilities. The performance accuracy was computed as

P(correct) with correct defined as $z_1 > z_2 \mid s_1$ and $z_2 > z_1 \mid s_2$ in the rule-preserving

switch condition and $z_1 > z_2 \mid s_2$ and $z_2 > z_1 \mid s_1$ in the rule-reversing switch condition.

We quantified the number of trials to reach criterion across four different switch

conditions: rule-preserving and rule-reversing in the full model; rule-preserving and

rule-reversing with RL silenced. We set a criterion value of 75% performance accuracy

for each trial condition to quantify the number of trials needed for the model to re-learn

after a switch. Because the mean performance accuracy increased approximately

monotonically while response conditions were held constant, we computed the first

time at which a given model run reached the criterion value as the time to relearn. If a

model run did not reach the criterion within 100 trials, we omitted it from this analysis.

1560

Variable	Value
W_{EE}	0.16
W_{EI}	0.2
W_{IE}	0.16
W_{II}	0.18
c_s	30
ρ	15
x_0	20
β	30
α	$1/(6N_E)$
α (lesion)	$1/(2N_E)$
γ	4e-7
σ	0.5
σ_x	$\sqrt{2}$
σ_r	0.75
τ_E	20
τ_I	10
θ_0	80

1561

1562

Quantification and statistical analysis

1563

Statistical details of experiments and analysis are described in figure legends and in

the main text. Details include statistical tests used, sample type and size as well as

definition of bar plots and error bars. In figure legends, standard error of the mean

(S.E.M.) is specified when plotted as error bars. Paired or unpaired t-tests were used

to assess significance of mean comparisons (implemented by Matlab functions "ttest"

and "ttest2", respectively). Normality tests were not performed systematically but

individual data points were plotted to visualize distributions. Wilcoxon signed rank test

were used to assess significance in paired median comparisons (implemented by

Matlab function "signrank"). Wilcoxon rank sum test was used for unpaired median

comparisons (implemented by Matlab function "ranksum"). Pearson coefficient of

correlation was used to compute correlations between two conditions (implemented

by Matlab function "corr"). Across all fits reported uncertainties (i.e. confidence

intervals) for best-fit parameter values were extracted from fit covariance matrices

(fitting was performed using Matlab function "nlinfit"). In order to assess the statistical

significance of multisensory modulation at each time bin, for each visuo-tactile

stimulus condition, and for each responsive neuron, we performed a bootstrap-t

procedure[64] with 2000 outer resamplings and 25 inner resamplings (over trials) in

order to obtain p values for the difference from zero of MI values.

1582

1583 Data and Code Availability

1584 The complete data set and Matlab analysis code will be made freely available at the

1585 CERN database, Zenodo.

1586

1587

1588

Acknowledgements

We thank Denis Jabaudon, Alan Carleton, Daniel Huber and members of the El-Boustani laboratory for discussions. This work was supported by the Swiss National Science Foundation: PCEFP3_181070 (S.E.-B.).

Author Information

M.G., G.M. and S.E.-B. conceived experiments. M.G. performed surgeries, viral infections, wide-field and two-photon imaging, behavioral experiments, and post-mortem analysis. M.G., G.M. and C.G.F. performed behavioral experiments with optogenetics. G.M. coded acquisition software and performed data analysis. M.P.G. and J.G. conceived the network model and performed simulations. M.G., G.M. and S.E.-B. wrote the manuscript.

Ethics Declarations

Authors declare no competing interests.

Corresponding author

Correspondence and requests for materials should be addressed to Sami El-Boustani (sami.el-boustani@unige.ch).

Supplementary Information

 Extended Data Fig. 1-14

Extended Data Figure 1

Extended Data Fig. 1. Cross-modal generalization in mice across different conditions.

a, Schematic of behavioral paradigm where switches occur between a tactile task and a visual task. Bottom tactile stimuli were rewarded upon licking in the first task. **b,** Left: Average task performance and conditional lick probabilities across sessions for mice (N=5 mice) undergoing a rule-preserving modality switch (dashed vertical line). Shaded areas and color code as in Fig. 1e. Right: detection (purple) and discrimination (green) performance distribution the session before and after the switch (two-sided paired t-test comparing days, *p<0.05, **p<0.01, ***p<0.001, N.S. Not significant). Performances are also tested against chance level (two-sided t-test, *p<0.05, **p<0.01, ***p<0.001, Blank: Not significant). **c,** Same as **b** but for a rule-reversing modality switch. **d,** Schematic of behavioral paradigm where switches occur between an auditory discrimination task with two pure tones (6 kHz and 12 kHz) and a visual task. The 6 kHz tone is always associated with a water reward. **e,** Same as **b** for an auditory to visual modality switch, where the bottom visual stimulus is associated to a reward. **f,** Same as **e** for an auditory to visual modality switch, where the top visual stimulus is associated to a reward. **g,** Schematic of behavioral paradigm where switches occur between a visual task and a tactile task with the top visual stimulus being the rewarded one. **h,** Same as **b** for mice undergoing a rule-preserving switch (N=5 mice). **i,** Same as **h** but for rule-reversing modality switches. **j,** Same as **g** with the bottom visual stimulus being the rewarded one. **k,** Same as **h** but for rule-preserving modality switch. **l,** Same as in **k** but for a rule-reversing modality switch. For cases with spatial rule-reversing modality switch or in absence of spatial prior (i.e. **c**,**e**,**f**,**i**,**l**), behavioral curves are shown for 6 consecutive sessions following the switch. We observed a small but significant drop in performance for switches from visual to tactile modalities under rule-preserving conditions (**h**,**k**) that was absent in the tactile to visual switches (**b**). This asymmetry might arise from generally higher initial performances in the visual task, the distinct nature of sensory representation (discrete for whiskers and continuous for visual inputs), or variations in circuit properties.

Extended Data Figure 2

1640

Extended Data Fig. 2. Performance of mice at the whisker discrimination task with and without whiskers in the piezo stimulators.

a, Conditional lick probabilities and performance during the whisker discrimination task averaged across expert mice (N=5 mice). Sessions were normalized between beginning (0%) and end (100%) before the average was performed. Color code is the same as in Fig. 1. During these sessions, mice performed the task for approximately one-third of the trials with whiskers present in the capillary tubes used to deliver tactile stimulations. Subsequently, the whiskers were removed while maintaining the tube positions, and the task continued for an equivalent duration. Finally, the whiskers were reinserted to confirm that mice were still motivated to perform the task. **b,** Comparison of detection (purple) and discrimination (green) performance during the 3 different phases of the sessions described in **a**. Comparisons are performed between consecutive phases (paired two-sided t-test, ***p<0.001, **p<0.01, *p<0.05) and with chance level (two-sided t-test with p=0.188 for detection and p=0.710 for discrimination during the Out phase). Mice were not able to perform the task in absence of direct whisker stimulations.

1656

Extended Data Figure 3

1657

Extended Data Fig. 3. Relearning the whisker discrimination task following a reversal of the spatial rule within the same modality.

a, Schematic of the behavioral paradigm for rule-reversing switch within the tactile modality. Once mice became experts in the whisker discrimination task, the spatial rule of the task was reversed. **b,** Left: Average task performance across sessions for mice exposed to the reversed reward association (N=7 mice). Shaded area: S.E.M. Vertical dashed line indicates the transition from one rule to the other. Right: detection (purple) and discrimination (green) performance distribution the session before and after the switch (paired two-sided t-test comparing days, *p<0.05, **p<0.01, ***p<0.001, N.S. Not significant). Performances are also tested against chance level (two-sided t-test, *p<0.05, **p<0.01, ***p<0.001, Blank: Not significant). **c,** Comparison of performance change relative to chance level after rule switch across sensory modalities (i.e. tactile to visual) or within the whisker task (N=10 mice for visuo-tactile and N=7 mice for tactile only, unpaired two-sided t-test, **p=0.0056). Mice maintained their responses to whisker stimulations significantly above chance level, yet their performance on the discrimination task fell significantly below chance (*p=0.0221). This suggests a reduced adaptability to a reversal of the spatial rule within the same modality, potentially due to a stronger habitual response linked to the previously rewarded whisker or an entrenched bias regarding the saliency of whisker stimulations.

1676

Extended Data Figure 4

1677

> **Extended Data Fig. 4. Performance level is maintained after switches from the whisker discrimination task to the visual task.**
>
> Comparison between performance in the whisker task the session before the switch and the performance in the visual task the session following the switch. This includes mice that experienced the switch, showcasing varying performance levels and rewarding either the top whisker (blue) or the bottom whisker (red). Performances before and after the switch were correlated (N=18 mice, Pearson coefficient: 0.68, p=0.0018) and close to the unity dashed line. Shaded area: 95% confidence interval. Mice exhibit consistent discrimination performance following modality switch, suggesting that cross-modal generalization may correspond to the initial sensorimotor learning level, or that shared influences, such as thirst-related motivation, impact both tasks equally.

Extended Data Figure 5

1689

1690

Extended Data Fig. 5. Average whisker and retinotopic maps with examples of single mouse visuo-tactile functional maps obtained in wide-field imaging.

a, Schematic of the protocols used to measure single whisker response maps (top) and retinotopic maps (bottom) in transgenic mice expressing GCaMP6f in layer 2/3 during wide-field imaging through a cranial window. **b,** Response map averaged across mice (N=41 mice) for C2 whisker stimulations. Projection of the Allen Mouse Brain Atlas is overlaid on top with areas names for reference. **c,** Same as in **b** for B2 whisker stimulations. **d,** Elevation phase map averaged over mice (N=50 mice). Pixels with low response amplitude are displayed in white. **e,** Azimuth phase map average over the same mice as in **d**. **f,** Retinotopic sign map computed based on the elevation and azimuth phase maps of **d** and **e**. **g,** Representative examples of visuo-tactile functional maps obtained from sparse noise protocol responses from 3 different mice (rows). From left to right: 1. Whisker preference map used to describe somatotopy in the vertical space with red colors indicating preference for the bottom whisker and blue colors indicating preference for the top whisker. 2. Visual position preference map used to describe retinotopy in vertical space. 3. Modality preference map describing the dominant modality for each pixel. 4. Spatial coherence map between retinotopic and somatotopic maps. 5. Multisensory modulation map indicating areas with responses enhancement or suppression for visuo-tactile stimuli.

Synchronous visuo-tactile stimulations

Delayed tactile stimulations (0.15 s)

Extended Data Figure 6

1710

Extended Data Fig. 6. Comparison of cortical response dynamics for synchronous or delayed visuo-tactile stimulations.

a, Average activation time course for each visually responsive area following stimulus onset (N=29 mice). Areas are ordered based on response onsets. White dashed line indicates stimulus onset. **b,** Same as **a** for responses in cortical areas evoked by whisker stimuli. **c,** Time course of the modality preference index across cortical areas with visuo-tactile responses. **d,** Average response time course in the primary visual cortex (V1) and the whisker primary somatosensory cortex (S1) highlighting an offset of about 0.15 s between cortical responses evoked by the two modalities. Colored dashed lines indicate response latency computed as the time point where the response curve reaches 80% of its peak amplitude. **e,** Distribution of response latencies for cortical areas with evoked visual (orange) or whisker tactile (magenta) responses. Colored dashed lines indicate the average latencies across areas. **f,** Time course of the multisensory modulation index across cortical areas with visuo-tactile responses. **g-l** Same as **a-f** for tactile stimuli presented with a delay of 0.15 s with respect to the visual stimuli to match response latencies in the cortical areas. Multisensory enhancement was observed when stimuli were synchronized or delayed though they were stronger when cortical responses had comparable latencies.

a Somatotopic mapping
b B2 whisker response map   C2 whisker response map
c Cranial window after injection

d Retinotopic mapping
e Elevation phase map   Azimuth phase map
f Cranial window after injection

g CTB retrograde markers in V1
h
i Dorsal-projecting neurons / Ventral-projecting neurons
j

Extended Data Figure 7

Extended Data Fig. 7. Functional mapping of vertical positions in the visual or whisker somatosensory cortex for anatomical tracing.

a, Schematic of the protocol for mapping whisker-responding areas in the dorsal cortex. Single whiskers are stimulated repeatedly during intrinsic optical signal imaging. **b,** Change in red light reflectance on the surface of the cortex measured after averaging several trials of whisker stimulations relative to baseline. The left image reveals the barrel location corresponding to B2 whisker stimulations and the right plot for C2 whisker stimulations. **c,** Image of the cranial window several weeks after injection of viral vectors to express GFP and tdTomato in two distinct parts of the whisker somatotopic map. Locations of B2 and C2 barrels in S1 defined in **b** are indicated with circles overlaid on top of the fluorescent signal where injections were performed. **d,** Schematic of the protocol for retinotopic mapping in areas of the dorsal cortex. Horizontal or vertical bars drift across the screen and they are textured with a flickering checkerboard pattern to elicit strong cortical responses at different phases corresponding to the retinotopic preference. **e,** Retinotopic phase maps for elevation and azimuth on the dorsal cortex. These maps are then used to identify a gradient across vertical dimension in an iso-profile along the azimuth dimension. Two retinotopically distinct locations along the vertical axis are identified and labeled for viral injection. **f,** Image of the cranial window four weeks after injection of viral vectors to express GFP and tdTomato in two distinct parts of the primary visual cortex retinotopic map. Locations where stereotaxic injections were performed corresponding to **e** are indicated with circles overlaid on top of the fluorescent signals. **g,** Cholera Toxin B subunit (CTB) conjugated to different Alexa dyes was injected in different locations of the retinotopic map in V1 (circles). After perfusing the brain and flattening the cortex, a map was reconstructed over the dorsal cortex combining all injection sites to reconstruct areas where neurons project to the region of V1 encoding stimuli on the top (blue) or bottom (red) part of the visual field. All cortices with CTB labels were registered to a common atlas before performing the analysis (5 injection sites from N=3 mice). A similar analysis was performed previously with coronal slices[29]. **h,** Flattened cortex with injections of CTB conjugated with two different dyes (Alexa555 and Alexa647) in the dorsal and ventral visuo-tactile area, respectively. Immunohistochemical localization of M2 AChR is used to identify V1 and the barrels in S1 for atlas registration. **I,** Example image obtained with confocal microscopy where dorsal-projecting neurons (magenta) and ventral-projecting neurons (green) are intermingled in V1. **j,** Fraction of neurons that projected to the ventral area exclusively (green), to the dorsal area exclusively (magenta) or to both (purple) (N=3 mice, unpaired two-sided t-test, **p=1.8×10^{-3}, n=230 neurons for ventral, n=277 neurons for dorsal and n=26 neurons for both out of 26 fields-of-view). Neurons that project to the dorsal and ventral visuo-tactile areas display minimal overlap, suggesting they may process distinct types of visual information.

Extended Data Figure 8

1767

Extended Data Fig. 8. Coverage of large portions of the cranial window for characterizing visuo-tactile responses at single-cell level using two-photon microscopy.

a-e, Functional maps are obtained and aligned with the Allen Mouse Brain Atlas for each individual mouse, ensuring precise localization of neurons within their respective areas. Subsequently, all maps are realigned to a common reference atlas to reconstruct wide-field maps. Visual stimulation protocols are used to obtain phase maps for elevation (**a**) and azimuth (**b**) to compute the retinotopic sign map (**c**). Additionally, whisker stimulations provide response maps for C2 (**d**) and B2 (**e**) whisker stimulations. **f,** Image obtained with wide-field imaging where blood vessels are well distinguished on the surface of the cortex. The atlas fitted with functional maps is overlaid on top in purple. For this mouse, seven fields-of-view were imaged using two-photon microscopy and positioned on the cortical surface by matching blood vessel patterns. **g,** Coverage of visual and tactile areas of the dorsal cortex with two-photon imaging across mice. Using the common atlas and information about all neuron's location, we can recreate a large-scale population using single-cell information. We could collect N=272 fields-of-view across N=25 mice. **h,** Fraction of neurons imaged with two-photon microscopy that were responsive to at least one stimulus condition. The areas of high responsivity are in accordance with wide-field imaging.

Extended Data Figure 9

Extended Data Fig. 9. Functional maps correspondence between wide-field and two-photon calcium imaging.

a, Left: Distribution of neurons imaged with two-photon microscopy with significant whisker tactile responses. Neurons are displayed on the common atlas and are color-coded based on their whisker preference (n=790 neurons). Middle: Large-scale functional map reconstructed from single neuron whisker preferences. Right: Somatotopic map for the whisker preference obtained with wide-field imaging. Pearson coefficient of correlation between the wide-field map and the two-photon reconstructed map: 0.847. **b,** Same as in **a** for neurons with significant visual responses (n=1,816 neurons, Pearson coefficient of correlation: 0.651). **c,** Same as in **a** displaying modality preference for all neurons with significant whisker tactile or visual responses (n=2,160 neurons, Pearson coefficient of correlation: 0.795). **d,** Same as in **a** displaying multisensory enhancement for all neurons with significant multisensory modulation. For this analysis, the multisensory modulation is computed as the variation between the most suppressed and most enhanced response across conditions for all neurons (n=2,497 neurons, Pearson coefficient of correlation: 0.4458).

Extended Data Figure 10

Extended Data Fig. 10. Distribution of functional cell-types across the dorsal cortex and their impact on cross-modal generalization.

a, Distribution of significantly responsive neurons across the dorsal surface of the cortex (from N=25 mice). Common atlas overlaid for reference. Color code indicates functional cell-type of each neuron (n=2,703 neurons). Neurons were classified in four categories: tactile neurons (magenta) responded only to whisker stimulations (n=567); visual neurons (orange) responded only to visual stimulations (n=1,593); visuo-tactile neurons (dark green) responded to both visual and whisker stimulations (n=213); gated neurons (light green) responded only to concomitant visuo-tactile stimulations (n=319). **b,** Example visuo-tactile receptive field for a visual neuron (top), a tactile neuron (middle) and a gated neuron (bottom). Color code is identical to Fig. 4b. Note that the gated neuron responds only when visual and whisker stimulations are presented together in the top location (spatially congruent condition). **c,** Decoding accuracy for the tactile task and the visual task as a function of the pool size. A Bayesian decoder was trained to discriminate the location of whisker stimulation (magenta) based on the neuronal responses of a pool of neurons. The same decoder was then used to decode the location of visual stimuli (orange), mimicking the modality switch experienced by mice in the behavioral experiment. Error bars: S.E.M. Only visuo-tactile neurons were able to achieve good generalization performance across modalities because of their functional properties. Tactile neurons could only afford good decoding performance in the whisker-based task but could not maintain it when switching to the visual task. Performance chance level is indicated with a gray dashed line at 50%. **d,** Comparison of single-neuron decoding accuracy distributions on tactile responses (magenta) and generalization to the visual ones (orange) for each functional cell-type class (paired two-sided t-test comparing tactile and visual task, *p<0.05, **p<0.01, ***p<0.001, N.S. Not significant). Decoding accuracies are also tested against chance level (two-sided t-test, *p<0.05, **p<0.01, ***p<0.001, N.S.: Not significant). Error bars: S.E.M.

Extended Data Figure 11

1832

Extended Data Fig. 11. Cross-modal generalization impairment following rule-preserving switches in relation to TeNT expression in different associative areas of the dorsal cortex.

a, Left: Example of TeNT-P2A-GFP expression pattern over the cranial window and corresponding estimate of expression coverage after blood vessels subtraction. Right: Effect of cortical silencing on cross-modal generalization from the tactile task to the visual task while preserving the spatial rule of the task. The discrimination performance dropped at chance level after the switch. **b-d,** Same as in **a** but for other example mice with cortical injections at various locations and the resulting impact on cross-modal generalization performance. Note that equal volumes of viral vectors were injected in each mouse; however, some expression patterns appear truncated at the edge of the window. **e,** Overall coverage across all experiments performed with this protocol. **f,** Example of correlation between the level of TeNT-P2A-GFP overlaps with a specific area (here RL or AL) and the corresponding drop of discrimination performance after switch. To obtain a more robust effect for the correlation, we computed the performance changes over 3 consecutive sessions after switch (N=22 mice, Pearson coefficient: -0.43, p=0.047 for RL; Pearson coefficient: -0.16, p=0.48 for AL).

Extended Data Figure 12

1850

Extended Data Fig. 12. Relearning after rule-reversing modality switch in mice expressing TeNT-P2A-GFP in the dorsal area RL.

a, Schematic of behavioral paradigm where the modality switches occur between a tactile task and a visual task in mice expressing TeNT-P2A-GFP in RL. The bottom whisker stimulus is associated with the reward initially while the top visual stimulus becomes associated to rewards after the modality switch. **b,** Average TeNT-P2A-GFP coverage in all mice where dorsal neurons were silenced (N=6 mice). **c,** Left: Average task performance and conditional lick probabilities across sessions for mice population in **b** (N=6 mice) switching from the tactile task to the visual task with a spatial rule reversal. Shaded area: S.E.M. Black dashed line indicates the switch between modalities. Right: detection (purple) and discrimination (green) performance distribution the session before and after the switch (paired two-sided t-test comparing days, *p<0.05, **p<0.01, ***p<0.001, N.S. Not significant). Performances are also tested against chance level (two-sided t-test, *p<0.05, **p<0.01, ***p<0.001, Blank: Not significant). Contrary to mice with intact cortical activity, mice with silenced RL area relearned the visual task faster after the rule-reversing modality switch, indicating a lack of prior on the spatial rule in these mice despite their previous experience with the tactile task.

Extended Data Figure 13

1868

Extended Data Fig. 13. Model performance on a rule-reversing modality switch.

a, Model schematic. The model consists of three networks for areas S1, V1, and RL (here only one network is shown for simplicity) that comprised excitatory and inhibitory units tuned to either bottom or top stimuli. Dashed lines denote plastic synapses, and their colors are consistent throughout the figure. The excitatory units in all networks project to the two decision variables z_1 and z_2. S1 was initially stimulated, and Bottom stimuli mapped to "Go" action. After the switch, V1 was stimulated, and Top stimuli were mapped to "Go" action. **b,** Weights shown for different time points (corresponding to the dashed lines in panels **c** and **d** as a function of the neuronal index. $W_{z_i,r_j}(t)$ denotes the weights from sensory neuron j to decision variable i at time t. t_0: initial timepoint; t_{sw}: time of the switch; t_{end}: final timepoint after the switch. At the time of the switch (t_{sw}), all regions have the same organization of weights such that the correct output is produced (including in V1, due to the reverberation of network dynamics). After the switch, areas RL and V1 have reorganized to drive the correct output. **c,** Subsample of weights from the Bottom-preferring units to z_1 (brown) and z_2 (teal) as a function of time in a simulation. **d,** Same as in **c** for Top-preferring units.

Extended Data Figure 14

1884

	Extended Data Fig. 14. Maximum average performance reached following modality switch across different conditions.
1885	**Extended Data Fig. 14. Maximum average performance reached following modality**
1886	**switch across different conditions.**
1887	Left: Maximum average performance reached in the discrimination task following the modality
1888	switch. Mice numbers are indicated for each group at the bottom of the bar. Mice that did not
1889	reach the expert criteria in Figure 7F are discarded from this plot. Error bars: S.E.M. Right:
1890	Surprise matrix computed from pairwise unpaired two-sided t-test between conditions. A
1891	hierarchical clustering based on cosine similarity was used to group conditions based on
1892	surprise values. Mice were clustered with respect to the maximum performance reached
1893	based on the nature of sensory stimulation used after switch (visual, auditory or optogenetic).