RBM39 shapes innate immunity through transcriptional and splicing control of key factors of the interferon response

Authors: Teng-Feng Li[1], Paul Rothhaar[1], Arthur Lang[1], Oliver Grünvogel[1], Ombretta Colasanti[1], Santa Mariela Olivera Ugarte[1], Jannik Traut[1], Antonio Piras[2], Nelson Acosta-Rivero[3], Vladimir Gonçalves Magalhães[4], Emely Springer[5], Andreas Betz[1], Hao-En Huang[1], Jeongbin Park[6], Ruiyue Qiu[7], Gnimah Eva Gnouamozi[3], Ann-Kathrin Mehnert[8], Viet Loan Dao Thi[8,9], Stephan Urban[3,9], Martina Muckenthaler[7], Matthias Schlesner[6,10], Dirk Wohlleber[5], Marco Binder[4], Ralf Bartenschlager[3,4,9], Andreas Pichlmair[2,11], Volker Lohmann[1,9]*

Affiliations:

[1] Department of Infectious Diseases, Molecular Virology, Section Virus-Host-Interactions, Medical Faculty Heidelberg, Heidelberg University, Heidelberg, Germany.

[2] Institute of Virology, School of Medicine, Technical University of Munich, Munich, Germany.

[3] Department of Infectious Diseases, Molecular Virology, Medical Faculty Heidelberg, Heidelberg University, Heidelberg, Germany.

[4] Division of Virus-Associated Carcinogenesis, German Cancer Research Center (DKFZ), Heidelberg, Germany.

[5] Institute of Molecular Immunology, University Hospital Klinikum rechts der Isar, Technical University of Munich, Munich, Germany.

[6] Bioinformatics and Omics Data Analysis, German Cancer Research Center (DKFZ), Heidelberg, Germany.

[7] Heidelberg University, Medical Faculty, Department of Pediatric Oncology, Hematology, Immunology and Pneumology, Heidelberg, Germany.

[8] Department of Infectious Diseases, Virology, Medical Faculty Heidelberg, Heidelberg University, Heidelberg, Germany.

[9] German Center for Infection Research (DZIF), Heidelberg Partner Site, Heidelberg, Germany.

[10] Biomedical Informatics, Data Mining and Data Analytics, Faculty of Applied Computer Science and Medical Faculty, University of Augsburg, Augsburg, Germany

[11] German Center for Infection Research (DZIF), Munich Partner Site, Munich, Germany.

*Corresponding author's email: Volker.Lohmann@med.uni-heidelberg.de

Author Contributions:
Conceptualization: VL, TL, OG, JT
Methodology: VL, OG, OC, NAR, GEG, AKM
Investigation: TL, AL, JT, SMOU, PR, VGM, APir, ES, RQ, AB, HEH, JP,
Visualization: TL, PR, OG, JT,
Funding acquisition: VL
Project administration: VL
Supervision: VL, AP, RB, MB, DW, MS, MM, SU, VLDT
Writing – original draft: TL, OC, VL
Writing – review & editing: all authors

Competing Interest Statement: Authors declare that they have no competing interests.

Classification: Immunology

Key Words: RBM39, IRF3, IFNs, splicing

This PDF file includes:
Main Text
Figures 1 to 7
Supplementary Materials and Mthods

Abstract

RNA-binding motif protein 39 (RBM39) is an RNA-binding protein involved in tumorigenesis, cell
metabolism, and development. Here, we performed a genome-wide CRISPR/Cas9 screen in two
liver-derived cell lines and identified RBM39 as a regulator of cell intrinsic innate immune responses.
The knockdown of *RBM39* or the treatment with Indisulam, an aryl sulfonamide drug targeting RBM39
for proteasomal degradation, strongly reduced the induction of interferon-stimulated genes (ISGs) in
response to double-stranded RNA (dsRNA) or viral infections upon sensing by toll-like receptor 3
(TLR3) or cytosolic RIG-I-like receptors. RNA sequencing (seq) and mass spectrometry identified that
transcription and/or splicing of the key pathway components IRF3, RIG-I, and MDA5 were affected by
RBM39 depletion. *RBM39* knockdown further restrained type I and type III IFN pathways, by reducing
expression of the type I IFN receptor subunit interferon alpha and beta receptor subunit 2 (IFNAR2),
type III IFN receptor subunit interleukin 10 receptor subunit beta (IL-10RB) and transcription factor
signal transducer and activator of transcription (STAT) 1 and 2. RBM39 overall orchestrates innate
immunity by regulating basal expression of key factors of the interferon response via transcription
and/or alternative splicing.

Significance

The function of RBM39 in tumorigenesis has been investigated intensively in the last decade, but its
immunological role is still largely unknown. In our study, we identified RBM39 as a regulatory factor of
cell intrinsic signaling via a CRISPR/Cas9 screen. Depletion of RBM39 impairs TLR3, RIG-I/MDA5,
and IFN pathways, and thus attenuates innate immune responses. Our omics analysis revealed that
RBM39 governs the basal expression of several key factors within these pathways, such as RNA
sensors RIG-I and MDA5, type I/III receptors, transcription factors IRF3, STAT1 and STAT2, via its
transcriptional and splicing function. Therefore, RBM39 might be a therapeutic target to modulate
innate immunity, e.g. in the context of autoimmune disorders.

Introduction

The cell intrinsic innate immune system belongs to the first line of host defense mechanisms
protecting against invading pathogens. When infected, a limited number of host Pattern Recognition
Receptors (PRRs) plays a vital role in recognizing several Pathogen-associated molecular patterns
(PAMPs), which activate innate immune responses through complex signaling cascades to eliminate
infections. Viral components, including viral DNA (*1, 2*), single-stranded RNA (ssRNA) (*3, 4*), double-
stranded RNA (dsRNA) (*5, 6*) and virus-encoded proteins (*7*), are sensed by distinct PRRs. The
recognition triggers the cell intrinsic innate immune response in infected cells. For instance, dsRNA, a
replication intermediate of many RNA viruses, is sensed by cytosolic retinoic acid-inducible gene
(RIG)-I-like receptors (RLRs, RIG-I and MDA5) (*5, 6, 8*) and endosomal Toll-like receptor 3 (TLR3)
(*9*). These PRRs then recruit the adaptor proteins mitochondrial antiviral signaling protein (MAVS)
and TIR domain containing adaptor molecule 1 (TRIF), respectively, activating a chain of kinases, E3
ubiquitin ligases and E2 ubiquitin-conjugating enzymes. This leads to nuclear factor kappa B subunit
1 (NF-κB) pathway activation and to production of pro-inflammatory cytokines or phosphorylating
IRF3, leading to its dimerization and translocation from cytoplasm into nucleus. Subsequently, IRF3
binds to responsive promoters and induces transcription of type I/type III interferons (IFNs) and
interferon stimulated genes (ISGs). IFNs act in an auto- and paracrine manner, binding to their
receptors, Interferon alpha and beta receptor (IFNAR) and Interferon lambda receptor (IFNLR), on the
cell surface, respectively, but signaling via a common cascade upon phosphorylation of signal
transducer and activator of transcription 1 and 2 (STAT1 and STAT2). Eventually, STATs form hetero-
or homodimers and translocate into nucleus to accelerate or initiate ISG expression in infected and
neighboring cells, respectively.

RNA-binding motif protein 39 (RBM39) (also known as HCC1.4 or CAPERα) was initially identified as
an autoantigen in a hepatocellular carcinoma patient (*10*) and has been found to be upregulated in
multiple cancer cells (*11-13*). It shares high similarity with the splicing factor U2AF 65 kDa subunit

(U2AF65), and contains an N-terminal serine/arginine (SR)-rich domain followed by two RNA-
recognition motifs (RRM) and a third C-terminal noncanonical RRM belonging to the U2AF homology
motif (UHM) family, which mediates the interaction with UHM-ligand motif (ULM) of U2AF65 (*14*).
Previous studies have demonstrated that RBM39 functions as a coactivator of activating protein-1
(AP-1), Estrogen Receptor 1(ERα) and Estrogen Receptor 2 (ERβ) (*15-17*) and that it is involved in
multiple biological processes, including cell cycle control (*18, 19*), cell metabolism (*19*), and
tumorigenesis (*20, 21*), possibly through its splicing activity. In recent years, aryl sulfonamide drugs,
such as Indisulam and E7820, were found to degrade RBM39 in a DDB1 and CUL4 associated factor
15 (DCAF15)-dependent manner and thus have been widely investigated in anticancer research (*22-
24*). While most studies on RBM39 so far focused on cancer cells, there is limited knowledge of its
function in immunity, with one *in vivo* study suggesting that Indisulam-mediated modulation of RBM39
enhances anti-tumor immunity (*25*).
119
In our study, RBM39 emerged as a prominent candidate through a CRISPR/Cas9-mediated screening
in two liver-based cell lines aimed at identifying novel factors associated with the TLR3 response in
hepatocytes. Our study reveals that RBM39 exerts a broader influence on innate immunity by
modulating the transcription and splicing processes involved in the basal expression of IRF3 and
other crucial factors in the IFN signaling pathway. Based on the findings, it is plausible to consider
that drugs targeting RBM39, such as Indisulam, may function as immunosuppressants in individuals
undergoing treatment.

127
128 **Results:**
129
130 **A genome-wide CRISPR/Cas9 Screen identifies novel factors contributing to TLR3 signaling.**
131 The initial aim of this study was the identification of novel factors contributing to the regulation of cell
132 intrinsic innate immune responses, in particular focusing on TLR3-mediated response in hepatocytes.
133
To establish a positive selection system, we developed a death reporter system expressing the
suicide gene truncated BH3 interacting domain death agonist (tBID) under the transcriptional control
of the interferon induced protein with tetratricopeptide repeats 1 (IFIT1) promoter (Fig. 1A). Upon
activation of the TLR3 pathway, tBID expression is induced, resulting in apoptosis (*26*). For the
screening we chose PH5CH (*27*), a non-neoplastic, immortalized hepatocyte cell line expressing all
PRRs responsive to dsRNA, and Huh7-Lunet-TLR3 (*28*), a hepatocellular carcinoma (HCC)-derived
cell line requiring ectopic expression of PRRs for efficient response to dsRNA. The tBID reporter was
transduced into both cell lines for stable expression. To evaluate the approach, TLR3-specific
stimulation was achieved by the addition of the synthetic dsRNA analog polyinosinic:polycytidylic acid
(poly(I:C)) to the cell culture supernatant, leading to efficient tBID-associated, TLR3-dependent cell
death only upon activation of the *IFIT1* promoter in both cell lines, in contrast to the empty vector (Fig.
S1A). Subsequently, we transduced the IFIT1-tBID cell lines with lentiviral vectors encoding Cas9
along with the GeCKOv2.0 human genome-wide single guide RNA (sgRNA) library, that includes a
total of 122,411 unique sgRNAs targeting 19,050 genes and 1,864 miRNAs (Fig. 1B). After selection
with puromycin, we added 50 µg/ml poly(I:C) to the medium to activate the TLR3 pathway, or PBS as
a control. As a consequence, cells with an intact TLR3 pathway were killed due to the induction of
tBID expression, whereas cells expressing guide RNAs targeting genes involved in TLR3 signaling
survived, resulting in the enrichment of such sgRNAs. We then analyzed the sgRNA population of
stimulated and non-stimulated cells at 72 hours post-stimulation by next-generation sequencing
(NGS), to identify target genes interfering with the TLR3 pathway.
154
Candidate genes were determined by being enriched beyond a cell-line-specific threshold calibrated
to yield about 2,000 overrepresented sgRNAs for each cell line. Due to the expected background of
the screen, we needed to narrow down the enriched genes to obtain primary screen hits. We filtered
primary candidate genes using a thresholding strategy that was calibrated to include three positive
controls (TIR-domain-containing adapter-inducing interferon-β (TRIF), TLR3 and unc-93 homolog B1
(UNC93B1)) for each cell line (*29-31*). A gene was considered as a primary hit in case 5 out of 6
different sgRNAs found enriched in the poly(I:C) treated samples of at least one replicate and the
presence of at least one sgRNA targeting a gene in 2 out of 3 replicates. After such filtering, 50
protein-coding genes were chosen for functional validation, whereas 5 microRNAs were excluded
despite matching our criteria (Table S1). The protein-coding genes were validated by siRNA
knockdown in Huh7-Lunet-TLR3 cells (Fig. 1C and D). However, knockdown of most candidates

resulted in unimpaired or even upregulated TLR3 responses (Fig. 1C), likely due to a high
background of surviving cells even in absence of sgRNAs (Fig. S1A), resulting in non-specific
enrichment. Only the knockdown of a few genes reduced the TLR3 response to levels comparable to
the positive control (siUNC93B1). The four candidates with strongest phenotype, RBM39, lysine
demethylase 2A (KDM2A), protein tyrosine phosphatase receptor type T (PTPRT), and cleavage and
polyadenylation factor II subunit (PCF11) were chosen for further validation in PH5CH cells.

Indeed, knockdown of these factors strongly attenuated innate immune responses in PH5CH cells as
well (Fig. 1E). However, PCF11 was not followed up due to its general role in termination of
transcription (32). PTPRT mRNA turned out not to be detectable in Huh7 or PH5CH cells, and the
overexpression of siKDM2A-resistant mutant in PH5CH cells failed to rescue the KDM2A phenotype
(Fig. S1B), arguing for off-target effects of these siRNAs, therefore RBM39 was chosen for further
studies.

RBM39 knockdown reduces IRF3- but not NF-κB-induced genes upon TLR3 activation
Since PH5CH was considered the most authentic among the available hepatocyte models due to its
non-neoplastic nature, we decided to continue here with an in-depth analysis of the specific role of
RBM39 on TLR3-IRF3 signaling (Fig. 2A).

First of all, an RBM39 siRNA-resistant mutant (RBM39.Esc) was able to rescue the IFIT1 mRNA
expression upon siRNA knockdown in both PH5CH (Fig. 2B) and Huh7-Lunet-TLR3 cells (Fig. S2A),
demonstrating that the phenotype indeed was mediated by RBM39 knockdown. To rule out that
RBM39 might have an exclusive impact on IFIT1 expression, which so far served as the sole readout
in our validation, we further analyzed modulation of other ISGs upon RBM39 knockdown and TLR3
stimulation, including Interferon-Stimulated Protein, 15 KDa (ISG15), MX dynamin like GTPase 1
(MxA) and C-X-C motif chemokine ligand 10 (CXCL10) in PH5CH cells. All tested ISGs showed
reduction in RBM39-silenced cells compared to the control, and RBM39.Esc overexpression again
rescued the knockdown effect (Fig 2 C-E), excluding the off-target effect of siRBM39 and suggesting
that RBM39 has a broad impact on various ISGs. To confirm this on protein level, we additionally
detected IFN-β secretion via ELISA upon RBM39 knockdown and observed a significant decline here
as well (Fig. 2F).

Considering that the activation of PRRs not only leads to production of various ISGs, but also triggers
NF-κB signaling, resulting in the production of cytokines and chemokines, we investigated a potential
involvement of RBM39 in the NF-κB pathway (Fig. 2G). However, the expression of two NF-κB-
induced genes, tumor necrosis factor alpha induced protein 3 (TNFAIP3) and interleukin 6 (IL6) was
not or only mildly affected in PH5CH cells upon poly(I:C) activation and RBM39 knockdown (Fig. 2H),
indicating a minor effect of RBM39 on the NF-κB pathway in this experimental setting.

Indisulam is a sulfonamide drug mediating the interaction between the E3 ligase DCAF15 and
RBM39, leading to the ubiquitination and proteasomal degradation of RBM39 (Fig. 2I) (22, 33, 34).
Indeed, Indisulam dramatically reduced RBM39 expression in PH5CH, resulting in significantly
attenuated IFIT1 induction upon TLR3 activation (Fig 2J), further emphasizing the role of RBM39 on
the TLR3 pathway.

Overall, our results suggest that RBM39 was involved in the direct induction of IFNβ and ISGs upon
stimulation with dsRNA, but had a minor influence on NF-κB signaling, suggesting that the IRF3
pathway was mainly affected.

RBM39 controls the RIG-I/MDA5 pathway

Given that Huh7.5 cells, a subclone of Huh7 (35), did not express functional PRRs, ectopic
expression of retinoic acid inducible gene I (RIG-I) and melanoma differentiation-associated gene 5
(MDA5) allowed us to identify the function of RBM39 in other key innate immune pathways governed
by IRF3 (Fig. 3A) (28). Notably, silencing of RBM39 decreased RIG-I and MDA5-mediated IFIT1
mRNA expression upon transfection of poly(I:C) (Fig. 3B and C). To validate our finding in non-
hepatic cell lines, we tested A549 cells, which are based on adenocarcinoma cells from human

alveolar basal epithelium. These cells endogenously express RIG-I and MDA5, but not TLR3, and are
regarded as innate immune-competent (*36*). Indeed, knockdown of *RBM39* in A549 cells reduced
IFIT1 induction in response to poly(I:C) similarly as observed for PH5CH and Huh7-Lunet (Fig. 3D).
Additionally, Indisulam induced RBM39 degradation also significantly reduced RIG-I/MDA5 responses
in A549 cells (Fig. 3E), collectively demonstrating that RBM39's involvement in innate immunity is
neither limited to hepatocytes, nor to TLR3 signaling.

So far, our study exclusively relied on stimulation of cell intrinsic innate immune responses using the
synthetic dsRNA analog poly(I:C). To investigate the function of RBM39 in innate immunity in more
physiological settings, we performed virus infection experiments with various viruses known to induce
strong innate immune responses mediated by IRF3.

Sendai virus (SeV) is a model virus sensed by RIG-I (*37*). Knockdown of *RBM39* in A549 cells indeed
resulted in a strong reduction of *IFIT1* mRNA expression, comparable to the one upon *MAVS*
knockdown. We further assessed viral RNA abundance to ensure that RMB39 depletion would not
inhibit viral replication, resulting in lower levels of viral stimuli. To this end, we found a slight,
comparable increase for both conditions (Fig. 3F). Indisulam was used as an alternative way to delete
RBM39 in A549 cells, consistently, it also attenuated SeV-induced *IFIT1* expression, again without
affecting viral RNA loads. Similar results were obtained for infection of A549 cells with rift valley fever
virus (RVFV) and vesicular stomatitis virus (VSV), HepG2/C3A cells with hepatitis E virus (HEV), and
HepG2-NTCP cells with hepatitis D virus (HDV) upon Indisulam treatment, resulting in reduced *IFIT1*
mRNA induction, in absence of inhibitory effects on viral RNA replication (Fig. S3A to D).

Altogether, our data demonstrate that RBM39 participated in the regulation of RIG-I and MDA5
pathways and substantially contributed to the cell intrinsic innate immune response to viral infections.

RBM39 regulates basal IRF3 expression to modulate innate immune responses

In order to elucidate the mechanism involved in RBM39-mediated regulation of IFN and ISG induction,
we firstly examined the phosphorylation of IRF3, a key event mediating IFN and ISG induction upon
PRR activation by dsRNA. As expected, phosphorylated-IRF3 (p-IRF3) was undetectable in non-
stimulated cells and strongly enhanced upon poly(I:C) stimulation. Intriguingly, *RBM39* knockdown not
only resulted in a remarkable decrease in p-IRF3, but also in significant reduction of total IRF3 protein
levels, suggesting a regulation of the constitutive expression of IRF3 (Fig. 4A).

To further validate the impact of RBM39 on IRF3 expression, we treated PH5CH cells with Indisulam.
Indeed, RBM39 protein levels were reduced in Indisulam-treated cells in a dose-dependent manner,
concomitant with lower IRF3 abundance and some cytopathic effect at higher concentrations (Fig. 4B).
In order to verify whether IRF3 depletion upon Indisulam treatment was mediated by RBM39, we made
use of an RBM39 mutant (G268V) reported to confer partial resistance to Indisulam (*22*).
Overexpression of RBM39 G268V partially rescued RBM39 Indisulam-mediated degradation and thus
restored IRF3 expression compared to cells expressing wild-type RBM39 (Fig. S4A). These results
confirmed that Indisulam affected IRF3 expression by targeting RBM39.

Similar results were obtained in A549 cells (Fig. S4B) and in primary human hepatocytes (PHH) (Fig.
4C), with the latter indicating that the RBM39-mediated IRF3 depletion also appeared in primary cells,
albeit requiring higher concentrations associated with substantial cytotoxicity (Fig. 4C). In contrast,
RBM39 was refractory to Indisulam treatment in primary mouse hepatocytes (PMH) (Fig. S4C),
probably due to the lack of expression of murine DCAF15 or resistance to Indisulam (*38*), in line with
the lack of cytotoxicity (Fig. S4C, right).

Above all, our results demonstrated that RBM39 governs basal expression of IRF3, thereby exerting a
crucial role in the activation of innate immune responses.

Global transcriptome, splicing and proteome analysis highlights the role of RBM39 in innate
immunity

We next aimed for a comprehensive proteomic and transcriptomic analysis, to identify other
components of innate immunity affected by RBM39, comparing the impact of siRNA- mediated
RBM39 knockdown and Indisulam treatment in PH5CH cells.
281

For the proteomic analysis we found 165 proteins in *RBM39* knockdown samples and 627 proteins in
Indisulam-treated samples with significant changes (fold change < 0.5 or > 2, p-value <0.05) (Fig. 5A
and B, Table S2). RBM39 was among the most significantly changed proteins, as expected (Fig. 5A
and S5B), whereas GAPDH signals were not significantly changed (Fig. S5A), arguing against a
strong impact of cytotoxic effects. IRF3 levels were significantly reduced in both conditions, even
though the changes upon RBM39 knockdown were mild (Fig. 5A and B, S5C). The generally lower
magnitude of effects upon RBM39 knockdown might be mediated by the difference in the time-frame
of RBM39 reduction, acting faster due to direct protein degradation by Indisulam treatment compared
to the time required for siRNA-mediated reduction of RBM39 abundance. Importantly, we identified
additional factors critically contributing to the cell intrinsic innate immune response, including RIG-I,
MDA5, STAT1 and STAT2 (Fig. 5A and B, S5D to G). Similar to IRF3, all these factors showed
significant changes in both conditions, with strong reduction in Indisulam-treated samples and mild
decline in RBM39 knockdown samples, except for RIG-I that only was significantly decreased in
Indisulam-treated samples (Fig. S5E).
296
Due to the complex, multifactorial mode of action of RBM39 in transcription and splicing, we analyzed
the RNAseq dataset in the context of differentially expressed genes (DEG) and differential transcript
usages (DTUs), indicative of changes in splicing, comparing knockdown of *RBM39* and Indisulam
treatment. Our analysis identified 4,007 DEGs (fold change < 0.5 or > 2, p-value <0.05) and 4,053
DTUs (p-value >0.05) in *RBM39* knockdown and Indisulam-treated samples. 1,570 DEGs and 1,977
DTUs overlapped in both conditions (39% and 49% of total events, respectively) (Fig. 5C-E, Table S3
and Table S4), arguing for the reliability of the analysis. Comparing with proteomic data, DEG
analysis revealed more factors with significant fold change and similar numbers as previous
transcriptomic studies (Fig. 5C and D, Table S3) (*21, 23, 39, 40*). Interestingly, *RBM39* RNA
abundance was significantly reduced upon knockdown (Fig. 5C), as expected, but strikingly increased
upon Indisulam treatment due to autoregulation of *RBM39* mRNA levels by protein abundance (Fig.
5D) (*41*). Importantly and in agreement with the proteomic data, we found significant changes in
factors involved in the cell intrinsic innate immune response, namely RIG-I, MDA5, STAT1, and
STAT2, but in addition, type I and type III IFN receptors.
311
312 In summary, proteomic and transcriptomic analysis consistently identified several key factors of the
313 interferon pathway beyond IRF3 that were regulated by RBM39.
314
315 **RBM39 governs the transcription and splicing of crucial innate immune factors**
316 Next, we aimed at elucidating whether reduction of mRNA and protein abundance of the key factors
317 of the IFN-response by RBM39 was mediated by alternative splicing or transcriptional regulation.
318
For IRF3, we found a non-significant 20% reduction *of IRF3* mRNA read counts upon knockdown of
RBM39 or Indisulam treatment (Fig. 6A, left), suggesting that transcriptional regulation was not the
main mode of action. This was confirmed by analyzing the impact of RBM39 knockdown on *IRF3*
mRNA by RT-qPCR (Figure S6A)*,* and on activity of the *IRF3* promoter (*42*) using luciferase reporter
plasmids (Fig. S6B, C) both showing a similar magnitude of reduction. In contrast, DTU analysis
revealed substantial, consistent changes in splicing of *IRF3* mRNA when RBM39 was suppressed
either by siRNA or by Indisulam treatment, with a significant decrease of the functional *IRF3-203*
isoform and an increased abundance of the particular isoform *IRF3-228*, which has been reported to
have an inhibitory effect on innate immune activation (*43*) (Fig. 6A, right panel). This data was
validated upon *RBM39* knockdown via RT-qPCR using primers specifically targeting each of the *IRF3*
isoforms that were detected in RNAseq, obtaining very similar results (Fig. S6D). To exclude the
contribution of additional factors in the TLR3 pathway potentially affected by RMB39, we expressed
IRF3 ectopically in PH5CH IRF3 KO cells. Indeed, overexpression of IRF3 restored the *IFIT1* mRNA
induction upon RBM39 knockdown and TLR3 stimulation (Fig. 6B, C), suggesting that IRF3 was the
sole determinant of the TLR3 response regulated by RBM39*.*
334
In addition to IRF3, the cytosolic dsRNA sensors RIG-I and MDA5 were found to be significantly
downregulated in the proteomic and transcriptomic analysis (Fig. 5/S5). DEG/DTU analysis revealed
that *RIG-I* expression was modulated both by transcription and splicing regulation, the latter resulting
in a lower abundance of the functional isoform 202 (Fig. 6D), whereas *MDA5* expression was mainly
decreased at the transcriptional level (Fig. 6E). To verify the consequence of RBM39 regulation on
the RIG-I/MDA5 pathways beyond IRF3 (Fig. 6F), we overexpressed IRF3 in A549 cells, thereby
excluding any contribution of TLR3. In contrast to the TLR3 pathway in PH5CH cells, IRF3 expression

did not suffice to restore *IFIT1* mRNA induction upon RBM39 knockdown and poly(I:C) transfection
(Fig. 6G), indicating additional contributions of *RIG-I* and *MDA5*. Consequently, induction of the NF-
kB-specific genes *TNFAIP3* and *IL6*, was significantly reduced upon RBM39 knockdown, when the
pathway was triggered via RIG-I/MDA5 (Fig. 6H, I).
346
347 Taken together, we found that basal expression of IRF3 and RIG-I/MDA5 was regulated by the
348 abundance of RBM39, with significant impact on the induction of cell intrinsic innate immune
349 responses.
350
351 **RBM39 regulates the IFN-JAK-STAT pathway**

352 Our proteomic and transcriptomic analysis further identified components of the IFN-JAK-STAT
353 pathway being regulated by RBM39, namely STAT1 and 2 and type I and III IFN receptors, the latter
354 were not detectable by proteomics likely due to low protein abundance. We therefore aimed to
355 understand the mode of regulation and the functional relevance for the IFN response.
356
The receptor for type I IFNs is a heterodimer consisting of IFNAR1 and IFNAR2. While *IFNAR1*
expression was upregulated only on the transcriptional level (Fig. S6E), *IFNAR2* mRNA abundance
was reduced and splicing was modulated upon RBM39 knockdown and Indisulam treatment (Fig. 7A).
The type III interferon receptor is a heterodimer as well, composed of IFNLR1 and interleukin 10
receptor subunit beta (IL10RB), and also here the expression of both subunits was regulated by
RBM39 in opposite directions (Fig. S6F and Fig. 7B), with an increase for *IFNLR1* (Fig. S6F) and a
decrease for *IL10RB* (Fig. 7B) RNA amounts, with minor or undetectable changes in the splicing
pattern, respectively.
STAT1 and STAT2, forming heterodimers upon activation of the type I and type III IFN responses,
were in contrast both reduced in protein expression levels upon depletion of RBM39 (Fig. S5F, G),
mediated by changes in RNA abundance and associated with modified splicing patterns (Fig. 7C, D).
368
369 To investigate the cumulative effect of up- or down-regulation of these factors on the respective IFN
370 response pathways (Fig. 7E), we silenced RBM39 in PH5CH cells and stimulated them with IFNα2
371 and IFNλ1 to activate type I and type III IFN pathways, respectively. Consequently, the type I and type
372 III IFN responses were both strongly impaired, as indicated by reduced IFIT1 induction (Fig. 7F and
373 G), suggesting that RBM39 also regulates IFN-JAK-STAT signaling.
374
375 All in all, our RNA-seq and proteome analysis identified key factors of the IFN response affected by
376 RBM39, such as IRF3, RIG-I, MDA5, STAT1 and STAT2 and type I/III receptors, arguing for a crucial
377 role of RBM39 in the regulation of innate immunity.
378
Discussion

Genome-wide CRISPR/Cas9 screens present a powerful tool to study host factors of innate immunity
mostly in immune cells, such as macrophages (*44*), dendritic cells (*45*), or in the more commonly used
Hela cells (*46*) and HEK 293T cells (*47*). In our study, we applied this approach in two liver cell lines,
Huh7 and PH5CH, and specifically targeted the TLR3 pathway by poly(I:C) supernatant feeding.
Eventually, we identified RBM39 as a novel factor that affects TLR3 signaling and has a broad impact
on the cell intrinsic innate immune response by regulating the basal expression of IRF3 and other
important pathway components.
387
RBM39 is a factor involved in splicing and transcriptional regulation of many genes, particularly
contributing to tumorigenesis (*20, 21, 48*), cell cycle control (*12, 19*), and metabolism (*19*), but so far
its role in innate immunity has not been shown. Paradoxically, a previous study indicated that porcine
reproductive and respiratory syndrome virus (PRRSV) infection upregulated porcine RBM39
expression, resulting in dephosphorylation of c-Jun, and subsequent downregulation of AP-1 signaling
(*49*). Additionally, studies have reported that some viruses regulate splicing factors like RBM10 and
U5 spliceosomal RNA (snRNA) as an evasion strategy (*50, 51*). While RBM39 has previously been
shown to interact with the viral NF-κB homolog v-Rel, inhibiting its activation, the interplay between
RBM39 and other human host NF-κB members like p50 and p105 has not been explored. Here, we
identified basal expression of *IRF3* as one of the key factors of cell intrinsic innate immune responses
being modulated by knockdown of *RBM39* and Indisulam-mediated RBM39 degradation. This
resulted in significantly impaired activation upon both dsRNA administration and virus infection. In
addition, we identified additional pathway components being modulated by RBM39, including the

PRRs RIG-I and MDA5, IL-10RB, IFNAR2, STAT1 and STAT2. These factors all contributed to the
reduced activation of the interferon response. Overall, this is supposed to result in an increased level
of viral replication. In our infection experiments RBM39 depletion only slightly stimulated viral RNA
abundance, most likely due to the limited timeframe, providing not sufficient time to establish more
pronounced phenotypes.
406
The complexity of regulatory functions mediated by RBM39 might explain why other transcriptomic
and proteomic analyses missed its role in innate immunity so far (*22, 23, 39, 52-55*). In the
transcriptomic and splicing analysis, knockdown of *RBM39* by siRNA and Indisulam treatment
showed a wide degree of overlap, arguing for the specificity of Indisulam towards RBM39 in the liver-
derived cell lines used in our study and in agreement with previous studies (*53, 56*). Overall, the
transcriptomic analysis was in our hands more prone to identify innate immune response factors
regulated by RBM39. While we could demonstrate that changes in IRF3, as well as RIG-I/MDA5
expression contributed to the induction of cell intrinsic innate immune response, the data were more
complex in the case of IFN treatment. Type I and type III IFNs bind to different receptors, but engage
identical signaling pathways (*57*). We observed reduced expression of *IL10RB*, *IFNAR2*, *STAT1* and
STAT2 but increased expression of *IFNAR1* and *IFNLR1*, all essential for signal transduction, with
STAT1 and STAT2 being involved in the response pathway to both type I and type III IFNs (*57*).
Importantly, we confirmed that the aforementioned changes of these factors in sum resulted in the
attenuation of both IFN pathways. In addition, further systemic effects of RBM39 ablation on innate
immune responses are possible due to the complex involvement of IL-10RB in several cytokine
receptor complexes, responding not only to IFNλ, but also to pro- and anti-inflammatory cytokines
including IL-10, IL-22 and IL-28/29 (*58, 59*).
424
RBM39 depletion was shown to have a strong anti-proliferative impact, which might explain our failure
to obtain RBM39 knockout clones in PH5CH cells (not shown). This growth inhibitory effect has put
forward RBM39 as a promising target for cancer therapy and contributed to numerous clinical trials
implementing aryl sulfonamides like Indisulam, albeit yet with limited success, likely due to the lack of
predictive markers (*60*). In contrast, Indisulam showed high activity in preclinical models like mice
engrafted with human tumors (*61*) and, more recently, in neuroblastoma models (*23, 61*), associated
with limited toxicity. Indisulam treatment of all human hepatic cells tested in our study, including two
cancer cell lines (HepG2 and Huh7), one non-neoplastic cell line (PH5CH), and PHH, resulted in a
significant degradation of RBM39, albeit at varying doses. Although RBM39 protein expression was
similar in all these cell lines, PHH could be less sensitive to Indisulam due to lower *DCAF15*
expression levels, critical for its mode of action. Importantly, RBM39 degradation was associated with
a certain degree of cytotoxicity in all of our cell culture models. In contrast, Indisulam treatment of
PMH had no impact on RBM39 and IRF3 expression and lacked obvious cytotoxicity. The different
Indisulam sensitivity of PHH and PMH therefore imposes particular caution in interpreting toxicity data
obtained in animal models. This is consistent with data showing only a minor effect of Indisulam
treatment on RBM39 expression in different organs of mice, including the liver, with significant
changes only in murine kidney and the engrafted human tumors (*22, 40, 61*).
442
The lack of Indisulam activity in PMH further precluded *in vivo* testing regarding immunosuppressive
activities *in vivo*. Currently available aryl sulphonamides might generally not be suitable for clinical
use as immunosuppressors, due to their focus on anti-cancer activity. However, substances targeting
RBM39 to proteasomal degradation based on alternative E3 ligases beyond DCAF15 might be
envisaged, as well as alternative tailored depletion strategies like molecular glues and proteolysis
targeting chimeras (PROTACs) (*62*). In light of our data defining RBM39 as a regulator of key
components of cell intrinsic innate immunity, it could be regarded as a candidate target to mitigate
excessive innate immune responses contributing to viral pathogenesis (*63*) or autoimmune diseases
(*64*).
452
453
454
455
Materials and Methods
Cell culture:
PH5CH cells were a generous gift from K. Shimotohno. Huh 7.5 cells were a kind gift from Charles
Rice. A549 cell were obtained from ATCC, A549 IFNAR/LR KO cells (*65*), Huh7-Lunet cells (*66*),

Huh7.5-RIG-I and Huh7.5-MDA5 (*28*), HepG2-NTCP (*67*) have been described before. HepG2/C3A
were obtained from ATCC. Primary human hepatocytes (PHH) were purchased from BioIVT and
cultured in William E media containing 1% (v/v) Glutamax, 1% (v/v) Non-Essential Amino Acids, 1%
(v/v) penicillin/streptomycin (all from Gibco, Germany), 0.2% (v/v) normocin (Invivogen, USA), 2%
(v/v) B27 (Gibco, Germany), 1% (v/v) N2 supplement (Gibco, Germany), 100 mM nicotinamide
(Sigma-Aldrich, USA), 1.25 mM N-acetylcysteine (Sigma-Aldrich, USA), 10 µM Y27632 (Peprotech,
USA), 1 µM A83-01 (Tocris, UK). Cells, unless specifically mentioned, were cultured in Dulbecco's
Modified Eagle Medium (Life Technologies, Darmstadt, Germany) supplemented with 10% fetal
bovine serum, nonessential amino acids (Life Technologies), 100 U/mL penicillin, and 100 ng/mL
streptomycin (Life Technologies) and cultivated at 37°C and 5% CO_2. PH5CH-KDM2A, PH5CH-
RBM39.Esc, PH5CH-RBM39 G268V, PH5CH-IRF3, A549-IRF3 cells were kept under selection
pressure of 1 mg/mL G418 (Geneticin) (Life Technologies, USA), Huh7-Lunet-TLR3, Huh7.5-RIG-I,
HepG2-NTCP cells were maintained under selective pressure of 1 µg/mL puromycin (Sigma-Aldrich,
Germany), Huh7.5-MDA5 cells were kept under the antibiotic pressure of 5 µg/mL blasticidin (Sigma-
Aldrich, Germany).
475
siRNA transfection：

The used siRNAs were purchased from Horizon Discovery, UK, unless specifically mentioned.
siMAVS was purchased from Sigma Aldrich, Germany. For siRNA transfection in a 6-well plate, cells
with 60%-70% confluency were seeded 16 hours prior to the experiment and 1.5 µM siRNA were
mixed with 9 µl Lipofectamine RNAiMax (Thermo Fisher Scientific, USA) in OptiMEM according to
manufacturer's instructions. After 20 min incubation at RT, the mixture was added into the medium.
Cells were incubated for 48 h and used for further experiments.
483
Drug treatments:

Poly(I:C) stimulation: Poly (I:C) (HMW) was used (Invivogen, USA) following the manufacturer's
instructions. To specifically trigger TLR3, 50 µg/ml poly (I:C) were added directly into the medium. For
TLR3, RIG-I and MDA5 stimulation, transfection was performed with 0.5 µg/ml poly (I:C) mixed with
Lipofectamine2000 (Thermo Fisher Scientific, USA) in OptiMEM (Gibco, USA) according to the
manufacturer's instructions. After 20 min incubation at room temperature (RT), the mixture was added
into the medium. Cells were stimulated for 6 h or 8 h, then harvested for RNA or protein extraction.
491
Indisulam treatment: Indisulam was utilized (Sigma Aldrich, Germany) solved in DMSO at a final stock
concentration of 10 mM. Indisulam, or same amount of DMSO as control, were added to the cell
medium at the indicated concentrations; cells were incubated for the indicated time and treated further
with either poly (I:C) or virus infection. At the indicated time points, cells were harvested for RNA
extraction or protein analysis. IFNs stimulation: IFNα2 and IFNλ1 were purchased from PBL, USA and
Peprotech, USA, respectively. 50 IU/ml IFN2α and 10ng/ml IFNλ1 were added to the medium. Cells
were stimulated for 24 h and collected for RNA extraction.
499
500 **Statistical analysis:**

501 Independent biological replicates are denoted with n-numbers. To test for significance, two-tailed
502 unpaired Welch's test, were performed using GraphPad Prism 8 software (GraphPad Software, La
503 Jolla, CA, USA). ∗p <0.05; ∗∗p <0.01; ∗∗∗p <0.001; ∗∗∗∗p <0.0001.
504

Acknowledgments: The authors thank Rahel Klein, Ulrike Herian for excellent technical assistance.
We thank Charles M. Rice for Huh7.5 cells and Kunitada Shimotohno for PH5CH cells. We
also need to thank the Genomics and Proteomics Core facility of the German Cancer
Research Centre for the RNA-seq sample measurement, as well as Dr Julia Mergner,
Proteomics Core Facility of the TranslaTUM. Additionally, we thank Zhenfeng Zhang, Julia
Welsch, Jun-gen Hu, Zhaozhi Sun for providing reagents, and Cong Si Tran and Minh-Thu
Pham for helpful discussions.
512
513 **Funding:** This work was supported by the Deutsche Forschungsgemeinschaft (DFG, German
514   Research Foundation) – project number 272983813 – TRR 179, to VL, MB, AP, DW, RB, MS,
515   SU and VLDT. VL received additional DFG funding, project number 519777725. TL was
516   supported by the China Scholarship Council. MB and VGM were supported by DFG

236528794. NAR is a holder of a Georg Forster Research Fellowship for Experienced Researchers from the Alexander von Humboldt Foundation.

Data and materials availability: Details related to materials and methods are provided in the Supplementary material. Transcriptomic data were deposited in the Sequence Read Archive (SRA) under accession PRJNA1028350. Further data supporting the findings of this study is available upon request. Please contact the corresponding author.

References and Notes

1. N. C. Di Paolo *et al.*, Virus binding to a plasma membrane receptor triggers interleukin-1 alpha-mediated proinflammatory macrophage response in vivo. *Immunity* **31**, 110-121 (2009).
2. A. Krug *et al.*, TLR9-dependent recognition of MCMV by IPC and DC generates coordinated cytokine responses that activate antiviral NK cell function. *Immunity* **21**, 107-119 (2004).
3. F. Heil *et al.*, Species-specific recognition of single-stranded RNA via toll-like receptor 7 and 8. *Science* **303**, 1526-1529 (2004).
4. S. A. Nicholas, V. V. Sumbayev, The involvement of hypoxia-inducible factor 1 alpha in Toll-like receptor 7/8-mediated inflammatory response. *Cell Res* **19**, 973-983 (2009).
5. D. Goubau *et al.*, Antiviral immunity via RIG-I-mediated recognition of RNA bearing 5 '-diphosphates. *Nature* **514**, 372-+ (2014).
6. L. Gitlin *et al.*, Essential role of mda-5 in type I IFN responses to polyriboinosinic:polyribocytidylic acid and encephalomyocarditis picornavirus. *Proc Natl Acad Sci U S A* **103**, 8459-8464 (2006).
7. E. A. Kurt-Jones *et al.*, Herpes simplex virus 1 interaction with Toll-like receptor 2 contributes to lethal encephalitis. *Proc Natl Acad Sci U S A* **101**, 1315-1320 (2004).
8. N. Gillich, Z. Zhang, M. Binder, S. Urban, R. Bartenschlager, Effect of variants in LGP2 on MDA5-mediated activation of interferon response and suppression of hepatitis D virus replication. *J Hepatol* **78**, 78-89 (2023).
9. M. Yamamoto *et al.*, Role of adaptor TRIF in the MyD88-independent toll-like receptor signaling pathway. *Science* **301**, 640-643 (2003).
10. H. Imai, E. K. Chan, K. Kiyosawa, X. D. Fu, E. M. Tan, Novel nuclear autoantigen with splicing factor motifs identified with antibody from hepatocellular carcinoma. *J Clin Invest* **92**, 2419-2426 (1993).
11. A. H. Sillars-Hardebol *et al.*, CSE1L, DIDO1 and RBM39 in colorectal adenoma to carcinoma progression. *Cell Oncol (Dordr)* **35**, 293-300 (2012).
12. I. Mercier *et al.*, CAPER, a novel regulator of human breast cancer progression. *Cell Cycle* **13**, 1256-1264 (2014).
13. Y. Chai *et al.*, Overexpression of HCC1/CAPERalpha may play a role in lung cancer carcinogenesis. *Tumour Biol* **35**, 6311-6317 (2014).
14. G. A. Stepanyuk *et al.*, UHM-ULM interactions in the RBM39-U2AF65 splicing-factor complex. *Acta Crystallogr D Struct Biol* **72**, 497-511 (2016).
15. D. J. Jung, S. Y. Na, D. S. Na, J. W. Lee, Molecular cloning and characterization of CAPER, a novel coactivator of activating protein-1 and estrogen receptors. *J Biol Chem* **277**, 1229-1234 (2002).
16. D. H. Dowhan *et al.*, Steroid hormone receptor coactivation and alternative RNA splicing by U2AF65-related proteins CAPERalpha and CAPERbeta. *Mol Cell* **17**, 429-439 (2005).
17. S. Mai *et al.*, Functional interaction between nonreceptor tyrosine kinase c-Abl and SR-Rich protein RBM39. *Biochem Biophys Res Commun* **473**, 355-360 (2016).
18. N. Faherty *et al.*, Negative autoregulation of BMP dependent transcription by SIN3B splicing reveals a role for RBM39. *Sci Rep* **6**, 28210 (2016).
19. Y. K. Kang *et al.*, CAPER is vital for energy and redox homeostasis by integrating glucose-induced mitochondrial functions via ERR-alpha-Gabpa and stress-induced adaptive responses via NF-kappaB-cMYC. *PLoS Genet* **11**, e1005116 (2015).
20. G. Huang, Z. Zhou, H. Wang, E. S. Kleinerman, CAPER-alpha alternative splicing regulates the expression of vascular endothelial growth factor(1)(6)(5) in Ewing sarcoma cells. *Cancer* **118**, 2106-2116 (2012).
21. E. Wang *et al.*, Targeting an RNA-Binding Protein Network in Acute Myeloid Leukemia. *Cancer Cell* **35**, 369-384 e367 (2019).
22. T. Han *et al.*, Anticancer sulfonamides target splicing by inducing RBM39 degradation via recruitment to DCAF15. *Science* **356**, (2017).

23. A. Nijhuis *et al.*, Indisulam targets RNA splicing and metabolism to serve as a therapeutic
strategy for high-risk neuroblastoma. *Nat Commun* **13**, 1380 (2022).
24. T. Uehara *et al.*, Selective degradation of splicing factor CAPERalpha by anticancer
sulfonamides. *Nat Chem Biol* **13**, 675-680 (2017).
25. S. X. Lu *et al.*, Pharmacologic modulation of RNA splicing enhances anti-tumor immunity. *Cell*
184, 4032-4047 e4031 (2021).
26. K. Huang *et al.*, Cleavage by Caspase 8 and Mitochondrial Membrane Association Activate
the BH3-only Protein Bid during TRAIL-induced Apoptosis. *J Biol Chem* **291**, 11843-11851
(2016).
27. M. Ikeda *et al.*, Human hepatocyte clonal cell lines that support persistent replication of
hepatitis C virus. *Virus Res* **56**, 157-167 (1998).
28. O. Colasanti *et al.*, Comparison of HAV and HCV infections in vivo and in vitro reveals distinct
patterns of innate immune evasion and activation. *J Hepatol*, (2023).
29. E. Meylan *et al.*, Cardif is an adaptor protein in the RIG-I antiviral pathway and is targeted by
hepatitis C virus. *Nature* **437**, 1167-1172 (2005).
30. L. Alexopoulou, A. C. Holt, R. Medzhitov, R. A. Flavell, Recognition of double-stranded RNA
and activation of NF-kappaB by Toll-like receptor 3. *Nature* **413**, 732-738 (2001).
31. K. Pelka *et al.*, The Chaperone UNC93B1 Regulates Toll-like Receptor Stability
Independently of Endosomal TLR Transport. *Immunity* **48**, 911-922.e917 (2018).
32. K. Kamieniarz-Gdula *et al.*, Selective Roles of Vertebrate PCF11 in Premature and Full-
Length Transcript Termination. *Mol Cell* **74**, 158-172.e159 (2019).
33. D. E. Bussiere *et al.*, Structural basis of indisulam-mediated RBM39 recruitment to DCAF15
E3 ligase complex. *Nat Chem Biol* **16**, 15-23 (2020).
34. T. C. Ting *et al.*, Aryl Sulfonamides Degrade RBM39 and RBM23 by Recruitment to CRL4-
DCAF15. *Cell Rep* **29**, 1499-1510 e1496 (2019).
35. K. J. Blight, J. A. McKeating, C. M. Rice, Highly permissive cell lines for subgenomic and
genomic hepatitis C virus RNA replication. *J Virol* **76**, 13001-13014 (2002).
36. Y. Li *et al.*, SARS-CoV-2 induces double-stranded RNA-mediated innate immune responses
in respiratory epithelial-derived cells and cardiomyocytes. *Proc Natl Acad Sci U S A* **118**,
(2021).
37. L. Martínez-Gil *et al.*, A Sendai virus-derived RNA agonist of RIG-I as a virus vaccine
adjuvant. *J Virol* **87**, 1290-1300 (2013).
38. B. V. Mistry *et al.*, Expression profiling of WD40 family genes including DDB1- and CUL4-
associated factor (DCAF) genes in mice and human suggests important regulatory roles in
testicular development and spermatogenesis. *BMC Genomics* **21**, 602 (2020).
39. S. Mai *et al.*, Global regulation of alternative RNA splicing by the SR-rich protein RBM39.
Biochim Biophys Acta **1859**, 1014-1024 (2016).
40. P. K. Puvvula *et al.*, Inhibiting an RBM39/MLL1 epigenomic regulatory complex with
dominant-negative peptides disrupts cancer cell transcription and proliferation. *Cell Rep* **35**,
109156 (2021).
41. S. Campagne *et al.*, Molecular basis of RNA-binding and autoregulation by the cancer-
associated splicing factor RBM39. *Nat Commun* **14**, 5366 (2023).
42. H. G. Xu, W. Ren, C. Lu, G. P. Zhou, Characterization of the human IRF-3 promoter and its
regulation by the transcription factor E2F1. *Mol Biol Rep* **37**, 3073-3080 (2010).
43. A. Y. Karpova, L. V. Ronco, P. M. Howley, Functional characterization of interferon regulatory
factor 3a (IRF-3a), an alternative splice isoform of IRF-3. *Mol Cell Biol* **21**, 4169-4176 (2001).
44. A. J. Chui *et al.*, N-terminal degradation activates the NLRP1B inflammasome. *Science* **364**,
82-85 (2019).
45. O. Parnas *et al.*, A Genome-wide CRISPR Screen in Primary Immune Cells to Dissect
Regulatory Networks. *Cell* **162**, 675-686 (2015).
46. R. Fang, Q. Jiang, X. Jia, Z. Jiang, ARMH3-mediated recruitment of PI4KB directs Golgi-to-
endosome trafficking and activation of the antiviral effector STING. *Immunity* **56**, 500-
515.e506 (2023).
47. R. Fang *et al.*, Golgi apparatus-synthesized sulfated glycosaminoglycans mediate
polymerization and activation of the cGAMP sensor STING. *Immunity* **54**, 962-975.e968
(2021).
48. J. Dutta, G. Fan, C. Gelinas, CAPERalpha is a novel Rel-TAD-interacting factor that inhibits
lymphocyte transformation by the potent Rel/NF-kappaB oncoprotein v-Rel. *J Virol* **82**, 10792-
10802 (2008).

49. Y. Song *et al.*, RBM39 Alters Phosphorylation of c-Jun and Binds to Viral RNA to Promote
PRRSV Proliferation. *Front Immunol* **12**, 664417 (2021).
50. B. Pozzi *et al.*, Dengue virus targets RBM10 deregulating host cell splicing and innate
immune response. *Nucleic Acids Res* **48**, 6824-6838 (2020).
51. S. Boudreault *et al.*, Reovirus mu2 protein modulates host cell alternative splicing by reducing
protein levels of U5 snRNP core components. *Nucleic Acids Res* **50**, 5263-5281 (2022).
52. E. Wang *et al.*, Targeting an RNA-Binding Protein Network in Acute Myeloid Leukemia.
Cancer Cell **35**, 369-384.e367 (2019).
53. T. C. Ting *et al.*, Aryl Sulfonamides Degrade RBM39 and RBM23 by Recruitment to CRL4-
DCAF15. *Cell Rep* **29**, 1499-1510.e1496 (2019).
54. Z. Dou *et al.*, Indisulam exerts anticancer effects via modulation of transcription, translation
and alternative splicing on human cervical cancer cells. *Am J Cancer Res* **13**, 2922-2937
(2023).
55. S. Coomar *et al.*, Overlaid Transcriptional and Proteome Analyses Identify Mitotic Kinesins as
Important Targets of Arylsulfonamide-Mediated RBM39 Degradation. *Mol Cancer Res*, Of1-
of11 (2023).
56. T. Uehara *et al.*, Selective degradation of splicing factor CAPERα by anticancer
sulfonamides. *Nat Chem Biol* **13**, 675-680 (2017).
57. H. M. Lazear, J. W. Schoggins, M. S. Diamond, Shared and Distinct Functions of Type I and
Type III Interferons. *Immunity* **50**, 907-923 (2019).
58. S. Mukhopadhyay *et al.*, Loss of IL-10 signaling in macrophages limits bacterial killing driven
by prostaglandin E2. *J Exp Med* **217**, (2020).
59. S. Mossner *et al.*, Synthetic interleukin 22 (IL-22) signaling reveals biological activity of
homodimeric IL-10 receptor 2 and functional cross-talk with the IL-6 receptor gp130. *J Biol
Chem* **295**, 12378-12397 (2020).
60. Y. Xu, A. Nijhuis, H. C. Keun, RNA-binding motif protein 39 (RBM39): An emerging cancer
target. *Br J Pharmacol* **179**, 2795-2812 (2022).
61. S. Singh *et al.*, Targeting the spliceosome through RBM39 degradation results in exceptional
responses in high-risk neuroblastoma models. *Sci Adv* **7**, eabj5405 (2021).
62. L. M. Luh *et al.*, Prey for the Proteasome: Targeted Protein Degradation-A Medicinal
Chemist's Perspective. *Angew Chem Int Ed Engl* **59**, 15448-15466 (2020).
63. M. M. Aboudounya, R. J. Heads, COVID-19 and Toll-Like Receptor 4 (TLR4): SARS-CoV-2
May Bind and Activate TLR4 to Increase ACE2 Expression, Facilitating Entry and Causing
Hyperinflammation. *Mediators Inflamm* **2021**, 8874339 (2021).
64. H. Kato, T. Fujita, RIG-I-like receptors and autoimmune diseases. *Curr Opin Immunol* **37**, 40-
45 (2015).
65. S. S. Burkart *et al.*, High-resolution kinetic characterization of the RIG-I-signaling pathway and
the antiviral response. *Life Sci Alliance* **6**, (2023).
66. P. Backes *et al.*, Role of annexin A2 in the production of infectious hepatitis C virus particles.
J Virol **84**, 5775-5789 (2010).
67. K. Giersch *et al.*, Hepatitis delta virus persists during liver regeneration and is amplified
through cell division both in vitro and in vivo. *Gut* **68**, 150-157 (2019).
68. O. Grünvogel *et al.*, DDX60L Is an Interferon-Stimulated Gene Product Restricting Hepatitis C
Virus Replication in Cell Culture. *J Virol* **89**, 10548-10568 (2015).
69. S. Wust, P. Schad, S. Burkart, M. Binder, Comparative Analysis of Six IRF Family Members
in Alveolar Epithelial Cell-Intrinsic Antiviral Responses. *Cells* **10**, (2021).
70. R. Patro, G. Duggal, M. I. Love, R. A. Irizarry, C. Kingsford, Salmon provides fast and bias-
aware quantification of transcript expression. *Nat Methods* **14**, 417-419 (2017).
71. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
72. M. Nowicka, M. D. Robinson, DRIMSeq: a Dirichlet-multinomial framework for multivariate
count outcomes in genomics. *F1000Res* **5**, 1356 (2016).
73. P. Shukla *et al.*, Cross-species infections of cultured cells by hepatitis E virus and discovery
of an infectious virus-host recombinant. *Proc Natl Acad Sci U S A* **108**, 2438-2443 (2011).

Fig. 1. A genome-wide CRISPR/Cas9 screen identifies potential TLR3 related factors.

(**A**) Schematic of the truncated BID (tBID) death reporter system. tBID expression and neomycin resistance (NeoR) are under control of the *IFIT1* promoter and the constitutive PGK promoter, respectively. Upon the stimulation by poly(I:C), the *IFIT1* promoter is activated via TLR3 signaling, inducing tBID and resulting in apoptosis. (**B**) Workflow of the CRISPR/Cas9 screen. PH5CH and Huh7 Lunet-TLR3 stably expressing the death reporter and Cas9 were transduced with the genome-wide lentiviral CRISPR sgRNA library and selected with puromycin for stable expression. TLR3 stimulation with poly(I:C) induced expression of tBID and thus apoptosis in case of intact TLR3 signaling. Then, surviving TLR3-deficient cells were enriched and collected for Next Generation Sequencing (NGS). (**C** to **E**) 50 candidate genes were selected and subjected to siRNA silencing for 48 h; then the cells were stimulated by 50 µg/ml poly(I:C) supernatant feeding for 8 h. *IFIT1* mRNA was measured by RT-qPCR. Knockdown of the candidates up-regulated (**C**) or down-regulated (**D**) *IFIT1* mRNA expression in Huh7 Lunet-TLR3 cells. Knockdown of *RBM39, PCF11, PTPRT* and *KDM2A* in PH5CH cells reduced *IFIT1* mRNA expression in PH5CH cells (**E**). siUNC93B1, siTLR3 and siTRIF were used as positive controls. Relative expression to siNT-treated samples was

707 normalized on GAPDH. Data are from three biological replicates (n = 2), error bars indicate standard
708 deviation (SD).
709

Fig. 2. RBM39 is crucial for the TLR3 pathway.

(**A**) Simplified schematic of the TLR3-IRF3 pathway. (**B** to **E**) RBM39 rescue experiment. PH5CH cells expressing RBM39.Esc or empty vector were transfected with siRBM39 or siNT/siTRIF as controls for 48 h, and then supernatant-fed with 50 µg/ml poly (I:C) for 6 h. *IFIT1* (**B**), *ISG15* (**C**), *MxA* (**D**) and *CXCL10* (**E**) mRNA was measured by RT-qPCR, RBM39 protein expression was measured by immunoblotting (**B**, right panel). (**F** to **H**) PH5CH cells were transfected with siRNA for 48h and then fed with 50 µg/ml poly(I:C) in the supernatant for 6 h. Secreted IFNβ protein (**F**) was measured by ELISA. Simplified schematic of the TLR3-NF-kB pathway (**G**). TNFAIP3 (**H**, left) and IL6 (**H**, right) mRNA was quantified by RT-qPCR. (**I**) The chemical structure of Indisulam. (**J**) PH5CH cells were treated with 1 µM Indisulam or DMSO as control for 48 h, and further stimulated with 50 µg/ml poly

721   (I:C) in the supernatant for 6 h. RBM39 degradation was measured by western blot (left). *IFIT1* mRNA
722   expression was measured by RT-qPCR (right). Relative expression to siNT control was shown,
723   mRNA fold change was normalized on *GAPDH*. Data are obtained from three biological replicates (n
724   = 3), error bars refer to SD. Statistical significance was assessed through Welsch's unpaired *t*-test.
725
726

Fig. 3. RBM39 participates in the RIG-I/MDA5 response.

(**A**) Simplified schematic of the RIG-I/MDA5-IRF3 pathway. (**B** to **D**) Huh7.5 cells with ectopic RIG-I
(**B**) or MDA5 (**C**) expression, and A549 cells (**D**) were transfected with siRBM39 or siNT/siMAVS as
control for 48 h and then transfected with 0.5 µg/ml poly(I:C) for 6 h. *IFIT1* mRNA was measured by
RT-qPCR and is shown relative to siNT. (**E**) A549 cells were treated Indisulam or DMSO as control for
48 h, and then transfected with 0.5 µg/ml poly (I:C) for 6 h. RBM39 degradation was determined by
immunoblotting. *IFIT1* mRNA was quantified by RT-qPCR. (**F** and **G**) A549 cells were transfected with
siRBM39 or siNT/siMAVS as control (**F**) or treated with 1 µM Indisulam or DMSO as control (**G**). 48
hours after treatment, cells were infected with Sendai virus (MOI= 1) for 24 h. *IFIT1* mRNA (left) and
Sendai virus P protein mRNA (right) were measured through qPCR. mRNA fold change was
normalized on *GAPDH*. Data are obtained from three biological replicates (n = 3), error bars refer to
SD. Statistical significance was assessed through Welsch's unpaired *t*-test.

740
741

Fig. 4. RBM39 controls the basal expression level of IRF3.

(**A**) PH5CH cells or PH5CH RBM39.Esc cells were transfected with siRBM39 or siNT/siTRIF as
controls for 48 h and then fed with 50 µg/ml poly (I:C) for 6 h. Phosphorylated IRF3 (p-IRF3), IRF3,
RBM39 and β-actin protein expression levels were assessed by western blot (left). Quantification of
three biological replicates was shown (right). (**B** and **C**) PH5CH cells (**B**) and primary human
hepatocytes (PHH) (**C**) were treated with increasing concentrations of Indisulam as indicated. IRF3,
RBM39 and β-actin protein expression levels were detected through immunoblotting (left). Cell
viability was measured via CellTiter-Glo luminescent cell viability assay (right). Quantification of three
independent experiments was expressed as average and is shown under each band. Protein
expression was normalized on β-actin, relative protein levels to siNT or DMSO control were shown.
Data shown are from three biological replicates (n = 3), error bars indicate SD. Statistical significance
was assessed through Welsch's unpaired *t*-test.

755
756

Fig. 5. Global proteomic, transcriptomic and splicing analysis upon modulation of RBM39 abundance.

PH5CH cells were treated with siRBM39 vs. siNT or Indisulam vs. DMSO after 48 h. The proteome was measured by mass-spectrometry. The transcriptome and splicing were measured through RNA-sequencing. (**A** and **B**) Volcano plot of the proteomic data for RBM39 knockdown (A) or Indisulam-treated (B) samples, the numbers of down- and up-regulated factors were shown under the plot. Factors with a fold change >2 or <0.5 and a p value <0.05 were highlighted with orange (A) or green (B). All factors with a p value <0.05 but lower than 2fold change are shown in black. IRF3, RBM39, RIG-I, MDA5, STAT1 and STAT2 were highlighted. (**C** and **D**) Volcano plot of DEGs in siRBM39- (C) and Indisulam-treated (D) samples, the numbers of down- and up-regulated factors were shown under the plot. DEGs with a fold change >2 or <0.5 and a p value <0.05 were highlighted with red (C) and blue (D). All factors with a p value <0.05 but lower than 2fold change are shown in black. IRF3, RBM39, RIG-I, MDA5, STAT1, STAT2, IFNAR1, IFNAR2, IFNLR1, IL10RB were highlighted. (**E**) RNA-seq data is presented as differential expressed gene (DEG) and different transcript usage

772   (DTU). The overview of DEG and DTU numbers in each condition and overlaps are indicated. RNA-
773   seq data are from three biological replicates (n = 3), proteomic data are from four replicates (n=4).
774   Proteomic data was analyzed via Perseus1.6.15.0. DEG and DTU analysis of individual genes were
775   performed using DESeq2 and DRIMseq, respectively.

776
777

Fig. 6. RBM39 controls basal expression of IRF*3* and *RIG-I and MDA5.*
(**A**) DEG and DTU analysis of *IRF3* mRNA. (**B**) Simplified schematic of the TLR3 pathway. (**C**) IRF3
was ectopically expressed in PH5CH IRF3 KO cells and cells were transfected with siRBM39 or
siNT/siTRIF. 48 h after the knockdown, cells were fed with 50 µg/ml poly (I:C) in the supernatant for 6
h. *IFIT1* mRNA expression levels were measured by RT-qPCR. (**D** and **E**) DEG and DTU analysis of
RIG-I (**D**) and *MDA5* (**E**) mRNA. (**F**) Simplified schematic of RIG-I/MDA5-IRF3 signaling. (**G**) A549
cells overexpressing IRF3 were transfected with siRBM39 or siNT/siMAVS as controls for 48 h. After
the knockdown, cells were transfected with 0.5 µg/ml poly (I:C) in the supernatant for 6 h. *IFIT1*
mRNA expression levels were measured by RT-qPCR. (**H**) Simplified schematic of the RIG-I/MDA5-
NF-κB pathway. (**I**) A549 cells were silenced with siRBM39 or siNT/siTRIF for 48 h and then
transfected with 0.5 µg/ml poly (I:C) for 6 h. *TNFAIP3* and *IL6* mRNA expression levels were
measured by RT-qPCR. Data shown are from three biological replicates (n = 3), error bars indicate
SD. Statistical significance was assessed through Welsch's unpaired *t*-test.

Fig. 7. RBM39 impacts on IFN-JAK-STAT signaling.

(**A** to **D**) DEG and DTU analysis of *IFNAR2* (**A**), *IL10RB* (**B**), *STAT1* (**C**) and *STAT2* (**D**) mRNA. (**E**) Simplified schematic of the type I and type III IFN pathways. (**F** and **G**) PH5CH cells were transfected with siRBM39 or siNT as control for 48 h and then stimulated with IFNα2 (**F**) or IFNλ1 (**G**) for 24 h. *IFIT1* mRNA was measured by qPCR. Relative expression to siNT control is shown, mRNA fold change was normalized on *GAPDH*. Data are obtained from three biological replicates (n = 3), error bars refer to SD. Statistical significance was assessed through Welsch's unpaired *t*-test.

Supplementary Materials

Supplementary materials and methods

Supplementary materials and methods

Primary mouse hepatocytes isolation:

C57BL6/N wild type mice (9-10 weeks, male) were housed in the specific-pathogen free (SPF) barrier at the Interfakultäre Biomedizinische Forschungseinrichtung (IBF) animal facility at Heidelberg University (Germany). Mice are maintained on a standard diet with ad libitum access to food and water under a constant light-dark cycle. Animal experiment was approved by the Regierungspräsidium Karlsruhe (T47/22). Mice were sacrificed, the abdomen was cleaned with ethanol and then opened expose inferior cava vein and portal vein. The inferior cava vein was Cannulated and washed with pre-warmed Liver Perfusion Medium (ThermoFisher, USA) for about 3-4 minutes until the liver became blood free. Afterwards, the liver was perfused with Liver Digest Medium (ThermoFisher, USA) until it was digested. The liver without gallbladder was transferred to a Petri dish. Cold Hepatocyte Wash Medium (ThermoFisher, USA) was added and the organ liberating cells were de-capsuled. Subsequently, the cells were sequentially passed through 100 μm and 70 μm cell strainers into a 50 mL falcon tube and centrifuged at 50g, 4°C for 2 minutes, to separate liver non-parenchymal cells (NPC) that remain in the supernatant. Next, hepatocytes (HC) pellet was washed with Hepatocyte Wash Medium (ThermoFisher, USA) once and re-suspended in William's E Medium, Glutamax Supplement (ThermoFisher, USA) with 4% FBS, 1% pen-strep and seeded 5×10^4 cells/cm^2 in a collagen-coated plate (Collagen I Rat Protein, Tail, ThermoFisher, USA).

CRISPR/Cas9 knockout screening:

For genome-wide CRISPR/Cas9 screen, PH5CH and Huh7-Lunet-TLR3 cells were transduced with lentiviral vectors containing the tBID death reporter and Cas9 (lentiCRISPRv2). Neomycin and puromycin were added for selection. After that, PH5CH and Huh7-Lunet-TLR3 cells stably expressing the tBID death reporter and Cas9 were transduced with lentiviral vectors containing the GeCKOv2.0 human genome-wide sgRNA library (Addgene, USA) at MOI=0.3. The GeCKOv2.0 human genome-wide sgRNA library includes a total of 122,411 unique sgRNAs targeting 19,050 genes (with 6 sgRNAs per gene) and 1,864 miRNAs (with 4 sgRNAs per miRNA) and a total number of 1,000 different non-targeting sgRNAs as controls.

After 24 hours, puromycin was added for selection. On day 7 post-transduction, 50ug/ml poly (I:C) or PBS as control were added to the medium to exclusively activate the TLR3 pathway. The experiments were performed in 2 independent biological replicates with 2 technical replicates each. Surviving cells were collected at 72 h post-stimulation, genomic DNA was extracted using NucleoSpin Blood L kit (Macherey-Nagel, Germany) according to manufacturer´s instructions. Amplicons were amplified, barcoded by PCR, and purified again via agarose gel separation using NucleoSpin Gel and PCR Clean-up kit (Macherey-Nagel, Germany), before being pooled and submitted to GATC Biotech

854 (Germany), for Illumina NGS (Illumina HiSeq, 50 bp single reads, 240 million reads total, 20% PhiX
855 DNA).
856
857 NGS data first was de-multiplexed to allocate the respective reads in the pooled population to each
858 sample. Reads were counted for each gRNA in every sample, and then normalized to the total
859 number of reads in this sample as well as the complexity sample taken one day after transduction of
860 the lentiviral library, yielding normalized read counts.
861
Mean values of normalized read counts of the 4 technical replicates were generated for each of the 4
experiments. Based on mean values, fold changes were calculated between poly(I:C)- and PBS-
treated samples. gRNAs with a higher fold change than a threshold were considered as primary hits.
The threshold was adjusted for each cell line to yield the about 2000 top-ranked gRNAs for each
experiment. Thresholds were 2.5x change in Huh-7-Lunet-TLR3 cells and 7x change in tPH5CH cells.
Preliminary hits were furthermore filtered for occurrences of at least 5 upregulated gRNAs per gene in
all datasets combined, and the occurrence of at least one gRNA per gene in at least 3 independent
datasets. This yielded 50 candidate genes and 5 candidate miRNAs.
870

871 **CRISPR/Cas9 knockout clones:**

872 To generate PH5CH IRF3 single knock-out (KO) clones, PH5CH cells were transduced with the
873 lentiCRISPRv2-IRF3 and lentiviral vectors encoding for Cas9 (see "Plasmids"). After puromycin
874 selection, cells were seeded into 96-well plates for screening of stable single clones.
875

Real time quantitative PCR:

Total RNA was isolated from cultured cells using the NucleoSpin RNA Plus kit (Macherey-Nagel,
Germany). RT-qPCR was performed as described before (*28*). For gene expression analysis, cDNA
as generated from RNA samples using High-Capacity cDNA Reverse Transcription Kit (Thermo
Fisher Scientific, USA), and was used for qPCR analysis with the 2x iTaq Universal SYBR Green
Supermix (Bio-Rad, Germany). Reactions were performed on a CFX96 Touch Real-Time PCR
Detection System (Bio-Rad, Germany) as follows: 95°C for 3 minutes, 95°C for 10 and 60°C for 30
seconds. mRNA fold change was calculated normalizing on Glyceraldehyde-3-phosphate
dehydrogenase (GAPDH) mRNA expression relative to untreated / uninfected / mock transfected
cells. For IRF3 isoforms mRNA measurement, cDNA was prepared by reverse transcription kit
(Thermo Fisher Scientific, USA) using oligo(dT)18 primers (Thermo Scientific, Germany). mRNA
expression fold change was normalized on GAPDH and Hydroxymethylbilane synthase
(HMBS) expression levels relative to siNT-transfected cells. Primers IRF3-228+203 were designed to
target both IRF3-228 and IRF3-203 stretches, other IRF3 primers were designed to target specific
exons or introns of individual isoforms. IRF3-203 mRNA expression was calculated by subtracting the
expression of IRF3-228 from that of IRF3-228 + IRF3-203. All qPCR primers are listed in Table S5.
892

Western blot:

Cells were lysed in protein lysis buffer (50mM TRIS HCl pH7.4, 150mM NaCl, 1% Triton-X 100,
protease inhibitor) on ice for 45 min. Cell lysates were then centrifuged at 14,000 g for 15 min at 4 °C
to exclude cell debris. Supernatants were collected and mixed with 6x Lämmli buffer (0.375M Tris-
HCl, 9% SDS, 50% Glycerol, 0.03% Bromophenol Blue, 9% β-mercaptoethanol), boiled at 95°C for 5
min and cooled down on ice for 2 min. Samples were separated by 10% SDS-PAGE and then blotted
onto polyvinylidene difluoride (PVDF) membranes (Merk Millipore, Germany). Membranes were then
blocked with 5% milk in PBST (0.05% Tween) or 5% BSA in TBST (0.01% Tween) and incubated with
the indicated antibodies. Membranes were imaged via the ECL Plus Western Blotting Detection
System (Pierce, GE Healthcare, Little Chalfont, UK) according to the instructions of the manufacturer.
Signal was recorded using the Advance ECL Chemocam Imager (Intas Science Imaging, Germany).
Quantification of western blot was performed by Fiji.
905

Lentivirus preparation and transduction:

5 x 10^6 HEK293T cells were seeded in 10 cm dishes 16 h before transfection. The next day, the target plasmid (5.14 µg) together with pCMV-Gag-Pol (5.14 µg) and pMD2-VSVG (1.71 µg) were mixed in 400µl OptiMEM (Gibco, USA). 364 µl of OptiMEM were separately mixed with 36µl polyethylenimine (PEI). The two mixtures were merged and incubated for 20 min at RT, and then added to the cells as previously described (*68*). After 2 days, lentivirus was collected and filtered at 0.45 µm filter (Sigma Aldrich, Germany) to exclude cell debris, and store at -80 °C or used for transduction. PH5CH-IRF3, A549-IRF3 stable cells were produced by transduction of lentiviral particles encoding for IRF3 under control of the Rosa26 promoter (IRF3 sgRNA resistant mutant) as previously described (*28*) in the abovementioned PH5CH IRF3 KO cells. PH5CH-KDM2A, PH5CH-RBM39.Esc, PH5CH-RBM39 G268V were generated by lentiviral transduction of particles containing corresponding plasmids.

Cell Viability Assay:

Cell viability upon transduction of tBID death reporter was assessed in 96-well plates through WST-1 assay (Roche, Switzerland) according to the manufacturers' instructions. Cell viability of Indisulam-treated cells was accessed by CellTiter-Glo® 2.0 Assay (Promega, Germany) according to the manufacturer's instructions.

ELISA:

Supernatants were collected from stimulated cells, IFNβ protein levels were measured by LumiKineTMXpress hIFN-β 2.0 (InvivoGen, Germany) according to the manufacturer's instructions.

Luciferase reporter assay:

pGL3-982 or pGL3-149 Firefly reporter, together with pRL-CMV Gaussia (Promega, USA) reporter as reference were transfected into PH5CH RBM39 knockdown cells using Lipofectamine2000 (Thermo Fisher Scientific, USA) according to the manufacturer's instructions. After 24 h, cells were lysed by addition of Luciferase lysis buffer (1% Triton X-100, 25mM Glycyl Glycin, 15mM MgSO4, 4mM EGTA, 10% Glycerol, pH7.8, 10mM DTT) and luciferase activity (Relative Ligh Units, RLU) was measured by a Mitras LB940 luminometer (Berthold Technologies, Germany). RVFV-RLuc assay was performed as described before (*69*). At 24 hours after infection, infected cells were washed once with PBS and immediately lysed with luciferase lysis buffer. Cell lysates were stored frozen at -80°C to ensure proper lysis and the Renilla luciferase signal was measured by injecting 300 µl of luciferase assay buffer (15 mM K$_3$PO$_4$ [pH7.8], 25 mM Glycyl-Glycin [pH 7.8], 15 mM MgSO$_4$, 4 mM EGTA) supplemented with 1.43 coelenterazine (102173, PJK, Germany) followed by read-out of the signal with a 480 nm high-sense filter (480m20BREThs, Berthold) using the Mitras2 multimode plate reader (LB942, Berthold).

RNA sequencing (RNA seq)

RNA-sequencing and RNA quality control was performed by the Genomics and Proteomics Core facility of the German Cancer Research Centre using the Illumina Hiseq NovaSeq 6000 with RNA seq reads in 2 x 100 bp format. For sample preparation, 1.5 x 10^5 cells were seeded in a 6-well plate. Cells were either transfected with siRBM39 or siNT as control, or treated with 1mM Indisulam or the same amount of DMSO as control. 48 hours after transfection, cells were lysed and total RNA was extracted via NucleoSpin RNA Plus kit (Macherey-Nagel, Germany) following the manufacturer's instructions. RNA was quantified using OD measurement at 260 nm.

Human reference transcriptome (Human Gencode v40) mapping and quantification of the transcripts were performed using the salmon package with GC bias correction (*70*). Differential expression analysis was performed using DESeq2 (*71*), and DTU analysis was performed with DRIMseq (*72*).

Virus infection:

For Sendai virus (SeV) infection, 6 x 10^4 A549 cells were transfected with siRNA RBM39 or treated with 1 µM Indisulam/DMSO. After 48 h, the cells were infected with SeV (MOI=1) in PBS with 0.3% BSA at room temperature (RT). One hour after infection, cells were washed with phosphate-buffered saline (PBS) and complete medium was added. Cells were collected after 24 h for further analysis.

For vesicular stomatitis virus (VSV) infection, 6 x 10^4 A549 cells were treated with 1 µM Indisulam or DMSO control for 48h. Afterwards, cells were infected with recombinant VSV-G (rVSV-G) (MOI=0.1) for 24 h. Cells were collected for further analysis.

For Rift Valley Fever Virus (RVFV) infection, 2 x 10^4 A549 cells were seeded into 24-well plates. The following day, cells were treated with 0.1 µM Indisulam or DMSO for 48 hours, and then inoculated with RVFV-RLuc at a MOI of 0.01 in DMEM with 2% FCS. After 24 hours of infection, cells washed once with PBS, lysed in 100 µl of luciferase lysis buffer, and frozen at -80 °C for further analysis.

For hepatitis E virus (HEV) (Genotype 3 Kernow C1 P6 accession number: JQ679013, a kind gift from Suzanne Emerson, NIH) (*73*) and hepatitis D virus (HDV) infection, 2.5 x 10^4 HepG2/C3A cells and HepG2-NTCP cells, respectively, were seeded in collagen (Corning, USA)-coated plates and treated with 0.1 µM Indisulam or same amount of DMSO for 48h. Subsequently, cells were infected with HEV (MOI=4) in MEM with 10% FBS or HDV (MOI=4) in DMEM with 1.5% DMSO, 4% PEG. 24 hours after infection, cells were washed with PBS. Cells were incubated for 5 days and collected for further analysis.

Mass spectrometry (MS)

PH5CH cells were treated with siRBM39 or Indisulam for 48 h, siNT and dimethyl sulfoxide (DMSO) were used as control. Afterwards, cells were harvested and lysed in 300 µl sodium dodecyl sulfate (SDS) lysis buffer (4% SDS, 10 mM dithiothreitol (DTT), 55 mM iodoacetamide (IAA), 50 mM Tris/HCl pH 7.5), and heat-inactivated (95°C for 10 min). The samples were sonicated (4°C, 15x 30 sec on and 30 sec off setting) and the protein lysate was precipitated using pre-chilled acetone (−20°C) at a final concentration of 80% acetone (v/v) overnight. Precipitates were pelleted and washed two times with pre-chilled 80% acetone (-20°C). Pellets were air dried at room temperature and resolubilized by adding 40 µl thiourea buffer (6 M urea, 2 M thiourea (U/T) in 10 mM HEPES, pH 8.0). Protein concentration was determined by BCA assay (Pierce) according to the manufacturer's instructions and adjusted to 50µg with thiourea buffer. Lysates were digested overnight (25°C, 16h, 1200 rpm) using LysC (1:50 w/w, enzyme/protein, Wako) and trypsin (1:50 w/w, enzyme/protein, Promega) per replicate. The samples were prepared for C18-stage tipping by adding 1/5 of loading buffer (10% acetonitrile (ACN), 3% trifluoroacetic acid (TFA)) to inactivate proteases (TFA) and enhance solubility (ACN) of samples. Peptides were desalted with 3-layer C18 stage tips (3M). Columns were equilibrated with 200 µl methanol (RT, centrifuge 300xg, 4min), washed twice with 200 µl washing buffer (0.5% acetic acid (AA)), loaded with the sample, washed twice with 200 µl washing buffer (0.5% AA) and eluted (80% acetonitrile, 0.5% AA). All steps were executed with 500 g at RT. The eluate was evaporated (45°C, vacuum) and resuspended in MS buffer (2% acetonitrile, 0.3% TFA). Peptide concentrations were measured at 280 nm (Nanodrop 2000, Thermo Scientific), and subsequently equalized to 0.25 µg/µl. Peptides were loaded onto a 2 cm trapping column (75 µm diameter; ReproSil-Pur C18-AQ 5 µm resin; Dr. Maisch) at 5 ul/min 0.1% formic acid (FA) in water using the UltiMate 3000 RSLCnano System (Thermo Fisher). Peptides were eluted onto a 42 cm analytical column (75 µm diameter; ReproSil-Pur C18-AQ 1.9 µm resin; Dr. Maisch) over a 120 min gradient at a flowrate of 300 nl/min and the column oven operating at 50 °C, using solvent B (0.1% FA, 5% DMSO in acetonitrile) starting from 4% to 32% over 110 min, then to 80% over 2 min, and kept at 80% for other 2 min, to 2% over 2min, and kept at 2% for 3 min. nanoLC solvent A was 0.1% FA, 5% DMSO in HPLC grade water.

Spectra were acquired with an Orbitrap Eclipse Tribrid mass spectrometer (Thermo Fisher Scientific) running Tune 3.5 and Xcalibur 4.5, and equipped with a nano-electrospray source (Thermo Fisher Scientific). Spray voltage was set to 2.1 kV, RF lens at 40 %, and heated capillary at 275 °C. The Eclipse mass spectrometer was operated in data-independent acquisition (DIA) and positive ionization mode. MS1 full scans (360–1300 m/z) were recorded at a resolution of 120,000 (Orbitrap) using a normalized automatic gain control (AGC) target value of 100% and maximum injection time of

1017   50 ms. Peptides were fragmented using higher energy collision induced dissociation (HCD) collision
1018   energy set to 30%. MS2 scans (200–1800 m/z) were acquired over a total of 40 DIA segments with
1019   variable isolation windows, and with a window overlap of 1 m/z. The scan resolution in the Orbitrap
1020   was set to 30,000 with a normalized AGC target value of 1000% and a maximum injection time of 54
1021   ms.
1022
Raw MS data files of the experiments conducted in data-independent acquisition (DIA) mode were
processed with DIA-NN version 1.8.1 (PMID 31768060). An in-silico predicted spectral library
composed of the human proteome (UniprotKB reference proteome, UP000005640, download Okt
2021) and common contaminants (MaxQuant contaminants.fasta) with trypsin as digestion enzyme
and one missed cleavage specified was generated in DIA-NN. Subsequently the acquired raw files
were processed in library-free mode using DIA-NN default settings and the match between runs
function enabled. The R package iq was used to generate MaxLFQ protein abundance output with a
Lib.Q.Value and Lib.PG.Q.Value filter setting of 0.01 (PMID 31909781). The ProteinGroups output file
was processed with Perseus (version 1.6.15.0). The intensities were expressed as Log2(x), imputed
(replacing missing values from normal distribution, with.0.3; down shift 1.8), and annotation included.
Statistical significance evaluated by "two-sample tests" (Student's T-test with permutation-based FDR
0.05 and 250 number of randomizations).
1035
1036

1037

Fig. S1. Efficiency of the tBID death reporter and the rescue of KDM2A.

(**A**) PH5CH cells (upper panel) or Huh7-Lunet cells, with or without ectopic TLR3 (lower panel), stably expressing IFIT1-tBID or empty-tBID death reporter as control were supernatant-fed with the indicated concentrations of poly(I:C), or a same volume of PBS, for the indicated time. Cell viability was determined at the indicated time points using the WST-1 assay. Data are from four independent biological replicates (n=4). (**B**) PH5CH expressing empty vector or *KDM2A* were transfected with the indicated siRNAs for 48 h. Cells were subsequently stimulated through poly(I:C) supernatant feeding. *IFIT1* mRNA was measured by qPCR. Data are from two biological replicates (n=2), error bars represent SD.

1047

1048

Fig. S2. Rescue of TLR3 response by ectopic RBM39 expression in Huh7 Lunet-TLR3 cells.

(**A**) Huh7 Lunet-TLR3 cells expressing RBM39.Esc or empty vector were transfected with siRBM39 or siNT/siTRIF as controls for 48 h and then fed with 50 µg/ml poly (I: C) in the supernatant for 6 h. *IFIT1* mRNA was measured by RT-qPCR (left), RBM39 expression was measured by western blot (right). Data are from three biological replicates (n = 3), error bars refer to mean ± SD. Statistical significance was assessed through Welsch's unpaired *t*-test.

1055

1056

Fig. S3. Indisulam treatment in different virus infection models.

(**A** to **D**) A549 cells (**A** and **B**), HepG2/C3A (**C**) cells and HepG3-NTCP cells (**D**) were pre-treated with Indisulam or same amounts of DMSO as control for 48 h, and subsequently infected with Luc-RVFV (MOI= 0.01) for 24 h (**A**), rVSV-G (MOI=0.1) for 24 h (**B**), HEV (MOI= 4) (**C**) or HDV (MOI=4) (**D**) for 5 days, respectively. *IFIT1* mRNA (left) and viral mRNA (right) were measured by qPCR. mRNA fold change was normalized on *GAPDH*. Data are derived from three biological replicates (n = 3), error bars refer to SD. Statistical significance was assessed through Welsch's unpaired *t*-test.

1064
1065

1066

Fig. S4. Impact of Indisulam-mediated RBM39 degradation on IRF3 in different cell lines.

(**A** to **D**) PH5CH-RBM39 and PH5CH-RBM39 G268V cells (**A**), A549 cells (**B**) and primary mouse
hepatocyte (PMH) (**C**) were treated with Indisulam at the indicated concentrations for 48 h. IRF3,
RBM39 and β-actin protein expression levels were measured by western blot. Quantification of three
independent experiments is shown as average number under the respective bands. (left) Cell viability
was measured via CellTiter-Glo luminescent cell viability assay (**B** and **C**, right). Data are from three
biological replicates (n = 3), error bars indicate SD.
1074

1075

Fig. S5. Analysis of individual protein expression levels from proteomics data

The protein level of GAPDH (**A**), RBM39 (**B**), IRF3 (**C**), RIG-I (**D**), MDA5 (**E**), STAT1 (**F**) and STAT2 (**G**) in siRBM39/siNT or Indisulam/DMSO-treated PH5CH samples. Data are derived from three biological replicates (n = 4), error bars refer to SD. Statistical significance was assessed through Welsch's unpaired *t*-test.

1081

Fig. S6. The transcription and splicing of *IRF3*, *IFNAR1* and *IFNLR1 mRNA*

(**A**) PH5CH cells were transfected with siRBM39 or siNT as controls. 48 h after transfection, *IRF3* and *RBM39* mRNA was measured by qPCR. (**B**) Schematic of the firefly luciferase reporter harboring the full-length IRF3 promoter (pGL3-982-firefly) or core IRF3 promoter (pGL3-149-firefly). (**C**) PH5CH cells were transfected with pGL3-982-firefly or pGL3-149-firefly, pGL3-CMV-Gaussia was used as reference. Luciferase activity of firefly was normalized on that of Gaussia. Relative luciferase activity (Relative Light Units, RLU) is shown. (**D**) Schematic of *IRF3* isoforms (left). Specific primers targeting different isoforms are shown as arrows. *IRF3* isoforms/total *IRF3* ratio in PH5CH cells transfected with siRBM39 or siNT as control were identified by qPCR. (**E** and **F**) DEG and DTU analysis of *IFNAR1* (**E**) and *IFNLR1* (**F**) mRNA. DEG and DTU analysis of individual genes were performed using DESeq2 and DRIMseq, respectively. Data shown are from three biological replicates (n = 3), error bars indicate SD. Statistical significance was assessed through Welsch's unpaired *t*-test.

 Table S5: Primers and antibodies

Primers for cloning	
Primer number	Sequence
#1	GATCATCGATGATCACTGGAATTTGATTCAGAAT
#2	GATCATCGATGATCCTTAAACCAATGGCAGAAGC
#3	GCATACTAGTCTAGTCCATCCCATTTCTGGCTAAG
#4	CACCGCTGGTGTCGCAGCTGGACC
#5	AAACGGTCCAGCTGCGACACCAGC
#6	GCAGGCGCGCCATGGAACCCGAAGAAGAAAGGATTCG
#7	TTTACTAGTTTAGCTGATCTTCTGTATCAGCTTC
#8	GCAGGCGCGCCATGGCAGACGATATTGATATTG
#9	TTTACTAGTTCATCGTCTACTTGGAACCAG
#10	CCTGCAGGCGCGCCATGGCAGACG
#11	GGAAGAGGTACATATGCTGCTGTTATCATTTTTCCTGCGAACCATCTACCATGCAATGCATTGACAGCAGCAATAGC
#12	GCTATTGCTGCTGTCAATGCATTGCATGGTAGATGGTTCGCAGGAAAAATGATAACAGCAGCATATGTACCTCTTCC
#13	CGTAGTTTACTAGTTCATCGTCTACTTGGAACCAGTAG
#14	CAAGCCTCAGACAGTGGTTC
#15	TCCGAGATCCAATTGACTGACTAACCAGGGCAGGATCCGTGGC
#16	GTTAGTCAGTCAATTGGATCTCGGACAACTGGAGGGCGTGGCC
#17	CGTAGTTTACTAGTTCAGCTCTCC
#18	CGGGGTACCAGAGTTAGGAGGGAGCCTC
#19	GGAAGATCTGCCCTTTTTTGGGTTTCC
Primers for RT-qPCR		
---	---	---
Gene name	Forward sequence	Reverse sequence
RBM39	GCAAGGACAGTCTTCTGTATGC	CGACGAACTCCACATAAGCAA
GAPDH	GAAGGTGAAGGTCGGAGTC	GAAGATGGTGATGGGATTTC
ISG15	CGCAGATCACCCAGAAGATCG	TTCGTCGCATTTGTCCACCA
MxA1	GTTTCCGAAGTGGACATCGCA	CTGCACAGGTTGTTCTCAGC
IFNB1	ATGACCAACAAGTGTCTCCTCC	GGAATCCAAGCAAGTTGTAGCTC
IL6	ACTCACCTCTTCAGAACGAATTG	CCATCTTTGGAAGGTTCAGGTTG
TNFAIP3	TCCTCAGGCTTTGTATTTGAGC	TGTGTATCGGTGCATGGTTTTA
IFIT1	GAAGCAGGCAATCACAGAAA	TGAAACCGACCATAGTGGAA
IFNLR1	CCCAAGGGTAAGAGCTTCGAT	CCTTCATATTTTACTGACATGGACAAG
IL10RB	TTGCTGTGGTGCGTTACAAG	CTTTCAGGTGCTGTGGAAGAGA
CXCL10	GGCATTCAAGGAGTACCTCTCTC	TGGACAAAATTGGCTTGCAGGA
HBMG1	TATGGCAAAAGCGGACAAGG	CTTCGCAACATCACCAATGGA
IRF3-Total	GACTTTTCCCAGCCAGACAC	TCCAGAATGTCTTCCTGGGTAT
IRF3-203+228	TACCCGGAGCTCCAAGACA	TTCCCATGGTCCGGCCTA
IRF3-228	CCCTTCATTGTAGGCTCCT	CACAGAACCAGAGGGCATAG
IRF3-222	CGACCTTCCATCGTAGGAGTT	GAAGTACTGCCTCCACCATTG

IRF3-202	AGCTACCCGGAGCTCCAA	CTCGGCCCAGGCCTAC
IRF3-219	CCTTCATTGTAGGTGAGTGAG	CCTGAGTTGTTAACCACTGTG
SeV-P	CAAAAGTGAGGGCGAAGGAGAA	CGCCCAGATCCTGAGATACAGA
VSV-P	TCTCGTCTGGATCAGGCGG	TGCTCTTCCACTCCATCCTCTTGG
HEV	GGTGGTTTCTGGGGTGAC	AGGGGTTGGTTGGATGAA
HDV	GCGCCGGCYGGGCAAC	TTCCTCTTCGGGTCGGCATG
Antibodies		
Target	product	company
RBM39	HCC1 Polyclonal Antibody	ThermoFisher Invitrogen
IRF3	IRF-3 (D6I4C) XP® Rabbit mAb	Cell Signaling
p-IRF3	Phospho-IRF-3 (Ser396) (4D4G) Rabbit mAb #4947	Cell Signaling
β-actin	Monoclonal Anti-b-Actin Clone AC-15	Sigma-Aldrich

1098

1099