Listen to yourself: How motivational interviewing shifts food choices and craving-related brain responses to healthier options.

Authors: Belina Rodrigues[a], Iraj Khalid[a], Benjamin Flament[a*], Solene Frileux[a*], Martine Rampanana[a*], Hippolyte Aubertin[a], Jean-Yves Rotge[a,b] Philippe Fossati[a,b], Leonie Koban[a,c], Liane Schmidt[a]

Affiliations:

[a] Control-interception-attention team, Institute du Cerveau et de la Moelle Epinière (ICM), UMR 7225, U1127, Institut National de la Santé et de la Recherche Médicale/Centre National de la Recherche Scientifique/Sorbonne Universités, Hôpital Pitié-Salpêtrière, 75013 Paris, France.

[b] Departement de Psychiatrie Adulte, Hôpital Pitié-Salpêtrière, Assistance Publique Hopitaux Publiques (APHP)

[c] Centre de Recherche en Neurosciences de Lyon (CRNL), CNRS, INSERM, Université Claude Bernard Lyon 1, Bron, France.

* equal contribution, listed in alphabetical order

Corresponding authors: Belina Rodrigues and Liane Schmidt,

E-mail: belina.rodrigues@icm-institute.org; liane.schmidt@icm-institute.org

Abstract

Changing one's habits is challenging. Motivational interviewing (MI) has been proposed as a communication-based approach to overcome this challenge. Here, we tested how change and sustain talk suggestions, elicited by MI, influenced value-based dietary decision-making and responses of a recently developed neurobiological craving signature (NCS) in healthy female adults. We found that following change talk, compared to sustain talk, participants' food choices and activity in the ventromedial prefrontal cortex were more driven by the healthiness and less by the tastiness of food. These findings were paralleled by lower NCS responses to tasty food after change compared to sustain talk. Further, following change talk, participants' body mass indices moderated the NCS decoding of healthy and tasty food choices. These results show that MI can shift value-based decision-making and reduce craving-related brain marker responses to highly palatable food items. The findings contribute to a better understanding of behavioral change toward healthier eating.

Keywords: motivational interviewing, behavioral change, value-based decision making, ventromedial prefrontal cortex, neurobiological craving signature

Main

Modifiable behaviors such as tobacco use, physical inactivity, unhealthy eating habits, and harmful alcohol consumption increase the risk of non-communicable diseases and account for a significant percentage of worldwide disability and mortality (Griswold et al., 2018; Afshin et al., 2019; Murray et al., 2020). In particular, unhealthy eating habits are associated with an increased risk of all-cause mortality (English et al., 2021) and carry a significant economic burden greater than smoking and physical inactivity (Scarborough et al., 2011). However, unhealthy eating behavior is rising worldwide (Imamura et al., 2015).

Given the consequences of unhealthy dietary habits, it is essential to understand the mechanisms of interventions designed to promote behavior change. In this study, we investigated the neurocognitive effects of a prominent communication-based approach to behavior change – motivational interviewing (MI) (Lundahl et al., 2010; Lundahl et al., 2013; VanBuskirk and Wetherell, 2014; Bully et al., 2015; Frost et al., 2018). MI is a collaborative clinical method, a goal-oriented style of communication focusing on the language of change. It aims to strengthen motivation and commitment to a specific goal. In more detail, MI elicits personal statements for change, called change talk, and deals with reasons against change, i.e., sustain talk, within an atmosphere of acceptance and compassion (Miller and Rollnick, 2013). Research on the associations between MI and behavioral outcomes showed that a higher proportion of change talk over sustain talk is associated with reduced risk-seeking behavior (Magill et al., 2018; Magill et al., 2019).

Previous work on the neural correlates of MI has focused on substance use disorders such as alcohol and cannabis use. These studies have provided convergent evidence that listening to change talk increased the recruitment of brain regions associated with cognitive regulation in terms of self-control among cannabis users (Feldstein Ewing et al., 2013) and dampened the activation of brain areas related to craving and reward when viewing alcohol-related cues (Feldstein Ewing et al., 2011). These findings suggest that MI might generate a cognitive shift towards self-control and a reduction in craving. However, there is a lack of consensus about the joint behavioral and neural effects when participants mentally transition between change and sustain talk during decision-making. Further, no previous studies have investigated whether MI can alter

responses in *a priori* and independently established brain markers of craving, such as the recently developed neurobiological craving signature (Koban et al., 2023).

Self-control propensities have been shown to play a crucial role in advantageous economic choices (Rangel, 2013). Models of economic choice propose that the decision process involves two stages (Rangel et al., 2008). During valuation, various attributes of choice alternatives are integrated into stimulus values, which approximate a decision-maker's preferences. Stimulus values are then compared during action selection to lead to a preferred choice. Findings from research on the neurobiology of valuation show that value is computed within the brain's reward and valuation system formed by the vmPFC, ventral striatum, and posterior cingulate cortex (for review, see Padoa-Schioppa (2011) and Bartra et al. (2013)). These value computations can be modulated by self-control, defined by the ability to forgo tasty and liked food in favor of healthier food to improve the quality of eating habits. Such value modulation has been reported to involve the activation of the dlPFC (Hare et al., 2011).

However, sticking to more self-controlled, healthier eating is tricky in everyday life. The vmPFC has also been shown to encode sensitivity to food stimuli and food craving (Tang et al., 2012; Volkow et al., 2017; Li et al., 2023) and has strong positive weights in a recently validated neural signature of food and drug craving – the NCS (Koban et al., 2023). Food cravings relate to eating and weight gain over time (Boswell and Kober, 2016). Motivational interviewing has been found effective for such situations with prominent cravings, for example, substance use disorders (Frost et al., 2018). However, its effectiveness for the self-control of dietary decision-making and on NCS responses is unknown.

Here, we adopted this value-based decision-making framework and used the neurobiological craving signature (NCS) to predict that listening to change versus sustain talk should (1) change food valuation during dietary decision-making by shifting the weights of healthiness and tastiness of food, (2) alter how the vmPFC as a central hub of the brain's valuation system encodes these attributes of food during economic choice formation, and (3) reduce craving related brain responses (as quantified by the NCS) to tasty food items.

Results

Behavioral results

We first tested whether change compared to sustain talk affected the value assigned to food stimuli, i.e., how much participants wanted to eat the food items. Two analogous linear mixed effects models (LME) of stimulus value were used: LME 1 was fitted to stimulus values assigned to healthy and tasty food. This model revealed that change talk significantly increased food value compared to sustain talk (main effect of change versus sustain talk: β = 0.05, SE = 0.02, 95%CI [0.006 – 0.08], t = 2.3, p =0.02). The model further showed a non-significant main effect of type of attribute (healthiness versus tastiness: β = -0.03, SE = 0.03, 95% CI [-0.09 – 0.027], t = -1.1, p = 0.29) and a significant interaction type of talk by attribute type (β = 0.06, SE = 0.02, 95% CI [0.03 – 0.09], t = 3.64, p =0.0004) such that after change talk (compared to sustain talk), stimulus value was greater for healthy food items. LME 2 was fitted to stimulus values assigned to unhealthy and untasty food. This model also found a significant interaction talk by attribute type (β = -0.06, SE = 0.02, 95% CI [-0.11 – -0.01], t = -2.5, p = 0.01), but a non-significant main effect of talk (β = -0.04, SE = 0.03, 95% CI [-0.09 – 0.012], t = -1.5, p = 0.14), and a significant main effect of attribute type (β = 0.3, SE = 0.03, 95% CI [0.20 – 0.34], t = 7.9, p = 1.3e-12). Both models were controlled for BMI.

To describe these interactions in more detail, posthoc t-tests and Figures 1a and 1b show that healthy food was chosen more often (t(31)= 4.1, p <.001, paired, two-tailed t-test, Cohen's d = 0.72), and unhealthy food was rejected more often (t(31) = -2.4, p = 0.02, paired, two-tailed t-test, Cohen's d = -0.42) after change talk than after sustain talk. This difference in stimulus values was non-significant for tasty foods (change versus sustain talk) (t(31) = -0.6, p = 0.55, paired, two-tailed t-test, Cohen's d = -0.1) nor for untasty foods (t(31)= 0.81, p = 0.42, paired, two-tailed t-test, Cohen's d = 0.14).

To further test how listening to change versus sustain talk influenced how much healthiness and tastiness information were integrated into stimulus values across all food choice trials, the individual beta values from a general linear model fit to stimulus value ratings were entered into a second-level analysis of variance (SI Table 1). Specifically, the ANOVA tested how much first-level interaction terms of interest (i.e.,

CTxTR, CTxHR, STxTR, and STxHR) differed as a function of the type of talk (change
versus sustain talk) and type of attribute (healthiness versus tastiness).

In line with food valuation-related analyses, the second-level random effect ANOVA
revealed a main effect of attribute type (F(1,124)=22.6, p<0.001, $\eta^2 = 0.14$), a small
borderline effect of talk (F(1,124)= 4,12, p=0.045, $\eta^2 = 0.02$), and a significant
interaction talk by attribute (F(1,124) =32.6, p<0.001, $\eta^2 = 0.15$) (Figure 1c). As shown
in Figure 1c, after listening to a change talk, healthiness predicted significantly more
food stimulus value than after listening to a sustain talk (CTxHR vs STxHR: t(31) =
3.9, p<0.001, Cohen's d = 0.7, paired, two-tailed t-test). In contrast, after listening to a
sustain talk, the tastiness predicted more food stimulus values than after listening to a
change talk (STxTR vs CTxTR: t(31)= 2.3, p=0.029, Cohen's d = 0.4, paired, two-tailed
t-test).

Figure 1. Behavioral results in N=32 participants. Boxplots display 95% CI for value
assigned after change (CT) and sustain (ST) talk to **(a)** food stimuli during healthy and tasty
food choices, **(b)** value assigned to food stimuli during unhealthy and untasty food choices,
and **(c)** the prediction of food stimulus value by tastiness and healthiness ratings across all
food choices. Change talk (and compared to sustain talk) increased the weight of healthiness
on the food choice which translated into choosing more often healthy and rejecting more often
unhealthy food. *** p<0.001, **p<0.01, * p<0.05 for a significant interaction attribute type
(healthiness versus tastiness) and talk type (change versus sustain talk) and paired, two-tailed
t-tests.

In summary, these complementary statistical models converged on the finding that
change talk compared to sustain talk increased the participant's propensity to consider
the healthiness more and the tastiness less during food valuation.

 <u>Effects of talk on food value computation within the prefrontal cortex</u>

At the time of food choice, significant activation within the brain's reward and valuation system encompassing the vmPFC, ventral striatum, and posterior cingulate cortex correlated significantly with food stimulus value (p_{FWE}<0.05 family-wise error correction on the peak and cluster level, Figure 2a, SI Table 2). This activation did not significantly differ between change and sustain talk conditions. We next tested whether listening to change versus sustain talk affected the integration of healthiness and tastiness within the value-encoding vmPFC region of interest. As shown in Figure 2b, the vmPFC-value encoding region of interest correlated stronger with healthiness at the time of food choice after listening to change compared to sustain talk (small volume correction, p_{FWE}<0.05). The vmPFC ROI correlation to tastiness did not differ between change and sustain talk conditions.

Figure 2. Brain responses to stimulus value and healthiness. (a) Stimulus value encoding in the brain. Statistical parametric maps show brain activation in correlation to stimulus value at the time of food choice. Significant voxels in yellow and orange survived a whole brain threshold of p_{FWE}<0.05, family-wise error corrected on the peak level, and are shown for display at p<0.001 uncorrected. **(b) Contrast between change versus sustain talk for healthiness encoding in the brain.** SPMs show activation of the vmPFC region of interest that correlated stronger to healthiness ratings at the time of food choice after the change than after sustain talk. Significant voxels in yellow survived small volume correction (SVC p_{FWE} < 0.05) within the vmPFC and are displayed at an uncorrected whole brain threshold of p<0.005. The boxplot shows 95% confidence intervals of beta values extracted from the peak activation in response to healthiness ratings within the vmPFC (MNI_{peak}=-8, 46, -8) for illustration. Colored dots correspond to individual participants. All SPMs are superimposed on the average structural brain image.

Prediction of food stimulus value by the neural craving signature (NCS)

We next tested whether the observed behavioral effects of change and sustain talk on healthy and tasty food choices were paralleled by differential responses to food stimulus values (i.e., choices) of the recently validated neurobiological craving signature (Koban et al., 2023).

A linear mixed effects model (LME 3), analogous to LMEs 1 and 2, was fit to the NCS responses to food stimulus value. The model revealed that overall NCS responses were higher for tasty food choices than healthy food choices (β = -1.9, SE = 0.63, 95% CI [-3.2 – -0.69], t = -3.1, p = 0.0025). The model also indicated a non-significant effect of talk (β = -0.07, SE = 0.32, 95% CI [-0.70 – 0.56], t = -0.21, p = 0.83) and, in line with our prediction, a significant interaction talk by attribute (β = 0.86, SE = 0.32, 95% CI [0.23 – 1.5], t = 2.7, p = 0.008). This interaction effect (see Figure 3b) was driven by a significant reduction in NCS responses to tasty food choices in the change, compared to the sustain talk condition (t(31) = -1.9, p = 0.03, one-tailed t-test, Cohen's d = -0.35). In addition, the NCS tended to respond more to healthy food choices following change talk than following a sustain talk (t(31) = 1.5, p = 0.07, one-tailed t-test, Cohen's d = 0.28).

Lastly, we explored the role of inter-individual differences in NCS responses to healthy and tasty food choices after change and sustain talk. In the change (but not in the sustain) talk condition, body mass index (BMI) correlated significantly and positively with the NCS responses to healthy food choices (Pearson's r = 0.46, p = 0.009). On the contrary, it correlated negatively with NCS responses to tasty food choices (r = -0.36, p = 0.04). This suggests that the effects of change talk on the NCS might be especially pronounced in participants with higher body mass.

Figure 3. **Neural craving signature prediction of food stimulus value in change and**
sustain talk conditions. (a) Pattern of the neural brain signature of craving (NCS) with
positive weights displayed in yellow and red and negative weights shown in blue (at p<0.05
uncorrected for display). **(b)** The boxplots depict 95% CI for the NCS response to food stimulus
value during healthy and tasty food choice trials. Error bars correspond to standard errors.
Horizontal bars indicate the median, and red points mean values. Individual points in black
indicate outlier participants. **(c)** Correlations between BMI and the neurobiological craving
signature (NCS) response to the value assigned to healthy and tasty food in the change talk
condition.

Discussion

Motivational interviewing is an evidence-based intervention technique in the context
of addiction and behavioral change more broadly. This study aimed to characterize
the effects of motivational interviewing on dietary decision-making and craving-related
brain responses to food choices. We combined fMRI with behavioral testing of value-
based dietary decision-making to contrast in a within-participant design two types of
suggestions, i.e., one in favor, the other against the change of eating habits elicited by
motivational interviewing.

In line with our predictions, change talk led to a greater weighting of the healthiness of
food items and a lower weighting of tastiness, an effect that was paralleled by
predefined craving-related responses in the neural craving signature (NCS) – a
recently established brain marker of food and drug craving (Koban, et al., 2023).
Moreover, the participant's body mass index moderated the NCS responses to tasty
and healthy food choices. These findings suggest an interaction between motivational
interviewing and body weight on craving-related brain circuits.

Previous research has looked at the neural mechanisms of motivational interviewing
in the context of alcohol and substance use (see Mackiewicz Seghete et al. (2022),

Grodin et al. (2019) or Feldstein Ewing et al. (2016) for examples). Here, we
investigated by using univariate and multivariate brain-behavior analyses an aspect
that has yet to be reported: food preferences, which were measured by an incentive-
compatible value-based decision-making task. Sticking to healthier food choices is
difficult, and eating-related conditions such as diabetes are a worldwide problem.
Here, we tested in hungry, normal to overweight participants the cognitive effects of
shifting between change and sustain talk in the laboratory context. Our findings in
healthy participants are therefore a proof of concept for motivational interviewing to
affect the basic mechanisms of value-based dietary decision-making. They further
align with an increased focus on prevention and the assumption that motivational
interviewing might also be helpful as a prevention strategy instead of an adjunct
treatment strategy (Foxcroft et al., 2016). Future work should test how the cognitive
shifting between change and sustain talk modulates food valuation and valuation-
related brain activation in populations struggling with food-addiction-like behavioral
habits and obesity.

Our behavioral finding of value modulation suggests that change talk might have
generated a form of dietary self-control (Hare et al., 2009; Hare et al., 2011;
Hutcherson et al., 2012; Harris et al., 2013; Tusche and Hutcherson, 2018; Schmidt
et al., 2018; Wilson et al., 2023). Previous studies of dietary decision-making and its
self-control have used instructions to incite participants to shift explicitly between
strategies, such as distancing from or focusing attention on the healthiness or
tastiness of food during incentive compatible dietary decision-making tasks (Hare et
al., 2009; Hare et al., 2011; Hutcherson et al., 2012; Harris et al., 2013; Tusche and
Hutcherson, 2018; Schmidt et al., 2018; Wilson et al., 2023). In line with this past work
participants of the current study performed a similar incentive compatible task in the
fasted, hungry state, which may have favored more indulgent decision-making under
sustain talk conditions, and at the same time may have also enhanced the conflict
between short-term taste and long-term health rewards of food under change talk
conditions. These two aspects are important ingredients to measure self-control
propensities during dietary decision-making. However, the current study also differed
from this past work in several, important and novel aspects. Change and sustain talk
was not always related to health or taste but to other aspects of food behavior (e.g.,
environmentally friendly food behavior). These reasons were unique and personal to

each participant. Moreover, participants were told to make food choices following their natural preferences and to remember the suggestions they listened to. The instructions also emphasized that both types of suggestions, change and sustain talk, were valid and that they should not favor one type of talk over another. Our findings thus suggest that patients and health professionals do not need to focus on the "health aspect" to elicit behavior change but instead on the personal reasons most relevant to a patient. This is not trivial, as not all patients are health-focused. While it remains an open question to what extent the observed behavioral effects generalize to real-world food choices, the scope of our study was to shed light on the primary cognitive and neural effects of shifting between change and sustain talk during economic food choice in a controlled laboratory setting.

The behavioral results were underpinned by the univariate finding that the vmPFC encoded food value during dietary decision-making irrespective of the type of talk. This result is in line with prior evidence that the vmPFC is a hub within the brain's valuation system that encodes expected and experienced rewards across different economic choice problems (Marsh et al., 2007; Forster and Brown, 2011; Rouault et al., 2019). On the behavioral level, the type of talk (i.e., change versus sustain talk) influenced how food preferences were formed, which was approximated by the integration of tastiness and healthiness into food stimulus values (i.e., food choices). In line with this behavioral finding, the value-encoding vmPFC region of interest correlated stronger to the healthiness of food at the time of choice following change talk than following sustain talk. This result aligns with findings from neuroeconomics that showed that the vmPFC computes the value signals that drive economic choices (Padoa-Schioppa, 2011). Past work from this stream of research has also found another brain region, the dorsolateral prefrontal cortex (dlPFC), to play a role in value modulation induced by cognitive regulation such as self-control during dietary decision-making (Hare et al., 2009; Hutcherson et al., 2012; Wilson et al., 2023). Here, the univariate brain analysis results were restricted to the vmPFC. We did not find evidence for the contribution of the dlPFC to dietary decision-making after change compared to sustain talk. Different experimental designs, participant samples, and contextual conditions might explain these odds. For example, the dlPFC might be more prominently recruited during economic choice formation when participants face a real-world conflict between change and sustain talk such as struggling with food addiction or obesity. Alternatively,

the dlPFC might also be more implicated in contexts where instructions to focus on specific attributes of food are more explicit.

Moreover, the modulatory effects observed by change and sustain talk on integrating healthiness and tastiness into stimulus value might be better explained and predicted by conjoint activation of more than one or two brain regions (Woo et al., 2017). Current multivariate, statistical approaches in the analysis of fMRI data use machine-learning algorithms to build predictive models – brain signatures to decode behavior from whole brain activation (Woo et al., 2017). Here, we used a recently validated neurobiological craving signature (NCS) that encompassed value and reward-related brain regions such as the vmPFC and ventral striatum, but also other regions such as the insula, anterior midcingulate cortex and temporal/parietal association areas (Koban et al., 2023). The NCS has been cross-validated to predict individual food and drug craving levels across different participant samples of drug and non-drug users to whom both drug and high palatable (i.e., tasty) food cues were presented. We showed that change versus sustain talk modulated NCS responses to healthy and tasty food choices. In more detail, similar to the effects of talk type on stimulus value, we found a significant interaction type of talk by attribute on the NCS response. NCS responses to *tasty* food choices were significantly reduced in the change compared to the sustain talk condition. This finding complements the effects of talk on stimulus value ratings, which was driven by a significant difference between change and sustain talk on *healthy* food choices. These differences might be explained by the fact that the NCS was trained to predict cravings for highly palatable food items and might, therefore, be more sensitive to tasty food choice trials in our data.

Moreover, the NCS responses to healthy and tasty food choices after change talk varied as a function of BMI. While these correlations should be interpreted with caution given the small sample size, they suggest that participants with higher BMI present greater NCS responses to healthy food choices and lower NCS responses to tasty food after change talk. Brain activation, especially within the brain's value and reward systems, which are a prominent part of the NCS, is influenced by weight status and hormones that signal energy states, such as leptin, which are directly related to body fat (Zanchi et al., 2017; Schmidt et al., 2021; Plassmann et al., 2022). Therefore, links between the brain, body composition, and energy-store signaling hormones could explain our correlational findings. Change talk could make craving- and reward-related

brain circuits more sensitive to such internal, afferent signals from the periphery. However, while these correlations were non-significant for sustain talk conditions, this might be due to the small sample size. Thus, these correlational results should be replicated in larger sample sizes that also include participants with obesity.

A few limitations need to be considered. Here, we report on the influence of client language on dietary decision-making. However, motivational interviewing is more than just eliciting change talk. It comprises two sets of skills: technical and relational (Miller and Rose, 2009). The technical skills, composed of open questions and reflections, elicit and strengthen change talk. The relational skills, comprising aspects such as empathy, ensure an atmosphere of acceptance and compassion. Both features aim at increasing the frequency and strength of change talk while handling sustain talk reasons (Miller and Rose, 2009; Frey et al., 2020). Interestingly, Magill et al. (2018) found a non-independence of technical and relational components of motivational interviewing, as the variance of the technical element was partly explained by varying levels of empathy. It has also been found that the relational aspect predicted more reflections of change, which, in turn, predicted more change talk (Villarosa-Hurlocker et al., 2019). These findings point to a synergistic effect of both sets of skills. However, the current study design combining value-based decision-making with fMRI did not allow for assessing the impact of change talk in conjunction with these relational components on real-world behavioral change propensities.

Additionally, the scope of our findings is restricted to the basic cognitive mechanisms of decision-making. In the real world, the medical appointment between patient and health professionals and the post-session processes may elicit many effects (Feldstein et al., 2011), which drive behavior change. Moreover, other behavior change approaches should be contrasted to motivational interviewing effects by future work to determine the specificity of basic cognitive mechanisms and how they might give rise to real-world behavioral change in the long term (Michie et al., 2011).

In conclusion, our findings indicate that MI leads to a shift in dietary decision-making paralleled by changes in specific brain circuits related to value-based decision-making and craving. These findings advance our understanding of the neurocognitive mechanism of communication-based approaches to behavior change, which may

have a place in regular medical appointments to facilitate sticking to healthier eating habits.

Methods

<u>Ethical considerations</u>

The study was approved by the local ethics committee and conducted in accordance with the Declaration of Helsinki. All participants provided written and informed consent.

<u>Study-pre-registration:</u>

This study was part of a still ongoing, larger-scale pre-registered study (https://clinicaltrials.gov/study/NCT05101863). It focused exclusively on the healthy participants (BMI<30) arm. A second deviation from the pre-registered protocol involved using a more recently validated neurobiological craving signature (Koban et al., 2023).

<u>Participants</u>

A total of 40 participants were recruited via public advertisement in the Paris area. The inclusion criteria were to self-identify as female, aged between 18 and 70 years old, right-handed, with normal to corrected-to-normal vision, no history of substance abuse or neurological or psychiatric disorder, no medication, and no MRI contraindication. Inclusion was limited to self-identified female participants to have a more homogenous group. Research on the physiology of appetite and eating behavior has shown that men and women differ in their food behavior (Rolls et al., 1991; Davy et al., 2006; Poldrack et al., 2008; Frank et al., 2010; Marino et al., 2011), such that women have been shown to display stronger brain activation to appetite-enhancing food stimuli (Rolls et al., 1991; Frank et al., 2010), despite a strong propensity for dietary self-restraint (Frank et al., 2010).

Eight participants were excluded from analysis due to problems with the MRI scanner (n=5), computer software to record the behavioral data (n=2), and the discovery of an fMRI contraindication on the day of the MRI visit (n=1) (Table 1).

415 **Table 1**. Socio-demographics and body composition of n=32 participants.

Mean age (SEM)	30 (1.82) years
Mean years of education after high school (SEM)	4.3 (0.25) years
High school graduates	3.1%
At least 2 years of college	3.1%
More than 2 years of college	93.8%
Body composition	
Mean BMI (SEM)	22.2 (0.49) kg/m^2
Range of BMI	17.0 – 27.3 kg/m^2
Mean body fat (SEM)	27.8 (1.3)%
Mean body water (SEM)	51.4 (4.6)%
Yale Food Addiction scale	
No Food addiction	93.8%
Food addiction	6.3%
HEI-2015	
Mean HEI-2015 (SEM)	45.6 (3.2)
HEI-2015 range (0-100)	13.7-78.8
Hunger ratings	
Mean hunger ratings at **baseline** (SEM)	3.6 (0.25) range (1 to 6.3)
Mean hunger ratings **after MRI** (SEM)	5.0 (0.19) range (2.7 – 6.7)
Mean IPAQ (SEM)	1688 (344) MET min/week

Hunger rating$_{before}$ vs hunger rating$_{after}$ = (t(30) = -8.2, p< 0.001, Cohen's d = -1.46, paired, two-tailed t-test.

<u>Experimental procedure</u>

The study was composed of two visits that took place one week apart.

<u>Visit 1</u>: Upon enrolment, sociodemographic data such as gender, age, and level and

years of education were collected. Then, each participant underwent a 24h dietary

recall to study food intake according to the manual adaptation of the Automated

Multiple-Pass Method (Raper et al., 2004). After obtaining a list of all food items eaten

on the previous day, the interviewer probed for frequently forgotten foods such as

savory snacks or candies. Then, the time and name of the eating occasion (e.g.,

breakfast) and a detailed description of the type of food item, cooking method, and other relevant aspects. Lastly, a final revision of the dietary intake took place (Raper et al., 2004). Whenever possible, visit 1 was scheduled so that seasonal changes (e.g., weekends, holidays such as Christmas) in the dietary habits would not be captured.

Following the 24h recall, all participants underwent a semi-structured brief (i.e., 20-30 minutes) motivational interviewing session, which was divided into four phases: The interviewer, who was a dietitian trained in MI, elicited the participants to reflect on the following: (1) Their dietary patterns and whether there was a desire for a change, and (2) the advantages and disadvantages of the behavior they would like to change. The experimenter then asked participants to (3) rate the importance and their readiness for change on a 10-point Likert scale and 4) verified again whether they still would like to change an aspect of their dietary habits. The brief motivational intervention was audio-recorded, and 5 change and 5 sustain talk statements were extracted for use during an fMRI session at the second visit one week later (for examples see SI Table 3). Phrases were chosen by agreement of two different raters trained in MI.

Visit 2:

A week after the first visit, participants underwent an fMRI session after overnight fasting. Before the fMRI session, and to ensure a balanced distribution of healthy, unhealthy, tasty, and untasty food stimuli across the two talk conditions, participants rated each food item in terms of healthiness and tastiness on a 4-point scale: strong no, no, yes, strong yes. Additionally, hunger ratings were measured on a 7-point Likert scale before and after the fMRI session. Importantly, these hunger ratings were averaged across the rating of three aspects of hunger: hedonic (i.e., How pleasant would it be to eat, now?), homeostatic (i.e., How much could you eat, now) and general (i.e., How hungry are you, now?).

To study the effect of change language, a modified version of a well-known value-based dietary decision-making task was presented to the participants (Hare et al., 2009; Hare et al., 2011; Hutcherson et al., 2012; Schmidt et al., 2018). It comprised 10 blocks of choice trials: Five blocks started with change talk statements and five with sustain talk statements. The order of change and sustain talk blocks was randomized. Each block counted 15 decision-making trials for 75 trials per talk condition. The blocks

started by instructing the participants to keep in mind the talk statement they will listen to. Then, a 15-second talk statement was played, and the corresponding verbatim could be read on the screen. Participants were asked to read and listen to their voices expressing the change or sustain talk statements, respectively. After listening and reading the suggestions they rated 15 food stimuli of varying tastiness and healthiness on whether they wanted to eat this food for real at the end of the experiment. Each choice trial started with the display of a food stimulus, and participants had three seconds to rate whether they wanted to eat the food item for real at the end of the experiment by using a 4-point Likert scale ranging from strong no to strong yes. Trials were separated by a jittered fixation cross (Figure 4). A short break was planned after 75 trials (i.e., 5 blocks) to reduce head movement. A brief practice session occurred outside the MRI beforehand, and a second practice session occurred within the MRI to adjust the headphones' volume.

Figure 4. **Dietary decision-making task.** Panels display screenshots of successive events within a change and sustain talk blocks. Each block started with a reminder to keep in mind the statement related to change/sustain current dietary habits. This information was followed by a change or sustain talk statement, which participants listened to and could also read on the screen. Following the statement, participants made 15 food choices. Food items with varying levels of healthiness and tastiness were presented. Each choice trial started with food displayed on the screen, and participants had to rate on a 4-point Lickert scale how much they wanted to eat the food for real at the end of the experiment. The choice was displayed on the screen when they hit a corresponding response button. A fixation cross-separated trials within each talk block. The duration of each event within these two example trials is displayed in seconds.

All food stimuli were selected from a database of 600 food pictures (contact Dr. Hilke

Plassmann for more information). The food pictures were presented on a computer

screen in form of high-resolution pictures (72 dpi). MATLAB and Psychophysics

Toolbox extensions (Brainard, 1997) were used for stimulus presentation and

response recording. Participants saw the food stimuli via a head-coil–based mirror and

indicated their responses using an fMRI compatible response box system. The hand

used to make choices and the display of the 4-point Likert scale (from left to right or

right to left) was counterbalanced across participants. The task was incentive

compatible, because one food was chosen out of chance for consumption at the end

of the experiment (Hare et al., 2009; Hare et al., 2011).

Body composition and questionnaires

After the fMRI session, anthropometric measures such as body fat percentage and

weight were obtained via Tanita® Body Composition Analyzer SC-240MA (Tanita

Corporation, Tokyo, Japan) and self-reported height. Body mass index (BMI) was

computed as Kg/m^2. The physical activity level was assessed via the short form of the

International Physical Activity Questionnaire (IPAQ-SF), which assesses the intensity

of physical activity and sitting time over the previous seven days. Estimates are given

in MET-min/week and time spent sitting (Craig et al., 2003). Participants filled in the

Yale Food Addiction Scale (YFAS) to measure signs of addictive-like eating behavior.

It comprises 35 items with eight response options ranging from "Never" to "Every Day."

Each question assesses one of the 11 diagnosis criteria defined originally for

substance abuse (e.g., tolerance, withdrawal, clinical Significance). The cut-off for

food addiction is a total score of at least two of these 11 diagnosis criteria and meeting

the criteria to be clinically significant (Gearhardt et al., 2016).

Dietary intake assessment

To ascertain the required nutrition-related variables, once the 24h-recall was collected,

energy intake (in Kcal), sodium and fatty acids quantities were ascertained through

Nutrilog SAS software (Nutrilog-Online 20231010). Additionally, to measure diet

quality, the healthy eating index (HEI) was computed (Krebs-Smith et al., 2018). The

HEI-2015 is based on the Dietary Guidelines for Americans 2015-2020 (for a

comparison between the American guidelines and French guidelines see SI table 4).

It has 13 items whose scoring follows adequacy- or moderation-based energy-adjusted intakes cut-offs, except for the fatty acids components. Each components weights differently into the final scoring (0-100, where higher scores indicate a healthier dietary pattern): total fruits (5 points), whole fruits (5 points), total vegetables (5 points), greens and beans (5 points), whole grains (10 points), dairy (10 points), total protein foods (5 points), fatty acids (polyunsaturated fatty acid plus monounsaturated fatty acid to saturated fatty acid ratio; (10 points), refined grains (10 points), sodium (10 points), added sugars (10 points), and saturated fats (10 points). Refined grains, sodium, added sugars, and saturated fats as components to be consumed in moderation have a reversed scoring (Krebs-Smith et al., 2018). The HEI score for the sample is shown in Table 1.

Imaging data acquisition

Brain images were collected on a 3 T Verio Siemens scan, using a 32-channel receive-only head coil. For the structural acquisition a whole-brain high-resolution T1-weighted structural scans were acquired for all participants with a magnetisation prepared rapid gradient echo (MPRAGE) sequence with the following parameters: TR = 2.4 s; TE = 2.05 ms; flip angle=9°, slice thickness=0.8mm. For the functional acquisition, a multi-band T2*-weighted multi-echoplanar images (mEPI) with BOLD contrast were acquired. To cover the whole brain with a repetition time of 1.25 seconds, the following sequence parameters were applied: echo times 14.8 ms; 33.4 ms and 52 ms; 48 slices collected in an interleaved manner; 3-mm slice thickness, FOV = 192 mm; flip angle = 68°, and an average of 646 volumes per participate session. Three echos were acquired to better accommodate for the compromise between spatial resolution and signal of the orbitofrontal context (Posse et al., 1999; Poser et al., 2006). To further minimise the signal drop out in the orbitofrontal cortex, an oblique acquisition orientation of 30° above the anterior–posterior commissure line was applied (Deichmann et al., 2003).

Imaging data pre-processing

Anatomical MPRAGE and functional mEPIs were processed using the Statistical Parametric Mapping – SPM12 – (Welcome Department of Imaging Neuroscience, Institute of Neurology, London, United Kingdom) using MATLAB version R2020b (The

MathWorks Inc., United States). Each participant's anatomical image was segmented into gray, white matter and cerebrospinal fluid using the SPM12 segmentation tool. The three echo images of each volume were summed into one EPI volume using the SPM 12 Image Calculator (Gowland and Bowtell, 2007; Poser and Norris, 2009; Kettinger et al., 2016). Then, the summed EPIs were spatially realigned and motion corrected, coregistered to the mean image, normalised to the Montreal Neurological Institute (MNI) space using the same transformation as the anatomical image, and spatially smoothed using a Gaussian kernel with a full-width-at-half-maximum of 8 mm.

Data analyses

Statistical tests were conducted with the Matlab Statistical Toolbox (Matlab 2020a, MathWorks), and JASP (JASP 0.16.1). Continuous variables were reported as mean and standard error of mean (SEM), and categorical variables as percentage.

Behavioral analyses

Linear mixed-effects models (LMEs) were fit to mean-centered stimulus value (SV, coded -2 for "strong no", -1 for "no", 1 for "yes" and 2 for "strong yes") ratings to test how much the type of talk (change talk coded 1, sustain talk coded -1) and the type of attribute (healthiness coded 1, tastiness coded -1) affected the valuation of food stimuli. Participant's tastiness and healthiness ratings were used to categorize trials into healthy, tasty, unhealthy, and untasty choice trials. In total, two LMEs were fit to (1) healthy food items and tasty food items (LME 1), and (2) unhealthy food items and untasty food items trials (LME 2), respectively, and the following equation (I.):

$$SV \sim 1 + attribute + talk + attribute \times talk + (1|participant)$$
$$+ (-1 + attribute|participant) + (-1 + talk|participant)$$

The model assumed random intercepts nested by participant number, and random slopes for the effect of type of talk (change versus sustain) and attribute (healthiness versus tastiness).

A similar third LME (LME 3) was fit to the neural craving signature responses to stimulus value (i.e., food choices) assigned to healthy and tasty food.

All LMEs were controlled for BMI. Post hoc, paired, one-tailed and two-tailed t-tests were used to determine the directionality of main effects and interactions.

To test how tastiness and healthiness attributes were integrated into stimulus value

following change and sustain talk statements, the mean-centered stimulus values (SV,

coded -2 for "strong no", -1 for "no", 1 for "yes" and 2 for "strong yes"), were fit to a

multilevel general linear model (GLM) following equation II:

II. $SV \sim \beta_0 + \beta_{Trial} + \beta_{HR}HR + \beta_{TR}TR + \beta_{ST}ST + \beta_{CT}CT + \beta_{ST \times HR}ST \times HR + \beta_{ST \times TR}ST \times$

$TR + \beta_{CT \times HR}CT \times HR + \beta_{CT \times TR}CT \times TR + \beta_{HR \times TR}HR \times TR + \epsilon$

On the first level analyses, the model included the following regressors: trial number

(β_{Trial}) to account for fatigue effects, health ratings (β_{HR}), and taste ratings (β_{TR}), and

two categorical indicators for sustain talk (β_{ST}) and change talk runs (β_{CT}). Additionally,

the model included four interaction terms: STxHR, CTxHR, STxTR, CTxTR. These

interactions tested for effects of healthiness and tastiness on stimulus value in trials

following change and sustain talk, respectively. Individual beta coefficients were then

fit into a second-level factorial analysis of variance (ANOVA) to test for main effects of

talk (change versus sustain talk), attribute rating (healthiness versus tastiness rating),

and the interaction talk type by attribute type. Post hoc, paired, two-tailed t-tests were

used to determine the directionality of interactions.

<u>Imaging analyses</u>

Analogous to the behavioral analyses, two multilevel general linear models (GLMs)

were fit to fMRI time series.

To detect the voxels involved in SV encoding, a first GLM included on the first level

four regressors of interest: (1) an indicator for type of talk with boxcar durations of 15

seconds, (2) choice onset regressor with a boxcar duration corresponding to reaction

times, (3) a parametric modulator at time of choice consisting of stimulus value, and

(4) missed trials with a boxcar duration of 3 seconds. The six realignment parameters

were added as regressors of non-interest to correct for head motion. Individual beta

images for this parametric modulator SV at the time of choice were then fit into a

second-level random effects analysis using one-sampled t-tests.

To identify the voxels involved in the encoding of healthiness and tastiness as a

function of the type of talk, a second GLM included nine regressors of interest: (1)

indicators for onset of change talk statements with boxcar durations of 15 seconds, (2)

choice onset regressors at time of food image display following change talk with

durations corresponding to the reaction time, two parametric modulator regressors at
time of food choice following change talk: (3) healthiness and (4) tastiness ratings, (5)
indicators for onset of sustain talk with boxcar durations of 15 seconds, (6) food choice
onset at time of food image display following sustain talk with a duration corresponding
to the reaction time, and two parametric modulator regressors: (7) healthiness and (8)
tastiness, (9) onset regressors for missed trials with boxcar duration of 3 seconds. The
6 realignment parameters were included as regressors of non-interest to control for
head movement. Then, individual beta images were fit into a second-level random
effects analysis using one-sampled t-tests, adding BMI as a covariate to the matrix.

<u>Definition of regions of interest (ROI) for small volume correction</u>
Regions of interest for small volume correction were functionally defined by the
significant, whole brain family wise error corrected activation within the vmPFC by
stimulus value across all trials (MNI = [-6, 50, -8], SI table 2).

<u>Responses of the neurobiological craving signature (NCS) to food stimulus value (i.e.,</u>
<u>choices)</u>

To assess how change and sustain talk altered craving-related brain responses (as
measured by the NCS, Koban et al., 2023), a general linear model (GLM 3) fit each
participant's BOLD signal. It included ten regressors of interest: (1) an indicator for
type of talk, (2) an onset regressor for tasty food choices following change talk, (3)
stimulus value as a parametric modulator of the tasty food choice onset regressor
following change talk, (4) an onset regressor for healthy food choices following change
talk, (5) stimulus value as a parametric modulator of the healthy food choice onset
regressor following change talk, (6) an onset regressor for tasty food choices following
sustain talk, (7) stimulus value as a parametric modulator of the tasty food choice
onset regressor following sustain talk, (8) an onset regressor for healthy food choices
following sustain talk, (9) stimulus value as a parametric modulator of the healthy food
choice onset regressor following sustain talk, and (10) onset regressors for missed
trials with boxcar duration of 3 seconds. The six realignment parameters were included
as regressors of non-interest to control for head movement.

NCS responses were quantified as the dot product between the contrast images of interest (i.e., on the parametric regressors 3 and 5 for tasty and healthy stimulus value at time of food choice) for each participant and the NCS weight map, plus a fixed intercept. This results in one scalar value per contrast image and participant, which can be further analyzed using standard statistical tests.

Data availability

Behavioral data is available on the study's OSF website.

Code availability

Code to analyse behavioral data and food decision making task scripts are available on study's OSF website. The NCS and code to apply it to any fMRI data (apply_ncs.m function) are publicly available on Github (https://github.com/canlab/Neuroimaging_Pattern_Masks/tree/master/Multivariate_signature_patterns/2022_Koban_NCS_Craving).

Acknowledgments

We thank Hilke Plassmann and Nathalie George for their insightful project discussion. We thank the CENIR team for their support and collaboration throughout the project. This project has received funding from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement n° 945298. BR is a Fellow of the Paris Region Fellowship Programme supported by the Paris Region. LK acknowledges funding from an ERC Starting Grant (SOCIALCRAVING, 101041087). However, the views and opinions expressed are those of the authors only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible. The funders had no role in study design, data analysis, manuscript preparation, or publication decisions. LS acknowledges funding from Fondation NRJ – Institute de France.

 Authors contributions

BR and LS conceived the original idea of the study and designed the study protocol and materials with advice from PF. BR, JYR, SF, IK, BF included participants. BR collected data with help from MR, BF, SF, and IK. BR and HA analyzed the data under the supervision of LS and advice from LK. BR and LS drafted the manuscript, and PF and LK contributed to the final text.

References

Afshin A et al. (2019) Health effects of dietary risks in 195 countries, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. The Lancet. doi:10.1016/S0140-6736(19)30041-8.

Bartra O, McGuire JT, Kable JW (2013) The valuation system: a coordinate-based meta-analysis of BOLD fMRI experiments examining neural correlates of subjective value. Neuroimage 76:412–427.

Boswell RG, Kober H (2016) Food cue reactivity and craving predict eating and weight gain: a meta-analytic review. Obes Rev 17:159–177.

Brainard DH (1997) The Psychophysics Toolbox. Spat Vis 10:433–436.

Bully P, Sanchez A, Zabaleta-del-Olmo E, Pombo H, Grandes G (2015) Evidence from interventions based on theoretical models for lifestyle modification (physical activity, diet, alcohol and tobacco use) in primary care settings: A systematic review. Prev Med 76 Suppl:S76-93.

Craig CL, Marshall AL, Sjostrom M, Bauman AE, Booth ML, Ainsworth BE, Pratt M, Ekelund U, Yngve A, Sallis JF, Oja P (2003) International physical activity questionnaire: 12-country reliability and validity. Med Sci Sports Exerc 35:1381–1395.

Davy SR, Benes BA, Driskell JA (2006) Sex differences in dieting trends, eating habits, and nutrition beliefs of a group of midwestern college students. J Am Diet Assoc 106:1673–1677.

Deichmann R, Gottfried J, Hutton C, Turner R (2003) Optimized EPI for fMRI studies of the orbitofrontal cortex. Neuroimage 19:430–441.

English LK et al. (2021) Evaluation of Dietary Patterns and All-Cause Mortality: A Systematic Review. JAMA Netw Open 4:e2122277.

Feldstein ESW, Filbey FM, Hendershot CS, McEachern AD, Hutchison KE (2011) Proposed Model of the Neurobiological Mechanisms Underlying Psychosocial Alcohol Interventions: The Example of Motivational Interviewing*. J Stud Alcohol Drugs 72:903–916.

Feldstein Ewing SW, Chung T, Caouette JD, Ketcherside A, Hudson KA, Filbey FM (2016) Orbitofrontal cortex connectivity as a mechanism of adolescent behavior change. Neuroimage. doi:10.1016/j.neuroimage.2016.12.076.

Feldstein Ewing SW, Filbey FM, Sabbineni A, Chandler LD, Hutchison KE (2011) How psychosocial alcohol interventions work: a preliminary look at what FMRI can tell us. Alcohol Clin Exp Res 35:643–651.

Feldstein Ewing SW, McEachern AD, Yezhuvath U, Bryan AD, Hutchison KE, Filbey FM (2013) Integrating brain and behavior: evaluating adolescents' response to a cannabis intervention. Psychology of Addictive Behaviors 27:510–525.

Forster SE, Brown JW (2011) Medial prefrontal cortex predicts and evaluates the timing of action outcomes. Neuroimage 55:253–265.

Foxcroft DR, Coombes L, Wood S, Allen D, Almeida Santimano NML, Moreira MT (2016) Motivational interviewing for the prevention of alcohol misuse in young adults. Cochrane Database Syst Rev 7:CD007025.

Frank S, Laharnar N, Kullmann S, Veit R, Canova C, Hegner YL, Fritsche A, Preissl H (2010) Processing of food pictures: influence of hunger, gender and calorie content. Brain Res 1350:159–166.

Frey AJ, Lee J, Small JW, Sibley M, Owens JS, Skidmore B, Johnson L, Bradshaw CP, Moyers TB (2020) Mechanisms of Motivational Interviewing: a Conceptual Framework to Guide Practice and Research. Prevention science : the official journal of the Society for Prevention Research. doi:10.1007/s11121-020-01139-x.

Frost H, Campbell P, Maxwell M, O'Carroll RE, Dombrowski SU, Williams B, Cheyne H, Coles E, Pollock A (2018) Effectiveness of Motivational Interviewing on adult behaviour change in health and social care settings: A systematic review of reviews. PLoS One 13:e0204890.

Gearhardt AN, Corbin WR, Brownell KD (2016) Development of the Yale Food Addiction Scale Version 2.0. Psychology of Addictive Behaviors 30:113–121.

Gowland PA, Bowtell R (2007) Theoretical optimization of multi-echo fMRI data acquisition. Phys Med Biol 52:1801–1813.

Griswold MG et al. (2018) Alcohol use and burden for 195 countries and territories, 1990–2016. A systematic analysis for the Global Burden of Disease Study 2016. The Lancet 392:1015–1035.

Grodin EN, Lim AC, MacKillop J, Karno MP, Ray LA (2019) An Examination of Motivation to Change and Neural Alcohol Cue Reactivity Following a Brief Intervention. Front Psychiatry 10:408.

Hare TA, Camerer CF, Rangel A (2009) Self-control in decision-making involves modulation of the vmPFC valuation system. Science 324:646–648.

Hare TA, Malmaud J, Rangel A (2011) Focusing attention on the health aspects of foods changes value signals in vmPFC and improves dietary choice. J Neurosci 31:11077–11087.

Harris A, Hare T, Rangel A (2013) Temporally dissociable mechanisms of self-control: early attentional filtering versus late value modulation. J Neurosci 33:18917–18931.

Hutcherson CA, Plassmann H, Gross JJ, Rangel A (2012) Cognitive regulation during decision making shifts behavioral control between ventromedial and dorsolateral prefrontal value systems. J. Neurosci. 32:13543–13554.

Imamura F, Micha R, Khatibzadeh S, Fahimi S, Shi P, Powles J, Mozaffarian D (2015) Dietary quality among men and women in 187 countries in 1990 and 2010. A systematic assessment. The Lancet Global Health 3:e132-e142.

Kettinger Á, Hill C, Vidnyánszky Z, Windischberger C, Nagy Z (2016) Investigating the Group-Level Impact of Advanced Dual-Echo fMRI Combinations. Front Neurosci 10:571.

Koban L, Wager TD, Kober H (2023) A neuromarker for drug and food craving distinguishes drug users from non-users. Nat Neurosci 26:316–325.

Krebs-Smith SM, Pannucci TE, Subar AF, Kirkpatrick SI, Lerman JL, Tooze JA, Wilson MM, Reedy J (2018) Update of the Healthy Eating Index: HEI-2015. J Acad Nutr Diet 118:1591–1602.

Li G, Hu Y, Zhang W, Wang J, Ji W, Manza P, Volkow ND, Zhang Y, Wang G-J (2023) Brain functional and structural magnetic resonance imaging of obesity and weight loss interventions. Mol Psychiatry 28:1466–1479.

Lundahl B, Moleni T, Burke BL, Butters R, Tollefson D, Butler C, Rollnick S (2013) Motivational interviewing in medical care settings: a systematic review and meta-analysis of randomized controlled trials. Patient Educ Couns 93:157–168.

Lundahl BW, Kunz C, Brownell C, Tollefson D, Burke BL (2010) A Meta-Analysis of Motivational Interviewing. Twenty-Five Years of Empirical Studies. Research on Social Work Practice 20:137–160.

Mackiewicz Seghete KL, Filbey FM, Hudson KA, Hyun B, Feldstein Ewing SW (2022) Time for a paradigm shift: The adolescent brain in addiction treatment. Neuroimage Clin 34:102960.

Magill M, Apodaca TR, Borsari B, Gaume J, Hoadley A, Gordon REF, Tonigan JS, Moyers T (2018) A meta-analysis of motivational interviewing process. Technical, relational, and conditional process models of change. J Consult Clin Psychol 86:140–157.

Magill M, Janssen T, Mastroleo N, Hoadley A, Walthers J, Barnett N, Colby S (2019) Motivational interviewing technical process and moderated relational process with underage young adult heavy drinkers. Psychol Addict Behav 33:128–138.

Marino M, Masella R, Bulzomi P, Campesi I, Malorni W, Franconi F (2011) Nutrition and human health from a sex-gender perspective. Mol Aspects Med 32:1–70.

Marsh AA, Blair KS, Vythilingam M, Busis S, Blair RJR (2007) Response options and expectations of reward in decision-making: the differential roles of dorsal and rostral anterior cingulate cortex. Neuroimage 35:979–988.

Michie S, van Stralen MM, West R (2011) The behaviour change wheel: A new method for characterising and designing behaviour change interventions. Implementation Science 6:42.

Miller WR, Rollnick S (2013) Motivational interviewing. Helping people change / William R. Miller and Stephen Rollnick. New York, NY: Guilford Press.

Miller WR, Rose GS (2009) Toward a theory of motivational interviewing. Am Psychol 64:527–537.

Murray C, Aravkin A, Zheng P, Abbafati C, Abbas K, Abbasi-Kangevari K (2020) Global burden of 87 risk factors in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. The Lancet 396:1223–1249.

Padoa-Schioppa C (2011) Neurobiology of economic choice: a good-based model. Annu Rev Neurosci 34:333–359.

Plassmann H, Schelski DS, Simon M-C, Koban L (2022) How we decide what to eat: Toward an interdisciplinary model of gut-brain interactions. Wiley Interdiscip Rev Cogn Sci 13:e1562.

Poldrack RA, Fletcher PC, Henson RN, Worsley KJ, Brett M, Nichols TE (2008) Guidelines for reporting an fMRI study. Neuroimage 40:409–414.

Poser BA, Norris DG (2009) Investigating the benefits of multi-echo EPI for fMRI at 7 T. Neuroimage 45:1162–1172.

Poser BA, Versluis MJ, Hoogduin JM, Norris DG (2006) BOLD contrast sensitivity enhancement and artifact reduction with multiecho EPI: parallel-acquired inhomogeneity-desensitized fMRI. Magn Reson Med 55:1227–1235.

Posse S, Wiese S, Gembris D, Mathiak K, Kessler C, Grosse-Ruyken M-L, Elghahwagi B, Richards T, Dager SR, Kiselev VG (1999) Enhancement of BOLD-contrast sensitivity by single-shot multi-echo functional MR imaging. Magn Reson Med 42:87–97.

Rangel A (2013) Regulation of dietary choice by the decision-making circuitry. Nat Neurosci 16:1717–1724.

Rangel A, Camerer C, Montague PR (2008) A framework for studying the neurobiology of value-based decision making. Nat Rev Neurosci 9:545–556.

Raper N, Perloff B, Ingwersen L, Steinfeldt L, Anand J (2004) An overview of USDA's Dietary Intake Data System. Journal of Food Composition and Analysis 17:545–555.

Rolls BJ, Fedoroff IC, Guthrie JF (1991) Gender differences in eating behavior and body weight regulation. Health Psychol 10:133–142.

Rouault M, Drugowitsch J, Koechlin E (2019) Prefrontal mechanisms combining rewards and beliefs in human decision-making. Nat Commun 10:301.

Scarborough P, Bhatnagar P, Wickramasinghe KK, Allender S, Foster C, Rayner M (2011) The economic burden of ill health due to diet, physical inactivity, smoking, alcohol and obesity in the UK: an update to 2006-07 NHS costs. J Public Health (Oxf) 33:527–535.

Schmidt L, Medawar E, Aron-Wisnewsky J, Genser L, Poitou C, Clément K, Plassmann H (2021) Resting-state connectivity within the brain's reward system predicts weight loss and correlates with leptin. Brain Commun 3:fcab005.

Schmidt L, Tusche A, Manoharan N, Hutcherson C, Hare T, Plassmann H (2018) Neuroanatomy of the vmPFC and dlPFC Predicts Individual Differences in Cognitive Regulation During Dietary Self-Control Across Regulation Strategies. J Neurosci 38:5799–5806.

Tang DW, Fellows LK, Small DM, Dagher A (2012) Food and drug cues activate similar brain regions. A meta-analysis of functional MRI studies. Physiol Behav 106:317–324.

Tusche A, Hutcherson CA (2018) Cognitive regulation alters social and dietary choice by changing attribute representations in domain-general and domain-specific brain circuits. Elife 7.

VanBuskirk KA, Wetherell JL (2014) Motivational interviewing with primary care populations: a systematic review and meta-analysis. J Behav Med 37:768–780.

Villarosa-Hurlocker MC, O'Sickey AJ, Houck JM, Moyers TB (2019) Examining the
influence of active ingredients of motivational interviewing on client change talk. J
Subst Abuse Treat 96:39–45.
Volkow ND, Wise RA, Baler R (2017) The dopamine motive system: implications for
drug and food addiction. Nat Rev Neurosci 18:741–752.
Wilson DJ, HajiHosseini A, Hutcherson CA (2023) Recruitment of dlPFC during
dietary self-regulation predicts the transience of regulatory effects. Soc Cogn
Affect Neurosci 18.
Woo C-W, Chang LJ, Lindquist MA, Wager TD (2017) Building better biomarkers:
brain models in translational neuroimaging. Nat Neurosci 20:365–377.
Zanchi D, Depoorter A, Egloff L, Haller S, Mählmann L, Lang UE, Drewe J, Beglinger
C, Schmidt A, Borgwardt S (2017) The impact of gut hormones on the neural
circuit of appetite and satiety: A systematic review. Neuroscience & Biobehavioral
Reviews 80:457–475.