Gene expression variation across genetically identical individuals predicts reproductive traits

Amy K. Webster[1,2], John H. Willis[1], Erik Johnson[1], Peter Sarkies[3], Patrick C. Phillips[1]

1. Institute of Ecology and Evolution, University of Oregon, Eugene, OR, USA

2. Department of Biological Science, Florida State University, Tallahassee, FL, USA

3. Department of Biochemistry, University of Oxford, Oxford, UK

ABSTRACT

In recent decades, genome-wide association studies (GWAS) have been the major approach to understand the biological basis of individual differences in traits and diseases. However, GWAS approaches have limited predictive power to explain individual differences, particularly for complex traits and diseases in which environmental factors play a substantial role in their etiology. Indeed, individual differences persist even in genetically identical individuals, although fully separating genetic and environmental causation is difficult in most organisms. To understand the basis of individual differences in the absence of genetic differences, we measured two quantitative reproductive traits in 180 genetically identical young adult *Caenorhabditis elegans* roundworms in a shared environment and performed single-individual transcriptomics on each worm. We identified hundreds of genes for which expression variation was strongly associated with reproductive traits, some of which depended on individuals' historical environments and some of which was random. Multiple small sets of genes together were highly predictive of reproductive traits, explaining on average over half and over a quarter of variation in the two traits. We manipulated mRNA levels of predictive genes to identify a set of causal genes, demonstrating the utility of this approach for both prediction and understanding underlying biology. Finally, we found that the chromatin environment of predictive genes was enriched for H3K27 trimethylation, suggesting that gene expression variation may be driven in part by chromatin structure. Together, this work shows that individual, non-genetic differences in gene expression are both highly predictive and causal in shaping reproductive traits.

MAIN TEXT

Introduction

Over the past two decades, genome-wide association studies (GWAS) have sought to identify the

genetic basis of individual differences driving traits and diseases. While there have been some notable

successes[1,2], GWAS analyses often require hundreds of thousands of individuals to identify genetic loci

that typically explain a small fraction of variation in traits and diseases[3], and the clinical utility of such loci is

often unclear[4,5]. Complex traits and diseases that are driven by a strong environmental component are

particularly problematic for GWAS, as these environmental inputs cannot be experimentally controlled in

humans[6,7]. This lack of control means that some loci may be spuriously identified and others left

undiscovered. More recent approaches, including transcriptome-wide association studies[8] (TWAS) and

epigenome-wide association studies[9] (EWAS) recognize the intermediate role of epigenetic regulation and

gene expression in driving complex traits and diseases. However, TWAS approaches focus specifically on

predicted genetic effects on expression to prioritize GWAS candidates and thus do not account for

potentially causal environmentally induced changes in gene expression. EWAS approaches, like GWAS

and TWAS, are performed in individuals for which genetic and environmental information are not

experimentally controlled, and thus it is unclear if identified epigenomic changes are driven by a genetic

difference or an environmental effect[10]. Overall, the major approaches used to uncover the etiology of

complex traits and diseases are limited in their ability to separate genetic and environmental effects. Here,

we show that even stochastic differences among individuals can be strongly predictive of complex

reproductive traits when analyzed through the lens of differences in gene expression.

Complete control of genetic background, especially as layered across different environment inputs,

is impossible within human populations. Model organism studies have therefore been invaluable in building

our understanding of environmentally induced effects on epigenetic regulation, gene expression, and traits

of interest in a controlled genetic background[11–17]. However, such analyses generally focus on bulk

populations (typically either to ensure ample material or to average differences across individuals within a

replicate for model organisms such as *Drosophila* and *C. elegans*[18,19]), or they use very few individuals

(typically due to difficulty sampling many larger organisms such as mice[20]), precluding knowledge of how

individual life trajectories may influence gene expression, epigenetic information, and traits of interest. In

recent years, it has become possible to generate individual transcriptomic profiles in *C. elegans* and

Drosophila, and methods to profile individuals in high throughput have improved[21–27]. This has revealed

that genetically identical individuals in a shared environment indeed differ in their gene expression profiles.

However, the effects that gene expression differences across genetically identical individuals may have on

complex fitness-related phenotypes of these same individuals -- which also vary substantially across

genetically identical individuals – remain largely unknown. To fully separate genetic and environmental

causation, we use an isogenic population of *C. elegans* to show that individuals in a common environment

at the same developmental stage exhibit substantial differences in two fitness-related reproductive traits.

By performing mRNA-seq on each individual worm, we identify hundreds of genes for which variation in

expression is strongly associated with reproductive trait variation, determine the extent to which expression

variation is driven by noise versus known environmental history, identify small sets of genes that are highly

predictive of both reproductive traits, and show experimentally that reducing mRNA levels of individual

genes causally affects progeny production. Finally, we show that predictive gene sets for reproductive traits

are enriched for genes with the chromatin modification H3K27me3, revealing that chromatin environment

defines which loci are subject to the individual variation in gene expression that drives variation in complex

traits.

Results

75   To determine if differences in gene expression levels correspond to differences in complex traits,

we measured reproductive traits and whole-animal mRNA transcriptomes for each of 180 isogenic *C.*

elegans in a shared environment at the same stage of early adulthood (Fig 1A). Two reproductive traits –

egg-laying onset (ELO) and the number of progeny produced in the first 24 hours of egg-laying (early

brood) – were measured for each individual. To generate additional phenotypic variation and allow us to

distinguish between gene expression noise and known environmental sources of variation, worms

experienced controlled environmental perturbations early in development in a two-factor design: they were

either born to a day 1 or day 3 adult parent (corresponding to the beginning and middle of egg-laying in *C.*

elegans, respectively), and, once hatched, they were subject to either a constant 20°C temperature or an

eight-hour shift to 25°C (both corresponding to typical rearing temperatures for *C. elegans*). This design is

ideal for at least two reasons. First, in standard worm rearing, worms that at the same developmental stage

in a shared environment have experienced some differences in their environmental histories that are

typically unknown to the observer because they appear physiologically identical. Some of these differences

in environmental history include factors that we directly controlled in this experiment, including the age of

each worm's parent and small fluctuations in rearing temperature. By explicitly incorporating this *historical*

environmental heterogeneity, we can distinguish between gene expression variability induced by

environmental history and gene expression variability driven by 'noise'. Noise encompasses both unknown

extrinsic factors (e.g., microenvironmental differences in food availability) and intrinsic factors (e.g.,

stochasticity in biochemical reactions). Second, with this approach we can ask whether sequencing adult

worms in a shared environment is a viable approach for phenotyping-by-transcriptomics and whether such

an approach depends on knowledge of the historical environment. In other words, if an adult worm is

picked off a plate and sequenced, can we determine when that worm began reproducing and how many

progeny it produced, without regard to its previous environment or its ancestors' environment? In our

experiment, isogenic adult worms in a common environment (with distinct historical environments)

exhibited a range of both ELO and early brood trait values (Fig. S1A), and variation in these traits was

partially driven by environmental history, consistent with prior work on parental age and rearing

temperature in *C. elegans*[27–29](Fig S1B). To determine if gene expression differences identified in single-

worm mRNA-seq were directly associated with reproductive traits measured in the same worms, we used a

linear mixed modeling approach and, remarkably, identified significant genes corresponding to each trait

(Fig 1B-C, Fig S2). Expression of 448 genes was strongly associated with early brood (Fig 1B-C,

Bonferroni p-value of 0.05, nominal p-value of 5.7×10^{-6}), while expression of 11 genes was strongly

associated with ELO (Fig S2). This model was agnostic to the environmental history of the worm, but

accounted for biological replicate, and we confirmed that these effects were not driven by mutations arising

in the population over the course of the experiment (see Methods). This reveals many genes, particularly

for early brood, for which expression differences across genetically identical individuals strongly associate

with the reproductive traits of these individuals. Importantly, this indicates that expression levels of

hundreds of genes provide useful information about the reproductive status of individual worms.

Figure 1. Differences in early brood across isogenic individuals are associated with expression levels of hundreds of genes.

A. Experimental design for single-individual mRNA-seq. P0 worms are allowed to lay progeny as Day 1 or Day 3 adults. F1 progeny are either subject to a constant temperature or an eight-hour shift to 25°C. Beginning in mid-larval stages all F1 worms experience the same environment. For each F1 worm, time to egg-laying onset is scored. 24 hours after egg-laying onset, each F1 worm is collected for single-individual mRNA-seq. F2 progeny laid on plates in the first 24 hours of egg-laying are counted to determine the early brood of each F1 worm.

B. Each of 8824 expressed genes was analyzed in a mixed model to assess the strength of its association with early brood. Significant genes are in pink (positively associated) or purple (negatively associated), non-significant genes are in black. Significance determined by a Bonferroni-corrected p-value cutoff of 0.05 (nominal p-value of 5.67×10^{-6}). Phenotypic and expression data for each worm is shown for the top two genes, *col-20* and *phb-1* with linear model fits in black and 95% confidence intervals in gray.

C. Heatmap of 448 brood-associated genes for all 180 individuals. Worms are sorted from left to right from lowest to highest early brood. Early brood data and environmental information is shown in the bars at the top. Genes are sorted into two clusters based on whether the association was positive or negative.

We next quantified the effect sizes on reproductive traits explained by the individual genes that

were significant in the previous analysis. Further, because of our experimental design, we also determined

the effect size and significance of the association after accounting for environmental history in our model,

allowing us to distinguish how much of the effect of gene expression on reproductive traits is gene

expression 'noise' and not driven by known environmental history. At the phenotypic level, differences in

early brood were partially driven by parental age but not affected by early-life temperature (Fig 1C and Fig

S1B). Differences in the timing of egg-laying onset were partially driven by both parental age and

Figure 2: Associations between gene expression and reproductive traits are partially driven by both noise and historical environment.

A. Two example path analyses showing the relationship between gene expression and early brood. In Model 1, gene expression of a particular gene, *puf-5*, drives differences in early brood and β_1 is the coefficient of the linear mixed model. β_1 represents the total effect of gene expression on early brood. In Model 2, β_2 represents the effect of *puf-5* on early brood that is independent of the historical environment.

B. Two example path analyses showing the relationship between gene expression and egg-laying onset, analogous to A. In Model 1, β_1 is the total effect of *grsp-4* on egg-laying onset, while in Model 2, β_2 represents the effect of *grsp-4* that is independent of historical environment.

C. The 448 genes with for which expression variation is significantly associated with early brood, ordered by the magnitude of their β_1 effect sizes. The absolute values of β_1 and β_2 are plotted upward and downward, respectively, for each gene.

D. The 11 genes for which expression variation is significantly associated with egg-laying onset across individuals, ordered by the magnitude of their β_1 effect sizes. The absolute values of β_1 and β_2 are plotted upward and downward, respectively, for each gene.

C-D. Legend indicates the $-\log_{10}$(p-value) of β_1 and β_2 for each gene.

E. GO Terms for the 448 genes associated with early brood with a q-value < 0.01 are shown. GO Terms for the two subsets of the 448 genes with overlapping GO Terms are also shown: 114 genes that have significant values of β_2 but parental age does not have a significant effect on expression ('noise' genes), and 97 genes that do not have significant values of β_2 but parental age does have a significant effect on expression (parental age genes).

temperature. Because temperature and parental age have effects on progeny phenotypes, we wanted to test the extent to which the association between gene expression and reproductive traits was driven by environmental history (parental age or early-life temperature) for the identified genes. Previous work on parental age found that maternal age affects vitellogenin gene expression to impact progeny phenotypes[27], but how other genes may be affected is not yet known. To determine effect sizes, we calculated the standardized beta coefficient for each gene using a linear mixed model without environmental history, in line with the analysis done in Figure 1 (β_1 in Fig. 2A-B). We then generated a linear mixed model that incorporates environmental history and calculated the standardized beta for the expression component of

the model, effectively isolating gene expression 'noise' that is not explained by environmental history (β_2 in

Fig. 2A-B). These two approaches to calculating standardized beta values for expression are illustrated as

path analyses in Fig. 2A-B. Model 1 considers expression alone and its association with brood, while Model

2 is a causal model that incorporates environmental history and its effects on gene expression. Equations

for these models, including the coefficients for $\beta_3 - \beta_6$, can be found in the methods. Values for all

coefficients can be found in Supplementary File 1. Here we primarily focus on standardized beta values β_1

and β_2, because these respectively represent 1) the total effect of expression on each trait and 2) the effect

of expression on each trait after controlling for historical environment. In Figure 2A, *puf-5* is shown as an

example gene that is part of the group of 448 genes significantly associated with brood. The β_1 value of

0.39 for *puf-5* is significant and indicates that a 1 standard deviation change in gene expression is

associated with a 0.39 standard deviation change in early brood. The β_2 value of 0.26 for *puf-5* is also

significant, meaning that after accounting for historical environment, a 1 standard deviation change in gene

expression is associated with a 0.26 standard deviation change in early brood. The standard deviation in

early brood is 20.3, so this indicates that these gene expression changes associated with *puf-5* account for

approximately 5-8 progeny produced in the first day of egg-laying, which is a sizable effect on fitness.

Similarly, in Figure 2B, *grsp-4* is shown as an example gene with a significant association with egg-laying

onset both before and after accounting for environmental history. The absolute values of β_1 and β_2 for all

significant genes are shown in Figure 2C-D. Early brood genes had a median absolute value of β_1 of 0.38

and a median absolute value of β_2 of 0.23, indicating that for a 1 standard deviation change in gene

expression for a typical significant gene, there was a corresponding 0.38 or 0.23 standard deviation

change, respectively, in early brood. Egg-laying onset genes had a median β_1 value of 0.35 and β_2 value of

0.31. Thus, while we are able to identify many more genes for early brood compared to egg-laying onset,

genes for both traits have largely comparable effect sizes. Among significant genes for both traits, β_2 values

were consistently lower than β_1 (Fig. 2C-D), suggesting some of the total effect size was driven by

environmental history rather than pure noise. Nonetheless, β_2 values were often still highly significant with

substantial explanatory power, indicating that gene expression noise does indeed explain differences in

reproductive traits. For egg-laying onset, 5 of 11 genes have highly significant values of β_2 (Bonferroni

corrected p-value of less than 0.05), though all are nominally significant. Of the 448 genes associated with

early brood, 157 genes have highly significant values of β_2, indicating that gene expression noise

independent of historical environment is driving associations for these genes.

In our previous analysis, we identified genes for which variation is associated with early brood or

egg-laying onset and determined which of these genes had expression variation that was not explained by

the historical environment (i.e., the 157 noise genes). Because of the unique structure of our dataset, we

can also directly ask whether the historical environment has effects on gene expression, regardless of

whether these expression changes lead to changes in our reproductive traits. To address this, we used a

negative binomial mixed model to identify genes that are differentially expressed due to parental age or

early-life temperature. We identified 186 genes with expression changes significantly affected by parental

age, and just 2 genes significantly affected by early-life temperature. 140 of the 186 genes differentially

expressed due to parental age in the previous generation overlapped with the original 448 genes for which

expression variation is associated with early brood, suggesting that the historical environment, i.e. parental

age, at least partially accounts for the gene expression changes that are associated with early brood. 43 of

the 186 genes also overlap with our 'noise' genes, meaning they exhibit gene expression variability

significantly explained by *both* historical environment and by noise; that is, these two sources of variation

operate together for these genes. Thus, among genes for which expression variation is associated with

early brood, we can gain insight on whether this variability in primarily driven by expression 'noise',

historical environmental perturbations, or both.

We next assessed Gene Ontology (GO) enrichments for genes associated with reproductive trait

variation. We focused on early brood genes because hundreds of genes were identified, giving us power to

detect enriched terms. We evaluated GO terms for all 448 genes as well as two subsets of the 448 genes:

1) genes that have a significant value of β_2 and a non-significant effect of parental age on gene expression

(114 genes, the 'noise' group) and 2) genes that have a non-significant value of β_2 and a significant value

of parental age on gene expression (97 genes, the 'parental age' group). Enrichments for all three gene

groups are shown in Figure 2E. We find that the majority of enriched terms for the 448 genes are also

enriched in either the noise group or parental age group, in a mutually exclusive manner (i.e. no term is

enriched in both subsets), suggesting that these two sources of gene expression variation may, to some

extent, be regulated by distinct processes. Indeed, the enriched parental age terms all impinge on

Figure 3. Multi-gene models are highly predictive of early brood and egg-laying onset at an individual level.

A. Total R^2 of gene expression principal components (PCs) in a multiple regression. PCs explain early brood (red), ELO (blue), and randomized data (dotted lines and gray shading).

B. Total R^2 of top genes in a multiple regression. Gene sets explain early brood (red), ELO (blue), and randomized data (dotted lines and gray shading).

A-B. Dotted lines represent average of randomized data iterations and gray shading indicates standard deviation.

C. Total R^2 of top 10 genes in a multiple regression identified in a train set and total R^2 of the same 10 genes used in a test set. Train and test sets randomly selected 500 times and top 10 genes identified in each iteration. Box plot center lines indicate median and box limits indicate upper and lower quartiles. An example of one iteration of train and test sets and corresponding model fits are shown for early brood.

D-E. Machine learning model elastic net regression and leave-one-out cross validation used to identify a predicted trait for each worm given transcriptomic profile and compared to true trait data. Linear fit used to determine R^2. Lines indicate linear fit, and gray indicates 95% confidence intervals.

D. Early brood model, E. ELO model

F. Proportion of 500 iterations from C in which a given gene is selected as one of the top 10 predictive genes for early brood, ordered by those selected from most to least often, and an inset showing the genes that are most frequently selected as predictive.

mitochondrial regulation, consistent with parental age affecting mitochondria-related gene expression in

progeny. Notable categories among enriched noise terms include cytoskeleton organization and female

gamete generation, with the latter category particularly relevant for affecting progeny production.

The analysis shown in Figure 2 provides a framework to understand individual gene expression as

useful both for prediction and understanding causal mechanisms. Toward the goal of using gene

expression profiles to predict phenotypes, we next asked if we could use *aggregate* gene expression

profiles for combinations of genes to predict the reproductive traits of individuals. For example, the β_1 value

of a single gene provides an indication of to what extent the phenotype is associated by that single gene. If

instead of generating individual regressions for each gene, we combined genes into a single model, would

we be able to better predict phenotypes? Because we are focused on prediction rather than the distinction

between expression noise and the historical environment in this analysis, we focus on an analysis that is

agnostic to environmental history, but we emphasize that all worms are genetically identical and collected

while in a shared environment at the same developmental stage. Principal component analysis (PCA)

provides one such aggregate profile which simplifies gene expression data to linearly independent factors

(PCs) that together explain the variation in the gene expression data and are ordered by how much

variation they explain (*i.e.*, PC1 explains the most variation, PC2 explains the second-most, etc.). We

therefore first asked whether ordered PCs together in a multiple regression explain more variation than

expected by chance. Indeed, gene expression PCs explained both ELO and early brood data substantially

better than randomized ELO and early brood data, indicating that individual gene expression profiles

provide a clear signal of phenotypic information (Fig. 3A). Because we have phenotypic and transcriptomic

data for 180 worms, and therefore up to 180 linearly independent PCs that can be added a multiple

regression model, the total R^2 eventually rises to 1 even for randomized data, as shown in Fig. 3A.

Therefore, the optimal number of PCs occurs at the inflection points of the graph, which is after only 7 PCs

for early brood (R^2 of 0.55) but 28 PCs for ELO (R^2 of 0.56). Thus, for early brood in particular, very few

PCs explain over half of the variation in the data and explain substantially more variation than for shuffled

phenotypic data. Therefore, whole transcriptomes are a viable approach to trait prediction in isogenic *C.*

elegans.

Because PCA incorporates data from all expressed genes, we next wanted to determine if a smaller

subset of genes could be similarly predictive of reproductive traits. We again used a multiple regression

approach in which we successively selected each gene by identifying the gene that explained the most

additional variance compared to the previous model (Fig. 3B). Genes that exhibit highly correlated

expression patterns, even if they are each individually associated with traits of interest, would not explain

much additional variance in a multiple regression approach. Therefore, this approach prioritizes identifying

genes that are most predictive in conjunction with other genes. We found relatively small sets of genes

predict ELO and early brood phenotypes very well, and do so significantly better than randomized trait

data, with diminishing returns for variance explained as the number of genes increases (Fig. 3B). Maximum

differences in the proportion of variance explained between trait data and randomized data occur in a

model with only 4 genes for the early brood analysis (R^2 difference of 0.433) and with 7 genes in the egg-

laying onset analysis (R^2 difference of 0.195). While less powerful than utilizing the entire transcriptome,

this result highlights that individual worm gene expression profiles consisting of only a handful of key

selected genes are highly informative for predicting the individual's traits.

We next wanted to develop a quantitative estimate of how much trait variation is explained by

aggregate gene expression profiles using two distinct approaches that prevent model over-fitting and would

therefore be more feasible to use for trait prediction in a new dataset. First, as an extension of the multiple

regression analysis identifying predictive genes, we randomly split the full dataset (consisting of trait data

and gene expression profiles) into a training set and test set. In the training set, we used the multiple

regression approach to identify the set of 10 genes that explained the most variance in the trait data. We

then asked how well this set of 10 genes explained variance in the test set and repeated this process 500

times, each time randomly splitting the data into training and test sets (Fig. 3C). We found that median

proportion of variance explained in the test sets was 0.55 (range 0.36 to 0.73) for early brood and 0.26

(range 0.07 to 0.49) for ELO, and an example of the fit of one of these iterations is also shown (Fig. 3C).

Thus, sets of only 10 genes explain over half of trait variation in early brood and over a quarter of variation

in ELO using test sets of data that were not used to select predictive genes. All 500 sets of 10 predictive

genes are available as part of Supplementary File 1. As a second approach, we used elastic net regression

and leave-one-out cross validation on the full set of expressed genes. In other words, we used a machine

learning approach designed to prevent overfitting to develop a model for all individuals except one, used

the model to predict the trait for the remaining individual, then repeated for all individuals. Consistent with

the multiple regression approach, the proportion of variance explained was 0.53 for early brood and 0.47

for ELO (Fig. 3D-E). Aggregate gene expression profiles identified via multiple approaches are thus highly

predictive of reproductive traits among isogenic individuals in a shared environment, reliably explaining

about half of reproductive trait variation among genetically identical individuals.

A key remaining question is whether predictive genes are causal to affect phenotypes and whether

similar underlying regulation underlies expression variation in predictive genes. To address causality, we

focused on the early brood trait because predictive genes explained a higher proportion of variance for this

trait compared to ELO. We used RNA interference (RNAi) to reduce the level of gene expression of

predictive genes and prioritized genes that were also in the 448 brood-associated genes, either highly

independent of parental age or dependent on parental age, and have an ortholog in humans. Within this

curated set, genes causally affected early brood in 5 of 7 cases compared to empty vector (Fig. 4A). We

followed up on one pair of genes from this screen, *puf-5* and *puf-7,* to ask whether perturbing the dose of

RNAi shows a dose response effect on brood. These genes have previously been shown to act

redundantly in fertility in *C. elegans*[30,31] and when knocked down simultaneously in our assay have a large

effect on brood without causing sterility (Fig. 4A). Consistent with the idea that differences in the level of

mRNA affect the early brood trait, the dose of RNAi corresponds to the effect size on brood (Fig. 4B). Thus,

this analysis reveals that the predictive genes also make excellent candidates as causal effectors of

complex traits and can affect these traits dependent on the amount of mRNA available.

Expression of predictive genes can be causally related to reproductive traits (Fig. 4A-B), but do

these genes share underlying regulation? The majority of *C. elegans* genes fall into one of two chromatin

domains with distinct levels of gene activity[32–34]. Regulated domains include genes that are expressed at

specific stages of development and are marked by the facultative heterochromatin modification H3K27me3.

In contrast, active domains include genes that are expressed broadly across cell types, including in the

germline, and marked by the euchromatic modification H3K36me3[32,34,35]. These two marks are generally

mutually exclusive, and marks are robust across several stages of development in *C. elegans*[34] despite

major changes in gene expression that occur during this time. Importantly, the robustness of these

chromatin marks throughout development means that we can use existing data to determine whether

genes in active or regulated domains are differentially associated with gene expression variation linked to

reproductive traits in our study. We therefore used genes previously classified as regulated or active to

gain insights into the epigenetic status of genes associated with and predictive of reproductive traits.

Because active genes are enriched for being expressed in the germline, we performed a two-factor

analysis that takes into account both tissue (soma or germline) and chromatin domain (regulated and

active). Among early brood-associated genes, germline genes are strongly positively associated with

brood, regardless of chromatin environment, but chromatin environment and tissue also exhibit a significant

Figure 4. Genes predictive of reproductive traits causally affect early brood and are enriched for H3K27me3

A. RNAi knockdown of selected predictive genes compared to empty vector (ev) and effects on early brood.

B. Dose response of *puf-5* and *puf-7* RNAi together with empty vector and effects on early brood.

A-B. Linear mixed model with RNAi as fixed effect and biological replicate as a random effect. Center lines, median; box limits, upper and lower quartiles. *p<0.05, ***p<0.001

C. Effect sizes on brood for genes categorized by tissue (soma or germline) and chromatin domain (active or regulated). Two-way ANOVA showed significant interaction between tissue and chromatin domain (p = 1.58 x 10^{-6}) and a significant main effect of tissue (p < 2 x 10^{-16}). Post-hoc Tukey tests corrected for multiple testing showed a significant difference between chromatin domains within somatic tissues (adjusted p = 0) but not germline tissues (adjusted p = 0.98). Within somatic genes, active somatic genes effects on average do not differ from 0 (p = 0.3, one-sample t-test with μ = 0), while regulated somatic genes on average have a significant negative association with brood (p < 2.2 x 10^{-16}, one-sample t-test with μ = 0). ***p<0.001

D. Histogram of all iterations identifying predictive gene sets and the proportion of each set in a regulated chromatin domain marked with H3K27me3 compared to randomly selected sets of genes from the same background set. Early brood iterations are shown in pink, ELO in blue, and the same gray control is shown for both histograms. The vertical lines are color-coded to indicate the median proportion of genes in regulated domains in the corresponding condition. Kolmogorov-Smirnov test used to determine statistical significance. ***p<0.001

interaction (Fig. 4C). Post-hoc pairwise comparisons reveal that this interaction is driven by differences in the chromatin environment in the soma. In particular, while active somatic genes have on average no effect on brood, regulated somatic genes are strongly negatively associated. Because regulated genes are typically expressed in a temporally controlled or tissue-specific manner in the soma, this may indicate that aberrant expression of these genes has negative consequences for reproduction, or similarly, that appropriate regulation and silencing of these genes promotes reproduction. To determine if epigenetic regulation may underlie variation in predictive genes for both traits overall, without regard to the direction of association, we used all gene sets identified from iterations of predictive sets (Fig. 3C,F) to determine whether these genes are more likely to be found in regulated domains. For both reproductive traits, predictive genes were strongly over-enriched for being in a regulated domain compared to randomly selected gene sets from the same background set (Fig. 4D). Genes in regulated domains are more likely to be variable across developmental stages and cell types[34], and we find in our data that genes with the most expression variation across individuals unexplained by experimental

factors are also more likely to be found in regulated domains (Fig. S3). Taken together, these results

suggest that regulated domains are subject to gene expression variation in genetically identical individuals

and that this variation is an important substrate underlying complex trait variation.

Discussion

Individuals differ from each other for a variety of traits over the course of their lives, but being able

to reliably predict these differences and understand their causal basis is a challenge. Gene expression is a

critical intermediate functional layer between genotype and phenotype, but the role that gene expression

plays in shaping individual organismal traits independently of genetic variation and environmental factors

has remained elusive. Here we effectively screened for genes that exhibit expression variation between

isogenic *C. elegans* individuals and for which this variation corresponds to two critical reproductive traits –

time to egg-laying onset and early brood. We identified hundreds of genes associated with these traits,

differentiated between expression variation that is dependent and independent of historical environmental

factors, and defined sets of highly predictive and causal genes. Predictive genes for both traits are

enriched for a particular chromatin modification – H3K27me3 – highlighting the possibility that chromatin

structure may underlie the potential for gene expression to vary between individuals. These findings build

upon past research and suggest key areas for future work.

With our framework, we identify individual differences in gene expression that are associated with

reproductive traits, and we parse whether these effects are primarily driven by noise or by known historical

environmental factors. For example, *puf-5* and *puf-7* expression is highly associated with early brood after

accounting for the known historical environment, and knockdown of these genes causally affects early

brood in a dose-dependent manner. On the other hand, genes like C34C12.8 exhibit an overall association

with early brood that is primarily driven by the effect of parental age on gene expression; nonetheless,

C34C12.8 is also causally linked to early brood. Thus, by experimentally manipulating and measuring

individuals, our approach is highly efficient in dissecting multiple sources of environmental variation --

including noise and known environmental perturbations – underlying multiple traits that would otherwise be

obscured by bulk experiments. In both cases, the mechanism(s) generating gene expression differences

across individuals remains unknown. For 'noise' genes in particular, whether differences in gene

expression are driven by extrinsic or intrinsic factors is unclear. While every effort was made to control the

environment for our experiment, isogenic individuals nonetheless exhibit some extrinsic differences in their

lives, such as how they choose to wander on a plate of food, that may to some extent impact gene

expression and reproduction. We further expect that some of the variation observed is intrinsic, based on

literature exploring the stochastic nature of transcription[36], as well as examples that suggest the amount of

phenotypic variation present among individuals of a given genetic background is itself genetically

controlled[37]. Continued experimentation to determine the extent to which intrinsic or extrinsic factors

regulate gene expression variability in this context -- where such variability is directly related to complex

phenotypes -- is of high interest for future work.

When genes expression differences are identified across individuals by single-worm mRNA-seq,

such differences could occur because individuals have different numbers of cells of a given type (e.g.,

germ cells) but each cell express genes at consistent levels, or alternatively, that individuals differ because

within each cell, additional copies of a gene are being transcribed while cell numbers themselves remain

constant. Of course, a combination of these scenarios is possible as well. *C. elegans* uniquely possess an

invariant somatic cell lineage[38], but can exhibit differences in cell number in the adult germline, including in

the primordial zone[39] or in the uterus [40]. Our results show that, in general, genes expressed in the germline

tend to be positively associated with early brood, while somatic genes tend to be negatively associated with

brood. Of somatic genes, genes marked with H3K27me3 are particularly predictive of early brood. Somatic

genes marked with H3K27me3 are also more likely to exhibit gene expression variability in general. Given

that the somatic lineage is invariant and somatic genes marked with H3K27me3 are the most variable in

expression and most predictive of early brood, it is unlikely that cell number differences are the major driver

of gene expression differences. While single-cell RNAseq has been performed in *C. elegans*, such analysis

requires thousands of individuals to be pooled to obtain single cells and is not technically possible at the

level of single individuals. Thus, to fully understand cell-type specific variation in single *C. elegans*, and

how these relate to individuals' traits, further innovation on this front is required.

We observed that differences in chromatin domains are associated with differences in gene

expression and traits across individuals. Regulated genes are defined by the histone modification

H3K27me3, a mark that is robust across developmental stages despite differences in gene expression,

while active genes are marked with H3K36me3 and are typically broadly expressed, including in the

germline. In our data, regulated genes are enriched both among genes with the most unexplained variation

and among genes for which variation is predictive of reproductive traits. In contrast, germline genes that

predominantly consist of genes in active domains are specifically positively associated with brood. Taken

together, one possible model is that regulated genes marked with H3K27me3 are more subject to inter-

individual variation than genes marked with H3K36me3, which may lead to differences in gene expression

without changing the levels of these marks. This is consistent with observations that regulated genes

exhibit a higher coefficient of variation across conditions and that increased expression of genes marked

with H3K27me3 is not associated with a change in histone post-translational modifications (*e.g.,* to an

active mark like H3K36me3)[41]. An alternative hypothesis is that the levels of the marks themselves vary

and that this leads to changes in gene expression across individuals. While it is clear that regulated genes

are more subject to inter-individual variability in expression, at this stage it is unknown whether this

variability is driven by differences in the level of chromatin marks. Performing analysis on histone

modifications genome-wide in individual worms, which is still technically challenging, will be a key next step

to determine whether differences in chromatin structure correlate with gene expression changes at an

individual level.

Individual gene expression differences underlying reproductive traits suggest that gene expression

differences may be important for a variety of other complex traits. Gene expression differences among

isogenic individuals in a shared environment have been previously documented as predictive of lifespan[42–]

[44]. In these studies, reporter gene expression for specific candidate genes predicted individuals' later

lifespan. Our results extend on such studies by identifying predictive genes in the context of reproductive

traits in a high-throughput manner, rather than relying on specific reporters. Our approach can in theory be

used directly in the context of other traits that occur throughout the life cycle, such as those related to

development, but must be used indirectly for time-to-death traits like lifespan or stress resistance. Because

a worm, once collected for RNA, cannot be subsequently phenotyped for traits like lifespan, using

predictive reporters for time-to-death traits in conjunction with transcriptomics will likely be a useful

approach for identifying novel regulators for a broad swath of traits.

Genetically identical individuals can have large differences in phenotype if gene expression varies

across those individuals. Our work illustrates that differences in gene expression can be causally linked to,

and are highly predictive of, important reproductive phenotypes among genetically identical individuals.

Assaying isogenic individuals for both organismal and molecular phenotypes enables us to uniquely

understand the basis of non-genetic variation underlying individual differences and provides an avenue for

high-throughput phenotyping by transcriptomics[17]. Given the major role of non-genetic factors in regulating

complex traits and diseases in humans, identification of these molecular drivers of variation may be a

powerful next step for advancing precision medicine.

Figure S1. Isogenic worms at the same developmental stage exhibit variability in egg-laying onset and

early brood that partially depends on previous environmental conditions

A. Univariate histograms of early brood and egg-laying onset trait values for all worms

B. Bivariate scatterplots of early brood and egg-laying onset trait for all worms. The same data is plotted

twice but color-coded according to two previous environmental perturbations (age of parent and early-life

temperature). Center lines for box plots indicate median; box limits indicate upper and lower quartiles.

Significance determined using a linear mixed model with environmental perturbation as a fixed effect and

biological replicate as a random effect, ***p<0.001, *p<0.05

Figure S2. Variation in 11 genes is significantly associated with egg-laying onset.

Each of 8824 expressed genes was analyzed in a mixed model to assess the strength of its association

with egg-laying onset. Significant genes are in red, non-significant genes are in black. Significance

determined by a Bonferroni-corrected p-value cutoff of 0.05 (nominal p-value of 5.67×10^{-6}). Phenotypic

and expression data for each worm is shown for two exemplar genes, *anmt-3* and *rnp-8*. Black lines from

linear model fit and gray shading indicates 95% confidence intervals.

Figure S3. Single-individual mRNA-seq reveals most and least variable genes after controlling for

environmental perturbations and biological replicate.

A. Log$_{10}$-transformed unexplained variance in mRNA-seq after controlling for environmental factors and

biological replicate is plotted against log$_{10}$-transformed average CPM across all worms and a loess fit line is

shown in black with a 95% confidence interval in gray. Each point is a different gene and genes are color-

coded according to their distance from the fit line. Genes in purple are most variable given their level of

expression, while genes in pink are the least variable. Exemplar genes with comparable expression levels

are shown on the right. Center lines for box plots indicate median; box limits indicate upper and lower

quartiles.

B. Variance Z-scores for genes with regulated domains (H3K27me3) compared to variance Z-scores for

genes with active domains (H3K36me3). Wilcoxon test, ***p < 2 x 10^{-16}

METHODS

Strains used

Strain N2-PD1073 was used for all experiments. N2-PD1073 is a subclone of the N2-derivative strain VC2010 that was generated in the process of assembling a new N2 reference. Worms were passaged on *E. coli* OP50 except during RNAi experiments, when worms were fed HT115 bacteria expressing dsRNA for genes of interest (see 'RNAi' section).

Experimental set-up for single-worm RNA-seq

Worms were well-fed at 20°C for several generations prior to beginning experiments. The experiment takes place over the course of 10 days. On day 1, for each of 5 biological replicates, P0 embryos and L4s were picked to separate 5 cm NGM plates with OP50 and kept at 20°C. Approximately 10-20 individuals were placed on each plate and several plates were picked for each stage, ensuring that worms do not experience dietary restriction. On day 4, P0 worms were either in the first day of adulthood ('young adult') or third day of adulthood ('older adult'). P0 young adults and older adults were singled to new plates, allowed to lay progeny for two hours at 20°C, then removed from the plate. Hour 0 of development for F1 worms is considered the end of this egg-laying window. F1 worms were then incubated at 20°C until day 5 of the experiment (undergoing approximately 18 hours of development as L1/L2 larvae). On day 5, approximately half of these larvae remained at 20°C while the other half were shifted to 25°C for 8 hours then shifted back to 20°C. On day 6 all worms remained at 20°C and were singled to individual NGM plates with OP50 to ensure they can be individually phenotyped for reproductive traits. On day 7, from 66-74 hours of development, egg-laying onset was assessed every hour by checking for embryos on each plate. Worms that died, bagged, or otherwise did not reach egg-laying onset in this time interval were not included in subsequent analysis, and all together this affected 6% of worms across all biological replicates. Time to egg-laying onset was noted for each worm. 24 hours after egg-laying onset, on day 8, each worm was collected for cell lysate generation for mRNA-seq (see 'Cell lysate generation' section of Materials and Methods). On day 10, the number of progeny laid on each plate during the 24 hours between egg-laying onset and collection were counted as late larvae/young adults to ensure all progeny were counted. All together, each F1 worm collected experienced two environmental perturbations (young adult

or older adult parent, and early-life constant temperature or temperature shift) and was measured for two

traits (egg-laying onset time and number of progeny produced within 24 hours) prior to mRNA-seq.

Cell lysate generation

Individual worms were collected for chemical lysis 24 hours after egg-laying onset (that is, worms

that reached egg-laying onset at 72 hours were collected at 96 hours, worms reaching egg-laying onset at

73 hours were collected at 97 hours, etc. for each hour). 200 μL of lysis mix, consisting of 187.2 μL Elution

buffer (10 mM Tris, 0.1 mM EDTA) and 12.8 μL Proteinase K, was made fresh every 2-3 hours on ice. 6 μL

of lysis mix was pipetted into each PCR tube according to how many worms needed to be collected in a

given hour. Each worm was picked into a 10 μL drop of dH2O for a quick wash to remove any excess

bacteria, then transferred to a second drop of dH2O. After this, each worm was picked into an individual

PCR tube with lysis mix. Immediately after worms are placed into tubes with lysis mix, we proceeded with

the lysis protocol (typically no more than a dozen worms were prepared at once). The lysis protocol is 65°C

for 10 min, 85°C for 1 min, and 4°C hold and is based on a previously published protocol[22]. After the lysis

protocol, worms were moved to ice and then stored at -80°C until mRNA-seq library preparation.

Preparation and sequencing of mRNA-seq libraries

The KAPA mRNA-seq preparation kit (KK8580, Roche Sequencing Solutions) was used to prepare

libraries. 5 μL of each cell lysate preparation described above, which contains total RNA, was used for

each sample. Reagents were used at 0.25X and kit instructions were followed to generate libraries. Library

concentrations were quantified using Qubit, and samples were pooled together in equimolar ratios in two

batches of 96 for sequencing. Custom dual-indexed adapters were designed by the University of Oregon

Genomics and Cell Characterization Core Facility and sequencing data was generated with the Illumina

NovaSeq 6000 at the same facility.

Analysis of single-worm mRNA-seq data

FASTQ files were aligned to WBcel235 version of the *C. elegans* genome using subread-align[45]

with flag -t 0. Bam files were counted using featureCounts with flag -T 5. 12 of 192 samples were filtered

out at this stage due to particularly low read counts (less than 1 million reads) and/or poor mapping quality (less than 70% of reads both mapped to the genome and assigned to a genomic feature). The remaining 180 samples had an average of 5.8 million reads (SD 3.4 million), and an average of 85.8% (SD 3.1%) of reads both mapping to the genome and assigned to a genomic feature. Resulting count data from these 180 samples was CPM-normalized in R using the package edgeR[46] and only protein-coding genes were included. Depending on the downstream analysis, genes were filtered in one of two ways for further analysis: 1) genes expressed at a level of CPM > 10 in at least one library (8824 genes) or 2) genes expressed at a level of CPM > 1 across all libraries (7938 genes). While these two sets are largely overlapping, the first way of filtering allows identification of genes that are not expressed at all in some samples but moderately expressed in others. However, as models become more complex, having some samples with 0 expression causes problems for model fitting and thus having a more stringently-defined set is more appropriate. To identify genes associated with early brood and egg-laying onset, we looped through the set of 8824 genes and tested two linear mixed models using the 'nlme' package[47] in R to test the effect of each gene on each of two reproductive phenotypes. In both cases the fixed effect was the normalized gene expression level in each worm and the random effect was biological replicate. For the first model, the dependent variable was early brood and for the second model the dependent variable was hours to egg-laying onset. To determine the effect of environmental perturbations on gene expression, we looped through each gene and fit a negative binomial mixed model to each gene. In this case, for each gene, fixed effects included environmental perturbations (parental age and early-life temperature), the random effect included biological replicate, and the dependent variable was gene expression level. For both sets of models, the summary() function was used to generate p-values, and a Bonferroni correction was used to determine genes with significant effects. Gene expression variation that was unexplained by the negative binomial (residual variance compared to the model) was used to determine the most and least variable genes in the data after controlling for environmental effects and biological replicate. Because variance increases with higher expression, we identified most and least variable genes by fitting a loess model to $\log_{10}$(unexplained variance) vs. $\log_{10}$(mean CPM) for the 7938 genes. Z-scores were calculated for each gene relative to the loess fit to determine genes that were most and least variable given their level of gene expression.

Path analysis and effect size calculations

Using the R package 'lme4'[48], for each individual i and biological replicate j, the following linear mixed models were fit for each gene k:

$(\text{Early brood})_{ij} = \beta_0 + \beta_1(\text{Expression level of gene } k) + u_j + e_{ij}$

$(\text{Early brood})_{ij} = \beta_0 + \beta_2(\text{Expression level of gene } k) + \beta_3(\text{Parental age}) + \beta_4(\text{Early-life temperature}) + u_j + e_{ij}$

$(\text{Egg-laying onset})_{ij} = \beta_0 + \beta_1(\text{Expression level of gene } k) + u_j + e_{ij}$

$(\text{Egg-laying onset})_{ij} = \beta_0 + \beta_2(\text{Expression level of gene } k) + \beta_3(\text{Parental age}) + \beta_4(\text{Early-life temperature}) + u_j + e_{ij}$

$(\text{Expression level of gene } k) = \beta_0 + \beta_5(\text{Parental age}) + \beta_6(\text{Early-life temperature}) + u_j + e_{ij}$

Parental age was binarized such that progeny of day 1 adults were given the value of 1 and progeny of day 3 adults were given the value of 0. Early-life temperature was binarized such that individuals experiencing a constant 20°C temperature throughout their lives were given the value of 1 and individuals that experienced a 25°C temperature shift were given the value of 0. All values of β_n were standardized such that a 1 standard deviation change in the independent variable represents a shift in the dependent variable corresponding to the value of β_n. u_j represents the random intercept for each biological replicate, and e_{ij} is the error term. Coefficients β_1 through β_6 are shown in a path analysis format in Figure 2A-B, and coefficients β_1 and β_2 are shown graphically in Figures 2C-D for all significant genes from Figure 1 and Figure S2. The values of all coefficients for all significant genes are in Supplementary File 1.

Gene Ontology (GO) analysis

GO Term analysis was performed using the R package 'clusterProfiler'[49]. The enrichGO function was used for each gene list with the following parameters: OrgDb = org.Ce.eg.db, keyType = "SYMBOL", ont = "BP", pAdjustMethod = "BH", qvalueCutoff = 0.05. Output for all three gene lists is included in Supplementary File 1. Figure 2E shows all enriched terms with a q-value < 0.01 for the list of 448 genes and any enriched terms that overlap with this set from that are enriched from the subsets of 97 parental age genes or 114 noise genes. The set of 97 genes was generated by identifying genes in the set of 448 that were significantly affected by parental age and did not have a significant value of β_2. The set of 114 genes was generated by identifying genes in the set of 448 that were not significantly affected by parental age and had a significant value of β_2.

Prediction analysis

Principal component analysis was performed on 7938 CPM-normalized genes (without regard to

reproductive phenotype data). CPM values for each gene were mean-normalized and log2-transformed

prior to use in the prcomp function in R. To determine whether principal components explained variance in

the reproductive traits, each principal component was successively added to a multiple linear regression

model, beginning with the principal component explaining the most variance in the gene expression data

(*i.e.,* PC1), to determine the total variance explained by cumulative principal components. As a control

comparison, phenotypic data was shuffled repeatedly (100x) and the same multiple regression analysis

was performed. This analysis was performed for both the early brood trait and egg-laying onset trait.

To identify a set of genes that explains variance in each trait, all 7938 genes were looped through in

a linear regression model to identify the single gene explaining the most variance. Then the remaining

7937 genes were looped to identify which gene explained the most additional variance when the first gene

is already included in the model. This was repeated until the total variance explained was close to 1. This

analysis was repeated for shuffled phenotypic data as a control comparison.

To determine how well sets of genes would be likely to work in a new dataset, we used two

approaches. First, we randomly split the data (including matched phenotypic data and gene expression

data) into two equal groups, the train set and test set, 500 times. Each time, for the train set, the protocol

described above was used to identify the top 10 genes that together explain the most variance in the data

using a multiple regression. This set of 10 genes was then used on the test set to determine how much

variation is explained when the genes were not selected because of how well they work in this set. The

typical variance explained in the test set provides an estimate of how well the genes should explain

variance in a new dataset. In a second machine learning approach, we used elastic net regression using

the glmnet package[50] in R with alpha = 0.5 and leave-one-out cross validation. That is, we trained the

model for all data points except for one, then used the model to predict that remaining data point, and

repeated for all data points.

Enrichment analysis

To determine whether brood-associated genes and variable gene sets were enriched for particular

chromatin environments and tissue specificities we used our previously generated annotation for *C.*

elegans transcriptional units[51]. Briefly, in this approach we used published data[32,35] to obtain tissue

specificity and chromatin environment for *C. elegans* regulatory elements genome wide and then used

bedtools2[52] to annotate regulatory elements to the nearest gene using the coordinates of transcriptional

units extracted from the Wormbase gff3 (version 279).

To determine if there was an interaction between tissue and chromatin environment for effects on

early brood (Fig. 4C), we restricted genes to those expressed in the soma or germline, and in either an

active or regulated chromatin domain. We then performed a two-way ANOVA to determine if these factors

affected the effect sizes for each gene on brood (the same effect sizes plotted in Fig. 1B), followed by post-

doc Tukey's tests for pairwise comparisons and one-sample t-tests. To determine if gene expression

variation differed by chromatin domain (Fig. S3B), we performed a Wilcoxon test comparing the variance z-

scores for regulated genes to the variance z-scores for active genes.

To determine whether predictive genes were enriched for regulated or active genes as shown in

Figure 4D, we used the 500 sets of 10 genes for each trait described in the 'Prediction analysis' section of

the methods that were used to predict reproductive phenotypes (Figure 3C). We filtered the chromatin

dataset described above to include only genes that were classified as either active or regulated. We then

merged this with the background of 7938 expressed genes from our gene expression analysis. Of 10

predictive genes, a median of 7 genes were either active or regulated for each of the 500 iterations and

included in subsequent analysis. For each iteration, the proportion of regulated genes was calculated. As a

control, 500 random sets of 10 genes were selected from our background set, merged with the filtered

chromatin data, and the proportion of regulated and active genes was determined. The distribution of the

proportion of predictive regulated genes for each trait was compared to the proportion of regulated genes

from random selected sets of genes using a Kolmogorov-Smirnov statistical test.

RNAi

RNAi was performed by feeding. Colonies of RNAi bacteria (*E. coli* HT115 containing a specific

RNAi construct) were individually grown in 5 mL LB with carbenicillin overnight (~16-18 hours) at 37°C with

shaking. RNAi bacteria was obtained from the Ahringer library[53] for empty vector (ev) and for genes of

interest: ZK1127.7 (*cin-4*), F54C9.8 (*puf-5*), B0273.2 (*puf-6/7*), R03G5.1 (*eef-1A.2*), C34C12.8, F13B10.1

(*tir-1*), T07C4.9 (*nex-2*), and B0416.6 (*gly-13*). Plates used for RNAi were made with NGM, carbenicillin,

and IPTG as described previously[54]. These plates were seeded with RNAi bacteria and allowed to dry and

grow for ~30 hours. L4 larvae were plated in groups of ~10 on these plates, then singled the next day (~16-

18 hours later) onto new RNAi plates prepared at the same time as the original batch of plates. Adult

worms were allowed to lay progeny for 24 hours, then adults were removed from the plates. If a worm died

during this time it was censored. The number of progeny laid by each adult was counted 2 days later,

resulting in an early brood measurement for these experiments. For the *puf-5* and *puf-7* dose response,

puf-5 and *puf-7* RNAi constructs were grown separately as described then pooled together 1:1. This mixed

RNAi was considered the full dose of *puf-5* and *puf-7* RNAi and was pooled with ev in the ratios shown in

Fig. 4C for the varying doses. Statistical analysis on early brood data was performed by comparing RNAi

treatment targeting a particular gene to empty vector for biological replicates within a batch. Linear mixed

effects models were used in the R package 'nlme'[55] with RNAi treatment as a fixed effect and biological

replicate as a random effect. The summary() function was used to generate p-values.

Validation of isogenic worms

We used the mRNA-seq data to validate that the frequency of genetic variants among the

essentially isogenic individual worms was very low. First, we used samtools[56] to generate coverage files

from the sorted bam files resulting from mRNA-seq data processing. We filtered coverage files for each

worm to include only nucleotides that had at least 20 reads mapping to that nucleotide. For each worm and

nucleotide, we determined whether the base call of that nucleotide was unambiguous (all 20+ reads have

the same call), ambiguous with two genotypes (exactly two different bases were called), or ambiguous with

more than two genotypes called. All worms with sufficient depth at a nucleotide were used to determine

putative homozygous or heterozygous variants. A particular nucleotide was included in subsequent

analysis if at least 10 worms had sufficient coverage at the site. This resulted in 8,361,705 unique sites and

a total of 768,738,255 sites across all worms. Candidate homozygous variants included nucleotides that

were called as one base unambiguously in at least one worm and called as a different base unambiguously

in other worms. Candidate heterozygous variants were identified by filtering ambiguous nucleotides with

two genotypes in which the frequency of each genotype was present at a frequency of between 0.3 and 0.7

and for which at least 10 other worms were unambiguously called a single genotype. These criteria reduce

the possibility that a heterozygous mutant is erroneously called if a site was merely subject to rare

sequencing errors. This resulted in a homozygous variant rate of 2.21×10^{-8} and a heterozygous variant

rate of 9.86×10^{-7}. While already quite low, these rates likely represent over-estimates because visual

inspection of bam files in some cases revealed ambiguous base calls. Over three quarters of sites (78.5%)

with a putative heterozygous variant were found only in a single worm, meaning that these rare variants

cannot account for an association with expression across 180 worms.

Statistical analysis

Statistical tests are described in detail in figure legends and in the corresponding section of the

methods. All individual data points are plotted in figures whenever possible.

Data availability statement

All raw mRNA-seq data is available at NCBI GEO accession GSE244875. All data used to generate

figures are available as part of Supplementary File 1.

Code availability statement

Scripts are available at https://github.com/amykwebster/SingleWormRNAseq and have been

preserved on Zenodo at time of publication at https://doi.org/10.5281/zenodo.15489998.

ACKNOWLEDGEMENTS

We thank Chien-Hui Chuang and the Bruce Bowerman lab for sharing RNAi clones. Funding was

provided by grants F32GM146402 to AKW and R35GM131838 to PCP. We thank members of the Phillips

lab for critical feedback. We would also like to thank WormBase.

REFERENCES

1.   Visscher, P.M., Wray, N.R., Zhang, Q., Sklar, P., McCarthy, M.I., Brown, M.A., and Yang, J. (2017). 10 years of GWAS discovery: biology, function, and translation. Am. J. Hum. Genet. *101*, 5–22. 10.1016/j.ajhg.2017.06.005.

2.   Edwards, A.O., Ritter, R., Abel, K.J., Manning, A., Panhuysen, C., and Farrer, L.A. (2005). Complement factor H polymorphism and age-related macular degeneration. Science *308*, 421–424. 10.1126/science.1110189.

3.   Nolte, I.M., van der Most, P.J., Alizadeh, B.Z., de Bakker, P.I., Boezen, H.M., Bruinenberg, M., Franke, L., van der Harst, P., Navis, G., Postma, D.S., et al. (2017). Missing heritability: is the gap closing? An analysis of 32 complex traits in the Lifelines Cohort Study. Eur. J. Hum. Genet. *25*, 877–885. 10.1038/ejhg.2017.50.

4.   Alsheikh, A.J., Wollenhaupt, S., King, E.A., Reeb, J., Ghosh, S., Stolzenburg, L.R., Tamim, S., Lazar, J., Davis, J.W., and Jacob, H.J. (2022). The landscape of GWAS validation; systematic review identifying 309 validated non-coding variants across 130 human diseases. BMC Med. Genomics *15*, 74. 10.1186/s12920-022-01216-w.

5.   Gallagher, M.D., and Chen-Plotkin, A.S. (2018). The Post-GWAS Era: From Association to Function. Am. J. Hum. Genet. *102*, 717–730. 10.1016/j.ajhg.2018.04.002.

6.   Barban, N., Jansen, R., de Vlaming, R., Vaez, A., Mandemakers, J.J., Tropf, F.C., Shen, X., Wilson, J.F., Chasman, D.I., Nolte, I.M., et al. (2016). Genome-wide analysis identifies 12 loci influencing human reproductive behavior. Nat. Genet. *48*, 1462–1472. 10.1038/ng.3698.

7.   Locke, A.E., Kahali, B., Berndt, S.I., Justice, A.E., Pers, T.H., Day, F.R., Powell, C., Vedantam, S., Buchkovich, M.L., Yang, J., et al. (2015). Genetic studies of body mass index yield new insights for obesity biology. Nature *518*, 197–206. 10.1038/nature14177.

8.   Wainberg, M., Sinnott-Armstrong, N., Mancuso, N., Barbeira, A.N., Knowles, D.A., Golan, D., Ermel, R., Ruusalepp, A., Quertermous, T., Hao, K., et al. (2019). Opportunities and challenges for transcriptome-wide association studies. Nat. Genet. *51*, 592–599. 10.1038/s41588-019-0385-z.

9.   Rakyan, V.K., Down, T.A., Balding, D.J., and Beck, S. (2011). Epigenome-wide association studies for common human diseases. Nat. Rev. Genet. *12*, 529–541. 10.1038/nrg3000.

10.   Battram, T., Gaunt, T.R., Relton, C.L., Timpson, N.J., and Hemani, G. (2022). A comparison of the genes and genesets identified by GWAS and EWAS of fifteen complex traits. Nat. Commun. *13*, 7816. 10.1038/s41467-022-35037-3.

11.   Torres-Garcia, S., Yaseen, I., Shukla, M., Audergon, P.N.C.B., White, S.A., Pidoux, A.L., and Allshire, R.C. (2020). Epigenetic gene silencing by heterochromatin primes fungal resistance. Nature *585*, 453–458. 10.1038/s41586-020-2706-x.

12.   Lismer, A., Dumeaux, V., Lafleur, C., Lambrot, R., Brind'Amour, J., Lorincz, M.C., and Kimmins, S. (2021). Histone H3 lysine 4 trimethylation in sperm is transmitted to the embryo and associated with diet-induced phenotypes in the offspring. Dev. Cell *56*, 671-686.e6. 10.1016/j.devcel.2021.01.014.

13.   Kaletsky, R., Moore, R.S., Vrla, G.D., Parsons, L.R., Gitai, Z., and Murphy, C.T. (2020). C. elegans interprets bacterial non-coding RNAs to learn pathogenic avoidance. Nature *586*, 445–451. 10.1038/s41586-020-2699-5.

14.   DiVito Evans, A., Fairbanks, R.A., Schmidt, P., and Levine, M.T. (2023). Histone methylation regulates reproductive diapause in Drosophila melanogaster. PLoS Genet. *19*, e1010906. 10.1371/journal.pgen.1010906.

15.   Werner, M.S., Loschko, T., King, T., Reich, S., Theska, T., Franz-Wachtel, M., Macek, B., and
Sommer, R.J. (2023). Histone 4 lysine 5/12 acetylation enables developmental plasticity of
Pristionchus mouth form. Nat. Commun. *14*, 2095. 10.1038/s41467-023-37734-z.

16.   Öst, A., Lempradl, A., Casas, E., Weigert, M., Tiko, T., Deniz, M., Pantano, L., Boenisch, U., Itskov,
P.M., Stoeckius, M., et al. (2014). Paternal diet defines offspring chromatin state and
intergenerational obesity. Cell *159*, 1352–1364. 10.1016/j.cell.2014.11.005.

17.   Webster, A.K., and Phillips, P.C. (2025). Epigenetics and individuality: from concepts to causality
across timescales. Nat. Rev. Genet. 10.1038/s41576-024-00804-z.

18.   Meyer, D.H., and Schumacher, B. (2021). BiT age: A transcriptome-based aging clock near the
theoretical limit of accuracy. Aging Cell *20*, e13320. 10.1111/acel.13320.

19.   Daines, B., Wang, H., Wang, L., Li, Y., Han, Y., Emmert, D., Gelbart, W., Wang, X., Li, W., Gibbs,
R., et al. (2011). The Drosophila melanogaster transcriptome by paired-end RNA sequencing.
Genome Res. *21*, 315–324. 10.1101/gr.107854.110.

20.   Li, B., Qing, T., Zhu, J., Wen, Z., Yu, Y., Fukumura, R., Zheng, Y., Gondo, Y., and Shi, L. (2017). A
Comprehensive Mouse Transcriptomic BodyMap across 17 Tissues by RNA-seq. Sci. Rep. *7*, 4200.
10.1038/s41598-017-04520-z.

21.   Chang, D., Serra, L., Lu, D., Mortazavi, A., and Dillman, A. (2021). A Revised Adaptation of the
Smart-Seq2 Protocol for Single-Nematode RNA-Seq. Methods Mol. Biol. *2170*, 79–99. 10.1007/978-
1-0716-0743-5_6.

22.   Serra, L., Chang, D.Z., Macchietto, M., Williams, K., Murad, R., Lu, D., Dillman, A.R., and
Mortazavi, A. (2018). Adapting the Smart-seq2 Protocol for Robust Single Worm RNA-seq. Bio
Protoc *8*. 10.21769/BioProtoc.2729.

23.   Pallares, L.F., Picard, S., and Ayroles, J.F. (2020). TM3'seq: A Tagmentation-Mediated 3'
Sequencing Approach for Improving Scalability of RNAseq Experiments. G3 (Bethesda) *10*, 143–
150. 10.1534/g3.119.400821.

24.   Lin, Y., Chen, Z.-X., Oliver, B., and Harbison, S.T. (2016). Microenvironmental Gene Expression
Plasticity Among Individual Drosophila melanogaster. G3 (Bethesda) *6*, 4197–4210.
10.1534/g3.116.035444.

25.   Werkhoven, Z., Bravin, A., Skutt-Kakaria, K., Reimers, P., Pallares, L.F., Ayroles, J., and de Bivort,
B.L. (2021). The structure of behavioral variation within a genotype. eLife *10*. 10.7554/eLife.64988.

26.   Wang, A.J., Wibisono, P., Geppert, B.M., and Liu, Y. (2022). Using single-worm RNA sequencing to
study C. elegans responses to pathogen infection. BMC Genomics *23*, 653. 10.1186/s12864-022-
08878-x.

27.   Perez, M.F., Francesconi, M., Hidalgo-Carcedo, C., and Lehner, B. (2017). Maternal age generates
phenotypic variation in Caenorhabditis elegans. Nature *552*, 106–109. 10.1038/nature25012.

28.   Gouvêa, D.Y., Aprison, E.Z., and Ruvinsky, I. (2015). Experience Modulates the Reproductive
Response to Heat Stress in C. elegans via Multiple Physiological Processes. PLoS ONE *10*,
e0145925. 10.1371/journal.pone.0145925.

29.   Hirsh, D., Oppenheim, D., and Klass, M. (1976). Development of the reproductive system of
Caenorhabditis elegans. Dev. Biol. *49*, 200–219. 10.1016/0012-1606(76)90267-0.

30.   Stumpf, C.R., Kimble, J., and Wickens, M. (2008). A Caenorhabditis elegans PUF protein family
with distinct RNA binding specificity. RNA *14*, 1550–1557. 10.1261/rna.1095908.

31.   Lublin, A.L., and Evans, T.C. (2007). The RNA-binding proteins PUF-5, PUF-6, and PUF-7 reveal
multiple systems for maternal mRNA regulation during C. elegans oogenesis. Dev. Biol. *303*, 635–
649. 10.1016/j.ydbio.2006.12.004.

32.   Jänes, J., Dong, Y., Schoof, M., Serizay, J., Appert, A., Cerrato, C., Woodbury, C., Chen, R.,
Gemma, C., Huang, N., et al. (2018). Chromatin accessibility dynamics across C. elegans
development and ageing. eLife *7*. 10.7554/eLife.37344.

33.   Ahringer, J., and Gasser, S.M. (2018). Repressive Chromatin in Caenorhabditis elegans:
Establishment, Composition, and Function. Genetics *208*, 491–511. 10.1534/genetics.117.300386.

34.   Evans, K.J., Huang, N., Stempor, P., Chesney, M.A., Down, T.A., and Ahringer, J. (2016). Stable
Caenorhabditis elegans chromatin domains separate broadly expressed and developmentally
regulated genes. Proc Natl Acad Sci USA *113*, E7020–E7029. 10.1073/pnas.1608162113.

35.   Serizay, J., Dong, Y., Jänes, J., Chesney, M., Cerrato, C., and Ahringer, J. (2020). Distinctive
regulatory architectures of germline-active and somatic genes in C. elegans. Genome Res. *30*,
1752–1765. 10.1101/gr.265934.120.

36.   Swain, P.S., Elowitz, M.B., and Siggia, E.D. (2002). Intrinsic and extrinsic contributions to
stochasticity in gene expression. Proc Natl Acad Sci USA *99*, 12795–12800.
10.1073/pnas.162041399.

37.   Yang, C.-H., Fagnocchi, L., Apostle, S., Wegert, V., Casaní-Galdón, S., Landgraf, K., Panzeri, I.,
Dror, E., Heyne, S., Wörpel, T., et al. (2022). Independent phenotypic plasticity axes define distinct
obesity sub-types. Nat. Metab. *4*, 1150–1165. 10.1038/s42255-022-00629-2.

38.   Sulston, J.E., and Horvitz, H.R. (1977). Post-embryonic cell lineages of the nematode,
Caenorhabditis elegans. Dev. Biol. *56*, 110–156. 10.1016/0012-1606(77)90158-0.

39.   Tolkin, T., and Hubbard, E.J.A. (2021). Germline Stem and Progenitor Cell Aging in C. elegans.
Front. Cell Dev. Biol. *9*, 699671. 10.3389/fcell.2021.699671.

40.   Mignerot, L., Gimond, C., Bolelli, L., Bouleau, C., Sandjak, A., Boulin, T., and Braendle, C. (2024).
Natural variation in the Caenorhabditis elegans egg-laying circuit modulates an intergenerational
fitness trade-off. eLife *12*. 10.7554/eLife.88253.

41.   Pérez-Lluch, S., Blanco, E., Tilgner, H., Curado, J., Ruiz-Romero, M., Corominas, M., and Guigó, R.
(2015). Absence of canonical marks of active chromatin in developmentally regulated genes. Nat.
Genet. *47*, 1158–1167. 10.1038/ng.3381.

42.   Rea, S.L., Wu, D., Cypser, J.R., Vaupel, J.W., and Johnson, T.E. (2005). A stress-sensitive reporter
predicts longevity in isogenic populations of Caenorhabditis elegans. Nat. Genet. *37*, 894–898.
10.1038/ng1608.

43.   Kinser, H.E., Mosley, M.C., Plutzer, I.B., and Pincus, Z. (2021). Global, cell non-autonomous gene
regulation drives individual lifespan among isogenic C. elegans. eLife *10*. 10.7554/eLife.65026.

44.   Mosley, M.C., Kinser, H.E., Martin, O.M.F., Stroustrup, N., Schedl, T., Kornfeld, K., and Pincus, Z.
(2024). Similarities and differences in the gene expression signatures of physiological age versus
future lifespan. Aging Cell, e14428. 10.1111/acel.14428.

45.   Liao, Y., Smyth, G.K., and Shi, W. (2013). The Subread aligner: fast, accurate and scalable read
mapping by seed-and-vote. Nucleic Acids Res. *41*, e108. 10.1093/nar/gkt214.

46.   Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics *26*, 139–140.
10.1093/bioinformatics/btp616.

47.   Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., and Team, R.C. (2007). Linear and nonlinear mixed
effects models. R package version.

48.   Bates, D., Maechler, M., Bolker, B., and Walker, S. (2015). Package "lme4." ….

49.   Yu, G., Wang, L.-G., Han, Y., and He, Q.-Y. (2012). clusterProfiler: an R package for comparing
biological themes among gene clusters. OMICS *16*, 284–287. 10.1089/omi.2011.0118.

50.   Friedman, J., Hastie, T., and Tibshirani, R. (2021). Package "glmnet." CRAN R ….

51.   Wilson, R., Le Bourgeois, M., Perez, M., and Sarkies, P. (2023). Fluctuations in chromatin state at
regulatory loci occur spontaneously under relaxed selection and are associated with epigenetically
inherited variation in C. elegans gene expression. PLoS Genet. *19*, e1010647.
10.1371/journal.pgen.1010647.

52.   Quinlan, A.R. (2014). BEDTools: The Swiss-Army Tool for Genome Feature Analysis. Curr. Protoc.
Bioinformatics *47*, 11.12.1-34. 10.1002/0471250953.bi1112s47.

53.   Kamath, R.S., and Ahringer, J. (2003). Genome-wide RNAi screening in Caenorhabditis elegans.
Methods *30*, 313–321. 10.1016/s1046-2023(03)00050-1.

54.   Ahringer, J. (2006). Reverse genetics. WormBook. 10.1895/wormbook.1.47.1.

55.   Van Willigen, B., and Maintainer, R. (2017). Package "nlme." Linear and ….

56.   Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
Durbin, R., and 1000 Genome Project Data Processing Subgroup (2009). The Sequence
Alignment/Map format and SAMtools. Bioinformatics *25*, 2078–2079. 10.1093/bioinformatics/btp352.