The chromatin accessibility dynamics during cell fate specifications in zebrafish early embryogenesis

Qiushi Xu[1,2,§], Yunlong Zhang[2,§], Wei Xu[3], Dong Liu[2], Wenfei Jin[2], Xi Chen[2,*], Ni Hong[2,*]

[1]Harbin Institute of Technology, Harbin, China
[2]Shenzhen Key Laboratory of Gene Regulation and Systems Biology, Department of Systems Biology, School of Life Sciences, Southern University of Science and Technology, Shenzhen 518055, Guangdong, China
[3]GMU-GIBH Joint School of Life Sciences, The Guangdong-Hong Kong-Macau Joint Laboratory for Cell Fate Regulation and Diseases, Guangzhou Medical University, Guangdong, China

[§]These authors contributed equally to this work

*Correspondences:

Xi Chen (chenx9@sustech.edu.cn)
Ni Hong (hongn@sustech.edu.cn)

Abstract

Chromatin accessibility plays a critical role in the regulation of cell fate decisions. Although gene expression changes have been extensively profiled at the single-cell level during early embryogenesis, the dynamics of chromatin accessibility at *cis*-regulatory elements remain poorly studied. Here, we used a plate-based single-cell ATAC-seq method to profile the chromatin accessibility dynamics of over 10,000 nuclei from zebrafish embryos. We investigated several important time points immediately after zygotic genome activation (ZGA), covering key developmental stages up to dome. The results revealed key chromatin signatures in the first cell fate specifications when cells start to differentiate into enveloping layer (EVL) and yolk syncytial layer (YSL) cells. Finally, we uncovered many potential cell-type specific enhancers and transcription factor motifs that are important for the cell fate specifications.

Introduction

In early embryogenesis, the totipotent zygote first enters the cleavage stage during which a series of synchronised mitotic divisions occur (1). The embryo is transcriptionally quiescent, and the progress is mainly driven by maternal factors. The main wave of zygotic genome activation (ZGA) when the genome of the embryo begins to make their own RNAs and proteins only occurs at later cell cycles (2). Starting from ZGA, embryonic cells become asynchronous along developmental trajectories. They undergo a series of complex events, such as cellular differentiation and migration, and eventually gives rise to different cell types with committed fates and distinct functions (3). Understanding how cells acquire specific fates is one of the key goals in developmental biology. Due to the heterogeneous and asynchronous nature of the developmental processes, single-cell genomics methods are necessary and serve as powerful approaches for the unbiased investigation of molecular profiles of each cell during development.

In zebrafish, recent single-cell RNA-seq (scRNA-seq) developmental atlases provided gene expression overviews of all major cell types from embryonic to adult stages (4–10). For example, the first two zebrafish scRNA-seq atlases (4, 5) described in great details the gene expression patterns during embryogenesis at unprecedented temporal and cellular resolution, revealing that cells exhibit potential plasticity during development and cell fate specification is a gradual and non-binary process. Furthermore, those transcriptomic profiling studies revealed stage-specific gene expression programs, branching points of emerging cell fates and key regulators during different developmental trajectories (4–10). While those datasets serve as invaluable resources to study organismal development, the investigation regarding the regulatory events underlying those transcriptomic and cell fate changes remains limited.

The progressive cell fate specifications are driven by the interaction between transcription factors and *cis*-regulatory elements located in regions such as promoters and enhancers (11). In general, chromatin around *cis*-regulatory elements needs to be in an accessible state to allow transcription factor binding. Therefore, genome-wide investigation of chromatin accessibility will provide insights on the molecular mechanisms of cell fate commitments. In recent years, several studies used the assay for transposase-accessible chromatin using sequencing (ATAC-seq) to interrogate the chromatin accessibility landscapes during zebrafish early embryogenesis (12–14).

Those studies generated detailed chromatin dynamics during early development,
demonstrating the gradual increase of chromatin accessibility around *cis*-regulatory
elements on a global scale.

To date, most chromatin accessibility studies during zebrafish early embryogenesis
have been performed at the bulk level (12–14). A limited number of single-cell
chromatin accessibility studies mainly focused on the post-gastrula stages or adult
tissues (15–17). Therefore, it is still not clear if and when cells become different at the
chromatin level during early embryogenesis. In this study, we used a plate-based
single-cell ATAC-seq (scATAC-seq) method (18, 19) to investigate the chromatin
accessibility states of 10,518 single embryonic nuclei from ZGA up to the dome stage.
Different accessible chromatin landscapes among cells begin to emerge when the
embryo is transitioning from the sphere to the dome stage. Distinct chromatin states
of the enveloping layer (EVL), the yolk syncytial layer (YSL) and the deep layer (DEL)
cells are clearly observed. Compared to DEL cells, the EVL and YSL cells obtain their
own signature profiles where specific distal regions that are putative enhancers change
from an inaccessible to accessible state. Overall, our time-course single-cell data
provides valuable information about molecular processes underlying the first cell fate
specification.

Materials & Methods

Zebrafish maintenance
The wild-type TU (Tübingen) zebrafish were maintained at 28°C with 14/10 hours of
light/dark cycles, respectively. Embryos were collected and transferred to the zebrafish
embryo medium (5 mM NaCl, 0.17 mM KCl, 0.33 mM $CaCl_2$, 0.33 mM $MgSO_4$, and
0.002% methylene blue) at 28°C immediately after fertilization. All animal
experiments were performed with approval from the Experimental Animal Welfare
Ethics Committee, Southern University of Science and Technology.

Embryo collection and cell dissociation
Embryos at different time points after fertilisation were collected, including 3 hours
post fertilisation (hpf) (the 1k-cell stage), 3.3 hpf (the high stage), 3.7 hpf (the oblong-
sphere stage), 4.1 hpf (the sphere-dome stage) and 4.3 hpf (the dome stage). Embryos
showing inconsistent developmental status were excluded. A total of 50 to 100
embryos in each stage were collected for one batch of experiment. To get embryonic
cells, chorions were removed by treating with 1 mg/mL Pronase E (Solarbio, P8360)

at room temperature for 3 min. Then the embryos were washed three times with 10

mL 1× PBS and manually deyolked using tweezers under a microscope, and cells were

dispersed by gentle pipetting. The dispersed cells were transferred to a 1.5 mL tube

and collected by centrifugation at 500g, 4°C for 5 minutes. The pellet was resuspended

in 1×PBS-0.5% bovine serum albumin (BSA), and the cell number was counted using

a C-Chip disposable hemocytometer (INCYTO, cat. no. DHC-N01) after stained with

trypan blue (Sigma-Aldrich, T8154).

Single-cell ATAC-seq

The scATAC-seq libraries were prepared using a plate-based workflow as described

previously (18, 19). Briefly, 50,000 cells were lysed in 50 μL RSB-DTN (10 mM Tris-

HCl, pH 7.4, 10 mM NaCl, 3 mM MgCl$_2$, 0.1% IGEPAL CA-630, 0.1% Tween-20 and

0.01% digitonin) on ice for 3 minutes. The lysate was then washed with 950 μL cold

RSB-T (10 mM Tris-HCl, pH 7.4, 10 mM NaCl, 3 mM MgCl2, and 0.1% Tween-20) and

centrifuged at 1,000g, 4°C for 10 minutes. The cell pellet was resuspended in 50 μL

tagmentation mix (33 mM Tris-acetate pH 7.8, 66 mM potassium acetate, 10 mM

magnesium acetate, 16% dimethylformamide, 0.01% digitonin, and 5 μL Tn5) and

incubated at 37 °C for 30 minutes with gentle shaking. The reaction was stopped by

adding 50 μL 2× stop buffer (10 mM Tris-HCl pH 8.0, 20 mM EDTA pH 8.0). Nuclei

were centrifuged at 500 g, 4 °C for 5 minutes and then resuspended in 500 μL 0.5× PBS-

0.5% BSA, stained by adding 1.5 μL 1 μg/μL DAPI and hold on ice. The single DAPI-

positive nucleus was sorted into each individual well of a 384-well plate containing 3

μL preloaded lysis buffer (11 mM Tris-HCl pH 8.0, 11 mM NaCl, 0.22% SDS, 10 μM

Nextra_S5_short primer, and 10 μM N7xx primer), and incubated at 65 °C for 15

minutes. After a brief centrifugation, 2 μL of 10% Tween-20 and 5 μL NEBNext High-

fidelity 2× PCR Master Mix (NEB, M0541) was added to each well sequentially. Pre-

amplification in each well was performed as follows: 72°C for 5 minutes; 98 °C for 1

minute; 10 cycles of 98 °C for 10 seconds, 63°C for 30 seconds, 72°C for 20 seconds. All

pre-amplified libraries in each well from the same 384-well plate were pooled together

and transferred into a 15-mL tube. The pooled pre-amplified libraries were purified by

the QIAquick PCR purification kit (Qiagen, 28104). Excessive primers were digested

by Thermolabile Exonuclease I (NEB, M0568) at 37 °C for 30 min, and then purified

by 1.5× VAHTS DNA clean beads (Vazyme, N411-01) and eluted in 20 μL of nuclease-

free water. The pooled pre-amplified libraries from each plate were mixed with 2.5 μL

scATAC_S5xx primer (10 μM), 2.5 μL Illumina P7 primer (10 μM), and 25 μL NEBNext

High-Fidelity 2× PCR Master Mix, followed by a final PCR for amplification: 98 °C for

1 minute; 12 cycles of 98 °C for 10 seconds, 63 °C for 30 seconds, 72 °C for 20 seconds.

The final libraries with plate-specific barcodes were purified by 1.2× VAHTS DNA

clean beads and eluted in 20 μL of nuclease-free water. Equimolar libraries from

different plates were pooled and sequenced on the Illumina HiSeq X-Ten machine

using standard sequencing scheme.

Data preprocessing

After sequencing, BCL files were converted to FastQ files using bcl2fastq (v.2.19) from

Illumina. Read 1 and Read 2 FastQ files were processed using a Snakemake pipeline

(20). In brief, 3′-end adapter sequence was trimmed with fastp (21). Then the trimmed

reads were mapped to UCSC danRer11 reference genome using hisat2 (22). Reads

with mapping quality less than 30 were removed by samtools (23). Duplicated reads

were removed using the MarkDuplicates function of the Picard tool

(http://broadinstitute.github.io/picard/). Reads from single cells were merged

together and deduplicated again using samtools and Picard to generate a merged BAM

file for each stage. Peak calling of the merged BAM files was performed using MACS2

(24). The peak-by-cell matrix was generated for each stage by counting the number of

reads within peaks from individual cells with bedtools (25). The software and

packages used in this pipeline can be found in the "Snakefile" provided in the GitHub

repository(https://github.com/dbrg77/scATAC_snakemake).

Quality control (QC)

To remove low-quality cells, we computed four quality control (QC) metrics of each

cell: (1) The number of unique nuclear fragments: the number of reads mapped to the

nuclear genome after deduplication; (2) Mapping rate: the overall alignment rate

calculated by hisat2; (3) Nucleosome signal: the ratio of mono-nucleosomal fragments

(147 - 294 bp) to nucleosome-free fragments (<147 bp) which was calculated by the

"NucleosomeSignal" function from the R package Signac (26); (4) Transcriptional

start site (TSS) enrichment score: a ratio between aggregated distribution of reads

centred on the TSS and the flanking regions, which was calculated by the

"TSSErichment" function of Signac. Based on the per-stage distribution of each QC

metric, the following thresholds were used for different stages to ensure that cells in

earlier stages with lower accessibility would not be removed: 3 hpf (unique nuclear

fragments > 1000 & mapping rate > 70 & TSS enrichment > 1), 3.3 hpf and 3.7hpf

(unique nuclear fragments > 2000 & mapping rate > 70 & TSS enrichment > 1.5), 4.1

hpf (unique nuclear fragments > 2000 & mapping rate > 75 & TSS enrichment > 1.5),

4.3 hpf (unique nuclear fragments > 2000 & mapping rate > 75 & TSS enrichment >

2).

Normalization and 2D visualisation

We used the "reduce" function of the R package GenomicRanges (27) to combine all intersecting peaks between stages and created a unified set of peaks. To generate a genomic blacklist, we used MACS2 to perform peak calling on a prior "input" dataset of ATAC-seq experiment where Tn5 was used to cut zebrafish naked genomic DNA (14). Peaks that intersected with the blacklist were removed. Based on the remaining peaks, we created a peak-by-cell count matrix. The count matrix was normalized them using the term frequency–inverse document frequency (TF–IDF) transformation. Then singular value decomposition (SVD) was applied to the matrix to obtain a lower-dimensional representation. Subsequently, the 2 - 30 latent semantic indexing (LSI) components were used to perform UMAP to visualise the final result in 2-dimension plots.

Data integration of different stages

To integrate data from all stages, we used BBKNN (28) for batch correlation. We created the combined peak-by-cell matrix with all stages and performed the LSI analysis using the same workflow described above. The top 20 LSI components were used as an input to the bbknn function from the Python package Scanpy (29). To integrate data for later stages (4.1 hpf and 4.3 hpf) for a detailed analysis of cell types, we used the "IntegrateEmbeddings" function of Signac to integrate data from 4.1 hpf (sphere-dome) and 4.3 hpf (dome) stages and projected them into a shared low-dimensional space. After that, new LSI-coordinates (rLSI) of cells were generated, which represented the low-dimensional cell embeddings after integration. We clustered sphere-dome- and dome-stage cells and visualised them using the "FindCluster" (resolution = 0.6) and RunUMAP functions from Signac.

Identification of peak modules

We used the non-negative matrix factorization (NMF) algorithm from Python scikit-learn toolkit (30) (version 1.1.0) to factorise the normalized peak-cell matrix into two matrices: a matrix H (Peaks by Peak modules), and a matrix W (Peak modules by Cells). The H matrix essentially groups peaks into peak modules such that peaks within the same peak module have similar patterns across single cells, that is, they co-vary among different cells. The number of peak modules is often much lower than the number of original peaks. This is reminiscent of the gene expression programs after performing NMF on a gene expression matrix, where genes that co-vary among cells are grouped into programs. The number of peak modules (k) was tested from 3 to 60, and the optimal number was determined to be 25. Modules with high usages in only

224 one batch were possibly caused by the batch effect and ignored for the subsequent
225 analyses. Monotonic increasing (modules 2, 3, 4 and 6) and decreasing module
226 (module 10) during embryo development were identified based on the usage
227 distribution.
228
Identification of differentially accessible sites
To identify differentially accessible sites between cell clusters, we performed peak
calling of each cell cluster in the 4.1 hpf (sphere-dome) and 4.3 hpf (dome) stages.
After combining the peaks and normalising the resulting matrix, we performed
differential accessibility (DA) test to determine the differential accessible sites and
marker peaks of each cell type by the "FindMakers" function of Seurat (min.pct = 0.25,
logfc.threshold = 0.5). Logistic regression (LR) method was chosen in this step. The
total number of fragments of each cell were used as a latent variable to mitigate the
effect of differential sequence depth on the result.
238
Motif analysis
For the *de novo* motif discovery, the 2,000 peaks from each peak module or EVL or YSL
marker peak were used as the input for the "findMotifsGenome.pl" function from
HOMER (31) with the parameter "-size 500" to look for the centre 500 bp of each peak.
To look at motif deviations across different cells, position frequency matrix (PFM) of
non-redundant vertebrates' transcription factors (TF) binding profiles form the
JASPAR database 2022 (32) version was used as the input to the "AddMotifs" function
of Signac. Then chromVAR (33) was invoked by the "RunChromVAR" function was
used to compute a per-cell motif activity score and created a motif-cell matrix. The
differential activity was computed using the "FindMarkers" function.
249
Gene activity score calculation
To quantify the gene activities of homeodomain transcription factors, the function
"GeneActivity" from Signac was used to generate a gene activity matrix of all
homeodomain transcription factors. In this process, gene coordinates were extracted
and extended to include the 2 kb region upstream of the transcriptional start site of
the gene. Then the number of fragments mapped to the extended region consisting of
the 2 kb proximal promoter and the gene body, were counted and treated as the gene
activity score for the gene. Finally, the function "AverageExpression" was used to get
the average values of the fragment counts for each cell type.
259
DNA methylation analysis

We used the publicly available WGBS data set (ArrayExpress, accession number E-
MTAB-12535) at the 1k stage to investigate the DNA methylation profiles around LTR
regions from module 10 peaks. BigWig files in danRer10 containing DNA methylation
values were downloaded. The UCSC genome browser utility "liftOver" was used to
convert the coordinates of LTR regions from danRer11 to danRer10. Then the function
"computeMatrix" from deepTools (34) were used to calculate methylation signals for
given regions with the following option: --skipZeros -a 1000 -b 1000 --binSize 50.
268
Co-accessibility analysis
The co-accessibility of peaks was estimated by Cicero (35). We calculated the co-
accessibility score for each peak pair using the "run_cicero" function. The peak pairs
with a co-accessibility score greater than 0.1 were retained. Next, we generated a co-
accessibility network using the "generate_ccans" function based on the co-accessibility
score framework. The result was visualized using the "LinkPlot" function from Signac.
275
Trajectory analysis
We used a diffusion map (36) to perform trajectory analysis of the 4.1 hpf (sphere-
dome) and 4.3 hpf (dome) stage cells. Scores for the top 12 rLSI components were
used as the input to compute the neighbourhood graph of cells. Then we generated a
diffusion map embedding using the "diffmap" function from Scanpy. To calculate the
pseudotime value of each cell, we set DEL1 as the root due to its lower accessibility,
and performed diffusion pseudotime (DPT) analysis using the "dpt" function. Finally,
we fitted a vector generalized additive model (VGAM) with a degree of freedom of 3
to show the dynamics of peak accessibility in pseudotime using the R package VGAM
(37).
286
287 # Results
288
Single-cell ATAC-seq reveal the accessible chromatin landscape from ZGA to dome
in zebrafish
To capture the earliest time point when embryonic cells become different from each
other, a plate-based scATAC-seq method previously developed in our labs (18, 19) was
used to profile the chromatin accessibility of each single nucleus in the zebrafish
embryos (**Figure 1A** and **Supplementary Figure 1A**). Several key developmental time
points immediately after ZGA were covered, spanning the 1k, high, oblong, sphere
and dome stages. A total of 11,520 nuclei from staged embryos were profiled and over
3 billion read pairs were obtained with an overall mapping rate of >75% to the

zebrafish genome (**Supplementary Figure 1B**). After quality control (see **Methods;**
Supplementary Figure 1C), 10,518 nuclei were used for further analysis
(**Supplementary Table 1**). After duplication removal, the median numbers of unique
fragments mapped to the nuclear genome range from ~26,000 to ~60,000
(**Supplementary Figure 1B**), which is at the higher end compared to the scATAC-seq
atlases in other stages and adult tissues (15–17). Peak calling using MACS2 from
merged single nuclei (see **Methods**) identified 446,501 open chromatin regions. Visual
inspections on a genome browser of the read pile-up from aggregated single nuclei
within the same time point indicated the data was of good quality (**Figure 1B** and
Supplementary Figure 1D). The aggregated single-nucleus profiles were also
comparable to a recent high-quality bulk ATAC-seq data in the corresponding stages
(14).
310
Following dimensionality reduction by the latent semantic indexing analysis(26, 38),
uniform manifold approximation and projection (UMAP) (39) was used to project the
data onto a two-dimensional space for visualisation per stage (**Figure 1C**). No
apparent separation of nuclei was observed from 3 to 3.7 hours post fertilisation (hpf),
indicating cells were mostly homogeneous at the chromatin level during the first few
stages following ZGA. Notably, discrete nuclei clusters started to emerge from 4.1 hpf
and onwards (**Figure 1C**). The observation suggested that cells began to acquire
distinct accessible chromatin profiles for the first time at the transition from the sphere
to the dome stage.
320
Key peak modules have different genomic features and dynamics during
development
To get an overview of the chromatin accessibility landscape during this developmental
period, a batch balanced k-nearest neighbour (BBKNN) method (28) was used to
remove batch effect and integrate all stages into a single UMAP space (**Figure 2A** and
Supplementary Figure 2A). Next, non-negative matrix factorisation (NMF) (40, 41)
was utilised to find peak modules that covary across single nuclei throughout the
development (**Figure 2B**). The first matrix from the NMF represents how much each
module is used by each nucleus. The usage of each peak module in single nuclei was
examined, and peak modules exhibiting high usage only in one batch were ignored
(**Figure 2B** and **Supplementary Figure 2B**). Eventually, five peak modules (peak
modules 2, 3, 4, 6 and 10) with consistent and reproducible usage across batches were
used for the downstream analysis (**Supplementary Table 2**).
334

Consistent with the notion that chromatin accessibility is gradually established during
ZGA in zebrafish (12–14) and other species (42–47), the overall scATAC-seq
enrichment around the transcriptional start sites (TSS) was increasing throughout the
time course (**Supplementary Figure 2C**). The usages of most peak modules (modules
2, 3, 4, and 6) were increasing during the developmental time course and reached
maximum in different subsets of cells at later stages (**Figure 2C**). The examination of
the aggregates of scATAC-seq reads showed peaks within these modules became
gradually accessible during the development process (**Figure 2D**). Interestingly, a
peak module (module 10) with decreasing usage was also identified (**Figure 2C**),
indicating certain genomic regions are turning into an inaccessible state following
ZGA (**Figure 2D**).
346
To further investigate the characteristics of the five peak modules, public ChIP-seq
experiments of important histone marks and histone variants across different
developmental stages in zebrafish were collected (48–50). A hidden Markov model,
ChromHMM (51), was used to categorise the zebrafish genome into eight different
states based on the ChIP-seq data, including various types of promoters and enhancers
(**Figure 3A**). The distributions of the five peak modules across the chromatin states
were quite different. Modules 2 and 3 were exclusively located in the constantly active
promoter regions (**Figure 3A**, ChromHMM state 1), and modules 4 and 6 were mainly
located in the enhancer regions (**Figure 3A**, ChromHMM states 3, 5 and 6).
Consistently, peaks from modules 2 and 3 were close to the TSS of annotated genes,
and most peaks from modules 4, 6 and 10 were located in the intergenic regions
(**Supplementary Figure 3A**). Notably, module 6 was preferentially located in the
dome-specific enhancers (ChromHMM state 6) and the post-ZGA stage enhancers
(ChromHMM state 5). Considering the usage of module 6 was most prominent in a
subset of cells during the late stages (**Figure 2C**), peaks within module 6 were likely
to play an important role in cell type specification (discussed later). Gene ontologies
using GREAT (52) also suggested distinct functions for genes associated with each
peak module. For example, the most enriched biological processes in modules 2 and 3
are translation and chromatin assembly (**Supplementary Figure 3B**), indicating these
processes are critical throughout early embryo development which is consistent with
the general knowledge (1).
368
One interesting observation was that module 10 was mainly located in the regions
where transposable elements (TEs) reside (**Figure 3A,** ChromHMM state 8). Visual
inspections on individual TEs showed the scATAC-seq reads covered the entire TE,
and the chromatin status around the TEs became gradually closed after ZGA (**Figure**

3B). A more detailed analysis on TE classes and families showed >80% of the top 2,000
peaks from module 10 overlapped long terminal repeats (LTR) (**Figure 3C**). Within
all LTRs from module 10, the vast majority were dictyostelium intermediate repeat
sequence (DIRS) (**Supplementary Figure 3C**). Intriguingly, a recent TE analysis (53)
of gene expression data (54) during zebrafish early embryogenesis indicated LTRs
were highly expressed immediately following ZGA, but the dynamic of the expression
was not concordant with the chromatin accessibility changes (**Figure 2C** and
Supplementary Figure 3D). These results together suggest a more complex
regulatory mechanism of LTRs from module 10.
382

De novo motif analysis (31) revealed different transcription factor motifs enriched
within the five peak modules (**Figure 3D**). While no prominent motifs were identified
in module 2, motifs bound by factors including Hsf1 (55) and Nfya (44) that were
important for ZGA reported in mice were more enriched in modules 3 and 4,
respectively (**Figure 3D**). In addition, several motifs were identified as specifically
enriched in peaks from module 10. When the DNA methylation profiles were
examined around these LTRs within module 10 at 3 hpf (1k stage) from a recent study
(56), they were quite different from DNA methylation profiles of typical open
chromatin regions (**Supplementary Figure 3E**). Therefore, the exact meaning and
functional implications of reduced chromatin accessibility of LTRs in module 10
during development need further investigation.
394

Distinct chromatin signatures of cell types emerge around the sphere and dome
stages
Since discrete cell populations started to show up around the 4.1 hpf (sphere-dome)
stage (**Figure 1C**) and cells from the 4.1 hpf and 4.3 hpf (dome) had similar chromatin
profiles (**Figure 2A**), all nuclei (a total of 4,376 nuclei) from these two late stages were
taken out for a more detailed analysis in order to identify emerging cell types.
Clustering on the data from the 4.1 hpf and the 4.3 hpf time points revealed four
different cell clusters (**Figure 4A** and **Supplementary Figure 4A**). Based on the genes
around the differentially accessible peaks in each cluster, two of the clusters were
identified as the enveloping layer (EVL) cells, a type of differentiated cells that covers
the outside of the embryo (57), and the yolk syncytial layer (YSL) cells which are an
extra-embryonic tissue that contributes to the patterning of the embryo (58) (**Figure**
4A). High chromatin accessibility around canonical marker genes, such as *krt18a.1*,
krt4, *cldne* and *gata3*, were observed specifically within the EVL cell cluster (**Figure 4B**,
Figure 4C and **Supplementary Figure 4B**). Consistent with previous studies that *afp4*
and *cyp11a1* were specifically expressed in the YSL cells during early embryogenesis

(59, 60), clear open chromatin peaks around the *afp4* and *cyp11a1* loci were observed
only in the YSL cell cluster (**Figure 4B** and **C**). The rest of the two cell clusters (**Figure
4A,** DEL1 and DEL2)constituted the majority of the cells and were most likely to be
the undifferentiated pluripotent cells in the deep layer (DEL) between EVL and YSL.
Although both were considered to be located in the deep layer, differences between
DEL1 and DEL2 were clearly observed. Many peaks, especially those from module 2,
showed higher accessibility in cells from the DEL2 cluster (**Supplementary Figure
4A**). These peaks were mainly associated with genes involved in translation, mitotic
cell cycle, mRNA metabolic process and RNA splicing (**Supplementary Figure 3B**),
suggesting DEL2 cells may have higher activity in those biological processes compared
to cells in the DEL1 cluster.

We also examined the expressions of the key marker genes around similar stages from
a scRNA-seq atlas of zebrafish development (4). The expression dynamics of EVL
marker genes were concordant with the chromatin accessibility changes during the
time course, where those genes remained silent before the dome stage and only started
to express in the EVL subset of cells in the dome stage (**Supplementary Figure 4C**).
Since the YSL cells were not identified in the scRNA-seq atlas (4), some marker genes
such as *afp4* were not detected. The YSL marker gene *cyp11a1* was detected as early as
the high stage, possibly due to the fact it is a maternal transcript (60). The discrepancy
of the YSL cells between scRNA-seq and scATAC-seq data was probably due to
technological difference (see **Discussion**).

To further explore the relationship among those clusters during development, nuclei
from the same cluster were first pooled and peak calling was performed for each
cluster aggregate (see **Methods**). Then the resulting cell-by-peak matrix were used for
the trajectory inference with diffusion pseudotime (DPT) (36). The DPT analysis split
the cells into three different trajectories (**Figure 4D**). The main trajectory from DEL1
to DEL2 represented the differentiation of pluripotent cells which ultimately would
lead to the formation of all other body parts of the zebrafish. Two branches emerged
from the main trajectory represented the EVL and YSL fate acquisition, respectively
(**Figure 4D**). The pseudotemporal ordering of cells also provided key dynamic
information of open chromatin sites along each branch (**Figure 4E**). For example, in
the EVL branch, the regulatory regions around key marker genes *krt4*, *krt18a.1* and
gata3 became fully open at the end of the pseudotime, but the *krt4* and *krt18a.1* loci
started to gain accessibility earlier than *gata3* (**Figure 4E**).

Newly identified cell-type specific enhancers and transcription factors drive the EVL and YSL cell fate acquisitions

We noticed the peak modules 2 and 6 identified by NMF reached the highest usage in the DEL2 and EVL cell clusters, respectively (**Supplementary Figure 4A**). In order to systematically find marker peaks of each cell cluster, differential accessibility analysis was performed among peaks called by the cell cluster aggregate in the previous section (**Supplementary Table 3**). While very few specific peaks were found for the DEL1 or DEL2 cell cluster, many novel marker peaks that were accessible only in the EVL or YSL cell cluster were identified (**Supplementary Table 3**). Compared to the eight-state ChromHMM model (**Figure 3A**), both the EVL and the YSL marker peaks were mainly located in the dome specific enhancer and the post-ZGA stage enhancer regions (**Figure 5A**). A more detailed investigation of ChIP-seq reads of key histone modifications (50) around the marker peaks indicated low level of H3K4me3 and high levels of H3K4me1 and H3K27ac around the centre of EVL and YSL marker peaks (**Figure 5B**). The H3K4me1 and H3K27ac signals around the marker peaks were observed at the dome stage and persisted in the 80% epiboly stage, indicating these newly identified marker peaks by our scATAC-seq data are putative enhancers that are important for the cell fate specification at the dome stage and remain active at least through the 80% epiboly stages.

To find out potential transcription factors that shape the chromatin profiles of the EVL and YSL cells, *de novo* motif discovery was performed on the two sets of marker peaks, respectively. The most enriched motif from the EVL marker peaks resembled the DNA binding consensus of the CP2 family, which could be bound by transcription factors such as Grhl and Tfcp2 (**Figure 5C**). A GC-rich motif that was similar to the Klf family and Sp (specificity protein) family consensus and a third one that resembled the IRF family motif were also found enriched within the EVL marker peaks (**Figure 5C**). Motif activity inferred by chromVAR (33) confirmed the results that the aforementioned motifs had the highest activity in the EVL cells (**Figure 5D**). The motif results point to the possibility that the CP2, Klf, Sp and Irf families might play an important role in shaping the chromatin state of EVL cells.

Two versions of the homeobox motifs were found over-represented within the YSL marker peaks (**Figure 5C** and **D**). The top motif, homeobox 1, was palindromic and consisted of the half-site TRATY with a 1-bp spacer, where R stands for A or G and Y stands for C or T (**Figure 5C**). The motif was previously shown to be bound by the mouse homeodomain transcription factors, such as Cphx1 and Phox2a/b (61–63), though no orthologs of the *Cphx1* gene was present in zebrafish. Another version of

the homeodomain motif, homeobox 2, contained the non-canonical core sequence TGAT followed by the canonical homeobox core TAAT (**Figure 5C**) (61, 64). The homeobox 2 motif was shown to be bound by transcription factors such as Pdx1 and Hoxa (61, 64–66). These results suggested the homeodomain transcription factors were important to establish the chromatin signature of the YSL cells. Interestingly, *dharma*, a homeodomain transcription factor expressed in the YSL has previously been shown to be important for the function of YSL cells (58, 67), and it can potentially bind to the half-site of the homeobox 1 motif (66).

In order to narrow down the key transcription factors that are important for the establishment of the EVL and YSL fates, the expression levels of all members from relevant transcription factor families (68) were checked using the scRNA-seq atlas of early zebrafish embryo development (4). Since YSL cells were not present in the scRNA-seq atlas, we first focused on the gene expression of relevant transcription factor families in EVL cells. Intuitively, a transcription factor must be expressed in the cell to be functional. Therefore, we checked expression levels of all transcription factors that can potentially bind to the three main motifs discovered from the EVL mark peaks (**Figure 5E**). Within the CP2 family, *grhl3* is the only member that could be detected, and it was specifically expressed in EVL cells at the dome stage (**Figure 5E**). Similarly, *klf17* had the highest expression level in EVL cells at the dome stage, though *klf2*, *klf6a* and *sp1* that are able to bind to similar DNA motif could also be detected at the dome stage (**Figure 5E**). In the Irf family, only the mRNA of *irf2a* could be detected, but its expression level went down since the high stage (**Figure 5E**). Previous functional studies implicated the importance of Grhl3 and Klf17 during the development of EVL cells (69, 70). Combined with our scATAC-seq data, the results indicated that those transcription factors promote EVL development and functions through the establishment of EVL specific chromatin profiles at enhancer regions. Next we systematically investigated the gene activity scores of all homeodomain transcription factors in the YSL cluster by measuring their promoter plus gene body accessibilities (35). A total of 73 homeodomain transcription factors were found to have much higher gene activity scores in the YSL cells compared to other cell clusters (**Figure 5F**, red box and **Supplementary Table 4**), including *dharma* and *mxtx2* that were previously shown to be important for the development of YSL (67, 71). Interestingly, *en1a*, a predicted paralog of mouse *Cphx1*, was also found to have higher gene activity score in the YSL cells (**Figure 5F**). This analysis suggests the 73 homeodomain transcription factors might be potentially expressed in the YSL cells and could possibly bind to the homeobox motifs enriched within the YSL marker peaks.

Finally, co-accessibility analysis (35) was performed to link distal regulatory elements
to their potential target genes. In this way, many putative distal enhancers were linked
to important marker genes in EVL and YSL cells, respectively (**Figure 6** and
Supplementary Figure 5). Many of the enhancers have at least one previously
described motif and reproducible H3K27ac signals from two independent studies (49,
50) (**Figure 6**). This demonstrated that our data have identified many potential cell-
type specific enhancers that may play critical roles in the cell fate specifications of EVL
or YSL cells.

Discussion

The acquisition of cell-type specific molecular signatures during early development is
a key feature in all multi-cellular organisms. Recent technological development has
allowed the interrogation of various molecular profiles at the single cell level,
including gene expression, chromatin accessibility and chromosome conformation etc
(72, 73). The techniques offer excellent opportunities to study cell fate specifications
during embryo development. In zebrafish, several single-cell atlases have been
generated in terms of gene expression (4–10) and chromatin accessibility (15–17).
They mainly focused on either the late stages of embryogenesis (mostly after gastrula)
or adult tissues, partly due to the fact that early embryo development is driven by
maternal factors in zebrafish (2).

In this study, we investigated the chromatin accessibility landscapes at the single-
nucleus level, with over 10,000 nuclei across five key stages that span the first hour
immediately following ZGA. We identified important open chromatin peak modules
with distinct sequence features during the developmental time course (**Figures 2** and
3). More importantly, the data suggested cells started to gain distinct chromatin
signatures when they progressed from the sphere to the dome stage. Cell-type specific
distal regulatory elements for EVL and YSL cells, respectively, were identified using
our scATAC-seq data (**Figures 4** and **5**). They were marked by low-level of H3K4me3
and high level of H3K4me1 and H3K27ac and contained important transcription factor
binding motifs (**Figures 5** and **6**), indicating those distal regulatory elements are
putative enhancers that are important for the EVL and YSL cell fate specifications.

The YSL cells were not present around the dome stage in the previous two scRNA-seq
atlases (4, 5). Since the YSL cells are present as nuclei in the syncytium on the surface

of the yolk cell (58), it is likely that the typical single-cell isolation procedures in the
scRNA-seq experiment would result in the loss of those nuclei during the experiments.
The ChIP-seq data of histone marks used to annotate the cell-type specific regulatory
regions were taken from bulk experiments at the dome stage (49, 50). Our single-cell
data suggested the ChIP signals of those histone marks might come from only a subset
of cells, EVL or YSL, at the dome stage, and the bulk ChIP-seq experiments seem to be
sensitive enough to enrich and uncover those signals (**Figure 5B** and **Figure 6**). With
the development of single-cell ChIP-like techniques (74–83), it is possible to
characterise potential cell-type specific enhancers more accurately in the future.

Our scATAC-seq data showed that EVL and YSL cells emerged and acquired distinct
chromatin profiles at 4.1 hpf (**Figure 1C** and **Figure 4A**), indicating those cells start to
differentiate and take specific cell fates. The classic cell fate commitment study from
Ho and Kimmel showed cells at the dome stage remain pluripotent and uncommitted
(84). When examined the chromatin status of key pluripotent genes, including *pou5f3*,
nanog, *sox19b*, in different cell clusters at the dome stage, all of them exhibit a
transcriptional permissive state with many open chromatin peaks covering the gene
loci (**Supplementary Figure 6**). These observations suggest that even EVL and YSL
cells start to obtain specific chromatin features that are distinct from DEL cells, their
cell fates are not fully committed yet at the dome stage.

In general, our data serve as a useful resource that provides an overview of the *cis*-
regulatory dynamics during the first cell fate specification of zebrafish early
embryogenesis. Combined with functional studies, gene expression profiles and other
data, it allows more detailed investigation of the mechanism underlying cell fate
decisions.

Data availability

The raw sequencing data, the processed peak by cell count matrix, the R object and
the meta data have all been deposited at ArrayExpress under the accession number E-
MTAB-13115.

Funding

This study was supported by National Key R&D Program of China (2021YFF1200900 to W.J, X.C and N.H), National Natural Science Foundation of China (32322019 to X.C, 32200509 to W.X), Guangdong Basic and Applied Basic Research Foundation (2023A1515011662 and 2022B1515120077 to X.C, 2023A1515011908 to N.H), Shenzhen Science and Technology Program (20220815094330000 to X.C).

Competing Financial Interests

None declared.

Acknowledgement

We thank all members from the Chen and Jin labs for the helpful discussion of the project. We also thank Dr. Zilong Wen for the helpful suggestions on the writing of the manuscript. We thank Xibin Lu for the excellent support of FACS. We acknowledge the assistance of SUSTech Core Research Facilities. The computational work was supported by Center for Computational Science and Engineering at Southern University of Science and Technology. Part of Figure 1a and Figure 4a were created with BioRender.com.

Author contributions

X.C and N.H conceived the project. Q.X, Y.Z, W.X, N.H and X.C designed the experiments. Q.X performed the experiments with the help from W.X and N.H. Y.Z, carried out the computational analysis with the help from X.C, Q.X and N.H. D.L, J.W, X.C and N.H supervised the entire project. All authors contributed to the writing.

Code availability

624 The code used for the data processing and analysis mentioned in the method section
625 is available on the GitHub repository
626 https://github.com/helianfeixing/Zebrafish_scATAC

References

1. Lee,M.T., Bonneau,A.R. and Giraldez,A.J. (2014) Zygotic Genome Activation During the
Maternal-to-Zygotic Transition. *Annu. Rev. Cell Dev. Biol.*, **30**, 581–613.

2. Schier,A.F. (2007) The Maternal-Zygotic Transition: Death and Birth of RNAs. *Science*, **316**,
406–407.

3. Burton,A. and Torres-Padilla,M.-E. (2014) Chromatin dynamics in the regulation of cell fate
allocation during early embryogenesis. *Nature Reviews Molecular Cell Biology*, **15**,
723–735.

4. Farrell,J.A., Wang,Y., Riesenfeld,S.J., Shekhar,K., Regev,A. and Schier,A.F. (2018) Single-cell
reconstruction of developmental trajectories during zebrafish embryogenesis.
Science, **360**, eaar3131.

5. Wagner,D.E., Weinreb,C., Collins,Z.M., Briggs,J.A., Megason,S.G. and Klein,A.M. (2018)
Single-cell mapping of gene expression landscapes and lineage in the zebrafish
embryo. *Science*, **360**, eaar4362.

6. Farnsworth,D.R., Saunders,L.M. and Miller,A.C. (2020) A single-cell transcriptome atlas for
zebrafish development. *Developmental Biology*, **459**, 100–108.

7. Jiang,M., Xiao,Y., E,W., Ma,L., Wang,J., Chen,H., Gao,C., Liao,Y., Guo,Q., Peng,J., *et al.*
(2021) Characterization of the Zebrafish Cell Landscape at Single-Cell Resolution.
Frontiers in Cell and Developmental Biology, **9**, 743421.

8. Sur,A., Wang,Y., Capar,P., Margolin,G. and Farrell,J.A. (2023) Single-cell analysis of shared
signatures and transcriptional diversity during zebrafish development Developmental
Biology.

9. Lange,M., Granados,A., VijayKumar,S., Bragantini,J., Ancheta,S., Santhosh,S., Borja,M.,
Kobayashi,H., McGeever,E., Solak,A.C., *et al.* (2023) Zebrahub – Multimodal Zebrafish
Developmental Atlas Reveals the State-Transition Dynamics of Late-Vertebrate
Pluripotent Axial Progenitors Developmental Biology.

10. Liu,C., Li,R., Li,Y., Lin,X., Zhao,K., Liu,Q., Wang,S., Yang,X., Shi,X., Ma,Y., *et al.* (2022)
Spatiotemporal mapping of gene expression landscapes and developmental
trajectories during zebrafish embryogenesis. *Developmental Cell*, **57**, 1284-1298.e5.

11. Spitz,F. and Furlong,E.E.M. (2012) Transcription factors: from enhancer binding to
developmental control. *Nat Rev Genet*, **13**, 613–626.

12. Liu,G., Wang,W., Hu,S., Wang,X. and Zhang,Y. (2018) Inherited DNA methylation primes
the establishment of accessible chromatin during genome activation. *Genome
Research*, **28**, 998–1007.

13. Veil,M., Yampolsky,L.Y., Grüning,B. and Onichtchouk,D. (2019) Pou5f3, SoxB1, and Nanog
remodel chromatin on high nucleosome affinity regions at zygotic genome activation.
Genome Res., **29**, 383–395.

14. Pálfy,M., Schulze,G., Valen,E. and Vastenhouw,N.L. (2020) Chromatin accessibility
established by Pou5f3, Sox19b and Nanog primes genes for activity during zebrafish
genome activation. *PLOS Genetics*, **16**, e1008546.

15. Yang,H., Luan,Y., Liu,T., Lee,H.J., Fang,L., Wang,Y., Wang,X., Zhang,B., Jin,Q., Ang,K.C., *et
al.* (2020) A map of cis-regulatory elements and 3D genome structures in zebrafish.
Nature, **588**, 337–343.

16. McGarvey,A.C., Kopp,W., Vučićević,D., Mattonet,K., Kempfer,R., Hirsekorn,A., Bilić,I.,
Gil,M., Trinks,A., Merks,A.M., *et al.* (2022) Single-cell-resolved dynamics of chromatin
architecture delineate cell and regulatory states in zebrafish embryos. *Cell Genomics*,
2, 100083.

17. Lin,X., Yang,X., Chen,C., Ma,W., Wang,Y., Li,X., Zhao,K., Deng,Q., Feng,W., Ma,Y., *et al.*
(2023) Single-nucleus chromatin landscapes during zebrafish early embryogenesis.
Sci Data, **10**, 464.

18. Chen,X., Miragaia,R.J., Natarajan,K.N. and Teichmann,S.A. (2018) A rapid and robust
method for single cell chromatin accessibility profiling. *Nat Commun*, **9**, 5345.

19. Xu,W., Wen,Y., Liang,Y., Xu,Q., Wang,X., Jin,W. and Chen,X. (2021) A plate-based single-cell
ATAC-seq workflow for fast and robust profiling of chromatin accessibility. *Nat Protoc*,
16, 4084–4107.

20. Köster,J. and Rahmann,S. (2012) Snakemake—a scalable bioinformatics workflow engine.
Bioinformatics, **28**, 2520–2522.

21. Chen,S., Zhou,Y., Chen,Y. and Gu,J. (2018) fastp: an ultra-fast all-in-one FASTQ
preprocessor. *Bioinformatics*, **34**, i884–i890.

22. Kim,D., Paggi,J.M., Park,C., Bennett,C. and Salzberg,S.L. (2019) Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology*,
37, 907–915.

23. Li,H., Handsaker,B., Wysoker,A., Fennell,T., Ruan,J., Homer,N., Marth,G., Abecasis,G.,
Durbin,R. and Subgroup,1000 Genome Project Data Processing (2009) The Sequence
Alignment/Map format and SAMtools. *Bioinformatics*, **25**, 2078–2079.

24. Zhang,Y., Liu,T., Meyer,C.A., Eeckhoute,J., Johnson,D.S., Bernstein,B.E., Nusbaum,C.,
Myers,R.M., Brown,M., Li,W., *et al.* (2008) Model-based Analysis of ChIP-Seq (MACS).
Genome Biology, **9**, R137.

25. Quinlan,A.R. and Hall,I.M. (2010) BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics*, **26**, 841–842.

26. Stuart,T., Srivastava,A., Madad,S., Lareau,C.A. and Satija,R. (2021) Single-cell chromatin
state analysis with Signac. *Nature Methods*, 10.1038/s41592-021-01282-5.

27. Lawrence,M., Huber,W., Gentleman,R., Morgan,M.T. and Carey,V.J. (2013) Software for
Computing and Annotating Genomic Ranges. *PLOS Computational Biology*, **9**.

28. Polański,K., Young,M.D., Miao,Z., Meyer,K.B., Teichmann,S.A. and Park,J.-E. (2020)
BBKNN: fast batch alignment of single cell transcriptomes. *Bioinformatics*, **36**, 964–
965.

29. Wolf,F.A., Angerer,P. and Theis,F.J. (2018) SCANPY: large-scale single-cell gene expression
data analysis. *Genome Biology*, **19**, 15.

30. Pedregosa,F., Varoquaux,G., Gramfort,A., Michel,V., Thirion,B., Grisel,O., Blondel,M.,
Prettenhofer,P., Weiss,R., Dubourg,V., *et al.* (2011) Scikit-learn: Machine Learning in
Python. *Journal of Machine Learning Research*, **12**, 2825–2830.

31. Heinz,S., Benner,C., Spann,N., Bertolino,E., Lin,Y.C., Laslo,P., Cheng,J.X., Murre,C., Singh,H.
and Glass,C.K. (2010) Simple Combinations of Lineage-Determining Transcription
Factors Prime cis-Regulatory Elements Required for Macrophage and B Cell Identities.
Molecular Cell, **38**, 576–589.

32. Castro-Mondragon,J.A., Riudavets-Puig,R., Rauluseviciute,I., Berhanu Lemma,R.,
Turchi,L., Blanc-Mathieu,R., Lucas,J., Boddie,P., Khan,A., Manosalva Pérez,N., *et al.*
(2022) JASPAR 2022: the 9th release of the open-access database of transcription
factor binding profiles. *Nucleic Acids Research*, **50**, D165–D173.

33. Schep,A.N., Wu,B., Buenrostro,J.D. and Greenleaf,W.J. (2017) chromVAR: inferring
transcription-factor-associated accessibility from single-cell epigenomic data. *Nat*
Methods, **14**, 975–978.

34. Ramírez,F., Ryan,D.P., Grüning,B., Bhardwaj,V., Kilpert,F., Richter,A.S., Heyne,S., Dündar,F.
and Manke,T. (2016) deepTools2: a next generation web server for deep-sequencing
data analysis. *Nucleic Acids Res*, **44**, W160–W165.

35. Pliner,H.A., Packer,J.S., McFaline-Figueroa,J.L., Cusanovich,D.A., Daza,R.M.,
Aghamirzaie,D., Srivatsan,S., Qiu,X., Jackson,D., Minkina,A., *et al.* (2018) Cicero
Predicts cis-Regulatory DNA Interactions from Single-Cell Chromatin Accessibility
Data. *Molecular Cell*, **71**, 858-871.e8.

36. Haghverdi,L., Büttner,M., Wolf,F.A., Buettner,F. and Theis,F.J. (2016) Diffusion pseudotime
robustly reconstructs lineage branching. *Nat Methods*, **13**, 845–848.

37. Yee,T.W. (2015) Vector Generalized Linear and Additive Models: With an Implementation
in R Springer New York, New York, NY.

38. Cusanovich,D.A., Daza,R., Adey,A., Pliner,H.A., Christiansen,L., Gunderson,K.L.,
Steemers,F.J., Trapnell,C. and Shendure,J. (2015) Multiplex single-cell profiling of
chromatin accessibility by combinatorial cellular indexing. *Science*, **348**, 910–914.

39. Becht,E., McInnes,L., Healy,J., Dutertre,C.-A., Kwok,I.W.H., Ng,L.G., Ginhoux,F. and
Newell,E.W. (2019) Dimensionality reduction for visualizing single-cell data using
UMAP. *Nature Biotechnology*, **37**, 38–44.

40. Cichocki,A. and Phan,A.-H. (2009) Fast Local Algorithms for Large Scale Nonnegative
Matrix and Tensor Factorizations. *IEICE Trans. Fundamentals*, **E92-A**, 708–721.

41. Févotte,C. and Idier,J. (2011) Algorithms for Nonnegative Matrix Factorization with the β-
Divergence. *Neural Computation*, **23**, 2421–2456.

42. Wu,J., Huang,B., Chen,H., Yin,Q., Liu,Y., Xiang,Y., Zhang,B., Liu,B., Wang,Q., Xia,W., *et al.*
(2016) The landscape of accessible chromatin in mammalian preimplantation
embryos. *Nature*, **534**, 652–657.

43. Gao,L., Wu,K., Liu,Z., Yao,X., Yuan,S., Tao,W., Yi,L., Yu,G., Hou,Z., Fan,D., *et al.* (2018)
Chromatin Accessibility Landscape in Human Early Embryos and Its Association with
Evolution. *Cell*, **173**, 248-259.e15.

44. Lu,F., Liu,Y., Inoue,A., Suzuki,T., Zhao,K. and Zhang,Y. (2016) Establishing Chromatin
Regulatory Landscape during Mouse Preimplantation Development. *Cell*, **165**, 1375–
1388.

45. Wu,J., Xu,J., Liu,B., Yao,G., Wang,P., Lin,Z., Huang,B., Wang,X., Li,T., Shi,S., *et al.* (2018)
Chromatin analysis in human early development reveals epigenetic transition during
ZGA. *Nature*, **557**, 256–260.

46. Gentsch,G.E., Spruce,T., Owens,N.D.L. and Smith,J.C. (2019) Maternal pluripotency
factors initiate extensive chromatin remodelling to predefine first response to
inductive signals. *Nat Commun*, **10**, 4269.

47. Bright,A.R., Van Genesen,S., Li,Q., Grasso,A., Frölich,S., Van Der Sande,M., Van
Heeringen,S.J. and Veenstra,G.J.C. (2021) Combinatorial transcription factor activities
on open chromatin induce embryonic heterogeneity in vertebrates. *The EMBO*
Journal, **40**, e104913.

48. Murphy,P.J., Wu,S.F., James,C.R., Wike,C.L. and Cairns,B.R. (2018) Placeholder
Nucleosomes Underlie Germline-to-Embryo DNA Methylation Reprogramming. *Cell*,
172, 993-1006.e13.

49. Zhang,B., Wu,X., Zhang,W., Shen,W., Sun,Q., Liu,K., Zhang,Y., Wang,Q., Li,Y., Meng,A., *et*
al. (2018) Widespread Enhancer Dememorization and Promoter Priming during
Parental-to-Zygotic Transition. *Molecular Cell*, **72**, 673-686.e6.

50. Bogdanović,O., Fernandez-Miñán,A., Tena,J.J., Calle-Mustienes,E. de la, Hidalgo,C.,
Kruysbergen,I. van, Heeringen,S.J. van, Veenstra,G.J.C. and Gómez-Skarmeta,J.L.
(2012) Dynamics of enhancer chromatin signatures mark the transition from
pluripotency to cell specification during embryogenesis. *Genome Research*, **22**, 2043–
2053.

51. Ernst,J. and Kellis,M. (2010) Discovery and characterization of chromatin states for
systematic annotation of the human genome. *Nature Biotechnology*, **28**, 817–825.

52. McLean,C.Y., Bristor,D., Hiller,M., Clarke,S.L., Schaar,B.T., Lowe,C.B., Wenger,A.M. and
Bejerano,G. (2010) GREAT improves functional interpretation of cis-regulatory
regions. *Nature Biotechnology*, **28**, 495–501.

53. Chang,N.-C., Rovira,Q., Wells,J., Feschotte,C. and Vaquerizas,J.M. (2022) Zebrafish
transposable elements show extensive diversification in age, genomic distribution,
and developmental expression. *Genome Res.*, **32**, 1408–1423.

54. White,R.J., Collins,J.E., Sealy,I.M., Wali,N., Dooley,C.M., Digby,Z., Stemple,D.L.,
Murphy,D.N., Billis,K., Hourlier,T., *et al.* (2017) A high-resolution mRNA expression
time course of embryonic development in zebrafish. *eLife*, **6**, e30860.

55. Christians,E., Davis,A.A., Thomas,S.D. and Benjamin,I.J. (2000) Maternal effect of Hsf1 on
reproductive success. *Nature*, **407**, 693–694.

56. Ross,S.E., Vázquez-Marín,J., Gert,K.R.B., González-Rajal,Á., Dinger,M.E., Pauli,A.,
Martínez-Morales,J.R. and Bogdanovic,O. (2023) Evolutionary conservation of
embryonic DNA methylome remodelling in distantly related teleost species. *Nucleic
Acids Research*, **51**, 9658–9671.

57. Kimmel,C.B., Ballard,W.W., Kimmel,S.R., Ullmann,B. and Schilling,T.F. (1995) Stages of
embryonic development of the zebrafish. *Dev. Dyn.*, **203**, 253–310.

58. Carvalho,L. and Heisenberg,C.-P. (2010) The yolk syncytial layer in early zebrafish
development. *Trends in Cell Biology*, **20**, 586–592.

59. Xiao,Q., Xia,J.-H., Zhang,X.-J., Li,Z., Wang,Y., Zhou,L. and Gui,J.-F. (2014) Type-IV Antifreeze
Proteins are Essential for Epiboly and Convergence in Gastrulation of Zebrafish
Embryos. *Int. J. Biol. Sci.*, **10**, 715–732.

60. Hsu,H.-J., Hsiao,P., Kuo,M.-W. and Chung,B. (2002) Expression of zebrafish cyp11a1 as a
maternal transcript and in yolk syncytial layer. *Gene Expression Patterns*, **2**, 219–222.

61. Berger,M.F., Badis,G., Gehrke,A.R., Talukder,S., Philippakis,A.A., Peña-Castillo,L.,
Alleyne,T.M., Mnaimneh,S., Botvinnik,O.B., Chan,E.T., *et al.* (2008) Variation in
Homeodomain DNA Binding Revealed by High-Resolution Analysis of Sequence
Preferences. *Cell*, **133**, 1266–1276.

62. Badis,G., Berger,M.F., Philippakis,A.A., Talukder,S., Gehrke,A.R., Jaeger,S.A., Chan,E.T.,
Metzler,G., Vedenko,A., Chen,X., *et al.* (2009) Diversity and Complexity in DNA
Recognition by Transcription Factors. *Science*, **324**, 1720–1723.

63. Newburger,D.E. and Bulyk,M.L. (2009) UniPROBE: an online database of protein binding
microarray data on protein-DNA interactions. *Nucleic Acids Research*, **37**, D77–D82.

64. Noyes,M.B., Christensen,R.G., Wakabayashi,A., Stormo,G.D., Brodsky,M.H. and
Wolfe,S.A. (2008) Analysis of Homeodomain Specificities Allows the Family-wide
Prediction of Preferred Recognition Sites. *Cell*, **133**, 1277–1289.

65. Liberzon,A. (2004) Role of intrinsic DNA binding specificity in defining target genes of the
mammalian transcription factor PDX1. *Nucleic Acids Research*, **32**, 54–64.

66. Weirauch,M.T., Yang,A., Albu,M., Cote,A.G., Montenegro-Montero,A., Drewe,P.,
Najafabadi,H.S., Lambert,S.A., Mann,I., Cook,K., *et al.* (2014) Determination and
Inference of Eukaryotic Transcription Factor Sequence Specificity. *Cell*, **158**, 1431–
1443.

67. Yamanaka,Y., Mizuno,T., Sasai,Y., Kishi,M., Takeda,H., Kim,C.-H., Hibi,M. and Hirano,T.
(1998) A novel homeobox gene, *dharma,* can induce the organizer in a non-cell-
autonomous manner. *Genes Dev.*, **12**, 2345–2353.

68. Shen,W.-K., Chen,S.-Y., Gan,Z.-Q., Zhang,Y.-Z., Yue,T., Chen,M.-M., Xue,Y., Hu,H. and
Guo,A.-Y. (2023) AnimalTFDB 4.0: a comprehensive animal transcription factor
database updated with variation and expression annotations. *Nucleic Acids Research*,
51, D39–D45.

69. Kotkamp,K., Mössner,R., Allen,A., Onichtchouk,D. and Driever,W. (2014) A Pou5f1/Oct4
dependent Klf2a, Klf2b, and Klf17 regulatory sub-network contributes to EVL and
ectoderm development during zebrafish embryogenesis. *Developmental Biology*,
385, 433–447.

70. Miles,L.B., Darido,C., Kaslin,J., Heath,J.K., Jane,S.M. and Dworkin,S. (2017) Mis-
expression of grainyhead-like transcription factors in zebrafish leads to defects in
enveloping layer (EVL) integrity, cellular morphogenesis and axial extension. *Sci Rep*,
7, 17607.

71. Xu,C., Fan,Z.P., Müller,P., Fogley,R., DiBiase,A., Trompouki,E., Unternaehrer,J., Xiong,F.,
Torregroza,I., Evans,T., *et al.* (2012) Nanog-like Regulates Endoderm Formation
through the Mxtx2-Nodal Pathway. *Developmental Cell*, **22**, 625–638.

72. Chen,X., Teichmann,S.A. and Meyer,K.B. (2018) From Tissues to Cell Types and Back:
Single-Cell Gene Expression Analysis of Tissue Architecture. *Annu. Rev. Biomed. Data
Sci.*, **1**, 29–51.

73. Preissl,S., Gaulton,K.J. and Ren,B. (2023) Characterizing cis-regulatory elements using
single-cell epigenomics. *Nat Rev Genet*, **24**, 21–43.

74. Grosselin,K., Durand,A., Marsolier,J., Poitou,A., Marangoni,E., Nemati,F., Dahmani,A.,
Lameiras,S., Reyal,F., Frenoy,O., *et al.* (2019) High-throughput single-cell ChIP-seq
identifies heterogeneity of chromatin states in breast cancer. *Nat Genet*, **51**, 1060–
1066.

75. Wang,Q., Xiong,H., Ai,S., Yu,X., Liu,Y., Zhang,J. and He,A. (2019) CoBATCH for High-
Throughput Single-Cell Epigenomic Profiling. *Molecular Cell*, **76**, 206-216.e7.

76. Ai,S., Xiong,H., Li,C.C., Luo,Y., Shi,Q., Liu,Y., Yu,X., Li,C. and He,A. (2019) Profiling
chromatin states using single-cell itChIP-seq. *Nature Cell Biology*, **21**, 1164–1172.

77. Xiong,H., Luo,Y., Wang,Q., Yu,X. and He,A. (2021) Single-cell joint detection of chromatin
occupancy and transcriptome enables higher-dimensional epigenomic
reconstructions. *Nature Methods*, **18**, 652–660.

78. Zhu,C., Zhang,Y., Li,Y.E., Lucero,J., Behrens,M.M. and Ren,B. (2021) Joint profiling of
histone modifications and transcriptome in single cells from mouse brain. *Nature
Methods*, 10.1038/s41592-021-01060-3.

79. Wu,S.J., Furlan,S.N., Mihalas,A.B., Kaya-Okur,H.S., Feroze,A.H., Emerson,S.N., Zheng,Y.,
Carson,K., Cimino,P.J., Keene,C.D., *et al.* (2021) Single-cell CUT&Tag analysis of
chromatin modifications in differentiation and tumor progression. *Nature
Biotechnology*, 10.1038/s41587-021-00865-z.

80. Bartosovic,M., Kabbe,M. and Castelo-Branco,G. (2021) Single-cell CUT&Tag profiles
histone modifications and transcription factors in complex tissues. *Nature
Biotechnology*, 10.1038/s41587-021-00869-9.

81. Rang,F.J., De Luca,K.L., De Vries,S.S., Valdes-Quezada,C., Boele,E., Nguyen,P.D.,
Guerreiro,I., Sato,Y., Kimura,H., Bakkers,J., *et al.* (2022) Single-cell profiling of
transcriptome and histone modifications with EpiDamID. *Molecular Cell*, **82**, 1956-
1970.e14.

82. Ku,W.L., Nakamura,K., Gao,W., Cui,K., Hu,G., Tang,Q., Ni,B. and Zhao,K. (2019) Single-cell
chromatin immunocleavage sequencing (scChIC-seq) to profile histone modification.
Nat Methods, **16**, 323–325.

83. Yeung,J., Florescu,M., Zeller,P., de Barbanson,B.A., Wellenstein,M.D. and van
Oudenaarden,A. (2023) scChIX-seq infers dynamic relationships between histone
modifications in single cells. *Nat Biotechnol*, 10.1038/s41587-022-01560-3.

84. Ho,R. and Kimmel,C. (1993) Commitment of cell fate in the early zebrafish embryo.
Science, **261**, 109–111.

Figure Captions

Figure 1 | Single-nucleus chromatin accessibility maps of zebrafish early
embryogenesis.
878
879   **Figure 2 | The dynamics of open chromatin peaks during early development.**
880
881   **Figure 3 | Distinct genomic features are associated with peaks from different peak**
882   **modules.**
883
884   **Figure 4 | Emerging different chromatin profiles among single cells when entering**
885   **the dome stage.**
886
887   **Figure 5 | Putative enhancers located in the marker peaks of EVL and YSL cells.**
888
889   **Figure 6 | Co-accessibility analysis links putative enhancers to their target genes.**
890
891

Figure 1 | Single-nucleus chromatin accessibility maps of zebrafish early embryogenesis. (**A**) Schematic view of the overall experimental workflow. (**B**) UCSC genome browser tracks of a genomic locus for all time points. The aggregated single-nucleus ATAC-seq profiles per time point was shown together with a previously published bulk ATAC-seq data around similar stages. (**C**) 2D UMAP visualisation of the single-nucleus ATAC-seq data per time point.

Figure 2 | The dynamics of open chromatin peaks during early development. (**A**) The BBKNN integrated UMAP of all single-nucleus ATAC-seq profiles coloured by time points. (**B**) Schematic view of the non-negative matrix factorisation (NMF) workflow to identify peak modules that covary across nucleus in the whole data. Only reproducible peak modules were used for the downstream analysis. Batch-related peak modules were ignored. (**C**) The embedding values of peak modules 2, 3, 4, 6 and 10 in each single nucleus shown as the violin plot (top) and on UMAP (bottom). (**D**) Browser tracks of the single-nucleus aggregate of ATAC-seq signals from typical examples of each peak module in (**C**).

Figure 3 | Distinct genomic features are associated with peaks from different peak modules. (**A**) ChromHMM analysis of the zebrafish genome using previously published ChIP-seq data sets. Peaks from peak modules 2, 3, 4, 6 and 10 were also compared to the chromatin states defined by the combinations of histone marks or variants. (**B**) Genome browser tracks showing two examples for peaks from peak module 10. (**C**) The distribution of the number of peaks in peak module 10 that overlapped the indicated transposable element class. Random was the background, which contained 2,000 randomly selected peaks from all ATAC-seq peaks. (**D**) HOMER *de novo* motif results from each peak module. The percentage of peaks containing the motif and the p-values were shown. The best match to known transcription factor motifs was shown at the right-hand side.

Figure 4 | Emerging different chromatin profiles among single cells when entering the dome stage. (**A**) UMAP visualisation of nucleus from the 4.1 hpf (sphere-dome) and the 4.3 hpf (dome) stages coloured by different clusters. Cluster annotations were indicated on the UMAP plot. A schematic view of the structure of the embryo was also shown at the top left. (**B**) The same UMAP plot as in (**A**), coloured by the normalised reads from the indicated gene locus. (**C**) Genome browser tracks showing the aggregated single-nucleus ATAC-seq signal from indicated cell clusters. Five examples were shown around the mark genes of EVL and YSL cells. (**D**) Diffusion pseudotime analysis of the nuclei from the 4.1 hpf (sphere-dome) and the 4.3 hpf (dome) stages, coloured by the pseudotime (left) or cell identity (right). (**E**) Heatmap representation of gene activity changes along three different branches. Important marker genes for EVL and YSL cells were indicated.

Figure 5 | Putative enhancers located in the marker peaks of EVL and YSL cells. (**A**) The comparison of the EVL and YSL marker peaks to the eight chromatin states described in **Figure 3A**. (**B**) Heatmap representation of the aggregated of scATAC-seq reads (left) or public ChIP-seq reads of indicated histone marks (right) around EVL and YSL marker peaks. Black arrows indicate the centre of the peak. 5 kb upstream and downstream of the peak were plotted. (**C**) HOMER *de novo* motif discovery results of the marker peaks of EVL and YSL cells. (**D**) The same UMAP plot as in **Figure 4A**, coloured by the deviation scores of indicated motifs calculated by chromVAR. (**E**) The mean expression levels of indicated genes in each stage from a previously published scRNA-seq atlas. All members from the CP2, the IRF, the KLF and the SP family were investigated. (**F**) Heatmap representation showing the gene activity scores of all homeobox transcription factors measured by scATAC-seq data in different cell clusters. Red box indicated the 73 homeodomain factors that have higher gene activity scores in the YSL cluster.

Figure 6 | Co-accessibility analysis links putative enhancers to their target genes. Examples of co-accessibility of peaks around EVL and YSL marker genes. The linkage strength analysed by Cicero was shown at the bottom. Cell-type specific open chromatin peaks with motifs from **Figure 5C** were highlighted. ChIP-seq data of H3K4me3, H3K4me3 and H3K27ac from two previous studies were also shown.

Supplementary Figure 1 | Quality control (QC) of the whole single-nucleus ATAC-seq experiment. (**A**) Examples of FACS for a typical experiment. FSC-A and SSC-A were used to remove cell debris. FSC-A and FSC-W were used to select only single nucleus. DAPI+ events were used to select only successfully permeabilised cells. (**B**) Median information of each QC metric in different stages. (**C**) QC cutoff used to remove failed wells in the experiment. (**D**) UCSC genome browser tracks from the indicated genomic locus showing an overview of the data quality.

Supplementary Figure 2 | The overview of the entire single nucleus ATAC-seq data set. (**A**) BBKNN integrated UMAP of all nucleus, coloured by experimental batches. (**B**) The module weights of each peak module grouped by experimental batches. (**C**) The global TSS enrichment scores of each time point.

Supplementary Figure 3 | Different genomic features and functions of each peak module. (**A**) The genomic distribution relative to annotated genes of each peak module. (**B**) GREAT analysis of enriched biological processes in each peak module. (**C**) The distribution of the number of peaks in peak module 10 that overlapped the indicated LTR family. Random was the background, which contained 2,000 randomly selected peaks from all ATAC-seq peaks. (**D**) The expressions of 519 LTRs in module 10 at 1k and dome stages. Data were taken from White *et al.* 2017 (54). (**E**) The DNA methylation levels (mCG/CG) at 3 hpf (1k stage) around indicated regions. 1 kb upstream and downstream were also plotted. Data were taken from Ross *et al.* 2023 (56).

Supplementary Figure 4 | Analysis of nucleus from 4.1 hpf (sphere-dome) and 4.3 hpf (dome) stages. (**A**) The same UMAP plot as in **Figure 4A**, coloured by time point (top left), experimental batches (top right) and module weights (bottom). (**B**) Browser tracks showing specific peaks around three EVL marker genes. (**C**) The expression levels of indicated genes from a previously published scRNA-seq atlas.

Supplementary Figure 5 | Examples of peak co-accessibility around two YSL marker genes. Cell-type specific open chromatin peaks with motifs from **Figure 5C** were highlighted. ChIP-seq data of H3K4me3, H3K4me3 and H3K27ac from two previous studies were also shown.

Supplementary Figure 6 | Browser tracks showing the aggregated single-nucleus ATAC-seq signal in each cell cluster around the three pluripotent genes.