# Cortical and behavioural tracking of rhythm in music: Effects of pitch predictability, enjoyment, and expertise

Anne Keitel*[1], Claire Pelofi[2,3], Xinyi Guan[3,4], Emily Watson[1], Lucy Wight[1,5], Sarah Allen[1], Iris Mencke[6,7], Christian Keitel[+1], Johanna Rimmele[+3,8]

[1]Psychology, University of Dundee, UK; [2]Department of Psychology, New York University, US; [3]Max Planck NYU Center for Language, Music, and Emotion, Frankfurt am Main, Germany, New York, NY, US; [4]Digital and Cognitive Musicology Lab, École Polytechnique Fédérale de Lausanne, Lausanne, Switzerland; [5]School of Psychology, Aston University, Birmingham, UK; [6]Department of Medical Physics and Acoustics, University of Oldenburg, Oldenburg, Germany; [7]Department of Music, Max-Planck-Institute for Empirical Aesthetics, Frankfurt, Germany; [8]Department of Cognitive Neuropsychology, Max-Planck-Institute for Empirical Aesthetics, Frankfurt, Germany

* Corresponding author:
Dr Anne Keitel
Psychology Division; School of Humanities, Social Sciences, and Law
University of Dundee
DD1 4HN
Dundee, UK

[+] indicates equal contributions from the two senior authors.

**Abbreviated title**: Pitch predictability, enjoyment and expertise affect rhythm tracking.

**Key words**: music perception, EEG, naturalistic music, pitch surprisal, musical expertise, top-down influences

This is a revised manuscript. An overview of the changes can be found here:
https://osf.io/wvsjr.

## Abstract

The cortical tracking of stimulus features (such as the sound envelope) is a crucial neural requisite of how we process continuous music. We here tested whether cortical tracking of the beat, typically related to rhythm processing, is modulated by pitch predictability and other top-down factors. Participants listened to tonal (high pitch predictability) and atonal (low pitch predictability) music while undergoing EEG, and we analysed their cortical tracking of the acoustic envelope. Interestingly, cortical envelope tracking was stronger while listening to atonal than tonal music, likely reflecting listeners' violated pitch expectations. Envelope tracking was also stronger with more expertise and enjoyment. Furthermore, we analysed cortical tracking of pitch surprisal (using IDyOM) and show that listeners' expectations for tonal and atonal music match those computed by the IDyOM model, with higher surprisal (prediction errors) for atonal than tonal music. Behaviourally, we measured participants' ability to tap along to the beat of tonal and atonal sequences in two experiments. In both experiments, finger-tapping performance was better in the tonal than the atonal condition, indicating a positive effect of pitch predictability on behavioural rhythm processing. Cortical envelope tracking predicted tapping performance for tonal music, as did pitch surprisal tracking for atonal music, indicating that conditions of high and low predictability might impose different processing regimes. We show that cortical envelope tracking, beyond reflecting musical rhythm processing, is modulated by pitch predictability, as well as musical expertise and enjoyment. Taken together, our results show various ways in which those top-down factors impact musical rhythm processing.

## Introduction

The cortical tracking of continuous auditory stimuli, such as music and speech, has been the topic of intense investigation in the past years (Keitel *et al.*, 2018; Peelle *et al.*, 2013; Tierney & Kraus, 2015). Cortical tracking usually refers to the neural signal matching slow amplitude fluctuations in the acoustic signal and is quantified by neural alignment to the stimulus envelope, thought to reflect processing of the rhythmic structure (Doelling & Poeppel, 2015; Gross *et al.*, 2013; Luo & Poeppel, 2007). Although mostly investigated in speech, recent findings suggest that processing of naturalistic music might rely on comparable mechanisms (Harding *et al.*, 2019; Sammler, 2020; Zuk *et al.*, 2021). Cortical tracking is influenced by numerous top-down factors, but their interaction and relative importance are poorly understood. For example, increased attention and listening effort generally leads to stronger speech tracking (Ding & Simon, 2012; Lesenfants & Francart, 2020; Rimmele *et al.*, 2015; Song & Iverson, 2018; Zion Golumbic *et al.*, 2013). Conversely, both speech and music tracking are enhanced with language and music proficiency and prior knowledge (Blanco-Elorrieta *et al.*, 2020; Cervantes Constantino & Simon, 2018; Di Liberto *et al.*, 2018; Doelling & Poeppel, 2015; Harding *et al.*, 2019). Other factors, such as the influence of enjoyment on the cortical tracking of music, have also recently elicited researchers' interest (Weineck *et al.*, 2022). Overall, any study of cortical tracking of rhythmic stimuli, needs to take into account stimuli and listener characteristics, which is one driver of the present study.

Recent studies on the cortical tracking of music have shown that the auditory cortex tracks not only the acoustic envelope but also melodic expectations, modelled as surprisal values (Abrams *et al.*, 2022; Di Liberto *et al.*, 2020a; Kern *et al.*, 2022; Marion *et al.*, 2021). These studies suggest that humans automatically process melodic expectations while listening to naturalistic, continuous stimuli (Pearce *et al.*, 2010). Here, we examine the cortical tracking of pitch surprisal, using music stimuli with different levels of pitch predictability, namely tonal and atonal music excerpts. Music that is composed according to (Western) tonal principles, has an intrinsic hierarchical pitch organisation (Lerdahl, 2019). Therefore, this compositional style results in far more predictable pitch sequences than atonal music (Mencke *et al.*, 2018), which is based on the compositional principle that all twelve tones within an octave are equiprobable. The few studies that have been conducted using atonal music show that the resulting lack of a hierarchical pitch organisation negatively affects memorisation (Schulze *et al.*, 2011), recognition (Cuddy *et al.*, 1981; Dibben, 1994; Dowling *et al.*, 1995) and the strength of melodic expectations (Ockelford & Sergeant, 2013) (for review, Mencke *et al.*, 2022; Mencke *et al.*, 2018; Vuvan *et al.*, 2014). Electrophysiological research suggests that weaker expectancies in atonal

music particularly affect later attention-related processing stages (Mencke *et al.*, 2021; Neuloh

& Curio, 2004). Taken together, atonal music seems to present specific perceptual challenges to

listeners, in particular related to melodic expectations.

In the context of musical rhythm perception, finger-tapping is often used as 'behavioural

tracking' measure (Harding *et al.*, 2019; Sammler, 2020) to assess rhythm skills (Fiveash *et al.*,

2022; Iversen *et al.*, 2015; Repp, 2005). The present study addresses the little-known relationship

between behavioural tracking (measured by finger-tapping) and cortical envelope tracking

(measured by electroencephalogram [EEG] recordings of listening participants) of musical

rhythm in the context of varying pitch predictability. While temporal predictability has been

shown to increase pitch discrimination performance (Herbst & Obleser, 2019; Jones *et al.*, 2002),

it is currently unclear whether pitch predictability affects the ability to behaviourally track

naturalistic musical rhythms.

In sum, we here investigated whether cortical tracking of the music envelope, beyond rhythm

processing, is modulated by pitch surprisal in two continuous, naturalistic stimulus conditions:

tonal (high pitch predictability) and atonal (low pitch predictability) music. In the main

experiment, we analysed participants' EEG during passive listening, focusing on cortical

envelope and surprisal tracking. We also investigated the role of enjoyment and musical

expertise for cortical envelope tracking. In both the main and follow-up replication experiment,

we used a behavioural measure of rhythm perception (finger-tapping) to analyse whether cortical

tracking is behaviourally relevant and whether pitch predictability influences behavioural

tracking. We expected that high pitch predictability in the tonal condition would be associated

with better behavioural rhythm tracking than low pitch predictability in the atonal condition (see

pre-registration: https://osf.io/qctpj). Due to complex and opposing effects of attention and

previous experience (e.g., Reetzke *et al.*, 2021) we did not hypothesise a priori on whether cortical

envelope tracking would be stronger in the tonal or atonal condition.

## Materials and methods

*Participants*

Twenty volunteers participated in the main study (14 female, 6 male; 18 to 26 years old;

$M = 20.95$, $SD = 1.88$). It was initially planned to test 24 participants (preregistration:

https://osf.io/qctpj), but data collection had to be halted due to the COVID-19 pandemic.

However, the sample size analysis was based on a previous study (Schulze *et al.*, 2011); $d = .64$,

$\alpha = .05$, $\beta = .80$; see preregistration) and yielded a desired sample size of $N = 21$, which was close

to being achieved. In addition, we tested further 52 participants in a behavioural follow up

experiment (see below). Participants in the main study were right-handed ($N$ = 19) or

ambidextrous (N = 1; Oldfield, 1971). Self-reports (Quick Hearing Check, Koike *et al.*, 1994)

indicated that 19 participants had normal hearing, while one reported a score that might suggest

slightly diminished hearing (score of 27/60, hearing test recommended from score 20). All

participants reported never having received a diagnosis of neurological/psychological disorders

or dyslexia. Participants rated their musical expertise on a scale from 1 to 3 ('none', 'some', 'a

great deal'; $M$ = 1.95, $SD$ = 0.76). Six participants reported no musical expertise. Most

participants ($N$ = 18) were unfamiliar with the musical stimuli, and although two reported

familiarity with the music, they could not name the piece nor composer.

The study was approved by the School of Social Sciences Research Ethics Committee at the

University of Dundee (approval number: UoD-SoSS-PSY-UG-2019-84) and adhered to the

guidelines for the treatment of human participants in the Declaration of Helsinki. Participants

were reimbursed for their time with £15.

*Musical stimuli*

Tonal and atonal polyphonic piano stimuli were used (see **Figure 1** top). For the tonal condition,

we used an excerpt from W. A. Mozart's *'Sonata No. 5 in G Major, K. 283'*. The excerpt was taken

from the second movement (*II. Andante*). The atonal piece was a manipulated version of this

excerpt, created by randomly shifting the pitch of each note from 1 to 9 semitones up or down

the musical scale (using GuitarPro v7.5) corresponding to 100 – 900 cents. Therefore, notes no

longer formed harmonic relationships, while the timing of each note remained the same. Overall,

the music in both conditions contained identical timbre, velocity, and rhythm. Each excerpt was

approximately five-minutes long (292 seconds) and had a standard 4/4 time signature. The tempo

of the pieces was 46 beats per minute (bpm), but because eighth note measures were

consistently used, the dominant beat was 92 bpm (**Figure 1**). This equalled a rate of 1.52 Hz (see

modulation spectrum in **Figure 2D**), and the beats were 652 ms apart. For the finger-tapping task,

unique two-bar segments from the same pieces were extracted per condition (18 segments, each

10.4 seconds long). All music pieces were presented at a sampling rate of 44,100 Hz. All stimuli

are available on the OSF server (https://osf.io/3gf6k/).

**Figure 1. Examples of sheet music and waveforms.** *Top:* Sheet music for two bars of the tonal
condition (original music: *Mozart's Sonata No. 5 in G Major, K. 283: II. Andante*) and the same bars in the
atonal condition. *Bottom:* Waveforms of the same bars in the tonal (green) and atonal (purple) condition,
including the music envelope. Grey bars represent the positions of the dominant beat with an inter-beat-
interval of 652 ms (1.52 Hz).

*Procedure and task*

Participants performed the EEG experiment in a quiet room. They sat comfortably, approximately

110 cm from a 'Benq' computer screen (22.65 × 13.39 inches; 1920 × 1080 pixels resolution). On-

screen instructions were presented in black, size 30 Consolas font, and displayed against a grey

background. Participants could adjust the volume of the sound to a comfortable level before the

start of the experimental blocks. Musical stimuli were presented using E-Prime 3.0 software

(Psychology Software Tools Inc., 2016), and listened to through high-quality wired headphones

(Creative, Draco HS880). Participants first passively listened to the 5-min tonal and atonal music

excerpts (randomised order). Participants started the music self-paced. A five-second

countdown was shown, before an 'X' appeared at music onset, on which participants fixated

throughout the music listening. After each music piece, participants rated how pleasant they had

found the music. We used a Visual Analog Scale, on which participants could rate their

enjoyment by drawing a vertical line between '*Not pleasant*' and '*Very pleasant*'.

After the passive listening blocks, participants performed a finger-tapping task to measure

behavioural rhythm tracking in the tonal and atonal conditions. 36 unique trials (18 per condition)

were presented in four blocks (two tonal and two atonal) of nine trials each. The order of blocks

and of trials within each block was randomised across participants. Each trial was started self-

paced and began with a visual presentation of the dominant beat (i.e., eighth notes). For this, an

'X' flashed four times at the beat frequency before the music started (see **Figure 2A**). The

dominant beat was presented visually and not acoustically, to not interfere with music

processing. Participants then tapped their right index finger on the outer 'Enter' key of a silent

keyboard to the dominant beat of the music. The length of the music segments (2 bars each)

required 16 finger taps per trial, resulting in approximately 288 taps per condition.

**Figure 2. Behavioural paradigm and stimulus properties. A)** Depiction of the trial time-course for the
behavioural tracking task. Before the music started for each 2-bar trial, the dominant beat frequency
(1.52 Hz) was indicated visually by flashing a fixation cross four times at that frequency. Participants
tapped their finger to the dominant beat of the music once the music started. **B)** Enjoyment ratings for
both 5-min tonal and atonal excerpts by all participants in the main experiment (N = 20). Overall,
participants rated the tonal condition as more pleasant/enjoyable than the atonal condition. **C)**
Modulation spectrum of both tonal and atonal excerpts. Thick lines indicate average values across 2-bar
segments, with shaded areas representing standard error of the mean. Beat/meter related frequencies
are indicated by arrows and dotted lines. **D)** Averaged amplitude values of beat-related frequencies (as
shown in C) for both tonal and atonal excerpts. **E)** Sensory dissonance, assessed via roughness. There
was a small difference in that the tonal condition showed larger roughness values than the atonal
condition.
*Notes:* Distribution plots show individual data points, box plots (including median, interquartile ranges
and minimum/maximum), and kernel density estimates.

*Replication of behavioural results*

To make sure that the behavioural effect found in the main experiment (more consistent finger-

tapping to tonal than atonal excerpts) was robust, we carried out a follow-up replication study.

All experimental procedures were ethically approved by the Ethics Council of the Max Planck

Society (Nr. 2017_12). The number of participants was $N$ = 52 (33 female, 19 male), and their age

ranged between 20 and 41 years ($M$ = 26.6, $SD$ = 5.3 years). Most participants were right-handed

($N$ = 43), some were left-handed ($N$ = 6) or ambidextrous ($N$ = 3). The procedure was identical to

the main experiment, with the exception that 4 bars were used for each trial, thus doubling the

time for finger-tapping per trial. This led to 10 unique tonal and 10 unique atonal trials, each 20.9

seconds long. Each trial required 32 finger taps, resulting in 320 taps per condition. Furthermore,

all tonal and atonal trials were presented in random order (in contrast to tonal and atonal blocks

as in the main experiment).

*Analysis of behavioural data*

The inter-tap-intervals of participants' keyboard taps for each trial were pre-processed in several

ways to clean the data. First, the first two finger taps (i.e., before 981 ms) at the beginning of the

trial were excluded from further analysis, to allow participants to hear two eighth notes to inform

their tapping. Trials with fewer than 50% of expected remaining inter-tap-intervals were excluded

(i.e., 6 necessary inter-tap-intervals in the original experiment, and 15 in the replication

experiment). Inter-tap-intervals of faster than 50 ms (indicating involuntary movements) and

slower than 3000 ms (indicating idling) were removed. Within each participant and condition,

trials with intervals of larger than 3 standard deviations from the mean were also excluded (Abel

*et al.*, 2009; Rovetti *et al.*, 2022; Zarate *et al.*, 2015). At the participant level, our criterion was to

exclude outlier data of larger than 3 standard deviations from the mean per condition ($N$ = 0 in

the original experiment, $N$ = 0 in the replication). During the replication experiment, three

participants misunderstood instructions and consistently tapped to fast 16th notes. These three

participants were excluded, resulting in 49 participants that were included in the final analyses.

Finger-tapping performance per trial, condition, and participant was quantified as the Median

Absolute Deviation (MAD), a robust measure of dispersion  (Leys *et al.*, 2013) that precisely

captures the variability in tapping timing. As the MAD is based on median values, it is less

affected by outliers than measures based on the mean, such as variance or the coefficient of

variation (Arachchige *et al.*, 2022). Enjoyment ratings on the visual analogue scales to both the

tonal and atonal excerpts were analysed on a scale between 0 and 100, in increments of 1 (a.u.,

see **Figure 2B**). We also preregistered analysing participants' tapping accuracy in addition to

their variability (https://osf.io/qctpj) but were unable to carry this out after the experiment had

concluded due to restrictions on access to the laboratory and to the equipment for necessary

latency measurements, as imposed by the start of the COVID-19 pandemic.

*EEG data acquisition and preprocessing*

The electroencephalogram (EEG) was recorded from 32 scalp electrodes, using a BioSemi

ActiveTwo system (sampling rate 512 Hz). Electrodes were placed according to the International

10-20 system. Electrodes with an offset of greater/less than ±20mV were adjusted. Ultimately,

electrode offset was always below an absolute value of 30mV before the experiment began.

Horizontal eye-movements were captured by two electro-oculographic (EOG) electrodes, placed

at the outer canthus of each eye. To capture vertical eye-movements and blinks, a further two

electrodes were positioned above and below the participants' left eye.

Pre-processing of the EEG data was conducted using FieldTrip (Oostenveld *et al.*, 2011) functions

in MATLAB 2021a (MathWorks Inc.). For both 5-minute excerpts used during passive listening, we

cut out epochs of 304 seconds (300 s stimulation time from music onset, plus additional 2-s

leading and trailing windows). Data were initially re-referenced to *Cz* and bandpass filtered

between 0.1 and 100 Hz (3rd order Butterworth filter, forward and reverse). Data were then visually

inspected using summary metrics (maximum value and z-value in each channel), and noisy

channels were removed and interpolated using triangulation. A maximum of 4 channels was

removed per participant (M = 2.47, *SD* = 0.91). Before ICA was conducted to identify blinks and

artifacts, data were re-referenced to average reference (Bertrand et al. 1985). On average,

*M* = 2.16 (*SD* = 0.69) components per participant were removed from the data.

*Music envelope pre-processing*

To analyse the tracking of the music signal, we extracted the wideband music envelope. We first

down-sampled each music excerpt to a sampling rate of 150 Hz (Keitel *et al.*, 2018). Acoustic

waveforms were then filtered into eight frequency bands (between 100 and 8,000 Hz, 3rd order

Butterworth filter, forward and reverse) that are equidistant on the cochlear frequency map

(Smith *et al.*, 2002). The signal in each of these eight frequency bands was *Hilbert*-transformed,

and the magnitude extracted, before they were averaged for the wideband music envelope,

which was used for further analyses.

*Pitch surprisal modelling*

Surprisal during music listening refers to how expected a certain musical event is. Some note

sequences are extremely prevalent across Western classical music, thus creating high

expectations and low surprisal for an audience listening to them. To provide a computational

account of music surprisal in the stimuli used, we rely on a model that learns statistical

regularities of music (Pearce, 2018). Based on a variable-order Markov model, IDyOM
(Information Dynamics Of Music; Pearce, 2005; Pearce & Wiggins, 2006) simulates listeners'
expectations while listening to music by collecting statistical structures of note sequences over
*n*-orders on a training corpus set. Here, the training corpus was composed of a collection of
Western folk songs (a subset of the Essen Folksong Collection containing 953 melodies), so as
to accurately model surprisal for typical Western listeners (Guan *et al.*, 2022; Kern *et al.*, 2022).
Specifically, the long-term component (LTM) of the model collects the sequence statistics over
*n*-orders of the training set while the short-term component (STM) dynamically collects the local
context over n-orders for each testing melody. For each note of the testing melodies, the model
outputs a probability distribution of pitch obtained from merging distributions obtained by the
STM and the LTM (for more details, see Pearce, 2005). By comparing the pitch 'ground truth' to
the probability predicted by the model, a surprisal value is obtained. Formally, the surprisal (or
information content) is the log-negative to the base 2 of the probability of the note. It essentially
represents the expectedness of each note given the STM (e.g., the local context) and LTM (e.g.,
the long-term exposure to a musical style or culture). If surprisal is high, prediction error is also
high, and vice versa (Pearce, 2018). The choice of IDyOM was motivated by numerous empirical
evidence that it can accurately model a listener's internal representation of musical regularities,
both using neural and behavioural data (Di Liberto *et al.*, 2020a; Gold *et al.*, 2019; Kern *et al.*,
2022; Pearce *et al.*, 2010; Pearce & Wiggins, 2012). Since IDyOM in its current development only
takes monophonic MIDI inputs, we reduced the complete score of each excerpt into a
monophonic version that contained the melody and the bass line. The pitch surprisal values for
each note were then used to build a continuous signal, with surprisal values making up the
amplitude for the duration of the respective note. This initial step function was smoothed by
convoluting it with a Gaussian filter (*sigma* = 50). The continuous surprisal signal was created to
have the same sampling rate as the EEG signal (150 Hz).

*Mutual information analysis*
The correspondence between the continuous EEG signal and envelope and surprisal signals (i.e.,
cortical envelope tracking and cortical surprisal tracking) was analysed using a Gaussian Copula
mutual information (MI) framework (Ince *et al.*, 2017; Keitel *et al.*, 2017). In this approach, which
is optimised for neurophysiological data, Gaussian copulas are used to normalise the
continuous, analytical signals (Ince *et al.*, 2017). The first 500 ms of the signals were removed
from analysis, to avoid contamination with strong transient evoked responses at the start of the
music. Mutual information (in bits) between the EEG signal and the music envelope was
computed with both signals filtered at the dominant beat frequency range (0.5 – 2 Hz). A

participant-specific stimulus-brain lag was included for the envelope tracking analyses, which

was based on the individual phase coherence (Harding *et al.*, 2019) peak at auditory electrode

Cz, which was averaged for slow frequencies between 1 and 12 Hz (before the narrow band-pass

filtering described above). Initial coherence values were computed for lags between 40 and

200 ms in steps of 20 ms.

Mutual information between the EEG signal and the pitch surprisal signals was computed with

signals bandpass filtered between 0.1 and 50 Hz. As the analysis sampling rate was 150 Hz,

filtering up to 50 Hz allowed for this frequency to be robustly reflected in the data (1/3 of the

sampling frequency). This wide range was chosen as no clear assumptions about a specific,

narrow-band frequency range could be made and prediction processes have been shown across

multiple frequency bands (Engel *et al.*, 2001; Morillon *et al.*, 2019). The surprisal-tracking

analysis included surprisal values for both the melody and bass lines jointly, capitalising on the

multivariate capabilities of the used MI approach (Ince *et al.*, 2017). Apart from the wider

frequency band, the analysis of surprisal tracking was equivalent to that of envelope tracking,

including the same individual stimulus-brain lags.

Each mutual information value was computed per participant, condition, and electrode. The

results of these analyses will be referred to as cortical (envelope or surprisal) 'tracking', and we

do not make assumptions about the underlying mechanisms (e.g., cortical entrainment) as these

are still debated (Alexandrou *et al.*, 2018; Keitel *et al.*, 2021; Obleser & Kayser, 2019).

*Statistical analyses*

To test the statistical significance of mutual information values for envelope and surprisal

tracking against chance, 3000 permutations were computed per participant, condition, and

electrode. Specifically, to create permuted data, we segmented the continuous

envelope/surprisal signals into 1-s segments and shuffled the segments randomly. This kept the

statistical properties of the signal but destroyed the temporal relationship between the music

and brain signals. MI was then computed between the brain signal and the 3000 shuffled

envelope/surprise signals. The group level mean was then tested against the 95th percentile of

the random group mean distribution, essentially implementing a one-sided randomisation test

at $p < .05$ (Brohl *et al.*, 2022).

For the comparison between the two conditions, *t*-values were computed using the real MI

values, as well as the 3000 MI values from the shuffled data. These real and permuted data were

then compared, again using a cluster-based permutation test, with a critical *t*-value of 2.1, which

represents the critical value of the *Student's t* distribution for 20 participants and a two-tailed
probability of *p* = .05 (Keitel *et al.*, 2018).

Pearson's correlations between cortical tracking and behavioural measures (tapping variability,
musical competency, and enjoyment) were computed between the behavioural measures and
the true MI values, as well as between the behavioural measures and the 3000 permuted MI
values. Before comparing the true *r*-values with the permutation distribution using cluster-based
permutation (as above), Pearson's *r*-values were transformed to be normally distributed using
Fisher's z-transformation (e.g., Gorsuch & Lehmann, 2010). For all cluster-based permutation
analyses, initial clusters were chosen at an alpha level of *p* < .05. As an indicator of effect sizes,
we either report *Cohen's d* for peak electrodes in the case of *t* or *r* values (Brohl & Kayser, 2021;
Lakens, 2013), or summed MI values within each significant cluster ($MI_{sum}$) (Keitel *et al.*, 2018).

To be able to draw evidence-led conclusions about the laterality of our results, we explicitly
tested for hemispheric lateralisation (Keitel *et al.*, 2020; Park & Kayser, 2019). Hemispheric
differences in cortical tracking have been theoretically assumed and occasionally been found
experimentally for the music envelope (Doelling & Poeppel, 2015; Zatorre, 2001). The participant-
specific results (e.g., MI values) were extracted for significant electrodes in one hemisphere and
in corresponding contralateral electrodes. We then averaged these values within each
hemisphere and calculated the between-hemispheres differences with a group-level, *Student's*
*t*-test (two-sided). *P*-values were corrected for multiple comparisons using FDR correction at the
level of 5% (Benjamini & Hochberg, 1995).

All tests are *two-tailed*, except for the comparison of finger-tapping variability between the tonal
and atonal condition. These comparisons are one-tailed, as we had an *a priori* directed
hypothesis that finger-tapping in the atonal condition would be more variable than in the tonal
condition (see preregistration, https://osf.io/qctpj).

## Results

*Differences between tonal and atonal music stimuli*

As intended, the pitch surprisal was overall higher for the atonal than the tonal stimuli (see **Figure
5B**). Pitch surprisal was computed for all notes, separately for melody and bass lines using
IDyOM (Pearce, 2005). The surprisal is estimated by comparing the pitch ground truth of a note
to its predicted value in the model's output distribution. For the melody line, pitch surprisal was
on average *M* = 3.02 (*SD* = 3.28) in the tonal condition, and *M* = 6.96 (*SD* = 3.78) in the atonal
condition (see **Figure  5B**). A *Student's t*-test confirmed that surprisal values were statistically
larger in the atonal than the tonal condition, $t(597) = -25.21$, $p < .001$, *Cohen's d* = -1.03. Likewise,

pitch surprisal in the bass line was higher in the atonal than in the tonal condition (tonal: *M* = 3.26,

*SD* = 2.72; atonal: *M* = 7.38, *SD* = 3.54; *t*(722) = -28.41, *p* < .001, *Cohen's d* = -1.06).

Furthermore, participants were asked to rate how pleasant they found listening to the tonal and

atonal music stimuli (on a scale effectively analysed from 0 to 100 a.u., **Figure 2B**). The

enjoyment ratings indicated that participants found the tonal excerpt more pleasant than the

atonal excerpt (tonal: *M* = 76.18, *SD* = 19.31; atonal: *M* = 42.53, *SD* = 25.99; *t*(19) = 6.47, *p* < .001,

*Cohen's d* = 1.45).

We also computed the modulation spectrum (Ding *et al.*, 2017) for both conditions between 0

and 12 Hz (**Figure 2C).** This showed several peaks at beat-related frequencies (i.e., subharmonic

and harmonics of 1.52 Hz). A comparison of average beat-related frequencies (c.f., Celma-

Miralles & Toro, 2019; Nozaradan *et al.*, 2012) for excerpts across both conditions showed no

statistical difference (tonal: *M* = 0.66, *SD* = 0.10; atonal: *M* = 0.63, *SD* = 0.06; *t*(17) = 1.02,

*p* = .322, *Cohen's d* = 0.24; **Figure 2D**). This indicates that the amplitudes of beat-related peaks

in the modulation spectrum were comparable in both conditions.

To assess whether the atonal condition shows any drastic perceptive pitch differences

compared with the tonal condition, we computed roughness as measure of sensory dissonance

(using the MIR toolbox; Lartillot *et al.*, 2008). The default frame length of 50 ms (25 ms overlap)

was used. The roughness in the tonal condition was slightly higher than in the atonal condition

(tonal: *M* = 82.85, *SD* = 155.73; atonal: *M* = 60.18, *SD* = 122.02; *t*(11674) = 15.25, *p* < .001,

*Cohen's d* = 0.14; **Figure 2E**). The effect was very small (according to *Cohen's d* conventions) but

implies at least that the atonal condition did not show more sensory dissonance than the tonal

condition.

*Behavioural tracking: Finger-tapping is more variable in the atonal than the tonal condition*

To test how pitch predictability influences behavioural rhythm tracking, we first analysed

differences in inter-tap-intervals between the tonal and atonal conditions. To mitigate the effect

of potential outliers on the group level (see **Figure 3**), we used a non-parametric approach. In the

Experiment 1, the median absolute deviation (MAD) of inter-tap-intervals in the tonal condition

was on average *M* = 29.72 ms (*SD* = 14.99 ms). In the atonal condition, the MAD was slightly

higher, on average *M* = 40.30 ms (*SD* = 35.66 ms). A direct comparison using a Wilcoxon Signed

Rank test indicated that tapping performance was significantly more variable in the atonal than

the tonal condition (difference = 10.6 ms, *Z* = -1.94, *p* = 0.026, *one-tailed*). Out of the 20

participants, 15 (75%) had more variable inter-tap-intervals when tapping in the atonal than in

the tonal condition.

## Behavioural tracking (fingertapping)

**Figure 3. Behavioural tracking results of the main experiment (*N* = 20) and the replication**
**experiment (*N* = 49)**. Shown is the Median Absolute Deviation (MAD) in ms for inter-tap-intervals in both
the tonal and atonal condition. Points indicate individual data for all participants, violin plots show kernel
density estimates, and boxplots show median interquartile ranges and minimum/maximum.

We performed the same analysis for the behavioural follow-up study, which had more than twice

as many participants, and in which individual trials were twice as long as in the original

experiment. The MAD of inter-tap-intervals in the tonal condition was on average $M$ = 42.56 ms

($SD$ = 46.24 ms). In the atonal condition, the MAD was again slightly higher, on average

$M$ = 52.42 ms ($SD$ = 67.33 ms). A Wilcoxon Signed Rank test confirmed that tapping performance

was significantly more variable in the atonal than the tonal condition (difference = 9.9 ms, $Z$ = -

1.78, $p$ = 0.037, *one-tailed*). Out of the 49 participants, 30 (61.2%) had more variable inter-tap-

intervals when tapping in the atonal than in the tonal condition. Together, the results of the

original and replication experiments indicate that there is a small but replicable effect of tonality

on finger-tapping variability: When listening to tonal music, participants tap to the beat more

consistently than when listening to atonal music.

*Cortical tracking of the music envelope*

We first analysed whether participants tracked the acoustic music envelope, band-pass filtered

around the dominant beat frequency, in the tonal and atonal condition compared with chance

level, using cluster-based permutation (**Figure 2A**). In the tonal condition, we found a large

positive cluster of 31 electrodes that tracked amplitude fluctuations significantly ($p$ < .001,

$MI_{sum}$ = 0.332). Equivalently, in the atonal condition, there was a positive cluster of 32 electrodes

that showed significant envelope tracking ($p$ < .001, $MI_{sum}$ = 0.438). There was no evidence for a

hemispheric lateralisation, neither in the tonal nor the atonal condition (both $p_{FDR}$ > 0.82). We

then directly compared envelope tracking in both conditions. This resulted in two negative

clusters, one over frontal electrodes ($p$ < .001, *Cohen's* $d_{peak}$ = -1.17, 7 electrodes) and one over

left occipital electrodes ($p$ = .003, *Cohen's d*$_{peak}$ = -2.09, 3 electrodes). These negative clusters

indicated that the acoustic music envelope was tracked stronger in the atonal than the tonal

condition.

*Envelope tracking for tonal music during passive listening predicts finger-tapping performance*

To test whether envelope tracking during passive listening predicted participants' behavioural

tracking of the beat (i.e., finger-tapping performance), we correlated the MI values per electrode

with participants' average tapping variation (MAD, **Figure 4A**). We found one negative cluster over

left-frontal electrodes that predicted tapping variance in the tonal condition ($p$ = .026, *Cohen's*

*d*$_{peak}$ = -1.02, 2 electrodes). These results indicate that participants who showed stronger

envelope tracking to tonal music had smaller variance (i.e., better performance) when tapping,

than participants with weaker envelope tracking. Envelope tracking when listening to atonal

music did not significantly predict tapping performance in the atonal condition. To compare this

relationship directly between the tonal and atonal condition, we entered the average MI values

of electrodes in the negative cluster as predictor into a regression model, with *tonality* as

additional predictor, an *envelope tracking × tonality* interaction term, and *finger-tapping*

*variability* (MAD) as outcome variable. This overall model was not significant ($F$(3,26) = 1.02,

$p$ = .395) and only explained 7.84% of the variance. The model yielded no main effects (both

$p$ > .32) and no interaction ($p$ > .21), which suggests that the effect of envelope tracking on finger-

tapping is small in the tonal condition, and not statistically different between the tonal and atonal

condition.

**Figure 4. Cortical tracking of acoustic music envelope and its relationship with behavioural measures. A)** Topography of cortical envelope tracking assessed through mutual information (in bits) for both conditions. The right topography shows *t*-values from a direct comparison between tonal and atonal music. **B)** Correlation between cortical envelope tracking and participants' finger-tapping performance. Envelope tracking predicted finger-tapping performance only in the tonal condition, in a left-frontal cluster. Here, stronger envelope tracking was associated with better performance (i.e., less tapping variability). **C)** Correlation between cortical envelope tracking and participants' self-reported musical expertise. Expertise predicted envelope tracking in both conditions, in a frontal (tonal) and fronto-right-lateral (atonal) cluster. Stronger envelope tracking was associated with more musical expertise. **D)**

Correlation between cortical envelope tracking and enjoyment. Enjoyment predicted envelope tracking in
the tonal condition in fronto-central and posterior electrodes. A regression model showed a significant
main effect of enjoyment with no significant interaction ($p$ = .06). Perceiving the music as more pleasant
was associated with stronger envelope tracking.
Note: Significant electrodes are highlighted with white circles.

*Expertise predicts envelope tracking for tonal and atonal music*

Several previous studies have found that musical expertise is associated with stronger neural

synchronisation to music (Di Liberto *et al.*, 2020b; Doelling & Poeppel, 2015; Harding *et al.*, 2019).

We therefore tested the relationship between participants' musical expertise and acoustic

envelope tracking. A large fronto-temporal cluster showed a significant positive correlation

between self-assessed musical competency and music tracking in the tonal condition ($p$ < .001,

*Cohen's $d_{peak}$* = 2.76, 7 electrodes). In the atonal condition, envelope tracking was also positively

predicted by a fronto-temporal cluster ($p$ = .004, *Cohen's $d_{peak}$* = 2.85, 4 electrodes). A regression

model predicting *envelope tracking* (averaged across the electrodes included in the significant

clusters reported above) from *tonality, musical expertise*, and their interaction, ($F$(3,26) = 12.83,

$p$ < .001) explained 51.7% of the variance. Only the main effect of *expertise* was significant

($t$ = 4.15, $p$ < .001; main effect of *tonality* and the interaction both $p$ > .59). These results indicate

that musical expertise is associated with enhanced tracking of the music envelope for both high

predictable (tonal) and low predictable (atonal) music but is unlikely to explain differences in

tracking between conditions.

*Enjoyment predicts envelope tracking for tonal music*

Participants also indicated how pleasant they found listening to the music after each condition

using visual analogue scales. These enjoyment ratings were significantly higher for tonal than

atonal music (*Cohen's d* = 1.45; see above and **Figure 2C**). Ratings were correlated with cortical

envelope tracking across participants. In the tonal condition, two significant clusters showed a

positive correlation between enjoyment and envelope tracking, one frontocentral cluster

($p$ = .026, *Cohen's $d_{peak}$* = 1.40, 3 electrodes [F3, FC1, Fz]) and one occipital cluster ($p$ = .004,

*Cohen's d* = 1.61; 4 electrodes [Oz, O2, PO4, P8]). No significant clusters emerged in the atonal

condition. A regression model predicting *envelope tracking* (averaged across the electrodes

included in the significant clusters reported above) from *enjoyment* ratings, *tonality* and their

interaction ($F$(3,26) = 2.92, $p$ = .047) explained 19.6% of the variance. More *enjoyment* increased

cortical tracking ($t$ = 2.28, $p$ = .029), whereas neither *tonality* ($t$ = 1.92, $p$ = .062), nor an interaction

of *tonality* with *enjoyment* affected tracking ($t$ = -1.03, p = .310).

*Cortical tracking of pitch surprisal*

Pitch surprisal was analysed using the IDyOM model (Pearce, 2005) in both conditions. As

expected, surprisal was higher for notes in the atonal than the tonal condition for both the melody

and bass lines (*Cohen's* $d_{peak}$ = -1.03 and *Cohen's* $d_{peak}$ = -1.06, respectively; see above and

**Figure 5B**). We first analysed whether pitch surprisal was tracked in both conditions above

chance level, in a multivariate analysis including surprisal in melody and bass lines. In the tonal

condition, pitch surprisal was tracked in a large bilateral cluster (**Figure 5A**; $p < .001$,

$MI_{sum}$ = 0.073; 16 electrodes). Likewise, in the atonal condition, pitch surprisal was tracked in a

bilateral electrode cluster ($p < .001$, $MI_{sum}$ = 0.051; 14 electrodes). Although pitch tracking in the

tonal condition appeared to be larger than in the atonal condition, directly comparing the tracking

of pitch surprisal between both conditions yielded no statistically significant clusters. Likewise,

although pitch surprisal tracking appeared larger in the right hemisphere, contrasting left- and

right-hemispheric cluster electrodes yielded no systematic lateralisation of cortical tracking

(tonal: $p_{FDR}$ =.27 and atonal: $p_{FDR}$ =.13, respectively).  These results suggest that listeners form

pitch expectations (i.e., prediction errors) comparable with the IDYOM model, and that pitch

surprisal is represented in the brain to a similar extent in tonal and atonal conditions.

**A** Pitch surprisal tracking

**B** Relationship with tapping performance

**C** Pitch surprisal (IDYOM)

**D** Surprisal tracking vs envelope tracking

**Figure 5. Cortical tracking of pitch surprisal performance. A)** Topography of cortical surprisal tracking assessed through mutual information (in bits) for both conditions. The right topography shows t-values from a direct comparison between tonal and atonal music. Significant electrodes are highlighted with white circles. **B)** Correlation between pitch surprisal tracking and participants' finger-tapping performance. Surprisal tracking predicted finger-tapping performance only in the atonal condition, in a right-frontal cluster. Here, stronger surprisal tracking was associated with worse performance (i.e., higher tapping variability). C) Pitch surprisal values for each note in the melody and bass lines of both tonal and atonal 5-min excerpts. Points indicate data for all notes, violin plots show kernel density estimates, and boxplots show median interquartile ranges and minimum/maximum. Pitch surprisal was higher for the atonal than the tonal condition. **C)** Relationship between cortical tracking of acoustic envelope and pitch surprisal.

*Surprisal tracking predicts finger-tapping performance in the atonal condition*

We also analysed whether the extent to which participants tracked pitch surprisal predicted

their finger-tapping performance. This correlation analysis yielded no significant clusters in

the tonal condition. However, in the atonal condition, the tracking of pitch surprisal was

positively correlated with finger-tapping performance in one frontocentral cluster (**Figure 5B**;

$p$ = .027, *Cohen's $d_{peak}$* = 1.11, 2 electrodes). Again, to be able to draw conclusions about

differences between the tonal and atonal condition, we entered the average MI values of the

positive cluster as predictor into a regression model, with *tonality* as additional predictor, a

*surprisal tracking × tonality* interaction term, and *finger-tapping variability* (MAD) as outcome

variable. The overall regression model ($F(3,36)$ = 7.26, $p$ < .001) explained 37.6% of

variance in finger-tapping variability. Neither the main effect of *surprisal tracking* ($p$ > .49) nor

the main effect of *condition* ($p$ > .16) reached significance. However, the interaction *surprisal*

*tracking × condition* was statistically significant ($t$ = 3.20, $p$ < .003). This interaction stemmed

from an effect, exclusive to the atonal condition, where participants who tracked the pitch

surprisal well, finger-tapped with higher variability than participants with relatively poor

cortical tracking.

*Relationship between envelope tracking and surprisal tracking*

Last, we were interested in the relationship between acoustic envelope tracking and pitch

surprisal tracking, because these measures have not previously been looked at together. We

used the average MI value per cluster (as seen in **Figure 4A** and **Figure 5A&B**) and participant in

a regression model with *acoustic envelope tracking* as outcome variable and *surprisal tracking,*

*tonality* and a *surprisal tracking × tonality* interaction term as predictors. The overall regression

model was not significant ($F(3,36)$ = 1.67, $p$ = .190) and explained 12.2% of the variance. No main

or interaction effect reached significance (all $p$-values > .295; **Figure 5D**). Thus, there seems to

be no systematic relationship between acoustic envelope tracking and pitch surprisal tracking.

## Discussion

533   In the present study we show that the cortical representation of naturalistic continuous music,

534   as measured through envelope tracking, reflects not only rhythm processing, but is also

535   modulated by pitch predictability, musical expertise, and enjoyment (**Figure 6**).

**Figure 6. Overview of results.**

*Pitch predictability affects rhythm processing as reflected in behavioural tracking*

Atonal music can be used to study predictive processing under high uncertainty contexts (Mencke *et al.*, 2022; Mencke *et al.*, 2018). While temporal predictability has been shown to increase pitch discrimination performance (Herbst & Obleser, 2019; Jones *et al.*, 2002), it is unclear whether long-term pitch predictability affects the ability to behaviourally follow the beat particularly in naturalistic musical stimuli. Crucially, in our main experiment and a replication study, we show that when listening to naturalistic music, pitch predictability (modelled on long-term statistics, which reflect exposure to a musical culture, and short-term melodic context) modulates the variability of finger-tapping to the beat. In the tonal condition with higher pitch predictability the finger-tapping performance was more consistent (less variable inter-tap-interval) compared with the atonal low predictability condition. The atonal music in our study contained a rhythmic structure that was identical to the tonal condition, but had generally lower pitch predictability, suggesting that this finding reflects a modulation of behavioural rhythm processing (i.e., finger-tapping to the beat) by pitch predictability. This is in line with and extends previous studies showing expectation effects on musical perception (Herbst & Obleser, 2019; Huron, 2008; Jones *et al.*, 2002; Pearce & Wiggins, 2006)

*Cortical tracking of the music envelope in tonal and atonal music*

At the neural level, the music envelope was tracked in our study for both tonal and atonal music (Figure 4A). The tracking was observed in both conditions with a centro-temporal topography, in accordance with previous reports suggesting auditory cortex generators of the envelope tracking

in speech (Gross et al., 2013; Luo & Poeppel, 2007) and music (Di Liberto et al., 2020b; Doelling

& Poeppel, 2015). Some studies reported a right lateralisation for music envelope tracking

(Doelling & Poeppel, 2015) in line with the asymmetric sampling in time theory (Poeppel, 2003;

Zatorre et al., 2002). The heterogeneous findings in the literature regarding whether a

hemispheric lateralisation is observed have been related to various top-down influences

(Assaneo et al., 2019; Flinker et al., 2019; Zatorre, 2022).

Atonal music was more strongly tracked at frontal and left parietal electrodes compared with

tonal music. Importantly, this was the case, although both conditions had an identical rhythmic

structure and there were no significant acoustic differences in the modulation spectrum (**Figure**

**2D**). We interpret this as evidence that pitch predictability influences neural rhythm tracking. Our

control analysis showed that even when the tracking of pitch intervals was partialled out, atonal

music was still tracked stronger than tonal music (**Figure S1**). Further, the atonal condition did

not carry higher sensory dissonance. These control analyses support the interpretation that it is

pitch predictability, and not low-level pitch differences, that affect rhythm tracking. Our results

from the neural data are also in line with our behavioural findings in that they suggest an effect of

pitch predictability on rhythm processing. Interestingly, Weineck *et al.* (2022) speculated that

more predictable music produces stronger neural synchronisation, which our results contradict.

However, their paradigm did not manipulate pitch predictability and results are therefore not

directly comparable. A predictive coding approach (Friston, 2010; Heilbron & Chait, 2018) could

provide a potential explanation for the observed effect: In the atonal condition, notes were

generally less predictable than in the tonal condition. In line with the assumption of expectation

suppression (Auksztulewicz & Friston, 2016), this likely led to stronger neural prediction errors,

which in turn might have resulted in stronger neural responses to the acoustic envelope (not

unlike a mismatch-negativity response, e.g., Koelsch *et al.*, 2019; Naatanen *et al.*, 2007).

Accordingly, Kern *et al.* (2022) showed that surprising notes elicit stronger neural responses

compared to predictable ones (see also Di Liberto *et al.*, 2020a).

*Increased envelope tracking in tonal music related to better tapping performance*

In the tonal condition, the cortical tracking of the musical envelope correlated with the

behavioural tracking, with higher cortical tracking being associated with an increased ability to

behaviourally follow the beat. The findings are in line with previous research showing a positive

correlation between behavioural performance and cortical tracking of speech (Schmitt *et al.*,

2022; Schubert *et al.*, 2023) and music (Doelling & Poeppel, 2015). The relationship between

cortical tracking and behavioural performance, however, might be more complex than this, as

suggested for speech (Howard & Poeppel, 2010; Pefkou *et al.*, 2017). No correlation was

observed in the atonal condition and furthermore when the correlation effects were tested in a

regression model that included both conditions and selected electrodes, no significant

interaction effect was observed. This makes any interpretation regarding differences between

tonal and atonal music difficult. A potential explanation is that under conditions of low pitch

predictability (as in the atonal condition) the positive relationship between envelope tracking and

behavioural tapping performance is confounded, perhaps due to the increased difficulty of trying

to (unsuccessfully) make predictions about the upcoming notes. In summary, increased

behavioural tracking was related to increased cortical tracking only in the tonal condition, albeit

the effect was small.

*Pitch surprisal is cortically tracked in tonal and atonal music*

We designed our stimuli so that the pitch predictability of the musical pieces was increased in

the atonal compared to the tonal condition (**Figure 5B**). Importantly, our human listeners

significantly tracked pitch surprisal in both conditions, which indicates that their pitch

expectations (or their prediction errors) were comparable with the IDyOM model. Further, there

were no significant condition differences in the cortical tracking of pitch surprisal (**Figure 5A**).

This suggests that participants' neural models of pitch surprisal matches the IDyOM

computations, and the notes in the atonal condition elicited not only more surprisal in the IDyOM

model, but also in listeners' neural response. Cortical tracking of pitch surprisal in natural music

has been rarely investigated. Two recent studies report melodic surprisal tracking in tonal music

that was localised to bilateral superior temporal and Heschl's gyri (amongst others), and

additionally showed either a central topography (using EEG/EcoG, Di Liberto *et al.*, 2020a) or a

broad fronto-temporal (and central) topography (using MEG, Kern *et al.*, 2022). Overall, we found

relatively widespread fronto-temporal tracking of pitch surprisal across conditions, which is in

line with the above results. We here show that listeners consistently track pitch surprisal not only

for high-predictable music, as previously shown, but also for low-predictable music.

*In atonal music lower pitch surprisal tracking is related to better tapping performance*

Interestingly, the surprisal tracking strength was only correlated with the behavioural rhythm

tracking performance in the atonal but not the tonal condition (as shown by the significant

interaction between condition and surprisal tracking). Participants who tracked the pitch

surprisal stronger, meaning they matched the high surprisal values from the IDyOM model, also

showed more variation (worse performance) in their tapping in the atonal condition. Prediction

tendencies have been suggested to vary across participants (Schubert *et al.*, 2023). Our measure

of surprisal tracking might reflect such a tendency, with some individuals being more or less

prone (or able) to make predictions. In the atonal condition with its high uncertainty, pitch

predictions might be less informative for rhythm processing, and listeners who tend to make
(stronger) predictions, which lead to high prediction errors, could have fewer resources to track
the envelope, and to perform well in the tapping task. The few behavioural studies that looked at
long-term pitch surprisal tracking have not related it to the rhythm processing performance (e.g.,
Kern *et al.*, 2022). Our results indicate that the negative effect of pitch surprisal tracking on
behavioural rhythm processing might only be expected when pitch predictability is low, and
prediction errors are high, as in the case of atonal music. The cortical tracking of pitch surprisal
was not systematically related to the cortical tracking of the acoustic envelope, at least not in
our sample of 20 participants that underwent the EEG part of the study.

*Enjoyment and musical expertise are related to cortical envelope tracking*
As expected based on the literature (Mencke *et al.*, 2022; Mencke *et al.*, 2018), the atonal music
condition was rated as less pleasant than the tonal music condition. The individually perceived
pleasure or enjoyment has a strong influence on our everyday music listening behaviour (c.f.,
Gold *et al.*, 2019). In the tonal condition, the enjoyment ratings correlated with the strength of
cortical music envelope tracking, and this pattern was not statistically different in the atonal
condition. The causal nature of this relationship remains unclear: Do listeners show stronger
acoustic envelope tracking because they find the music more enjoyable, or do they find it more
enjoyable because they have better acoustic envelope tracking? Interestingly, a previous study
that investigated whether enjoyment influences neural synchronisation to music did not find a
significant effect (Weineck *et al.*, 2022). The discrepancy with our findings might be due to
differences in experimental paradigms, music choices, quantification of music tracking and
analytical methods. If future studies replicate our finding, this suggests that the individual
preference of listeners should be taken into account when measuring envelope tracking.

Additionally, musical expertise was correlated positively with the cortical envelope tracking of
the music pieces at a cluster of frontal and right temporal electrodes for both the tonal and atonal
condition, with more expertise being related to stronger tracking. Our finding is in line with
previous reports of effects of musical expertise on cortical envelope tracking (Di Liberto *et al.*,
2020b; Doelling & Poeppel, 2015; Harding *et al.*, 2019; but see Weineck *et al.*, 2022 for a null
effect). We here extend these previous results by showing that stronger envelope tracking by
individuals with more musical expertise also extends to atonal music with low pitch
predictability. The effect of musical expertise on auditory processing of music has been related
to increased auditory-motor coupling after musical training (Du & Zatorre, 2017; Rimmele *et al.*,
2021).

Another, not mutually exclusive, interpretation for the effects of both enjoyment and expertise,

could be that both are associated with generally more attention to the musical stimuli. That is,

more musical expertise, as well as more enjoyment of the music, could result in unspecific

increases of attention, which in turn could increase the signal-to-noise ratio of acoustic envelope

tracking (e.g., Ding & Simon, 2012; Keitel *et al.*, 2011; Rimmele *et al.*, 2015). To disentangle

contributions of attention and other variables, future studies could implement measures of

attention or listening effort into their paradigms.

## Conclusion

Our findings suggest that tracking of the envelope of naturalistic music, beyond rhythmic

processing, is modulated by top-down factors such as pitch predictability, musical expertise,

and enjoyment. In addition to the rhythm, musical pitch surprisal is tracked for both low and high

predictable music. This supports the view that long-term musical pitch predictability is

processed in the brain and used to facilitate rhythm processing. For tonal, more predictable

music, the ability to make valid pitch predictions seems to facilitate the ability to behaviourally

follow the rhythm. For atonal music, the reduced pitch predictability results in stronger acoustic

envelope tracking than for tonal music, possibly related to higher prediction errors. These higher

prediction errors also seem to come at the cost of finger-tapping performance, as individuals

with stronger pitch surprisal tracking show worse behavioural rhythm tracking. Overall, our

findings indicate that rhythm processing interacts with non-rhythmic stimulus properties, in our

case pitch surprisal, and listeners' characteristics such as music expertise and enjoyment.

## Acknowledgments

AK is supported by the Medical Research Council [grant number MR/W02912X/1]. AK, CK and JR

are members of the Scottish-EU Critical Oscillations Network (SCONe), funded by the Royal

Society of Edinburgh (RSE Saltire Facilitation Network Award to CK and AK, Reference Number

1963). JR is supported by the Max Planck Institute for Empirical Aesthetics. IM is funded by the

Deutsche Forschungsgemeinschaft (DFG, German Research Foundation; project number

510788453). JR, CP and XG are supported by the Max Planck NYU Center for Language, Music,

and Emotion (CLaME). We thank Roddy Easson for providing the 'tapping person' drawing.

## **References**

Abel, S., Dressel, K., Bitzer, R., Kummerer, D., Mader, I., Weiller, C., & Huber, W. (2009). The separation of processing stages in a lexical interference fMRI-paradigm. *Neuroimage*, *44*(3), 1113-1124. https://doi.org/10.1016/j.neuroimage.2008.10.018

Abrams, E. B., Vidal, E. M., Pelofi, C., & Ripollés, P. (2022). Retrieving musical information from neural data: how cognitive features enrich acoustic ones. Ismir 2022 Hybrid Conference,

Alexandrou, A., Saarinen, T., Kujala, J., & Salmelin, R. (2018). Cortical entrainment: what we can learn from studying naturalistic speech perception. *Language, Cognition and Neuroscience*, 1-13.

Arachchige, C., Prendergast, L. A., & Staudte, R. G. (2022). Robust analogs to the coefficient of variation. *J Appl Stat*, *49*(2), 268-290. https://doi.org/10.1080/02664763.2020.1808599

Assaneo, M. F., Rimmele, J. M., Orpella, J., Ripolles, P., de Diego-Balaguer, R., & Poeppel, D. (2019). The Lateralization of Speech-Brain Coupling Is Differentially Modulated by Intrinsic Auditory and Top-Down Mechanisms. *Front Integr Neurosci*, *13*, 28. https://doi.org/10.3389/fnint.2019.00028

Auksztulewicz, R., & Friston, K. (2016). Repetition suppression and its contextual determinants in predictive coding. *cortex*, *80*, 125-140. https://doi.org/10.1016/j.cortex.2015.11.024

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate - a Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B-Methodological*, *57*(1), 289-300. <Go to ISI>://WOS:A1995QE45300017

Blanco-Elorrieta, E., Ding, N., Pylkkänen, L., & Poeppel, D. (2020). Understanding requires tracking: noise and knowledge interact in bilingual comprehension. *Journal of cognitive neuroscience*, *32*(10), 1975-1983.

Brohl, F., & Kayser, C. (2021). Delta/theta band EEG differentially tracks low and high frequency speech-derived envelopes. *Neuroimage*, *233*, 117958. https://doi.org/10.1016/j.neuroimage.2021.117958

Brohl, F., Keitel, A., & Kayser, C. (2022). MEG Activity in Visual and Auditory Cortices Represents Acoustic Speech-Related Information during Silent Lip Reading. *eNeuro*, *9*(3). https://doi.org/10.1523/ENEURO.0209-22.2022

Celma-Miralles, A., & Toro, J. M. (2019). Ternary meter from spatial sounds: Differences in neural entrainment between musicians and non-musicians. *Brain Cogn*, *136*, 103594. https://doi.org/10.1016/j.bandc.2019.103594

Cervantes Constantino, F., & Simon, J. Z. (2018). Restoration and efficiency of the neural processing of continuous speech are promoted by prior knowledge. *Frontiers in systems neuroscience*, *12*, 56.

Cuddy, L. L., Cohen, A. J., & Mewhort, D. J. (1981). Perception of structure in short melodic sequences. *J Exp Psychol Hum Percept Perform*, *7*(4), 869-883. https://doi.org/10.1037//0096-1523.7.4.869

Di Liberto, G. M., Lalor, E. C., & Millman, R. E. (2018). Causal cortical dynamics of a predictive enhancement of speech intelligibility. *Neuroimage*, *166*, 247-258.

Di Liberto, G. M., Pelofi, C., Bianco, R., Patel, P., Mehta, A. D., Herrero, J. L., . . . Mesgarani, N. (2020a). Cortical encoding of melodic expectations in human temporal cortex. *Elife*, *9*, e51784.

Di Liberto, G. M., Pelofi, C., Shamma, S., & de Cheveigné, A. (2020b). Musical expertise enhances the cortical tracking of the acoustic envelope during naturalistic music listening. *Acoustical Science and Technology*, *41*(1), 361-364.

Dibben, N. (1994). The cognitive reality of hierarchic structure in tonal and atonal music. *Music Perception*, *12*(1), 1-25.

Ding, N., Patel, A. D., Chen, L., Butler, H., Luo, C., & Poeppel, D. (2017). Temporal modulations in speech and music. *Neurosci Biobehav Rev*, *81*(Pt B), 181-187. https://doi.org/10.1016/j.neubiorev.2017.02.011

Ding, N., & Simon, J. Z. (2012). Emergence of neural encoding of auditory objects while listening
to competing speakers. *Proc Natl Acad Sci U S A*, *109*(29), 11854-11859.
https://doi.org/10.1073/pnas.1205381109
Doelling, K. B., & Poeppel, D. (2015). Cortical entrainment to music and its modulation by
expertise. *Proceedings of the National Academy of Sciences*, *112*(45), E6233-E6242.
Dowling, W. J., Kwak, S., & Andrews, M. W. (1995). The time course of recognition of novel
melodies. *Perception & Psychophysics*, *57*, 136-149.
Du, Y., & Zatorre, R. J. (2017). Musical training sharpens and bonds ears and tongue to hear
speech better. *Proc Natl Acad Sci U S A*, *114*(51), 13579-13584.
https://doi.org/10.1073/pnas.1712223114
Engel, A. K., Fries, P., & Singer, W. (2001). Dynamic predictions: oscillations and synchrony in
top-down processing. *Nat Rev Neurosci*, *2*(10), 704-716.
https://doi.org/10.1038/35094565
Fiveash, A., Bella, S. D., Bigand, E., Gordon, R. L., & Tillmann, B. (2022). You got rhythm, or more:
The multidimensionality of rhythmic abilities. *Atten Percept Psychophys*, *84*(4), 1370-
1392. https://doi.org/10.3758/s13414-022-02487-2
Flinker, A., Doyle, W. K., Mehta, A. D., Devinsky, O., & Poeppel, D. (2019). Spectrotemporal
modulation provides a unifying framework for auditory cortical asymmetries. *Nat Hum
Behav*, *3*(4), 393-405. https://doi.org/10.1038/s41562-019-0548-z
Friston, K. (2010). The free-energy principle: a unified brain theory? *Nat Rev Neurosci*, *11*(2), 127-
138. https://doi.org/10.1038/nrn2787
Gold, B. P., Pearce, M. T., Mas-Herrero, E., Dagher, A., & Zatorre, R. J. (2019). Predictability and
uncertainty in the pleasure of music: a reward for learning? *Journal of Neuroscience*,
*39*(47), 9397-9409.
Gorsuch, R. L., & Lehmann, C. S. (2010). Correlation coefficients: Mean bias and confidence
interval distortions. *Journal of Methods and Measurement in the Social Sciences*, *1*(2),
52-65.
Gross, J., Hoogenboom, N., Thut, G., Schyns, P., Panzeri, S., Belin, P., & Garrod, S. (2013). Speech
rhythms and multiplexed oscillatory sensory coding in the human brain. *PLoS Biol*,
*11*(12), e1001752. https://doi.org/10.1371/journal.pbio.1001752
Guan, X., Ren, Z., & Pelofi, C. (2022). py2lispIDyOM: A Python package for the information
dynamics of music (IDyOM) model. *Journal of Open Source Software*, *7*(79), 4738.
Harding, E. E., Sammler, D., Henry, M. J., Large, E. W., & Kotz, S. A. (2019). Cortical tracking of
rhythm in music and speech. *Neuroimage*, *185*, 96-101.
Heilbron, M., & Chait, M. (2018). Great Expectations: Is there Evidence for Predictive Coding in
Auditory Cortex? *Neuroscience*, *389*, 54-73.
https://doi.org/10.1016/j.neuroscience.2017.07.061
Herbst, S. K., & Obleser, J. (2019). Implicit temporal predictability enhances pitch discrimination
sensitivity and biases the phase of delta oscillations in auditory cortex. *Neuroimage*, *203*,
116198. https://doi.org/10.1016/j.neuroimage.2019.116198
Howard, M. F., & Poeppel, D. (2010). Discrimination of speech stimuli based on neuronal
response phase patterns depends on acoustics but not comprehension. *J Neurophysiol*,
*104*(5), 2500-2511. https://doi.org/10.1152/jn.00251.2010
Huron, D. (2008). *Sweet anticipation: Music and the psychology of expectation*. MIT press.
Ince, R. A., Giordano, B. L., Kayser, C., Rousselet, G. A., Gross, J., & Schyns, P. G. (2017). A
statistical framework for neuroimaging data analysis based on mutual information
estimated via a gaussian copula. *Hum Brain Mapp*, *38*(3), 1541-1573.
https://doi.org/10.1002/hbm.23471
Iversen, J. R., Patel, A. D., Nicodemus, B., & Emmorey, K. (2015). Synchronization to auditory and
visual rhythms in hearing and deaf individuals. *Cognition*, *134*, 232-244.
https://doi.org/10.1016/j.cognition.2014.10.018

Jones, M. R., Moynihan, H., MacKenzie, N., & Puente, J. (2002). Temporal aspects of stimulus-
driven attending in dynamic arrays. *Psychol Sci*, *13*(4), 313-319.
https://doi.org/10.1111/1467-9280.00458
Keitel, A., Gross, J., & Kayser, C. (2018). Perceptually relevant speech tracking in auditory and
motor cortex reflects distinct linguistic features. *PLoS Biol*, *16*(3), e2004473.
Keitel, A., Gross, J., & Kayser, C. (2020). Shared and modality-specific brain regions that mediate
auditory and visual word comprehension. *Elife*, *9*. https://doi.org/10.7554/eLife.56972
Keitel, A., Ince, R. A., Gross, J., & Kayser, C. (2017). Auditory cortical delta-entrainment interacts
with oscillatory power in multiple fronto-parietal networks. *Neuroimage*, *147*, 32-42.
https://doi.org/10.1016/j.neuroimage.2016.11.062
Keitel, C., Obleser, J., Jessen, S., & Henry, M. J. (2021). Frequency-Specific Effects in Infant
Electroencephalograms Do Not Require Entrained Neural Oscillations: A Commentary
on Koster et al. (2019). *Psychol Sci*, *32*(6), 966-971.
https://doi.org/10.1177/09567976211001317
Keitel, C., Schroger, E., Saupe, K., & Muller, M. M. (2011). Sustained selective intermodal
attention modulates processing of language-like stimuli. *Exp Brain Res*, *213*(2-3), 321-
327. https://doi.org/10.1007/s00221-011-2667-2
Kern, P., Heilbron, M., de Lange, F. P., & Spaak, E. (2022). Cortical activity during naturalistic
music listening reflects short-range predictions based on long-term experience. *eLife*,
*11*. https://doi.org/10.7554/eLife.80935
Koelsch, S., Vuust, P., & Friston, K. (2019). Predictive Processes and the Peculiar Case of Music.
*Trends Cogn Sci*, *23*(1), 63-77. https://doi.org/10.1016/j.tics.2018.10.006
Koike, K. J., Hurst, M. K., & Wetmore, S. J. (1994). Correlation between the American-Academy-
of-Otolaryngology-Head-and-Neck-Surgery 5-minute hearing test and standard
audiological data. *Otolaryngology-Head and Neck Surgery*, *111*(5), 625-632. <Go to
ISI>://A1994PR79400014
Lakens, D. (2013). Calculating and reporting effect sizes to facilitate cumulative science: a
practical primer for t-tests and ANOVAs. *Front Psychol*, *4*, 863.
https://doi.org/10.3389/fpsyg.2013.00863
Lartillot, O., Toiviainen, P., & Eerola, T. (2008). A Matlab Toolbox for Music Information Retrieval.
In C. Preisach, H. Burkhardt, L. Schmidt-Thieme, & R. Decker (Eds.), *Data Analysis,
Machine Learning and Applications*. Springer-Verlag.
Lerdahl, F. (2019). *Composition and cognition: Reflections on contemporary music and the
musical mind*. University of California Press.
Lesenfants, D., & Francart, T. (2020). The interplay of top-down focal attention and the cortical
tracking of speech. *Scientific reports*, *10*(1), 1-10.
Leys, C., Ley, C., Klein, O., Bernard, P., & Licata, L. (2013). Detecting outliers: Do not use standard
deviation around the mean, use absolute deviation around the median. *Journal of
experimental social psychology*, *49*(4), 764-766.
Luo, H., & Poeppel, D. (2007). Phase patterns of neuronal responses reliably discriminate speech
in human auditory cortex. *Neuron*, *54*(6), 1001-1010.
https://doi.org/10.1016/j.neuron.2007.06.004
Marion, G., Di Liberto, G. M., & Shamma, S. A. (2021). The Music of Silence: Part I: Responses to
Musical Imagery Encode Melodic Expectations and Acoustics. *Journal of Neuroscience*,
*41*(35), 7435-7448.
Mencke, I., Omigie, D., Quiroga-Martinez, D. R., & Brattico, E. (2022). Atonal Music as a Model for
Investigating Exploratory Behavior. *Frontiers in Neuroscience*, *16*, 793163.
Mencke, I., Omigie, D., Wald-Fuhrmann, M., & Brattico, E. (2018). Atonal Music: Can Uncertainty
Lead to Pleasure? *Front Neurosci*, *12*, 979. https://doi.org/10.3389/fnins.2018.00979
Mencke, I., Quiroga-Martinez, D. R., Omigie, D., Michalareas, G., Schwarzacher, F., Haumann,
N. T., . . . Brattico, E. (2021). Prediction under uncertainty: Dissociating sensory from

cognitive expectations in highly uncertain musical contexts. *Brain Res*, *1773*, 147664.
https://doi.org/10.1016/j.brainres.2021.147664
Morillon, B., Arnal, L. H., Schroeder, C. E., & Keitel, A. (2019). Prominence of delta oscillatory
rhythms in the motor cortex and their relevance for auditory and speech perception.
*Neuroscience            &            Biobehavioral            Reviews*.
https://doi.org/https://doi.org/10.1016/j.neubiorev.2019.09.012
Naatanen, R., Paavilainen, P., Rinne, T., & Alho, K. (2007). The mismatch negativity (MMN) in basic
research of central auditory processing: a review. *Clin Neurophysiol*, *118*(12), 2544-2590.
https://doi.org/10.1016/j.clinph.2007.04.026
Neuloh, G., & Curio, G. (2004). Does familiarity facilitate the cortical processing of music
sounds? *Neuroreport*, *15*(16), 2471-2475. https://doi.org/10.1097/00001756-
200411150-00008
Nozaradan, S., Peretz, I., & Mouraux, A. (2012). Selective neuronal entrainment to the beat and
meter embedded in a musical rhythm. *J Neurosci*, *32*(49), 17572-17581.
https://doi.org/10.1523/JNEUROSCI.3203-12.2012
Obleser, J., & Kayser, C. (2019). Neural Entrainment and Attentional Selection in the Listening
Brain. *Trends in cognitive sciences*, *23*(11), 913-926.
https://doi.org/0.1016/j.tics.2019.08.004
Ockelford, A., & Sergeant, D. (2013). Musical expectancy in atonal contexts: Musicians'
perception of "antistructure". *Psychology of Music*, *41*(2), 139-174.
Oldfield, R. C. (1971). The assessment and analysis of handedness: the Edinburgh inventory.
*Neuropsychologia*, *9*(1), 97-113. http://www.ncbi.nlm.nih.gov/pubmed/5146491
Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J. M. (2011). FieldTrip: Open source software for
advanced analysis of MEG, EEG, and invasive electrophysiological data. *Comput Intell*
*Neurosci*, *2011*, 156869. https://doi.org/10.1155/2011/156869
Park, H., & Kayser, C. (2019). Shared neural underpinnings of multisensory integration and trial-
by-trial perceptual recalibration in humans. *Elife*, *8*.
Pearce, M. T. (2005). *The construction and evaluation of statistical models of melodic structure*
*in music perception and composition* https://openaccess.city.ac.uk/id/eprint/8459
Pearce, M. T. (2018). Statistical learning and probabilistic prediction in music cognition:
mechanisms of stylistic enculturation. *Ann N Y Acad Sci*, *1423*(1), 378-395.
https://doi.org/10.1111/nyas.13654
Pearce, M. T., Ruiz, M. H., Kapasi, S., Wiggins, G. A., & Bhattacharya, J. (2010). Unsupervised
statistical learning underpins computational, behavioural, and neural manifestations of
musical expectation. *Neuroimage*, *50*(1), 302-313.
Pearce, M. T., & Wiggins, G. A. (2006). Expectation in Melody: The Influence of Context and
Learning. *Music Perception*, *23*(5), 377–405. https://doi.org/10.1525/mp.2006.23.5.377
Pearce, M. T., & Wiggins, G. A. (2012). Auditory expectation: the information dynamics of music
perception and cognition. *Top Cogn Sci*, *4*(4), 625-652. https://doi.org/10.1111/j.1756-
8765.2012.01214.x
Peelle, J. E., Gross, J., & Davis, M. H. (2013). Phase-locked responses to speech in human
auditory cortex are enhanced during comprehension. *Cereb Cortex*, *23*(6), 1378-1387.
https://doi.org/10.1093/cercor/bhs118
Pefkou, M., Arnal, L. H., Fontolan, L., & Giraud, A. L. (2017). theta-band and beta-band neural
activity reflects independent syllable tracking and comprehension of time-compressed
speech. *J Neurosci*, *37*(33), 7930-7938. https://doi.org/10.1523/JNEUROSCI.2882-
16.2017
Poeppel, D. (2003). The analysis of speech in different temporal integration windows: cerebral
lateralization as 'asymmetric sampling in time'. *Speech Communication*, *41*(1), 245-255.
https://doi.org/10.1016/S0167-6393(02)00107-3

Reetzke, R., Gnanateja, G. N., & Chandrasekaran, B. (2021). Neural tracking of the speech
envelope is differentially modulated by attention and language experience. *Brain Lang*,
*213*, 104891. https://doi.org/10.1016/j.bandl.2020.104891
Repp, B. H. (2005). Sensorimotor synchronization: a review of the tapping literature. *Psychon Bull*
*Rev*, *12*(6), 969-992. https://doi.org/10.3758/bf03206433
Rimmele, J. M., Kern, P., Lubinus, C., Frieler, K., Poeppel, D., & Assaneo, M. F. (2021). Musical
Sophistication and Speech Auditory-Motor Coupling: Easy Tests for Quick Answers. *Front*
*Neurosci*, *15*, 764342. https://doi.org/10.3389/fnins.2021.764342
Rimmele, J. M., Zion Golumbic, E., Schroger, E., & Poeppel, D. (2015). The effects of selective
attention and speech acoustics on neural speech-tracking in a multi-talker scene.
*Cortex*, *68*, 144-154. https://doi.org/10.1016/j.cortex.2014.12.014
Rovetti, J., Copelli, F., & Russo, F. A. (2022). Audio and visual speech emotion activate the left
pre-supplementary motor area. *Cogn Affect Behav Neurosci*, *22*(2), 291-303.
https://doi.org/10.3758/s13415-021-00961-2
Sammler, D. (2020). Splitting speech and music. *science*, *367*(6481), 974-976.
https://doi.org/10.1126/science.aba7913
Schmitt, R., Meyer, M., & Giroud, N. (2022). Better speech-in-noise comprehension is associated
with enhanced neural speech tracking in older adults with hearing impairment. *cortex*,
*151*, 133-146. https://doi.org/10.1016/j.cortex.2022.02.017
Schubert, J., Schmidt, F., Gehmacher, Q., Bresgen, A., & Weisz, N. (2023). Cortical speech
tracking is related to individual prediction tendencies. *Cereb Cortex*, *33*(11), 6608-6619.
https://doi.org/10.1093/cercor/bhac528
Schulze, K., Jay Dowling, W., & Tillmann, B. (2011). Working memory for tonal and atonal
sequences during a forward and a backward recognition task. *Music Perception*, *29*(3),
255-267.
Smith, Z. M., Delgutte, B., & Oxenham, A. J. (2002). Chimaeric sounds reveal dichotomies in
auditory perception. *Nature*, *416*(6876), 87-90. https://doi.org/10.1038/416087a
Song, J., & Iverson, P. (2018). Listening effort during speech perception enhances auditory and
lexical processing for non-native listeners and accents. *Cognition*, *179*, 163-170.
Tierney, A., & Kraus, N. (2015). Neural entrainment to the rhythmic structure of music. *J Cogn*
*Neurosci*, *27*(2), 400-408. https://doi.org/10.1162/jocn_a_00704
Vuvan, D. T., Podolak, O. M., & Schmuckler, M. A. (2014). Memory for musical tones: the impact
of tonality and the creation of false memories. *Front Psychol*, *5*, 582.
https://doi.org/10.3389/fpsyg.2014.00582
Weineck, K., Wen, O. X., & Henry, M. J. (2022). Neural synchronization is strongest to the spectral
flux of slow music and depends on familiarity and beat salience. *eLife*, *11*.
https://doi.org/10.7554/eLife.75515
Zarate, J. M., Tian, X., Woods, K. J., & Poeppel, D. (2015). Multiple levels of linguistic and
paralinguistic features contribute to voice recognition. *Sci Rep*, *5*, 11475.
https://doi.org/10.1038/srep11475
Zatorre, R. J. (2001). Neural specializations for tonal processing. *Ann N Y Acad Sci*, *930*, 193-210.
https://doi.org/10.1111/j.1749-6632.2001.tb05734.x
Zatorre, R. J. (2022). Hemispheric asymmetries for music and speech: Spectrotemporal
modulations and top-down influences. *Front Neurosci*, *16*, 1075511.
https://doi.org/10.3389/fnins.2022.1075511
Zatorre, R. J., Belin, P., & Penhune, V. B. (2002). Structure and function of auditory cortex: music
and speech. *Trends Cogn Sci*, *6*(1), 37-46.
http://www.ncbi.nlm.nih.gov/pubmed/11849614
Zion Golumbic, E. M., Ding, N., Bickel, S., Lakatos, P., Schevon, C. A., McKhann, G. M., . . .
Schroeder, C. E. (2013). Mechanisms underlying selective neuronal tracking of attended

speech      at      a      "cocktail      party".      *Neuron*,      *77*(5),      980-991.
946              https://doi.org/10.1016/j.neuron.2012.12.037
947    Zuk, N. J., Murphy, J. W., Reilly, R. B., & Lalor, E. C. (2021). Envelope reconstruction of speech
948              and music highlights stronger tracking of speech at low frequencies. *PLoS Comput Biol*,
949              *17*(9), e1009358. https://doi.org/10.1371/journal.pcbi.1009358

# Supplemental analysis

To explore whether pitch intervals influence acoustic envelope tracking (and our result that

envelope tracking is stronger in the atonal than the tonal condition), we performed a conditional

MI analysis (Brohl *et al.*, 2022; Ince *et al.*, 2017). Here, all parameters were identical to the main

analysis of acoustic envelope tracking (**Figure 4a**), but we included pitch intervals as the to-be-

partialled-out signal in both tonal and atonal conditions.

To extract pitch values, we first used the MIDI toolbox (Eerola & Toiviainen, 2004) for melody and

bass lines of the tonal and atonal conditions. The difference between subsequent MIDI pitch

values was then used to obtain a measure of pitch intervals, and to create a continuous signal

where each pitch interval was as long as the corresponding note (equivalent to how the surprisal

signal was created). These pitch interval signals were then normalised using Gaussian Copulas,

before including them in the conditional MI analysis. This means each MI computation (for tonal

and atonal conditions) included the EEG signal, the envelope signal, and the to-be-partialled out

pitch interval signals for both the melody and bass lines.

With pitch intervals in melody and bass lines partialled out, the tracking of the acoustic envelope

yielded similar results as the initial tracking analysis (**Figure S1A**). When compared against

chance, the envelope was tracked significantly in both tonal and atonal conditions. In the tonal

condition, we found a large positive cluster of 31 electrodes that tracked amplitude fluctuations

significantly ($p$ = .002, $MI_{sum}$ = 0.325). Equivalently, in the atonal condition, there was a positive

cluster of 32 electrodes that showed significant envelope tracking ($p$ < .002, $MI_{sum}$ = 0.430). A

direct comparison showed that envelope in the atonal condition was tracked stronger than in the

tonal condition in two clusters, one over frontal electrodes ($p$ = .002, Cohen's $d_{peak}$ = -1.21, 6

electrodes) and one over left occipital electrodes ($p$ = .002, Cohen's $d_{peak}$ = -2.17, 3 electrodes).

This suggests that, with pitch intervals partialled out, the acoustic envelope is still tracked above

chance level, and this tracking is still stronger in the atonal condition.

**A** Envelope tracking - pitch intervals partialled out

**B** Envelope tracking w./w.o. pitch intervals

**Figure S1. Conditional MI analysis of envelope tracking with pitch intervals partialled out. A)**
Topography of cortical envelope tracking assessed through mutual information (in bits) for both
conditions, with partialled out pitch intervals in the melody and bass lines. The right topography shows *t*-values from a direct comparison between tonal and atonal music. **B)** Envelope tracking for all electrodes,
averaged across participants, for the tonal and atonal conditions, and for the original MI analysis
(tracking of envelope) and the conditional MI analysis (tracking of envelope, pitch intervals partialled
out). MI values are systematically lower when pitch intervals are partialled out, but this effect is not
different for the tonal or atonal condition.

To directly compare MI values across tracking analyses (with and without pitch intervals
partialled out), we averaged MI values for each electrode across participants (**Figure S1B**) and
performed a repeated measures 2x2 ANOVA with factors *tracking measure* (envelope tracking vs
conditional envelope tracking) and *tonality* (tonal vs atonal). Both main effects were significant
(*tracking measure*: $F(1,31) = 129.93$, $p < .001$, $\eta_p^2 = .807$; *tonality*: $F(1,31) = 31.39$, $p < .001$,
$\eta_p^2 = .503$). The MI values were lower when the tracking of pitch intervals was partialled out (*tonal*:
$M_{Env} = 0.011$, $SD = 0.006$, $M_{Env-Pitch} = 0.010$, $SD = 0.006$; *atonal*: $M_{Env} = 0.014$, $SD = 0.006$, $M_{Env-Pitch} = 0.013$, $SD = 0.006$). MI values were higher in the atonal than the tonal condition, reflecting
the difference shown in **Figure S1A**. Crucially, the interaction *tracking measure × tonality* was not
significant ($F(1,31) = 1.51$, $p > .227$, $\eta_p^2 = .047$), which implies that pitch-interval tracking did not
affect the conditions differently.

In sum, while there is a small but highly consistent decrease in MI when the tracking of pitch

intervals is partialled out, this effect is not different in tonal and atonal conditions. Our previous

finding that the acoustic envelope in the atonal condition is tracked stronger than in the tonal

condition, therefore also holds true in the conditional MI analysis. This means that the found

differences in envelope tracking are unlikely to be due to low-level differences in the tracking of

pitch intervals.

**References**

Brohl, F., Keitel, A., & Kayser, C. (2022). MEG Activity in Visual and Auditory Cortices
Represents Acoustic Speech-Related Information during Silent Lip Reading. *eNeuro*,
*9*(3). https://doi.org/10.1523/ENEURO.0209-22.2022
Eerola, T., & Toiviainen, P. (2004). *MIDI Toolbox: MATLAB Tools for Music Research*.
University of Jyväskylä.
Ince, R. A., Giordano, B. L., Kayser, C., Rousselet, G. A., Gross, J., & Schyns, P. G. (2017).
A statistical framework for neuroimaging data analysis based on mutual information
estimated via a gaussian copula. *Hum Brain Mapp*, *38*(3), 1541-1573.
https://doi.org/10.1002/hbm.23471

1014

1015