1 **TITLE:**
2 Viromes vs. mixed community metagenomes: choice of method dictates interpretation of
3 viral community ecology
4

5 **AUTHORS:**
6 James C. Kosmopoulos[1,2], Katherine M. Klier[1,3], Marguerite V. Langwig[1,3], Patricia Q.
7 Tran[1], and Karthik Anantharaman[1,4*]
8
[1] Department of Bacteriology, University of Wisconsin-Madison, Madison, Wisconsin,
USA
[2] Microbiology Doctoral Training Program, University of Wisconsin-Madison, Madison,
Wisconsin, USA
[3] Freshwater and Marine Sciences Program, University of Wisconsin-Madison, Madison,
Wisconsin, USA
[4] Department of Integrative Biology, University of Wisconsin-Madison, Madison,
Wisconsin, USA
[*] Correspondence: karthik@bact.wisc.edu
18

ABSTRACT

Background
Viruses, the majority of which are uncultivated, are among the most abundant biological entities on Earth. From altering microbial physiology to driving community dynamics, viruses are fundamental members of microbiomes. While the number of studies leveraging viral metagenomics (viromics) for studying uncultivated viruses is growing, standards for viromics research are lacking. Viromics can utilize computational discovery of viruses from total metagenomes of all community members (hereafter metagenomes) or use physical separation of virus-specific fractions (hereafter viromes). However, differences in the recovery and interpretation of viruses from metagenomes and viromes obtained from the same samples remain understudied.

Results
Here, we compare viral communities from paired viromes and metagenomes obtained from 60 diverse samples across human gut, soil, freshwater, and marine ecosystems. Overall, viral communities obtained from viromes were more abundant and species rich than those obtained from metagenomes, although there were some exceptions. Despite this, metagenomes still contained many viral genomes not detected in viromes. We also found notable differences in the predicted lytic state of viruses detected in viromes vs metagenomes at the time of sequencing. Other forms of variation observed include genome presence/absence, genome quality, and encoded protein content between viromes and metagenomes, but the magnitude of these differences varied by environment.

Conclusions
Overall, our results show that the choice of method can lead to differing interpretations of viral community ecology. We suggest that the choice of whether to target a metagenome or virome to study viral communities should be dependent on the environmental context and ecological questions being asked. However, our overall recommendation to researchers investigating viral ecology and evolution is to pair both approaches to maximize their respective benefits.

KEYWORDS:
Virome, metagenome, viral ecology, differential abundance

INTRODUCTION

Viruses exist in all known ecosystems and infect cells from all domains of life. As the most abundant biological entity on Earth [1,2], viruses significantly impact the ecology and evolution of their hosts [3,4], play pivotal roles in microbial community succession [5], contribute to community-wide metabolic processes [6–8], and are a source of novel therapies being used to combat a worldwide antimicrobial resistance crisis [9,10]. Advances in these areas have been enabled by large-scale investigations into entire communities of viruses which have revealed tremendous amounts of previously unknown virus diversity in human [11–13] and environmental [14–18] systems. Since their hosts largely have not been isolated, these investigations have utilized viral metagenomics (viromics) to examine thousands of viral genomes from DNA/RNA sequence data extracted directly from host-associated and environmental samples. While the number of studies using viromics has been growing in the past decade [18–20], the sampling and analytical methods used vary greatly [20,21]. Although there have recently been efforts to establish standards for analyzing viruses from sequence data [19–21], standards in extraction methodologies are still largely lacking.

There are two ways to identify genomic sequences of viral communities. First, one can sequence metagenomes of a mixed microbial community (hereafter metagenomes). Second, virus-like particles (VLPs) can be separated from a sample to enrich for viral community DNA prior to sequencing (hereafter viromes). Both methods involve computational approaches to identify viral sequences after sequencing, but they each have their own benefits and drawbacks. For instance, viromes do not offer the host context that metagenomes can [22,23]. Thus, investigations into virus☐host relationships can benefit from the use of metagenomes. On the other hand, predicting virus☐host relationships from metagenomes alone remains difficult and can often only be achieved for a fraction of viral genomes [22,23]. Furthermore, rare, low-abundance viruses are diverse and have significant impacts on their communities [24–26]. These viruses are often not detected in metagenomes because viruses represent a small fraction of the mixed community [27]. However, they are detectable in viromes because viruses and other forms of protected environmental DNA represent the majority of sequences in these samples [27,28]. It has also been argued that active viruses exist mostly in an intracellular state and therefore metagenomes are more likely to be appropriate to study viral communities [29,30]. However, the high rates of viral lysis and virion production that have been widely observed [31] might suggest that sequences captured in viromes could better reflect the active viral community. Overall, most studies of viral ecology typically use either method depending on their scope and environmental context.

Although most viral ecology studies have typically utilized either viromes or metagenomes, only a few have leveraged both methods. For example, in an agricultural soil ecosystem, the cumulative richness of viruses in viromes was orders of magnitude greater than that of metagenomes [27]. In a seasonally anoxic freshwater lake, viromes were richer in viruses than metagenomes [6] but the magnitude of this difference was much smaller than that of the soil study. Viral community composition in the freshwater

111 lake was also mostly influenced by sample type (viromes or metagenomes) [6], while
112 human gut viral communities were mostly influenced by the individual human host
113 rather than sample method [32]. These studies offer novel insights into the viral and
114 prokaryotic community composition of their respective ecosystems, but they remain to
115 be synthesized together into a broader context of method application.
116
The few existing studies that leverage paired viromes and metagenomes have largely
paid attention to community-level differences in viruses assembled from each approach,
but it remains unknown whether or how this influences the interpretation of ecology and
evolution, and the abundance of viruses at the genome level. While differences in
genome contiguity and assembly quality between viromes and metagenomes have
been discussed [33,34], focused comparisons of viral genomes assembled from
viromes versus metagenomes are lacking. Similarly, since the gene content of viruses
can vary greatly both within and between populations [35–37], existing community-level
comparisons of viromes and metagenomes are unable to highlight any gene-level
differences between the two methods.
127
Here, we directly compare paired viromes and metagenomes from multiple samples
obtained from four different environments: a freshwater lake, the global oceans, the
human gut microbiome, and soil. After using the same, standardized analytical workflow
for every sample and across each environment, we compared viral sequence yields,
genome presence/absence, viral genome quality, and virus gene differential abundance
between viromes and metagenomes. Last, we discuss the unique insights offered by
each approach and suggest when to apply viromes, metagenomes, or both methods
when studying viral communities in different environmental contexts.
136
137 **METHODS**
138
Data acquisition
In an effort to compare paired viromes and mixed community metagenomes from a
variety of environments, we obtained sequence reads from publicly available studies.
We searched for short-read collections that met the following criteria: (1) both viromes
and metagenomes must have been generated for the same biological samples, (2)
neither virome nor metagenome samples underwent whole-genome or multiple-
displacement amplification, and (3) metadata were available that allowed virome and
metagenome pairs originating from the same biological sample to be identified, or read
filenames made it otherwise clear.
148
Among the datasets that met the criteria, we chose collections of paired viromes and
metagenomes to represent four vastly different environments: a freshwater lake, marine
water columns from the global oceans, the human gut microbiome, and soil. Raw reads
from virome and metagenome libraries sequenced from water column samples of Lake
Mendota, Wisconsin, USA [6] were chosen to represent a freshwater environment.
Reads from soil samples of an agricultural field in Davis, California, USA [27] were
chosen to represent a soil environment. Fecal sample sequence reads of a cohort in

Cork, Ireland [11] were chosen to represent human gut samples. Finally, reads from the
Tara Oceans database were obtained to represent marine samples [38,39].
158
Marine, soil, and human gut reads were obtained from NCBI GenBank [40] using
SRAtoolkit (hpc.nih.gov/apps/sratoolkit.html) from BioProjects PRJEB1787 (marine
metagenomes), PRJEB4419 (marine viromes), PRJNA545408 (soil viromes and
metagenomes) and PRJNA646773 (human gut viromes and metagenomes). For the
Tara Oceans marine samples, we obtained reads for the <0.22 µm fractions of samples
for viromes and the 0.22-3.0 µm fractions for metagenomes (Figure 1A), and read
libraries were removed if there was no counterpart library available from the same
sample station and depth for the other size fraction. Freshwater virome and
metagenome reads were obtained directly by the first author of the study, and can also
be found at the JGI Genome Portal under Proposal ID 506328. For all environments, all
read libraries obtained were composed of paired-end Illumina reads. A detailed
description of the data sources for this study and relevant information can be found in
Supplementary Table 1.
172
Sequence read quality control and assembly
Freshwater samples were previously sequenced by the Department of Energy Joint
Genome Institute (DOE JGI) and thus sequence reads underwent quality control (QC)
and were assembled into contigs within the DOE JGI metagenome workflow [41]. To
reduce biases that could have been introduced by different QC and assembly methods,
read QC and metagenome assembly were performed following the same assembly
workflow with the same sequence of software (and versions), commands, and
parameters as JGI (Figure 1B). Briefly, raw reads from marine, soil, and human gut
samples underwent quality filtering and trimming with BBDuk and BBMap using
rqcfilter.sh which were then error-corrected with bbcms. Filtered, error-corrected reads
were split into separate mates and singletons using reformat.sh, and the resulting read
pairs were imported to metaSPAdes v3.13.0 [42] for assembly. Read lengths and counts
at each step of QC were obtained with readlen.sh from the BBTools suite
(sourceforge.net/projects/bbmap/) and assembly statistics were obtained for samples
from all environments using metaQUAST v5.2.0 [43] which were parsed in R [44] and
plotted using ggplot2 [45] to generate Figure 2.
189
Virus identification, mapping, binning, quality assessment, and taxonomic
assignment with ViWrap
For every sample, ViWrap v1.2.1 [46] was run (Figure 1B) with the assembled sample
contigs and filtered reads using the parameter "--identify_method vb" to only use
VIBRANT v1.2.1 [47] to identify viral contigs, as well as the options "--input_length_limit
10000" and "--reads_mapping_identity_cutoff 0.90" to adhere to established
recommended minimum requirements for virus detection [20]. In accordance with these
standards for virus detection, only viral contigs of at least 10 kb were retained for
downstream analyses. After using VIBRANT to identify viral contigs, ViWrap mapped
reads to the input assembly using Bowtie2 v2.4.5 [48]. Read recruitment to all
assembled contigs at least 10 kb was calculated using SAMtools v1.17 [49] using the
read mapping files generated by Bowtie2. Read recruitment statistics were then filtered

to only include the viral contigs with a length of at least 10 kb identified by VIBRANT.
Additionally, ViWrap used the resulting coverage files to bin viral contigs into vMAGs
with vRhyme v1.1.0 [50].
205
In this study, both binned viral contigs and unbinned singletons are together referred to
as vMAGs. The quality, completeness, and redundancy of the resulting vMAGs were
assessed with CheckV v1.0.1 [51] by ViWrap. ViWrap then grouped vMAGs within
samples into genus-level clusters with vConTACT2 v0.11.0 [52] and then into species-
level clusters with dRep v3.4.0 [53]. ViWrap assigned taxonomy to vMAGs by aligning
proteins with DIAMOND v2.0.15 [54] to NCBI RefSeq viral proteins [55], the VOG HMM
database v97 [56], and IMG/VR v4.1 high-quality vOTU representative proteins [57].
Summary statistics on the number of viral contigs, read recruitment, vMAGs, taxonomy,
and genome quality gathered by ViWrap for each sample were parsed in R and plotted
using ggplot2 to generate Figure 2, Figure S2, Figure S3, and Figure S4.
216

Predicting the lytic state of vMAGs
ViWrap provides a prediction of the lytic state for all vMAGs it identifies [46], i.e.,
whether a vMAG is likely to represent a lytic virus, a lysogenic virus, an integrated
prophage flanked with cellular DNA, or not determined. ViWrap makes these
determinations based on a combination of annotation results from VIBRANT and
binning results from vRhyme. Possible predictions by ViWrap include "lytic scaffold",
"lytic virus", "lysogenic scaffold", "lysogenic virus", and "integrated prophage". ViWrap
handles instances when vRhyme bins multiple integrated prophage sequences or lytic
and integrated prophage sequences together by splitting the vMAG back into individual
scaffolds to avoid retaining potentially contaminated bins (see
github.com/AnantharamanLab/ViWrap). Furthermore, the distinction made by ViWrap
between "scaffold" and "virus" depends on the genomic context of the contigs in a
vMAG [50] and the estimated completion of a vMAG [51]. Here, we simplified these
predictions using a custom python script and did not distinguish between predictions on
the "virus" or "scaffold" level and used the results predicted by ViWrap to label vMAGs
as "lytic", "lysogenic", or "integrated prophage".
233

vMAG presence/absence analysis
Although ViWrap employed dRep to dereplicate vMAGs into species-level clusters at
95% ANI within samples, species representative vMAGs were still redundant between
samples after running ViWrap on each. To dereplicate vMAGs across all samples, an
additional ANI-based approach was taken. Redundant vMAGs from each sample were
gathered and dereplicated using dRep v3.4.3 [53] with a minimum genome length of 10
kb in addition to the options "-pa 0.8 -sa 0.95 -nc 0.85" to set the ANI thresholds for
primary and secondary clusters to 80% and 95%, respectively, and to require a
minimum covered fraction of 85%, as recommended by established benchmarks for
viral community analyses [20]. The parameters "-comW 0 -conW 0 -strW 0 -N50W 0 -
sizeW 1 -centW 0" were also used when running dRep so the resulting species
representative vMAGs were simply the largest vMAGs in each cluster.
246

Bowtie2 mapping indices were created from fasta files containing all representative
vMAGs from each environment, separately, to be used in competitive alignments. For
each environment, filtered reads from every sample were separately mapped to the
environment's mapping index using Bowtie2 v2.5.1 with default parameters to perform
an end-to-end alignment and report single best matches at a minimum of 90% identity.
The resulting alignment files were sorted and indexed using SAMtools v1.17 [49].
Sorted and indexed files were used with CoverM v0.6.1 (github.com/wwood/CoverM) to
obtain covered fraction (genome breadth) statistics at the vMAG level for reads mapping
with at least 90% identity. A minimum breadth threshold of 75% was used to establish
the detection of a vMAG in each read sample in accordance with previously established
recommendations [20]. Lists of unique representative vMAG IDs determined to be
present in samples in this way were used to generate Figure 3 and Figure S4 with the R
package eulerr (CRAN.R-project.org/package=eulerr) [58,59]. Labels for Figure 3 were
manually edited for clarity.
261
Virus genome assembly comparison
To address a preexisting notion that metagenomes typically result in truncated or less-
complete viral genome assemblies than viromes [21,27,60], we identified vMAGs
shared between viromes and metagenomes. Using our previously generated dRep
results, we identified pairs of vMAGs that met the following criteria: (1) one vMAG was
assembled from a virome and the other a metagenome, (2) each vMAG in the pair was
placed in the same species-level cluster, (3) both vMAGs were assembled from the
same sample source, (4) the virome-assembled vMAG was a single contig and
predicted by CheckV to be complete, and (5) the metagenome-assembled vMAG was
predicted by CheckV to be incomplete.
272
A single pair was chosen among the resulting candidates based on their respective
lengths. Each genome was then subjected to noncompetitive mapping of filtered reads
from the virome and metagenome of the same sample source. This resulted in four read
mapping files: virome reads mapped to the virome-assembled vMAG, virome reads
mapped to the metagenome-assembled vMAG, metagenome reads mapped to the
virome-assembled vMAG, and metagenome reads mapped to the metagenome-
assembled vMAG. For each file, the read depths d at each genome position were
obtained using SAMtools v1.17 [49] with the option "depth", and then $\log_{10}$ normalized
by the total number of reads in the sample n in hundreds of millions to obtain a
normalized read depth.
283

$$\text{normalized read depth} = \log_{10} \frac{d}{(n \cdot 10^{-8})}$$

284
The two vMAGs were aligned using Mauve [61] and BLASTn v2.5.0 from the BLAST+
suite [62] to identify regions in the virome-assembled genome that were missing from
the metagenome-assembled genome, as well as gaps and alternate sequences. This
revealed the metagenome-assembled vMAG in the pair to be on the opposite strand as
the virome-assembled vMAG, so downstream analyses of this vMAG were performed
on its reverse-complement. Finally, each vMAG in the chosen pair was reannotated for

gene predictions and function using Pharokka v1.4.1 [63] with default settings. The
resulting read depths by genome position and unassembled regions were plotted using
ggplot2 and arrows representing gene prediction coordinates were added with gggenes
v0.5.1 (wilkox.org/gggenes) to generate Figure 4. Highlighted regions and coloring for a
selection of genes of interest were added manually to Figure 4.

Differential abundance of viral proteins
We sought to identify protein-coding viral genes that were differentially abundant across
virome and metagenome assemblies. For each environment (both viromes and
metagenomes), we combined all nucleotide sequences of protein-coding genes
predicted by Prodigal [64] that were encoded on viral contigs >10 kb identified by
VIBRANT into a database of redundant gene sequences. These databases were then
dereplicated, separately by environment, using MMseqs2 v14.7e284 [65]. We used the
command "mmseqs easy-search" to estimate pairwise average nucleotide identities
(ANI) for all genes in each database, with parameters "--min-seq-id 0.95 -c 0.80 –cov-
mode 1" to only retain alignments with minimum ANI of 0.95 and a minimum aligned
fraction to the target sequence of 0.80. A clustered graph was generated from the
pairwise ANI estimates using mcl with mcxload v14-137 [66] to obtain gene clusters,
and the longest gene within each cluster was chosen to be the cluster's dereplicated
representative. Bowtie2 mapping indices were separately generated from the four
databases of dereplicated gene representatives of each environment. For each
environment, filtered reads from all samples were mapped to the Bowtie2 index of
dereplicated genes corresponding to the same environment, using the same
parameters and filtering steps as in the vMAG presence/absence analysis above.

Tables of raw mapped read counts for each dereplicated gene representative were
obtained for each environment using CoverM. These tables were used to build negative
binomial generalized models of gene counts with DESeq2 [67] to infer genes that were
differentially abundant across viromes and metagenomes for each environment,
separately. The extraction method (virome or metagenome) and sample source were
included as factors in the models for each environment, and the DESeq2 workflow
employed Wald tests to compare the counts between viromes and metagenomes. For
each test, the resulting $\log_2$ fold changes reported by DESeq2 were shrunken using the
function "lfcShrink" with adaptive Student's *t* prior shrinkage estimators. We used a
false-discovery rate adjusted *P*-value cutoff of 0.05 for the Wald test results as well as a
minimum shrunken $\log_2$ fold change of 0.58 (corresponding to a minimum fold change
of 1.5) as requirements to determine if a given gene was enriched in either virome or
metagenome samples of a given environment. The results were visualized using
ggplot2 to generate Figure 5A.

PHROG [68] functional predictions for all dereplicated gene representatives were
obtained by running Pharokka v1.4.1 [60] on each dereplicated gene database. The
resulting PHROG annotations and functional categories were mapped back to the
DESeq2 significant genes to obtain the presence of PHROG functional categories in
each enrichment (virome or metagenome). The relative abundance of PHROG
categories among all genes in each enrichment group was calculated and plotted with

ggplot2 to generate Figure 5B. To assess the over- or underrepresentation of any
PHROG category within either enrichment group, we performed hypergeometric tests
on the genes assigned to each enrichment group for every environment, separately,
using the function "phyper" from the stats R package [44]. The resulting *P*-values were
false-discovery rate adjusted, and significant results were plotted using ggplot2 to
generate Figure 5C.
RESULTS
Table 1:

Figure 1. Sampling and analytical approaches used to generate metagenomes, viromes, and vMAGs.
(A) Overview of sampling approaches to generate viromes and metagenomes. Viromes were sequenced from
a size fraction below 0.22 μm or from a virus-like particle fraction achieved from ultracentrifugation [11,27].
Metagenomes were sequenced using one of two main approaches: DNA from the bulk sample was extracted
and sequenced, allowing the recovery of DNA from prokaryotes, viruses, and other microbes. Alternatively,
after filtering a sample to isolate virus-like particles in the <0.22 μm fraction, other studies extracted and
sequenced DNA from the remaining >0.22 μm fraction that did not pass through the filter [6,38,39]. (B)
Overview of metagenome/virome assembly and virus identification methods to obtain viral metagenome-
assembled genomes (vMAGs). (C) Overview of methods for the vMAG presence/absence analysis. Figure

345 **Sources of data used in this study.**

Environment	Sample origin	Source	Virus enrichment approach	# of virome-metagenome sample pairs used	Sample design
Human gut	Fecal samples; Cork, Ireland	Shkoporov et al., 2019 [11]	0.45 µm filtration, ultracentrifugation, & polyethylene glycol (PEG) precipitation	10	Individuals, timepoint
Freshwater	Oxic & anoxic water columns; Lake Mendota, Madison, WI, USA	Tran et al., 2023 [6]	0.22 µm filtration & $FeCl_3$ precipitation	14	Water column depth, timepoint
Marine	Tara Oceans	Pesant et al., 2015; Sunagawa et al., 2015 [38,39]	0.22 µm filtration & $FeCl_3$ precipitation	21	Water column depth, geographic location
Soil	Tomato field; Davis, CA, USA	Santos-Medellin et al., 2021 [27]	Amended 1% potassium citrate (AKC) resuspension, 0.22 µm filtration	15	Soil amendment, plot, timepoint

Viromes were successful in enriching for viral sequences
Sequencing depth within and between viromes versus metagenomes varied (Figure
2A). Freshwater and human gut viromes had a significantly higher sequencing depth
than metagenomes, while marine metagenomes had a higher sequencing depth than
viromes (Figure 2A). There was no difference in depth between viromes and
metagenomes of soil samples (Figure 2A). Because of this observed variation in
sequencing depth, results hereafter were normalized to sequencing depth unless
otherwise specified. Reads from viromes of all environments mapped back to their
assembled contigs (>10 kb) at a significantly higher rate than metagenomes (Figure
2B). Strikingly, soil viromes recruited upward of 25% of filtered reads while all soil
metagenomes recruited less than <1% of filtered reads. Further inspection of soil
metagenome assembly statistics revealed a median N50 <3,000, even when only
calculating statistics for contigs >2,000 bp (Figure S1). The poor read recruitment of the
soil metagenome assemblies is likely a result of the poor contiguity of the assemblies
arising from high community complexity in soils [69,70].

Although the differences between viromes and metagenomes with respect to
sequencing depth and read recruitment varied by environment, viromes from all
environments had reads mapping to viral contigs at a greater rate than metagenomes
(Figure 2C). All assemblies (metagenomes and viromes) except for the human gut had
a greater proportion of viral to nonviral contigs (Figure 2D). Moreover, viromes from all
environments except for the human gut had a higher total number of viral contigs than
metagenomes (Figure S2A). Marine and soil viromes had a higher total number of

vMAGs than metagenomes (Figure S2B). When considering only "high-quality" vMAGs
that are estimated to represent complete or near-complete viral genomes [51], viromes
from all environments had a greater yield than metagenomes (Figure S2C). Similarly,
after dereplicating vMAGs to species-level clusters within samples, viromes had a

Figure 2. Read recruitment and the enrichment of viral sequences were higher in viromes than metagenomes. Points indicate an individual metagenome/virome assembly. Significance was inferred by Wilcoxon rank sum test: ns $p > 0.05$; * $p \leq 0.05$; ** $p \leq 0.01$; *** $p \leq 0.001$; **** $p \leq 0.0001$. (A) While virome samples yielded significantly more read pairs after quality filtering in freshwater and human gut samples, marine metagenomes had greater sequencing depth than viromes, and there was no difference in soil samples. (B) With a minimum alignment identity cutoff of 90%, filtered read pairs from all environments mapped back to assembled contigs >10 kb at a significantly higher rate than metagenomes. (C) In all tested environments, virome assemblies contained more read pairs mapping to viral contigs as a proportion of all quality-filtered read pairs (mapped or unmapped) than metagenome assemblies. (D) All tested environments except human gut samples contained a greater proportion of viral contigs to all assembled contigs >10 kb.

374 higher viral species richness than metagenomes among marine and soil assemblies.
375 However, there was no difference in viral species richness between methods among
376 freshwater and human gut assemblies (Figure S2D).
377
The abundance of lytic and lysogenic viruses in viromes* vs. *metagenomes
varied
Among human gut assemblies, there was no significant difference between the number
of lytic vMAGs from viromes compared to metagenomes, while freshwater, marine, and
soil assemblies had a higher number of lytic vMAGs in viromes compared to
metagenomes (Figure S3A). In contrast, there was no difference in the number of
lysogenic vMAGs between viromes and metagenomes of freshwater and human gut
assemblies, while marine and soil viromes contained significantly more lysogenic
vMAGs than metagenomes (Fig S3B). Freshwater metagenomes contained significantly
more vMAGs predicted to represent integrated prophage (Figure S3C). Integrated
prophage vMAGs were found in viromes across all four environments (Figure S3C).
Strikingly, marine and soil viromes contained significantly more integrated prophage
vMAGs than metagenomes (Figure 3C). Closer inspection revealed that soil
metagenomes did not contain any vMAGs predicted to represent integrated prophages
at all. Given that the total number of vMAGs generated from marine and soil
metagenomes was so low compared to their viromes (Figure S2B), these striking
differences are explained by the low virus richness in these metagenomes overall. Last,
while there was a small observable increase in the normalized number of integrated
prophages in human gut metagenomes, these differences were not significant (Figure
S3C).
398
Viromes and metagenomes have unique and shared vMAGs
Dereplication and read mapping yielded 24,761 unique species-representative vMAGs
in freshwater assemblies, 18,331 in marine assemblies, 9,039 in soil assemblies, and
2,271 in human gut assemblies, with a total of 54,402 unique vMAGs identified across
all environments (Figure 3A). Of this total, 2,539 were found only in metagenome
assemblies, 32,601 were found only in virome assemblies, and 19,262 were found in
both (Figure 3B). Overall, virome assemblies from all four environments contained more
unique vMAGs than metagenome assemblies (Figure 3C). Soil virome assemblies
contained nearly all vMAGs detected in soil metagenomes, except for a single vMAG
found unique to soil metagenomes (Figure 3C). Notably, more vMAGs were detected in
both viromes and metagenomes of freshwater and human gut samples than were
detected in either method, alone (Figure 3C).
411
We also examined the presence and absence of vMAGs in viromes and metagenomes
separated by their predicted lytic state. More lytic vMAGs (Figure 3D), lysogenic vMAGs
(Figure 3E), and integrated prophages (Figure 3F) were detected in viromes than
metagenomes for all environments. However, freshwater assemblies had more lytic
vMAGs detected in both methods than lytic vMAGs present in only one method (Figure
3D). Similarly, the human gut had more lysogenic vMAGs and integrated prophages
present in both methods than those present in only one method (Figure 3E-F). However,
the patterns of detection for integrated prophages may have been caused by virome

reads originating from excised lysogenic/temperate virus genomes that had mapped to
metagenome vMAGs integrated in host DNA.
422

Figure 3. vMAGs assembled from viromes were not detected in most metagenome samples. Euler diagrams generated using eulerr (CRAN.R-project.org/package=eulerr) [58,59] with IDs of unique species-level vMAGs detected in the labeled category; quantities within areas are given beneath labels. An individual vMAG was marked as detected in a virome/metagenome if reads from the virome/metagenome mapped to the contigs in the vMAG with a minimum breadth of 75% across the entire vMAG. (A) Total number of vMAGs in each environment, regardless of method. (B) All vMAGs and environments, separated by method. (C) All vMAGs, separated by environment and method. (D) Predicted lytic vMAGs, separated by environment and method. (E) Predicted lysogenic vMAGs, separated by environment and method. (F) Predicted integrated prophage vMAGs, separated by environment and method.

Virome assembly resulted in a more complete viral genome

Past arguments in favor of utilizing virome extractions to study viral communities have
cited a tendency to assemble more complete viral genomes with greater depth than
those assembled from metagenomes [21,27,60]. To test this, we identified the same
species vMAG from a virome and from a metagenome. The virome-assembled viral

genome was nearly 38 kb in length with 70 gene predictions (Figure 4, Table S2), and
was predicted to be complete by CheckV [51] due to the presence of direct terminal
repeats. The metagenome-assembled viral genome, however, was predicted by
CheckV to be incomplete and was nearly 5 kb shorter than the virome assembly and
contained only 57 gene predictions (Figure 4, Table S2).
434
The missing regions in the metagenome-assembled viral genome spanned both ends of
the contig (Figure 4). These regions covered eleven genes with unknown functions that
were present in the virome but not the metagenome assembly, as well as the first 527
bases of a phage portal protein (Figure 4, Table S2). Additionally, the virome-assembled
viral genome contained a 130 bp region spanning two genes predicted to encode a
hypothetical protein and a tail protein (Figure 4, Table S2). This 130 bp region was
absent from the metagenome assembly, resulting in a single, fused gene prediction for
a phage tail protein (Figure 4, Table S2). The only region we identified in the
metagenome-assembled viral genome that was absent from the virome assembly was a
single 3 bp sequence over the portal protein (Table S2). Finally, although this genome
was incompletely assembled from the metagenome, metagenome reads mapped over
the entire length of the virome-assembled genome (Figure 4, Table S3). Virome reads
also mapped to both assemblies of the same genome with a depth up to two orders of
magnitude greater than metagenome reads (Figure 4, Table S3).
449
450

Figure 4. An incomplete metagenome-assembled viral genome was complete in its corresponding virome. A single-contig, complete viral genome identified from a virome assembly was detected but was incompletely assembled in the sample's corresponding metagenome. Areas highlighted in gray represent regions in the virome-assembled genome that were absent from the metagenome-assembled genome. Reads yielded from the virome and metagenome of the same sample source were each mapped to both versions of the genome assembly. Arrows along the x-axis represent predicted genes that are colored by the extraction method of their genome's origin, except for a selection of genes of interest that are colored by their functional

Viral genes are differentially abundant across viromes and metagenomes

We identified a total of 414,780 protein-coding viral genes after dereplication across all environments and extraction methods. Of these, 13,099 proteins came from human gut assemblies, 206,127 from freshwater assemblies, 116,900 from marine assemblies, and 78,654 from soil assemblies (Table 2, Table S4). Out of all dereplicated genes, a total of 72,082 unique genes were differentially abundant across extraction methods (Wald test *P* <0.05, FDR adjusted) (Table 2, Table S4). Only 55 of these genes were from the human gut, while 64,999 genes were from freshwater samples, 5,722 from marine samples, and 1,306 from soil samples (Table 2, Table S4). Using a minimum fold change cutoff of ±1.5, we found that 67,521 of the differentially abundant genes were enriched in either virome or metagenome samples (Table 2, Table S4, Figure 5A). The remaining 4,561 genes were differentially abundant but did not meet the minimum fold change of 1.5 (Table 2, Table S4, Figure 5A). We did not identify any genes that were enriched in either virome or metagenome samples from the human gut (Table 2, Figure 5A). However, 37,683 and 25,328 genes were enriched in viromes and metagenomes from freshwater samples, respectively (Table 2, Table S4, Figure 5A). Among marine samples, only 222 genes were enriched in viromes whereas 3,265 were enriched in metagenome samples (Table 2, Table S4, Figure 5A). Finally, 432 genes were enriched in soil viromes and 591 were enriched in soil metagenomes (Table 2, Table S4, Figure 5A).

Table 2. Number of genes throughout the differential abundance (DA) workflow.

Environment	Number of genes before dereplication	Number of genes after dereplication (% of before)	Differentially abundant genes (% of dereplicated)	Virome-enriched genes (% of DA)	Metagenome-enriched genes (% of DA)
Human gut	8.39×10^4	1.31×10^4 (16%)	55 (0.004%)	0	0
Freshwater	1.02×10^6	2.06×10^5 (20%)	6.50×10^4 (32%)	3.77×10^4 (58%)	2.53×10^4 (39%)
Marine	6.75×10^5	1.17×10^5 (17%)	5.72×10^3 (4.9%)	222 (3.9%)	3.27×10^3 (57%)
Soil	4.42×10^5	7.87×10^4 (18%)	1.31×10^3 (1.7%)	432 (33%)	591 (45%)
Total	**2.22×10^6**	**4.15×10^5 (19%)**	**7.21×10^4 (17%)**	**3.83×10^4 (53%)**	**2.92×10^4 (40%)**

To predict potential functions for the differentially abundant genes enriched in either viromes or metagenomes, we used PHROG [68] functional categories predicted by Pharokka [63]. Out of the 67,521 unique genes enriched in viromes or metagenomes across all environments, Pharokka assigned PHROG functional categories to a total of 11,115 genes (16%), 6,247 in viromes and 4,868 in metagenomes (Table S4). Because predicted PHROG functional categories were largely present in both virome- and metagenome-enriched genes across the three environments (Figure 5B), we performed hypergeometric tests on enriched genes from each environment to determine whether any functional categories were over or underrepresented in viromes or metagenomes. We found nine PHROG categories that were significantly over- or underrepresented between viromes and metagenomes across freshwater, marine, and soil samples

(hypergeometric test *P* <0.05, FDR adjusted) (Figure 5C, Table S5). Generally, genes
encoding viral structural proteins such as head-tail connectors, packaging proteins, and
tail proteins were underrepresented in metagenomes and overrepresented in viromes
across freshwater and soil samples, while marine samples displayed the opposite
pattern (Figure 5C, Table S5). Integration and excision coding genes were
overrepresented in freshwater and marine metagenomes but underrepresented in
freshwater viromes (Figure 5C, Table S5). Conversely, lysis genes were
underrepresented in freshwater metagenomes and overrepresented in viromes, but
were overrepresented in marine metagenomes.
494

Figure 5. Protein-coding viral genes are differentially abundant across viromes and metagenomes and have predictable functions. (A) Differential abundance of protein-coding viral genes as inferred by DESeq2 [67]. Points indicate unique, dereplicated protein-coding viral genes that were annotated from viral contigs assembled from the environment indicated by the panel labels. Enrichment of a given gene in virome or metagenome samples was determined if the resulting fold change was at least 1.5. (Wald test *P* <0.05, FDR adjusted). No protein-coding viral genes were determined to be significantly enriched in the virome or metagenome human gut assemblies. (B) Relative abundance and (C) over/underrepresentation of PHROG [68] functional categories assigned to differentially abundant genes displayed in (A) (hypergeometric test *P* <0.05, FDR adjusted). Categories without an arrow in a given environment/method were not significantly over or underrepresented in

495
496
497

DISCUSSION

The sequencing of whole virus communities in recent years has resulted in an explosion
of known viral diversity and viral community ecology studies [12,13,16,17,57,71].
Assembly of virus communities can be achieved either by sequencing extracted DNA
from the total, mixed community of prokaryotes, eukaryotes, and viruses within a
sample to generate metagenomes. Viral communities can also be assembled by
enriching for virus-like particle DNA during extraction to generate viromes. Although
viromes can generally offer a more focused view of viruses in a sample compared to
metagenomes [33], the consequences of choosing one sampling method over the other
have been relatively unexplored and limited to individual study ecosystems [5,6,27].
Here, we applied the same analytical methods to collections of paired virome and
metagenome sequence reads to directly infer the unique and shared results gained
from each sample method. We assembled, annotated, and analyzed 60 pairs of viromes
and metagenomes across four different environments and found that the similarities and
differences between each method varied across environments.

Viromes, by design, typically allow more viral species and genome coverage to be
obtained compared to metagenomes [33]. In support of this, virome assemblies here
generally contained more viral contigs, more binned vMAGs, higher species richness,
and greater read recruitment to vMAGs. Interestingly, there were some exceptions
among freshwater and human gut samples. We observed no difference in the number of
vMAGs or in viral species richness between viromes and metagenomes of the human
gut or freshwater. There was additionally no difference in the number of viral contigs
from the human gut.

While there have been a handful of studies in the past that have examined viral
community data resulting from viromes in comparison to metagenomes [6,11,27,72,73],
even fewer have taken a closer look at specific genome-level differences that result
across the two methods. While we only focused on one viral species in this context, we
found that a virome assembly resulted in a more complete viral genome with greater
sequencing depth than the genome assembled from a metagenome of the same
sample. Notably, the metagenome sample contained reads that mapped over the entire
length of the complete version of the genome. Although some viral genomes may be
incompletely assembled in metagenomes, their full sequences may be assembled if the
metagenome reads are mapped to a higher quality virome assembly or reference
genome.

Freshwater and marine metagenome samples used here were recovered from >0.22
µm size fractions, while human gut and soil metagenomes were unfiltered by particle
size. Considering this, any observed differences between viromes and metagenomes
from freshwater and marine assemblies may have been driven by the approach used to
generate the metagenomes. On the other hand, differences (or lack thereof) between
viromes and metagenomes from soil and human gut assemblies may have been driven
by the low abundance of viral DNA relative to nonviral DNA in bulk, unfiltered samples.

Nonetheless, both freshwater and marine metagenomes contained substantial numbers
of viral contigs and vMAGs despite efforts to filter the viral fraction. Furthermore, there
were striking differences between viromes and metagenomes from soil samples, as well
as in human gut samples to a lesser extent, both of which did not have their viral
fraction filtered from the metagenome fraction. Altogether, this highlights the importance
of utilizing enrichment techniques that are tailored to the environment of interest and the
research questions being asked.
551
Whether the purpose is to assign taxonomy [74], reveal mechanisms to avoid host
defenses [75], identify auxiliary metabolic genes [76], or investigate mobile reservoirs
for antimicrobial resistance genes [77,78], obtaining functional gene predictions is a
critical step in analyses of viral communities. However, it can be quite challenging to
assign functional predictions to viral genes annotated from metagenomic environmental
data due to their large sequence diversity and the undercharacterization of viruses.
Thus, annotating genes in complex viral communities often reveals a substantial
amount of viral "dark matter" represented as genes with no known function that encode
"hypothetical" proteins [23,79,80]. This challenge was indeed present here, as we could
obtain functional predictions for only 16% of genes enriched in viromes or
metagenomes. Nonetheless, we identified several functional categories across the three
environments where genes were differentially abundant.
564
Our results show that one's choice of extraction method does indeed influence the
identification of gene families, but the significance and magnitude of differences vary
between environments. We found an overrepresentation of integration and excision
genes in freshwater and marine metagenomes with an underrepresentation in
freshwater viromes. However, lysis genes were underrepresented in freshwater
metagenomes and overrepresented in freshwater viromes. This is consistent with our
observations that freshwater metagenomes contained a greater number of integrated
prophage vMAGs than viromes. On the other hand, this contrasts with our observation
that there was no difference in the proportion of lysogenic vMAGs between freshwater
viromes and metagenomes, and that marine viromes contained more lysogenic and
integrated vMAGs than metagenomes. Regardless of the exact mechanism(s), as a
consequence, the choice between viromes and metagenomes can significantly
influence one's interpretation of viral communities based on gene annotations.
578
579 **CONCLUSIONS**
580
In many contexts, viromes revealed more viral sequences and diversity than
metagenomes. Hence, extracting viromes may be more advantageous than
metagenomes when studying viral communities (Table 3). However, a noticeable
number of viruses were detected only in metagenomes in all four environments tested
here. Thus, we recommend that researchers investigating viral communities extract both
viromes and mixed-community metagenomes in pairs from the same biological
samples, when possible (Table 3). However, if one is restricted to using just one
method, viromes present the better option for virus-focused studies in most
environments.

Table 3. Recommendations for choosing extraction methods depending on research context.

Context	Recommended method(s)	Rationale
Viral community dynamics, overall virus diversity, assembly of uncultivatable virus genomes	Virome	Viromes generally contained more viral species and greater viral sequence enrichment than metagenomes.
Bacterial/archaeal communities, no interest in viruses	Metagenomes	Viromes are unnecessary to the study of just the cellular members of communities.
Fast-growing, highly dynamic communities, and/or lytic viruses	Virome	Assuming viral lysis is prevalent due to the present biotic or abiotic conditions, viromes will enrich for lytic viruses.
Slow-growing, low-biomass communities, and/or integrated viruses	Metagenomes	Assuming lysogeny is prevalent due to the present biotic or abiotic conditions, detecting viruses integrated in the host genome require metagenomics.
Host□virus interactions	Paired viromes & metagenomes	Metagenomes are necessary to provide any host context. While metagenomes alone can yield some viral genomes, viromes are also recommended to maximize viral genome assembly.
Maximization of total virus diversity	Paired viromes & metagenomes	Both viromes and metagenomes resulted in the assembly of viral genomes not detected in the other method. Utilizing both methods can maximize the detection and assembly of as many viral genomes as possible.

ABBREVIATIONS

VLP: Virus-like particle

ANI: Average nucleotide identity

PEG: Polyethylene glycol

vMAG: Viral metagenome-assembled genome

AKC: Amended 1% potassium citrate

vOTU: Viral operational taxonomic unit

DA: Differentially abundant

DECLARATIONS

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The datasets analyzed during the current study are available in the following repositories: Freshwater, originally presented by Tran et al. [6] and deposited to the JGI Genome Portal under Proposal ID 506328; Marine, originally presented by Pesant et al. [38] and Sunagawa et al. [39] and deposited to the NCBI Sequence Read Archive under BioProject accessions PRJEB1787 and PRJEB4419; Human gut, originally presented by Shkoporov et al. [11] and deposited to the NCBI Sequence Read Archive under BioProject accession PRJNA545408; Soil, originally presented by Santos-Medellin et al. [27] and deposited to the NCBI Sequence Read Archive under BioProject accession PRJNA646773. All scripts and intermediate files to reproduce the figures and tables presented here are available at github.com/jamesck2/ViromeVsMetagenome.

Competing interests
The authors declare that they have no competing interests.

Funding
This research was supported by National Institute of General Medical Sciences of the National Institutes of Health under award number R35GM143024, and by the National Science Foundation under grant numbers DBI2047598 and OCE2049478.

Authors' contributions
Conceptualization, J.C.K. and K.A.; Methodology, J.C.K. and K.A.; Software, J.C.K.; Validation, J.C.K.; Formal Analysis, J.C.K.; Investigation, J.C.K., K.M.K., M.V.L., P.Q.T., and K.A.; Resources, K.A.; Data curation, J.C.K., P.Q.T., and K.A.; Writing — Original Draft, J.C.K., K.A. ; Writing — Review & Editing, J.C.K., K.M.K., M.V.L., P.Q.T., and K.A.; Visualization, J.C.K.; Supervision, K.A.; Project Administration, K.A.; Funding Acquisition, K.A.

Acknowledgments
We are thankful to all authors of the studies that originally generated and distributed the data analyzed here. We also gratefully acknowledge the insights provided by Cody Martin during this study.

REFERENCES
1. Wommack KE, Colwell RR. Virioplankton: Viruses in Aquatic Ecosystems [Internet]. MICROBIOLOGY AND MOLECULAR BIOLOGY REVIEWS. 2000. Available from: https://journals.asm.org/journal/mmbr
2. Cesar Ignacio-Espinoza J, Solonenko SA, Sullivan MB. The global virome: Not as big as we thought? Curr Opin Virol. Elsevier B.V.; 2013. p. 566–71.
3. Stern A, Sorek R. The phage-host arms race: Shaping the evolution of microbes. BioEssays. 2011. p. 43–51.
4. Kosmopoulos JC, Campbell DE, Whitaker RJ, Wilbanks EG. Horizontal Gene Transfer and CRISPR Targeting Drive Phage-Bacterial Host Interactions and Coevolution in "Pink Berry" Marine Microbial Aggregates. Vives M, editor. Appl Environ Microbiol [Internet]. 2023;89. Available from: https://journals.asm.org/doi/10.1128/aem.00177-23

5. Santos-Medellín C, Blazewicz SJ, Pett-Ridge J, Emerson JB. Viral but not bacterial community succession is characterized by extreme turnover shortly after rewetting dry soils. bioRxiv. 2023;

6. Tran PQ, Bachand SC, Peterson B, He S, Anantharaman K. Viral impacts on microbial activity and biogeochemical cycling in a seasonally anoxic freshwater lake. bioRxiv [Internet]. 2023; Available from: https://doi.org/10.1101/2023.04.19.537559

7. Hurwitz BL, U'Ren JM. Viral metabolic reprogramming in marine ecosystems. Curr Opin Microbiol. Elsevier Ltd; 2016. p. 161–8.

8. Kieft K, Zhou Z, Anderson RE, Buchan A, Campbell BJ, Hallam SJ, et al. Ecology of inorganic sulfur auxiliary metabolism in widespread bacteriophages. Nat Commun. 2021;12.

9. Fujimoto K, Kimura Y, Shimohigoshi M, Satoh T, Sato S, Tremmel G, et al. Metagenome Data on Intestinal Phage-Bacteria Associations Aids the Development of Phage Therapy against Pathobionts. Cell Host Microbe. 2020;28:380-389.e9.

10. Gordillo Altamirano FL, Barr JJ. Phage Therapy in the Postantibiotic Era. Clin Microbiol Rev [Internet]. 2019;32. Available from: http://cmr.asm.org/

11. Shkoporov AN, Clooney AG, Sutton TDS, Ryan FJ, Daly KM, Nolan JA, et al. The Human Gut Virome Is Highly Diverse, Stable, and Individual Specific. Cell Host Microbe. 2019;26:527-541.e5.

12. Camarillo-Guerrero LF, Almeida A, Rangel-Pineros G, Finn RD, Lawley TD. Massive expansion of human gut bacteriophage diversity. Cell. 2021;184:1098-1109.e9.

13. Shah SA, Deng L, Thorsen J, Pedersen AG, Dion MB, Castro-Mejía JL, et al. Expanding known viral diversity in the healthy infant gut. Nat Microbiol. 2023;8:986–98.

14. Paez-Espino D, Zhou J, Roux S, Nayfach S, Pavlopoulos GA, Schulz F, et al. Diversity, evolution, and classification of virophages uncovered through global metagenomics. Microbiome. 2019;7.

15. Gregory AC, Zayed AA, Conceição-Neto N, Temperton B, Bolduc B, Alberti A, et al. Marine DNA Viral Macro- and Microdiversity from Pole to Pole. Cell. 2019;177:1109-1123.e14.

16. Gaïa M, Meng L, Pelletier E, Forterre P, Vanni C, Fernandez-Guerra A, et al. Mirusviruses link herpesviruses to giant viruses. Nature [Internet]. 2023; Available from: https://www.nature.com/articles/s41586-023-05962-4

17. Hillary LS, Adriaenssens EM, Jones DL, McDonald JE. RNA-viromics reveals diverse communities of soil RNA viruses with the potential to affect grassland ecosystems across multiple trophic levels. ISME Communications. 2022;2.

18. Roux S, Emerson JB. Diversity in the soil virosphere: to infinity and beyond? Trends Microbiol. 2022;30:1025–35.

19. Roux S, Adriaenssens EM, Dutilh BE, Koonin E V., Kropinski AM, Krupovic M, et al. Minimum information about an uncultivated virus genome (MIUVIG). Nat Biotechnol. 2019;37:29–37.

20. Roux S, Emerson JB, Eloe-Fadrosh EA, Sullivan MB. Benchmarking viromics: an *in silico* evaluation of metagenome-enabled estimates of viral community composition and diversity. PeerJ. 2017;5:e3817.

21. Kieft K, Anantharaman K. Virus genomics: what is being overlooked? Curr Opin Virol. Elsevier B.V.; 2022.

22. Roux S, Camargo AP, Coutinho FH, Dabdoub SM, Dutilh BE, Nayfach S, et al.
iPHoP: An integrated machine learning framework to maximize host prediction for
metagenome-derived viruses of archaea and bacteria. PLoS Biol. 2023;21.

23. Roux S, Hallam SJ, Woyke T, Sullivan MB. Viral dark matter and virus–host
interactions resolved from publicly available microbial genomes. Elife. 2015;4.

24. Holmfeldt K, Solonenko N, Shah M, Corrier K, Riemann L, VerBerkmoes NC, et al.
Twelve previously unknown phage genera are ubiquitous in global oceans. Proc Natl
Acad Sci U S A. 2013;110:12798–803.

25. Pascoal F, Costa R, Magalhães C. The microbial rare biosphere: Current concepts,
methods and ecological principles. FEMS Microbiol Ecol. Oxford University Press; 2021.

26. Garin-Fernandez A, Pereira-Flores E, Glöckner FO, Wichels A. The North Sea goes
viral: Occurrence and distribution of North Sea bacteriophages. Mar Genomics.
2018;41:31–41.

27. Santos-Medellin C, Zinke LA, ter Horst AM, Gelardi DL, Parikh SJ, Emerson JB.
Viromes outperform total metagenomes in revealing the spatiotemporal patterns of
agricultural soil viral communities. ISME Journal. 2021;15:1956–70.

28. Lücking D, Mercier C, Alarcón-Schumacher T, Erdmann S. Extracellular vesicles are
the main contributor to the non-viral protected extracellular sequence space. ISME
Communications. 2023;3:112.

29. Forterre P. Manipulation of cellular syntheses and the nature of viruses: The virocell
concept. Comptes Rendus Chimie. Elsevier Masson SAS; 2011. p. 392–9.

30. López-Pérez M, Haro-Moreno JM, Gonzalez-Serrano R, Parras-Moltó M,
Rodriguez-Valera F. Genome diversity of marine phages recovered from Mediterranean
metagenomes: Size matters. PLoS Genet. 2017;13.

31. Breitbart M, Bonnain C, Malki K, Sawaya NA. Phage puppet masters of the marine
microbial realm. Nat Microbiol. 2018;3:754–66.

32. Chen C, Yan Q, Yao X, Li S, Lv Q, Wang G, et al. Alterations of the gut virome in
patients with systemic lupus erythematosus. Front Immunol. 2023;13.

33. Roux S, Matthijnssens J, Dutilh BE. Metagenomics in Virology. Encyclopedia of
Virology. Elsevier; 2021. p. 133–40.

34. Trubl G, Hyman P, Roux S, Abedon ST. Coming-of-Age Characterization of Soil
Viruses: A User's Guide to Virus Isolation, Detection within Metagenomes, and Viromics.
Soil Syst. 2020;4:23.

35. Dion MB, Oechslin F, Moineau S. Phage diversity, genomics and phylogeny. Nat
Rev Microbiol. 2020;18:125–38.

36. Mavrich TN, Hatfull GF. Bacteriophage evolution differs by host, lifestyle and
genome. Nat Microbiol. 2017;2:17112.

37. Gregory AC, Zayed AA, Conceição-Neto N, Temperton B, Bolduc B, Alberti A, et al.
Marine DNA Viral Macro- and Microdiversity from Pole to Pole. Cell. 2019;177:1109-
1123.e14.

38. Pesant S, Not F, Picheral M, Kandels-Lewis S, Le Bescot N, Gorsky G, et al. Open
science resources for the discovery and analysis of Tara Oceans data. Sci Data.
2015;2:150023.

39. Sunagawa S, Coelho LP, Chaffron S, Kultima JR, Labadie K, Salazar G, et al.
Structure and function of the global ocean microbiome. Science (1979). 2015;348.

40. Sayers EW, Bolton EE, Brister JR, Canese K, Chan J, Comeau DC, et al. Database
resources of the national center for biotechnology information. Nucleic Acids Res.
2022;50:D20–6.
41. Clum A, Huntemann M, Bushnell B, Foster B, Foster B, Roux S, et al. DOE JGI
Metagenome Workflow. Segata N, editor. mSystems [Internet]. 2021;6:D723–33.
Available from: https://journals.asm.org/doi/10.1128/mSystems.00804-20
42. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. MetaSPAdes: A new versatile
metagenomic assembler. Genome Res. 2017;27:824–34.
43. Mikheenko A, Saveliev V, Gurevich A. MetaQUAST: evaluation of metagenome
assemblies. Bioinformatics. 2016;32:1088–90.
44. R Core Team. R: A Language and Environment for Statistical Computing [Internet].
Vienna, Austria; 2020. Available from: https://www.R-project.org/
45. Wickham H. ggplot2: Elegant Graphics for Data Analysis. New York: Springer-
Verlag; 2016.
46. Zhou Z, Martin C, Kosmopoulos JC, Anantharaman K. ViWrap: A modular pipeline to
identify, bin, classify, and predict viral–host relationships for viruses from metagenomes.
iMeta [Internet]. 2023; Available from:
https://onlinelibrary.wiley.com/doi/10.1002/imt2.118
47. Kieft K, Zhou Z, Anantharaman K. VIBRANT: automated recovery, annotation and
curation of microbial viruses, and evaluation of viral community function from genomic
sequences. Microbiome. 2020;8:90.
48. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat
Methods. 2012;9:357–9.
49. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve
years of SAMtools and BCFtools. Gigascience. 2021;10.
50. Kieft K, Adams A, Salamzade R, Kalan L, Anantharaman K. vRhyme enables
binning of viral genomes from metagenomes. Nucleic Acids Res. 2022;50:e83–e83.
51. Nayfach S, Camargo AP, Schulz F, Eloe-Fadrosh E, Roux S, Kyrpides NC. CheckV
assesses the quality and completeness of metagenome-assembled viral genomes. Nat
Biotechnol. 2021;39:578–85.
52. Bin Jang H, Bolduc B, Zablocki O, Kuhn JH, Roux S, Adriaenssens EM, et al.
Taxonomic assignment of uncultivated prokaryotic virus genomes is enabled by gene-
sharing networks. Nat Biotechnol. 2019;37:632–9.
53. Olm MR, Brown CT, Brooks B, Banfield JF. dRep: a tool for fast and accurate
genomic comparisons that enables improved genome recovery from metagenomes
through de-replication. ISME J. 2017;11:2864–8.
54. Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using
DIAMOND. Nat Methods. 2015;12:59–60.
55. O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, et al.
Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion,
and functional annotation. Nucleic Acids Res. 2016;44:D733–45.
56. Grazziotin AL, Koonin E V., Kristensen DM. Prokaryotic Virus Orthologous Groups
(pVOGs): a resource for comparative genomics and protein family annotation. Nucleic
Acids Res. 2017;45:D491–8.
57. Camargo AP, Nayfach S, Chen IMA, Palaniappan K, Ratner A, Chu K, et al. IMG/VR
v4: an expanded database of uncultivated virus genomes within a framework of

extensive functional, taxonomic, and ecological metadata. Nucleic Acids Res.
2023;51:D733–43.

58. Wilkinson L. Exact and Approximate Area-Proportional Circular Venn and Euler
Diagrams. IEEE Trans Vis Comput Graph. 2012;18:321–31.

59. Micallef L, Rodgers P. eulerAPE: Drawing Area-Proportional 3-Venn Diagrams Using
Ellipses. PLoS One. 2014;9:e101717.

60. Roux S, Matthijnssens J, Dutilh BE. Metagenomics in Virology. Encyclopedia of
Virology. Elsevier; 2021. p. 133–40.

61. Darling ACE, Mau B, Blattner FR, Perna NT. Mauve: Multiple Alignment of
Conserved Genomic Sequence With Rearrangements. Genome Res. 2004;14:1394–
403.

62. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al.
BLAST+: architecture and applications. BMC Bioinformatics. 2009;10:421.

63. Bouras G, Nepal R, Houtak G, Psaltis AJ, Wormald P-J, Vreugde S. Pharokka: a
fast scalable bacteriophage annotation tool. Bioinformatics. 2023;39.

64. Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal:
prokaryotic gene recognition and translation initiation site identification. BMC
Bioinformatics. 2010;11:119.

65. Steinegger M, Söding J. Clustering huge protein sequence sets in linear time. Nat
Commun. 2018;9:2542.

66. Van Dongen S. Graph Clustering Via a Discrete Uncoupling Process. SIAM Journal
on Matrix Analysis and Applications. 2008;30:121–41.

67. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol. 2014;15:550.

68. Terzian P, Olo Ndela E, Galiez C, Lossouarn J, Pérez Bucio RE, Mom R, et al.
PHROG: families of prokaryotic virus proteins clustered using remote homology. NAR
Genom Bioinform. 2021;3.

69. Anantharaman K, Brown CT, Hug LA, Sharon I, Castelle CJ, Probst AJ, et al.
Thousands of microbial genomes shed light on interconnected biogeochemical
processes in an aquifer system. Nat Commun. 2016;7:13219.

70. Fierer N. Embracing the unknown: disentangling the complexities of the soil
microbiome. Nat Rev Microbiol. 2017;15:579–90.

71. Sunagawa S, Acinas SG, Bork P, Bowler C, Babin M, Boss E, et al. Tara Oceans:
towards global ocean ecosystems biology. Nat Rev Microbiol. Nature Research; 2020.
p. 428–45.

72. Santos-Medellín C, Blazewicz SJ, Pett-Ridge J, Firestone MK, Emerson JB. Viral
but not bacterial community successional patterns reflect extreme turnover shortly after
rewetting dry soils. Nat Ecol Evol. 2023;

73. ter Horst AM, Santos-Medellín C, Sorensen JW, Zinke LA, Wilson RM, Johnston ER,
et al. Minnesota peat viromes reveal terrestrial and aquatic niche partitioning for local
and global viral populations. Microbiome. 2021;9.

74. Moreno-Gallego JL, Reyes A. Informative Regions In Viral Genomes. Viruses.
2021;13:1164.

75. Gao Z, Feng Y. Bacteriophage strategies for overcoming host antiviral immunity.
Front Microbiol. 2023;14.

76. Shaffer M, Borton MA, McGivern BB, Zayed AA, La Rosa SL, Solden LM, et al.
DRAM for distilling microbial metabolism to automate the curation of microbiome
function. Nucleic Acids Res. 2020;48:8883–900.
77. Moon K, Jeon JH, Kang I, Park KS, Lee K, Cha C-J, et al. Freshwater viral
metagenome reveals novel and functional phage-borne antibiotic resistance genes.
Microbiome. 2020;8:75.
78. Strange JES, Leekitcharoenphon P, Møller FD, Aarestrup FM. Metagenomics
analysis of bacteriophages and antimicrobial resistance from global urban sewage. Sci
Rep. 2021;11:1600.
79. Hurwitz BL, Sullivan MB. The Pacific Ocean Virome (POV): A Marine Viral
Metagenomic Dataset and Associated Protein Clusters for Quantitative Viral Ecology.
PLoS One. 2013;8:e57355.
80. Brum JR, Ignacio-Espinoza JC, Kim EH, Trubl G, Jones RM, Roux S, et al.
Illuminating structural proteins in viral "dark matter" with metaproteomics. Proc Natl
Acad Sci U S A. 2016;113:2436–41.
853

854 **ADDITIONAL FILES**
855 **Additional file 1. Supplementary data and tables.** Includes Tables S1-S5 as
856 referenced in the main manuscript text. File format: .xlsx.
857 **Additional file 2. Supplementary text and figures**. Includes supplementary methods,
858 supplementary results, Figures S1-S6, and associated references. File format: .docx.