Emergent effects of synaptic connectivity on the dynamics of global and local slow waves in a large-scale thalamocortical network model of the human brain

Brianna M Marsh[1,2†], **M. Gabriela Navas-Zuloaga**[1†], **Burke Q Rosen**[2], **Yury Sokolov**[1], **Jean Erik Delanois**[1,3], **Oscar C González**[1], **Giri P. Krishnan**[1], **Eric Halgren**[2,4], **Maxim Bazhenov**[1,2]

*For correspondence:
mbazhenov@ucsd.edu (MB)

[†]These authors contributed equally to this work

[1]Department of Medicine, University of California, San Diego ; [2]Neuroscience Graduate Program, University of California, San Diego ; [3]Department of Computer Science and Engineering, University of California, San Diego ; [4]Departments of Radiology and Neuroscience, University of California, San Diego

---

Abstract   Slow-wave sleep (SWS), characterized by slow oscillations (SO, <1Hz) of alternating active and silent states in the thalamocortical network, is a primary brain state during Non-Rapid Eye Movement (NREM) sleep. In the last two decades, the traditional view of SWS as a global and uniform whole-brain state has been challenged by a growing body of evidence indicating that SO can be local and can coexist with wake-like activity. However, the understanding of how global and local SO emerges from micro-scale neuron dynamics and network connectivity remains unclear. We developed a multi-scale, biophysically realistic human whole-brain thalamocortical network model capable of transitioning between the awake state and slow-wave sleep, and we investigated the role of connectivity in the spatio-temporal dynamics of sleep SO. We found that the overall strength and a relative balance between long and short-range synaptic connections determined the network state. Importantly, for a range of synaptic strengths, the model demonstrated complex mixed SO states, where periods of synchronized global slow-wave activity were intermittent with the periods of asynchronous local slow-waves. Increase of the overall synaptic strength led to synchronized global SO, while decrease of synaptic connectivity produced only local slow-waves that would not propagate beyond local area. These results were compared to human data to validate probable models of biophysically realistic SO. The model producing mixed states provided the best match to the spatial coherence profile and the functional connectivity estimated from human subjects. These findings shed light on how the spatio-temporal properties of SO emerge from local and global cortical connectivity and provide a framework for further exploring the mechanisms and functions of SWS in health and disease.

Author Summary

Slow Wave Sleep (SWS) is a primary brain state displayed during Non-Rapid Eye Movement (NREM) sleep. While previously thought of as homogenous waves of activity that sweep across the entire

brain, modern research has suggested a more nuanced pattern of activity that can vary between local and global slow wave activity. However, understanding how these states emerge from small scale neuronal dynamics and network connectivity remains unclear. We developed a biophysically realistic model of the human brain capable of generating SWS-like behavior, and investigated the role of connectivity in the spatio-temporal dynamics of these slow waves. We found that the overall strength and a relative balance between long and short-range synaptic connections determined the network behavior - specifically, models with relatively weaker long-range connectivity resulted in mixed states of global and local slow waves. These results were compared to human data, and we found that models producing mixed states provided the best match to the network behavior and functional connectivity of human subject data. These findings shed light on how the spatio-temporal properties of SWS emerge from local and global cortical connectivity and provide a framework for further exploring the mechanisms and functions of SWS in health and disease.

Introduction

Internal states of the brain rapidly and profoundly influence sensation, cognition, emotion, and action (*Lee and Dan, 2012*; *Pfaff et al., 2008*; *Steriade and McCarley, 2013*). Dramatic change of internal brain state occurs at the transition between wakefulness and sleep (*Campbell and Tobler, 1984*; *Leung and Mourrain, 2018*; *Steriade and McCarley, 2013*). The main electrophysiological hallmarks of slow wave sleep (SWS) and rapid eye movement (REM) sleep are shared across mammals. During the SWS, the brain dynamics is dominated by oscillations between an active (Up) and a silent (Down) states, the activity called slow oscillation (SO, <1Hz) (*Steriade et al., 1993a*,b). Alterations of these sleep dynamics have significant consequences for brain function and are at the core of many neuropsychiatric disorders (*Winkelman and Lecea, 2020*).

Until recently, sleep/wake transitions or transitions between different stages of sleep were thought to occur uniformly throughout the cortex. Recent evidence in rodents, primates and humans points to the existence of electrographic events resembling local sleep during behavioral wakefulness (*Vyazovskiy et al., 2011*; *Hung et al., 2013*; *Quercia et al., 2018*; *Engel et al., 2016*; *van Kempen et al., 2021*). During wakefulness, local groups of cortical neurons tend to fall silent for brief periods, as they do during SWS. Similarly, during behavioral sleep, transient local wake-like activity can intrude SWS (*Nobili et al., 2011*; *Peter-Derex et al., 2015*). SWS and REM sleep can also coexist in different cortical areas (*Soltani et al., 2019*; *Funk et al., 2016*; *Bernardi et al., 2018*). Homeostatic sleep pressure can affect cortical activity locally (*Siclari and Tononi, 2017*). Spindles and slow waves, may increase locally after learning (*Eschenko et al., 2006*; *Pugin et al., 2015*), suggesting experience dependent regulation. Together these results suggest an intriguing idea that local slow-waves may have a functional significance enabling brain to process and learn in a parallel way. However, the underlying mechanisms of local vs global slow-wave dynamics as well as factors controlling transitions between them are unknown.

We previously developed computer models of sleep SO (*Timofeev et al., 2000*; *Bazhenov et al., 2002*), including transitions between awake and sleep states (*Krishnan et al., 2016*). However, the highly detailed and computationally intensive nature of these models have prevented up-scaling to investigate slow wave dynamics at the whole-brain scale. Recently, several whole-brain models of slow-wave sleep incorporating biologically grounded connectivity from probabilistic diffusion MRI tractography and providing a good fit to human sleep recordings were proposed (*Cakan et al., 2022*; *Goldman et al., 2023*). These models investigated the role of long-range connections in SO brain-wide propagation. However, the mean-field neural-mass representation of cortical regions in these models provides only limited insights on how local (within a few milimeters) and long-range connectivity and activity interact to affect initiation and propagation of sleep slow waves.

To bridge this gap between micro and macro-scale mechanisms, we have developed a multi-scale, whole-brain, thalamocortical network model with biologically grounded cortical connectivity that exhibits the essential activity states of wake and slow wave sleep. The model consists of 10,242

cortical columns per hemishpere, each containing spiking pyramidal (PY) and inhibitory (IN) neurons arrayed in 6 layers, and 642 thalamic cells of each of four types (thalamo-cortical (TC) and reticular (RE), belonging to either core or matrix). To investigate the role of connectivity in emerging SO, we systematically modified the density, range, and relative strength of cortical connections, as well as distance-dependent synaptic delays. The model dynamics were compared to the spatial coherence profiles obtained from human subjects. Our study revealed that a characteristic balance of long and short-range connectivity is necessary to match the data and enable the emergence of complex mixed states of local and global SO (*Nir et al., 2011*; *Bernardi et al., 2018*; *Vyazovskiy et al., 2009*). It predicts, that changes in synaptic weights, which likely occur over the course of sleep (*Timofeev and Chauvette, 2017*), may enable transitions between global synchronized and local SO.

Results

Connectivity

We start with a brief overview of the model (Figure 1, see Methods for details). The model includes 10,242 cortical columns positioned uniformly across the surface of one hemisphere, corresponding to the vertices in the ico5 cortical reconstruction reported in (*Rosen et al., 2019*). The medial wall includes 870 of these vertices, so all analyses for this one-hemisphere model were done on the remaining 9372 columns. Each column has 6 cortical layers (L2, L3, L4, L5a, L5b and L6), with 1 pyramidal cell and 1 inhibitory cell per layer. Cortical neurons are modeled using computationally efficient map-based (*Komarov et al., 2018*) spiking PY and IN neurons. Intra-columnar connections follow the canonical cortical circuit (Figure 1A), while long-range cortical connectivity (Figure 1B) is based on diffusion MRI (dMRI) tractography (*Rosen and Halgren, 2021*) of the Human Connectome Project (HPC) young adult dataset (*Van Essen et al., 2013*), organized into the 180 parcels of the HCP multimodal atlas (*Glasser et al., 2016*)). The originating and terminating layers of long range connections are assigned according to the parcel's relative hierarchical positions (Figure 1D-E). Column-wise connectivity is obtained by applying the parcel-wise relation between connection length and probability to synthesized intercolumnar distances (Figure 1C, see Diffusion MRI Guided Connection Probability for details). The resulting connectivity (Figure 1G) retains the parcel-wise structure from the data (*Rosen and Halgren, 2021*) (Figure 1F, Pearson's correlation r = 0.81, p < 0.0001), with an approximately exponential distribution between connection frequency and length (see Figure 1C). The connection delays, which are proportional to the geodesic connection lengths, follow a similar distribution. The thalamus is simulated with layers of thalamocortical (TC) and reticular thalamic (RE) neurons (*Bazhenov et al., 1998a*,b, *2002*), and consists of the core and the matrix (*Bonjean et al., 2011*, *2012*), which selectively project to the neurons from different cortical layers satisfying the outlined cortico-thalamo-cortical loop (*Shepherd and Yamawaki, 2021*). Following (*Krishnan et al., 2016*) the model included the effects of neuromodulators to induce transitions between Wake state and SWS.

In the following sections, we first discuss "baseline" awake and sleep state dynamics of the model. Next we discuss how sleep dynamics are affected by altering network connectivity and reveal conditions for coexistence of local and global SO. We finally present analysis of human recordings and compare different model dynamics to human data using coherence analysis across frequency and distance and functional connectivity analysis. In addition to the basic characteristics of SO frequency, amplitude, and propagation speed, network effects are primarily described in terms of 3 key metrics: *synchrony*, *spread*, and *participation*. Synchrony refers to the timing of the Up state onsets and offsets across all neurons participating in a slow wave. Spread refers to how far the slow wave travels across the cortex. Finally, participation refers to the percent of neurons that participate in a slow wave within the area of spread.

Figure 1. Model architecture. A) Connectivity between neurons in the same column follows the canonical circuit. Arrows indicate directed connections between layers. Neurons are never connected to themselves. B) Connectivity diagram between different cortical columns and thalamocortical cells in the core (TC) or matrix (TCa). C) Exponential decay of connection probability with distance between columns. Previously reported inter-parcel dMRI connectivity (*Rosen and Halgren, 2021*) shows an exponential relation with fiber distance. Column-wise connection probability is obtained by applying the dMRI exponential function to inter-column fiber distances (synthesized from their corresponding geodesic distances) and then adding the residual inter-parcel dMRI connectivity, estimated by regressing out the exponential trend (see Diffusion MRI Guided Connection Probability for details). D) Estimated myelination index (*Glasser and Essen, 2011*; *Burt et al., 2018*) throughout the cortex. A hierarchical index inversely proportional to this myelination index is assigned to each of the 180 cortical parcels designated in the HCP-MMP atlas (*Glasser et al., 2016*). E) Excitatory corticocortical connections belong to one of 6 classes based on the hierarchical index of the pre- and post-synaptic neurons: within the same column, within the same parcel, weakly or strongly feedforward (from lower to higher hierarchical index) and weakly or strongly feedback (from higher to lower hierarchical index). Based on experimental reports on connectivity (*Markov et al., 2013*; *Rockland, 2019*) weights are scaled according to the connection type by the factor shown in the matrices. F) Parcel-wise connectivity from *Rosen and Halgren* (*2021*). G) Generated model connectivity, which retains the parcel-wise structure from data in panel F (Pearson's correlation r = 0.81, p < 0.0001).

Wake State Characterization

Wake state (Figure 2A) was achieved by applying biologically relevant changes in key parameter values (Table 4), based on our previous work (*Krishnan et al., 2016*). The resultant network shows a consistent average firing rate of individual neurons at about 17 Hz. Given the broad distribution of firing rates reported in awake state (*Miyawaki et al., 2019*; *Thomas et al., 2020*), we believe this model is appropriate. There is indeed experimental evidence of average firing rates in awake state being 15 Hz (*Vyazovskiy et al., 2009*), 20-50 Hz (*Thomas et al., 2020*), or 5-16 Hz (*McKillop et al., 2018*). However, it should be noted that different mean firing rates can be achieved with minimal adjustments to key parameters, namely AMPA and GABA strength. The power spectral density (Figure 2a.5) revealed a distinct $1/f$ phenomenon, as is common for in vivo recordings in awake state (*Donoghue et al., 2020*). Individual neurons showed irregular tonic firing (Figure 2a.1), with baseline membrane voltage around -60 mV (Figure 2a.4). The firing activity was not synchronized (Figure 2a.3), which was reflected in the low voltage LFP (Figure 2a.2). Overall, the network revealed properties representative of biological awake state activity.

Slow Wave Characterization

To attain a network displaying SWS-like activity from the wake state baseline, several cellular and neurochemical parameters were adjusted according to the methodology presented in (*Krishnan et al., 2016*). Namely, we increased cortical AMPA and GABA conductances as well as the leak current and slow hyperpolarizing currents in all cortical neurons (see Methods, *Table 4*). The resulting baseline model of SWS (Figure 2B) revealed synchronized and regular oscillations (~0.27 Hz) between silent (Down) and active (Up) states (Figure 2b.2) in all cortical excitatory and inhibitory neurons (Figure 2b.1). This network dynamics was organized as the waves of spiking activity propagating across the entire cortex - slow-waves. (This can be seen in Supplemental Video S1.) The slow-waves were further highly synchronized across layers (Supplementary Figure S1). Differences between layers include a relatively less active L4. This is probably due to L4 receiving within-column connections from only L6, while other layers have inputs from two or more other layers (see Figure 1E).

In general, the network spent more time in the Down states, and the Up state onsets and offsets were well synchronized (see narrow onset and offset histograms in Figure 2b.5). This pattern of longer Down states is typical for recordings under urethane anesthesia (*Steriade et al., 1993a*), while Down states may be shorter in non-anesthetized animals (*Steriade et al., 2001*). Nevertheless, the frequency of SOs was within the range of human data (*Csercsa et al., 2010*). The latency maps in Figure 2D demonstrate how waves of activity travel from a single initiation site, which varied between Up states with no detectable dependency on the previous ones. Each Up state involved >97% of all excitatory neurons. In this baseline model, slow waves revealed both high spread and high participation.

Cortical inhibitory, thalamocortical, and reticular thalamic neurons all showed similar patterns of oscillations (Supplementary Figure S2), as previously shown in *Bazhenov et al.* (*2002*). While thalamic cells followed cortical oscillations, they were not directly involved in the generation of SO. In fact, simulations with no thalamic inputs to the cortex maintained cortical SO (Supplementary Figure S5). This is in line with experimental results showing that SO survives extensive thalamic lesions (*Steriade et al., 1993a*), and is present in neocortical slices (*Sanchez-Vives and McCormick, 2000*) and isolated cortical slabs (*Timofeev et al., 2000*). Nevertheless, removing the thalamus reduced synchrony of cortical slow waves (see Supplementary Information: Effect of thalamus on SO synchronization), in agreement with in vivo data (*Lemieux et al., 2014*).

Figure 2. Baseline model activity in awake and sleep states. A) Wake state. a.1) Membrane voltages of all neurons from layer 2 over 5 seconds of simulation; a.2) Layer average voltage over time, simulating LFP; a.3) Activity of two representative neurons from layer 2, both showing irregular tonic firing; a.4) Voltage histogram of all neurons over the whole simulation time, note approximately -60mV peak representing baseline membrane voltage; a.5) Power spectral density (PSD) of the average voltage revealed a distinct $1/f$ phenomenon typical for *in-vivo* recordings. B) Slow-wave dynamics. b.1) Membrane voltages of all neurons (excluding medial wall) from layer 2 over 30 seconds of simulation, revealed synchronized bands of activity during Up states; b.2) Average voltage of all cortical layers (*dashed black line*) and layer 2 neurons (*solid blue line*, excluding medial wall) over time, with nearly identical behavior. *Red triangles* above and below the trace mark global Up and Down states, respectively, from layer 2 (coincident with global Up and Down states from the average of layers); a.3) Activity of two representative neurons from layer 2, both showing synchronized transitions between Up and Down states; a.4) Voltage histogram of all neurons over the whole simulation time, revealed the characteristic bi-modal distribution caused by Up and Down state alternations during SWS. Dashed vertical lines labeled V^- and V^+ indicate the voltage thresholds used to detect Down states and Up states, respectively; b.5) Distribution of the Up state onsets and offsets for all neurons over the whole simulation time. Narrow histograms indicate highly synchronous initiation and termination of the Up states. C) Zoom into the first Up state from panel b.1, with neurons sorted from earliest to latest onset time, and a single cell voltage from panel b.2, showing the transition from Down to Up state, steady firing during the Up state, and transition to Down state. D) Latency map, calculated as the onset delay of each neuron with respect to the earliest onset, for each Up state in panel b.2 (see Onset/offset Detection in Methods for details). The percent of active neurons during each Up state is shown below the corresponding latency map. Up states involve nearly every neuron in the cortex within 300ms from its initiation time.

Connection Density

In the next two sections, we test effects of alternating the intracortical connection density and the maximal radius of connections on SO properties. Importantly, in these simulations we scaled the strengths of remaining synaptic connections to compensate for the lost connections and to ensure an equivalent amount of net activation per neuron (see Modifying Network Structure for details).

In the baseline model, synaptic connectivity was set to match biological data (see Methods). Next, to test the effects of connection density between regions, we gradually changed the density of connections, by applying a probability P to retain each possible connection (Figure 3A-B) regardless of the distance between neurons, thus keeping the connection distance distribution unchanged (Figure 3C). Remaining synapses were scaled to keep total synaptic input per neuron unchanged.

Setting the probability from 100% down to 50% (i.e., ablating a random half of all connections) had minimal effect on the slow waves, indicating the model's robustness to even significant loss of individual connections as long as the total strength of excitatory synaptic input per neuron remains unchanged. Slow waves exhibited a similar propagation pattern with only a slightly reduced amplitude (approx. 92% of baseline SOs) and speed (Figure 3D, F). The synchrony of the Up state offsets, however, was impaired; the distribution of offset timings across neurons became much wider (Figure 3d.5, f.5) compared to baseline model (Figure 2b.5). Further decreasing density below 30% caused a considerable drop in amplitude and participation (Figure 3f.1, f.4). In the extreme case of only 10% of connections remaining ($P = 0.1$, Figure 3E) (see Supplementary Information: Effects of connections density, range and delays and Figure S6), there was continued loss of synchrony in Up state offsets and onsets, slow waves became less frequent (down to 0.17 Hz), with lower amplitude (50% of baseline), and lower participation (only 70% of neurons involved per Up state) (Figure 3F). Nevertheless, slow waves still propagated through the all brain regions on the macro scale, showing that long range synchrony is still possible (if slightly impaired) with only very few connections in place as long as those connections are sufficiently strong and long-range connections are present. Below this level of remaining connections, slow waves no longer occurred and the network was largely silent.

Connection Range

To determine the specific contribution of long-range connections, we varied the maximum connection radius; any connections longer than a specified distance R were set to 0 (Figure 4A). Again, the strengths of remaining connections were scaled up to compensate for reduction in the total number of connections and to keep total synaptic input per cell unchanged (see Modifying Network Structure for details). In contrast to the density reduction (see Figure 3C), this alteration did shift the connection distance distribution (Figure 4C).

The maximum connection length in baseline model was 226.1 mm, but most connections (approximately 80%) were maintained within 100 mm (Figure 4B). A connection radius as small as 5 mm, which includes second-order neighbors of a given neuron, resulted in only approximately 10% of connections still intact. The alterations of the maximum connection length led to significantly different activity patterns than in the network with a spatially uniform drop to the same percentage of connections (Connection Density). The network with a maximum connection radius of 5 mm (Figure 4D) showed no reduction in the frequency of the slow waves, but both the onsets and offsets of the slow waves revealed dramatic reduction in synchrony. Additionally, there was a substantial drop in slow wave propagation speed, while changes in connection density had minimal effect on speed (compare Figure 4F and Figure 3F). Despite all this, characteristic properties of global slow waves (high spread and high participation) found in the baseline model were maintained.

However, reducing radius below 5mm (i.e., not including 2nd degree neighbors) caused a dramatic drop in global oscillation amplitude, akin to a phase transition from global to local oscillations. With a connection radius of 2.5 mm (Figure 4E), including only first order neighbors of each cell, global slow waves were lost in favor of local slow waves (Figure 5, low spread, high participation). Note that this radius retains only 3.7% of connections, where a spatially uniform drop to any

Figure 3. Effect of connection density on SO dynamics. A) Example target cortical column (*yellow*) with all columns connected to it through any layer (*green*) for different connection densities. Density is reduced by decreasing P, which denotes the probability of preserving a connection from the original dMRI-based connectivity. With $P = 1$ all connections are preserved, while for $P = 0.5$, $P = 0.3$ and $P = 0.1$ each connection is preserved with a 50%, 30% or 10% probability, respectively. B) Percent of the original number of connections retained for different values of P. C) Distribution of connection distances, or lengths, for different values of P. Red horizontal lines indicate medians, bottom and top box edges indicate the 25th and 75th percentiles, respectively, whiskers extend to the most extreme data points not considered outliers, and outliers are plotted individually using the '+' marker symbol. Note, connection density is reduced uniformly across all lengths. D-E) Individual analysis of SO dynamics for different values of P. Subpanels as in Figure 2. F) Summary of the effect of reducing connection density on the global SO frequency and amplitude as well as the standard deviation of the onset/offset delays (i.e. the width of the onset/offset histograms in d.5 / e.5)

Figure 4. Effect of connection range on SO dynamics. A) Example target column (*yellow*) with all its connected columns (*green*). The *blue* area indicates the connection range for the corresponding radius R. No connections are made in the *black* region outside the radius. $R = 226mm$ encompasses all the original dMRI-derived connections. B) Percent of the original number of connections preserved for different values of R. C) Distance distribution for different values of R, with *inset* zooming into radii below 10mm. R imposes a maximum connection length and truncates the distance distribution accordingly. Red horizontal lines indicate medians, bottom and top box edges indicate the 25th and 75th percentiles, respectively, whiskers extend to the most extreme data points not considered outliers, and outliers are plotted individually using the '+' marker symbol. D-E) SO analysis for $R = 5mm$ and $R = 2.5mm$. Subpanels as in Figure 2. F) Summary of the effect of reducing connection density on the global SO frequency and amplitude as well as the standard deviation of the onset/offset delays (i.e. the width of the onset/offset histograms in d.5 / e.5)

Figure 5. Local SO for small connection radius ($R = 2.5$mm). A) Ten cortical areas with a 5mm radius, that were used to calculate local dynamics. B) Average membrane voltage of layer II neurons, as in Figure 4e.2. C-E) For each region in (A), subpanels show (C) the single-cell voltage for two neurons in the area, (D) the local field potential (LFP) for the 5mm area, and (E) heatmap of individual voltages of all neurons in the area. Note that even within these very small regions, there are still subgroups of independently synchronized neurons. F) Latency map, calculated as the onset delay of each neuron with respect to the earliest onset, for each Up state identified in Figure 4e.2. Note that even the most global of slow waves has very low participation, around 35%, and can be seen to consist of many extremely small initiation sites that generally do not spread.

less than 10% of connections removed slow waves entirely. The local slow waves were generated
independently in few selected areas, and occasionally showed regional propagation but never trav-
elled across the full hemisphere (Figure 5F). This resulted in reduced global SO amplitude as long-
range synchrony between locally oscillating neural populations was impaired (Figure 5C-D). This
model can further be seen in Supplemental Video S3. While a large fraction of the cortex was not

participating in SO, the active regions (such as region 6, Figure 5) were consistent across Up states.
A connectivity analysis revealed that these regions correspond to the network areas with the high-
est node degree (see Supplementary Figure S9). The same regions exhibited local slow waves for
different simulations with the same connectivity parameters but different random seeds for net-
work generation, implying that these regions are not random but emerge from the underlying
structure imposed by the dMRI connectivity data.

As a whole, this analysis revealed that long range connections (longer than 5 mm) are essential
for the large-scale synchrony of the slow waves, but having only local connections (less than 5
mm) is still sufficient to make slow waves propagating over entire cortex (as long as remaining
connections strength is scaled up to maintain total synaptic input per cell). Even with only first
order neighbors, the network still maintained slow wave initiation, synchrony, and participation
with local propagation.

Connection Delay

The model implements synaptic delays scaled by geodesic distance between neurons with a maxi-
mum delay of 40 ms. The delay for the majority of synapses is under 2 ms because the vast majority
of connections are relatively short range. To test the effects of delay times, we changed the delays
in two ways (Figure S7): 1) by changing the maximum delay and scaling all others proportionally,
and 2) by setting one uniform delay to all connections.

Setting the maximum delay to be smaller (all the way down to 0.1 ms) had little to no effect
on slow wave amplitude, frequency, or synchrony. Setting the maximum delay to be larger also
had minimal effect on network activity until biologically implausible delays were implemented (≥ 1
second). Even at this extreme the network retained strong global slow waves, albeit with slightly
lower amplitude, longer frequency, and de-synchronized onsets and offsets. Setting a uniform
delay of 10 ms (where most delays were previously under 2 ms) further left the network behavior
largely unchanged. Only when the uniform delay was increased to 30 ms did we observe a loss
of amplitude and synchrony (Figure S7). Since the delay times here shown to have an effect on
network behavior are clearly beyond the range of biological plausibility, these results indicate that
synaptic delays are not a primary determinant of slow wave spatio-temporal properties.

Connection Strength

To probe the effects of absolute connection strength on the network activity, we varied the base
connection strength between all cortical pyramidal neurons (Figure 6A, C). In contrast to exper-
iments with changing connectivity density or radius, where remaining synapses were scaled to
keep total synaptic input constant, this changed the total synaptic input per cell. Weight reduc-
tions or increments by a factor of 4 or more caused the slow oscillations to be barely detectable or
undetectable (note the low amplitude at the extremes of Figure 6c.1). This was a result of low ac-
tivity in the case of weights reduction, and of sustained Up state (Down state absence) in the case
of weights increase. In general, within the detectable range, the SO amplitude, frequency, and
participation decreased when synaptic strength was decreased (Figure 6C). Note that the model
revealed an overall robustness to weight variation: the slow-wave propagation was minimally al-
tered by strength increases or decreases by a factor of 2.

To further investigate the role of long range connectivity, we next performed N times synaptic
weight reductions on only "long-range connections", i.e., connections over a defined minimum
radius R_{th} (Figure 6B,D). We varied the scaling factor N from 2 to 7 and radius threshold R_{th} from
2.5 mm to 50 mm. (Figure 6B illustrates the case N=5, R_{th}=10mm). Again, we did not scale the
total synaptic input per cell, so these alternations affected balance of excitation and inhibition in
the network. This experiment can be interpreted to model the effect of stronger synaptic inputs
nearest to the cell body (*Hawkins and Ahmad, 2016*; *London and Segev, 2001*).

When connections over R_{th}=10mm were significantly weakened (N=5) (Figure 6B, Figure 7), the
model displayed a number of clearly identifiable local slow waves originating and reliably travel-

Figure 6. Effect of cortical excitatory synaptic strength on the network dynamics. A) Reduction of all cortical connections by 3x. B) Reduction of only connections longer than 10 mm by 5x. All subpanels as in Figure 2. C) Summary of changes in frequency, amplitude, speed, participation, and onset/offset distribution spread when all connections are reduced as shown in (A). Increasing all weights shows little effect while decreasing all weights shows decreased amplitude and participation, with increases in onset/offset distributions (decreases in synchrony). The frequency of slow waves also becomes more variable. Note, Increasing all weights 5x results in the ablation of slow waves due to constant Up state, i.e., SO characteristics cannot be meaningfully quantified. D) Summary of changes in SO characteristics when only long range connections are reduced as shown in (B). Across all plots, 5 mm is seen to be an inflection point (or "elbow") where network activity changes.

ing around the hemisphere in different directions (Figure 7F). Occasionally, these local slow waves would coincide to produce a global slow wave in the LFP (see Figure 6b.2, where red downward triangles mark the four detectable global UP states analyzed in Figure 7F). In this model, the local populations of neurons would regularly synchronize producing local slow waves but the global synchrony was lost in many cases except for a subset of all events when global slow waves were generated (Figure 7B, E). This was further confirmed by running longer 150 sec simulation (see Supplemental Video S4) that revealed multiple transitions between local and global synchrony states. With reduced spread and high participation, the local slow wave behavior resembles the results from simulations with exclusively local connectivity (see Connection Range), but stems from the more biologically plausible assumption of weaker (rather than nonexistent) long range connectivity. Indeed, the propagation patterns in Figure 7 is similar to the proposed pattern of biological sleep including a mix of Type I (global) and Type II (local) slow waves, as suggested by (*Siclari et al., 2014*; *Nir et al., 2011*; *Bernardi et al., 2018*; *Funk et al., 2016*; *Vyazovskiy et al., 2007*, *2011*; *Riedner et al., 2007*).

Importantly, not all the cortical regions were equally likely to generate local slow waves. As in simulations with only local connections, the network areas with the highest node degree (see Supplementary Figure S9) produced almost regular local slow-waves (e.g., region 6, (Figure 7)), while some other (e.g., region 9, (Figure 7)) displayed low frequency irregular spiking with only rare Up state-like events. This again suggests that the active, i.e., generating local SO, areas emerge from the underlying structure imposed by the dMRI-based connectivity.

Local vs Global Slow-Waves

Given above analysis, we next sought to quantify the global v.s. local slow waves in the model as a function of synaptic strength. For each model, we calculated the regularity in each of the 180 parcels, where regularity is defined as the inverse of approximate entropy. The mean and variance of this distribution are then plotted against each other, as shown in Figure 8A. We define global models as models with high regularity, local models as those with low regularity, and mixed states falling in between. The Intact Score of each model, defined as the percentage of remaining synaptic strength compared to the base model, is further predictive of the regularity and network behavior (Figure 8B). Two groups of models in the middle of distribution with Intact Score 30-50% and regularity 4-12, were identified as mixed states models. In Figure 8C, ten random parcel LFPs from representative global, mixed, and local models show demonstrably different behavior - the global model parcels are all in synchrony, while the local model parcels can be seen to transit into Up states largely independently. The mixed model shows a combination of these behaviors, i.e., transitions to Up state sometimes occurred independently and sometimes in sync across parcels. This group predominately includes models with long-range connectivity scaled down beyond certain radius.

Coherence Analysis

To validate our model, we conducted a comparative analysis with experimental data. Specifically, we analyzed coherence across 180 cortical parcels. We calculated coherence in successive narrow-band signals across all parcel pairs to generate a snapshot of coherence across both frequency and distance. We then took the average of the coherence matrices in the SO range (<2 Hz), scaled the values by parcel-to-parcel distances, and masked the model data to match the sparsity of the experimental data. These modified 2D coherence matrices were then used to calculate 2 metrics of functional connectivity: Percent connected and number of communities. Percent connected reports the percentage of parcel pairs with a coherence greater than a given threshold, while the number of communities is determined using Louvain community detection.

Subsequently, we fit the full coherence matrices (across all distances and frequencies) with an exponential model. From this model, we estimated the spatial length constant (λ) and the scaling term (α). These four metrics were used to compare the functional connectivity and coherence

Figure 7. Mixed global (Type I) and local (Type II) Slow Waves: Connections greater than 10mm reduced 5-fold. A) Ten cortical areas with a 5mm radius, that were used to calculate local dynamics. B) Average membrane voltage of layer II neurons. C-E) For each region in (A), subpanels show (C) the single-cell voltage for two neurons in the area, (D) the local field potential (LFP) for the 5mm area, and (E) heatmap of individual voltages of all neurons in the area. This demonstrates that while there is some alignment, up states are not strongly global. F) Latency map, calculated as the onset delay of each neuron with respect to the earliest onset, for each Up state. Note, that while some Up states are global (characteristic examples are marked by blue lines), others are comprised of several independent but coinciding local events (examples are marked by yellow lines).

Figure 8. Quantification of Global vs Local Slow Waves. The labels show the radius and factor of strength reduction in each model - e.g., 5/4 indicates a model where connections longer than 5 mm are reduced by a factor of 4. A) For each model, the Regularity of each parcel is calculated; the mean and variance of these distributions predict the overall network behavior, with global models showing high Regularity and local models showing low Regularity. The group of models in the middle of the distribution, boxed in orange, represents Mixed states. B) Mean Regularity is plotted against the Intact Score (total percent of synaptic strength relative to the global base model). Note that the total level of synaptic strength is itself predictive of network behavior - the same mixed models fall in the orange box in the middle of the distribution. C) Example traces of 10 random parcels in representative Global, Mixed, and Local models show different levels of synchrony across the brain.

³³² profiles of different biophysical models against in vivo data. Full details of coherence analysis can
³³³ be found in the Methods under Coherence Analysis.

Empirical coherence

³³⁵ The experimental data collection is fully described in Experimental Data. Briefly, continuous stereo-
³³⁶ electroencephalography telemetry data were collected for 236 patients with focal epilepsy. Af-
³³⁷ ter significant data cleaning to ensure removal of epileptic activity and subsequent sleep scoring,
³³⁸ NREM sleep data were selected for analysis. In Figure 9, the yellow star indicates the functional
³³⁹ connectivity metrics (Figure 9A) and the values of λ and α obtained by fitting an exponential model
³⁴⁰ (Figure 9B), based on in vivo data.

Model coherence

³⁴² To perform a comparable model analysis, we first averaged all the cell voltages per parcel (all layers,
³⁴³ excitatory and inhibitory neurons) and added noise with signal-to-noise ratio (SNR) = 5 to mimic
³⁴⁴ the signal that would be recorded by a depth electrode, as in experimental data (***Ball et al., 2009***;
³⁴⁵ ***Suarez-Perez et al., 2018***). We then performed coherence analysis in an identical manner to the
³⁴⁶ experimental data. After the full coherence matrices were computed, they were further masked to
³⁴⁷ fit the sparsity of the experimental data before metrics of functional connectivity were computed.

³⁴⁸ The functional connectivity metrics reported the total percent of connected parcels, and the
³⁴⁹ number of communities as determined by Louvain community detection (Figure 9A). Global mod-
³⁵⁰ els were highly connected, showing a low number of distinct communities. Conversely, local mod-
³⁵¹ els showed very sparse connectivity, with multiple distinct communities detected. Mixed models
³⁵² showed moderate values on both scales. The experimental data point can be seen to fall in the
³⁵³ middle of the distribution, surrounded by mixed models.

³⁵⁴ We then compared the exponential fits of the full coherence landscapes in the SO range (less
³⁵⁵ than 2 Hz) using the mean α and the first λ value (at 0.5 Hz); these two describe the peak coher-
³⁵⁶ ence and rate of decay. When plotted against each other (Figure 9B), all the mixed state models
³⁵⁷ revealed intermediate α and low λ values. The experimental data point can be seen to fall in close
³⁵⁸ proximity to the mixed state models. In particular, models in which connections longer than 10
³⁵⁹ mm were scaled down by a factor of 5, 6, or 7 were consistently very close to the experimental
³⁶⁰ data point; this is also true for the model where connections longer than 5 mm are scaled down by
³⁶¹ a factor of 4 (Figure 9A.2,B.2). We found that the global slow wave model, that was initially set as a
³⁶² baseline model, was very far off. Note that different metrics identified slightly different variations
³⁶³ of the models as the best match to the data (compare identified regions in Figure 8 and Figure
³⁶⁴ 9); nevertheless, we found high overlap between different measurements, all indicating that the
³⁶⁵ models generating mixed states are in good agreement with in vivo data.

³⁶⁶ Figure 9C,D,E shows coherence profiles obtained for in vivo data, selected mixed states model
³⁶⁷ (model with a greater than 10 mm reduction, reduced 5-fold) and baseline global state model. Both
³⁶⁸ the experimental data and the model displaying mixed local/global slow waves exhibit a sharp peak
³⁶⁹ of coherence at low frequencies, which quickly tapers off with increased distances between parcels
³⁷⁰ (Figure 9C,D). Notably, this pattern is not observed in the global slow wave baseline model (Figure
³⁷¹ 9E), where only a slight drop in coherence is seen as the distance between parcels increases. This
³⁷² is expected, given the unrealistic levels of slow wave synchrony across the entire network in this
³⁷³ model. Additionally, the global slow wave model shows very high levels of coherence (nearing 1,
³⁷⁴ indicating perfect coherence) at low frequencies.

³⁷⁵ In summary, these results suggest that mixed states models, particularly those with a 5-10 mm
³⁷⁶ radius of stronger local connections, are the best match for experimental data.

Figure 9. Functional Connectivity and Coherence Analysis of Experimental v.s. Model data. Coherence analysis was performed on all data sets according to the procedure described in Coherence Analysis. A) The parcel-to-parcel coherence matrices are first averaged in the slow wave frequency band and scaled by parcel-to-parcel distances to determine functional connectivity via percent of connected parcels and number of communities (determined by Louvain community detection). B) Full (unaveraged, unscaled) coherence matrices are then fitted with an exponential function to determine the full shape of the coherence landscape across distance and frequency. The resultant first (0.5 Hz) Lambda and mean Alpha in the slow wave frequency range (<2 Hz) are taken to describe the shape of the coherence landscape, and plotted in (B). (A.1 and B.1) show zoom-ins of each respective plot; the labels show the radius and factor of strength reduction - e.g., 10/5 indicates a model where connections longer than 10 mm are reduced by a factor of 5. This shows that the models with primarily mixed slow waves are the closest fit to experimental results across all 4 metrics. Full 3D coherence plots across distance and frequency are shown for the experimental data (B), the 10mm / 5 model (C), and the global slow wave base model (D), showing that the global slow wave model has a fundamentally different shape (lacking a dependence of coherence on distance) and much higher levels of coherence at low frequencies

Discussion

Traditionally, brain states have been categorized into wakefulness, slow-wave sleep (SWS), and rapid eye movement (REM) sleep, assuming their global occurrence across brain regions. How-

380 ever, recent large-scale neural recordings have revealed the rich heterogeneity of neural dynam-
381 ics on a brain-wide scale, demonstrating that sleep and wake signatures can coexist in different
382 brain regions, influencing behavior with spatial and temporal specificity. While local circuit mech-
383 anisms of slow waves — alternations of positive/negative EEG/LFP waves, also called slow oscilla-
384 tion (SO) — have been well described, the factors leading to complex heterogeneous brain states
385 with local slow waves remain unknown. Here, we present a whole-brain thalamocortical network
386 model based on human dMRI connectivity data capable of generating awake states and sleep states
387 with biologically realistic cortical slow waves. The model was applied to make specific predictions
388 about the effects of long- and short-range connectivity on the spatio-temporal dynamics of the
389 slow waves. We identified intracortical synaptic connectivity properties that allow the coexistence
390 of local and global slow waves and proposed mechanisms for transitions between local sleep and
391 global uniform sleep states. The model predictions were compared to human data to validate and
392 fine-tune the models.

393 A prominent example of brain states heterogeneity is local sleep (*Vyazovskiy et al., 2011*; *Hung*
394 *et al., 2013*; *Quercia et al., 2018*; *Engel et al., 2016*; *van Kempen et al., 2021*). During wakefulness,
395 local groups of cortical neurons tend to fall silent for brief periods, as they do during SWS. Further-
396 more, during behavioral sleep, transient local wake-like activity can intrude SWS (*Nobili et al., 2011*;
397 *Peter-Derex et al., 2015*). It is worth noting that SWS and REM sleep can also coexist in different
398 cortical areas (*Soltani et al., 2019*; *Funk et al., 2016*; *Bernardi et al., 2018*). Additionally, the pro-
399 portion of time allocated to wakefulness, SWS, and REM states varies significantly across cortical
400 areas (*Soltani et al., 2019*; *Nazari et al., 2023*), highlighting the diverse predisposition of different
401 regions to support these states.

402 From a mechanistic perspective, the basic need for local sleep can possibly be explained by
403 homeostatic sleep pressure, that accumulates throughout the day, shaped by the duration and
404 quality of preceding alertness. This pressure is evident in the intensity of SWS in the cortex, indi-
405 cating the brain's need for recuperation (*Wilckens et al., 2018*). Homeostatic sleep pressure can
406 influence cortical functions in specific areas (*Siclari and Tononi, 2017*). For example, spindles and
407 slow waves may intensify in regions engaged in learning (*Eschenko et al., 2006*; *Pugin et al., 2015*),
408 suggesting that sleep pressure is regulated based on recent cognitive activities. Notably, after mo-
409 tor skill training, there can be a localized increase in SWS within the motor cortex during rest (*Huber*
410 *et al., 2004*; *Korman et al., 2007*; *Hanlon et al., 2009*), and a visual perception learning task can lead
411 to a rise in the initiation of slow waves in the lateral occipital cortex (*Mascetti et al., 2013*).

412 Why brain areas that are more actively involved in sensory processing during the day require
413 more SWS is a challenging question. We can speculate that it may be related to the well-characterized
414 function of SWS in memory consolidation (*Rasch and Born, 2013*). The slow oscillation (SO) repeat-
415 edly resets the thalamocortical network during the Down phase and temporally groups sleep spin-
416 dles, which are associated with long-term potentiation (LTP)-like processes and learning, during
417 the Up phase (*Mölle et al., 2009*; *Rosanova and Ulrich, 2005*). These observations have led to the
418 hypothesis that SO provides a global temporal frame within the cortex and between brain regions
419 (e.g., hippocampus, thalamus, striatum) for offline memory processing and reactivation through
420 the strengthening of neuronal circuits (*Isomura et al., 2006*; *Ji and Wilson, 2007*; *Pfaff et al., 2008*;
421 *Rasch et al., 2007*; *Wierzynski et al., 2009*). Local sleep may thus enable cortical networks to aug-
422 ment memory consolidation in specific brain regions that need it the most.

423 While specific mechanisms of slow-wave sleep (SWS) generation have been explored in many
424 local circuit models (*Timofeev et al., 2000*; *Bazhenov et al., 2002*; *Compte et al., 2003*; *Hill and*
425 *Tononi, 2005*; *Destexhe, 2009*; *Crunelli and Hughes, 2010*; *Levenstein et al., 2019*), only a few very
426 recent studies (*Cakan et al., 2022*; *Goldman et al., 2023*) have proposed simulating realistic long-
427 range cortical connectivity to test its effect on the spatio-temporal properties of SO. While these
428 models take an important step in utilizing global brain connectivity data to study SWS, they ap-
429 ply a mean-field neural-mass representation of cortical regions, which limits their ability to study
430 how the interaction of local (within a few millimeters) and long-range connectivity affects spatio-

	Density ⇓	Radius ⇓	Delay ⇑	Strength ⇓	Long Range Strength ⇓
Frequency	⇓	-	↓	⇓	-
Amplitude	⇓	⇓	↓	⇓	⇓
Speed	-	⇓	-	-	⇓
Synchrony	⇓	⇓	↓	⇓	⇓
Participation	⇓	-	-	-	-
Spread	Global	Local	Global	Local	Local

Table 1. Summary of results for each network manipulation. Decreasing connection density resulted in decreased slow wave frequency, amplitude, synchrony, and spread, but slow waves remained global. Decreasing connection radius resulted in decreased slow wave amplitude, speed, participation, and spread. Increasing delays resulted in effects are seen only for manipulations far past biological plausibility, as indicated with a single down arrow for frequency, amplitude, and participation. Decreases in synaptic strength caused reduced amplitude, synchrony, frequency and participation. Finally, decreasing the strength of only long range connections resulted in traveling local slow waves with decreased amplitude, speed, and synchrony.

---

temporal SWS dynamics.

Here, we developed a large-scale thalamocortical network model based on spiking excitatory

and inhibitory neurons in thalamus and cortex, organized in 6 cortical layers and thalamic core

and matrix subsystems, with connectivity based on Human Connectome Project (HCP) diffusion-

MRI (dMRI) tractography- based connection probabilities and HCP-derived degree of grey-matter

myelination (*Rosen and Halgren, 2021*; *Van Essen et al., 2013*). To test the contribution of specific

elements of connectivity to SO properties, we then systematically modified selected elements of

the network connectivity: connection density and radius, synaptic delays and strength.

While we inferred connectivity profiles from dMRI data, predicting the strength of synaptic con-

nections is much harder problem, especially considering that the proposed model is still a signif-

icant scaling down of the biological brain. Therefore, a substantial effort of this project was to

explore the strength of connections to identify the SWS regimes that match those known from the

literature and characterized in our data. We found that altering the balance of excitation and inhibi-

tion had a significant impact on the synchronization properties of slow waves. While our baseline

model produced unrealistically synchronized global slow-waves, reducing the strength of "long-

range" excitatory connectivity resulted in models with mixed global and local slow waves, similar

to results reported in vivo (*Nir et al., 2011*; *Funk et al., 2016*). This occurred in models where con-

nections longer than 5-10 mm were reduced four- to five-fold compared to the baseline model, to

mimic the high density and strength of local connections in vivo (*Hawkins and Ahmad, 2016*; *Lon-*

don and Segev, 2001). Interestingly, this aligns with the results presented by (*Destexhe et al., 1999*),

which show an approximately 7mm radius within which slow waves are highly synchronous. Fur-

thermore, (*Riedner et al., 2007*) and (*Vyazovskiy et al., 2007*) have reported that the occurrence of

multi-peak slow waves, where each peak has a different origin, increases over the course of sleep,

which was proposed to result from a gradual net decrease in the strength of cortico-cortical connec-

tions (*Vyazovskiy et al., 2009*) (however, see (*Timofeev and Chauvette, 2017*)). Human recordings

(*Siclari et al., 2014*; *Bernardi et al., 2018*) reported two types of slow waves: Type I - global and

more frequently occurring earlier in sleep, and Type II - local slow waves found later in sleep. A

recent study suggests that global slow waves (referred to as SO) are responsible for the consolida-

tion of new memories, while local events (referred to as delta oscillations) lead to forgetting (*Kim*

et al., 2019).

To validate the model, we compared models with different synaptic connectivity profiles to

human recordings during NREM sleep. This was done via a coherence analysis to determine cor-

respondence of signals across distance and frequency and functional connectivity analysis to eval-

uate percentage of parcel pairs with a coherence greater than a given threshold and the number

of communities. We found the experimental data to closely match the mixed state models, i.e.,

models showing coexistence of local and global slow-waves. Importantly, we also found that relatively modest changes of synaptic connectivity in these models lead to the models with only local or global slow-waves. Taking these results together, our model predicts that changes of the cortical synapses strength over night, may explain more frequent occurrence of global slow waves earlier in sleep. However, a more extensive analysis of synaptic connectivity in the model and analysis of in vivo data are needed to confirm this prediction.

Several studies have revealed a link between long-range intracortical connectivity and the properties of sleep slow waves. In (*Kurth et al., 2017*), it was shown that sleep slow oscillations propagate over longer distances with increasing age, and cortical SO propagation was positively correlated with intrahemispheric myelin content. Additionally, it was found that human subjects with a steeper rising slope of the slow wave exhibited higher axial diffusivity in the temporal fascicle and frontally located white matter tracts (*Piantoni et al., 2013*). These findings suggest that the profiles of sleep oscillations reflect not only the synaptic-level dynamics of the neuronal network but also the microstructural properties of its structural foundation, the white matter tracts. Factors such as normal aging (*Tomasi and Volkow, 2012*) and Alzheimer's Disease (AD) (*Liu et al., 2014*) impact long-range connectivity. Specifically, in patients with severe Alzheimer's Disease, functional connectivity was notably reduced between regions separated by greater distances, and this loss of long-distance connectivity was associated with a less efficient global and nodal network topology (*Liu et al., 2014*). Alongside well-documented observations of reduced slow wave density in aging (*Mander et al., 2013*) and Alzheimer's Disease (*Katsuki et al., 2022*), these findings align with our model predictions about the critical role of balancing local and long-range connectivity for maintaining characteristic slow-wave dynamics.

The sleep SO is hypothesized to be a cortical rhythm (*Steriade et al., 1993a*; *Sanchez-Vives and McCormick, 2000*; *Timofeev et al., 2000*; *Timofeev and Steriade, 1996*). In line with these results, cortical SO were maintained in the model even when thalamic input was removed. However, other studies have shown that the normal pattern of the neocortical SO requires thalamic inputs to synchronize activity across large cortical areas (*Lemieux et al., 2014*). Here, we found that removing the thalamus reduced synchrony of slow-waves in the mixed-state (but not the global) SO model, suggesting its role in long-range synchronization of cortical activity. A deeper study into thalamus involvement in SO synchrony will be a highly relevant avenue for future experimental and modeling studies. Moreover, despite not driving the SO rhythm that constitutes the object of this study, the thalamus allows this model to capture a range of human brain dynamics beyond SO. Indeed, natural extensions of this model include investigating sleep spindles, in which the thalamus is heavily involved (*Steriade et al., 1993b*, *1990*), or the cortical impacts of thalamic stimulation, which constitute an active field of research in neuroscience (*Neudorfer et al., 2024*; *Schüller et al., 2024*).

Although our model design achieves high computational efficiency considering the size and complexity of the network, the number of model parameters and simulation time do not easily allow a systematic fitting of the model output to electrophysiological brain signals, as it has been done with smaller computational models (*Cakan et al., 2022*). It is possible that optimization techniques such as surrogate model optimization or simulation based inference may be used on a subset of parameters in the model with the addition of biological constraints on possible values. Future studies may wish to explore this, but care will have to be taken to maintain biological plausibility. Nevertheless, the use of realistic parameters for individual neuron dynamics in wake and sleep stages (*Bazhenov et al., 2002*; *Krishnan et al., 2016*) as well as experimentally grounded cortical connectivity (*Rosen and Halgren, 2021*) allowed us to reproduce the essential characteristics of SWS activity in human brain.

Importantly, we have now established a biologically relevant model of network dynamics that can be used for a multitude of other applications. For example, this model could be used to study possible mechanisms of sleep disorders by testing the fit of various hypothesis-driven perturbations of the model to relevant experimental data. In that context, one particularly important application of the model is using it to reveal mechanisms of sleep alterations in aging and Alzheimer's

Disease. Beyond the realm of sleep, this model could be used to study the response of a healthy
brain to various types of traumatic brain injury or neurodegenerative diseases that may alter neu-
ral connectivity. Finally, this model could further provide preliminary insight on the network effects
of drugs that act on specific neurotransmitters, either globally or in a targeted location.

In summary, we present a whole-hemisphere thalamocortical network model constrained by
realistic local and long range connectivity based on human dMRI data and implementing effects
of neuromodulators to account for sleep-wake transitions. The model displays both a biologically
realistic awake state with characteristic random neuron firing as well as Type I and Type II slow
wave sleep. The critical results of our study lie in revealing the separate contributions of struc-
tural connectivity (i.e., connectivity matrix) and functional connectivity (i.e., synaptic connectivity
strength): while structural connectivity can alter the characteristics of individual slow waves (syn-
chrony, spread, participation), it is primarily the proper balance of connectivity strength between
local and global connections that leads to the mixed cortical states, where periods of synchronized
global SO are intermittent with the periods of asynchronous local slow-waves. Importantly, only the
models generating such intermittent states show spatio-temporal SO characteristics that match hu-
man recordings. Comparing the models to empirical data further supported the fundamental role
of connectivity strength in generating biophysically realistic network behavior via mixed-state mod-
els. Furthermore, the model provides both the scale and the level of detail required to investigate
how large-scale sleep dynamics arise from cellular and circuit level activity.

Methods

Model neurons

Map-based cortical neurons

The model has 10,242 cortical columns uniformly positioned across the surface of one hemisphere,
one for each vertex in the ico5 cortical reconstruction reported in *Rosen et al.* (*2019*). The medial
wall includes 870 of these vertices, so all analyses for this one-hemisphere model were done on the
remaining 9372 columns. Each column has 6 cortical layers (L2, L3, L4, L5a, L5b and L6). For each
layer in each column, one excitatory (PY) and one inhibitory (IN) neuron were simulated. To allow
for large-scale network simulations we modeled cortical neurons with a model based on difference
equations (MAP) (*Rulkov, 2002*; *Rulkov et al., 2004*) which has a number of distinct numerical ad-
vantages: the common problem of selecting the proper integration scheme is avoided since the
model is already written in the form needed for computer simulations. The model parameters
can be adjusted to match experimental data (*Rulkov et al., 2004*; *Bazhenov et al., 2008*; *Rulkov*
and Bazhenov, 2008). Map-based models sample membrane potential using a large discrete time
step compared to the widely used conductance-based models described by ordinary differential
equations, and still capture neural responses from these models. Importantly, map-based models
replicate spiking activity of cortical fast-spiking, regular-spiking and intrinsically bursting neurons
(*Rulkov et al., 2004*). Variations of the map model have enabled the simulation of large-scale brain
networks with emergent oscillatory activity (*Rulkov and Bazhenov, 2008*), including realistic sleep
spindles (*Rosen et al., 2019*) and cortical slow oscillations (*Komarov et al., 2018*). In our study, we
used the map model modification proposed by *Komarov et al.* (*2018*), which implemented nonlin-
ear dynamical bias, activity dependent depolarizing mechanisms, and slow hyperpolarizing mech-
anisms capturing the effects of leak currents, the Ca2+ dependent K+ currents and the persistent
Na+ currents, which were found to be critical for simulating up/down state transitions during SO.
Parameter values were initially set to the reference values in *Komarov et al.* (*2018*), which were
shown to produce biologically realistic neural behavior during SO, and then tuned with small varia-
tions to maintain adequate neural responses in our comparatively larger-scale network. The map
model equations are described below, and all parameter values specific to this study are reported
in Table 2.

The activity of individual pyramidal PY cells was described in terms of four continuous variables

sampled at discrete moments of time n : x_n, dictating the trans-membrane potential $V_n = 50x_n - 15$; y_n, representing slow ion channel dynamics; u_n, describing slow hyperpolarizing currents; and k_n, determining the neuron sensitivity to inputs. The activity of IN cells, which biologically have faster spiking dynamics, was modeled using only variable x_n and fixing the other variables to constant values. Dynamical PY variables evolve according to the following system of difference equations, where $n = 1, 2, ...$ indexes time steps of 0.5ms in size (as suggested in *Rulkov and Bazhenov* (*2008*)):

$$\text{spike generation} \begin{bmatrix} x_{n+1} = f_{PY}(x_n, y_n + \beta_x I_{total}(x_n, u_n), k_n) \\ y_{n+1} = y_n - \mu_1 \cdot (x_n + 1) + \mu_2 \cdot (\sigma p_L + \beta_y I_{total}(x_n, u_n)), \end{bmatrix} \tag{1}$$

$$\text{adaptation} \begin{bmatrix} u_{n+1} = \gamma \cdot (u_n + \delta) \end{bmatrix} \tag{2}$$

$$\text{sensitivity to input} \begin{bmatrix} k_{n+1} = k_n + H(x_n + 0.5) \cdot (K_1 - k_n) + H(-(x_n + 1)) \cdot (K_0 - k_n)) \end{bmatrix} \tag{3}$$

where function $H(w)$ denotes the Heaviside step function, which takes the value 0 for $w \le 0$ and 1 otherwise. The reduced model for IN cells is given by

$$x_{n+1} = f_{IN}\left(x_n, y^* + \psi(\beta_x I_{syn})\right) \tag{4}$$

$$\psi(w) = \begin{cases} -1 & , \ w < -1 \\ w & , \ -1 \le w \le 1 \\ 1 & , \ 1 < w \end{cases} \tag{5}$$

where y^* is a fixed value of y_n. We describe each of the system's components below. The model's parameters and their values are summarized in *Table 2*.

Spike-generating Systems (1) and (4): Variables x and y model the fast neuron dynamics, representing the effects of the fast Na$^+$ and K$^+$ currents responsible for spike generation (*Hodgkin and Huxley, 1952*). The nonlinear function f_{PY}, modified by *Komarov et al.* (*2018*) from *Rulkov* (*2002*), determines the spike waveform as

$$f_{PY}(x_n, w, k_n) = \begin{cases} \dfrac{\alpha}{1 - x_n} + w & , \ x_n < -0.5 \ \wedge \ w < w_0, \\ \dfrac{\alpha}{1 - x_n} + w_0 + (w - w_0)k_n & , \ x_n < -0.5 \ \wedge \ w \ge w_0, \\ 1 & , \ -0.5 \le x_n < 1, \\ -1 & , \ x \ge 1 \end{cases} \tag{6}$$

where α and w_0 are constant parameters that control the non-linearity of the spike-generating mechanism. Parameters $\mu_1, \mu_2 < 1$ modulate the slower updating of variable y with respect to variable x, and parameter σ dictates the neuron resting potential as $(1 - \sigma)$. The scaling constants β_x, β_y modulate the input current I_{total} for variables x and y, respectively. The neuron's membrane potential is calculated as $V_n = 50x_n - 15$ and a spike is generated if and only if $V_n > 0.1$. In simulations, the quantity $\beta_x I_{total}(x_n, u_n)$ was manually prevented from reaching 0 by imposing a lower bound of 10^{-4}.

For inhibitory neurons IN, function f_{IN} in (4) is given by

$$f_{IN}(x_n, w) = \begin{cases} \dfrac{\alpha}{1 - x_n} + w & , \ x_n \le 0, \\ \alpha + w & , \ 0 < x_n \le \alpha + w \ \wedge \ x_{n-1} \le 0, \\ -1 & , \ \text{otherwise} \end{cases} \tag{7}$$

A spike is generated if and only if $0 < x_n \le \alpha + y^* + \psi(\beta_x I_{syn}) \ \wedge \ x_{n-1} \le 0$.

588 **Adaptation Equation** (2): The slow variable u_n represents the effects of the slow hyperpolariz-
589 ing potassium currents (I_D, explained below) reducing neural excitability over the course of the Up
590 state and involved in Up state termination. The value of γ is taken as 0.99 for $x_n < -1$ and 0.995
591 otherwise. Parameter $\delta = 0.24$ has an effect only if $-0.5 < x_n < 1$; we assume $\delta = 0$ for other values
592 of x_n.

593 **Sensitivity Equation** (3): The variable k_n reduces neuron sensitivity to inputs during Up states
594 (i.e. during high spiking activity) (*Steriade et al., 2001*) by regulating the spike nonlinearity func-
595 tion f_{PY}. This gives a net phenomenological representation of the combined effects of high-level
596 synaptic activity, increase in voltage-gated hyperpolarizing currents, and/or adaptation of fast Na$^+$
597 channels. It adopts the value K_0 (low sensitivity) when voltage x_n crosses threshold -0.5 (i.e. Up
598 state) and switches to $K_1 > K_0$ (high sensitivity) when $x_n < -1$ (i.e. cell leaves Up state). This pre-
599 vents over-excitation and provides a physiological firing frequency (*Komarov et al., 2018*).

600 **Computation of currents**: The total current I_{total} is a sum of external applied DC current (I_{DC}),
601 synaptic inputs (I_{syn}), persistent Na$^+$ current (I_{Nap}), slow hyperpolarizing currents (I_D), and leak
602 currents (I_{leak}):

$$I_{total} = I_{DC} + I_{syn} + I_{Nap} + I_D + I_{leak}. \tag{8}$$

603 with the last three dictated primarily by p_{Nap} (maximal conductance of persistent Na$^+$ current), p_D
604 and p_L (strength of leak current), respectively:

$$I_{Nap} = \frac{p_{Nap}}{1 + \exp\left(-20(x_n + 0.99)\right)} \tag{9}$$

$$I_D = -p_D u_n (x_n + 1.2) \tag{10}$$

$$I_{leak} = -p_L (x_n - \underbrace{(\sigma p_L - 1)}_{\text{baseline potential}}) \tag{11}$$

605 The persistent Na$^+$ current contributes to initiating and maintaining Up states (*Timofeev et al.,*
606 *2000*). It is modeled in I_{Nap} through a sigmoidal activation function depolarizing the neuron with
607 strength ρ_{Nap} at high voltage values. The effect of slow hyperpolarizing currents contributing to Up
608 state termination is represented by I_D, which is regulated by the dynamic variable u_n (described
609 above). Lastly, ρ_L represents the effect of potassium leak current, and here it is used to model
610 the hyperpolarization of cortical neurons during sleep (*Bazhenov et al., 2002*). For each neuron i,
611 the synaptic current I_{syn}^i is the sum of individual AMPA, GABA$_A$ and NMDA synaptic currents. The
612 individual synaptic current of type X from neuron j to neuron i has the general form

$$g_X^{ij}(n)\left(V_X - V_n^i\right)$$

613 for AMPA and GABA synapses, and

$$g_X^{ij}(n)\varphi_n\left(V_X - V_n^i\right)$$

614 for NMDA synapses, with an additional modulator φ_n,

$$\varphi_n = \left(1 + e^{-\frac{V_n^i + 25}{12.5}}\right)^{-1}.$$

615 Here $g_X^{ij}(n)$ represents conductance at time n for directed connections from neuron j to neuron i.
616 Dynamic synaptic conductances were described by the first-order activation schemes:

$$g_X^{ij}(n+1) = \underbrace{\gamma_{g_X} g_X^{ij}(n)}_{\text{decay}} + \underbrace{d_X^{ij}(n) \frac{G_X^{syn}}{m_X^i} H\left(V_n^j - V_\theta\right)}_{\text{activity-dependent depression}} + \underbrace{\frac{G_X^{mini}}{m_X^i} \eta_n^{mini}(\mu_X)}_{\text{miniature postsynaptic potentials}} \tag{12}$$

617 with $d(n)$ governed by

$$d_X^{ij}(n+1) = \begin{cases} \gamma_{d_x}^{dep} d_X^{ij}(n) & , V_n^j > V_\theta \ (j \text{ spikes}) \\ 1 + \gamma_{d_x}^{rec}(d_X^{ij}(n) - 1) & , \text{otherwise} \end{cases} \tag{13}$$

618 for AMPA and GABA$_A$ and

$$d_X^{ij}(n+1) = \begin{cases} d_X^{ij}(n) & , V_n^j > V_\theta(j \text{ spikes}) \\ 1 - (1 - \gamma_{d_X}^{rec})\left(1 - d_X^{ij}(n)\right) & , \text{otherwise} \end{cases} \tag{14}$$

for NMDA type synapses. Here, γ_{g_X} determines the decay rate of synaptic conductance and G_X^{syn}
regulates the overall synaptic strength. To keep total synaptic input per cell constant, coefficients
G_X^{syn} and G_X^{mini} are scaled by the number m_X^i of synapses of type X incoming to neuron i. The de-
pression variable d_X is dependent on whether the pre-synaptic neuron j spikes at time n or not,
that is, if the voltage of neuron j surpasses a threshold V_θ. Lastly, miniature post-synaptic poten-
tials are only present in AMPA and GABA synapses. Variable η_n^{mini} is a spontaneous event from a
random Poisson process with mean frequency μ_X, and G_X^{mini} is the constant by which conductance
increases in an event ($\eta_X^{mini} = 1$), which produces miniature post-synaptic potentials ("minis") in neu-
ron i (*Stevens, 1993*). Following *Bazhenov et al.* (*2002*), the frequency μ_X is modulated in time to
account for the reduction in mini frequency after Up states (*Timofeev et al., 2000*), so that

$$\mu_X = \bar{\mu}_X \left(0.2 + \frac{0.8}{1 + e^{-0.1(n-n_0^j-350)}}\right)$$

where n_0^j is the time of the most recent pre-synaptic spike. This modulation reduces the mini rate
to 20% of its maximal value $\bar{\mu}_X$ after a pre-synaptic spike, and then allows recovery to $\bar{\mu}_X$ with time.

Hodgkin-Huxley thalamic neurons
The thalamus was modeled using a network of core (specific) and matrix (non-specific) nuclei, each
consisting of thalamic relay (TC) and reticular (RE) neurons. We simulated 642 thalamic cells of
each of these four types. TC and RE cells were modeled as single compartments with membrane
potential V governed by the Hodgkin-Huxley kinetics, so that the total membrane current per unit
area is given by

$$I = C_m \frac{dV}{dt} = -g_L(V - E_L) - I_{int} - I_{syn},$$

where C_m is the membrane capacitance, g_L is the non-specific (mixed Na$^+$ and Cl$^-$) leakage conduc-
tance, E_L is the reversal potential. These parameters (*Table 3*) were unchanged from (*Bazhenov*
et al., 1998a), where they were determined based on previous modeling studies grounded in ex-
perimental observations (*McCormick and Huguenard, 1992*; *Destexhe et al., 1994*). Specifically, tha-
lamic reticular and relay neurons included intrinsic properties necessary for generating rebound
responses which were found to be critical for spindle generation (*Bazhenov et al., 2000*; *Destexhe*
et al., 1996). Furthermore, I_{int} is a sum of active intrinsic currents, and I_{syn} is a sum of synaptic
currents.
Intrinsic currents for both RE and TC cells included a fast sodium current, I_{Na}, a fast potassium
current, I_K, a low-threshold Ca^{2+} current, I_T, and a potassium leak current, $I_{KL} = g_{KL}(V - E_{KL})$.

Name	Description	PY value	IN value
β_x	x_n input current scaling	0.133	0.05
α	spike nonlinearity parameter	3.65	3.8
w_0	spike nonlinearity parameter	-2.819	-
y^*	fixed y_n value for IN neurons	-	-2.93
μ_1	y_n timescale relative to x_n	0.0008	-
μ_2	y_n timescale relative to x_n	0.002	-
σ	determines neuron resting potential	-0.4	-
β_y	y_n input current scaling	1	-
γ	strength of adaptation	0.99/0.995	-
δ	adaptation parameter	0.24/0	-
K_1	high sensitivity value	0.25	-
K_0	low sensitivity value	0.0025	-
p_{Nap}	persistent Na$^+$ current strength	0.15	-
p_D	slow hyperpolarizing current strength	4	-
p_L	leak current strength	0.5	-
γ_{g_X}	decay rate of AMPA/GABA$_A$ synaptic conductance	0.995	-
G^{syn}_{AMPA}	AMPA synaptic strength	$7.3 \cdot 10^{-5}$	$2.3 \cdot 10^{-4}$
G^{syn}_{GABA}	GABA$_A$ synaptic strength	$5 \cdot 10^{-5}$	-
G^{syn}_{NMDA}	NMDA synaptic strength	$6.7 \cdot 10^{-10}$	-
G^{mini}_{AMPA}	mini strength for AMPA synapses	$1.6 \cdot 10^{-6}$	
G^{mini}_{GABA}	mini strength for GABA$_A$ synapses	-	0
$\bar{\mu}_X$	mini frequency for AMPA/GABA$_A$ synapses	500	
$\gamma^{dep}_{d_X}$	time scale of synaptic depression	0.05	-
$\gamma^{rec}_{d_X}$	time scale of synaptic recovery	0.005	-
$\gamma^{rec}_{d_{NMDA}}$	time scale of NMDA synaptic recovery	$5 \cdot 10^{-5}$	-
V_{AMPA}	reversal potential for AMPA synapses	0	-
V_{GABA}	reversal potential for GABA$_A$ synapses	-70	-
V_{NMDA}	reversal potential for NMDA synapses	0	-

Table 2. Map-based model parameter definitions and values. Parameters γ and δ may take one of the two values shown (see text). Parameters were initially set to the values in (*Komarov et al., 2018*) and then fine tuned to maintain biologically reasonable spiking behavior in the current larger-scale network. Sub-index X stands for "AMPA" or "GABA". For all parameters regarding synapses, the PY and IN columns denote the post-synaptic neuron

Name	Description	TC value	RE value	Units
C_m	membrane capacitance	1		$\mu F/cm^2$
g_L	leakage conductance	0.01	0.05	mS/cm^2
E_L	reversal potential	-70	-77	mV
S	cell area	$1.43 \cdot 10^{-4}$	$2.9 \cdot 10^{-4}$	cm^2

Table 3. Parameter definitions and values for conductance-based models are based on previous modeling of single-cell dynamics (*Bazhenov et al., 1998a*, *2002*; *McCormick and Huguenard, 1992*; *Destexhe et al., 1994*)

Parameter	Wake	Sleep
p_L	0.15	0.5
p_D	1.6	4.0
G_{AMPA}^{syn} (PY→PY)	$2 \cdot 10^{-5}$	$7.3 \cdot 10^{-5}$
G_{AMPA}^{syn} (PY→IN)	$6.3 \cdot 10^{-5}$	$2.3 \cdot 10^{-4}$
G_{GABA}^{syn} (IN→PY)	$3 \cdot 10^{-5}$	$5 \cdot 10^{-5}$
g_{KL} (TC)	0.007	0.012
g_{KL} (RE)	0.022	0.015

Table 4. Parameter changes between sleep and wake stages

647 For TC cells, an additional hyperpolarization-activated cation current, I_h, was also included. The
648 expressions and parameter values for each of these currents are given in *Bazhenov et al.* (*1998a*,
649 2002); *Krishnan et al.* (*2018*); *Rosen et al.* (*2019*).

650   Synaptic currents (GABA, NMDA and AMPA) were modeled by first-order activation schemes
651 (*Destexhe et al., 1994*), using the general form

$$I_{syn} = g_{syn}[O](V - E_{syn}),$$

where g_{syn} is the maximal conductance, $[O](t)$ is the time-dependent fraction of open channels, and
E_{syn} is the reversal potential. Our previous studies show specific equations for all synaptic currents
(*Bazhenov et al., 2002*; *Chen et al., 2012*), as well as a detailed description of O(t) based on first-
order activation schemes (*Bazhenov et al., 1998b*; *Timofeev et al., 2000*).

Wake to sleep transition

The transition from wake to slow wave sleep stage was accomplished via tuning of several key
parameters. The primary tunable parameters of the map model cortical neurons are p_D and p_L,
which control the strength of the slow hyperpolarizing currents I_D (similar to a calcium current) and
the potassium leak currents I_{leak} as described in Equations (10) and (11). Both of these parameters
significantly increased from wake to sleep. We further increased the strength of AMPA and GABA
synaptic currents, as well as the strength of TC potassium leak currents g_{KL}, similar to the changes
described in *Krishnan et al.* (*2016*) (see *Table 4*).

Network Structure

Our whole hemisphere computational model has realistic local and long-range cortical connectiv-
ity. Connections within the column follow those described in the canonical cortical circuit (Figure
1A). Additionally, connections within 0.1 mm are strengthened by a factor of 5 to mimic in-vivo
increased strength of proximal synapses (*London and Segev, 2001*; *Hawkins and Ahmad, 2016*). In-
hibition is local: IN cells project only to PY cells in the same layer, with connections only in their
own column, 1st and 2nd degree neighbors. The strength of inhibitory synapses is 2 times larger
within the local column than outside. Finally, miniature postsynaptic potentials on intracortical
AMPA synapses are modeled as a Poisson process (as in *Krishnan et al.* (*2018*)).

Thalamocortical (TC) and reticular (RE) neurons are modeled using Hodgkin-Huxley dynamics
(*Hodgkin and Huxley, 1952*), with 2 subtypes of each to represent matrix and core neurons. The
matrix TC neurons have a 45 mm (and 80 mm) fanout radius to cortical excitatory (and inhibitory)
neurons while core TC neurons have 12 mm and 13 mm radii, respectively. Within-thalamic connec-
tions have 11 mm radii. There are 642 neurons of each of these 4 types (thalamic matrix, thalamic
core, reticular matrix, reticular core).

Hierarchically Guided Laminar Connectivity

Long-range connectivity in the model is primarily based on the cortical parcellation into 180 areas
proposed by the Human Connectome Project (HCP)-MMP1.0 atlas (*Van Essen et al., 2013*; *Glasser*

682 *et al., 2016*). Each parcel was assigned a hierarchical index inversely proportional to its estimated
683 bulk myelination (*Glasser and Essen, 2011*; *Burt et al., 2018*). Excitatory cortical connections (PY-
684 PY) were then split into 6 classes according to the pre- and post-synaptic hierarchical index: within
685 the local column, within the local parcel (or between contralateral homologs), strongly (>50th per-
686 centile) or weakly (≤50th percentile) feedforward (from lower to higher hierarchical index), and
687 strongly or weakly (>, ≤ 50th percentile respectively) feedback (from higher to lower hierarchi-
688 cal index). Based on previous connectivity reports (*Markov et al., 2013*; *Rockland, 2019*), synaptic
689 weights for these connections were scaled by the factors shown in Figure 1 D.

Diffusion MRI Guided Connection Probability

691 The connection probability was further scaled based on previous diffusion MRI (dMRI) connectivity
692 studies (*Rosen and Halgren, 2021*). These observations reveal an exponential decay relationship
693 between inter-parcel dMRI connectivity and fiber distance (length constant $\lambda = 23.4$mm, scaling
694 parameter $\beta = 0.17$). In order to obtain a probability of connection at the scale of columns, ap-
695 proximate intercolumnar fiber distances were estimated from geodesic distances. For parcel cen-
696 troids, intra-hemispheric dMRI streamline distances and their corresponding geodesic distances
697 were found to be related to a first approximation by a linear rational function

$$F(x) = \frac{(p1 * x + p2)}{(x + q1)},$$

698 with $p1 = 295.6, p2 = -2256$, and $q1 = 69.4$, where x denotes the geodesic distance. Column-
699 wise fiber distances were estimated by applying this function to intercolumnar geodesic distances.
700 Thus, a distance-dependent probability of connection between columns was obtained by applying
701 the dMRI exponential relation to column-wise fiber distances (Figure 1 E). Moreover, the residual
702 parcel-wise dMRI connectivity (not distance related, obtained by regressing out the exponential
703 trend) was added back to intercolumnar connections based on the columns' parcels. This distance-
704 independent connectivity accounts for the functional specialization of each parcel. Lastly, con-
705 duction delays were set to be proportional to fiber tract distances, with the longest connection
706 (~226mm) having an assigned delay of 40ms.

Analysis

Modifying Network Structure

709 We here attempted to create the most biologically realistic model and connectivity possible. How-
710 ever, there is no ground truth to dictate the absolute strength of connection between two individ-
711 ual neurons. While rules exist in our architecture to modify relative strength based on biological
712 connectivity, the base connection strength to be modified is a completely free (and incredibly im-
713 portant) parameter. We first set the base strength level to be high enough to generate Type I slow
714 waves which quickly synchronize across the entire brain. This makes slow wave property analysis
715 more straightforward and less prone to spatial bias when modifying further network parameters,
716 as is the goal of this paper. Hence, we use this global Type I slow wave model for all subsequent
717 analysis aside from analyzing the effect of connection strength itself.

718 In the case of modulating connection density and range, the synaptic strengths were scaled up
719 proportionally to the loss of connections to retain sufficient activity levels in the network. The scal-
720 ing mechanism works on an individual neuron basis. For each neuron, the weights of all synapses
721 of each type (AMPA, GABA, etc.) are scaled by the total number of synapses of that type. Thus, af-
722 ter removing connections, the weights are divided by a smaller quantity, which results in stronger
723 connections. This approach allows us to test the relative effects of connectivity, e.g., local vs global,
724 without dramatic changes in the network dynamics.This allowed us to isolate the effects of patterns
725 of connection apart from changes in total synaptic input. The ablation of connections in these trials
726 was accomplished by setting synaptic connection strengths to 0, removing all functional connectiv-
727 ity.

Local Field Potentials, Amplitude and Frequency

In Figures 2,4,3, the local field potential (LFP) was obtained as the average voltage of Layer 2 at each timestep scaled by a factor of 20. In Figures 5,S6,7, the LFP for each local region was further smoothed using a running mean with 20ms window.

Layer LFPs were used to detect cortical Up states. A peak detection algorithm (MATLAB's islocalmax function) was implemented for this purpose, requiring a minimum peak separation of 500ms and a minimum prominence of half the difference between the maximum and minimum values of the signal. The amplitude of each Up state was defined as the difference between the corresponding peak and the lowest point before the next peak. The inverse of each inter-peak interval was used to approximate the oscillation frequency. Average amplitudes and frequencies across all Up states of a simulation are reported in the summary plots of Figures 4,3,S7, 6.

Onset/offset Detection

In order to characterize how SO spread through the cortex, we identified the points in time when each neuron transitions from Down state to Up state (onset) and viceversa (offset). Our Up state detection algorithm follows *Komarov et al.* (*2018*), based on *Volgushev et al.* (*2011*): we set two thresholds, $V^+ = -65$mV and $V^- = -68$mV (see dashed vertical lines in voltage histogram of Figure 2 B), and label any activity above V^+ as an Up state and any activity below V^- as a Down state. This initial detection was further refined by merging any two Up states (or Down states) that were less than 100ms apart, and then removing any remaining Up state (or Down state) lasting less than 100ms, in a similar approach to *Volgushev et al.* (*2011*).

To identify onsets and offsets for a particular global Up state, we considered the time window between consecutive minima of the average voltage signal around that Up state (e.g. see black upward triangles in average voltage signal of Figure 2 B). For each neuron, we found the first time (if any) in this window when it reached an Up state (onset), and then the first time after the onset when it reached a Down state (offset). The minimum onset was then subtracted from all onsets so that the earliest neuron would have an assigned time of 0 and all other would have positive delays, or *latencies*. Similarly, the earliest offset was subtracted from all offsets.

In order to evaluate the synchrony of active and silent states, we constructed aggregated histograms of the onset and offset times across all Up states of each simulation. We used the standard deviation of the histograms as an measure for synchrony, with narrow histograms indicating highly synchronized global Up states.

Latency and participation

Latency maps are shown in Figure 2 F (also Figure S6 F, Figure 5 F and Figure 7 F) for each global Up state, representing the delay for activation of each cell after the earliest onset in that Up state. The 3D representation of latencies allows for a visual identification of the source (or sources) of each global Up state, characterized by clusters of adjacent neurons with near-zero latencies.

The percent of participating columns in each Up state was defined as the number of columns with onset delays greater than or equal to 0 over the total number of columns not belonging to the medial wall. Non-participating columns are shown in white in the latency maps for each Up state. The average percent of active columns per Up state is reported in the summary plots in Figure 3F,Figure 3F, Figure S7 and Figure 6C-D.

Speed of propagation

To obtain speed of propagation of each Up state, we first identified columns with onset delays under 10ms to get an approximate area of origin. We then selected the column with minimal sum of (geodesic) distances to all columns with onset delays under 10ms, excluding the medial wall. This column was identified as the origin of the Up state. We considered only an area of 150mm radius around the origin to compute propagation speed, in order to avoid interference from waves originating later in other cortical regions. A linear regression was performed between

776 the onset delays of columns in the area and their distances to the origin. For regressions with an

777 $R^2 > .25$, the speed of propagation was defined as the slope of the regression line. The average

778 propagation speed over all Up states is reported in summary plots. Speed is not reported for

779 simulations in which there is not a linear relation between onset delay and distance from the origin,

780 that is, simulations in which regressions for all Up states yield $R^2 \leq .25$.

Experimental Data

781

782 Patients and intracranial recordings

783 Continuous stereo-electroencephalography telemetry and structural imaging were obtained for

784 298 patients undergoing pre-resection intracranial monitoring for phamaco-resistant focal epilepsy

785 at University of California, San Diego Medical Center, La Jolla CA, the Cleveland Clinic, Cleveland

786 OH, Oregon Health and Science University Hospital, Portland OR, Massachusetts General Hospital,

787 Boston MA, Brigham and Women's Hospital, Boston MA, the Hospital of the University of Pennsyl-

788 vania, Philadelphia PA, and The University of Alabama Birmingham Hospital, Birmingham AL. These

789 clinical data are recorded as standard of care and written informed consent, approved by each insti-

790 tution's review board (The Institutional Review Board of the University of California, San Diego, The

791 Oregon Health and Sciences University Institutional Review Board, The University of Pennsylvania

792 Institutional Review Board, The University of Alabama, Birmingham Institutional Review Board for

793 Human Use, The Institutional Review Board at the Cleveland Clinic Foundation, The Mass General

794 Brigham Institutional Review Boards), was obtained from each patient before the data were used

795 for research purposes. Recordings were collected with a Nihon Kohen Neurofax (the Cleveland

796 Clinic), Cadwell Arc (Oregon Health and Science University Hospital), or Natus Quantum amplifier

797 (all other clinical sites) and acquired with a sampling frequency of at least 500 Hz. After preliminary

798 inspection, 27 patients with prior resections, highly abnormal sleep patterns, or whose data were

799 over-contaminated with excessive epileptiform activity or technical artifacts were excluded from

800 analysis. A further 35 patients were excluded from further analysis after channel selection and

801 quality control procedures, described below, resulted in them having with fewer than two clean

802 bipolar channel pairs. After quality control, 236 patients (126 female, 101 male, 9 unknown, 36.5

803 $\pm$ 13.3 (mean $\pm$ stdev.) years old, 22 unknown age) were included in the final analysis.

804 Channel localization and inter-channel distance

805 As part of standard of care, a pre-operative T1-weighted high-resolution structural MRI and inter-

806 operative CT scan were obtained for each patient. SEEG contacts were localized as previoulst de-

807 scribed (*Dickey et al., 2022*). Briefly, the post-implant CT volume was co-registered to the pre-

808 implant MR volume in standardized 1mm isotropic FreeSurfer space with the general registration

809 module (*Johnson et al., 2007*) in 3D Slicer (*Fedorov et al., 2012*) and each contact's position was de-

810 termined by manually marking the contact centroid as visualized in the co-registered CT volume.

811 Cortical surfaces were reconstructed from the MR volume with the FreeSurfer recon-all pipeline

812 (*Fischl, 2012*). Each transcortical bipolar channel was assigned to the white-gray surface vertex

813 nearest to the midpoint of the adjacent contacts. Spherical folding-based surface registration (*Fis-*

814 *chl et al., 1999*) was used to map the individual subjects' bipolar channel vertices to the standard-

815 ized fsaverage ico7 (*Fischl, 2012*) and 32k_FS_LR grayordinate (*Marcus et al., 2013*) atlases as well as

816 to parcels of the HCP-MMP parcellation (*Glasser et al., 2016*). The HCP-MMP parcel assignments of

817 each channel were used to determine the approximate white matter fiber length of the connection

818 between them, using the mean values for a cohort of 1,065 healthy subjects we have previously

819 reported (*Rosen and Halgren, 2021*).

820 Telemetry preprocessing and channel selection

821 For each patient, an average of 185.3 $\pm$ 112.0 hours of raw telemetry were analyzed. Data were

822 anti-aliased filtered at 250 Hz and resampled to 500 Hz. To remove line noise, a notch filter was

823 applied at 60, 120, 180, and 240 Hz. On each electrode shaft, data were re-referenced to a bipolar

scheme by taking the difference of potentials in adjacent contacts, yielding 30,769 total channels.
To ensure that bipolar channels were independent, contact pairs were selected such that no two
pairs shared a contact, eliminating just under half of all channels from further consideration. Peaks
in the delta band (0.5-2 Hz) signal during high-delta periods were identified and the peri-delta-peak
high-gamma band (70-190 Hz) envelope obtained. Transcortical contact pairs were identified by
(1) having a midpoint close to the gray-matter–white-matter interface (<5 mm), (2) having a high
anticorrelation between the delta-band amplitude and the high gamma band (70-190 Hz) enve-
lope during delta-band peaks, indicative of non-pathological Down states with a surface-negative
LFP deflection that quieted spiking, and (3) having high delta-band amplitude with and average
delta peak of at least 40 μV, indicative of robust inversion in adjacent contacts. Only contacts with
5 mm pitch were analyzed to standardize absolute bipolar amplitudes and contact pairs located
deep in the white matter or in subcortical gray matter were excluded from analysis. All channels
were visually inspected to ensure they contained broadly normal non-pathological waveforms such
as pathological delta activity. This procedure yielded 4,903 candidate channels which underwent
NREM scoring, described below, yielding 9.2 $\pm$ 7.8 hours of putatively clean scored NREM sleep.
After scoring, which excluded continuous periods of artifactual or pathological signals, remaining
epileptogenic activity was detected with spike template convolution and gross artifacts, with am-
plitudes > 300 μV. Channels which contained persistent automatically detected interictal spikes or
gross artifacts were removed from further analysis, yielding 3,665 channels. Periods containing in-
terictal spikes or gross artifacts in any channel were excluded from analysis in all channels. Finally,
after cross-spectral matrices were obtained for each subject as described below, channels with
autospectral power more than 3 standard deviations above the pooled channelwise mean in any
examined frequency were excluded, leaving 3,588 channels and 37,123 within-patient bivariate
channel pairs in the final analysis.

NREM selection

Clean periods of sleep were first manually classified based on ultradian delta power. Within man-
ually identified nightly periods of sleep, NREM sleep was automatically identified by the presences
of slow oscillations (SOs) in a manner derived from standard clinical procedures (*Iber et al., 2007*).
The density of SOs was ascertained in each 30-second frame and NREM marked for the frame if at
least 35% of the frame contained SOs in at least one bipolar channel. The reduction in the number
of channels and increase in the percent of frame threshold relative to the clinical standard of 20%
of the frame across scalp channels was to accommodate the increased spatial heterogeneity of
sleep graphoelements observed intracranially, relative to the scalp (*Mak-McCully et al., 2017*). For
this purpose, slow waves were automatically detected with the method described in (*Mak-McCully*
et al., 2017), briefly, consecutive delta-band zero-crossings occurring within 0.23-3 s were detected
and the top 20% of peaks were retained as SOs. Only periods without detected interictal spikes or
gross artifacts in any bipolar channel were included in subsequent analysis.

Coherence Analysis
Mock Experimental Data
Coherence analysis was performed on both experimental and model data in an identical manner
following preprocessing. The model data was preprocessed by averaging the voltages from all
cells belonging to each of the 180 parcels (all 6 layers, both excitatory and inhibitory) to mimic
what would be recorded by an SEEG electrode. Additionally, white gaussian noise was added to
each of the 180 parcel averaged signals. This was done to mimic experimental noise picked up
from both other non-neural bodily signals (i.e., muscle movements) and noise from the recording
electrode itself. *Ball et al.* (*2009*) demonstrated that ECOG arrays show an SNR ranging from 2.22
to 4.37 in relation to eye blink movement artifacts. *Suarez-Perez et al.* (*2018*) further calculated
the SNR of slow wave sleep Up states (signal) v.s. Down states (noise); they found low frequencies
(less than 30 Hz) to have a steady SNR of 4-9, depending on the type of electrode used. At higher

frequencies, the SNR dropped severely. With these studies in mind, we implemented an SNR of 5
to account for both body and electrode noise that our model data lacks.

Signal processing and epoch sampling

A frequency resolved map of each patient's (or simulation's) neural covariability was achieved by
estimating the complex cross-spectral matrix among bipolar SEEG channels (for empirical data)
or model parcel averages. Narrow-band signals were extracted with a series of second-order res-
onator filters (matlab's iirpeak) with peaks and 3 dB attenuation bandwidths shown in Table S1. For
each frequency the complex analytic signal was extracted via the Hilbert transform. Calculating the
variance-covariance matrix of analytic signals yields the cross-spectral matrix. Cross-spectral ma-
trices were normalized into magnitude-squared coherence by squaring the complex modulus of
each cross-pairwise cross-spectrum and dividing by the product of the constituent autospectra.

For experimental data, an adaptive procedure was used to obtain stable representative esti-
mates of spontaneous covariability for each patient. Continuous minutes of telemetry were itera-
tively sampled at random, without replacement, from artifact-free NREM sleep. Sampled minutes
of telemetry were appended and prepended with 1.5 seconds of data for filtering and analytic sig-
nal extraction. These data were removed prior to covariance estimation to avoid edge artifacts.
For each frequency, the coherence matrices were obtained as described above for the incoming
minute, $_{n+1}\Sigma_{n+1}$, the cumulatively sampled telemetry, $_1\Sigma_{n+1}$, and, for the second sampled minute on-
ward, the prior cumulatively sampled telemetry, $_1\Sigma_n$. The correlation matrix distance (CMD) (*Herdin*
et al., 2005), was calculated between $_1\Sigma_n$ and $_1\Sigma_{n+1}$, and between $_1\Sigma_n$ and $_{n+1}\Sigma_{n+1}$. Starting with the
third sampled minute, incoming minutes were rejected if the CMD between $_1\Sigma_n$ and $_{n+1}\Sigma_{n+1}$, was
more than 3 standard deviations greater than average CMD between $_1\Sigma_n$ and $_{n+1}\Sigma_{n+1}$ for all previ-
ously sampled minutes. The coherence matrix was considered to have converged when the CMD
between $_1\Sigma_n$ and $_1\Sigma_{n+1}$ was less than 1×10^{-5} and the cumulative cross-spectral matrix was then re-
tained. To dilute the influence of the first sampled minute, this procedure was repeated ten times,
and the mean of the resulting cross-spectral matrices was used for subsequent analyses.

Statistical analyses and model fitting

The effect of inter-parcel fiber distance on coherence was assessed by fitting a linear mixed-effects
model: log(coherence) distance + 1 + (1|patient), which accounted for the fixed effects of the pre-
dictor variable distance and an intercept term, as well the random effect intercept term grouped
by individual (this term was excluded for simulated data). Because of the natural log transforma-
tion of the response variable coherence, the linear model is equivalent to an exponential model for
the untransformed distance predictor. Therefore, we report the modeled slopes and intercepts as
length constants (λ) and scaling terms (α, maximum coherence), respectively. Separate fits were
made for each narrowband frequency. Because the parameters of the exponential trend appeared
to change at distances greater that 100 mm in experimental, especially for higher frequencies, sep-
arate models were fit to electrode pairs 0-100 mm distant, >100 mm, as well as to all pairs. Despite
having twice as many parameters, the two-domain model was favored over the one-domain model
by both Bayesian and Akaike information criteria for all frequencies and behavioral states. Only
the 0-100 mm model parameters are reported here for both experimental and simulation data.
All mixed effects linear models were fitted with matlab's fitlme. To avoid the inherent skewness
coherence coefficient distributions, coherence values were transformed into z-statistics (*Fisher,*
1921) before model fitting or averaging for display then reverted into coherence coefficients after
model fitting or averaging.

Finally, we take the 0.5 Hz lambda and the mean <2 Hz alpha values to describe the key rate
of decrease and max coherence in the slow wave range as the primary characteristics of each
coherence landscape to be compared across experimental and model data (Figure 9).

Acknowledgments

This research was supported by the National Science Foundation Graduate Research Fellowship Program under Grant No. DGE-2038238, and the National Institutes of Health (NIH) (grants 1R01MH117155, 1R01NS109553, 1R01MH125557). Data were provided [in part] by the Human Connectome Project, WU-Minn Consortium (Principal Investigators: David Van Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH Institutes and Centers that support the NIH Blueprint for Neuroscience Research; and by the McDonnell Center for Systems Neuroscience at Washington University.

References

Ball T, Kern M, Mutschler I, Aertsen A, Schulze-Bonhage A. Signal quality of simultaneously recorded invasive and non-invasive EEG. NeuroImage. 2009 Jul; 46(3):708–716. https://linkinghub.elsevier.com/retrieve/pii/S1053811909001827, doi: 10.1016/j.neuroimage.2009.02.028.

Bazhenov M, Rulkov NF, Timofeev I. Effect of synaptic connectivity on long-range synchronization of fast cortical oscillations. Journal of Neurophysiology. 2008 Sep; 100(3):1562–1575. doi: 10.1152/jn.90613.2008.

Bazhenov M, Timofeev I, Steriade M, Sejnowski T. Spiking-Bursting Activity in the Thalamic Reticular Nucleus Initiates Sequences of Spindle Oscillations in Thalamic Networks. Journal of Neurophysiology. 2000 Aug; 84(2):1076–1087. https://www.physiology.org/doi/10.1152/jn.2000.84.2.1076, doi: 10.1152/jn.2000.84.2.1076.

Bazhenov M, Timofeev I, Steriade M, Sejnowski TJ. Cellular and Network Models for Intrathalamic Augmenting Responses During 10-Hz Stimulation. Journal of Neurophysiology. 1998 May; 79(5):2730–2748. https://journals.physiology.org/doi/full/10.1152/jn.1998.79.5.2730, doi: 10.1152/jn.1998.79.5.2730, publisher: American Physiological Society.

Bazhenov M, Timofeev I, Steriade M, Sejnowski TJ. Computational Models of Thalamocortical Augmenting Responses. Journal of Neuroscience. 1998 Aug; 18(16):6444–6465. https://www.jneurosci.org/content/18/16/6444, doi: 10.1523/JNEUROSCI.18-16-06444.1998, publisher: Society for Neuroscience Section: ARTICLE.

Bazhenov M, Timofeev I, Steriade M, Sejnowski TJ. Model of Thalamocortical Slow-Wave Sleep Oscillations and Transitions to Activated States. The Journal of Neuroscience. 2002 Oct; 22(19):8691–8704. https://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.22-19-08691.2002, doi: 10.1523/JNEUROSCI.22-19-08691.2002.

Bernardi G, Siclari F, Handjaras G, Riedner BA, Tononi G. Local and Widespread Slow Waves in Stable NREM Sleep: Evidence for Distinct Regulation Mechanisms. Frontiers in Human Neuroscience. 2018; 12. https://www.frontiersin.org/articles/10.3389/fnhum.2018.00248.

Bonjean M, Baker T, Lemieux M, Timofeev I, Sejnowski T, Bazhenov M. Corticothalamic Feedback Controls Sleep Spindle Duration In Vivo. Journal of Neuroscience. 2011 Jun; 31(25):9124–9134. https://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.0077-11.2011, doi: 10.1523/JNEUROSCI.0077-11.2011.

Bonjean M, Baker T, Bazhenov M, Cash S, Halgren E, Sejnowski T. Interactions between Core and Matrix Thalamocortical Projections in Human Sleep Spindle Synchronization. The Journal of Neuroscience. 2012 Apr; 32(15):5250–5263. https://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.6141-11.2012, doi: 10.1523/JNEUROSCI.6141-11.2012.

Burt JB, Demirtaş M, Eckner WJ, Navejar NM, Ji JL, Martin WJ, Bernacchia A, Anticevic A, Murray JD. Hierarchy of transcriptomic specialization across human cortex captured by structural neuroimaging topography. Nature Neuroscience. 2018 Sep; 21(9):1251–1259. doi: 10.1038/s41593-018-0195-0.

Cakan C, Dimulescu C, Khakimova L, Obst D, Flöel A, Obermayer K. Spatiotemporal Patterns of Adaptation-Induced Slow Oscillations in a Whole-Brain Model of Slow-Wave Sleep. Frontiers in Computational Neuroscience. 2022; 15. https://www.frontiersin.org/articles/10.3389/fncom.2021.800101.

Campbell SS, Tobler I. Animal sleep: A review of sleep duration across phylogeny. Neuroscience & Biobehavioral Reviews. 1984 Sep; 8(3):269–300. https://linkinghub.elsevier.com/retrieve/pii/014976348490054X, doi: 10.1016/0149-7634(84)90054-X.

Chen JY, Chauvette S, Skorheim S, Timofeev I, Bazhenov M. Interneuron-mediated inhibition synchronizes neuronal activity during slow oscillation. The Journal of Physiology. 2012; 590(16):3987–4010. https://onlinelibrary.wiley.com/doi/abs/10.1113/jphysiol.2012.227462, doi: 10.1113/jphysiol.2012.227462, _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1113/jphysiol.2012.227462.

Compte A, Sanchez-Vives MV, McCormick DA, Wang XJ. Cellular and Network Mechanisms of Slow Oscillatory Activity (<1 Hz) and Wave Propagations in a Cortical Network Model. Journal of Neurophysiology. 2003 May; 89(5):2707–2725. https://www.physiology.org/doi/10.1152/jn.00845.2002, doi: 10.1152/jn.00845.2002.

Crunelli V, Hughes SW. The slow (<1 Hz) rhythm of non-REM sleep: a dialogue between three cardinal oscillators. Nature Neuroscience. 2010 Jan; 13(1):9–17. https://www.nature.com/articles/nn.2445, doi: 10.1038/nn.2445, number: 1 Publisher: Nature Publishing Group.

Csercsa R, Dombovári B, Fabó D, Wittner L, Erőss L, Entz L, Sólyom A, Rásonyi G, Szűcs A, Kelemen A, Jakus R, Juhos V, Grand L, Magony A, Halász P, Freund TF, Maglóczky Z, Cash SS, Papp L, Karmos G, et al. Laminar analysis of slow wave activity in humans. Brain. 2010 Sep; 133(9):2814–2829. https://doi.org/10.1093/brain/awq169, doi: 10.1093/brain/awq169.

Destexhe A, Bal T, McCormick DA, Sejnowski TJ. Ionic mechanisms underlying synchronized oscillations and propagating waves in a model of ferret thalamic slices. Journal of Neurophysiology. 1996 Sep; 76(3):2049–2070. https://journals.physiology.org/doi/abs/10.1152/jn.1996.76.3.2049, doi: 10.1152/jn.1996.76.3.2049, publisher: American Physiological Society.

Destexhe A, Contreras D, Sejnowski TJ, Steriade M. A model of spindle rhythmicity in the isolated thalamic reticular nucleus. Journal of Neurophysiology. 1994 Aug; 72(2):803–818. https://www.physiology.org/doi/10.1152/jn.1994.72.2.803, doi: 10.1152/jn.1994.72.2.803.

Destexhe A. Self-sustained asynchronous irregular states and Up-Down states in thalamic, cortical and thalamocortical networks of nonlinear integrate-and-fire neurons. Journal of Computational Neuroscience. 2009 Dec; 27(3):493–506. doi: 10.1007/s10827-009-0164-4.

Destexhe A, Contreras D, Steriade M. Spatiotemporal Analysis of Local Field Potentials and Unit Discharges in Cat Cerebral Cortex during Natural Wake and Sleep States. The Journal of Neuroscience. 1999 Jun; 19(11):4595–4608. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6782626/, doi: 10.1523/JNEUROSCI.19-11-04595.1999.

Dickey CW, Verzhbinsky IA, Jiang X, Rosen BQ, Kajfez S, Stedelin B, Shih JJ, Ben-Haim S, Raslan AM, Eskandar EN, Gonzalez-Martinez J, Cash SS, Halgren E. Widespread ripples synchronize human cortical activity during sleep, waking, and memory recall. Proceedings of the National Academy of Sciences. 2022 Jul; 119(28):e2107797119. https://pnas.org/doi/full/10.1073/pnas.2107797119, doi: 10.1073/pnas.2107797119.

Donoghue T, Haller M, Peterson EJ, Varma P, Sebastian P, Gao R, Noto T, Lara AH, Wallis JD, Knight RT, Shestyuk A, Voytek B. Parameterizing neural power spectra into periodic and aperiodic components. Nature Neuroscience. 2020 Dec; 23(12):1655–1665. https://www.nature.com/articles/s41593-020-00744-x, doi: 10.1038/s41593-020-00744-x, number: 12 Publisher: Nature Publishing Group.

Engel TA, Steinmetz NA, Gieselmann MA, Thiele A, Moore T, Boahen K. Selective modulation of cortical state during spatial attention. Science. 2016 Dec; 354(6316):1140–1144. https://www.science.org/doi/full/10.1126/science.aag1420, doi: 10.1126/science.aag1420, publisher: American Association for the Advancement of Science.

Eschenko O, Mölle M, Born J, Sara SJ. Elevated Sleep Spindle Density after Learning or after Retrieval in Rats. Journal of Neuroscience. 2006 Dec; 26(50):12914–12920. https://www.jneurosci.org/content/26/50/12914, doi: 10.1523/JNEUROSCI.3175-06.2006, publisher: Society for Neuroscience Section: Articles.

Fedorov A, Beichel R, Kalpathy-Cramer J, Finet J, Fillion-Robin JC, Pujol S, Bauer C, Jennings D, Fennessy F, Sonka M, Buatti J, Aylward S, Miller JV, Pieper S, Kikinis R. 3D Slicer as an image computing platform for the Quantitative Imaging Network. Magnetic Resonance Imaging. 2012 Nov; 30(9):1323–1341. https://linkinghub.elsevier.com/retrieve/pii/S0730725X12001816, doi: 10.1016/j.mri.2012.05.001.

Fischl B. FreeSurfer. Neuroimage. 2012; 62(2):774–781. ISBN: 1053-8119 Publisher: Elsevier.

Fischl B, Sereno MI, Tootell RBH, Dale AM. High-resolution intersubject averaging and a coordinate system for the cortical surface. Human Brain Mapping. 1999; 8(4):272–284. https://onlinelibrary.wiley.com/doi/10.1002/(SICI)1097-0193(1999)8:4<272::AID-HBM10>3.0.CO;2-4, doi: 10.1002/(SICI)1097-0193(1999)8:4<272::AID-HBM10>3.0.CO;2-4.

Fisher RA. On the" Probable Error" of a Coefficient of Correlation Deduced from a Small Sample. Metron. 1921; 1:3–32.

Funk CM, Honjoh S, Rodriguez AV, Cirelli C, Tononi G. Local slow waves in superficial layers of primary cortical areas during REM sleep. Current biology : CB. 2016 Feb; 26(3):396–403. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4747819/, doi: 10.1016/j.cub.2015.11.062.

Glasser MF, Coalson TS, Robinson EC, Hacker CD, Harwell J, Yacoub E, Ugurbil K, Andersson J, Beckmann CF, Jenkinson M, Smith SM, Van Essen DC. A multi-modal parcellation of human cerebral cortex. Nature. 2016 Aug; 536(7615):171–178. https://www.nature.com/articles/nature18933, doi: 10.1038/nature18933.

Glasser MF, Essen DCV. Mapping Human Cortical Areas In Vivo Based on Myelin Content as Revealed by T1- and T2-Weighted MRI. Journal of Neuroscience. 2011 Aug; 31(32):11597–11616. https://www.jneurosci.org/content/31/32/11597, doi: 10.1523/JNEUROSCI.2180-11.2011, publisher: Society for Neuroscience Section: Articles.

Goldman JS, Kusch L, Aquilue D, Yalçınkaya BH, Depannemaecker D, Ancourt K, Nghiem TAE, Jirsa V, Destexhe A. A comprehensive neural simulation of slow-wave sleep and highly responsive wakefulness dynamics. Frontiers in Computational Neuroscience. 2023 Jan; 16:1058957. https://www.frontiersin.org/articles/10.3389/fncom.2022.1058957/full, doi: 10.3389/fncom.2022.1058957.

Hanlon EC, Faraguna U, Vyazovskiy VV, Tononi G, Cirelli C. Effects of Skilled Training on Sleep Slow Wave Activity and Cortical Gene Expression in the Rat. Sleep. 2009 Jun; 32(6):719–729. https://doi.org/10.1093/sleep/32.6.719, doi: 10.1093/sleep/32.6.719.

Hawkins J, Ahmad S. Why Neurons Have Thousands of Synapses, a Theory of Sequence Memory in Neocortex. Frontiers in Neural Circuits. 2016 Mar; 10:23. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4811948/, doi: 10.3389/fncir.2016.00023.

Herdin M, Czink N, Ozcelik H, Bonek E. Correlation Matrix Distance, a Meaningful Measure for Evaluation of Non-Stationary MIMO Channels. In: *2005 IEEE 61st Vehicular Technology Conference*, vol. 1 Stockholm, Sweden: IEEE; 2005. p. 136–140. http://ieeexplore.ieee.org/document/1543265/, doi: 10.1109/VETECS.2005.1543265.

Hill S, Tononi G. Modeling Sleep and Wakefulness in the Thalamocortical System. Journal of Neurophysiology. 2005 Mar; 93(3):1671–1698. https://journals.physiology.org/doi/full/10.1152/jn.00915.2004, doi: 10.1152/jn.00915.2004, publisher: American Physiological Society.

Hodgkin AL, Huxley AF. A quantitative description of membrane current and its application to conduction and excitation in nerve. The Journal of Physiology. 1952 Aug; 117(4):500–544. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1392413/, doi: 10.1113/jphysiol.1952.sp004764, publisher: Wiley-Blackwell.

Huber R, Felice Ghilardi M, Massimini M, Tononi G. Local sleep and learning. Nature. 2004 Jul; 430(6995):78–81. https://www.nature.com/articles/nature02663, doi: 10.1038/nature02663, number: 6995 Publisher: Nature Publishing Group.

Hung CS, Sarasso S, Ferrarelli F, Riedner B, Ghilardi MF, Cirelli C, Tononi G. Local experience-dependent changes in the wake EEG after prolonged wakefulness. Sleep. 2013; 36(1):59–72. https://academic.oup.com/sleep/article-abstract/36/1/59/2656920, publisher: Oxford University Press.

Iber C, Ancoli-Israel S, Chesson A, Quan S. Manual for the Scoring of Sleep and Associated Events: Rules, Terminology and Technical Specifications. American Academy of Sleep Medicine. 2007; 59.

Isomura Y, Sirota A, Özen S, Montgomery S, Mizuseki K, Henze DA, Buzsáki G. Integration and Segregation of Activity in Entorhinal-Hippocampal Subregions by Neocortical Slow Oscillations. Neuron. 2006 Dec; 52(5):871–882. https://linkinghub.elsevier.com/retrieve/pii/S0896627306008221, doi: 10.1016/j.neuron.2006.10.023.

Ji D, Wilson MA. Coordinated memory replay in the visual cortex and hippocampus during sleep. Nature Neuroscience. 2007 Jan; 10(1):100–107. https://www.nature.com/articles/nn1825, doi: 10.1038/nn1825.

Johnson H, Harris G, Williams K. BRAINSFit: mutual information rigid registrations of whole-brain 3D images, using the insight toolkit. Insight J. 2007; 57(1):1–10.

Katsuki F, Gerashchenko D, Brown RE. Alterations of sleep oscillations in Alzheimer's disease: A potential role for GABAergic neurons in the cortex, hippocampus, and thalamus. Brain Research Bulletin. 2022 Sep; 187:181–198. https://linkinghub.elsevier.com/retrieve/pii/S0361923022001654, doi: 10.1016/j.brainresbull.2022.07.002.

van Kempen J, Gieselmann MA, Boyd M, Steinmetz NA, Moore T, Engel TA, Thiele A. Top-down coordination of local cortical state during selective attention. Neuron. 2021; 109(5):894–904. https://www.cell.com/neuron/pdf/S0896-6273(20)30995-8.pdf, publisher: Elsevier.

Kim J, Gulati T, Ganguly K. Competing Roles of Slow Oscillations and Delta Waves in Memory Consolidation versus Forgetting. Cell. 2019 Oct; 179(2):514–526.e13. https://linkinghub.elsevier.com/retrieve/pii/S0092867419309596, doi: 10.1016/j.cell.2019.08.040.

Komarov M, Krishnan G, Chauvette S, Rulkov N, Timofeev I, Bazhenov M. New class of reduced computationally efficient neuronal models for large-scale simulations of brain dynamics. Journal of Computational Neuroscience. 2018 Feb; 44(1):1–24. http://link.springer.com/10.1007/s10827-017-0663-7, doi: 10.1007/s10827-017-0663-7.

Korman M, Doyon J, Doljansky J, Carrier J, Dagan Y, Karni A. Daytime sleep condenses the time course of motor memory consolidation. Nature Neuroscience. 2007 Sep; 10(9):1206–1213. https://www.nature.com/articles/nn1959, doi: 10.1038/nn1959, number: 9 Publisher: Nature Publishing Group.

Krishnan GP, Chauvette S, Shamie I, Soltani S, Timofeev I, Cash SS, Halgren E, Bazhenov M. Cellular and neurochemical basis of sleep stages in the thalamocortical network. eLife. 2016 Nov; 5:e18607. https://elifesciences.org/articles/18607, doi: 10.7554/eLife.18607.

Krishnan GP, Rosen BQ, Chen JY, Muller L, Sejnowski TJ, Cash SS, Halgren E, Bazhenov M. Thalamocortical and intracortical laminar connectivity determines sleep spindle properties. PLOS Computational Biology. 2018 Jun; 14(6):e1006171. https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1006171, doi: 10.1371/journal.pcbi.1006171, publisher: Public Library of Science.

Kurth S, Riedner BA, Dean DC, O'Muircheartaigh J, Huber R, Jenni OG, Deoni SCL, LeBourgeois MK. Traveling Slow Oscillations During Sleep: A Marker of Brain Connectivity in Childhood. Sleep. 2017 Sep; 40(9). https://academic.oup.com/sleep/article/doi/10.1093/sleep/zsx121/3953857, doi: 10.1093/sleep/zsx121.

Lee SH, Dan Y. Neuromodulation of Brain States. Neuron. 2012 Oct; 76(1):209–222. https://linkinghub.elsevier.com/retrieve/pii/S0896627312008203, doi: 10.1016/j.neuron.2012.09.012.

Lemieux M, Chen JY, Lonjers P, Bazhenov M, Timofeev I. The Impact of Cortical Deafferentation on the Neocortical Slow Oscillation. The Journal of Neuroscience. 2014 Apr; 34(16):5689–5703. https://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.1156-13.2014, doi: 10.1523/JNEUROSCI.1156-13.2014.

Leung LC, Mourrain P. Sleep: Short Sleepers Should Keep Count of Their Hypocretin Neurons. Current Biology. 2018 May; 28(9):R558–R560. https://linkinghub.elsevier.com/retrieve/pii/S0960982218303075, doi: 10.1016/j.cub.2018.03.006.

Levenstein D, Buzsáki G, Rinzel J. NREM sleep in the rodent neocortex and hippocampus reflects excitable dynamics. Nature Communications. 2019 Jun; 10(1):2478. https://www.nature.com/articles/s41467-019-10327-5, doi: 10.1038/s41467-019-10327-5, publisher: Nature Publishing Group.

Liu Y, Yu C, Zhang X, Liu J, Duan Y, Alexander-Bloch AF, Liu B, Jiang T, Bullmore E. Impaired Long Distance Functional Connectivity and Weighted Network Architecture in Alzheimer's Disease. Cerebral Cortex. 2014 Jun; 24(6):1422–1435. https://academic.oup.com/cercor/article-lookup/doi/10.1093/cercor/bhs410, doi: 10.1093/cercor/bhs410.

London M, Segev I. Synaptic scaling in vitro and in vivo. Nature Neuroscience. 2001 Sep; 4(9):853–854. https://www.nature.com/articles/nn0901-853, doi: 10.1038/nn0901-853, number: 9 Publisher: Nature Publishing Group.

Mak-McCully RA, Rolland M, Sargsyan A, Gonzalez C, Magnin M, Chauvel P, Rey M, Bastuji H, Halgren E. Coordination of cortical and thalamic activity during non-REM sleep in humans. Nature Communications. 2017 May; 8(1):15499. https://www.nature.com/articles/ncomms15499, doi: 10.1038/ncomms15499.

Mander BA, Rao V, Lu B, Saletin JM, Lindquist JR, Ancoli-Israel S, Jagust W, Walker MP. Prefrontal atrophy, disrupted NREM slow waves and impaired hippocampal-dependent memory in aging. Nature Neuroscience. 2013 Mar; 16(3):357–364. https://www.nature.com/articles/nn.3324, doi: 10.1038/nn.3324.

Marcus DS, Harms MP, Snyder AZ, Jenkinson M, Wilson JA, Glasser MF, Barch DM, Archie KA, Burgess GC, Ramaratnam M, Hodge M, Horton W, Herrick R, Olsen T, McKay M, House M, Hileman M, Reid E, Harwell J, Coalson T, et al. Human Connectome Project informatics: Quality control, database services, and data visualization. NeuroImage. 2013 Oct; 80:202–219. https://linkinghub.elsevier.com/retrieve/pii/S1053811913005776, doi: 10.1016/j.neuroimage.2013.05.077.

Markov NT, Ercsey-Ravasz M, Van Essen DC, Knoblauch K, Toroczkai Z, Kennedy H. Cortical High-Density Counterstream Architectures. Science. 2013 Nov; 342(6158):1238406. https://www.science.org/doi/full/10.1126/science.1238406, doi: 10.1126/science.1238406, publisher: American Association for the Advancement of Science.

Mascetti L, Muto V, Matarazzo L, Foret A, Ziegler E, Albouy G, Sterpenich V, Schmidt C, Degueldre C, Leclercq Y, Phillips C, Luxen A, Vandewalle G, Vogels R, Maquet P, Balteau E. The Impact of Visual Perceptual Learning on Sleep and Local Slow-Wave Initiation. The Journal of Neuroscience. 2013 Feb; 33(8):3323–3331. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6619511/, doi: 10.1523/JNEUROSCI.0763-12.2013.

McCormick DA, Huguenard JR. A model of the electrophysiological properties of thalamocortical relay neurons. Journal of Neurophysiology. 1992 Oct; 68(4):1384–1400. https://www.physiology.org/doi/10.1152/jn.1992.68.4.1384, doi: 10.1152/jn.1992.68.4.1384.

McKillop LE, Fisher SP, Cui N, Peirson SN, Foster RG, Wafford KA, Vyazovskiy VV. Effects of Aging on Cortical Neural Dynamics and Local Sleep Homeostasis in Mice. Journal of Neuroscience. 2018 Apr; 38(16):3911–3928. https://www.jneurosci.org/content/38/16/3911, doi: 10.1523/JNEUROSCI.2513-17.2018, publisher: Society for Neuroscience Section: Research Articles.

Miyawaki H, Watson BO, Diba K. Neuronal firing rates diverge during REM and homogenize during non-REM. Scientific Reports. 2019 Jan; 9:689. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6345798/, doi: 10.1038/s41598-018-36710-8.

Mölle M, Eschenko O, Gais S, Sara SJ, Born J. The influence of learning on sleep slow oscillations and associated spindles and ripples in humans and rats. European Journal of Neuroscience. 2009 Mar; 29(5):1071–1081. https://onlinelibrary.wiley.com/doi/10.1111/j.1460-9568.2009.06654.x, doi: 10.1111/j.1460-9568.2009.06654.x.

Nazari M, Karimi Abadchi J, Naghizadeh M, Bermudez-Contreras EJ, McNaughton BL, Tatsuno M, Mohajerani MH. Regional variation in cholinergic terminal activity determines the non-uniform occurrence of cortical slow waves during REM sleep in mice. Cell Reports. 2023 May; 42(5):112450. https://linkinghub.elsevier.com/retrieve/pii/S2211124723004618, doi: 10.1016/j.celrep.2023.112450.

Neudorfer C, Kultas-Ilinsky K, Ilinsky I, Paschen S, Helmers AK, Cosgrove GR, Richardson RM, Horn A, Deuschl G. The role of the motor thalamus in deep brain stimulation for essential tremor. Neurotherapeutics. 2024 Apr; 21(3):e00313. https://www.sciencedirect.com/science/article/pii/S1878747923020299, doi: 10.1016/j.neurot.2023.e00313.

Nir Y, Staba RJ, Andrillon T, Vyazovskiy VV, Cirelli C, Fried I, Tononi G. Regional slow waves and spindles in human sleep. Neuron. 2011 Apr; 70(1):153–169. doi: 10.1016/j.neuron.2011.02.043.

Nobili L, Ferrara M, Moroni F, De Gennaro L, Russo GL, Campus C, Cardinale F, De Carli F. Dissociated wake-like and sleep-like electro-cortical activity during sleep. NeuroImage. 2011 Sep; 58(2):612–619. https://www.sciencedirect.com/science/article/pii/S1053811911006501, doi: 10.1016/j.neuroimage.2011.06.032.

Peter-Derex L, Magnin M, Bastuji H. Heterogeneity of arousals in human sleep: A stereo-electroencephalographic study. NeuroImage. 2015 Dec; 123:229–244. https://www.sciencedirect.com/science/article/pii/S1053811915006771, doi: 10.1016/j.neuroimage.2015.07.057.

Pfaff D, Ribeiro A, Matthews J, Kow L. *Concepts and Mechanisms of Generalized Central Nervous System Arousal*. Annals of the New York Academy of Sciences. 2008 May; 1129(1):11–25. https://nyaspubs.onlinelibrary.wiley.com/doi/10.1196/annals.1417.019, doi: 10.1196/annals.1417.019.

Piantoni G, Poil SS, Linkenkaer-Hansen K, Verweij IM, Ramautar JR, Van Someren EJW, Van Der Werf YD. Individual Differences in White Matter Diffusion Affect Sleep Oscillations. The Journal of Neuroscience. 2013 Jan; 33(1):227–233. https://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.2030-12.2013, doi: 10.1523/JNEUROSCI.2030-12.2013.

Pugin F, Metz AJ, Wolf M, Achermann P, Jenni OG, Huber R. Local increase of sleep slow wave activity after three weeks of working memory training in children and adolescents. Sleep. 2015 Apr; 38(4):607–614. doi: 10.5665/sleep.4580.

Quercia A, Zappasodi F, Committeri G, Ferrara M. Local Use-Dependent Sleep in Wakefulness Links Performance Errors to Learning. Frontiers in Human Neuroscience. 2018; 12. https://www.frontiersin.org/articles/10.3389/fnhum.2018.00122.

Rasch B, Born J. About Sleep's Role in Memory. Physiological Reviews. 2013 Apr; 93(2):681–766. https://journals.physiology.org/doi/full/10.1152/physrev.00032.2012, doi: 10.1152/physrev.00032.2012, publisher: American Physiological Society.

Rasch B, Büchel C, Gais S, Born J. Odor Cues During Slow-Wave Sleep Prompt Declarative Memory Consolidation. Science. 2007 Mar; 315(5817):1426–1429. https://www.science.org/doi/10.1126/science.1138581, doi: 10.1126/science.1138581.

Riedner BA, Vyazovskiy VV, Huber R, Massimini M, Esser S, Murphy M, Tononi G. Sleep Homeostasis and Cortical Synchronization: III. A High-Density EEG Study of Sleep Slow Waves in Humans. Sleep. 2007 Dec; 30(12):1643–1657. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2276133/.

Rockland KS. What do we know about laminar connectivity? NeuroImage. 2019 Aug; 197:772–784. doi: 10.1016/j.neuroimage.2017.07.032.

Rosanova M, Ulrich D. Pattern-Specific Associative Long-Term Potentiation Induced by a Sleep Spindle-Related Spike Train. The Journal of Neuroscience. 2005 Oct; 25(41):9398–9405. https://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.2149-05.2005, doi: 10.1523/JNEUROSCI.2149-05.2005.

Rosen BQ, Krishnan GP, Sanda P, Komarov M, Sejnowski T, Rulkov N, Ulbert I, Eross L, Madsen J, Devinsky O, Doyle W, Fabo D, Cash S, Bazhenov M, Halgren E. Simulating human sleep spindle MEG and EEG from ion channel and circuit level dynamics. Journal of Neuroscience Methods. 2019 Mar; 316:46–57. https://www.sciencedirect.com/science/article/pii/S0165027018303108, doi: 10.1016/j.jneumeth.2018.10.002.

Rosen BQ, Halgren E. A Whole-Cortex Probabilistic Diffusion Tractography Connectome. eneuro. 2021 Jan; 8(1):ENEURO.0416–20.2020. https://www.eneuro.org/lookup/doi/10.1523/ENEURO.0416-20.2020, doi: 10.1523/ENEURO.0416-20.2020.

Rulkov NF, Timofeev I, Bazhenov M. Oscillations in Large-Scale Cortical Networks: Map-Based Model. Journal of Computational Neuroscience. 2004 Sep; 17(2):203–223. http://link.springer.com/10.1023/B:JCNS.0000037683.55688.7e, doi: 10.1023/B:JCNS.0000037683.55688.7e.

Rulkov NF. Modeling of spiking-bursting neural behavior using two-dimensional map. Physical Review E. 2002 Apr; 65(4):041922. https://link.aps.org/doi/10.1103/PhysRevE.65.041922, doi: 10.1103/PhysRevE.65.041922, publisher: American Physical Society.

Rulkov NF, Bazhenov M. Oscillations and Synchrony in Large-scale Cortical Network Models. Journal of Biological Physics. 2008 Aug; 34(3-4):279–299. http://link.springer.com/10.1007/s10867-008-9079-y, doi: 10.1007/s10867-008-9079-y.

Sanchez-Vives MV, McCormick DA. Cellular and network mechanisms of rhythmic recurrent activity in neocortex. Nature Neuroscience. 2000 Oct; 3(10):1027–1034. https://www.nature.com/articles/nn1000_1027, doi: 10.1038/79848, number: 10 Publisher: Nature Publishing Group.

Schüller T, Huys D, Kohl S, Visser-Vandewalle V, Dembek TA, Kuhn J, Baldermann JC, Smith EE. Thalamic deep brain stimulation for tourette syndrome increases cortical beta activity. Brain Stimulation. 2024 Mar; 17(2):197–201. https://www.sciencedirect.com/science/article/pii/S1935861X24000123, doi: 10.1016/j.brs.2024.01.011.

Shepherd GMG, Yamawaki N. Untangling the cortico-thalamo-cortical loop: cellular pieces of a knotty circuit puzzle. Nature Reviews Neuroscience. 2021 Jul; 22(7):389–406. https://www.nature.com/articles/s41583-021-00459-3, doi: 10.1038/s41583-021-00459-3.

Siclari F, Bernardi G, Riedner BA, LaRocque JJ, Benca RM, Tononi G. Two Distinct Synchronization Processes in the Transition to Sleep: A High-Density Electroencephalographic Study. Sleep. 2014 Oct; 37(10):1621–1637. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4173919/, doi: 10.5665/sleep.4070.

Siclari F, Tononi G. Local aspects of sleep and wakefulness. Current Opinion in Neurobiology. 2017 Jun; 44:222–227. https://linkinghub.elsevier.com/retrieve/pii/S0959438816302410, doi: 10.1016/j.conb.2017.05.008.

Soltani S, Chauvette S, Bukhtiyarova O, Lina JM, Dubé J, Seigneur J, Carrier J, Timofeev I. Sleep–Wake Cycle in Young and Older Mice. Frontiers in Systems Neuroscience. 2019; 13. https://www.frontiersin.org/articles/10.3389/fnsys.2019.00051.

Steriade M, Nunez A, Amzica F. Intracellular analysis of relations between the slow (< 1 Hz) neocortical oscillation and other sleep rhythms of the electroencephalogram. Journal of Neuroscience. 1993 Aug; 13(8):3266–3283. https://www.jneurosci.org/content/13/8/3266, doi: 10.1523/JNEUROSCI.13-08-03266.1993, publisher: Society for Neuroscience Section: Articles.

Steriade M, Timofeev I, Grenier F. Natural Waking and Sleep States: A View From Inside Neocortical Neurons. Journal of Neurophysiology. 2001 May; 85(5):1969–1985. https://journals.physiology.org/doi/full/10.1152/jn.2001.85.5.1969, doi: 10.1152/jn.2001.85.5.1969, publisher: American Physiological Society.

Steriade M, Jones EG, Llinás RR. Thalamic oscillations and signaling. Thalamic oscillations and signaling, Oxford, England: John Wiley & Sons; 1990. Pages: xiii, 431.

Steriade M, McCormick DA, Sejnowski TJ. Thalamocortical Oscillations in the Sleeping and Aroused Brain. Science. 1993 Oct; 262(5134):679–685. https://www.science.org/doi/10.1126/science.8235588, doi: 10.1126/science.8235588, publisher: American Association for the Advancement of Science.

Steriade MM, McCarley RW. Brainstem Control of Wakefulness and Sleep. Springer Science & Business Media; 2013. Google-Books-ID: X5juBwAAQBAJ.

Stevens CF. Quantal release of neurotransmitter and long-term potentiation. Cell. 1993 Jan; 72:55–63. https://www.sciencedirect.com/science/article/pii/S0092867405800285, doi: 10.1016/S0092-8674(05)80028-5.

Suarez-Perez A, Gabriel G, Rebollo B, Illa X, Guimerà-Brunet A, Hernández-Ferrer J, Martínez MT, Villa R, Sanchez-Vives MV. Quantification of Signal-to-Noise Ratio in Cerebral Cortex Recordings Using Flexible MEAs With Co-localized Platinum Black, Carbon Nanotubes, and Gold Electrodes. Frontiers in Neuroscience. 2018 Nov; 12:862. https://www.frontiersin.org/article/10.3389/fnins.2018.00862/full, doi: 10.3389/fnins.2018.00862.

Thomas CW, Guillaumin MC, McKillop LE, Achermann P, Vyazovskiy VV. Global sleep homeostasis reflects temporally and spatially integrated local cortical neuronal activity. eLife. 2020 Jul; 9:e54148. https://doi.org/10.7554/eLife.54148, doi: 10.7554/eLife.54148, publisher: eLife Sciences Publications, Ltd.

Timofeev I, Grenier F, Bazhenov M, Sejnowski TJ, Steriade M. Origin of Slow Cortical Oscillations in Deafferented Cortical Slabs. Cerebral Cortex. 2000 Dec; 10(12):1185–1199. https://doi.org/10.1093/cercor/10.12.1185, doi: 10.1093/cercor/10.12.1185.

Timofeev I, Steriade M. Low-frequency rhythms in the thalamus of intact-cortex and decorticated cats. Journal of Neurophysiology. 1996 Dec; 76(6):4152–4168. https://journals.physiology.org/doi/abs/10.1152/jn.1996.76.6.4152, doi: 10.1152/jn.1996.76.6.4152, publisher: American Physiological Society.

Timofeev I, Chauvette S. Sleep slow oscillation and plasticity. Current Opinion in Neurobiology. 2017 Jun; 44:116–126. https://www.sciencedirect.com/science/article/pii/S0959438817300417, doi: 10.1016/j.conb.2017.03.019.

Tomasi D, Volkow ND. Aging and functional brain networks. Molecular Psychiatry. 2012 May; 17(5):549–558. https://www.nature.com/articles/mp201181, doi: 10.1038/mp.2011.81.

Van Essen DC, Smith SM, Barch DM, Behrens TEJ, Yacoub E, Ugurbil K. The WU-Minn Human Connectome Project: An overview. NeuroImage. 2013 Oct; 80:62–79. https://www.sciencedirect.com/science/article/pii/S1053811913005351, doi: 10.1016/j.neuroimage.2013.05.041.

Volgushev M, Chauvette S, Timofeev I. Long-range correlation of the membrane potential in neocortical neurons during slow oscillation. In: *Progress in Brain Research*, vol. 193 Elsevier; 2011.p. 181–199. https://linkinghub.elsevier.com/retrieve/pii/B9780444538390000120, doi: 10.1016/B978-0-444-53839-0.00012-0.

Vyazovskiy VV, Olcese U, Hanlon EC, Nir Y, Cirelli C, Tononi G. Local sleep in awake rats. Nature. 2011 Apr; 472(7344):443–447. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3085007/, doi: 10.1038/nature10009.

Vyazovskiy VV, Olcese U, Lazimy YM, Faraguna U, Esser SK, Williams JC, Cirelli C, Tononi G. Cortical Firing and Sleep Homeostasis. Neuron. 2009 Sep; 63(6):865–878. https://linkinghub.elsevier.com/retrieve/pii/S0896627309006370, doi: 10.1016/j.neuron.2009.08.024.

Vyazovskiy VV, Riedner BA, Cirelli C, Tononi G. Sleep Homeostasis and Cortical Synchronization: II. A Local Field Potential Study of Sleep Slow Waves in the Rat. Sleep. 2007 Dec; 30(12):1631–1642. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2276140/.

Wierzynski CM, Lubenov EV, Gu M, Siapas AG. State-Dependent Spike-Timing Relationships between Hippocampal and Prefrontal Circuits during Sleep. Neuron. 2009 Feb; 61(4):587–596. https://linkinghub.elsevier.com/retrieve/pii/S0896627309000786, doi: 10.1016/j.neuron.2009.01.011.

Wilckens KA, Ferrarelli F, Walker MP, Buysse DJ. Slow-Wave Activity Enhancement to Improve Cognition. Trends in Neurosciences. 2018 Jul; 41(7):470–482. https://www.cell.com/trends/neurosciences/abstract/S0166-2236(18)30064-X, doi: 10.1016/j.tins.2018.03.003, publisher: Elsevier.

Winkelman JW, Lecea LD. Sleep and neuropsychiatric illness. Neuropsychopharmacology. 2020 Jan; 45(1):1–2. https://www.nature.com/articles/s41386-019-0514-5, doi: 10.1038/s41386-019-0514-5.

Supplementary Information

Figure S1. Neuronal activity in all 6 layers during the Global SO model from Figure 2. Voltage traces in mV for two different cells are shown on the left panels, with individual spikes marked under each trace in the same color.

Figure S2. Activity of pyramidal, inhibitory and thalamic RE and TC neurons in baseline model in Figure 2B.Voltage traces in mV for two different cells are shown on the left panels, with individual spikes marked under each trace in the same color.

Figure S3. Activity of cortical cells in all layers in mixed model.Voltage traces in mV for two different cells are shown on the left panels, with individual spikes marked under each trace in the same color.

Figure S4. Activity of pyramidal, inhibitory, and thalamic cells in the mixed model. Voltage traces in mV for two different cells are shown on the left panels, with individual spikes marked under each trace in the same color.

Figure S5. Activity of pyramidal and inhibitory neurons in the Activity by Layer and Cell Type: Mixed local/global model in Figure S4, with the thalamus isolated from the cortex. Voltage traces in mV for two different cells are shown on the left panels, with individual spikes marked under each trace in the same color.

Figure S6. Local activity for $P = 0.1$, with 90% connections removed. A) Ten cortical areas with a 5mm radius, that were used to calculate local dynamics. B) Average membrane voltage of layer II neurons, as in Figure 3e.2. C-E) For each region in (A), subpanels show: (C) the single-cell voltage for two neurons in the area , (D) the local field potential (LFP) for the 5mm area, and (E) heatmap of individual voltages of all neurons in the area. Up states are largely synchronized across all 5 regions. F) Latency map for each Up state in the $P = 0.1$ simulation. Even with very sparse connectivity, Up states spread to the whole cortex. Participation was reduced uniformly to about 70% compare to nearly 100% participation when all connections are present (compare to Figure 2D)

Figure S7. Effect of Synaptic Delay, either via setting a uniform delay (A) or scaling the max delay (B) on SO dynamics. (A and B) Summary plots of frequency, amplitude, and onset / offset spread. Decreasing the max scaled delay to 0.1 ms (from 40 ms) has no effect, while increasing the max scaled delay up to 1 second shows only minor changes in frequency frequency and amplitude but notably increased onset/offset synchrony. Low uniform delays similarly showed no effect, with loss of amplitude and synchrony only seen at 30 ms (most delays were previously under 2 ms).

Figure S8. Activity maps and graph properties for the full-connectivity, Global SO network in Figure 2. Rows from top to bottom show (1) the percent of simulation time that each pyramidal cell in layer 2 spends in an Up state; (2) the mean onset/offset delay across Up states for each pyramidal

cell in layer 2; (3) the number of strongly/weakly connected components in the structural connectivity graph, constructed with each cortical column as a node and each directed edge weight as the number of synaptic connections from one column to another; (4) the normalized in/out-degree for each node in the graph; and (5) the normalized in/out-degree centrality for each node in the graph, defined as the number of synaptic connections to/from that column (see Graph Properties for details). Initiation zones (low mean onset delay) at the front and back and have generally lower in-degree and out-degree than areas in the middle.

Figure S9. Activity maps and graph properties for the $R = 2.5$mm network in Figure 5. Rows from top to bottom show (1) the percent of simulation time that each pyramidal cell in layer 2 spends in an Up state; (2) the mean onset/offset delay across Up states for each pyramidal cell in layer 2; (3) the number of strongly/weakly connected components in the structural connectivity graph, constructed with each cortical column as a node and each directed edge weight as the number of synaptic connections from one column to another; (4) the normalized in/out-degree for each node in the graph; and (5) the normalized in/out-degree centrality for each node in the graph, defined as the number of synaptic connections to/from that column (see Graph Properties for details). Regions with high percent time active have high in/out-degree and centrality

Table S1. Peaks and 3 dB attenuation bandwidths for the second-order resonator filters used to extract narrow-band signals.

Video S1. The base global slow wave sleep model over 30 seconds of simulated time. The top plot shows Layer 2 cell voltages across the cortex, while the bottom plot shows the average voltage trace.

Video S2. The slow wave sleep model with a connection density of P = 0.1 over 30 seconds of simulated time. The top plot shows Layer 2 cell voltages across the cortex, while the bottom plot shows the average voltage trace.

Video S3. The slow wave sleep model with a connection range of R = 2.5 mm over 20 seconds of simulated time. The top plot shows Layer 2 cell voltages across the cortex, while the bottom plot shows the average voltage trace.

Video S4. The slow wave sleep model with connections longer than 10mm reduced 5-fold over 105 seconds of simulated time. The top plot shows Layer 2 cell voltages across the cortex, while the bottom plot shows the average voltage trace.