1   **Graphical Abstract**
2

3
4
5
6   <u>**Highlights**</u>
7   1. In patients with hepatocellular carcinoma (HCC), tertiary lymphoid structures (TLS) are
8   induced by neoadjuvant immunotherapy and are associated with favorable clinical
9   outcomes.
10
11   2. TLS within the same tumor demonstrate extensive sharing of expanded granzyme K and
12   granzyme B-expressing CD8[+] T effector memory clonotypes, but the B cell repertoires of
13   individual TLS are almost wholly distinct, consistent with independent germinal center
14   reactions.
15
16   3. Within areas of viable tumor, mature TLS are characterized by high expression of CD21
17   and CD23, BCL6[+] germinal center B cells, and close interactions between DCLAMP[+]
18   mature dendritic cells and CXCR5[-]CXCR3[+] CD4 T peripheral helper cells within a T cell
19   zone adjacent to the B cell follicle.
20
21   4. Within areas of tumor regression, an involuted TLS morphology is identified that is notable
22   for dissolution of the B cell germinal center, retention of the T cell zone, and increased T
23   cell memory.
24

25
26

Immune landscape of tertiary lymphoid structures in hepatocellular carcinoma (HCC) treated with neoadjuvant immune checkpoint blockade

Daniel H. Shu[1,2], Won Jin Ho[1,2], Luciane T. Kagohara[1,2], Alexander Girgis[1,2,3], Sarah M. Shin[1], Ludmila Danilova[1], Jae W. Lee[4], Dimitrios N. Sidiropoulos[1,2], Sarah Mitchell,[1] Kabeer Munjal[1], Kathryn Howe[1], Kayla J. Bendinelli[1], Hanfei Qi[1], Guanglan Mo[1], Janelle Montagne[1,2], James M. Leatherman[1], Tamara Y. Lopez-Vidal[1,2], Qingfeng Zhu[2,4], Amanda L. Huff[1,2], Xuan Yuan[3], Alexei Hernandez[1], Erin M. Coyne[1], Neeha Zaidi[1,2], Daniel J. Zabransky[1,2], Logan L. Engle[5,6,7], Aleksandra Ogurtsova[5,6,7], Marina Baretti[1], Daniel Laheru[1,2], Jennifer N. Durham[1], Hao Wang[1], Robert Anders[2,4], Elizabeth M. Jaffee[1,2,7], Elana J. Fertig[1,2,3,8*], Mark Yarchoan[1,2,7*]

[1] Department of Oncology, Sidney Kimmel Comprehensive Cancer Center, Johns Hopkins University School of Medicine, Baltimore, Maryland.
[2] Convergence Institute, Johns Hopkins University, Baltimore, Maryland.
[3] Department of Biomedical Engineering, Johns Hopkins University School of Medicine, Baltimore, Maryland.
[4] Department of Pathology, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA.
[5] Department of Dermatology, Johns Hopkins University School of Medicine, Baltimore, Maryland.
[6] The Mark Foundation Center for Advanced Genomics and Imaging, Johns Hopkins University, Baltimore, Maryland.
[7] Bloomberg~Kimmel Institute for Cancer Immunotherapy and Sidney Kimmel Comprehensive Cancer Center, Johns Hopkins University, Baltimore, Maryland.
[8] Department of Applied Mathematics and Statistics, Johns Hopkins University Whiting School of Engineering, Baltimore, Maryland.

* Corresponding authors:
Mark Yarchoan (mark.yarchoan@jhmi.edu)
Elana J. Fertig (ejfertig@jhmi.edu)

Keywords: neoadjuvant immunotherapy, tertiary lymphoid structures, hepatocellular carcinoma, immune checkpoint inhibitors, single cell multiomics, imaging mass cytometry, single cell RNA, single cell TCR,

ABSTRACT

Neoadjuvant immunotherapy is thought to produce long-term remissions through induction of antitumor immune responses before removal of the primary tumor. Tertiary lymphoid structures (TLS), germinal center-like structures that can arise within tumors, may contribute to the establishment of immunological memory in this setting, but understanding of their role remains limited. Here, we investigated the contribution of TLS to antitumor immunity in hepatocellular carcinoma (HCC) treated with neoadjuvant immunotherapy. We found that neoadjuvant immunotherapy induced the formation of TLS, which were associated with superior pathologic response, improved relapse free survival, and expansion of the intratumoral T and B cell repertoire. While TLS in viable tumor displayed a highly active mature morphology, in areas of tumor regression we identified an involuted TLS morphology, which was characterized by dispersion of the B cell follicle and persistence of a T cell zone enriched for ongoing antigen presentation and T cell-mature dendritic cell interactions. Involuted TLS showed increased expression of T cell memory markers and expansion of CD8[+] cytotoxic and tissue resident memory clonotypes. Collectively, these data reveal the circumstances of TLS dissolution and suggest a functional role for late-stage TLS as sites of T cell memory formation after elimination of viable tumor.

INTRODUCTION

Immune checkpoint blockade (ICB) therapy has revolutionized the treatment of metastatic solid tumors, offering to a subset of patients the potential for sustained remissions beyond what was previously possible with chemotherapy alone.[1,2] For patients with early stage, non-metastatic disease, the role for ICB and ideal timing of its administration remains an area of intense clinical investigation. Recent clinical data in patients with melanoma suggest that neoadjuvant immunotherapy, in which ICB is administered prior to curative-intent resection of the primary tumor, may produce superior long-term outcomes compared to immunotherapy given after surgery.[3,4] Preclinical data suggest that these improved outcomes may be attributable to an elevated and sustained tumor-specific immune response that occurs when immunotherapy is initiated with the primary tumor *in situ*.[5,6] However, in human subjects it is not known where or by what means the establishment of immunological memory occurs.

Tertiary lymphoid structures (TLS), organized collections of B and T cells that can arise within solid tumors, have been associated with favorable responses to neoadjuvant ICB,[7–13] and it is hypothesized that TLS play a mechanistic role in promoting effective antitumor immunity. However, understanding of the structure, constituent immune populations, and life cycle of TLS in this treatment setting remains limited by the rarity of neoadjuvant clinical trial specimens, particularly in solid tumor types where the successes of immunotherapy have been modest, and paucity of animal models for TLS in cancer.[14,15] Thus patient samples from neoadjuvant clinical trials provide a unique opportunity to investigate the contribution of TLS to the development of antitumor immunity.

We previously reported an association between TLS and pathologic response in a phase 1 trial of patients with locally advanced HCC who received neoadjuvant nivolumab and cabozantinib.[8] Here, we evaluated the clinical and immunological characteristics of TLS in an expanded cohort of patients with locally advanced hepatocellular carcinoma (HCC) treated with neoadjuvant ICB. We find evidence that neoadjuvant ICB induces the formation of intratumoral TLS, and that high TLS density following neoadjuvant therapy is associated with superior pathologic response to treatment and relapse-free survival. Using a multiomics approach employing imaging mass cytometry, bulk TCR and BCR sequencing of microdissected TLS, and paired single cell RNA

and TCR sequencing, we identify key differences in the spatial and immunological landscape of TLS in areas of viable and nonviable tumor that suggest that the contribution of TLS to antitumor immunity in tumors treated with neoadjuvant immunotherapy varies significantly according to morphological stage and circumstance.

RESULTS

Neoadjuvant ICB in HCC induces intratumoral TLS

To determine the clinical significance of TLS in patients with HCC treated with neoadjuvant ICB, we identified patients from the Johns Hopkins Liver Cancer Biorepository who had undergone surgical resection of their primary tumor after receiving neoadjuvant ICB-based therapy for locally advanced HCC. In total, 19 patients were identified who received treatment between October 2019 and January 2022 (**Extended Data Table 1**). 11/19 (57.9%) were male, 13/19 (68.4%) had tumors with moderately differentiated histology, and 11/19 (57.9%) had a history of viral hepatitis. No patients had active viral hepatitis at the time of surgery. 14/19 (73.6%) received anti-PD-1 plus an oral tyrosine kinase inhibitor, 3/19 (15.8%) received anti-PD-1 monotherapy, 1/19 (5.2%) received combination anti-PD-1 and anti-CTLA-4 monoclonal antibody, and 1/19 (5.2%) received combination anti-PD-1/anti-CTLA-4 and oral TKI prior to resection of the primary tumor.

Since TLS are known to occur in treatment naïve HCC,[16] we first attempted to determine if TLS were present in the tumors of patients prior to receiving neoadjuvant ICB. 7/19 (36.8%) of patients had undergone pre-treatment fine needle biopsies prior to initiation of neoadjuvant therapy and no intratumoral TLS were identified in these specimens. Given the limited assessment of the tumor microenvironment provided by fine needle biopsy, we next identified a second cohort of HCC patients treated at our institution who had undergone surgical resection without receiving prior systemic therapy, which would serve as a control cohort. 17 patients were identified who had received upfront surgical resection for HCC between 2017 and 2022, from which 3 patients were excluded due to small tumor volume, poor tissue quality, or HCC etiology not represented by the treatment cohort. The 14 remaining patients (**Extended Data Table 2**), were similar to the neoadjuvant treatment cohort by age, sex, histologic grade, and etiology (**Table 1**).

Evaluation of TLS density in the two cohorts was performed by CD20 staining of resected FFPE tumor (**Fig. 1a**). TLS, which we defined as $CD20^+$ lymphoid aggregates with diameter greater than 150 µm, were classified as either peritumoral or intratumoral according to their location relative to the interface between tumor and normal adjacent parenchyma (**Fig. 1b**). TLS were observed in 7/14 (50%) of untreated tumors and 12/19 (63.2%) treated tumors. No significant difference was identified in total TLS density (0.08 ± 0.09 TLS/mm^2 versus 0.05 ± 0.10 TLS/mm^2, P = 0.42) or peritumoral TLS density (0.03 ± 0.05 TLS/mm^2 versus 0.04 ± 0.1 TLS/mm^2, P = 0.73) (**Extended Fig. 1a**), but intratumoral TLS density was significantly increased in treated patients compared to untreated controls (0.05 ± 0.08 TLS/mm^2 versus 0.01 ± 0.02 TLS/mm^2, P = 0.05) (**Fig. 1c**). In untreated tumors, the majority of TLS were peritumoral, whereas in neoadjuvant treated tumors the majority were intratumoral (**Extended Data Fig 1b-c**). Taken together, these data suggest that neoadjuvant ICB induces the formation of intratumoral TLS.

High intratumoral TLS density after neoadjuvant ICB is associated with superior pathologic response and disease-free survival

We next set out to determine if there were an association between high TLS density after neoadjuvant ICB and three clinically meaningful endpoints: pathologic response to treatment, relapse free survival, and overall survival. Tumors treated with neoadjuvant ICB were reviewed

and assigned to a pathologic response category (non-response [NR], partial pathologic response [pPR], or major or complete pathologic response [MPR/CR]) according to percent residual viable tumor at the time of surgery.[17] 8/19 (42.1%) patients had a major or complete pathologic response, of which 2 had CR and 6 had MPR; 8/19 (42.1%) had a partial pathologic response (pPR); and 3/19 (15.8%) had non-response (NR). Intratumoral TLS density was significantly increased in tumors with MPR/CR compared to tumors with pPR (P = 0.000246), NR (P = 0.0129), or untreated patients (P = 0.000142) by Tukey's HSD test. In addition, total TLS density was also increased in tumors with MPR/CR compared to tumors with pPR (P = 0.00144), NR (P = 0.02), and untreated tumors (P = 0.00694) (**Fig. 1d** and **Extended Data Fig. 1d**). No significant difference was observed in peritumoral TLS density across pathologic response groups or untreated tumors (**Extended Data Fig. 1e**). Additional pathologic assessment was also performed according to the Immune Related Pathologic Response Criteria (irPRC), a set of categorical histopathologic criteria developed for standardized pathologic assessment of the regression bed of neoadjuvant immunotherapy treated solid tumors.[17] Using these criteria, we also observed a significant association between the presence of intratumoral TLS and MPR/CR (P = 0.02), while no significant association was detected between peritumoral TLS and MPR/CR (P = 0.38) (**Extended Data Table 3**). Thus, both a quantitative assessment of TLS density and a categorical evaluation of individual pathologic features suggested that intratumoral TLS density may be most correlated with response to treatment.

We next examined relapse free survival and overall survival in the treated cohort, excluding the untreated cohort from analysis lack of follow up data for the majority of the cohort. Significantly longer relapse free survival after surgery was observed in treated patients in the upper tertile of intratumoral TLS density compared to patients in the middle and lower tertiles *(P = 0.021) (***Fig. 1e***). At a median follow up of 38 months for patients in the upper tertile of intratumoral TLS density and 32 months for patients in the middle and lower tertiles, median RFS was not reached in the upper tertile and 9.1 months in the middle and lower group. RFS at 30 months was 100% and 38.5% (95% CI, 19.3% to 76.5%), respectively. No significant difference in overall survival (OS) was observed between the two groups (P = 0.24) (**Fig. 1f**), but at 30 months OS was 100% in the upper tertile and 76.9% (95% CI, 57.1% to 100%) in the middle and lower tertiles. In addition, we observed a trend toward improved RFS for patients in the upper tertile of total TLS density compared to the middle and lower tertiles (P = 0.13) (**Extended Data Fig. 2a**). OS was not significantly different (P = 0.28) (**Extended Data Fig. 2b**), but no deaths were observed in the upper tertile of total TLS density while three deaths were observed in the middle and lower tertiles. With respect to peritumoral TLS density, no difference was observed in relapse free survival (P = 0.56) or overall survival (p = 0.23) when comparing the upper tertile to the middle and lower tertiles (**Extended Data Fig. 2c-d**). Notably, in this cohort MPR/CR, which was closely associated with high TLS density, was also associated with superior RFS (P = 0.025) (**Extended Data Fig. 2e**). No significant difference was observed in OS (P = 0.16) (**Extended Data Fig. 2f**), but no deaths were observed in the MPR/CR group while three deaths were observed in the pPR/NR group. In addition, we also evaluated outcomes according to sex and previous viral HBV or HCV infection and identified no significant differences in RFS or OS (**Extended Data Fig. 2g-l**). Finally, to compare the different clinical covariates, we used the Bayesian information criterion[18] to quantify the strength of each parameter in predicting relapse free survival or death after neoadjuvant ICB and surgical resection. The strongest predictors of relapse free survival by BIC analysis were intratumoral TLS density and pathologic response (**Extended Data Table 4**).

High TLS density after neoadjuvant ICB is associated with increased T and B cell activation and an expanded intratumoral T and B cell repertoire

To identify differences in gene expression between tumors with high and low TLS density in this treatment context, we performed bulk RNA sequencing from FFPE surgical resection specimens. Tissue sections were collected from 14 tumors in the neoadjuvant treatment group, of which 2 samples were excluded after quality control. The resultant 12 samples were designated as TLS high (n = 5) or TLS low (n = 7) according to total TLS density relative to the mean total TLS density of the treatment group. Here, total TLS density was used rather than intratumoral or peritumoral TLS density since bulk sequencing of FFPE tissue blocks did not have spatial resolution to account for these differences. By principal component analysis, the 5 TLS high tumors and 1 TLS low tumor clustered separately from the remaining TLS low tumors (**Fig. 2a**). Differential expression analysis using the R package DESeq2 identified 814 differentially expressed genes (DEG), defined as having fold change in the TLS high group greater than 2 times that of the TLS low group and false discovery rate less than 0.05 (**Fig. 2b-c** and **Extended Data Table 5)**.

Compared to TLS low tumors, TLS high tumors demonstrated significant overexpression of multiple genes belonging to the Gene Ontology Biological Pathways gene sets for T and B cell activation, cytokine production, and antigen presentation, including *CTLA4*, *IL7R*, *IL6*, the B cell activating factor BAFF(*TNFSF13B*) and its receptors BAFF-R (*TNFRSF13C*) and TACI (*TNFRSF13B*), and the T cell-derived cytokine *IL17C*. TLS high tumors displayed significantly greater expression of *CCL19*, a chemokine involved in T-cell and B-cell migration to secondary lymphoid organs, and *CXCR5*, the receptor for the B-cell chemoattractant *CXCL13*. TLS high tumors also demonstrated increased expression of multiple B-cell related genes such as the B cell antigen CD79 (*CD79A* and *CD79B*), CD20 (*MS4A1*), and Fc Receptor Like A protein (*FCRLA*) which is highly expressed in germinal center B cells.[19] In addition, we found increased expression of immunoregulatory genes including *IL10*, *IL17REL*, and the integrin αvβ8-mediated *ITGB8*, which mediates TGF-beta-1 activation on the surface of regulatory T cells.[20,21] We also identified significantly increased expression of the gene encoding the germinal center regulatory protein EBI2 (*GPR183*), *DOCK10*, which regulates CD23 expression and sustains B-cell lymphopoiesis in secondary lymphoid tissue,[22] and *WDFY4*, a mediator of dendritic cell cross presentation.[23]

Gene set enrichment analysis for human gene sets in the MSigDB collections further identified significant enrichment in TLS high tumors of pathways associated with increased adaptive immune response, including Hallmark pathways for allograft rejection and inflammatory response, and multiple pathways related to T and B cell receptor activation (**Fig. 2d-e** and **Extended Data Table 6**). Consistent with these findings, TLS high tumors also displayed increased expression of the 12-chemokine gene signature which has previously been found in association with TLS formation in multiple solid tumor types (**Extended Data Fig. 3a**).[24] Taken together, these bulk gene expression data demonstrate that tumors with high TLS density display significantly higher levels of T and B cell activation compared to TLS low tumors.

To determine if TLS density was associated with differences in the adaptive immune repertoire, we used the Personalis ImmunoID NeXT platform to extract immunoglobulin heavy chain (IGH), TCRβ, and TCRα repertoire data from bulk RNA sequencing data. Statistical power was limited by the small sample size, but in tumors with high TLS density there were a significant increase in total number of immunoglobulin heavy chain (IGH) clones (P = 0.02), unique clonotypes (P = 0.029), and repertoire diversity (P = 0.043) by Wilcoxon rank sum test (**Extended Data Fig. 4a-c**). In addition, we identified a trend toward increased median number of total clones, unique clonotypes, repertoire diversity in the TCRα (P = 0.29, 0.18, and 0.18, respectively, by Wilcoxon rank sum test) (**Extended Data Fig. 4d-f**) and TCRβ repertoires (P = 0.22, 0.095, and 0.095, respectively, by Wilcoxon rank sum test) (**Extended Data Fig. 4g-i**). Overall, these findings suggest that high TLS density is associated with an expansion of the B and T cell repertoire in HCC treated with neoadjuvant immunotherapy.

In areas of tumor regression, an involuted TLS morphology is found that displays dissolution of the B cell germinal center, retention of the T cell zone, and increased expression of T cell memory markers

Based on these data, we hypothesized that a distinctive immunological process may occur in tumors with high intratumoral TLS density and major or complete pathologic response that contributes to long-term disease-free survival. To evaluate this hypothesis, we performed histologic examination of tumors with both viable tumor and extensive tumor regression beds. In viable tumor, the predominant phenotype observed was the canonical 'mature' stage of TLS characterized by a CD20+ B cell germinal center surrounded by CD4+ and CD8+ T cells.[25] TLS of this morphology showed characteristically high expression of the follicular dendritic cell marker CD21 and the proliferation marker Ki67. In contrast, in areas of tumor regression bed we observed an 'involuted' TLS morphology characterized by CD20+ B cells in a halo-like ring surrounding a central core of CD4+ and CD8+ T cells. CD21 and Ki67 expression were low to absent (**Fig. 3a**). To confirm that this involuted morphology was not an artifact of sectioning, we performed serial sectioning and anti-CD20 staining of FFPE tissue sections and confirmed the absence of a dense B cell core as is seen in mature TLS (**Extended Data Fig. 5a**). These involuted TLS were highly associated with tumors with complete pathologic response, and in several tumors were found in series (**Extended Data Fig 5b**), suggesting a shared lymphatic supply. No TLS of this morphology were detected in untreated tumors.

Given the location of the latter morphology within areas of nonviable tumor and the dispersed appearance of B cells in these lymphoid aggregates, we hypothesized that this morphology may represent TLS undergoing shutdown of the germinal center.[26,27] To characterize the features of this stage of TLS, we developed a 38-marker imaging mass cytometry antibody panel. Markers included in this panel were selected to identify different T cell subsets (CD3, CD4, CD8, FOXP3, CXCR3, CXCR5, ICOS), B cells subsets (CD20, BCL6, AID, CD138), follicular dendritic cells (CD21, CD23), dendritic cells (CD11c, DC-LAMP, CCR7), high endothelial venules (PNAd), macrophages (CD68), fibroblasts (Podoplanin [PDPN], αSMA), and tumor (CK). We included markers for T cell activation and exhaustion (CD25, CD69, CD137, PD-1, LAG3, TOX), co-stimulatory or antigen presenting molecules (CD86, HLA-DR), and markers of cell proliferation (Ki67) (**Extended Data Tables 7 and 8**). FFPE sections were obtained from the tumors of 9 patients treated with neoadjuvant ICB, in 8 of which the involuted morphology was identified, and, after whole-slide staining, 31 regions of interest (ROI) were captured by laser ablation (**Extended Data Fig. 6a**) from which 38 TLS areas (*n* = 20 mature and 18 involuted) were identified.

Consistent with previously published data, imaging mass cytometry of mature TLS (**Fig. 3b**) demonstrated dense B cell follicle-like structures surrounded by peripherally located CD4+ and CD8+ T cells with associated high endothelial venules (HEV) with a cuboidal morphology.[28,29] In multiple HEV, we observed CD8+ T cells in transit through these structures (**Fig. 3b, far left inset**). Mature TLS were also notable for an extensive CD21 and CD23 follicular dendritic cell network (**3b, middle left inset**), a distinct T cell zone with densely concentrated DCLAMP+ mature dendritic cells in close contact with T cells (**3b, middle right inset**), and a dense PDPN+ stromal network, similar to the fibroblastic reticular cell networks seen in secondary lymphoid organs[15,30] (**3b, far right inset**). Involuted TLS (**Fig. 3c**) showed no detectable HEV with a cuboidal morphology, scattered CD21 and CD23 expression, and diminished PDPN expression, consistent with attenuation of the TLS structure. Notably, the center of involuted TLS demonstrated apparent persistence of the T cell zone with co-location of DCLAMP+ mature dendritic cells and CD4+ and CD8+ T cells.

Further quantitative analysis of these structures supported these initial observations. After cell
segmentation, we identified 61,371 single cells which were assigned to 16 distinct cell clusters
(**Fig. 3d-f** and **Extended Data Fig. 6b-c**). In mature TLS we observed significantly higher density
of a BCL6^{high} population of B cells (B_BCL6^{high}), which was consistent with a germinal center B
cell population (P = 0.00023). In mature TLS, this cluster localized to the center of the B cell
germinal center in close proximity to CD21$^+$CD23$^+$ follicular dendritic cells and demonstrated high
expression of HLADR, a marker of antigen presentation, and CD86, a B cell activation marker,
consistent with an activate B cell population. A second B cell cluster was identified on the
periphery of the B cell follicle which displayed lower expression of BCL6 (B_BCL6^{low}) and
decreased expression of HLADR and CD86. This cluster was also found in significantly higher
density in mature TLS (P = 0.038). In contrast, in involuted TLS we observed significantly
increased density of a third B cell cluster (B_AID+) (P = 0.0026), which was characterized by high
expression of activation-induced cytidine deaminase (AID), the B cell enzyme which drives
somatic hypermutation and class switch recombination. AID is induced by BCR cross-linking and
has a half-life of 2.5 hours in the nucleus and 18-20 hours in its cytoplasmic form,[31,32] thus this
population may correspond to B cells undergoing somatic hypermutation or memory B cells which
had recently undergone immunoglobulin class switching, the latter of which we felt to be more
likely given the context. No difference was observed in plasma cell densities between the two
morphologies.

In the T cell compartment, we identified a single cytotoxic CD8$^+$ T cell population (Tc) and two
major CD4 T helper populations, a CD4$^+$CXCR5$^-$CXCR3$^+$ T peripheral helper (Tph) cluster, which
was located around the peripheral of the B cell germinal center in mature TLS and at the center
of involuted TLS, and a CD4$^+$CXCR3$^-$ T helper (Th_CXCR3^{low}) clusters. In location and marker
expression, Tph in these data were consistent with CD4$^+$ Tph that have been identified in patients
with autoimmune disease, where they are thought to play a T follicular helper (Tfh)-like role in
promoting pathogenic B cell responses in non-lymphoid tissue.[33–35] In contrast to changes
observed in the B cell compartment, no significant difference was in density of CD4$^+$ T
cell clusters or the cytotoxic Tc cluster.

Clustering analysis also identified a CD4$^+$CD57$^+$ cluster (Th_CD57$^+$) within the germinal center of
mature TLS, which may provide help to B cells and induce class switch recombination,[36–38] a
cluster of CD4$^+$FOXP3$^+$ regulatory T cells (Th_FOXP3$^+$), a cluster of cytotoxic T cells (Tc); CD4$^+$
and CD8$^+$ T cells defined by high expression of granzyme B (GZMB$^+$ T cell); proliferating T and B
cells defined by high expression of Ki67 (Proliferating T and B); a macrophage cluster with high
expression of CD68; a mature dendritic cell cluster defined by the presence of high expression of
DCLAMP and CCR7[39]; a high endothelial venule (HEV) cluster defined by expression of the
protein peripheral node addressin (PNAd); and a tumor cluster with high expression of Cytokeratin
(CK) and PDL1. In mature TLS compared to involuted TLS, there was significantly higher density
of proliferating T and B cells (P = 0.022), HEV (P = 0.00028), and tumor (P = 0.0026). On the
other hand, density of mature dendritic cells was increased in involuted TLS (P = 0.025).

To further evaluate the spatial relationships between different cell types in the mature and
involuted morphologies, we performed nearest neighbor analysis of the top 2 most frequent cell
neighbors for each cell cluster (**Fig. 3g**). As in previous the above analyses, neighborhood
analysis showed that the primary differences in spatial relationships between the two
morphologies occurred in B cell clusters. In particular, in the mature morphology BCL6^{high} germinal
center B cell cluster were first and second nearest neighbors for themselves, consistent with a
highly concentrated germinal center. In contrast, in the involuted TLS morphology the most
common first neighbor of this cluster was the BCL6^{low} cluster, consistent with a more dispersed
germinal center in the involuted TLS. On the other hand, Tph were the most common non-self

neighbors for GZMB T cells, mature dendritic cells, proliferating T cells, and FOXP3[+] Tregs in both mature and involuted TLS, suggesting that the spatial relationships of these clusters was preserved across the two morphologies despite changes occurring in the B cell germinal center. Network analysis, which we used to visualize the average distances between cell clusters and cell cluster abundance across the two TLS morphologies, demonstrated similar changes to the two B cell clusters (B_BCL6[high] and B_BCL6[low]) occupying the germinal center and preservation of spatial relationships between mature dendritic cells, Tph, FOXP3[+] T cells, proliferating T and B cells, and GZMB[+] T cells is mature and involuted TLS (**Fig. 3h**). Overall, these neighborhood and network analyses suggested that while the B cell germinal center appeared to undergo dissolution in involuted TLS, the T cell zone was preserved.

Finally, evaluation of individual marker expression by cluster supported these observations regarding persistence of the T cell zone (**Fig. 3i**). In the mature dendritic cell cluster, expression of CCR7, HLADR, and CD86 were significantly increased, implying ongoing antigen presentation in these structures, and both the Tph and Tc clusters demonstrated increased expression of markers of antigen experience, including CD45RO, CD25, PD1, and TOX expression While the precise role of TOX in T peripheral helper cells is not established, TOX2 has previously been shown to be involved in the establishment of durable GC Tfh memory.[40] Taken together, these data suggest that the involuted morphology may be a site of persistent antigen presentation by mature dendritic cells, which drive the formation of antigen-experienced memory T cell populations.

Expanded T cell clonotypes are shared across TLS within a tumor, while B cell repertoires of individual TLS are highly distinct

Based on these data, we next sought to determine whether there were differences in the T and B cell repertoires of TLS of these two morphologies. We microdissected 38 individual TLS (32 mature and 6 involuted) from 7 treated tumors and performed bulk sequencing using the Adaptive ImmunoSEQ TCRβ and IGH assays (**Fig. 4a, Extended Data Fig. 7a, Extended Data Table 9**). After filtering to remove repertoires with low counts, 35 TCRβ repertoires and 32 IGH repertoires were analyzed. Across all samples, the repertoire size was variable with a mean total TCRβ clonotypes of 7171±8472 (**Extended Data Fig. 7b**). Singleton clonotypes comprised 68.7±13.4% of the TCRβ repertoire in all TLS sampled. In mature TLS, singleton clonotypes comprised 72.02±9.83% of the T cell repertoire, while in involuted TLS the singleton compartment constituted 48.98±16.1%. Across TLS microdissected from the same tumors, a mean of 32.3±12.3% of unique TCRβ clonotypes could be identified in other TLS from the same tumor. TCRβ clonotypes identified in all TLS from the same tumor were highly expanded, while those found in only one TLS were predominantly singletons (**Fig. 4b-c and Extended Data Fig. 7c-h**), suggesting a high degree of T cell trafficking as well as significant local T cell repertoire diversity at each individual TLS.

Across all microdissected TLS, the mean total number of IGH clonotypes was 922±1188 (**Extended Data Fig. 8a**). Singleton clonotypes comprised 95.3±3.9% of IGH repertoire of all TLS sampled. In mature TLS, singleton clonotypes comprised 95.7%±4.0 of the IGH repertoire, while in involuted TLS the singleton compartment constituted 93.4±3.1%. IGH repertoire sharing was significant lower across TLS microdissected from the same tumor (P = 7.6e-15), with only 6.7±5.6% of unique IGH clonotypes of each TLS detected in other TLS from the same tumor (**Fig. 4d-f** and **Extended Figure 7b-g**). These B cell repertoire characteristics are consistent with highly distinct, independent germinal center reactions.

In three patients (P12, OT1, and OT6) in which mature and involuted TLS were present in the same tissue block, we compared the immune repertoires of these two morphologies. TCRβ clonality was significantly increased in mature TLS compared to involuted TLS (P = 0.023) (**Fig. 4g**), although this difference was primarily observed in a single patient OT6 (**Extended Data Fig. 7i**). No difference was observed in IGH clonality (**Extended Data Fig. 7g**), but the IGH repertoire of involuted TLS did demonstrate a significantly higher number of V gene substitutions, a surrogate for somatic hypermutation (P < 2.22e-16) (**Fig. 4h** and **Extended Fig. 8i**). Taken together, these comparisons suggest that B cell populations in involuted TLS have undergone greater antigen-driven positive selection, consistent with a late-stage germinal center, and that there is associated T cell repertoire contraction and clonal expansion at these structures.

Given the extensive sharing of expanded T cell clonotypes observed within TLS from the same tumor, we also evaluated the peripheral blood to determine the extent of T cell trafficking between TLS and peripheral blood. We performed TCRβ sequencing of pre- and post-treatment peripheral blood mononuclear cells from 5 of the 7 patients whose TLS were microdissected. In TLS from these 5 patients, a mean of 44.0±8.4% of unique TCRβ clonotypes and 52.7±8.5% of total clonotypes in TLS were also identified in post-treatment peripheral blood. Similar overlap was observed between TLS repertoires and pre-treatment peripheral TCRβ repertoires, where a mean of 40.4±11.5% of unique TCRβ clonotypes and 48.7±11.0% total clonotypes in TLS were identified in pre-treatment peripheral blood (**Extended Data Fig. 9a-b**). In 3 of the 5 patients, 13 unique TCRβ were significantly expanded after neoadjuvant treatment and 9/13 (69.2%) were detected in at least one TLS (**Extended Data Table 10**). Together, these data provide evidence for a high degree of overlap between T cells within TLS and T cells in the peripheral blood.

The top expanded T cell clonotypes in mature TLS are cytotoxic granzyme K and granzyme B-expressing CD8$^+$ T cells

To further characterize T and B cell populations identified in TLS, we performed single cell RNA/TCR/BCR sequencing of post-treatment peripheral blood from the 7 patients from whose tumors TLS were microdissected. Sequencing of tumor infiltrating lymphocytes (TIL) was also performed for all 7 patients, but in only one sample (patient OT6) was sequencing data of sufficient quality for further analysis. Peripheral blood and TIL samples were processed by fluorescence-activated cell sorting (FACS) after labeling with antibodies to CD3 and CD19. After pre-processing and filtering to remove low quality sequencing data, 28,694 single cells were identified in the peripheral blood and 620 in the TIL. After performing preliminary cluster annotation using a reference annotated dataset, we attempted to match the TCRβ and IGH CDR3 amino acid sequences identified in microdissected TLS with sequences identified in the single cell dataset. TCRβ in the microdissection and single cell datasets were successfully matched (described below), but no matching IGH were identified between the bulk sequencing performed on microdissected TLS and single cell sequencing data, therefore B cells were excluded from subsequent analysis.

The resultant 23,172 T cells in the post-treatment peripheral blood samples and 562 T cells in the TIL of patient OT6 were clustered into 16 distinct cell clusters based on expression of canonical genes associated with specific T cell subsets. The CD4 compartment of the single cell dataset was divided into a naïve CD4$^+$ T cell cluster (CD4 Naïve) expressing high levels of *CCR7* and *LEF1*; a CD4 Naïve-like cluster (CD4 Naïve-like) characterized by expression of *CCR7 and TCF7*; a CD4 T central memory cluster (CD4 TCM) with high expression of *LTB and S100A4*; a CD4 T peripheral helper cluster (Tph) characterized by low expression of *CXCR5,* high expression of *CXCR3,* and high expression of ICOS; and two CD4 T effector memory clusters notable for high expression of granzyme K (CD4 TEM_GZMK) and granzyme B (CD4 TEM_GZMB). CD8 T cells

were divided into the following clusters: a naïve cluster (CD8 Naïve) highly expressing *CD8B*, *CCR7*, *LEF1*; a CD8 T central memory cluster (CD8 TCM) with elevated expression of *CD8B and LINC02446*; two CD8 T effector memory clusters distinguished by high expression of granzyme K (CD8 TEM_GZMK) and high expression of granzyme B (CD8 TEM_GZMB); and a CD8 tissue resident memory-like cluster (CD8 TRM) with increased expression of *NR4A2*, *DUSP2*, and *ZNF683.* In addition, we identified a CD4 regulatory T cell cluster (Treg) with high expression of *FOXP3* and *RTKN2*; an NK-T cell cluster (NK-T) highly expressing *PRF1* and *GZMB*; a double negative T cell cluster (dnT) with high expression of *SYNE* and *MALAT1*; a gamma delta T cells (gdT) with high expression of *TRDV2* and *TRGV9*; and mucosal invariant T cells cluster (MAIT) highly expressing *KLRB1* and *SLC4A10* (**Fig. 5a-c** and **Extended Data Fig. 10a-b**). In the TIL of patient OT6, 11 of 16 clusters were identified: CD4 Naïve-like, CD4 TCM, CD4 Tph, CD4 TEM_GZMK, CD8 TCM, CD8 TEM_GZMK, CD8 TEM_GZMB, CD8 TRM, Treg, dnT, and MAIT **(Extended Data Fig. 11a-c**).

19,546/23,172 (84.3%) single cells in the peripheral blood dataset and 346/562 (61.6%) cells in the TIL had a partial or completely sequenced TCRαβ chain identified by single cell TCR sequencing, of which there were 15,016 and 256 unique TCRs, respectively (**Extended Data Table 11**). In the peripheral blood single cell dataset, clonal expansion was most strongly associated with the CD4 TEM_GZMB (Odds Ratio 31.48, P < 0.001) and CD8 TEM_GZMB clusters (Odds Ratio 17.99, P < 0.001) by Fisher's Exact test (**Fig. 5d** and **Extended Data Table 12**). In TIL, clonal expansion was most strongly associated with the CD8 TEM_GZMB (Odds Ratio 3.49, P = 0.003), CD8 TRM (Odds Ratio 2.57, P = 0.041), and CD8 TEM_GZMK (Odds Ratio 1.94, P = 0.013) clusters (**Extended Data Fig. 11d and Extended Data Table 12).**

T cells belonging to the CD8 TEM_GZMK cluster were notable for increased expression of the gene associated with cytotoxicity, including Granzyme K (*GZMK*) and the chemokine ligand CCL5 (*CCL5*) and decreased expression of Granulysin (*GNLY*). The CD8 TEM_GZMB clusters also demonstrated hallmarks of cytotoxicity, including elevated expression of granzyme B (*GZMB*) and granzyme H (*GZMH*), as well as elevated expression of perforin (*PRF1*), and *GNLY* (**Extended Data Fig. 10c-d**).[41] *GZMK* expression was low in the latter cluster. These two transcriptional phenotypes were consistent with T progenitor exhausted and cytotoxic/terminally differentiated CD8 T cell states, respectively, which have been previously identified in the peripheral blood and tumors of patients treated with ICB.[41] Consistent with this identity, in the TIL from patient OT6, both *GZMK* and *GZMB* expressing CD8 clusters showed increased expression of multiple T cell exhaustion markers, including *PDCD1*, *CTLA4*, *LAG3*, *TIGIT*, and *TOX*, which were more highly expressed in CD8 TEM_GZMB compared to CD8 TEM_GZMK. *NKG7* and *CCL5*, which are associated with cytotoxic CD8 T cells, were also increased in both clusters, with higher expression in the GZMB high cluster (**Extended Data Fig 11h-i**). Notably, both clusters in the single cell TIL from patient OT6 demonstrated elevated expression of the B cell chemoattractant *CXCL13*, which has been associated with tumor-specific T cells in single cell sequencing studies of TIL from patients treated with ICB and is involved in the formation of TLS.[42,43]

No CD4+ T cell cluster was observed in the single cell data that displayed a transcriptional phenotype consistent with a CD4+ T follicular helper (Tfh) population, which are defined by high expression of CXCR5, CXCR3, and ICOS. Instead, we identified a CD4+ T cell cluster consistent with the T peripheral helper cluster identified by imaging mass cytometry. In the peripheral blood, cells belonging to this cluster demonstrated low CXCR5 expression, increased CXCR3 expression, and elevated expression of CTLA4, TIGIT, and TOX (**Extended Data Fig. 10e**). Cells belonging to this cluster in the single cell TIL demonstrated increased expression of *CXCL13*, *ICOS*, *PD1*, *MAF*, *TOX* and high expression of multiple exhaustion markers including *CTLA4*, *LAG3*, *TIGIT*, *HAVCR2*, and *TNFRSF18 (GITR)* (**Extended Data Fig 11j**). Based on these data

518 and the imaging mass cytometry above, we conclude that Tph constitute a major CD4$^+$ T cell
519 subset found in TLS in the context of neoadjuvant immunotherapy.
520
Approximately one-third, or 6349/19546 (32.5%), of single cells with a TCRβ in the peripheral T
cell dataset and 199/346 (77.7%) single cells with a TCRβ in the TIL were identified in at least
one TLS (**Extended Data Table 11**). TCRβ identified in TLS were detected in all clusters of the
peripheral and TIL single cell dataset. Among peripheral blood T cells, the clusters most strongly
associated with TLS were CD4 TEM_GZMB (Odds Ratio 10.72, P < 0.001), CD8 TEM_GZMB
(Odds Ratio 9.73, P < 0.001), and CD8 TEM_GZMK (Odds Ratio 9.46, P < 0.001) (**Fig. 5e-f** and
Extended Data Table 13), suggesting that TLS specifically promote the trafficking of effector
memory CD4$^+$ and CD8$^+$ T cell populations from the peripheral blood to tumor. In the TIL, no
cluster was significantly correlated with TLS but the Treg cluster was inversely correlated with
presence in TLS (Odds Ratio 0.22, P = 0.003) (**Extended Data Fig. 11e-f** and **Extended Data
Table 13**). The proportion of all TCRβ identified in TLS that were matched to single cell data was
low overall (2,908/135,909 unique clonotypes or 2.1%), but a higher proportion of clonotypes were
successfully matched for the most expanded clones, with 369/1359 (27.2%) of the top 1% of
TCRβ that had been identified in TLS being successfully matched to the single cell data, and
63/137 (46%) of the top 0.1% of TCRβ (**Extended Data Table 14**). Thus, this approach, while
providing a limited view of singleton TCRβ identified in TLS, could be used to provide additional
transcriptional information about expanded T cell populations trafficking to TLS.
538
We next used these data to infer the transcriptional phenotype of T cells trafficking through TLS.
Since our single cell dataset contained peripheral blood and TIL data, we first evaluated the
correlation between peripheral blood and TIL phenotypes, and the reliability of using data from
single cell sequencing from one compartment to infer the properties of single cells with the same
TCRβ in the other compartment. To carry this out, we examined the correlation between single
cell cluster identity for TCRβ which were found in both peripheral blood and TIL in patient OT6. In
total, 16 unique TCRβ sequences were present in both peripheral blood and TIL (**Extended Data
Fig. 11a**). 7/16 TCRβ clonotypes had the same single cell identity for all cells with the same TCRβ
in the peripheral blood and TIL, and an additional 5/16 TCRβ had the same cluster identity
assigned to at least half of the cells in both peripheral blood and TIL. In only 4/16 TCRβ were the
cluster identities of cells with the same TCRβ entirely discordant (**Extended Data Fig**. **11b).**
Based on these data, we concluded that the peripheral blood transcriptional phenotype closely
recapitulates the cluster assignment of TIL, and thus both identities may be used to determine a
putative phenotype for T cells identified in TLS by TCRβ sequencing. These findings were
consistent with previous work demonstrating that in circulating TILs, gene signatures of effector
functions, but not terminal exhaustion, reflect those observed in the tumor.[44]
555
Across all 7 patients, the majority of T cells identified by matching of the TCRβ were GZMK and
GZMB expressing CD8 T effector memory cells, but we also observed CD4 TEM_GZMK, CD4
CTL, and CD4 T peripheral helper clusters among the putative phenotypes of T cells trafficking
through TLS (**Fig. 5g**). Notably, in the involuted TLS from the tumor of patient OT6, where we had
previously noted significant increase in clonality relative to mature TLS, clonal expansion was
greatest in the of CD8 TEM_GZMK, CD8 TEM_GZMB, and CD8 TRM clusters (**Fig. 5h** and
Extended Data Fig. 11k). Overall, these data provide single cell resolution to the top expanded
clonotypes in TLS and show that highly expanded T cell populations in TLS are CD8$^+$ T cell
effector memory, which may undergo clonal expansion and repertoire contraction in concert with
expansion of resident memory populations in areas of tumor regression.
566
567 **DISCUSSION**
568

Neoadjuvant immunotherapy aims to use the primary tumor as a source of antigens to enhance antitumor immunity and prevent cancer recurrence after surgery.[4] Preclinical and clinical data suggest that this approach induces more durable immunologic memory than adjuvant immunotherapy alone,[3] but the mechanism by which this occurs and the contribution of TLS to this process are not well understood. The data presented here show that in HCC treated with neoadjuvant immunotherapy, intratumoral TLS are associated with superior pathologic responses and improved relapse free survival. These findings are consistent with data reported in other solid tumors treated with ICB,[9–12] as well as studies of the prognostic significance of intratumoral TLS in treatment-naïve early stage HCC treated with surgical resection.[16] In tumors with high TLS density and significant regression of the tumor, we further identified an involuted morphology of TLS in areas of nonviable tumor whose location, histologic, and immunologic features, and similarity to late stage germinal centers observed in murine secondary lymphoid organs,[26] are consistent with a terminal stage of the TLS life cycle. Using laser capture microdissection, bulk immune repertoire sequencing, and matched single cell sequencing, we identify and characterize expanded T cell populations trafficking through TLS and find evidence for significant immune repertoire changes associated with TLS dissolution.

While TLS are thought to mature from a loosely organized lymphoid aggregate to a CD21[+] primary follicle and reach full maturity as a CD21[+]CD23[+] secondary follicle,[16,45,46] which have distinct T and B cell zones,[25] the circumstances of TLS resolution are not known.[47] These data suggest that TLS dissolution may be driven by elimination of tumor and may occur dyssynchronously, with dissolution of the B cell germinal center accompanied by persistence of a T cell zone enriched for interactions between DCLAMP[+]CCR7[+]HLADR[+] mature dendritic cells and CD4[+] and CD8[+] T cells. Furthermore, the changes observed in T cell repertoire at these structures, including increase in clonality and expansion of cytotoxic and tissue resident memory-like CD8[+] T cell clonotypes, suggest that late-stage TLS may play a functional role in supporting the contraction and memory phase of the intratumoral adaptive immune response through persistent antigen presentation in the T cell zone (**Fig. 6**). Such a role would also be consistent with recent data suggesting that tonic antigenic stimulation drives programs of T cell residency in tumors, and would identify a specific place where such interactions may occur.[48]

Notably, in neoadjuvant treated tumors we did not detect a CXCR5[+]CXCR3[+]CD4[+] T follicular helper population, which have been reported in tumor associated TLS.[49,50] Rather, in both the imaging mass cytometry analysis and single cell datasets we identified a CXCR5[-]CXCR3[+] CD4[+] T cell population, which expressed CXCL13 in the single cell TIL of patient OT6 and was also detectable in post-treatment peripheral blood single cell sequencing. CXCR5[-]CXCL13-producing CD4[+] T cells have been identified in untreated human breast cancer under the name TFHX13[51,52] and in autoimmune disease, where they are termed CD4[+] T peripheral helper cells and have been shown to provide help to B cells in an IL-21 dependent manner.[34,35,53] This population was present in both the mature and involuted morphologies observed in these patients. Additional studies are required to determine whether this population of CD4[+] T cells provides help to B cells in this treatment setting, and to determine their role in involuted TLS, where they were found in association with mature dendritic cells.

We recognize several limitations of the findings reported here. First, samples were obtained from a single institution and may not represent the full diversity of HCC etiologies and subtypes. Additionally, while the untreated cohort in this study was similar to the treatment cohort in age, sex, and HCC etiology and received treatment at the same primary institution, we cannot exclude the possibility that the different pathologic findings between the untreated and neoadjuvant cohorts arose as a consequence of differences in the patient populations rather than treatment status. We infer the transcriptional phenotype of T cells infiltrating TLS based on matching TCRs

identified by microdissection with peripheral T cells and/or TIL subjected to single cell sequencing.
In our data, we demonstrate a correlation between the cluster identity for 16 TCRs shared
between the peripheral blood and TIL of patient OT6, and others have reported correlation
between gene signatures of effector functions in circulating TILs and tumor;[44] however, it is
possible that the phenotype of these T cells are not fully conserved outside of TLS. Finally, our
analyses throughout are limited by small sample size. However, clinical samples from
neoadjuvant studies are rare, and as we have shown here even small samples can provide
important insights into the constituent immune populations in tumors arising in human subjects.
628
Finally, these data raise several important questions which future studies should address. First,
the role of FOXP3[+] T follicular regulatory cells, which have been shown to regulate germinal
center reactions,[54,55] is not completely understood in this context and this population could not be
resolved in our imaging mass cytometry analysis. Second, while dispersion of the B cell follicle
was associated with attenuation of the PDPN[+] fibroblastic reticular cell and the CD21[+]CD23[+]
follicular dendritic cell network in our data, other changes occurring in the fibroblast populations
involved in this process remain unclear and it is not known what permits retention of mature
dendritic cells within the T cell zone. Third, since all tumors in which involuted TLS were observed
were treated with neoadjuvant ICB and these structures were not seen in untreated tumors, the
contribution of therapy to their formation cannot be established from our data. In murine studies,
both the PD-1 and CTLA-4 pathways have been shown to be involved in regulation of T follicular
regulatory populations,[56–58] suggesting that therapeutic blockade may affect the dynamics of
germinal center formation and dissolution. Nonetheless, these data shed light on the
circumstances of TLS resolution and suggest that this terminal stage, about which nothing was
previously known, may play a functional role in the formation of intratumoral T cell memory after
elimination of viable tumor.
645
646 **METHODS**
647
Study design
The aim of this study was to characterize tertiary lymphoid structures (TLS) in patients with
hepatocellular carcinoma (HCC) treated with neoadjuvant ICB-based therapy prior to surgical
resection of the primary tumors. To understand the clinical significance of TLS, we analyzed TLS
density in treated patients and untreated controls and correlated TLS density in treated patients
against pathologic response and post-surgical clinical outcomes. We performed bulk RNA
sequencing of tumors with high and low TLS density after neoadjuvant treatment to understand
the gene expression programs of tumors with high TLS density. We then characterized the
morphological and functional properties using bulk immune repertoire sequencing of TCRβ and
IGH, imaging mass cytometry, matched single cell TCR and RNA sequencing of peripheral blood
and tumor infiltrating lymphocytes.
659
Patient identification and data collection
Patients were identified for inclusion in this study if they received surgical resection for locally
advanced, non-metastatic hepatocellular carcinoma after neoadjuvant ICB-based therapy
between October 1, 2019 and January 31, 2022 at the Johns Hopkins Sidney Kimmel Cancer
Center. Retrospective chart review was performed to collect clinical data from the electronic
medical record regarding age at surgery, sex, date of resection, HCC etiology, histologic grade of
tumor, neoadjuvant treatment, relapse free survival, and overall survival. A cohort of untreated
control patients who had undergone surgical resection for HCC without prior systemic treatment
were also identified via a search of the electronic medical record. Review of the electronic medical
record was performed to confirm absence of prior systemic treatment. For both cohorts, histologic
grade was based on pathologic assessment at the time of resection if there was discordance with

grade reported for pre-treatment biopsy. Patients in both cohorts were excluded from analysis if
there was evidence of active hepatitis B (defined by a positive HBsAg or detectable HBV DNA)
prior to surgery. Patients were excluded from the control group if the etiology of their HCC was
not represented in the treatment group (e.g. hepatic adenoma and hereditary hemochromatosis).
This study was conducted in accordance with the Declaration of Helsinki and was approved by
the Johns Hopkins University Institutional Review Board (IRB00149350, IRB00138853,
NA_00085595). Informed consent or waiver of consent was obtained from all patients. Treated
patients identified with the letter P were accrued as participants in the phase I clinical trial
NCT03299946.[8]
680

Histopathologic assessment of TLS density and pathologic response
Evaluation of pathologic response was performed by a hepatopathologist (RA). Pathologic
response designations were assigned according to percent residual viable tumor in surgically
resected tumors. Complete response (CR) was defined as 0% residual viable tumor, major
pathologic response (MPR) as less than 10% residual viable tumor, partial pathologic response
(pPR) as 10-90% residual viable tumor, and non-response (NR) as >90% residual viable tumor.[17]
12 of the 19 patients had previously undergone assignment of pathologic response according
binary categorization of major or complete pathologic response versus non-response,[8] and for
this group non-responders were categorized as NR or pPR as described above. To determine
TLS density, formalin fixed paraffin embedded (FFPE) tumors were sectioned, mounted on glass
slides, and stained with anti-CD20 antibody as described below. Whole slide images were
obtained at 0.49 µm per pixel using the Hamamatsu NanoZoomer. The presence of CD20
positivity was determined by digital image analysis software (HALO v3.0.311 Indica Labs), with
TLS defined as CD20 positive cell aggregates greater than 150µm in diameter located among
tumor cells or at the invasive margin in areas of viable and nonviable tumor. TLS density was
determined by calculating the number of TLS per mm^2 of viable and nonviable tumor. TLS were
classified as peritumoral if they were found within 200 µm of the interface between normal
adjacent parenchyma and tumor and intratumoral if they were found within the tumor or tumor
regression bed.
700

Survival analyses
The Kaplan–Meier method was used to estimate relapse-free survival (PFS) and overall survival
(OS). Relapse free survival (RFS) was defined as the time from surgical resection to radiographic
relapse. Overall survival (OS) was defined as the time from surgical resection to death from any
cause. If a patient was not known to have had either event, RFS and OS were censored at the
last date of known healthcare contact. RFS and OS analyses were limited to patients treated with
neoadjuvant therapy and were not performed in the untreated controls due to limited follow up in
this cohort. Survival analyses using the Kaplan-Meier method were performed using the R
package survminer. Bayesian information criterion (BIC) analysis was performed using the R
package stats. A linear regression model was used to evaluate the effect of each marker,
dichotomized by the mean, as a predictor of each distance measure. For each binary outcome,
logistic regression was employed, with each marker treated as continuous. A meaningful
difference in BIC between the two models is 2 at a minimum, and a difference between 5-10 and
above 10 is considered to be strong and very strong, respectively.[59]
715

Immunohistochemistry
Automated single and dual staining was performed on the Leica Bond RX (Leica Biosystems).
Single staining for CD20 was employed for determination of TLS density. Dual staining for CD3
and CD21, CD8 and CD4, Ki67 and CD20 was performed prior to laser capture microdissection
of TLS. Slides were baked and dewaxed online followed by antigen retrieval for 20 min at 100°C.
Endogenous peroxidase was blocked using Peroxidase block (Refine Kit) followed by Protein

block (X090930-2, Agilent Technologies Inc., Santa Clara, CA). Primary antibodies were applied
at room temperature. Detection was performed using the Bond Polymer Refine Kit (DS9800, Leica
Biosystems). For dual staining, a second round of antigen retrieval was performed for 20 min at
95°C followed by application of a second primary antibody. Detection of the second primary
antibody was performed using the Bond Polymer Red Refine Kit (DS9390, Leica Biosystems).
Slides were counterstained, baked and coverslipped using Ecomount (5082832, Biocare Medical,
Walnut Creek, CA). Antigen retrieval buffers and concentrations of all antibodies are listed in
Extended Data Table 15. Antibodies were diluted to appropriate working concentration using
Antibody Diluent (S302283-2, Agilent Technologies Inc).
731
Bulk RNA sequencing and TCRβ/BCR immune repertoire profiling of FFPE tumor
RNA was extracted from FFPE tumor from the treatment cohort and sequenced using the
commercial platform ImmunoID NeXT with 200 million paired end reads (150 base pair). Reads
were aligned in accordance with the Personalis Cancer RNA pipeline and transcript per million
(TPM) values were extracted.[60] Bulk RNA sequencing was performed on 14 tumors in two
batches. No batch correction was applied due to lack of clear batch-to-batch differences by
principal component analysis. 2 samples were excluded due to poor sequencing depth, as defined
by a median of the log2 transformed count data being equal to 0 for those samples. The remaining
12 samples were filtered to include only genes for which the sum of raw counts across all samples
was greater than 1. Variance stabilizing transformation was performed on the resultant data and
differentially expressed genes were identified using DESeq2.[61] Genes with an adjusted P value
of < 0.05, and a minimum log2 fold change of 1 were considered differentially expressed. Pathway
analysis was performed using the R package fsgsea to identify biologically enriched pathways
from the MSigDB hallmark gene sets.[62,63] For pathway analyses, adjusted P values of < 0.05 were
considered statistically significant. TCRβ and BCR repertoire profiling was performed using the
ImmunoID NeXT transcriptome, which provides augmented (approximately a 100x increase over
a standard transcriptome) coverage of TCRβ and BCR.[60,64] Clones were identified using MiXCR[65]
and repertoire analysis was performed using the R package immunarch.[66] Clonality was
calculated as 1-Shannon's equitability[67] with clonality values ranging from 0-1, with 0 indicating
equal representation of all clones within a repertoire and 1 being a repertoire consisting of only
one clone.
753
IMC staining and acquisition
IMC Staining was done as previously described.[8,68] Briefly, formalin-fixed paraffin-embedded
(FFPE) resected liver tissue sections were baked, deparaffinized in xylene, then rehydrated in an
alcohol gradient. Slides were incubated in Antigen Retrieval Agent pH 9 (Agilent PN S2367) at
96 °C for 1 hour then blocked with 3% BSA in PBS for 45 min at room temperature followed by
overnight staining at 4°C with the antibody cocktail. Antibodies, metal isotopes, and their titrations
are listed in **Extended Data Table 8**. Images were acquired using a Hyperion Imaging System
(Standard BioTools) at the Johns Hopkins Mass Cytometry Facility. Upon image acquisition,
representative images were visualized and generated through MCD™ Viewer (Standard
BioTools).
764
IMC data analysis
Images were segmented into a single-cell dataset using the publicly available software pipeline
based on CellProfiler, ilastik, and HistoCAT.[69–72] Since multiple images contained more than one
TLS, images were subset for distinct TLS regions by manual gating using FlowJo™v10.9.0
Software (BD Life Sciences), which identified the xy coordinates of cells belonging to distinct
lymphoid aggregates (**Extended Data Fig. 6b-c**). This resulted in 38 unique TLS matching either
the mature (*n* = 20) or involuted morphology (*n* = 18). The resulting 61,371 single cells were
clustered using FlowSOM[73] into metaclusters, which were manually annotated into final cell types.

Density of each cell type was determined by calculating the number of cells per unit area as
determined by ImageJ v1.53.[74] For network visualization, the mean distance between each cell
type was computed and visualized using the R package qgraph.[75] Neighborhood analysis was
performed by using data generated by HistoCAT summarizing the top neighboring cell types for
every cell type.
778
Laser capture microdissection and TCRβ/IGH sequencing of microdissected TLS
10-14 μm serial tissue sections were obtained from formalin-fixed paraffin embedded (FFPE)
tumor tissue blocks and mounted on UV activated PEN membrane glass slides (Applied
Biosystems Cat. No. LCM0522) with additional 4-μm tissue sections cut every 150 μm for staining
with hematoxylin and eosin (H&E) and dual IHC for CD3/CD21, CD8/CD4, and Ki67/CD20 as
described above. Stained sections were scanned at 20x objective equivalent (0.49 μm pixel^{-1}) on
a digital slidescanner (Hamamatsu Nanozoomer) in advance of microdissection and annotated
using NDP.view2 viewing software in order to identify areas for microdissection. On the day of
microdissection, unstained tissue sections mounted on PEN membrane slides were
deparaffinized using xylene and graded alcohol washes and stained with H&E. Laser capture
microdissection of individual TLS was performed on the LMD 7000 system (Leica) and genomic
DNA was extracted using the Qiagen QIAamp DNA FFPE Tissue Kit following the manufacturer's
protocol (Qiagen). DNA concentrations were quantified with a Qubit 4 Fluorometer using the Qubit
dsDNA high sensitivity assay (Invitrogen). Sequencing of the TCRβ and IGH CDR3 regions was
performed using the immunoSEQ platform (Adaptive Biotechnologies).[76,77] TCRβ and IGH
repertoire data were downloaded from the Adaptive ImmunoSEQ analyzer web interface after
filtering to remove non-productive reads. After exclusion of repertoires with fewer than 500 TCRβ
clones and 50 IGH clones, subsequent analysis was performed using the R package immunarch[66]
and the Python package Change-O,[78] which was used to assign clonal families to IGH data. J
gene, and greater than 90% identical CDR3 sequence according to nucleotide hamming distance.
Clonality was calculated as described above using 1-Shannon's equitability. To compare clonality
across multiple TLS from the same tumor, we used the median clonality of 1000 iterations of
downsampling to the number of productive CDR3 sequences in the smallest TCRβ or IGH
repertoire for that patient.[79]
803
Peripheral blood and fresh tumor collection and processing
Processing of peripheral and cryopreservation was completed as previously described.[8] Fresh
tumor tissue was diced with a sterile scalpel and dissociated in 0.1% collagenase in RPMI 1640
for 60 minutes at 37ºC using the gentleMACS OctoDissociator (Miltenyi Biotec) according to the
manufacturer's instructions. Supernatant was collected and centrifuged at 1500 rpm for 10
minutes. Supernatant was removed and discarded, and the cell pellet was resuspended in ACK
Lysing buffer (Quality Biological, cat# 118-156-721) and incubated at room temperature for 5
minutes before centrifugation at 1500 rpm for 10 minutes. Cells were resuspended in PBS,
counted using a manual hematocytometer, and cryopreserved in 10% DMSO/AIM-V freezing
media.
814
Single cell RNA/TCR/BCR-sequencing
For all 7 patients from whose tumors TLS were microdissected, single cell sequencing was
obtained for peripheral blood T and B cells isolated by Fluorescent Activated Cell Sorting (FACS).
For 6 of 7 patients, the peripheral blood sample was obtained following completion of neoadjuvant
ICB and prior to surgical resection; for 1 of the patients, the peripheral blood sample was drawn
4 weeks after resection. In addition, in the latter patient, single cell sequencing was performed for
tumor infiltrating T and B cells isolated by FACS from tumor specimen. Cryopreserved PBMC and
tumor suspension were thawed and washed with pre-warmed RPMI with 10% FBS. Cells were
resuspended 0.04% BSA in PBS and stained with a viability marker (Zombie NIR, BioLegend)

and Fc block (Biolegend, Cat. no. 422302) for 10 minutes at room temperature in the dark. Cells
were then stained with antibodies against CD3 (FITC, clone HIT3a), for 20 minutes on ice and
CD19 (PE/dazzle, clone SJ25C1) (**Extended Data Table 16**). After staining, viable CD3$^+$ and
CD19$^+$ cells were sorted into 0.04% BSA in PBS using a BD FACS Aria II Cell Sorter at a 4:1
ratio. Sorted cells were counted and resuspended at a concentration of 1000 cells per µl. The
single-cell library preparations for gene expression and V(D)J were performed with the Chromium
Next GEM Single Cell 5' GEM Kit v2 (10x Genomics) and Chromium Single Cell V(D)J
Amplification Kit (human TCR) (10x Genomics), respectively. The cells were partitioned into
nanoliter-scale Gel Beads in-emulsion (GEMs) and cells were barcoded. The cDNA synthesis
and amplification was performed prior to sample split for the gene expression and for V(D)J
libraries. Single cell libraries were sequenced on an Illumina NovaSeq instrument using 2 × 150-
bp paired end sequencing. 5′ VDJ libraries were sequenced to a depth of 5,000 reads per cell. 5′
DGE libraries were sequenced to a depth of 50,000 reads per cell.
837
Single cell data pre-processing, quality control, clustering and integration
Cell Ranger v6.1.2 was used to demultiplex FASTQ reads, perform sequence alignment to the
GRCh38 transcriptome, and extract unique molecular identifier (UMI) barcodes. Single cell gene
expression matrices were analyzed using the R package Seurat v4.1.1 as a single Seurat object.
Cells were filtered to include only cells with less than 25% mitochondrial RNA content and
between 200 and 4000 genes detected. For single-cell VDJ sequencing, only cells with full-length
sequences were retained. Raw count data were normalized using the Seurat function
SCTransform to normalize raw count data to a Gamma-Poisson Generalized Linear Model,
perform variance stabilization, identify highly variable features, and scale features.[80,81] Cells were
projected into their first 50 principal components using the RunPCA function in Seurat, and further
reduced into a 2-dimensional visualization space using the RunUMAP function. Initial cell cluster
identification was performed using the Seurat function FindClusters at a resolution of 0.7. Initial
cell type assignment was performed by reference mapping to the human PBMC dataset
associated with the R package Azimuth.[82] Cluster identities were then manually assigned by
identification of differentially expressed genes using the MAST hurdle model as implemented in
the Seurat FindAllMarkers function with a log fold change threshold of 0.25 and minimum
fractional expression threshold of 0.25.[83] Integration of single cell TCR-seq and BCR-seq data
into the scRNA-seq data was performed using the R package scRepertoire.[84] For each patient,
TCRβ sequences identified in single cell data were compared against TCRβ identified in
microdissected TLS to identify T cells present in TLS. In cases where single cells with the same
TCR occupied multiple clusters, a putative transcriptional phenotype was assigned to a T cell in
the TLS repertoire according to the most common T cell subset to which the single cells with the
same TCR belonged. No matches were identified between IGH sequences identified by Adaptive
sequencing and IGH sequences in the single cell dataset, and thus we excluded B cells in the
single cell dataset from further analysis.
863
Data availability
Bulk RNA-seq, single cell RNA/TCR-seq data, and imaging mass cytometry data from this study
are deposited in dbGap under *** and the Gene Expression Omnibus (GEO) under accession
number GSE ***. Bulk TCRβ and IGH data from microdissected TLS are available on the Adaptive
ImmunoSEQ web analyzer portal at ***. All other relevant data are available from the
corresponding authors upon request.
870
871 **Code availability**
872 All custom code used to generate the results in this study has been deposited in a GitHub
873 repository at https://github.com/FertigLab/HCCTLS.
874

Acknowledgements

This work was supported by the Johns Hopkins SPORE in Gastrointestinal Cancer and the National Institutes of Health (NIH) U01CA253403 and U01CA212007 grants to M.Y. and E.J.F. Additional research support was provided by the Breeden-Adams Foundation and Conquer Cancer to M.Y.; the Johns Hopkins School of Medicine J. Mario Molina Physician Scientist Fund, Linda Rubin Fellowship, and Bernice Garchik Fellowship to D.S; and the Maryland Cancer Moonshot Research Grant to the Johns Hopkins Medical Institutions to D.J.Z. We thank Theinmozhi Arulraj for reviewing the final manuscript.

Author contributions

D.S., M.Y., and E.J.F. conceived and designed this study. D.S., L.K., M.Y., and L.D. performed data analysis and interpreted results. Q.Z. and R.A. performed the pathologic review. D.S., K.M., Q.Z., and R.A. performed the histologic analysis. All authors assisted with the data analysis, provided valuable discussion, and reviewed and edited the final manuscript draft. D.S. and M.Y. wrote the manuscript with input from all the authors.

Competing interests

M.Y. reports grant/research support from Bristol-Myers Squibb, Incyte, Genentech (to Johns Hopkins) and honoraria from Genentech, Exelixis, Eisai, AstraZeneca, Replimune, Hepion, and equity in Adventris Pharmaceuticals. E.J.F is on the Scientific Advisory Board of Viosera/Reistance Bio, is a paid consultant for Merck and Mestag Therapeutics, and receives research funds from Abbvie. W.J.H. has received patent royalties from Rodeo/Amgen and is the recipient of grants from Sanofi, NeoTX, and CirclePharma. He has received speaking/travel honoraria from Exelixis and Standard BioTools. E.M.J. reports grant/research support from the Lustgarten Foundation, Break Through Cancer, Genentech, Bristol-Meyers Squibb; honoraria from Achilles, DragonFly, Parker Institute, Cancer Prevention and Research Institute of Texas, Surge, HDT Bio, Mestag Therapeutics, Medical Home Group; and equity in AbMeta Therapeutics and Adventris Pharmaceuticals. D.J.Z. reports grant/research support from Roche/Genentech.

REFERENCES

1. Garon EB, Hellmann MD, Rizvi NA, et al. Five-Year Overall Survival for Patients With Advanced Non–Small-Cell Lung Cancer Treated With Pembrolizumab: Results From the Phase I KEYNOTE-001 Study. *J Clin Oncol Off J Am Soc Clin Oncol*. 2019;37(28):2518-2527. doi:10.1200/JCO.19.00934

2. Larkin J, Chiarion-Sileni V, Gonzalez R, et al. Five-Year Survival with Combined Nivolumab and Ipilimumab in Advanced Melanoma. *N Engl J Med*. Published online September 27, 2019. doi:10.1056/NEJMoa1910836

3. Patel SP, Othus M, Chen Y, et al. Neoadjuvant–Adjuvant or Adjuvant-Only Pembrolizumab in Advanced Melanoma. *N Engl J Med*. 2023;388(9):813-823. doi:10.1056/NEJMoa2211437

4. Topalian SL, Forde PM, Emens LA, Yarchoan M, Smith KN, Pardoll DM. Neoadjuvant immune checkpoint blockade: A window of opportunity to advance cancer immunotherapy. *Cancer Cell*. Published online August 17, 2023. doi:10.1016/j.ccell.2023.07.011

5. Liu J, Blake SJ, Yong MCR, et al. Improved Efficacy of Neoadjuvant Compared to Adjuvant Immunotherapy to Eradicate Metastatic Disease. *Cancer Discov*. 2016;6(12):1382-1399. doi:10.1158/2159-8290.CD-16-0577

6. Liu J, O'Donnell JS, Yan J, et al. Timing of neoadjuvant immunotherapy in relation to surgery is crucial for outcome. *Oncoimmunology*. 2019;8(5):e1581530. doi:10.1080/2162402X.2019.1581530

7. Chalabi M, Fanchi LF, Dijkstra KK, et al. Neoadjuvant immunotherapy leads to pathological responses in MMR-proficient and MMR-deficient early-stage colon cancers. *Nat Med*. 2020;26(4):566-576. doi:10.1038/s41591-020-0805-8

8. Ho WJ, Zhu Q, Durham J, et al. Neoadjuvant cabozantinib and nivolumab convert locally advanced hepatocellular carcinoma into resectable disease with enhanced antitumor immunity. *Nat Cancer*. Published online July 29, 2021:1-13. doi:10.1038/s43018-021-00234-4

9. Cabrita R, Lauss M, Sanna A, et al. Tertiary lymphoid structures improve immunotherapy and survival in melanoma. *Nature*. 2020;577(7791):561-565. doi:10.1038/s41586-019-1914-8

10. Petitprez F, de Reyniès A, Keung EZ, et al. B cells are associated with survival and immunotherapy response in sarcoma. *Nature*. 2020;577(7791):556-560. doi:10.1038/s41586-019-1906-8

11. Helmink BA, Reddy SM, Gao J, et al. B cells and tertiary lymphoid structures promote immunotherapy response. *Nature*. 2020;577(7791):549-555. doi:10.1038/s41586-019-1922-8

12. Vanhersecke L, Brunet M, Guégan JP, et al. Mature tertiary lymphoid structures predict immune checkpoint inhibitor efficacy in solid tumors independently of PD-L1 expression. *Nat Cancer*. 2021;2(8):794-802. doi:10.1038/s43018-021-00232-6

13. Italiano A, Bessede A, Pulido M, et al. Pembrolizumab in soft-tissue sarcomas with tertiary lymphoid structures: a phase 2 PEMBROSARC trial cohort. *Nat Med*. Published online May 26, 2022. doi:10.1038/s41591-022-01821-3

14. Joshi NS, Akama-Garren EH, Lu Y, et al. Regulatory T Cells in Tumor-Associated Tertiary Lymphoid
Structures Suppress Anti-tumor T Cell Responses. *Immunity*. 2015;43(3):579-590.
doi:10.1016/j.immuni.2015.08.006

15. Rodriguez AB, Peske JD, Woods AN, et al. Immune mechanisms orchestrate tertiary lymphoid
structures in tumors via cancer-associated fibroblasts. *Cell Rep*. 2021;36(3):109422.
doi:10.1016/j.celrep.2021.109422

16. Calderaro J, Petitprez F, Becht E, et al. Intra-tumoral tertiary lymphoid structures are associated
with a low risk of early recurrence of hepatocellular carcinoma. *J Hepatol*. 2019;70(1):58-65.
doi:10.1016/j.jhep.2018.09.003

17. Cottrell TR, Thompson ED, Forde PM, et al. Pathologic features of response to neoadjuvant anti-PD-
1 in resected non-small-cell lung carcinoma: a proposal for quantitative immune-related pathologic
response criteria (irPRC). *Ann Oncol*. 2018;29(8):1853-1860. doi:10.1093/annonc/mdy218

18. Claeskens G, Hjort NL. *Model Selection and Model Averaging*. Cambridge University Press; 2008.
doi:10.1017/CBO9780511790485

19. Santiago T, Kulemzin SV, Reshetnikova ES, et al. FCRLA is a resident endoplasmic reticulum protein
that associates with intracellular Igs, IgM, IgG and IgA. *Int Immunol*. 2011;23(1):43-53.
doi:10.1093/intimm/dxq456

20. Lainé A, Labiad O, Hernandez-Vargas H, et al. Regulatory T cells promote cancer immune-escape
through integrin αvβ8-mediated TGF-β activation. *Nat Commun*. 2021;12(1):6228.
doi:10.1038/s41467-021-26352-2

21. Dodagatta-Marri E, Ma HY, Liang B, et al. Integrin αvβ8 on T cells suppresses anti-tumor immunity in
multiple models and is a promising target for tumor immunotherapy. *Cell Rep*. 2021;36(1):109309.
doi:10.1016/j.celrep.2021.109309

22. García-Serna AM, Alcaraz-García MJ, Ruiz-Lafuente N, et al. Dock10 regulates CD23 expression and
sustains B-cell lymphopoiesis in secondary lymphoid tissue. *Immunobiology*. 2016;221(12):1343-
1350. doi:10.1016/j.imbio.2016.07.015

23. Theisen DJ, Davidson JT, Briseño CG, et al. WDFY4 is required for cross-presentation in response to
viral and tumor antigens. *Science*. 2018;362(6415):694-699. doi:10.1126/science.aat5030

24. Coppola D, Nebozhyn M, Khalil F, et al. Unique Ectopic Lymph Node-Like Structures Present in
Human Primary Colorectal Carcinoma Are Identified by Immune Gene Array Profiling. *Am J Pathol*.
2011;179(1):37-45. doi:10.1016/j.ajpath.2011.03.007

25. Schumacher TN, Thommen DS. Tertiary lymphoid structures in cancer. *Science*. 375(6576):eabf9419.
doi:10.1126/science.abf9419

26. Jacobsen JT, Hu W, R. Castro TB, et al. Expression of Foxp3 by T follicular helper cells in end-stage
germinal centers. *Science*. 2021;373(6552):eabe5146. doi:10.1126/science.abe5146

27. Arulraj T, Binder SC, Meyer-Hermann M. Investigating the Mechanism of Germinal Center
Shutdown. *Front Immunol*. 2022;13. Accessed March 4, 2023.
https://www.frontiersin.org/articles/10.3389/fimmu.2022.922318

28. Milutinovic S, Abe J, Godkin A, Stein JV, Gallimore A. The Dual Role of High Endothelial Venules in
Cancer Progression versus Immunity. *Trends Cancer*. 2021;7(3):214-225.
doi:10.1016/j.trecan.2020.10.001

29. Asrir A, Tardiveau C, Coudert J, et al. Tumor-associated high endothelial venules mediate
lymphocyte entry into tumors and predict response to PD-1 plus CTLA-4 combination
immunotherapy. *Cancer Cell*. 2022;40(3):318-334.e9. doi:10.1016/j.ccell.2022.01.002

30. Astarita JL, Cremasco V, Fu J, et al. The CLEC-2–podoplanin axis controls fibroblastic reticular cell
contractility and lymph node microarchitecture. *Nat Immunol*. 2015;16(1):75-84.
doi:10.1038/ni.3035

31. Aoufouchi S, Faili A, Zober C, et al. Proteasomal degradation restricts the nuclear lifespan of AID. *J
Exp Med*. 2008;205(6):1357-1368. doi:10.1084/jem.20070950

32. ZAN H, CASALI P. Regulation of Aicda expression and AID activity. *Autoimmunity*. 2013;46(2):83-101.
doi:10.3109/08916934.2012.749244

33. Yoshitomi H, Ueno H. Shared and distinct roles of T peripheral helper and T follicular helper cells in
human diseases. *Cell Mol Immunol*. 2021;18(3):523-527. doi:10.1038/s41423-020-00529-z

34. Rao DA, Gurish MF, Marshall JL, et al. Pathologically expanded peripheral T helper cell subset drives
B cells in rheumatoid arthritis. *Nature*. 2017;542(7639):110-114. doi:10.1038/nature20810

35. Bocharnikov AV, Keegan J, Wacleche VS, et al. PD-1[hi]CXCR5[−] T peripheral helper cells promote B cell
responses in lupus via MAF and IL-21. *JCI Insight*. 2019;4(20). doi:10.1172/jci.insight.130062

36. Kim JR, Lim HW, Kang SG, Hillsamer P, Kim CH. Human CD57+ germinal center-T cells are the major
helpers for GC-B cells and induce class switch recombination. *BMC Immunol*. 2005;6:3.
doi:10.1186/1471-2172-6-3

37. Marinova E, Han S, Zheng B. Germinal Center Helper T Cells Are Dual Functional Regulatory Cells
with Suppressive Activity to Conventional CD4+ T Cells. *J Immunol*. 2007;178(8):5010-5017.
doi:10.4049/jimmunol.178.8.5010

38. Palmer BE, Blyveis N, Fontenot AP, Wilson CC. Functional and Phenotypic Characterization of
CD57+CD4+ T Cells and Their Association with HIV-1-Induced T Cell Dysfunction1. *J Immunol*.
2005;175(12):8415-8423. doi:10.4049/jimmunol.175.12.8415

39. Hong W, Yang B, He Q, Wang J, Weng Q. New Insights of CCR7 Signaling in Dendritic Cell Migration
and Inflammatory Diseases. *Front Pharmacol*. 2022;13. Accessed January 30, 2023.
https://www.frontiersin.org/articles/10.3389/fphar.2022.841687

40. Horiuchi S, Wu H, Liu WC, et al. Tox2 is required for the maintenance of GC TFH cells and the
generation of memory TFH cells. *Sci Adv*. 2021;7(41):eabj1249. doi:10.1126/sciadv.abj1249

41. van der Leun AM, Thommen DS, Schumacher TN. CD8+ T cell states in human cancer: insights from
single-cell analysis. *Nat Rev Cancer*. 2020;20(4):218-232. doi:10.1038/s41568-019-0235-4

42. Caushi JX, Zhang J, Ji Z, et al. Transcriptional programs of neoantigen-specific TIL in anti-PD-1-
treated lung cancers. *Nature*. 2021;596(7870):126-132. doi:10.1038/s41586-021-03752-4

43. Liu B, Zhang Y, Wang D, Hu X, Zhang Z. Single-cell meta-analyses reveal responses of tumor-reactive
CXCL13+ T cells to immune-checkpoint blockade. *Nat Cancer*. 2022;3(9):1123-1136.
doi:10.1038/s43018-022-00433-7

44. Lucca LE, Axisa PP, Lu B, et al. Circulating clonally expanded T cells reflect functions of tumor-
infiltrating T cells. *J Exp Med*. 2021;218(4):e20200921. doi:10.1084/jem.20200921

45. Posch F, Silina K, Leibl S, et al. Maturation of tertiary lymphoid structures and recurrence of stage II
and III colorectal cancer. *Oncoimmunology*. 2017;7(2):e1378844.
doi:10.1080/2162402X.2017.1378844

46. Siliņa K, Soltermann A, Attar FM, et al. Germinal Centers Determine the Prognostic Relevance of
Tertiary Lymphoid Structures and Are Impaired by Corticosteroids in Lung Squamous Cell
Carcinoma. *Cancer Res*. 2018;78(5):1308-1320. doi:10.1158/0008-5472.CAN-17-1987

47. Arulraj T, Binder SC, Robert PA, Meyer-Hermann M. Germinal Centre Shutdown. *Front Immunol*.
2021;12. Accessed March 4, 2023.
https://www.frontiersin.org/articles/10.3389/fimmu.2021.705240

48. Gavil NV, Scott MC, Weyu E, et al. Chronic antigen in solid tumors drives a distinct program of T cell
residence. *Sci Immunol*. 2023;8(84):eadd5976. doi:10.1126/sciimmunol.add5976

49. Garaud S, Dieu-Nosjean MC, Willard-Gallo K. T follicular helper and B cell crosstalk in tertiary
lymphoid structures and cancer immunotherapy. *Nat Commun*. 2022;13(1):2259.
doi:10.1038/s41467-022-29753-z

50. Gutiérrez-Melo N, Baumjohann D. T follicular helper cells in cancer. *Trends Cancer*. 2023;9(4):309-
325. doi:10.1016/j.trecan.2022.12.007

51. Gu-Trantien C, Loi S, Garaud S, et al. CD4[+] follicular helper T cell infiltration predicts breast cancer
survival. *J Clin Invest*. 2013;123(7):2873-2892. doi:10.1172/JCI67428

52. Gu-Trantien C, Migliori E, Buisseret L, et al. CXCL13-producing TFH cells link immune suppression
and adaptive memory in human breast cancer. *JCI Insight*. 2017;2(11):e91487, 91487.
doi:10.1172/jci.insight.91487

53. Argyriou A, Wadsworth MH, Lendvai A, et al. Single cell sequencing identifies clonally expanded
synovial CD4+ TPH cells expressing GPR56 in rheumatoid arthritis. *Nat Commun*. 2022;13(1):4046.
doi:10.1038/s41467-022-31519-6

54. Chung Y, Tanaka S, Chu F, et al. Follicular regulatory T cells expressing Foxp3 and Bcl-6 suppress
germinal center reactions. *Nat Med*. 2011;17(8):983-988. doi:10.1038/nm.2426

55. Linterman MA, Pierson W, Lee SK, et al. Foxp3+ follicular regulatory T cells control the germinal
center response. *Nat Med*. 2011;17(8):975-982. doi:10.1038/nm.2425

56. Sage PT, Francisco LM, Carman CV, Sharpe AH. The receptor PD-1 controls follicular regulatory T
cells in the lymph nodes and blood. *Nat Immunol*. 2013;14(2):152-161. doi:10.1038/ni.2496

57. Sage PT, Paterson AM, Lovitch SB, Sharpe AH. The coinhibitory receptor CTLA-4 controls B cell
responses by modulating T follicular helper, T follicular regulatory, and T regulatory cells. *Immunity*.
2014;41(6):1026-1039. doi:10.1016/j.immuni.2014.12.005

58. Wing JB, Ise W, Kurosaki T, Sakaguchi S. Regulatory T cells control antigen-specific expansion of Tfh
cell number and humoral immune responses via the coreceptor CTLA-4. *Immunity*. 2014;41(6):1013-
1025. doi:10.1016/j.immuni.2014.12.006

59. Raftery AE. Bayesian Model Selection in Social Research. *Sociol Methodol*. 1995;25:111-163.
doi:10.2307/271063

60. Abbott CW, Boyle SM, Pyke RM, et al. Prediction of Immunotherapy Response in Melanoma through
Combined Modeling of Neoantigen Burden and Immune-Related Resistance Mechanisms. *Clin
Cancer Res Off J Am Assoc Cancer Res*. 2021;27(15):4265-4276. doi:10.1158/1078-0432.CCR-20-
4314

61. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data
with DESeq2. *Genome Biol*. 2014;15(12):550. doi:10.1186/s13059-014-0550-8

62. Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide
expression profiles | PNAS. Accessed June 12, 2023.
https://www.pnas.org/doi/10.1073/pnas.0506580102

63. Korotkevich G, Sukhov V, Budin N, Shpak B, Artyomov MN, Sergushichev A. Fast gene set
enrichment analysis. Published online February 1, 2021:060012. doi:10.1101/060012

64. Levy E, Milani P, Bartha G, et al. 67 B-cell receptor heavy chain repertoire profiling using an
augmented transcriptome. *J Immunother Cancer*. 2020;8(Suppl 3). doi:10.1136/jitc-2020-
SITC2020.0067

65. Bolotin DA, Poslavsky S, Mitrophanov I, et al. MiXCR: software for comprehensive adaptive
immunity profiling. *Nat Methods*. 2015;12(5):380-381. doi:10.1038/nmeth.3364

66. ImmunoMind Team. immunarch: an R package for painless bioinformatics analysis of T-cell and B-
cell immune repertoires. Zenodo https://doi.org/10.5281/zenodo.3367200 (2019).

67. Montagne JM, Zheng XA, Pinal-Fernandez I, et al. Ultra-efficient sequencing of T Cell receptor
repertoires reveals shared responses in muscle from patients with Myositis. *EBioMedicine*.
2020;59:102972. doi:10.1016/j.ebiom.2020.102972

68. Zabransky DJ, Danilova L, Leatherman JM, et al. Profiling of syngeneic mouse HCC tumor models as a
framework to understand anti–PD-1 sensitive tumor microenvironments. *Hepatology*.
2023;77(5):1566. doi:10.1002/hep.32707

69. Zanotelli VRT, Bodenmiller B. ImcSegmentationPipeline: A pixelclassification based multiplexed
1081   image segmentation pipeline. Published online September 14, 2017. doi:10.5281/zenodo.3841961

1082   70. Carpenter AE, Jones TR, Lamprecht MR, et al. CellProfiler: image analysis software for identifying
1083   and quantifying cell phenotypes. *Genome Biol*. 2006;7(10):R100. doi:10.1186/gb-2006-7-10-r100

1084   71. Berg S, Kutra D, Kroeger T, et al. ilastik: interactive machine learning for (bio)image analysis. *Nat*
1085   *Methods*. 2019;16(12):1226-1232. doi:10.1038/s41592-019-0582-9

1086   72. Schapiro D, Jackson HW, Raghuraman S, et al. histoCAT: analysis of cell phenotypes and interactions
1087   in multiplex image cytometry data. *Nat Methods*. 2017;14(9):873-876. doi:10.1038/nmeth.4391

1088   73. FlowSOM: Using self-organizing maps for visualization and interpretation of cytometry data - Van
1089   Gassen - 2015 - Cytometry Part A - Wiley Online Library. Accessed June 14, 2023.
1090   https://onlinelibrary.wiley.com/doi/full/10.1002/cyto.a.22625

1091   74. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. *Nat*
1092   *Methods*. 2012;9(7):671-675. doi:10.1038/nmeth.2089

1093   75. Epskamp S, Cramer AOJ, Waldorp LJ, Schmittmann VD, Borsboom D. qgraph: Network Visualizations
1094   of Relationships in Psychometric Data. *J Stat Softw*. 2012;48:1-18. doi:10.18637/jss.v048.i04

1095   76. Robins HS, Campregher PV, Srivastava SK, et al. Comprehensive assessment of T-cell receptor beta-
1096   chain diversity in alphabeta T cells. *Blood*. 2009;114(19):4099-4107. doi:10.1182/blood-2009-04-
1097   217604

1098   77. Carlson CS, Emerson RO, Sherwood AM, et al. Using synthetic templates to design an unbiased
1099   multiplex PCR assay. *Nat Commun*. 2013;4:2680. doi:10.1038/ncomms3680

1100   78. Gupta NT, Vander Heiden JA, Uduman M, Gadala-Maria D, Yaari G, Kleinstein SH. Change-O: a
1101   toolkit for analyzing large-scale B cell immunoglobulin repertoire sequencing data. *Bioinformatics*.
1102   2015;31(20):3356-3358. doi:10.1093/bioinformatics/btv359

1103   79. Venturi V, Kedzierska K, Tanaka MM, Turner SJ, Doherty PC, Davenport MP. Method for assessing
1104   the similarity between subsets of the T cell receptor repertoire. *J Immunol Methods*.
1105   2008;329(1):67-80. doi:10.1016/j.jim.2007.09.016

1106   80. Hafemeister C, Satija R. Normalization and variance stabilization of single-cell RNA-seq data using
1107   regularized negative binomial regression. *Genome Biol*. 2019;20(1):296. doi:10.1186/s13059-019-
1108   1874-1

1109   81. Ahlmann-Eltze C, Huber W. glmGamPoi: fitting Gamma-Poisson generalized linear models on single
1110   cell count data. *Bioinformatics*. 2020;36(24):5701-5702. doi:10.1093/bioinformatics/btaa1009

1111   82. Hao Y, Hao S, Andersen-Nissen E, et al. Integrated analysis of multimodal single-cell data. *Cell*.
1112   2021;184(13):3573-3587.e29. doi:10.1016/j.cell.2021.04.048

1113   83. Finak G, McDavid A, Yajima M, et al. MAST: a flexible statistical framework for assessing
1114   transcriptional changes and characterizing heterogeneity in single-cell RNA sequencing data.
1115   *Genome Biol*. 2015;16:278. doi:10.1186/s13059-015-0844-5

1116   84. Borcherding N, Bormann NL, Kraus G. scRepertoire: An R-based toolkit for single-cell immune
1117   receptor analysis. *F1000Research*. 2020;9:47. doi:10.12688/f1000research.22139.2

1118

    Table 1 | Characteristics of the Patients, According to Treatment Status

	Neoadjuvant (N=19)	Untreated (N=14)	Total (N=33)
Age at surgery — yr			
Mean (SD)	64±10	66±9.0	65±9.6
Median (range)	65 (41-79)	66 (49-84)	65 (41-84)
Sex — no. (%)			
Male	11 (58)	11 (79)	22 (67)
Female	8 (42)	3 (21)	11 (33)
Histologic grade — no. (%)			
Poorly differentiated	3 (16)	1 (7)	4 (12)
Moderately differentiated	13 (68)	8 (57)	21 (64)
Well differentiated	3 (16)	5 (36)	8 (24)
Etiology — no. (%)			
HBV	3 (16)	1 (7)	4 (12)
HCV	7 (37)	6 (43)	13 (39)
HBV/HCV	1 (5)	1 (7)	2 (6)
NASH	2 (11)	3 (21)	5 (15)
ETOH	1 (5)	1 (7)	2 (6)
Unknown	5 (26)	2 (14)	7 (21)
Neoadjuvant Treatment — no. (%)			
anti-PD1 + TKI	14 (74)	0 (0)	14 (42)
anti-PD1	3 (16)	0 (0)	3 (9)
anti-PD1 + anti-CTLA4	1 (5)	0 (0)	1 (3)
anti-PD1 + anti-CTLA4 + TKI	1 (5)	0 (0)	1 (3)
None	0 (0)	14 (100)	14 (42)

1120
1121
1122

1123
1124

Fig. 1 | Neoadjuvant ICB induces intratumoral TLS, which are associated with superior pathologic response and relapse free survival. a, Workflow for TLS density analysis. **b,** Representative images of formalin fixed paraffin embedded (FFPE) HCC tumors stained with anti-CD20 antibody. Annotations indicate boundary between tumor/tumor regression bed and adjacent normal parenchyma (red), extension of boundary by 200 μm (yellow), intratumoral TLS (arrow), and peritumoral TLS (arrow head). Inset shows representative TLS at high magnification. Scale bar, 1mm. **c,** Box-and-whisker plots showing intratumoral TLS density in patients with locally advanced HCC treated with neoadjuvant ICB (n = 19) and untreated controls (n = 14). **d,** Boxplot-and-whisker plots showing intratumoral TLS density in untreated (n = 14) and neoadjuvant treated tumors, divided according to pathologic response (n = 19). For each box-and-whisker plot, the horizontal bar indicates the median, the upper and lower limits of the boxes the interquartile range, and the ends of the whiskers 1.5 times the interquartile range. **e-f,** Kaplan-Meier curves showing relapse free survival (**e**) and overall survival (**f**) for patients with HCC in the highest tertile (purple) compared to the middle and lowest tertiles (green) of intratumoral TLS density after neoadjuvant ICB. Statistical significance was determined by two-tailed t-test (**c**), one-way ANOVA followed by Tukey's honest significant difference (HSD) test (**d**), and log-rank test (**e** and **f**).

Fig. 2 | High TLS density is associated with increased T and B cell activation in HCC treated with neoadjuvant ICB. a, Principle component analysis of bulk RNA-sequencing of resected HCC tumors treated with neoadjuvant ICB (n = 12), divided according to TLS density relative to mean density of the neoadjuvant treatment group. **b,** Heatmap showing differentially expressed genes (DEG) with a log2 fold change > 1 and P < 0.05 between tumors with high (n = 5) and low (n = 7) TLS density. Annotation rows indicate TLS group, HCC etiology, treatment, response, relapse, and TLS density. Annotation columns at right identify DEG belonging to Gene Oncology Biological Pathways gene sets for T cell activation, B cell activation, Cytokine production, and Dendritic Cell Antigen Processing and Presentation. **c,** Volcano plot showing differentially expressed genes between tumors with high and low TLS density. Vertical dotted lines represents log2 fold change greater than or less than 1. Horizontal dotted line indicates adjusted P value of 0.05. 4 outlier genes are excluded from the plot for the purposes of visualization. **d,** Gene set enrichment analysis showing differentially enriched gene sets from the HALLMARK database between tumors with high and low TLS density. **e,** Barcode plots showing enrichment scores for the Gene Ontology Biological Pathways gene sets for T cell activation, B cell activation, and Dendritic Cell Antigen Processing and Presentation.

Fig. 3 | Identification of divergent TLS morphologies and cellular spatial relationships in viable tumor and tumor regression bed. a, Representative formalin-fixed, paraffin embedded (FFPE) neoadjuvant ICB-treated tumor stained with hematoxylin and eosin (H&E) showing divergent TLS morphologies ("mature" and "involuted") in viable residual viable tumor and regression bed. Dotted line shows boundary between residual viable tumor and regression bed. Blue arrows indicate mature TLS and red arrows indicate involuted TLS. Scale bar, 2.5mm. Higher magnification images of representative mature and involuted TLSare shown on right with serial sections stained with dual immunohistochemistry staining for CD20 (magenta) and Ki67 (brown), CD3 (magenta) and CD21 (brown), and CD4 (magenta) and CD8 (brown). Scale bars, 250 µm. **b-c,** Representative images of mature (**b**) and involuted (**c**) TLS obtained by imaging mass

cytometry. Insets show higher magnification images of CD8$^+$ T cells trafficking through high
endothelial venules (**b, far left**), an extensive CD21$^+$CD23$^+$ follicular dendritic cell network in the
mature morphology (**b**, **middle left**) compared to scant CD21$^+$ and CD23$^+$ in the involuted
morphology (**c**, **middle left**), close interactions between T cells and DCLAMP$^+$ mature dendritic
cells in the T cell zone adjacent to the germinal center (**b, middle right**), and high podoplanin
expression in the germinal center of the mature TLS (**b**, **far right**). Scale bars, 100 µm. **d,**
Heatmap showing average IMC marker expression in annotated cell clusters identified from
90,344 single cells from 38 TLS (*n* = 20 mature, *n* = 18 involuted). **e,** Composition of mature and
involuted TLS regions by cell type as a percentage of total cells per TLS. **f**, Box-and-whisker plots
showing cell cluster density in mature versus involuted TLS. For each box-and-whisker plot, the
horizontal bar indicates the median, the upper and lower limits of the boxes the interquartile range,
and the ends of the whiskers 1.5 times the interquartile range. **g,** Nearest neighbor analysis with
rows indicating individual clusters in mature and involuted TLS and columns corresopnding to first
and second most common neighbors. **h,** Network analysis for cell clusters in mature and involuted
TLS. Node size corresponds to the proportion of total cells for each TLS type occupied by each
cluster. Edge length represents the shortest distance between cell clusters and thickness
corresponds to the number of measurements for each TLS type. **i**, Box and violin plots showing
expression of mature dendritic cell markers (CD11c, CCR7, DCLAMP, HLADR, and CD86) in the
mature DC cluster and markers of T cell activation and exhaustion (CD45RO, CD25, CD69,
CD137, LAG3, PD1, and TOX) in the T peripheral helper (Tph) and cytotoxic T cell (Tc) clusters,
by TLS morphology. Statistical significance was determined by pairwise two sample Wilcoxon test
(**f** and **g**).
1287
1288

Fig. 4 | Expanded T cell clonotypes are shared across TLS within a tumor, while B cell repertoires of individual TLS are highly distinct. a, Workflow for T and B cell repertoire profiling of microdissected TLS (n = 30 mature and 5 involuted) from 7 patients. **b** and **d,** Upset plots showing overlap in unique TCRβ (**b**) and IGH (**d)** clonotypes across microdissected TLS from the same patient (P02). Barplots in gray and annotation row indicate distinct groups of clonotypes shared between different TLS. Top stacked barplots indicate composition of groups according to clonal expansion. Bottom right stacked barplots indicate total number of unique TCRβ or IGH clonotypes identified at each TLS according to degree of clonal expansion. **c** and **e,** Alluvial plots tracking the top 10 TCRβ (**c**) or IGH (**e**) clonotypes from TLS # 1 of patient P02 across all TLS microdissected from the patient's tumor. **f,** Box-and-whisker plot comparing the percentage of the TCRβ or IGH repertoire of each TLS that is shared with other TLS from the same tumor. **g,** Box-and-whisker plots comparing TCRβ clonality (as determined by Normalized Shannon Entropy) in mature and involuted TLS microdissected from patients P12, OT1, and OT6. Each point represents the TCRβ of an individual TLS. **h,** Violin plots comparing number of somatic hypermutations in IGH of mature and involuted TLS microdissected from patients P12, OT1, and OT6. Individual data points (not shown) represent individual IGH sequences. Statistical significance was determined by two-tailed t test (**f-h**).

Fig. 5 | Cytotoxic granzyme K and granzyme B-expressing CD8 T cells are highly represented in TLS. a, Uniform Manifold Approximation and Projection (UMAP) of 23,172 T cells identified by single cell RNA/TCR/BCR sequencing of CD3$^+$CD19$^+$ FACS-sorted peripheral blood from HCC patients treated with neoadjuvant ICB (n = 7). **b,** Barplot showing number of single cells per cluster. **c,** Violin plots showing expression of subset specific marker genes across clusters. **d-e,** UMAPs showing clonality of single cells with an associated T cell receptor sequence (**d**) and single cells with a TCRβ identified in microdissected TLS (**e**). **f,** Stacked barplot showing proportion of each single cell cluster identified in TLS. **g,** Inferred transcriptional phenotype of TCRβ clonotypes in microdissected TLS with a matching TCRβ in single cell sequencing of post-treatment peripheral blood (n = 7) or tumor infiltrating lymphocytes (n = 1). **h,** Inferred transcriptional phenotype of TCRβ clonotypes in mature and resolving TLS of patient OT6.

Fig. 6 | TLS structure and function in viable tumor and tumor regression bed in tumors treated with neoadjuvant checkpoint blockade. Mature TLS in viable tumor display a highly organized germinal center with close interactions between germinal center B cells and CD21[+] follicular dendritic cells, a T cell zone characterized by CD4[+] T peripheral helper cells in close proximity to mature dendritic cells, and cytotoxic CD8[+] T cells trafficking to the tumor via high endothelial venules. In areas of tumor regression, an involuted TLS morphology is found which displays dissolution of the B cell germinal center and persistence of Tph-DC interactions in the T cell zone, increased T cell memory marker expression, and clonal expansion of cytotoxic and tissue resident memory CD8[+] T cells.

Extended Data Table 1 | Clinical Characteristics of HCC cohort treated with neoadjuvant ICB. Characteristics of treated patients. Each row represents a single patient and columns indicate age at surgery, sex, HCC etiology, histologic grade, treatment regimen, pathologic response, and whether the patient suffered relapsed or death.

Patient	Age at surgery	Sex	Etiology	Histologic grade	Treatment	Response	Relapse	Death
P01	66	M	HCV	Moderately differentiated	α-PD1 + TKI	NR	No	No
P02	72	M	HCV	Moderately differentiated	α-PD1 + TKI	MPR	No	No
P03	76	F	Unknown	Moderately differentiated	α-PD1 + TKI	MPR	No	No
P07	64	F	HBV/HCV	Moderately differentiated	α-PD1 + TKI	pPR	Yes	Yes
P08	65	M	HCV	Moderately differentiated	α-PD1 + TKI	MPR	Yes	No
P09	47	M	HBV	Moderately differentiated	α-PD1 + TKI	CR	No	No
P10	41	F	HBV	Moderately differentiated	α-PD1 + TKI	pPR	Yes	Yes
P11	56	F	Unknown	Well differentiated	α-PD1 + TKI	pPR	No	No
P12	69	M	HCV	Moderately differentiated	α-PD1 + TKI	MPR	No	No
P13	74	F	Unknown	Poorly differentiated	α-PD1 + TKI	NR	No	No
P14	79	M	Unknown	Moderately differentiated	α-PD1 + TKI	pPR	Yes	No
P15	49	M	NASH	Moderately differentiated	α-PD1 + TKI	NR	Yes	Yes
OT1	68	M	ETOH	Poorly differentiated	α-PD1	MPR	No	No
OT2	71	F	Unknown	Well differentiated	α-PD1	pPR	No	No
OT3	63	M	HCV	Moderately differentiated	α-PD1 + TKI	pPR	Yes	No
OT4	69	F	HCV	Well differentiated	α-PD1 + TKI	pPR	Yes	No
OT5	62	M	NASH	Moderately differentiated	α-PD1	pPR	Yes	No
OT6	61	M	HCV	Moderately differentiated	α-PD1 + α-CTLA4 + TKI	MPR	No	No
OT7	56	F	HBV	Poorly differentiated	α-PD1 + α-CTLA4	CR	No	No

1388 **Extended Data Table 2 | Clinical characteristics of untreated HCC cohort.** Characteristics of
1389 untreated patients. Each row represents a single patient and columns indicate age at surgery,
1390 sex, HCC etiology, histologic grade, and treatment.
1391

Patient	Age at surgery	Sex	Etiology	Histologic grade	Treatment
C2	61	M	HCV	Moderately differentiated	none
C3	60	M	HBV/HCV	Well differentiated	none
C5	75	M	HCV/NASH	Well differentiated	none
C6	63	F	HCV	Moderately differentiated	none
C7	61	M	HCV/ETOH	Well differentiated	none
C8	57	M	NASH	Well differentiated	none
C9	65	M	HBV/ETOH	Moderately differentiated	none
C10	66	M	HCV	Well differentiated	none
C11	69	F	HCV	Moderately differentiated	none
C12	71	M	NASH	Moderately differentiated	none
C13	78	M	Unknown	Moderately differentiated	none
C14	66	M	Unknown	Moderately differentiated	none
C15	84	F	NASH	Poorly differentiated	none
C16	49	M	ETOH	Moderately differentiated	none

1392

1393

 Extended Data Table 3 | Immune Related Pathologic Response Criteria Scoring for HCC tumors treated with neoadjuvant anti-PD-1.

Feature, *n* (%)	MPR/CR (*n* = 8)	pPR/NR (*n* = 11)	P value†
Tumor			
Fibrosis			
Immature fibrosis	8 (100%)	7 (63.6%)	0.103
Mature fibrosis	8 (100%)	7 (63.6%)	0.103
Neovascularization	1 (12.5%)	0 (0%)	0.421
Cholesterol clefts	4 (50.0%)	2 (18.2%)	0.319
Granulomas	3 (37.5%)	2 (18.2%)	0.603
Foamy histiocytes	5 (62.5%)	4 (36.4%)	0.37
Giant cells	1 (12.5%)	2 (18.2%)	1
Hemosiderin macrophages	7 (87.5%)	5 (45.5%)	0.147
Calcification	1 (12.5%)	1 (9.1%)	1
Lymphoid aggregates	8 (100%)	8 (72.7%)	0.228
Tertiary lymphoid structures	6 (75.0%)	2 (18.2%)	0.0237
Dense plasma cells	2 (25.0%)	1 (9.1%)	0.546
Peritumor			
Lymphoid aggregates	7 (87.5%)	8 (72.7%)	0.603
Tertiary lymphoid structures	4 (50.0%)	3 (27.3%)	0.377
Dense plasma cells	1 (12.5%)	0 (0%)	0.421

1396   † Fisher's Exact test
1397

1398 **Extended Data Table 4 | Bayesian Information Criteria results for predicting relapse and**
1399 **death following surgical resection in HCC treated with neoadjuvant ICB.** Rows represent
1400 different BIC calculations and columns indicate outcomes and variables evaluated and calculated
1401 BIC.
1402

Outcome	Variable	BIC
Relapse	Total TLS density	43.04
Relapse	Peritumoral TLS density	45.4
Relapse	Intratumoral TLS	37.61
Relapse	Pathologic Response	40.18
Relapse	Sex	45.58
Relapse	Prior HCV	45.68
Relapse	Prior HBV	45.63
Death	Total TLS density	15.99
Death	Peritumoral TLS density	15.63
Death	Intratumoral TLS	15.99
Death	Pathologic Response	14.89
Death	Sex	17.2
Death	Prior HCV	17.85
Death	Prior HBV	16.22

1403

1404 **Extended Data Table 5 | Differentially expressed genes in TLS high and TLS low tumors.**
1405 Each row represents a single gene, and columns provide mean of normalized counts for all
1406 samples, log2 fold change in mRNA expression, Wald statistic, Wald test p-value, and Benjamini
1407 Hochberg adjusted p-values.
1408
1409 significantGenes.csv
1410
Extended Data Table 6 | Gene Set Enrichment Results showing enriched pathways in TLS
high tumors compared to TLS low tumors. Each tab corresponds to a different gene set in the
human MSigDB. Each row represents a single pathway, and columns provide the name of the
pathway, enrichment p-values, a Benjamini Hochberg adjusted p-value, the expected error for the
standard deviation of the P-value logarithm, enrichment score, enrichment score normalized to
mean enrichment of random samples of the same size, size of the pathway after removing genes
not present, and a vector with indexes of leading-edge genes that drive the enrichment.
1418
1419 pathwayAnalysisResultsCombined.xlsx
1420
1421

Extended Data Table 7 | Summary of antibodies selected for imaging mass cytometry.

1423

Lymphocyte	
CD45	Pan-leukocyte
CD45RA	Naïve lymphocyte
CD45RO	Antigen-experienced lymphocyte
CD3	T cell
CD4	T helper
CD8 (CD8a)	Cytotoxic T cell
CD20	B cell
CD21	Follicular dendritic cell and B cell
CD23	Follicular dendritic cell and B cell
CD138	Plasma cell
ICOS	T follicular helper
CXCR3	T follicular vs peripheral helper
CXCR5	T follicular vs peripheral helper
CCR7	Central and effector memory T cell, mature dendritic cell
FOXP3	Treg
Tumor/Structural	
PNAd	Lymphatic vessel
aSMA	Myofibroblast
Podoplanin	Fibroblast
Vimentin	Mesenchymal
Cytokeratin	Epithelial/Tumor
Functional	
CD25 (IL2R)	Activated/memory T cell
CD69	Activated/memory T cell
CD137 (4-1BB)	T cell activation
HLADR	APC, Activated T cell
PD1	T cell activation/exhaustion
PDL1	Macrophage, activated T/B, dendritic cell, epithelial
TOX	T cell activation/exhaustion
LAG3	T cell activation/exhaustion
GZMB	CTL, NK
Ki67	Proliferation Marker
BCL6	B cell GC regulator
AID	B cell somatic hypermutation
Myeloid	
CD11c	cDC1, cDC2
DCLAMP	Mature dendritic cell
DCSIGN	Macrophage and dendritic cell
CD57	NK cell, T cell senescence
CD68	Macrophage
CD86	DC, Langerhans cell, macrophage, B cell, other APC

1424

1425 **Extended Data Table 8 | Summary of metal-conjugated antibodies used for imaging mass**
1426 **cytometry.** Rows indicates a different stain, and columns indicate the metal, antigen, clone,
1427 dilution factor, source, and whether the antibody was custom-conjugated for the current study.
1428

Mass	Metal	Antigen	Clone	Dilution Factor	Source	Custom
89	Y	CD45	D9M8I	125	Cell Signaling Technology®	X
96-104	Ru	Counterstain			Electron Microscopy Sciences	
113	In	PNAd	MECA-79	250	Biolegend®	X
115	In	AID	mAID-2	125	eBioscience™	X
141	Pr	αSMA	1A4	500	Standard BioTools™	
142	Nd	Podoplanin	D2-40	125	Biolegend®	X
143	Nd	Vimentin	D21H3	500	Standard BioTools™	
144	Nd	CD11c	EP1347Y	250	Abcam	X
145	Nd	CD45RO	UCHL1	250	Biolegend®	X
146	Nd	CXCR3	IC6	100	AbboMax	X
147	Sm	CD69	EPR21814	250	Abcam	X
148	Nd	Pan-Keratin	C11	125	Standard BioTools™	
149	Sm	CD25 (IL2R)	SP176	250	Abcam	X
150	Nd	PD-L1	E1L3N	125	Cell Signaling Technology®	X
151	Eu	CXCR5	51505	250	Novus Biologicals	X
152	Sm	DC-LAMP	1010E1.01	125	Novus Biologicals	X
153	Eu	Tox/Tox2	E6I3Q	250	Cell Signaling Technology®	X
154	Sm	CD57	HNK-1	250	Cell Signaling Technology®	X
155	Gd	Foxp3	PCH101	100	Standard BioTools™	
156	Gd	CD4	EPR6855	125	Standard BioTools™	
158	Gd	ICOS	D1K2T™	250	Cell Signaling Technology®	X
159	Tb	CD68	KP1	100	Standard BioTools™	
160	Gd	Syndecan 1 (CD138)	IHC138	125	Cell Signaling Technology®	X
161	Dy	CD20	H1	125	Standard BioTools™	
162	Dy	CD8α	C8/144B	250	Standard BioTools™	
163	Dy	CD21	Bu32	250	Biolegend®	X
164	Dy	BCL-6	K112-91	250	BD Pharmingen™	X
165	Ho	PD1	EPR4877(2)	250	Abcam	X
166	Er	CD45RA	HI100	250	Standard BioTools™	
167	Er	Granzyme B	D6E9W	125	Cell Signaling Technology®	X
168	Er	Ki-67	B56	250	Standard BioTools™	
169	Tm	CD23	MRQ-57	125	Cell Marque™	X
170	Er	CD3ε	Polyclonal, C-terminal	125	Standard BioTools™	
171	Yb	Lag3	17B4	125	GeneTex	X
172	Yb	CD137 (4-1BB)	D2Z4Y	250	Cell Signaling Technology®	X
173	Yb	DC-SIGN	DCN46	62.5	Ionpath	X
174	Yb	HLA-DR	LN3	250	Standard BioTools™	
175	Lu	CD86	E2G8P	125	Cell Signaling Technology®	X
176	Yb	CCR7	EPR23192-57	250	Cell Signaling Technology®	X
191	Ir	DNA 1			Standard BioTools™	
193	Ir	DNA 2			Standard BioTools™	
195	Pt	Plasma Membrane 2	1A36	250	Standard BioTools™	
196	Pt	Plasma Membrane 3	1A37	250	Standard BioTools™	
198	Pt	Plasma Membrane 4	1A38	250	Standard BioTools™	

1429

Extended Data Table 9 | Characteristics of microdissected TLS. Each row indicates a different
patient with HCC treated with neoadjuvant immunotherapy. Columns indicate number of TLS
microdissected per patient according to location (tumor or normal adjacent) and morphology
(mature or involuted). Sample number shown differs from the final number of TCRβ and IGH
repertoires analyzed due to filtering to remove TCRβ repertoires with fewer than 500 clones and
IGH repertoires with fewer than 50 clones.
1436

	Tumor		Normal adjacent	
Patient	Mature	Involuted	Mature	Involuted
P02	2		4	
P03	5	1		
P07			5	
P08	5			
P12	3	1		
OT1	4	1		
OT6	4	3		
Total	*23*	*6*	9	

1437

Extended Data Table 10 | Expanded TCRβ clones in peripheral blood after neoadjuvant

ICB.

Patient	TCRβ sequence	pre-PBMC count	post-PBMC count	pre-PBMC freq	post-PBMC freq	pValue	fold change	present TLS
P02	CASSSLSDNYGYTF	1412	3064	0.011764608	0.025459289	2.1E-13	2.16	1
P02	CAISLDRGGEAFF	314	687	0.002616209	0.005708398	2.8E-08	2.18	1
P02	CASKPLVWNTGELFF	0	26	0	0.000216038	2.1E-06	Inf	0
P07	CASSEPQGQLTEAFF	0	44	0	0.000261729	2.3E-08	Inf	1
P12	CASSFGTSRRSEFF	9	127	6.6405E-05	0.000723144	7.8E-13	10.89	1
P12	CAISVDRGYSGANVLTF	9	114	6.6405E-05	0.000649121	1.9E-11	9.78	1
P12	CASSFLETQYF	16	124	0.000118053	0.000706062	1.8E-09	5.98	1
P12	CAWSRAAGGPNEQFF	63	250	0.000464835	0.001423512	5.4E-08	3.06	1
P12	CASSPGLAGDEQYF	1	33	7.37833E-06	0.000187904	7.4E-06	25.47	0
P12	CASSYTVGEYNEQFF	9873	17927	0.072846265	0.102077189	5.1E-05	1.40	1
P12	CASSLDAGASSYNSPLHF	35	125	0.000258242	0.000711756	7.1E-05	2.76	1
P12	CASSPEGQIRETQYF	1501	2954	0.011074875	0.016820216	8.5E-05	1.52	0
P12	CASSSDGAYLGTEAFF	0	21	0	0.000119575	1.0E-04	Inf	0

Extended Data Table 11 | TCR repertoire characteristics of peripheral blood and TIL single cell RNA/TCR-seq data

Peripheral blood

Cluster	Cells, *n*	Cells with TCR, *n*	Cells with expanded TCR, *n* (%)	Cells with TCRβ in TLS, *n* (%)
CD4 Naive	615	590	47 (8)	16 (2.7)
CD8 Naive	848	807	88 (10.9)	11 (1.4)
MAIT	315	217	108 (49.8)	129 (59.4)
CD4 Naive-like	8580	7224	806 (11.2)	604 (8.4)
CD4 TCM	2677	2531	464 (18.3)	767 (30.3)
CD4 Tph	1088	988	113 (11.4)	188 (19)
CD4 TEM_GZMK	1271	1086	384 (35.4)	702 (64.6)
CD4 TEM_GZMB	942	698	660 (94.6)	601 (86.1)
CD8 TCM	892	785	164 (20.9)	202 (25.7)
CD8 TEM_GZMK	1906	1676	1147 (68.4)	1390 (82.9)
CD8 TEM_GZMB	2551	1963	1682 (85.7)	1569 (79.9)
CD8 TRM	212	91	41 (45.1)	52 (57.1)
NK-T	75	32	25 (78.1)	22 (68.8)
Treg	801	712	106 (14.9)	70 (9.8)
dnT	267	116	23 (19.8)	23 (19.8)
gdT	132	30	7 (23.3)	3 (10)
Total	23172	19546	5865 (30)	6349 (32.5)

TIL

Cluster	Cells, *n*	Cells with TCR, *n*	Cells with expanded TCRs, *n* (%)	Cells with TCRβ in TLS, *n* (%)
CD4 Naive-like	38	11	3 (27.3)	8 (72.7)
CD4 TCM	28	14	1 (7.1)	10 (71.4)
CD4 Tph	31	29	7 (24.1)	24 (82.8)
CD4 TEM_GZMK	76	48	10 (20.8)	40 (83.3)
CD8 TCM	53	10	3 (30)	7 (70)
CD8 TEM_GZMK	194	147	68 (46.3)	121 (82.3)
CD8 TEM_GZMB	37	28	16 (57.1)	24 (85.7)
CD8 TRM	36	25	14 (56)	18 (72)
Treg	22	17	3 (17.6)	8 (47.1)
dnT	28	3	1 (33.3)	3 (100)
MAIT	19	14	10 (71.4)	10 (71.4)
Total	562	346	136 (39.3)	273 (78.9)

Extended Data Table 12 | Association of peripheral blood and TIL single cell clusters with clonal expansion.°†

Peripheral blood

Cluster	P value	OR	95%_CI_Lower	95%_CI_Upper
CD4 TEM_GZMB	< 0.001	31.48	25.72	38.83
CD8 TEM_GZMB	< 0.001	17.99	16.17	20.04
NK-T	< 0.001	10.27	4.73	23.62
Treg	< 0.001	9.02	5.42	16.23
CD8 TEM_GZMK	< 0.001	6.27	5.63	6.97
MAIT	0.398	0.82	0.54	1.22
CD8 TRM	0.389	0.73	0.35	1.39
gdT	0.614	0.59	0.12	1.93
CD4 TEM_GZMK	< 0.001	0.4	0.32	0.51
dnT	0.002	0.34	0.13	0.73
CD8 TCM	< 0.001	0.28	0.2	0.38
CD4 TCM	< 0.001	0.13	0.1	0.17
CD8 Naive	< 0.001	0.08	0.04	0.14
CD4 Naive	< 0.001	0.05	0.02	0.12
CD4 Naive-like	< 0.001	0.04	0.03	0.05
CD4 Tph	< 0.001	0.04	0.01	0.07

TIL

Cluster	P value	OR	95%_CI_Lower	95%_CI_Upper
Treg	0.03	Inf	1.2	Inf
CD8 TEM_GZMB	0.003	3.49	1.45	8.32
CD8 TRM	0.056	2.25	0.88	5.49
CD8 TEM_GZMK	0.013	1.94	1.12	3.36
MAIT	0.518	1.44	0.32	5.19
CD8 TCM	0.689	1.19	0.12	6.83
CD4 Tph	0.159	0.39	0.07	1.32
CD4 TEM_GZMK	< 0.001	0.13	0.02	0.53
CD4 TCM	0.046	0	0	1.05
CD4 Naive-like	0.131	0	0	1.4
dnT	1	0	0	8.64

°Clonal expansion defined as greater than 5 cells per unique TCRαβ in the peripheral blood dataset and greater than 2 in TIL.

† Fisher's Exact test

Extended Data Table 13 | Association of peripheral blood and TIL single cell clusters with detection in TLS.†

Peripheral blood

Cluster	P value	OR	95%_CI_Lower	95%_CI_Upper
CD4 TEM_GZMB	< 0.001	14.12	11.35	17.72
CD8 TEM_GZMK	< 0.001	12.65	11.08	14.48
CD8 TEM_GZMB	< 0.001	10.66	9.49	12
Treg	< 0.001	4.59	3.57	5.97
NK-T	< 0.001	4.58	2.08	10.85
CD4 TEM_GZMK	< 0.001	4.15	3.64	4.73
MAIT	< 0.001	3.09	2.33	4.11
CD8 TRM	< 0.001	2.79	1.8	4.34
CD4 TCM	0.012	0.89	0.81	0.98
CD8 TCM	< 0.001	0.71	0.6	0.84
dnT	0.003	0.51	0.31	0.82
CD4 Tph	< 0.001	0.47	0.4	0.56
gdT	0.006	0.23	0.04	0.75
CD4 Naive-like	< 0.001	0.1	0.1	0.11
CD4 Naive	< 0.001	0.06	0.03	0.09
CD8 Naive	< 0.001	0.03	0.01	0.05

TIL

Cluster	P value	OR	95%_CI_Lower	95%_CI_Upper
CD8 TCM	0.445	1.63	0.26	7.35
CD4 TCM	0.505	1.52	0.34	5.48
MAIT	0.505	1.52	0.34	5.48
CD8 TRM	0.444	1.5	0.51	3.96
CD4 Naive-like	0.706	1.42	0.24	6.11
CD4 Tph	0.812	0.76	0.22	2.15
CD4 TEM_GZMK	0.567	0.72	0.28	1.66
CD8 TEM_GZMK	0.23	0.7	0.39	1.22
CD8 TEM_GZMB	0.472	0.6	0.15	1.84
Treg	0.003	0.22	0.07	0.66
dnT	1	0	0	9.1

† Fisher's Exact test

Extended Data Table 14 | Match rate of TLS TCRβ in single cell sequencing of post-treatment peripheral blood and TIL.

Peripheral Blood

Patient	Total TCRβ, *n* (%)	Top 10% of TCRβ, *n* (%)	Top 1% of TCRβ, *n* (%)	Top 0.1% of TCRβ, *n* (%)
P02	274/11783 (2.3)	130/1178 (11)	52/118 (44.1)	8/12 (67)
P03	114/10872 (1)	39/1087 (3.6)	15/109 (13.8)	2/11 (18)
P07	624/11623 (5.4)	276/1162 (24)	61/116 (52.6)	11/12 (92)
P08	742/33681 (2.2)	374/3368 (11)	123/337 (36.5)	20/34 (59)
P12	422/7025 (6)	161/702 (23)	38/70 (54.3)	6/7 (86)
OT1	58/2740 (2.1)	39/274 (14)	9/27 (33.3)	1/3 (33)
OT6	674/58185 (1.2)	237/5818 (4.1)	71/582 (12.2)	15/58 (26)
Total	2908 / 135909 (2.1)	1256/13589 (9.2)	369/1359 (27.2)	63/137 (46)

TIL

Patient	Total TCRβ, *n* (%)	Top 10% of TCRβ, *n* (%)	Top 1% of TCRβ, *n* (%)	Top 0.1% of TCRβ, *n* (%)
OT6	196/58185 (0.34)	135/5818 (2.3)	76/582 (13.1)	24/58 (41)

1477 **Extended Data Table 15 | Summary of antibodies used for immunohistochemistry.** Each
1478 row indicates a different stain, and columns indicate the target, antigen retrieval buffer used,
1479 clone, vendor, product ID and concentration. For dual IHC stains, first and second antibody are
1480 indicated.
1481

Target	Antigen retrieval buffer	First antibody			Second antibody		
		Clone	Vendor	Conc. (µg/mL)	Clone	Vendor	Conc. (µg/mL)
CD20	low pH sodium citrate	L26	Leica	0.114			
CD21/CD3	low pH sodium citrate	2G9	Leica	4.075	LN10	Leica	0.213
CD8/CD4	high pH EDTA	4B11	Leica	0.114	ER204	Millipore Sigma	0.096
Ki67/CD20	high pH EDTA	MM1	Leica	0.42	L26	Leica	0.19

1482

1483 **Extended Data Table 16 | Summary of antibodies used for FACS.**
1484

Antibody	Clone	Vendor	Dilution
Live/Dead	Zombie NIR	Biolegend	1:1000
CD3-FITC	HIT3a	Biolegend	1:100
CD19-PE/dazzle	SJ25C1	Biolegend	1:200
Fc Block	N/A	Biolegend	1:100

1485

1486

 EXTENDED DATA

Extended Data Fig. 1 | TLS density in HCC tumors treated with neoadjuvant ICB and untreated controls. a, Box-and-whisker plots showing total and peritumoral TLS density in patients with locally advanced HCC treated with neoadjuvant ICB (n = 19) and untreated controls (n = 14). **b-c,** Stacked barplots showing proportion of TLS comprised of peritumoral verus intratumoral TLS location neoadjuvant treated and untreated HCC tumors (**b**) and by patient (**c**). Labels indicate proportion of total TLS comprised of peritumoral or intratumoral TLS. In **c**, patients with no observed TLS are not shown. **d-e,** Box-and-whisker plots showing total (**d**) and peritumoral (**e**) TLS density in untreated (n = 14) and neoadjuvant treated tumors, divided according to pathologic response (n = 19). Statistical significance was determined by two-tailed t-test (**a**) and one-way ANOVA followed by Tukey's honest significant difference (HSD) test (**d** and **e**).

Extended Data Fig. 2 | Relapse free survival and overall survival in HCC cohort treated with
neoadjuvant ICB, according to clinical covariates. a—l, Kaplan-Meier curves showing relapse
free survival and overall survival after surgical resection for HCC patients treated with neoadjuvant
ICB (n = 19), according to total TLS density (**a** and **b**), peritumoral TLS density (**c** and **d**),
pathologic response (**e** and **f**), sex (**g** and **h**), prior hepatitis C (HCV) infection (**i** and **j**), and prior
hepatitis B (HBV) infection (**k** and **l**). Statistical significance was determined by log-rank test.
1510

Extended Data Fig. 3 | High TLS density after neoadjuvant ICB is associated with increased expression of the 12-chemokine TLS gene signature. a, Heatmap showing expression of the 12-chemokine gene signature in tumors with high TLS density ($n = 5$) and low TLS density ($n = 7$). Annotation rows indicate TLS group, HCC etiology, neoadjuvant treatment, pathologic response, relapse, and TLS density.

1546

1547
1548
1549
Extended Data Fig. 4 | HCC tumors with high TLS density after neoadjuvant ICB have
expanded T and B cell repertoires compared to tumors with low TLS density. Box-and-
whisker plots showing the total clones, unique clonotypes, and effective number of clonotypes
(i.e. true diversity index) for the immunoglobulin heavy chain (IGH) **(a-c),** TCRα (**d-f**), and TCRβ
(**g-i**) repertoires of HCC tumors with high and low TLS density after neoadjuvant ICB. For each
box-and-whisker plot, the horizontal bar indicates the median, the upper and lower limits of the
boxes the interquartile range, and the ends of the whiskers 1.5 times the interquartile range.
Statistical significance was determined by Wilcoxon rank sum test.

1558

1559
1560
Extended Data Fig. 5 | Involuted TLS in an HCC tumor with complete pathologic response
after neoadjuvant ICB (OT7). a, Serial FFPE sections of an involuted TLS stained with anti-
CD20 antibody (brown). Numbered images indicate the order in which the sections were cut from
the tissue block. Scale bar, 250 μm. **b,** Representative images of multiple involuted TLS (red
arrows) stained with hematoxylin and eosin (H&E), anti-CD20 (magenta) and anti-Ki67 (brown)
(right middle), anti-CD3 (magenta) and anti-CD21 (brown) (middle right), and anti-CD4 and anti-
CD8 (bottom right).

Extended Data Fig. 6 | Characterization of divergent TLS morphologies in viable tumor and tumor regression bed by imaging mass cytometry. a, Imaging mass cytometry workflow. **b-c**, Dot plots showing representative mature (**b**) and involuted (**c**) TLS, colored according to cluster assignment of individual cells after cell segmentation. (**d**) Box-and-whisker plots showing cell cluster density in mature versus involuted TLS for CXCR3^{low} CD4 T cells, CD57$^+$ CD4 T cells, Macrophages, and Stroma. For each box-and-whisker plot, the horizontal bar indicates the median, the upper and lower limits of the boxes the interquartile range, and the ends of the whiskers 1.5 times the interquartile range. Statistical significance was determined by pairwise two sample Wilcoxon test (**d**).

Extended Data Fig. 7 | TCRβ repertoire features of microdissected TLS. a, Representative images showing method of identification and microdissection of individual TLS. Image on left shows HCC tumor stained with hematoxylin and eosin (H&E) at low magnification. Insets show higher magnification of staining with H&E, anti-CD20 (magenta) and anti-Ki67 (brown), anti-CD3 (magenta) and anti-CD21 (brown), anti-CD4 and anti-CD8 (bottom right), and corresponding pre- and post-microdissection images. Scale bar, 1mm. **b,** Barplot showing total clone count across all microdissected TLS. **c-h,** Representative upset plots showing overlap in TCRβ clonotypes across microdissected TLS from patients P03 (**c**), P07 (**d**), P08 (**e**), P12 (**f**), OT1 (**g**), and OT6 (**h**). For each upset plot, barplots in gray and row below indicate number of overlapping clonotypes between different combinations of TCRβ repertoires. Stacked barplots at top indicate repertoire composition of different groups of TCRβ and at bottom right indicate total number of unique TCRβ clonotypes identified in each TLS, colored according to clonal expansion. Intersections with fewer than 20 unique clonotypes are not shown. **i**, Dotplot showing TCRβ repertoire clonality (as determined by Normalized Shannon Entropy) for matched mature and involuted TLS. Statistical significance was determined by two-tailed t test (**i**).

none

Extended Data Fig. 8 | IGH repertoire features of microdissected TLS. a, Stacked barplot showing IGH repertoire composition across all TLS. **b-f,** Representative upset plots showing overlap in unique IGH clonotypes across microdissected TLS from patients P03 (**b**), P07 (**c**), P08 (**d**), P12 (**e**), OT1 (**f**), and OT7 (**g**). Bottom barplots and annotation row indicate number of overlapping clonotypes between different TLS repertoires. Top stacked barplots indicate clonal composition of overlapping ("public IGH") and nonoverlapping ("Private IGH"). Bottom right stacked barplots indicate total number of unique IGH clonotypes identified at each TLS and overall clonal composition. **h,** Dotplot showing IGH repertoire clonality (as determined by Normalized Shannon Entropy) for microdissected TLS, according to TLS morphology. Statistical significance was determined by two-tailed t test (**h** and **i**).

Extended Data Fig. 9 | TLS display a high degree of T cell repertoire overlap with pre- and post-treatment peripheral blood. a-b, Barplots showing proportion of unique TCRβ clonotypes at each TLS that also identified in matched pre-treatment (**a**) and post-treatment (**b**) peripheral blood.

Extended Data Fig. 10 | Single cell sequencing of post-treatment peripheral blood. a,
UMAPs showing gene expression of *CD3E*, *CD4*, *CD8A*, *CCR7*, *SELL*, *GZMK*, *PDCD1*, *CXCL13*,
TOX, and *ZNF683* across all single cells sequenced from post-treatment peripheral blood of 7
HCC patients treated with neoadjuvant ICB. **b,** Heatmap showing gene expression of the top 3
differentially expressed genes per cluster. Rows represent single genes and columns represent
individual cells. Annotation bar indicates cluster identity, whether each cell had a sequenced TCR,
the clonality of the TCR, and whether the TCR was identified in microdissected TLS from the
same patient. Clusters were downsampled to 75 cells per cluster for visualization. **c-e,** Volcano
plots showing differentially expressed genes in the CD8 TEM_GZMK (**b**),CD8 TEM_GZMB (**c**),
and CD4 Tph (**d**) clusters compared to all other cells. Vertical dotted lines indicates a fold change
of greater or less than 1.4 and horizontal line indicates a P value of 0.05. Labeled genes in **c** and
d indicate genes with the highest differential expression. Labeled genes in **e** indicate genes known
to be highly expressed in CD4 Tph.
1739

g

CD4 Naive-like CD4 TCM CD4 Tph CD4 TEM_GZMK CD8 TCM CD8 TEM_GZMK CD8 TEM_GZMB CD8 TRM Treg dnT MAIT

Identity
TCR
Clonal expansion
TCR in TLS

SELL
MALAT1
SESN3
GPR183
IL7R
FOS
CXCL13
PKM
MAF
TLE4
RUNX3
CCL5
GZMK
CCL4
NKG7
GZMH
GZMB
ZNF683
KLRK1
ANXA1
IL2RA
TBC1D4
BIRC3
NEAT1
HNRNPH1
DYNC1H1
KLRB1
MT2A
HSPA1A

TCR
No
Yes

Clonal expansion
Hyperexpanded (100 < X <= 500)
Large (20 < X <= 100)
Medium (5 < X <= 20)
Single (0 < X <= 1)
Small (1 < X <= 5)

TCR in TLS
No
Yes

CD8 TEM_GZMK versus non-CD8 TEM_GZMK

CD8 TEM_GZMB versus non-CD8 TEM_GZMB

CD4 Tph versus non-CD4 Tph

h

NS Log$_2$ FC p-value p-value and log$_2$ FC

$-Log_{10} P$

Log$_2$ fold change

CCL5, GZMK, GZMA, CD8A, CCL4, CST7, CD8B, DUSP2, IL7R, HLA-A, NKG7, TNFSF9, CD74, PTPRCAP, SELL, COTL1, CCL4L2, KLF2, LTB, TMEM123, MSB4X, GPR183, SPOCK2, DUSP4, ITGAE, NEAT1, RORA, MALAT1, RPS27, KLRB1, SNX9

total = 275 genes

i

NS Log$_2$ FC p-value and log$_2$ FC

$-Log_{10} P$

Log$_2$ fold change

NKG7, GZMH, GZMB, CCL5, IL7R, PTMS, IFNG, PRF1, KLRD1, KLRK1, CD8A, LAG3, TXNIP, PLCA4

total = 173 genes

j

NS Log$_2$ FC p-value p-value and log$_2$ FC

$-Log_{10} P$

Log$_2$ fold change

CTLA4, CXCL13, TNFRSF18, PKM, IL2RG, CD8A, SNX9, MAF, ANXA1, TNFRSF4, TP53INP1, RPL21, ITGB2, SPOCK2, CD8B, RPL38, CD4, DUSP4, PRDM1, RPL27A, GNA15, BTLA, GRGN, TFRC, CD69, RPL12, TSHZ2, CD7, RPL23A, UCP2, GPRIN3, PDCD1

total = 400 genes

k

Mature   Involuted

Proportion

CD8 TEM_GZMK
CD8 TEM_GZMB
CD8 TRM
dnT
MAIT
unmatched

TLS #1 TLS #3 TLS #4 TLS #5 TLS #6 TLS #7 TLS #8

Top 15 TCRβ Clonotypes per TLS

Extended Data Fig. 11 | Single cell sequencing of post-treatment TIL from patient OT6. a, Uniform Manifold Approximation and Projection (UMAP) for 562 T cells identified by single cell RNA/TCR/BCR sequencing of CD3$^+$CD19$^+$ FACS-sorted tumor infiltrating lymphocytes. **b,** Barplot showing number of single cells per cluster. **c,** Violin plots showing expression of subset specific marker genes across clusters. **d-e,** UMAPs showing clonality of single cells with an associated T cell receptor sequence (**d**) and single cells with a TCRβ identified in microdissected TLS (**e**). **f,** Stacked barplot showing proportion of each single cell cluster identified in TLS. **g,** Heatmap showing gene expression of the top 3 differentially expressed genes per cluster. Rows represent single genes and columns represent individual cells. Annotation bar indicates cluster identity, whether each cell had a sequenced TCR, the clonality of the TCR, and whether the TCR was identified in microdissected TLS from the same patient. **h-j,** Volcano plots showing differentially expressed genes in the CD8 TEM_GZMK (**h**),CD8 TEM_GZMB (**i**), and CD4 Tph **(j)** clusters compared to all other cells. Vertical dotted lines indicates a fold change of greater or less than 1.4 and horizontal line indicates a P value of 0.05. **k,** Inferred transcriptional phenotype of the top 15 TCRβ clonotypes in mature and involuted TLS of patient OT6.

a

TCRβ CDR3aa	*n* cells in peripheral blood	*n cells in* tumor
CAIKGGLAEETQYF	2	4
CASAQQDTDTQYF	2	2
CASGWDRRNTEAFF	1	1
CASRDQGGQTQYF	2	2
CASRSLTGQGAVRETYEQFF	2	2
CASSDSSGRADTQYF	12	5
CASSFDPYEQYF	12	2
CASSLGLGQNTGELFF	1	1
CASSLVGGSGANVLTF	1	2
CASSQVREGTQYF	1	1
CASSVGGDLGEAFF	1	1
CASTPYPSGRRNEQFF	1	1
CAWSVSPPGEQYF	1	1
CSAPSRDFRNSPLHF	2	2
CSARGTREPYEQYF	4	2
CSVSPESAGTRETQYF	1	1

b

Extended Data Fig. 12 | Cluster annotation of single cells with shared TCRβ in post-treatment peripheral blood and TIL (*n* = 16) from patient OT6. a, Shared TCRβ identified in both PBMC and TIL for patient OT6. Rows indicate different TCRβ clonotype and columns provide the complementarity determining region 3 (CDR3) amino acid sequence and number of cells with the TCRβ CDR3 amino acid sequence in peripheral blood and TIL, respectively. **b,** Single cell cluster identities of shared TCRβ according to unique CDR3 and compartment where the TCR was identified. Piecharts are colored according to the cluster identities of all cells with the same TCRβ. The radius of each piechart is proportional to the total number of cells in which each TCRβ was identified (square root of *n* cells divided by eight).